



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

5821

C213

The Hopkins Library
presented to the
Yeland Stanford Junior University
by Timothy Hopkins.

PAIDROMUS

STUDIOSA LECTIO

ET

1888



PRODROMUS
SYSTEMATIS NATURALIS
REGNI VEGETABILIS.

PARS V.

On trouve aux mêmes adresses :

- DE CANDOLLE.** — *Prodromus systematis naturalis regni vegetabilis*; in-8. Tomi I, II, III, IV. 1824 à 1830..... 63 fr.
- Collection de Mémoires pour servir à l'Histoire du règne végétal.
- Premier Mémoire : sur la famille des *Mélastomacées*; in-4. avec 10 Planches. 1828..... 10 fr.
- Second Mémoire : sur la famille des *Crassulacées*; in-4. avec 13 Pl. 1828. 10 fr.
- Troisième et quatrième Mémoires : sur la famille des *Onagracees* et des *Paronichées*; in-4. avec 9 Planches. 1829..... 8 fr.
- Cinquième Mémoire : sur la famille des *Ombellifères*; in-4. avec 19 Pl. 1829. 15 fr.
- Sixième Mémoire : sur la famille des *Loranthacées*; in-4. avec 12 Pl. 1830. 10 fr.
- Septième Mémoire : sur la famille des *Valérianiées*; in-4. avec 4 Pl. 1832. 5 fr.
- Huitième Mémoire : sur la famille des *Cactées*; in-4. avec 12 Pl. 1834. 10 fr.
- L'auteur n'ayant pu, ni dans le *Prodromus*, ni dans le *Cours de Botanique* qu'il publie, donner sur certains points de la science les développemens qui lui paraissent nécessaires pour fixer l'attention des botanistes, a cru devoir publier séparément une série de Mémoires servant d'éclaircissement aux deux ouvrages ci-dessus mentionnés, et dont dix formeront un volume. Cet ouvrage, accompagné de planches, et de même format que les *Mémoires du Musée d'Histoire naturelle*, sera donc, à proprement parler, le *Commentaire du Prodromus*.
- Chaque Mémoire se vend séparément.
- *Regni vegetabilis Systema naturale*; sive *Ordines, Genera et Species Plantarum secundum methodi naturalis normas digestarum et descriptarum*. Volumen primum : *Sistens Prolegomena et Ordines quinque, nempe Ranunculaceas, Dilleniaceas, Magnoliaceas, Anonaceas et Menispermicas*, in-8. de 568 p. 1818. 12 fr.
- Volumen secundum : *Sistens Ordines sex, nempe Barberidas, Podophyllas, Nymphaeaceas, Papaveraceas, Fumariaceas et Cruciferas*; in-8. de 750 pages. 1821..... 15 fr.
- *Astragalologia, nempe Astragali, Biserrulæ et Oxytropidis Historia*; cum 50 tab. æneis; 1 vol. petit in-folio. 1802..... 54 fr.
- Le même ouvrage, grand in-folio, papier vélin superfin double..... 96 fr.
- JAUME SAINT-HILAIRE.** — *Exposition des Familles naturelles et de la Germination des Plantes*; 4 vol. gr. in-8. avec 117 planches en noir. 1805..... 36 fr.
- Le même ouvrage, 2 vol. in-4. Planches en couleur, papier ordinaire. 96 fr.
- Le même ouvrage, 2 vol. in-4. Planches en couleur, papier vélin..... 192 fr.
- HUMBOLDT (A.) et BONPLAND.** — *Plantes équinoxiales recueillies au Mexique, aux Andes de la Nouvelle-Grenade et au Pérou*; par MM. de Humboldt et Bonpland, avec leur description; 17 livraisons formant 1 vol. in-folio, avec 144 Pl. (Ouvrage complet)..... 500 fr.
- HOOKE (J.) et GREVILLE (R. K.).** — *Icones filicum : ad eas potissimum Species illustrandas delineatæ, quæ hactenus vel in herbariis delinuerunt prorsus incognitæ, vel saltem nondum per icones botanicas innotuerunt; cum tabulis æneis*. Le texte en latin, avec 240 planches; 2 vol. in-fol. Londres, 1731. (Ouvrage complet.) Les planches en noir..... 360 fr.
- Le même ouvrage, les planches en couleur..... 600 fr.
- WALLICH (N.)**, surintendant du Jardin botanique de Calcutta. — *Plantæ Asiaticæ rariores*, ou Descriptions et figures d'un choix de Plantes inédites de l'Inde orientale, 3 vol. in-fol. avec 300 planches très soigneusement coloriées; le texte en langue latine. Londres, 1830 à 1832. (Ouvrage complet)..... 900 fr.

PRODROMUS
SYSTEMATIS NATURALIS
REGNI VEGETABILIS,

SIVE

ENUMERATIO CONTRACTA

ORDINUM, GENERUM, SPECIERUMQUE PLANTARUM

HUC USQUE COGNITARUM, JUXTA METHODI NATURALIS NORMAS
DIGESTA;

AUCTORE AUG. PYRAMO DE CANDOLLE.

PARS QUINTA:

Sistens Calycereas et Compositarum tribus priores.

PARISIIS,

Sumptibus Sociorum TREUTTEL ET WÜRTZ,
viâ dictâ de Lille, n° 17.

VENITQUE IN EORUMDEM BIBLIOPOLIO,

ARGENTORATI.

M. DCCC. XXXVI.



HD 605

PRODROMUS

SYSTEMATIS UNIVERSALIS

REGNI VEGETABILIS.

Ordo CI. CALYCERÆ *R. Brown, diss. publ. lect. Londini febr. 1816. trans. soc. lin. Lond. v. XII (1817). p. 132-142. DC. theor. elem. ed. 2. p. 246. L. C. Rich. mem. mus. h. nat. 6. p. 77. Less. in linnæâ 1831. p. 258. — Boopideæ Cass. diss. publ. lect. Paris. aug. 1816. journ. phys. oct. 1816. opusc. phyt. 2. p. 344.*

Flores densè capitati ut in Compositis super receptaculum (ramuli apicem) sessiles, cincti involucro polyphylo, squamis basi inter se concretis, paleis bracteolisve inter flores è receptaculo ortis. Calycis tubus ovario adnatus, limbus in lobos 5 sæpius dispare partitus. Corolla regularis infundibuliformis, tubo gracili longo, lobis 5 trinerviis, nervis lateralibus submarginalibus, areolis 5 infra stamina sitis et iis alternis notata, æstivatione loborum valvatâ. Stam. 5 monadelphæ et synantherea, antheris supernè distinctis appendice apicilari destitutis. Stylus lævis filiformis apice subclavatus indivisus. Stigma capitellatum subhispidulum. Achænium lobis calycinis sæpè induratis et spinescentibus coronatum, 1-spermum. Semen albumine carnosio donatum. Embryo axilis rectus inversus radiculâ nempè superâ, cotyledonibus 2 planiusculis. — Ordo omninò medius inter Dipsaceas et Compositas, prioribus affinis ob seminis embryonem inversum intra albumen carnosum, posterioribus ob staminâ synanthereâ, nervationem loborum corollæ ferè similem, et habitum simillimum. — Circa characteres præcipuos et locum ordinis in serie naturali, omninò consentiunt cl. Brown et Cassini, qui eodem ferè tempore ordinem egregiè admiserunt.

I. GAMOCARPHA. — Boopidis sp. Poepp. Less.

Involucrum dentatum, constans squamis circ. 12 longè concretis. Fimbrillæ recept. acutæ in alveolas concretæ. Flores omnes fertiles. Calycis lobi 5 elliptici obtusiusculi. Cor. 5-fidæ 10-nerviæ. Antheræ inclusæ exalatae ecaudatæ, filamentis apice distinctis. Stylus longè exsertus tenuis apice vix capitatus. Fruct. ign. — Herba Chilensis Valerianam Saliuncam aut globulariæfoliam habitu referens.

1. G. POEFFIGII \mathcal{L} in Chili summis montibus. Boopis? Alpina Poepp! pl. exs. n. 823. Less. in linnæa 1831 p. 258. Herba digitalis glaberrima. Folia ad col- lum adgregata lineari-subspathulata integerrima scapi ferè dimidiam longitudi- nem æquantia. Rhizoma cylindraceum subrepens. (v. s.)

- II. BOOPIS Juss. ann. mus. 2. p. 350. t. 58. f. 2. Cass. journ. phys. aug. 1818. p. 114. opusc. phyt. 2. p. 355. L. C. Rich. diss. cal. p. 51. t. 2. mem. mus. h. n. 6. p. 78. t. 11.

Involucrum constans squamis 7-8 ad medium concretis, dentibus sæpè interjectis. Receptaculum parvum convexum paleolis filiformibus apice subdilatis inter flores onustum. Flores omnes fertiles subconsimiles. Calycis lobi ovario breviores membranacei integri aut inciso-dentati. Cor. tubus gracilis, limbus campanulatus ultra medium 5-fidus, longè exsertus simplex. Achenium hialatum? — Herbæ erectæ perennes. Folia alterna pinnatipartita. Capitula hemisphærica terminalia.

1. B. ANTHEMOIDES (Juss! ann. mus. 2. p. 350. t. 58. f. 2), tota glabra, foliis pinnato-partitis, lobis capillaceis. \mathcal{L} in Bonariâ, ad Rio de la Plata et Rio-Grande. L. C. Rich. l. c. cum ic. Cass. 5. dict. suppl. 28 op. phyt. 2. p. 356. Herba 6-8 pollic. (v. s.)
2. B. LEUCANTHEMA (Poepp! pl. exs. n. 774), caule villis laxis longisque sublanato, foliis pinnatipartitis, lobis linearibus acutiusculis. \mathcal{L} in Chili legit el. Poepp. Less. in linnæa 1831 p. 259. Herba 2-3 polli. longa. Capitula dimidio minora quam in priore cui similis. Cor. albæ 4-fidæ. (v. s.)

- III. CALYCERA Rich. diss. cal. in mem. mus. 6. p. 77. t. 10 et segr. p. 50. t. 1. Cass. dict. 6. suppl. 36. journ. phys. aug. 1818. p. 113. op. phyt. 2. p. 353. — Calicera Cav. icon. 4. p. 34. t. 358.

Involucrum squamis 5-7 vix basi concretis inæqualibus constans. Receptaculum subglobosum paleis tenuissimis setaceo-subulatis inter flores onustum. Flores fertiles dissimiles et promiscui. Cal. lobi in aliis indurati valdè elongati corniformes, in alteris squamacei ovales obtusi. Corollæ tubus gracilis, limbus obconico-elongatus infra incisuras coarctatus vix semi-5-fidus. — Herbæ perennes Chilenses. Folia integra aut grossè dentata. Capitula terminalia.

1. C. CAVANILLESII (Rich. l. c.), glaberrima, foliis dentibus grossis distantibus ferè semi-pinnatifidis, paleis supernè angusto-subulatis. \mathcal{L} in Chili. C. her- bacea Cav. l. c. Cass. l. c. Involucrum dicitur polyphyllum. Folia radicalia longissima petiolata.
2. C. BALSAMITÆFOLIA (Rich. l. c.), canle subpiloso, foliis dentibus grossis distantibus ferè pinnatifidis ad margines ciliato-puberulis, bracteolis supernè lineari-lanceolatis. \mathcal{L} in Chili. Boopis balsamitæfolia Juss. ann. mus. 2 p. 350. Cass. dict. 5. suppl. 28. Scabiosa sympaganthera R. et Pav. fl. per. 1. p. 49 t. 76. f. 2. Caulis bipedalis. (v. s. ex h. Dombey.)

- IV. ACICARPHA Juss. ann. mus. 4. p. 347. Cass. dict. 1. suppl. 3a. L. C. Rich. diss. cal. in mém. mus. 6. p. 77. t. 12. et segr. p. 31. t. 3. — *Cryptocarpa* Brown. trans. soc. lin. 12. p. 131. non Raddi. — *Sommea* Bory ann. gen. sc. phys. (1820) 6. p. 92. t. 87. — *Echinolema* Jacq. f. — *Acanthosperma* Arrab. fl. flum. 8. t. 152.

Involucri squamæ 5 basi concretæ aut liberæ. Receptaculum oblongum. Paleolæ parvæ vix conspicuæ. Flores numerosi, centrales styli abortu steriles, laterales hermaphroditi ovariiis densè concretis Cal. lobi 5, post anthesin excrescentes indurati conici spiniformes. Cor. limbo infundibuliformi. — Herbæ. Folia indivisa. Capitula oppositifolia. — Lobi calycini olim pro paleis receptaculi male habiti, undè nomen minùs aptum.

1. *A. TRIBULOIDES* (Juss. ann. mus. 2. p. 348. t. 58. f. 1), glabra erecta, foliis oblongo-cuneatis, sinuato-dentatis obtusis, inferioribus spathulatis, superioribus basi latiore amplexicaulibus, stylo longissimè exserto. ①? in Bonariâ. L. C. Rich. diss. cal. p. 18. t. 2. mem. mus. 6. p. 78. t. 11. *Cryptocarpa* tribuloides Cass. dict. 12. p. 85. atl. dict. sc. nat. ic. opusc. phyt. ic. Pedunculi axillares dicuntur, potiùs ramuli apice capituliferi. (v. s.)
2. *A. SPATHULATA* (Brown. comp. 129), glabra decumbens, foliis spathulatis basi in petiolum attenuatis mucronatis integris aut pauci-dentatis glaucescentibus, stylo brevissimè exserto. 5 in arenosis maritimis Brasiliz, ad Rio-de-Janeiro, Sanctam-Catharinam, Bahiâ. L. C. Rich. diss. cal. p. 51. t. 3. mem. mus. 6. p. 78. t. 12. *Cryptocarpa* spathulata Cass. l. c. *Sommea* Calciatrapa Bory ann. gen. sc. ph. 6. p. 92. t. 87. *Echinolema* arenarium Jacq. f! in litt. *Acanthosperma* littorale Arrab. fl. flum. 8. t. 152. *Bupthalmum* Bonariense Pers. ench. 2. p. 474? Capitula duplò quàm in priore majora. (v. s.)
β. glauca, erecta glaberrima glauca, foliis obovato-cuneatis grossè dentato-incisis attenuatis in petiolum limbo duplò ferè longiorem, stylo brevissimè exserto. ①? in Brasiliâ. Simillima *A. spathulata* sed erecta nec procumbens, foliorum petioli duplò triplòve longiores et involucri squamæ 5 nullo modo basi concretæ sed omninò liberæ! Radix in horto exilis perpendicularis herbam annuam indicat. An mera varietas, aut herba primo anno florens? (v. v. c. in h. Genev. et s. sp.)
3. *A. PROCUMBENS* (Less. in linnæâ 1831. p. 527), procumbens, multicaulis glaberrima, foliis sessilibus glaucis membranaceis obtusis mucronulatis integerrimis, invol. squamis 6 patentibus. 2? in Brasiliâ. Calycis lobi indurato-corniculati minores quàm *A. spathulata*.
4. *A. EUPHYROIDES* (Less. l. c.), erecta glabra dichotoma, foliis membranaceis reticulato-venosis obtusis subdentatis, capitulis terminalibus, involucri corollæ que 4-lobis. 2? in Brasiliz uliginosis.
5. *A. LANATA* (Lag. ex Pers. ench. 2. p. 438) caule lanato, foliis glabris linearibus dente uno alterove incisis, summis integerrimis. — in Novâ Hispaniâ. Species vix palmaris, non satùs nota.

Ordo CII. COMPOSITÆ *Vaill. act. acad. Par. 1718-1721. Linn. ord. nat. 1737. n. 21. 1764. n. 49. Berckh. exp. car. fl. comp. 1761. 4°. Adans. fam. (1763). 2. p. 103. DC. fl. fr. (1805). 4. p. 1. ann. mus. h. nat. Par. 16. p. 190. bot. gall. 1. p. 258. Brown. obs. comp. in trans. linn. soc. (1817). 12. p. 76. Kunth nov. gen. et sp. am. (1820). 4. p. 1. — Syngenesia polygamia Linn. syst. sex. 1737. cl. xix. Meese class. syng. 1768. 8°. — Compositifloræ Gaertn. fruct. (1791). 2. p. 353. — Actinophytum, Glossariphytum, et Siphoniphytum Neck. elem. vol. 1. n. I. II. et III. — Synanthereæ L. C. Rich. in Marth. cat. h. bot. 1801. p. 85. Cass. opusc. phyt. 1826. v. 1. p. 1. dict. sc. nat. (1827). v. 51. p. 443. Bartl. ord. nat. p. 120. — Compositæ seu Synanthereæ Cass. dict. sc. nat. (1818). 10. p. 131. Less. syn. 1832. in-8°. — Flosculosæ, Semiflosculosæ et Radiatæ Tourn. inst. (1694). cl. 12. 13. et 14. — Cynarocephalæ, Cichoraceæ et Corymbiferæ *Vaill. l. c. Juss. gen. pl. (1789). p. 168.**

DIAGN. Flores capitati seu glomerati. Corollæ nevramphipetalæ, nempè nervis in tubo ad sinus tendentibus, in lobis (per æstivationem valvatis) nervis marginalibus donatæ. Antheræ in tubum concretæ. Semen exalbuminosum. Embryo erectus.

CAR. FRUCT. Calyx gamosepalus ovario adhærens, tubo nunc ovario æquali (1) nunc ovarium superante (2), limbo seu pappo nunc omninò nullo aut ad marginem minimum reducto, rarissimè foliaceo, nunc sæpiùs scarioso, integro, dentato, lobato aut sæpiùs evadente in squamellas paleaceas seu setas piliformes, simplices, ramosas, denticulatas aut plumosas, uni-bi- aut multi-seriales. — Corolla summo calycis tubo inserta, gamopetala, nevramphipetala nempè petalis nervis duobus margini proximis donatis undè sæpiùs in tubo nervi prima fronte 5 (10 geminatim concreti) à basi ad sinus loborum tendentes et in lobis 10 nempè in quoque lobo marginales, adjectis hinc indè nervis accessoriis medium petalis occupantibus, in paucissimis solis ferè obviis. Tubus longitudine varius, à basi ad filamentorum liberationem productus. Faux sæpiùs ampliata à filamentorum liberatione ad separationem loborum : lobi sæpiùs 5 rariùs 4 aut 3 aut 2, per

(1) Tunc pappus seu limbus calycis abortivus incongruè dicitur *sessilis* seu potiùs *ructus erostratus*.

(2) Tunc pappus dicitur incongruè *stipitatus* seu potiùs *fructus rostratus*.

æstivationem valvati, nunc omnes æquales, nunc subæquales aut irregulares inæquales seu palmati, nunc bilabiati (aut 3 in labio exter. 2 in interiore, aut 4 in exter. 1 in inter.), nunc fissurâ ad latus interius productâ in ligulam 5-dentatam planam concreti. — Stamina 5, rariùs 4, in floribus fœmineis neutrisve nulla aut rudimentaria; filamenta lobis corollæ alterna et ideò cum nervis tubi congruentia, tubo corollæ adnata, supernè libera, aut inter se omninò distincta aut in tubum concreta et ideò monadelphæ, versus apicem articulata, articulo superiore antherifero connectivi locum tenente. Antheræ erectæ uniseriales in tubum stylo perforatum concretæ (*syngenesæ* seu *synantheræ*) lineares, biloculares, rimâ longitudinali intùs dehiscèntes, 4-valves, valvis extroflexis, apice in appendicem seu alam formâ et magnitudine in diversis variam, sæpiùs à parte polliniferâ consistentiâ distinctam, desinentes: basi sæpè in caudas duas longitudine et formâ subvariâs productæ. Pollen globosum aut ellipticum echinulatum aut læve. — Ovarium 1-carpellare calyci adhærens, 1-ovulatum. Stylus teres aut rariùs basi tumidus (tunc bulbosus dictus) in floribus masculis sæpiùs (ubi adest) simplicissimus, in fœmineis et hermaphroditis supernè bifidus; rami (seu vulgo stigmata) supernè plani, subtùs convexi, nunc omninò liberi nunc inter se plùs minùs concreti. Glandulæ stigmaticæ (stigmata proprio sensu) biseriales, ad paginam superam ramorum styli sitæ, seriebus continuis marginalibus scabriusculis magis minùsve prominulis distinctisve. Pili pollinis collectores rigidi in florum hermaphroditorum parte styli superâ, aut in ejus ramis ad latus inferius aut ad apicem siti, in floribus fœmineis neutrisve subnulli.

Fructus seu achænium constat calycis tubo, pericarpio, seminisque spermodermio inter se plùs minùs arcè concretis et embryonem includentibus. Achænium ideò 1-loculare 1-spermum, supra receptaculum articulatam sæpiùs sessile rariùs stipitatum, basi areolâ basilari aut laterali instructum, apice erostre aut rostratum, pappo suprâ descripto superatum, disco epigyno nectarium centrale cum ovario continuum styloforme aut alveolare cingente terminatum. Semen funiculo brevissimo imæ fructus parti adfixum erectum basi latere funiculo opposito affixum. Endoplevra seu spermodermii lamina interior (albumen Less.) crassiuscula tenera diaphana funiculo bifido percursa. Embryo erectus, rectus exorhizus.

dicotyledoneus, radiculâ brevi rectâ, plumulâ plerumque inconspicuâ, cotyledonibus planis dorso sæpè subconvexis, rarissimè curvatis, casu interdum ternis.

INFL. Flores collecti in *capitulum* (1) aut in *glomerulum* (2). Capitulum constat floribus pluribus super receptaculum (3) seu rami apicem nunc planum nunc conicum elongatumve, exterioribus seu inferioribus præcocioribus, intimis sensim serioribus. Flores aut omnes hermaphroditi et tunc capitula *homogama*, aut exteriores neutri seu fœminei, interiores hermaphroditi seu masculi et tunc capitula *heterogama*, aut omnes in eadem plantâ aut masculi aut fœminei et tunc capitula nunc *monoïca*, nempe floribus masculis et fœmineis, nunc *heterocephala* nempe in eadem plantâ alia mascula alia fœminea, nunc *dioïca* nempe in eadem plantâ omnia mascula seu omnia fœminea. Capitula aut corollis omnibus tubulosis donantur et tunc *discoïdea* seu *flosculosa* dicuntur, aut corollis omnibus ligulatis et tunc *ligulata* seu olim *semiflosculosa*, aut corollis in margine seu radio ligulatis in centro seu disco tubulosis et tunc *radiata* nuncupantur, aut omnibus bilabiatis et tunc *falso-discoïdea*, aut margine ligulatis centro bilabiatis et tunc *falso-radiata* seu *radiatifor-mia*. Capitula discoïdea et falso-discoïdea exhibent interdum flores marginales cæteris similes sed majores, et tunc *coronata* dicuntur.

Involucrum (4) constat *squamis* (5) plurimis, uni-aut pluri-serialibus, liberis aut (ubi uniserialibus) margine coalitis, siccis scariosis coriaceis carnosissimis spinosis foliosisve, sæpè in squamam propriam et appendicem distinctis, seriebus æqualibus aut sæpius inæqualibus imbricatis aut calyculatis aut variè et subinordinate elongatis.

Glomerulus constat capitulis uni-aut paucifloris involucri proprio instructis, dense et variè aggregatis, involucri gene-

(1) *Flos compositus* Veterum. *Calathis* Mirb. *Anthodium* Ehr. *Cephalanthium* Rich.

(2) *Capitulum* Cass.

(3) *Phoranthium* Rich. *Clinanthium* Cass. *Rachis* Less.

(4) *Calyx communis* Veterum. *Perigynanda communis* Neck. *Periclinium* Mirb. *Periphoranthium* Rich.

(5) Plurimi squamas vocant foliola, sed squamarum terminus aptior mihi videtur 1°. Quia vox foliolum sensu diversissimo jam gaudet in foliis compositis. 2°. Quia squamæ involucri omnino repræsentant squamas gemmarum et præcipuè gemmarum floriferarum. 3°. Quia jam in pluribus aliis ordinibus partes involucri sunt dictæ squamæ.

rali cinctis, super receptaculum generale insidentibus, sæpius centralibus præcocioribus, exterioribus senioribus aut inordinate florentibus.

Receptaculum capituli nunc *paleaceum*, nempe *paleis* (1) involucris squamis similibus ad latus cujusque floris externum sitis, super basin articulatis et in folia vera hinc inde monstrosè expansis donatum, nunc *semipaleaceum* nempe *paleis* in parte exteriori tantum donatum, nunc *epaleaceum* seu *nudum* nempe *paleis* omnino orbatam. In aliis capitulis (seu fortè glomerulis arctissime cinctis) receptaculum dicitur nunc *fimbrilliferum*, nempe ubi flos singulus ex omni parte cinctus est alveolâ basi squamosâ apice sæpè in laminas setasve irregulares desinente: nunc *alveolatum* nempe ubi alveolæ breves in setas non productæ sed sæpè denticulatæ: nunc tandem *areolatum* ubi nempe areola pentagona (alveolæ rudimentum) singuli floris basin circumdat. Fortè receptacula plus minus paleacea flores verè spicatos et capitula proprio sensu designant, et receptacula fimbrillifera aut alveolata indicant capitula arctissime cincta proprio sensu flores umbellatos et quorum alveolæ involucelli rudimenta repræsentant. Clariùs dicam capitula aut more spicæ contractæ aut more umbellæ contractæ disposita.

VEG. Herbæ sæpius perennes, aut frutices aut rariùs (præsertim in insulis) arbores, per totum orbem terrarum obviæ et decimam ferè partem regni vegetabilis comprehendentes. Folia alterna aut opposita formâ incisuris aut aspectu diversissima, semper simplicia nec composita. Caulis (ubi simplex) aut rami capitulo glomerulove terminati. Rami sæpius corymbosi, centrales præcociores. Folia sub capitulis glomerulisve sita à cæteris sæpè diversa et floralia seu bracteantia dicta, nunc gradatim in squamas mutata nunc abruptè squamis multò majora. Corollæ nunc xanthicæ nunc cyanicæ (2) in eodem capitulo aut *homochromæ* seu concolores aut *heterochromæ* seu discolores; ubi discolores ligulæ semper cyanicæ et tubuli xanthici! imò tubuli xanthici cum ligulati monstrosè evadunt, cyanici fiunt! Succus proprius varius, interdum lacteus. Sapor sæpius amarus, adstringens et aromaticus.

(1) Aliis *Bracteolæ* dictis. Vox *paleæ* mihi servanda videtur quia notissima et quia squamæ extimæ receptaculi seu accessorizæ à plurimis et quidem commodè bracteolæ dicuntur.

(2) Vid. Physiol. veget. 2. p. 907.

Obs. Ordo omnium naturalissimus et vastissimus et ideo difficillimus, à Tournefortio facilè sed artificialiter, à Vaillantio naturalius sed difficilius, à Linnæo eleganter sed arbitrariò, ab Adansonio nimis vagè, à Gærtnero nimis carpologicè, à Cassinio strenuè et philosophicè sed sæpè obscurè, à Lessingio lucidè et simpliciter dispositus, summam adhuc botanicorum attentionem requirit. An capitula alveolis (nec paleis) donata, ritè (Albertiniâ præsertim suadente) cum glomerulis veris consocianda? An pappus qui evidenter in plerisque generibus à limbo calycis variè abortivo provenit, fortè in quibusdam (v. gr. Dipterocome) etiam ex pericarpio proprio elongato provenire posset? An pars subcarnosa embryonem circumdans albuminis veri aut endoplevræ locum tenet? An fila è basi styli orta et per partem istam carnosam descendunt basin embryonis attingunt et fœcundationis principium conducunt? Quare in capitulis heterogamis flores masculi si adsint semper centrales et fœminei semper marginales? Cur corollæ tubulosæ xanthicæ in seriem cyanicam transeunt cum ligulatæ aut ampliatae evadunt? An achæonii rostrum ubi adest, à calycis tubo tantum aut etiam è pericarpio vero producto formatur?

CONSPECTUS ORDINIS.

* *Tubulifloræ*. Nempè floribus hermaphroditis tubulosis regulariter 5-(rarius 4-) dentatis.

Trib. I. VERNONIACEÆ. Stylus fl. herm. cylindræus, ramis sæpius subulatis elongatis, rarius obtusis brevibus, semper æqualiter et longiusculè hispidis. Stigmatis series antè medios styli ramos desinentes.

Trib. II. EUPATORIACEÆ. Stylus fl. herm. cylindræus, ramis longis subclavellatis extus supernè puberulo-papillois. Stigmatis series parum prominulæ antè medios styli ramos sæpius desinentes.

Trib. III. ASTEROIDEÆ. Stylus fl. herm. cylindræus, ramis linearibus extus subplanis et supernè æqualiter et minutè puberulis. Stigmatis series prominulæ usque ad originem pubis exterioris circiter extensæ.

Trib. IV. SENECTIONIDEÆ. Stylus fl. herm. cylindræus, ramis linearibus apice penicillatis nunc truncatis nunc ultra penicillum, in comum brevem aut appendiculum elongatum angustum et hispidulum productis. Stigmatis series latiusculæ et prominulæ usque ad penicillum extensæ.

Trib. V. CYNARÆÆ. Stylus fl. herm. supernè nodoso-incrassatus et sæpè ad nodum penicillatus, ramis nunc concretis nunc liberis extùs puberulis. Stigmatis series haud prominulæ apicem ramorum attingentes et ibi confluentes.

** *Labiatifloræ*. Nempè floribus hermaphroditis sæpissimè bilabiatis.

Trib. VI. MUTISIAEÆ. Stylus fl. herm. supernè cylindraceus aut subnodosus, ramis sæpiùs obtusis truncatisve extùs valdè convexis et in parte superiore pube minutâ æquali (rariùs nullâ) vestiti.

Trib. VII. NASSAUVIAEÆ. Stylus fl. herm. nunquam nodoso-incrassatus, ramis linearibus longiusculis truncatis apiceque solo penicillatis.

*** *Ligulifloræ*. Nempè floribus omnibus hermaphroditis ligulatis.

Trib. VIII. CICHORACEÆ. Stylus supernè cylindraceus, ramis longiusculis obtusiusculis æqualiter pubescenti-hirtellis. Stigmatis series antè medios styli ramos desinentes.

N. B. Confer ad frontem tribùs cujusque subtribuum divisionumque conspectum methodicum et ad calcem totius ordinis 1°. indicem generum dichotomum, et ad tyronum faciliorem usum, artificialem: 2°. indicem nominum alphabeticum.

Trib. I. VERNONIAEÆ *Less. in linnæd* 1829. p. 241. 1830. p. 134. 1831. p. 625. *syn. p.* 145. — *Vernoniæ* (excl. sect. 2 et 3) et *Tagetineæ* *Pectideæ* *Cass. tabl. II. XX opusc. I. p.* 333. 2. p. 206. *dict. 20. p.* 384. 36. p. 19. 57. p. 359.

Stylus supernè cylindraceus, ramis æqualiter hispidis in plurimis longis subulatis in paucissimis brevibus obtusisve; stigmatis glandulæ dispositæ in series prominulas angustas antè medios styli ramos desinentes. Corolla 5-fida sæpiùs regularis interdum obscurè bilabiata aut palmata. Pollen globosum echinulatum. — Capitula uni-pauci- aut multiflora, sæpiùs homogama discoidea, floribus omnibus hermaphroditis aut in unico genere dioicis, rariùs heterogama radiata floribus radii ligulatis foemineis. Recept. alveolatum fimbriiferumve aut nudum, in paucissimis (dubiis?) paleaceum. Flores cyanici (purpurascens, cœrulescentes aut albi) nec ferè unquam xanthici nisi in *Pectidearum* subtribu. — Tribus, jam rectè suadentibus cl. Cassini et Lessing, media inter *Cichoraceas* et *Eupatorineas*, sed etiam mediantibus *Pectideis* ad *Senecioneas*-*Tagetineas* affinis.

CONSPECTUS TRIBUS.

Subtrib. I. VERNONIEÆ. Capitula discoidea homogama. (Gen. I-XLVIII.)

Div. I. EUVERNONIEÆ. Antheræ ecaudatæ. Involucrum non compressum polyphyllum. (Gen. I-XXXV.)

Subdiv. 1. *ETHULIÆ*. Pappo nullo aut simplici coroniformi. (Gen. I-V.)

Subdiv. 2. *HETEROCOMEÆ*. Pappo uni- bi- aut pluriseriali, omnibus aut saltem interiore setiformi aut angustissime ligulato. Capitulis non fasciculatis. (Gen. VI-XXVI.)

Subdiv. 3. *ALBERTINIEÆ*. Pappo prioris subdiv. Capitulis sæpius oligocephalis in glomerulum fasciculatis. (Gen. XXVII-XXXV.)

Div. II. *ELEPHANTOPEÆ*. Antheræ ecaudatæ. Involucrum compressum, squamis alternis conduplicatis. (Gen. XXXVI-XXXVIII.)

Div. III. *ROLANDREÆ*. Capitula 1-flora. Invol. oligophyllum (1-5), squamis sæpius concretis. Antheræ ecaudatæ. (Gen. XXXIX-XLIV.)

Div. IV. *BOJERIEÆ*. Antheræ bicaudatæ. (Gen. XLV-XLVIII.)

Subtrib. II. *PECTIDEÆ*. Capitula radiata heterogama. (Gen. XLIX-LVIII.)

Div. I. *LIABEÆ*. Cor. disci regulariter 5-fidæ. Styli rami elongati. Folia eglandulosa eciliata. (Gen. XLIX-LIV.)

Div. II. *EUPECTIDEÆ*. Cor. disci sæpè irregulares. Styli rami abbreviati. Folia glandulosa basi sæpius ciliata. (Gen. LV-LVIII.)

Subtrib. I. *VERNONIÆ*. *Cass. trib. XX. gen. paucis excl. aut add.*

Capitula discoidea nec radiata. Styli rami elongati acuminati. — Folia alterna rarius opposita. Flores albi, cœrulei, aut purpurei.

Div. I. *EUVERNONIÆ*. — *Vernoniæ* prototypæ *Cass. tabl. xx. sect. iv.* — *Vernoniæ* pleræque *Less. in linnæâ 1830. p. 136. syn. 146. excl. Liabeis et Pectideis.*

Capitula discoidea multiflora (in paucissimis pauciflora imò 1-flora), Antheræ ecaudatæ. Involucrum teres multiseriale imbricatum. Recept. nudum aut alveolatum imò fimbrilliferum nec forsitan unquam vere paleaceum (excl. fortè *Heterocoma*?) Cor. tubulosa regularis aut subpalmata. Styli rami semisubulati. — Folia alterna (in paucissimis opposita).

Subdiv. I. *ETHULIÆ*.

Pappo nullo aut coroniformi uniseriali. Capitulis uni- aut multifloris, distinctis nec in glomerulum concretis.

I. *ADENOCYCLUS* *Less. in linnæâ 1829. p. 337. f. 39-41. syn. p. 151.*

Capitulum 1-florum. Involucrum oblongo-cylindraceum imbricatum, squamis coriaceis siccis 1-nerviis interioribus multò longioribus. Recept. punctiforme. Cor. regularis, limbo à tubo haud distincto profundè 5-fido, lobis acuminatis parte integrâ glabrâ brevioribus. Anth. inclusæ. Filam. glabra lævia. Achænium obconicum breve sulcatum glaberrimum, nectario alveolari, disco epigyno orbe glandularum notato. Pappus 0. — Frutex Odontolomæ habitu. Rami juniores petiolique subcanescentes. Folia alterna petiolata ovato-oblonga acuminata sub-

integerrima glabra. Corymbi terminales pluriès dichotomi condensati. Cor. rosææ.

- I. *A. CONDENSATUS* (Less. l. c.). 5 in insulâ Trinitatis. Eupatorium condensatum Reich. in Sieb! fl. exs. trin. n° 73. (v. s.)

II. *ODONTOLOMA* H. B. et Kunth. nov. gen. 4. p. 43. t. 319. Juss. dict. 35. p. 395. Cass. tabl. xx. n. 46. Less. in linnæâ 1829. p. 337. syn. p. 147. non Pers.

Capitulum 1-florum. Invol. cylindraceum imbricatum, squamis siccis 1-nerviis interioribus longioribus. Recept. punctiforme. Cor. regularis glabra 5-fida, lobis acuminatis parte integrâ brevioribus. Fil. lævia. Styli rami semi-subulati. Achæmium breve glaberrimum, nectario alveolari. Pappus 1-serialis coroniformis coriaceus brevis apice inæqualiter et acutè multifido-dentatus. — Arbuscula, ramis fuscescenti-tomentosis, foliis sparsis petiolatis ovatis integerrimis acuminatis suprà glabris, subtùs pulverulento-tomentosis. Corymbi terminales subsessiles ramosissimi. Capitula fasciculato-congesta. Cor. albidæ?

- I. *O. ACUMINATUM* (H. B. et Kunth. l. c.). 5 in convalli Caracasana. Eupatorium cornifolium Willd. herb. ex Less.

III. *OIOSPERMUM* Less. in linnæâ 1829. p. 339. f. 69. 71. 72. 78. syn. p. 148. — *Ethulia* sp. Nees. *Calydermi* sp. Spreng.

Capitulum multiflorum æqualiflorum. Involucrum imbricatum patens teres, squamis scariosis glabris acutis, basi foliis inæqualibus capitulo longioribus cinctum. Recept. latum nudum. Cor. regularis, limbo 5-fido à tubo vix distincto, lobis acuminatis parte integrâ brevioribus. Anth. inclusæ. Filam. lævia. Achæmium apice rotundatum glandulosum multicosatum, disco epigyno parvo. Pappus o. — Herba ramosa glabriuscula adscendens basi repens. Folia alterna petiolata ovato-lanceolata utrinquè attenuata serrata. Capitula solitaria longè pedunculata oppositifolia. Cor. pallida.

- I. *O. INVOLUCRATUM* (Less. l. c.), caule glabrinsculo, foliis petiolatis ovatis membranaceis glabris, corollâ hispidiusculâ, achænio turbinato-ovato pubescenti-hispidulo longiore. (†) in Brasiliæ arenosis circa San Pedro de Alcantara. *Ethulia* involucrata Nees et Mart. nov. act. nat. cur. 12. p. 3. *Amphirephis psilocarpa* Nees et Mart. ibid. p. 4. *Calydermos repens* Spreng. nov. entd. 1. p. 278. obs. ex ipso (syst. 3. p. 458) manca. (v. s. comm. à Mus. reg. Berol.)

2. *O? WIGHTIANUM*, caule villosissimo, foliis sessilibus ovatis acuminatis basi attenuatis utrinquè (præcipuè in nervis) villosissimis, corollâ glabrâ, achænio gracillimo brevior. — In Indiâ orientali legit cl. Wight. *Amphirephis Wightiana* Wall! cat. comp. n. 68. Stylus ex antherarum tubo non exsertus simplex glaber fortè abortivus. Flores in capitulo circiter 100 gracillimi. An genus proprium? (v. s. com. ab ampl. cœtu merc. angl. Ind. or.)

IV. *SPARGANOPHORUS* Vaillant. act. soc. par. 1719. p. 368. Gærtn. fr. 2. p. 396. t. 165. Cass. dict. 50. p. 71. Less. in linn. 1829. p. 335. syn. p. 147. non Lag. nec Michx. — *Struchium* P. Brown. jam. 312. t. 34. f. 2. Juss. gen. 184. — *Athenæa* Adans. fam. 2. p. 121. non Schreb. — *Ethulia* sp. Lin.

Capitulum multiflorum æqualiflorum. Involucrum teres imbricatum squarrosum, squamis siccis. Recept. nudum. Cor. regularis 3-fida rarisimè 4-fida, lobis acuminatis, tubo et fauce subcontinuis brevioribus. Antheræ inclusæ. Filam. lævia. Styli rami semi-subulati. Achæmium obliquè obpyramidatum 3-5-costatum glabrum glandulosum, collo apicali magno. Pappus 1-serialis brevis coroniformis crenulatus cartilagi-

neus. — Herbæ annuæ glabræ ramosæ. — Folia alterna elliptica utrinquæ acuminata brevè petiolata serrata. Capitula ad axillas sessilia.

1. *S. VAILLANTII* (Gærtn. l. c.), foliis distanter serratis. (I) in subhumidis, fossis et ripis graminosis, Jamaica, Sancti-Thomasii, Porto-Ricci, Guadalupæ, Guayaquil, Brasilæ; nec ut videtur in Indiâ orient. ut dixit Vaill. l. c. *Strachium* Brown. l. c. Sparg. *Vaillantii* et Sparg. *Strachium* Pers. ench. 2. p. 398. Steud. nomen. phan. n. 801. excl. syn. Adams. et R. Br. *Ethulia sparganophora* Lin. sp. 1171. Arrab. fl. flum. 8. t. 90. *Ethulia Strachium* Swartz. ! flor. ind. occ. 1297. *Strachium Americanum* Poir. dict. 7. p. 475 (v. s.)
2. *S. AFRICANUS* (Steud. nom. phan. 801) foliis obtusè repandis dentatisve. (I) in Africâ Ovariensi et Senegalensi. *Strachium Africanum* Beauv ! fl. Ow. et Ben. 1. p. 80. t. 48. Herba priori simillima nec forsân specificè distinguenda. Capitula in utrâque variant solitaria gemina aut in axillis 3-4. (v. s.)

† *Species exclusæ.*

- S. AGERATOIDES* Lag. = *Ageratum cælestinum*.
S. OBTUSIFOLIUS Lag. = *Ageratum conyzoides*.
S. VERTICILLATUS Mich. = *Sclerolepis verticillatus*.
S. SONCHIFOLIUS Spr. = *Cephalophorus sonchifolius*.

V. ETHULIA Cass. dict. 15. p. 7. et 487. Less. in linn. 1829. p. 336. f. 26. 27. 35. et 43. syn. p. 148. non Gærtn. — *Kahiria* Forsk. pl. æg. 153. — *Ethulia* spec. Lin. — *Pirarda* Adans. fam. 2. p. 499 non Cass. — *Leighia* Scop. non Cass.

Capitulum multiflorum æqualiflorum. Invol. teres, pluriseriale, squamis subfoliaceis acutis. Recept. nudum. Cor. regularis, limbo campanulato 5-fido, laciniis acuminatis parte integrâ brevioribus. Anth. inclusæ. Styli rami semisubulati. Achænium obpyramidatum 4-quetrum 4-costatum glabrum glandulosum apice truncatum 4-5-gonum. Pappus minimus coroniformis carnosus integerrimus. — Herbæ ramosæ erectæ. Folia alterna oblonga utrinquæ attenuata serrata aut integra pellucido punctata. Capitula Pisi magnitudine in corymbum disposita. Cor. rosæ aut purpureæ.

* *Genuinæ, foliis non decurrentibus.*

1. *E. CONYZOIDES* (Linn. sp. 1171 Lin. f. decad. 1. t. 1), ramis corymbosis subdivergentibus. Foliis elliptico-lanceolatis sæpius grossè serratis. (I) Cor. purpureæ. Variat foliis grossè serratis rarissimè subintegerrimis plus minùs distinctè pellucido-glandulosis.
 - a. *Indica*, caulibus foliisque glabriusculis. — in Indiâ orientali ad Borruguya, Barsi, etc. (Roxb ! Wall !) *Eth. ramosa* Roxb ! herb. (v. s. sp. et v. c.)
 - β. *Africana*, caulibus foliisque adpressè villosis. — In Ægypto ad ripas Nili (Forsk ! Delil !) in Senegaliâ (Rouss !) in paludosis Albredæ ad ripas flum. Gambie (Perrot !). *Kahiria* Forsk. pl. æg. descr. 153. *Eupatorium areolatum* Perrottet ! in litt. Herba major et magis villosa quàm var. α : fortè specificè distinguenda ? Specimina glabriora et subintegerrima. In Africâ australi et orient. ad Omsamwubo et Omsanculo legit cl. Drege. n. 5065 ! (v. s. sp.)
2. *E. GRACILIS* (Delil. in Caill. voy. t. 64 et in descr. pl. Caill. p. 44. t. 3. f. 5), ramis strictis paniculatis, foliis lanceolatis integerrimis aut parcè serratis. (I) In Sennar ad El-Zerebyn (Cailland et Acerbi !) Cor. rosæ. Herba piloso-asperula. Folia distinctè pellucido-punctata. (v. s. comm. à cl. Acerbi.)
3. *E. ANGUSTIFOLIA* (Bojer ! in litt. 1835) ramis corymbosis apice adpressè pubescenti-villosis, foliis elongato-lineari-lanceolatis subintegerrimis, invol. squa-

mis obtusiusculis. ① in ins. Madagascar ad agros cultos legit cl. Bojer. Cor. purpureæ. Duabus prioribus similis, sed capitula, etiamsi numero florum simili 20-25, composita, duplò minora et invol. squamæ subobtusæ nec acuminatæ. (v. s. comm. à cl. inv.)

** *Spuriæ, foliis decurrentibus.*

4. E? *GARIEPINA*, suffruticulosa ramosissima glabra, foliis oblongo-linearibus integris aut remotè serratis in alas angustas integerrimas decurrentibus, capitulis ad apices ramorum glomerato-capitatis. 5 in Africâ australi ad flumen Gariep legit cl. Drege n. 2721! Habitus E. gracilis, sed caulis suffrutescens. Invol. squamæ acutæ. Capitula fortè dioica. Styli fl. masc. ? simplices, fl. scem. bifidi, ramis hispidulis. Achænia minima ex statu immaturo calva. An genericè separanda? Ex capitulis junioribus non asserere queo. (v. s.)
5. E? *BRASILIENSIS* (Link. enum. hort. Berol. 2. p. 310) caule alato, foliis lanceolatis acutis serratis pubescentibus decurrentibus, floribus subcorymbosis. 7 in Brasiliâ! †.

†† *Species exclusæ.*

- E. *AGERATOIDES* Spreng. = *Alomia ageratoides*.
 E. *BIDENTIS* Linn. = *Flaveria contrayerba*.
 E. *AURICULATA* Thunb. = *Dichrocephala Capensis*.
 E. *DIVARICATA* Linn. = *Epaltes divaricata*.
 E. *INVOLUCRATA* Nees et Mart. = *Oiospermum involucratum*.
 E. *INTEGRIFOLIA* Don. = *Cotula*.
 E. *PANICULATA* Houtt. = *Hippia integrifolia*.
 E. *REPANDA* Nee. = *Cephalophora glauca*.
 E. *SPARGANOPHORA* Linn. = *Sparganophorus Vaillantii*.
 E. *STRUCHIUM* Swartz. = *Sparganophorus Vaillantii*.
 E. *TOMETOSA* Linn. = *Artemisia lavandulæfolia*.
 E. *UNIFLORA* Michx. = *Sclerolepis verticillatus*.

VI. *HERDERIA* Cass. dict. 6o. p. 586 et 599.

Capitulum multiflorum discoideum homogamum. Recept. planum nudum. Invol. campanulatum biseriale, squamis oblongis æqualibus foliaceis. Cor. 5-fida, lobis extus glandulosus. Stigmata ut in *Ethuliâ*. Achænia tri-rariùs-tetraquetra prismatica in invol. densè conferta, glabra, angulis nerviformibus, annulo cupulari apice coronata. Pappus biformis uniserialis, paleis 8-10, crassis oblongis obtusis, imò interdum emarginatis ciliatis, demùm stellato-patentibus constans, et setis 2-3 elongatis rigidis scabris ex angulis achæniis ortis constans. — Herbæ Senegalenses ripariæ, ramosæ sæpè ramosissimæ. Folia alterna obovata obtusa apice subdenticulata. Capitula terminalia solitaria. Flores purpureo-violacei, Genus affine *Ethuliæ* sed pappo facilè diversum.

1. *H. TRUNCATA* (Cass. l. c.). ①? 7? secus flumen Senegal legit cl. Perrottet et quatuor varietates aut species observavit.

α. *leptorhiza*, radice exili simplici caule erectiusculo foliis obovato oblongis, nascentibus ramulisque subvillosis adultis glabris. ①? in demissis et argillosis secus flumen Senegal in regione Walo post fluv. inundationem mense decembri. Fortè sequentis ætas junior (v. s.)

β. *adscendens*, radice obliquâ fibrillosâ caule adscendente ramoso foliis obovato oblongis, nascentibus ramulisque subvillosis, adultis glabris. 7? in argillosis cum priore. (v. s.)

γ. *multicaulis*, radice crassâ sublignosâ tereti verticali, caulibus è collo

innumeris diffusis ramosis obovatis, foliis ramulisque junioribus subvillosis, adultis subglabris 2? in inundatis argillosis cum prioribus. (v. s.)
 .. *villosa*, tota etiam adulta villosissima ferè lanata. 2? cum priori (v. s. omnes comm. à cl. Perrottet.)

Subdiv. 2. HETEROCOMEÆ.

Pappo uni- bi- aut rariùs pluri-seriali, interiore semper setiformi, exter. setiformi squamellato aut rarissimè coroniformi; capitulis uni- bi- aut multifloris distinctis nec in glomerulum concretis.

VII. PACOURINA *Aubl. guian.* 2. p. 800. t. 316. *Kunth nov. gen.* 4. p. 23. — *Pacourinopsis Cass. bull. philom.* 1817 septemb. p. 151. *dict. sc. nat.* 37. p. 212. *Less. in linnæa* 1829. p. 318. — *Pacourina* sect. *Vernonia Less. in linnæa* 1831. p. 680. *Meisteria Scop. introd. non Gmel. nec Raesch. — Haynea Willd. spec.* 3. p. 1787. non *Schum.*

Capitulum globosum, multi- et æqualiflorum. Involucrum floribus subæquale imbricatum, squamis latis ovali-oblongis parallelinerviis margine membranaceis, exterioribus foliaceis spinâ terminatis, inter. siccis obtusis. Recept. planum nudum carnosum. Cor. regularis, tubo brevi, fauce cylindricâ, 5-lobis subacuminatis parte integrâ brevioribus. Anth. inclusæ. Styli rami glabriusculi. Achænia cylindracea longè glabra, cortice crasso enervio. Pappus brevis pluriserialis, pilis rigidis crassiusculis denticulatis, interioribus basi subconcretis. — *Herbæ glabræ. Caulis teres. Folia alterna ovalia aut oblonga in petiolum marginatum subamplexicaulem attenuata. Capitula oppositifolia aut lateralialia? sessilia solitaria. Cor. purpureæ. — Genus à Vernoniâ diversissimum habitu toto, achæniî formâ et pappo.*

1. *P. EDULIS* (Aubl. l. c.), foliis ovali-lanceolatis dentibus setiformibus distantibus hinc indè serratis, capitulis oppositifoliis, involucri squamis exterioribus obtusis spinulâ mucronatis. — In *Cayennâ*. Folia utrinquè glandulis innumeris opacis notata. *Pacourinopsis integrifolia Cass! Dict.* 37. p. 212. *Less. in linnæa* 1829. p. 318. *Vernonia* n° 142. *Less. in linnæa* 1831. p. 681. *Haynea edulis Willd. l. c.* (v. s. comm. à mus. Par. et sem. mat. in herb. Desf.)
2. *P. CRASSIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 30.) foliis oblongis acuminatis grossè denticato-spinosis, capitulis lateralibus? involucri squamis exterioribus spinâ terminatis 2? in humidis prope *Guaïaquil Quitenaïum*. *Pacourinopsis dentata Cass. l. c. Vernonia* n° 143. *Less. in linnæa* 1831. p. 681.

+ *Species exclusa.*

Haynea pedunculata Spr. = Vernonia pedunculata.

VIII. HETEROCOMA *DC. ann. mus.* 16. p. 191. t. 7. *choix mém.* 2. p. 34. t. 3. *Less. in linnæa* 1829. p. 339 *syn. p.* 150. *Serratulæ spec. Peri.*

Capitulum multiflorum æqualiflorum. Involucrum campanulatum subterse imbricatum, squamis coriaceis siccis plurinerviis, interioribus sublongioribus subpungentibus. Recept. paleis involucri squamas referentibus caducis onustum, et alveolis i-floris ad palearum axillis notatum! Cor. regularis, tubo tereti, laciniis 5 linearibus tubo longioribus. Anth. subexsertæ. Styli rami semi-subulati. Achænium oblongo-cylindraceum glabrum, disco epigyno magno, nectario alveolari. Pappus biserialis, serie exteriori brevi coronæformi cartilagineâ crenatâ, interiori setis filiformibus apice serratis longis rigidis caducis. — Suffrutices *Brasiliani*. Folia alterna. Capitula ad axillas sessilia. Fortè capitulum

potius dicendum esset glomerulus capitulis 1-floris ad bractearum axillas ortis constans: paleæ recept. potius bractæ: alveolæ involucri proprii rudimentum?

1. *H. ALBIDA* (DC. 1. c. deser. et icon.), foliis semi-amplexicaulibus oblongo-lanceolatis utrinquè tomento adpresso brevissimo albidis, corollis glabris. 5 in Brasiliâ Less. in linn, 1829. p. 340. f. 50. 51. 52. Cuius foliis semi-amplexicaulibus, etc. Vell. in Roëm. script. lus. et br. 141. *Serratula albida* DC. in Pers. ench. 2. p. 391. (v. s.)

† *Species exclusæ.*

H. BIFRONS DC. = *Chronopappus bifrons*.

H. PEDUNCULATA Steud. = *Hololepis pedunculata*.

- IX. *VERNONIA* Schreb. gen. 2. p. 541. Swartz fl. Ind. occ. 3 p. 1317. t. 16. Lessing in linnæâ 1829. p. 244 syn. p. 146. (excl. spec. plur.) *Vernonia*, *Ascaricida*, *Gymnanthemum*, *Oliganthes*, *Lepidaploa* et verosimiliter *Achyrocoma* et *Centrapalus* Cass. dict. — *Vernonia* et *Pollalesta* H. B. et Kunth.

Capitulum pluri- seu multiflorum æqualiflorum. Involucrum imbricatum floribus brevius, squamis interioribus longioribus. Recept. nudum aut rarius alveolato-subfimbriolatum. Cor. regularis 5-fida, lobis parte integrâ subæqualibus. Filam. lævia. Achænium callo basilari cartilagineo, disco epigyno magno. Pappus sæpius biserialis, serîe interiore setiformi externam sæpè paleaceam longè superante, rarius biserialis æqualis. — Herbæ erectæ aut frutices imò ferè arbores habitu vario. Folia alterna (in paucissimis opposita) sæpè glandulosa. Inflorescentia varia sæpè scorpioidea. Flores in capitulis numero varii, in unicâ solitarii, in plurimis 3-4, in plerisque 6. Cor. purpureæ, rosææ aut albæ, nunquam flavæ, ferè in omnibus glabræ. Genus vastissimum aut fortè dividendum, aut plerisque sequentibus augendum! Habitum præcipuè secutus sum in distributione genericâ admissâ forsân ulteriori examini subijciendâ.

Sect. I. *ACILEPIS* Don prod. fl. nep. 169.

Capitula multiflora sessilia. Invol. squamæ multiseriales imbricatæ subpungentes. Recep. alveolatum. Achænia angulato-villosa. Pappus biserialis, serîe externâ brevissimâ. Herbæ Indicæ.

1. *V. RIGIOPHYLLA*, caule tereti pubero, foliis obovato-oblongis basi attenuatis apice acutis serrulatis utrinquè glandulosi subtus reticulatis secus nervos rufo et brevè velutinis, corymbi capitulis 1-3 sessilibus, involucri squamis sublineari-bus valdè acuminatis glabris, exterioribus confertis brevissimis, achænio dense villoso. 2/ in Indiâ orient. Nepauliâ ad Bechiaeo et Hethaura legit (cl. Wallich. *V. rigida* Wall! cat. et herb. n. 2925. comp. 35. non Swartz. Folia ferè Decanuri asperi sed invol. et fruct. diversus. Pappi series externa interiori quintuplò brevior. (v. s. comm. ab hon. cœt. merc. angl. Ind. orient.)
2. *V. TERRES* (Wall! cat. et herb. n. 2926. comp. 36), canle tereti simplici glandulosi pilis scabrido, foliis rigidis sessilibus obovatis oblongiave brevè acuminatis grossè serratis utrinquè glandulosi subtus reticulatis secus nervos scabridis, capitulis axillaribus sessilibus solitariis ultimis subconfertis, involucri squamis oblongo-linearibus acutis subpungentibus multiserialibus, externis brevissimis, achænio pubescente. 2/ in Indiâ orient. *Acilepis squarrosa* Don prod. 169. *Vern. squarrosa* Less. in linnæâ 1831. p. 627. non p. 678. Pappi series ext. interiori 4° brevior paucisetosa.
 - a. *angustior*, foliis oblongis subovatis. Eupatorium pauciflorum Wall! ad Taogdong et Promæ. (v. s.)

β. *lactor*, foliis latius obovatis. Onopordon teres Ham. ad Sakanaghur.
(v. s. comm. ab hon. coet. merc. angl. Ind. or.)

Sect. II. HOLOLEPIS. — *Hololepis* DC. ann. mus. 16 p. 190. rec. mem.
p. 34. Cass. dict. 21. p. 308. Hayneæ sp. Spreng. — Vernoniæ
sp. Less.

Capitulum multiflorum. Involucrum squamæ siccae coriaceæ adpressæ imbricatæ; exteriores breves ovatæ, interiores seorsim elongatæ. Recept. latum planum alveolatum fimbriiferum. Achænium glaberrimum. Pappus paleis piliformibus rigidis æqualibus barbellulatis numerosis. — Herbae (forsan basi suffruticosæ) Americanæ erectæ crassæ ramosæ. Folia alterna petiolata elliptica penninervia integerrima suprâ glaberrima, subtus sublepidoto-canescens. Rami axillares foliis longiores apice r-cephali erecti.

§. 1. *Pedunculis 1-cephalis nudis aut immediate sub involucri folia floralia 4 bracteantia gerentibus.*

3. V. PEDUNCULATA (DC. ann. mus. 16. p. 190. t. 6. rec. mem. p. 35. t. 2. sub Hololepis), involucrum squamis glaberrimis, corollæ lobis glabris, receptaculi fimbriis hinc inde elongatis et achænium superantibus 2? in Brasiliâ. Serratula pedunculata DC. in Pers. ench. 2. p. 391. Hayneæ pedunculata Spreng. syst. 3. p. 391. H. pedunculata Cass. l. c. Pappus rufescens. (v. s.)

4. V. INVOLUCRATA (Less. in Linnæâ 1829. p. 246. et 1831 p. 627 excl. syn.), involucrum squamis extus margine ciliatis, corollæ lobis apice barbatis, receptaculi fimbriis brevibus 2? in Brasiliâ tropicâ. Pappus flavescens. Invol. squamæ 3-nerviæ dicuntur nec ut in priore enerviæ aut multinerviæ dicendæ.

§. 2. *Capitulis ad apicem plurimis aggregatis inter folia sessilibus, invol. squamis in acumen subspinosum aut molle desinentibus.*

5. V? PROTEOPSIS, tota capitulis exceptis adpressæ sericeo-argentea, caule herbaceo tereti striato usque ad apicem foliato, foliis semiamplexicaulibus elongato-lanceolatis acuminatis integerrimis summis basi multinerviis, capitulis ad apicem aggregatis breviter pedunculatis multifloris, invol. campanulati squamis in acumen longum subspinosum desinentibus. 2? in serrâ de Piedade prov. Min. Ger. legit. cl. Martius. Proteopsis argentea Mart. et Zucc! herb. Achænia glaberrima angulatâ. Pappi setæ 2-seriales lineari-acuminatæ ciliatæ caducæ inter se similes. An Decaneuro an Hololepidi affinis? (v. s. in h. acad. Reg. Monac.)

6. V. ANDRIEUXII, caule frutescente tereti araneoso et piloso-glanduloso, foliis oppositis petiolatis elliptico-lanceolatis utrinque acuminatis argutè et remotè serratis hinc inde sublobatis suprâ breviter scabridis subtus dense cano-tomentosis, capitulis ternis pedunculatis, invol. squamis inermibus multiserialibus elongatis squaroso-subtortis, achænio adpressæ pubescente. 3 in Mexico inter Tehuantepec et flum. Guaracoalcos septemb. flor. legit. cl. Andrieux pl. exs. n. 269! Pappus rufescens, ext. squamellatus brevis. Petioli 1; poll. longi. Folia 4-5 poll. longa, sesquipollicem lata. Species pulchra priori habitu affinis sed fortè caract. nimis aliena. (v. s. comm. cl. Andrieux.)

§. 3. *Ericoideæ, nempe fruticosa, ramis usque ad flores folia conferta gerentibus, apice capitula cylindracea gerentibus.*

7. V. BUXOIDES (Less. in Linnæâ 1829. p. 247. 1831. p. 629), fruticulosa, ramulis apice angulatis tomentosis, foliis sessilibus confertis obovatis integerrimis coriaceis suprâ glabris lucidis, subtus dense tomentosis, ramis floridis ad apicem usque foliosis, capitulis cylindraceis 7-floris, invol. pauciserialis squamis siccis 3-ner-

- vile obtusis, achenio multicostrato. $\frac{5}{8}$ in Brasiliâ tropicâ. Cor. glabrâ purpureâ? Pappus flavescens. Folia 3-4 lin. longa, 2 lin. lata.
8. *V. ERICOIDES* (Less. l. c. 247. excl. syn. et Less. in Linnæâ 1831. p. 629), fruticosa, ramis confertis teretibus tomentosis, foliis sessilibus 1-nerviis coriaceis confertis linearibus integerrimis suprà glabris subtus tomentosis, ramulis ad capitula usque foliosis, capitulis cylindraceis multifloris, invol. squamis oblongis acutis basi trinerviis, achenio multicostrato, pappo biseriali, paleis aequalibus ciliatis. $\frac{5}{8}$ in Brasiliâ tropicâ (Less.), in monte Itambe da Villa do Principe prov. Minarum (Mart!) *Vernonia alpicola* Mart! herb. Cor. purpureâ? glabrâ. Pappus flavescens. (v. s. in h. Acad. reg. Monac.)
9. *V. OLEASTER*, caule fruticoso ramoso tereti, ramis cano-tomentosis, foliis oblongis subsessilibus integerrimis 1-nerviis subobtusis suprà virescentibus dense velutinis subtus cano-tomentosis, capitulis ad apices ramorum circ. 3 sessilibus congestis 20-floris, invol. multiserialis squamis lineari-lanceolatis acuminatis subtomentosis longè ciliato-barbatis. $\frac{5}{8}$ in Serra de Piedade prov. Minas-Geraes Brasiliæ legit. cl. Martins. *Albertinia Oleaster* Mart! herb. Folia pollicem longa, 2-3 lin. lata. Achenium glaberrimum. Pappus biserialis setis caducissimis albidis longitudine aequalibus! Receptaculum punctatum. (v. s. in herb. Reg. Monac.)
10. *V. PSEUDO-MYRTUS* (S. Hil. voy. Diam. 1. p. 94 et 367), caule fruticuloso erecto ramoso tereti, ramis apice angulatis tomentosis, foliis confertis parvis petiolatis lanceolatis 1-nerviis obtusis suprà nigro-punctulatis, subtus albo-tomentosis, capitulo terminali solitario inter folia sessili, invol. cylindrici squamis linearibus obtusis ciliatis, achenio glabro 10-costato, pappi seriebus aequalibus, paleaceis barbatis supernè capillaceis. $\frac{5}{8}$ in Brasiliâ Serra de S. Antonio. Cor. pallide cæruleæ aut. purpureæ. Fol. 4-5 lin. longa, sesquilineam lata, petiolo 1-2 lin. Pappus rufescens.

Sect. III. LEPTOSPERMOIDES.

Capitula solitaria terminalia multiflora. Achenia sericeo-villosa. Pappi series inæquales, externæ paleaceæ brevi. — Species Americanæ et Madagascarienses.

11. *V. ADENOPHYLLA* (Mart! herb.), caule fruticuloso ramoso tereti basi folioso inter folia villosa cæt. glabro, foliis sessilibus obovato-oblongis integerrimis obtusis vix mucronulatis penninerviis nervis utrinque prominulis utrinque glabris subtus crebrè glandulosi basi sparsè piligeris, ramis subaphyllis subtilissimè striatis 1-cephalis, capitulo 20-38-floro, invol. hemisphaerici squamis oblongis obtusis dorso nigro-puberulis pappum æquantibus, cor. glabrâ glanduliferâ, achenio striato sericeo-villosa pappi ser. ext. angustè paleaceæ acutâ brevi. $\frac{5}{8}$ in altis Serro Frio prov. Minar. Gener. Brasiliæ legit. cl. Martius. Species distinctissima vix semipedalis, habitu ferè *Vernoniæ* fruticulosæ. Folia 8-10 lin. longa, 3-4 lata. Pedunc. 2 pollic. Cor. purpureæ. (v. s. in h. Acad. reg. Monac.)
12. *V. LEPTOSPERMOIDES* (Mart! herb.), fruticosa, ramis teretibus suboppositis, junioribus tomentosis, adultis glabris, foliis confertis sessilibus coriaceis integerrimis, junioribus utrinque molliter hirsutis, adultis suprà glabris subtus velutinis, nervo medio prominente, capitulis 9-floris ad apices ramulorum solitariis inter folia sessilibus, invol. squamis paucis oblongis obtusiusculis enerviis ciliatis dorso apice puberis. $\frac{5}{8}$ in campis editis prov. Minarum generalium Brasiliæ ad Tejuco, Serro Frio, legit cl. Martins. Achenia obovata substriata rufa glabriuscula. Pappus biserialis stramineus, exter. brevis, interior quadruplò longior scaber. Affinis ex descr. videtur *V. ericoidi* et *buxoidi*, sed pappi ext. brevi à sectione differt. (v. s. in h. Acad. reg. Monac.)
13. *V. POLYGALÆFOLIA* (Less. in Linnæâ 1831. p. 628. non Licht.), caule fruticoso diffuso, ramis teretibus pube brevissimâ stellato-lepidotâ supernè pubescentibus, foliis brevissimè petiolatis coriaceis oblongo-obovatis obtusis integerrimis sparsè lepidotis, capitulis 1-3 terminalibus sessilibus 9-floris, invol. cylindracei squamis imbricatis acuminatis dorso albo-pilosis longè ciliatis, achenio piloso, pappi ser. ext. brevi capillari. $\frac{5}{8}$ in montibus altis prov. Emirnæ insulæ Ma-

- dagascar. *Vernonia procumbens* Bojer! in litt. 1835. Folia 4-7 lin. longa, $\times \frac{1}{2}$ -4 lin. lata; cor. lobi revoluti; limbus à tubo non distinctus. Stylus bulbosus. Pappus ferrugineus. (v. s. comm. à cl. Bojer.)
14. *V. GLUTINOSA*, fruticosa, ramis teretibus junioribus brevissimè velutinis folisque superuè glutinosis, foliis obovatis ovalibusve basi cuneatis subpetiolatis tenè quintuplinerviis reticulatis rigidulis integerrimis, capitulis 1-3 sessilibus terminalibus, invol. squamis lanceolato-linearibus subviscosis, achænio sericeo-villoso, pappo multiseriali, setis exterioribus paucis paulò brevioribus. $\frac{3}{4}$ in insulâ Madagascar propè urbem Tannanarivou, legit. cl. Bojer. *Vernonia pauciflora* Bojer! in litt. non Less. Frutex 4-5-pedalis. Folia juniora rufescentia. Pappus rutilans invol. paulò longior. Flores in spec. meo inexplicati. (v. s. comm. à cl. Bojer.)
15. *V. ARGYROPHYLLA* (Less. in linnæâ 1831. p. 627), ramis ad apicem usquè foliatis, foliis approximatis coriaceis brevissimè petiolatis obovatis acutis suprà demum nitidis subtus argenteo-tomentosis, capitulis terminalibus solitariis, invol. campanulati squamis siccis muticis acuminatis extus pilosis, achæniis villosis, pappo pluriseriali, ser. ext. breviori latiori. $\frac{3}{4}$ in Brasiliâ. Cor. glabræ purpureo-violaceæ.
16. *V. MONTICONA* (Mart! herb.), fruticosa, ramis teretibus glabris, foliis sessilibus erectis coriaceis integerrimis oblongis margine latè revolutis suprà junioribus sparsè puberulis demum glabris punctulatis subtus adpressè tomentosis, capitulis 5-floris oblongo-cylindricis 2-5 ad apices ramulorum aggregatis in summis axillis aut apice ipso sessilibus, invol. squamis siccis rufis glabris acuminatis. $\frac{3}{4}$ in monte Itambe prov. Minarum Brasiliæ, legit. cl. Martius. Habitus *Lychnophoræ*, sed capitula distincta nec verè glomerata. Achænium pubescens oblongum. Pappi exter. paleæ latæ acuminatæ, inter. duplò triplòve longiores setiformes, omnes albæ. An *Chrysocoma* adpressa Arrab. fl. flum. 8. t. 35? (v. s. in h. Acad. reg. Monac.)

Sect. IV. *VANILLOSMA* Less. in linnæâ 1831. p. 630. — Vern. axillifloræ Less. in linnæâ 1829. p. 251.

Capitula pauciflora aggregata fasciculata, fasciculis axillaribus folio suo brevioribus. — Frutices Americani. Folia petiolata penninervia sæpissimè subtus incana.

§. 1. *Achæniis glabris.*

* *Capitulis sessilibus.*

17. *V. SPLENDENS* (Less. in linnæâ 1829. p. 252. et 1831. p. 630), fruticosa, caule tereti glabro, ramis flexuosis vix apice petiolisque subtomentosis, foliis petiolatis coriaceis oblongis integerrimis suprà lucidis subtus tomento albo adpresso, capitulis spicato-congestis sessilibus 3-floris, invol. squamis coriaceis obtusis 1-nerviis ciliatis extus pubescentibus, achænio glaberrimo multicostato, pappi serie exter. brevi. $\frac{3}{4}$ in Brasiliâ tropicâ, ad sepes circa Rio-Janeiro. Pappus argenteus. Conyza lucida Spreng. neu. entd. 2. p. 142. Conyza splendens Spreng. syst. 3. p. 509. *Carpholobus sessiliflorus* Schott. in Spreng. cur. post. 409. *Vanillosma argyrophyllum* Mart! herb. Variat foliis ovatis oblongis lanceolatisve, paginâ super. venis subprominulis reticulatâ. Pappi ser. ext. multisetosa, inter. triplò brevior. (v. s.)
18. *V. ISOTRICHA*, caule fruticoso tereti glabro, ramis flexuosis junioribus petiolisque velutino-tomentosis, foliis petiolatis ellipticis breviter acuminatis integerrimis suprà glabris lucidis subtus velutino-tomentosis, capitulis 3-floris ad axillas foliorum sessilibus 7-8 glomeratis, invol. squamis obtusis intimis glabrescentibus, achænio glaberrimo pappi setis omnibus æqualibus. $\frac{3}{4}$ in fruticetis circa Rio de Janeiro legit. cl. Lhotsky. *V. splendenti* similis, sed pappus isochætus, achænium dimidio brevius, invol. glabriora paulò longiora, folia superne juniora subreticulata adulta lævia, petioli $\frac{5}{8}$ lin. longi. Folia 2-3 poll. longa, pollicem sesquipollicemve lata. (v. s.)

19. *V. ZEPHYRA* (Less. in *linnæa* 1829. p. 252), fruticosa, ramis angulatis tomento adpresso fulvo petiolisque 4-gonis obiectis, foliis coriaceis oblongis acuminatis integerrimis pellucidoglandulosis glabris subtus leprosis, capitulis paucis spicato-glomerulatis 2-3-floris, invol. squamis subrotundis siccis 1-nerviis, achenio glaberrimo glanduloso multicostato, pappi serie externâ caducâ brevi. 5 in *Brasiliâ tropicâ*. Pappus argenteus.
20. *V. ROTUNDIFOLIA* (Less! in *linnæa* 1829. p. 254. 1831. p. 631), fruticosa, ramis teretibus sulcato-striatis petiolisque cano-tomentosis, foliis petiolatis coriaceis elliptico-rotundis retusis, suprà glabris nervis venisque impressis, subtus densè hirsuto-velutinis nervis prominulis, capitulis 3-4 petiolo brevioribus glomeratis 3-floris, invol. squamis ovatis acutis plurinerviis glabris ciliatis, achenio glaberrimo multicostato, pappi paleis exterioribus paucis et interioribus æqualiter angustis. 5 in *Brasiliâ tropicâ*. Cor. purpureæ glabræ. Pappus argenteus. *Vanillosma firmum* Mart! herb. (v. s.)
21. *V. PYRIFOLIA*, fruticosa glaberrima, ramulis apice angulatis, foliis breviter petiolatis ovalibus oblongisve obtusis rigidis suprà nitidis, subtus reticulato-venosis, capitulis in axillis sessilibus congestis 3-floris, invol. squamis oblongis obtusis, achenio glaberrimo, pappi 3-4-seriali, serie ext. interiore multò brevior. 5 in *Brasiliæ* prov. Rio de Janeiro, leger. cl. Martins et Vauthier (coll. n° 342!) *Vanillosma pyrifolium* et *V. obtusum* Mart! herb. Folia variant. 4 poll. longa, vix pollicem lata, et 2-3 poll. longa, $\frac{1}{2}$ lata, petiolo 3-6 lin. sub lente vix subpuberulo. (v. s.)
22. *V. OXYPHYLLA*, fruticosa, ramis teretibus junioribus petiolisque pube ferrugineâ velutinis, foliis longè petiolatis lanceolatis longè acuminatis integerrimis, suprà glabris, subtus in nervis subhirsutis, utrinquè nervulis elevatis reticulatis, capitulis ad axillas plurimis sessilibus confertis 3-floris, invol. squamis ext. dorso velutinis intimis longioribus glabriusculis subobtusis, achenio glaberrimo pappo longiore, pappi setis ext. brevibus. 5 in sylvis propè Rio-de-Janeiro legit cl. Martins. *Vanillosma acuminatum* Mart! herb. ex parte. Petiolus 9 lin. longus, folia 5 poll. longa, æquipoll. lata. Affinis *V. pyrifoliæ*. (v. s. in h. Acad. Monac.)
23. *V. NOTATA* (Less. in *linnæa* 1829. p. 256), fruticosa subacandens, ramis angulatis tomento adpresso cinereis, foliis petiolatis rigidis lanceolatis integerrimis suprà glaberrimis subtus lepidoto-tomentosis, capitulis fasciculatis petiolo subæqualibus 3-floris, involucri squamis siccis 1-nerviis apice maculâ tomentosa notatis, achenio glaberrimo, pappi serie ext. paulò, int. brevior. 5 in *Brasiliâ*. Cor. glabræ purpureæ. Pappus argenteus. Inflorescentiæ rami 1-3-cephali. (v. s. comm. à Mus. reg. Berol.)
24. *V. AXILLARIS* (Less! in *linnæa* 1829. p. 256), fruticosa, ramis teretibus petiolisque fulvo-tomentosis, foliis oblongo-ellipticis subintegerrimis acutis coriaceis adultis suprà glabris, junioribus suprà et omnibus subtus lanato-tomentosis, capitulis cylindraceis circ. 6-floris ad axillas foliorum glomeratis sessilibus, invol. foliolis coriaceis ovato-oblongis glabris, achenio oblongo subtrigono glaberrimo. 5 in *Brasiliâ tropicâ*. Pappi straminei series externa caduca brevis, multò magis quàm interior lata. (v. s. comm. à Mus. reg. Bav.)
25. *V. MACROPODA*, fruticosa, ramulis petiolisque striato-angulatis densè velutinis, foliis ovalibus utrinquè subattenuatis rariter subdenticulatis suprà adultis glabris, subtus cano-velutinis nervis elevatis reticulatis, capitulis ad axillas 3-5 sessilibus petiolo brevioribus circ. 20-floris, invol. obovatis squamis siccis dorso velutinis imbricatis, ext. latè ovatis, int. elongatis, achenio glaberrimo, pappi serie ext. brevi paulò quam inter. lata. 5 in *Brasiliâ*, sed locus proprius mihi ign. Petioli 12-15 lin. longi. Folia 5-6 poll. longa, 2-3 lata. (v. s.)

* * *Capitulis plurimis pedunculo ramoso insidentibus.*

26. *V. RAMIFLORA* (Less! in *linnæa* 1829. p. 255), fruticosa scandens, ramis teretibus junioribus cinereo-velutinis, foliis petiolatis coriaceis ovalibus subacutis subintegerrimis suprà glabris lucidis, subtus lepidoto-cinereis albidisve, fasciculis

subumbellatis axillaribus, capitulis cylindraceis pedunculatis 2-3-floris invol. squamis ovatis acutis subciliatis adpressis, achænio oblongo glabro, pappi serie externâ paucisetosâ inferiore triplô brevior. ζ in Brasiliâ tropicâ. Cor. purpureæ glabræ. Pappus argenteus. (v. s.)

27. *V. CAPITELLATA*, fruticosa, ramis subangulatis tenuissimè velutinis, foliis petiolatis ellipticis utrinquè attenuatis suprâ glabris, subtus petioloque tenuissimè subvelutinis utrinquè nervulis elevatis reticulatis, pedunculis ad axillas fol. brevissimis ramoso-corymbosis polycephalis subvelutinis, capitulis ovatis 3-floris, invol. squamis ext. brevibus obtusis dorso velutinis, int. oblongis glabresculis, achænio glaberrimo, pappo brevior, pappi setis ext. brevibus. ζ in Brasiliâ (prope Rio-de-Janeiro?) legit cl. Martins. Vanillosma acuminatum Mart! herb. ex parte. Certissimè ab oxyphyllâ diversâ! Proximè ad *V. tereticaulem* accedit. Achænia capituli 3 sæpè conglutinata. Petioli 6 lin. longi. Limbi 3 poll. longi 15 lin. lati. An *Chrysocoma verticillata* Arrab. fl. flum. 8. t. 26? (v. s. in h. Acad. reg. Monac.)

28. *V. TERETICAULIS*, caule fruticoso, ramis jnnioribus teretibus pubescenti-velutinis, foliis petiolatis ellipticis subacutis basi sæpius inæquilateraliter integerrimis suprâ glabris, subtus petiolisque velutino-tomentosis, pedunculis in axillis plurimis umbellatis petiolo brevioribus apice ramosis, capitulis ovatis 4-5-floris subvelutinis, squamis ovatis acutis, achænio gracili glaberrimo, pappi serie ext. paucisetosâ brevi. ζ in Americâ calidiore? (Nee!), ad Guayaquil (Haenk!) Affinis *V. Poeppigianæ*, sed differt achænio longiore non concolato vix striato, pappo albisimo nec subrufescente, foliis paulò minoribus et magis coriaceis, etc. Folia infer. interdum ovata. Petioli 5-6 lin. longi. (v. s. in herb. Haenk.)

29. *V. BLANCHETIANA*, fruticosa, ramis subsæcudentibus teretibus lepidoto-velutinis cinereis, foliis petiolatis ellipticis subacutis integerrimis suprâ glabris, subtus petioloque lepidoto-velutinis, pedunculis in axillis confertis ramosis polycephalis, capitulis ovatis 5-floris, squamis glabris subciliatis ovatis obtusis ext. brevissimis, achænio glabro multicostato, pappi serie ext. multisetosâ brevi. ζ in Brasiliâ circa Bahiâ legit cl. Blanchet. Petioli 9 lin. longi. Folia 5-6 poll. longa, 2 $\frac{1}{2}$ -3 lata, basi sæpè inæqualia. Affinis *V. Poeppigianæ*, sed rami teretes nec apice angulati, folia minùs acuminata, invol. minora. Flores 5, nec 7-8, etc. (v. s. comm. à cl. Blanchet.)

30. *V. POEPPIGIANA*, fruticosa, ramis tenniter velutinis apice angulatis, foliis petiolatis ellipticis acuminatis suprâ glabris, subtus petiolo et in basi nervi medii paginæ super. subadpressè lepidoto-velutinis subrufescentibus, pedunculis ramosis axillaribus petiolo brevioribus fasciculatò-polycephalis, invol. ovatis glabris 7-8-floris, squamis siccis ovatis, intimis subacutis, achænio glaberrimo, pappi setis brevioribus paucis. ζ in Peruvîâ legit cl. Poeppig. Pl. ign. n° 23 et 1425. Petioli 9 lin. longi, Folia 5-6 poll. longa, 2-2 $\frac{1}{2}$ lata, basi sæpè inæqualia. Species *V. ramifloræ* accedens, sed capit. ovata (nec cylindracea) pluriflora. (v. s. comm. à cl. Poeppig.)

31. *V. UMBELLATA* (Less! in linnæâ 1829. p. 255. 1831. p. 631), fruticosa scandens, ramis teretibus glabris vix apice pubescentibus, foliis petiolatis lanceolatis utrinquè acuminatis integerrimis suprâ glabris, subtus densè albo-tomentosis, fasciculis umbellatis axillaribus, capitulis pedicellatis cylindraceis 7-9-floris invol. squamis subscariosis subrotundis glabris internis oblongo-linearibus, achænio glabro, pappi serie externâ brevi caducâ. ζ in Brasiliâ tropicâ. *Conyza ramiflora* Spreng. syst. 3. p. 509. Cor. purpureæ glabræ. Pappus argenteus. (v. s.)

32. *V. LUNDIANA* (Less! in linnæâ 1831. p. 631), caule arboreo ramis dependentibus striatis tomentoso-lepidotis, foliis petiolatis oblongo-ellipticis utrinquè obtusissimis integerrimis suprâ lævibus, subtus densissimè tomentoso-lepidotis, capitulorum fasciculis axillaribus petiolo brevioribus, capitulis plus quàm 7-floris, invol. cylindracei squamis obtusis apicè subpubescentibus, achænio glaberrimo. ζ in Brasiliâ ad Serra de Mar, prope novum Friburgum, colonianu

Helvetiorum. Arbor 20 ped. octobri florens, odorem Philadelphici coronarii spirans. (Lund!) Cor. albidæ. Petioli 12-15 lin. longi. Folia 8-10 poll. longa, 5 et parum ultra lata. (v. s. comm. à cl. Lund.)

§. 2. *Achæniis pilosis.* — *Species à Vanillosmâ subulienæ et inter se fortè vix cognatæ!*

33. V. VAUTHIERANA, caule fruticoso ramosissimo tereti, ramulis angulatis pulverulento-incanis foliis breviter petiolatis lanceolatis acuminatis basi cuneatis integerrimis suprà adultis glabris, subtùs pube adpressissimâ canescentibus, capitulis ad axillas fol. 3-5 breviter pedicellatis confertis 10-floris, invol. squamis oblongis subacutis dorso præsertim ad apicem pubescentibus, cor. lobis apice villosis, achænio turbinato brevi sericeo-villosissimo, pappi serie ext. paleis ovatis acutis inter. multò brevioribus. 5 in Brasiliæ prov. Minarum gener. legit. cl. Vauthier. coll. n° 336. Folia cum petiolo ferè 3 poll. longa, pollicem lata.

♀. *angustifolia*, foliis lineari-lanceolatis. 5 cum priore. Vauth! coll. exs. n° 337. (v. s. comm. à cl. Vauthier.)

34. V. ELÆAGNOIDES (H. B. et Kunth nov. gen. 4. p. 42), fruticosa, ramis teretibus tenuissimè sericeo-pubescentibus, foliis brevissimè petiolatis coriaceis elongato-lanceolatis obtusis integerrimis suprà glaberrimis lucidis, subtùs sericeo-tomentosis, capitulis ad axillas glomerulatis campanulatis 15-20-floris, invol. squamis coriaceis 3-nerviis ovatis acutis extùs pilosis, achænio oblongo piloso, pappi serie ext. brevi latiusculâ. 5 in montosis prope fluvium Cuellar Novo-Granatensium. Less. in linnæâ 1829. p. 254. 1931. p. 631. Cor. glabræ, Pappus stramineus.

35. V. CANDOLKEI (Mart! herb. sub Vanillosma), caule fruticoso tereti, ramisque subangulatis cano-tomentosis foliis breviter petiolatis lanceolatis oblongisve subacutis penninerviis suprà glabris reticulato-venulosis, subtùs petioloque cano-tomentosis, pedunculis axillaribus petiolo longioribus ramoso-corymbosis incanis, capitulis oblongis 7-floris, invol. squamis acutis subpuberulis, cor. lobis tenuissimis glabris, stylo breviter bifido lobis subobtusis, achænio pilosiusculo, pappi setis ext. paucis brevioribus. 5 in campis editis ad Rio de Contas prov. Bahiensis, legit. ill. Martins. Folia majora 3 poll. longa, 12-15 lin. lata, petiolo 3-lin. Species distinctissima fortè propriæ sectionis? (v. s. in h. Acad. reg. Monac.)

36. V? GAUDICHAUDIANA, fruticosa, ramis teretibus adultis glabris junioribus vix subcanescentibus, foliis petiolatis ovato-lanceolatis acuminatis integerrimis membranaceis suprà glabris, subtùs adpressè cano-tomentosis, capitulis axillaribus 8-10 umbellulatis pedicellatis, pedicellis vix petiolo longioribus parcè ramosis, invol. ovatis squamis glabris subobtusis 5 circa Rio de Janeiro legit. cl. Gaudichaud. Flores fructusque ignoti, undè species dubia. Accedit hinc ad V. tereticanlem illinc ad V. Candolkei, sed diversa videtur. Petiol. 4 lin. longus; limb. fol. 3 poll. longus, pollicem latus. (v. s. comm. à cl. inv.)

SECT. V. STROBICALYX, Blum. in litt.

Invol. squamæ imbricatæ ovatæ obtusæ brevissimæ et floribus multò superatæ. Involucrum 3-5-florum teretiusculum. Pappus 1-serialis ant setis ext. paucis calyculatus. — Arborea Indicæ aut Africanæ. Folia petiolata plana penninervia.

37. V. CELEBICA, arborea, foliis longè petiolatis oblongo-ellipticove lanceolatis basi subinæqualiter attenuatis cuspidatis integerrimis glabris cum nervis subtùs petiolo ramulisque pubescenti-velutinis, corymbo terminali divaricato, capitulis 3-4-floris, invol. squamis imbricatis obtusis ciliolatis; achænio teretiusculo substriato pubero invol. circ. longitudine. 5 in sylvis insulæ Celebes. Eupatorium Celebicum Blum! bijdr. 903. Leucomeris celebica Blum! in litt. Affinis V. arboreæ, sed pappus duplex, ext. brevis. Folia 5-poll. longa sesqui-poll. lata, petiolo pollicari. (v. s. comm. à cl. Blum.)

38. *V. JAVANICA*, arborea, foliis longè petiolatis ellipticis acuminatis basi rotundatis integerrimis suprâ glabris, subtus reticulatis ramulisque fusco-tomentosis, corymbo terminali paniculato divaricato, ramis lateralibus elongatis, invol. squamis ovali-oblongis obtusis imbricatis dorso villosis. \S in sylvis altioribus insulæ Javæ ubi dicitur *Hambirung*. Eupatorium Javanicum Blum! bijdr. 903. Lecommeris Javanica Blum! in litt. Arbor 50-60 ped. Folia petiolo sesquipoll., limbo 6 poll. longo, $2\frac{1}{2}$ 3 poll. lato. Paniculæ rami laterales 3 poll. et ultrâ longæ. Pappi series ext. paucisetosa vix ulla (v. s. comm. à cl. Blum.)

β ? *conferta*, foliis, oblongo-lanceolatis acuminatis basi attenuatis et subtus ad marginem subreplicatis suprâ glabris, subtus junioribus cum ramis et pedunculis densè rufo-velutinis, adultis glabriusculis, paniculâ confertâ, ramis nempe ferè à basi ramulosis, invol. squamis extus densè velutinis. \S in Javâ. Specimen habeo à cl. Blume cum var. α sub nomine Eup. Javanici mixtum, sed meo sensu diversum. (v. s.)

39. *BLUMIANA*, arborea, ramis teretibus glabris in paniculâ pulverulento- et rufo-velutinis, foliis petiolatis oblongo-lanceolatis basi breviter apice longè acuminatis integerrimis utrinquè glabris, paniculâ corymbosâ compositâ, invol. squamis ovali-oblongis obtusis dorso subvillosis. \S in montanis Karang Javæ occidentalia. Lecommeris glabra Blum! in litt. Pappus sordidus, serie ext. brevi paucisetâ. Folia petiolo 4-5 lin. longo, limbo 3-4 poll. longo, 10-14 lin. lato. (v. s. comm. à cl. Blume.)

40. *V. ARBOREA* (Ham! in trans. lin. soc. lond. 14. p. 218), arborea, ramis teretibus subtriatis glabriusculis, foliis petiolatis ovato-lanceolatis acuminatis integerrimis paucinerviis glabris exceptis nervo pag. infer. et petiolo subvelutino-furfuraceis, cymæ terminalis ramosissimæ paniculæformis rachi angulatâ ramisque furfuraceis, invol. squamis paucis ovato-oblongis demum patentibus, achænio turbinato glabro, pappi setis 1-serialibus æqualibus. \S in sylvis Nepauliæ inferioris et Camrupæ (Ham. tr.) et ad Goalpura Beng. (Ham. herb.) Conyza arborea Wall! cat. n° 2. Cor. purpureâ 6-7 in capitulo unico, sed vidi achænia 3 in unicum concreta! (v. s. comm. ab hon. cœt. merc. Anglorum Ind. or.) *Mogol* Bengalensium.

41. *V. ELÆAGNIFOLIA*, caule ramoso tereti pubescenti-albido, foliis brevè petiolatis lanceolatis acuminatis integerrimis suprâ glabris, subtus villis adpressis sericeis argenteis, paniculâ ramosissimâ, ramulis ultimis capitula 3-5 gerentibus, involucri pauciflori squamis ovatis paucis marginè membranaceis subciliatis, achænio glanduloso glabro, pappi serie externâ paucisetosâ brevi. \mathcal{L} \S in Indiâ orient. ad Moalmyne. Wallich. Conyza elæagnifolia Wall! cat. n° 3041. comp. 151. Eupatorium elæagnifolium hort. calc. 1797. Pappus argenteus. (v. s. comm. ab hon. cœtu merc. angl. Ind. or.)

42. *V. ELLIPTICA* (DC. in Wight, contrib. bot. ind. p. 1), caule scandente ramoso tereti subpubescente, foliis breviter petiolatis exactè ellipticis integerrimis suprâ glabris, subtus villis adpressis sericeis argenteis, paniculâ ramosissimâ divaricatâ ramis ultimis capitula 3-5 pauciflora gerentibus, invol. ovatis squamis ovatis subacutis glabris. \mathcal{L} \mathcal{L} in Ind. orient. montibus Neelgerry legit. cl. Wight. Vernonia sp. scandens Wight! herb. Ex foliis et inflorescentiâ valdè accedit ad *V. elæagnifoliam* sed car. gener. ex statu specimin. imperfecto non vidi. (v. s. in herb. Wight.)

43. *V. SECUNDIFOLIA* (Bojer! in litt. 1835), fruticulosa glaberrima, ramis teretibus flexibilibus subscaudentibus, foliis petiolatis secundis ellipticis basi cuneatis apice mucronatis serratis coriaceis nitidis, corymbo composito subfastigiato, pedunculis axillaribus folio paulò longioribus apice corymbiferis, capitulis brevè pedicellatis 5-floris, invol. squamis linear-oblongis vix acutis, achæniis glabris, pappi serie ext. brevi paucisetâ. \S ad margines sylvarum prov. Emirrensis ius. Madagascar legit cl. Bojer. Cor. purpureâ. Invol. apices subpurpurascentes. Pappus albus. Folia. $1\frac{1}{2}$ poll. longa, 9 lin. lata. (v. s. comm. à cl. Bojer.)

44. *V. GLANDULOSA*, fruticulosa altè scandens, ramis angulatis, foliis petiolatis

elliptico-oblongis utrinquē acutis integerrimis utrinquē pube minimā sparsā puberulis subtus glandulas parvas sessiles gerentibus, capitalis ad apices ramulorum circ. 3 breviter pedicellatis, invol. squamis ovatis obtusiusculis ciliatis, acheniis angulatis glabris, pappi serie externā paucisetā brevi. 5 in sylvis Bé-fououn dictis prov. Bé-tani-mena Madagascar insula super arbores scandens. Vern. scandens Bojer! in litt. 1835, non Wall. Flores flavi in invol. 7. Folia 4 poll. longa, pollicem lata. Pappus flavescens rigidulus. (v. s. comm. à cl. inv.)

45. *V. DOWNIANA*, fruticosa, ramulis teretibus pube ferrugineā pedunculis petiolis nervoque medio foliorum subtomentosis, foliis petiolatis ellipticis utrinquē acutis penninerviis nervo excepto glabris, paniculā elongatā terminali laxā aphyllā, invol. squamis 3-4-seriatis siccis glabris ext. brevibus ovatis, int. oblongis, achenio glabro striato-angulato, pappi albiissimi serie ext. brevi setiformi. 5 in Sierra Leona legit cl. Don. Eupatorium paniculatum herb. Lindl. Folia superiora (cum petiolis 4. lin.) 4 poll. longa, 15-18 lin. lata. Cor. non vidi (v. s. in h. Lindl.)

Sect. VI. TRIANTHÆA.

Capitula 3-flora rariùs 4-flora oblonga in corymbum dichotomum disposita.

46. *V. TRINIFLORA* (Less. in linnæa 1831 p. 633), caule fruticoso, ramulis angulatis brevissimè tomentosis, foliis petiolatis membranaceis lanceolatis integerrimis suprā nitidis, subtus lepidoto-tomentosis, paniculā terminall dichotomā, capitulis breviter pedicellatis oblongo-cylindraceis 3-floris, invol. squamis arcetè imbricatis glabris ciliatis ovatis acutiusculis, achenio costato glabro, pappi ser. ext. paleaceā brevissimā, int. lingulatā. 5 in ins. Madagascar. Folia 3-3½ poll. longa, pollicem lata, petiolo 3 lin.
47. *V. TRIFLOSCULOSA* (H. B. et Kunth nov. gen. 4. p. 40), fruticosa, ramis teretibus glabris, foliis petiolatis lanceolato-oblongis apice basique acutiusculis integerrimis membranaceis suprā glabriusculis subtus pubescentibus, cymā paniculato-ramosisimā foliatā, capitulis sessilibus fasciculatis 3-floris, invol. cylindracei squamis acutis ciliatis, ext. minimis, cor. glabrā, achenio canescente, pappi serie ext. brevi. 5 in montibus Mexicanis prope Acaguizola et in calidis prov. Vera-Cruz. Gymnanthemum congestum Cass. dict. 20. p. 110. et 26. p. 20. Pappus albidus. Cor. alba ex Kunth, flavescens ex Cass.
48. *V. DICHOCARPEA* (Less. in linnæa 1829. p. 269. 1831. p. 634. non Spreng.), arborea, ramis teretibus fulvo-tomentosis, foliis petiolatis coriaceis ellipticis acutis integerrimis suprā glabris, subtus petiolisque fulvo-tomentosis, cymis terminalibus subcorymbosis confertis, capitulis cylindraceis 3-floris, invol. squamis siccis 1-nerviis extimis ovatis tomentosis, intimis linearibus glabris, achenio multicostato subpilosisculo, pappi serie externā paleaceā brevi, interioris paleis tortis. 5 in Novā Andalusā juxta Caripe. Pollalesta vernonioides H. B. et Kunth nov. gen. 4. p. 36. t. 321. Oliganthes triflora Cass. dict. 36. p. 18. Eupatorium cydoniaefolium Willd. herb. ex Less. Cor. glabræ albæ! glandulosæ. Pappus flavescens.
49. *V. FIMBRILLIFERA* (Less. in linnæa 1829. p. 109), fruticosa, ramis striatis tomentoso-pubescentibus, foliis ovato-lanceolatis acuminatis in petiolum attenuatis subdenticulatis suprā glabris, subtus glandulosis pubescentibusque, cymā paniculeformi terminali ramosissimā aphyllā, capitulis pedicellatis 8-4-floris, involucri pauciserialis squamis ovalis pubescentibus, receptaculi fimbriis setiformibus, achenio hispido, pappi setis inæqualibus. 5 in ins. Bourboniæ. Pappus rufescens. Cor. flavæ dicuntur, undè genus esset dubium, sed cl. Cassini flavos dixit flores ferè omnes decoloratos. Gymnanthemum fimbrelliferum Cass. dict. 20. p. 109. Eupatorium coriaceum Vahl symb. 3. p. 95. non Spreng.
50. *V. PANICULATA*, fruticosa, ramis teretibus cano-tomentosis, foliis breviter petiolatis ovalibus integerrimis supernè puncta resinosa (glandulose?) conferta gerentibus glabris, subtus petiolisque cano-tomentosis, paniculā maximā

myriocephalâ, capitulis ad apices ramulorum congestis oblongis, invol. squamis oblongis mucronatis extus araneosis, intus et intimis glabris, fl. 1-3 p. . . 3 in Mexico ad San-Dionysio in ditione Oaxaca legit cl. Andrieux pl. exs. n. 270! Capitula speciminis mei floribus fructibusque orbata, et ideo de genere imò de sectione dubito, sed ob habitum speciosum præterire nolui. (v. s.)

Sect. VII. TEPHRODES. — Isomeria *Don in Wight. herb.*

Invol. squamæ imbricatæ lanceolato-lineares acuminatæ rarius acutæ subtrinerviæ disco breviores, ext. breves, int. sensim longiores demum reflexo-patentes. Achæniū teres villosū. Pappus ext. brevissimū regularis, int. 1-serialis albus tenuis. — Herbarum gerontogæ pleræque annuæ cineræ (undè nomen). — Nomen Isomeriæ optimum cum discrimen erga Conyzam designabat, non servandum cum Vernoniæ sectio indicanda. — Si quis vellet omnes subsequentes species in 1 aut 2 coalescere, non valdè repugnarem.

51. V. *PHYSALIFOLIA* (DC. in Wight contrib. bot. ind. p. 2), caule herbaceo erecto ramoso striato, foliis petiolatis latè ovatis repando-dentatis obtusis suprà pubescentibus subtus cauleque cinereo-tomentosis, corymbo laxo, capitulis pedicellatis, invol. squamis lanceolato-linearibus dorso villosis vix acutis. ①? in Indiâ orient. Maderaspatanâ. Conyza cinerea C. Wall! cat. n. 3008. comp. 118. Conyza incana Heyn. herb. Affinis cineræ, sed distincta videtur ex invol. squamis non acuminatis. (v. s. in herb. Wight. et h. Cur. merc. Ind.)

β. *glabrescens*, foliis tenuioribus et magis glabratis. — Cum priori. Transit ad sequentem. (v. s.)

52. V. *CINEREA* (Less. in linnæâ 1829. p. 291. 1831. p. 673), caule herbaceo erecto ramoso pilis brevibus confertis subcanescente, foliis petiolatis, infimis subrotundis, cæteris obovato-oblongis subdentatis subtus præsertim cinereo-canescens, corymbo laxo dichotomo, capitulis pedunculatis, invol. squamis lanceolato-linearibus acuminatis subtrinerviis extus pilosisculis. ① in Indiâ orient. continentalis et insularis partibus ferè omnibus, et in Africa æquinoctiali, in ruderatis secus vias etc. Conyza cinerea Linn. 1208. Wall! cat. n. 3903 et 7137. Serratula cinerea Roxb! cat. calc. 60. Isomeria cinerea Wight! herb. Chrysocoma violacea Schum. gain: 158. Chrysocoma purpurea Forst. in h. Ups. ex Less. Conyza mollis Willd. spec. 3. p. 1924 ex Less. Rumph. amb. 6. t. 14. f. 1. Burm. Zeyl. t. 96. f. 1. Pluk. alm. t. 243. f. 3. ic. omnes mal. Herba valdè varians, cuius forma præcipuæ hic subjungendæ:

β. *rotundifolia* (DC. in Wight contrib. bot. ind. p. 2), foliis etiam superioribus subrotundis obtusè repando-dentatis. ① in regione Maderaspatanâ frequens ad littora maris. Cacalia rotundifolia Willd. phyt. 11. p. 6. t. 1. Wight! herb. Conyza cinerea Wall! cat. n. 118. D. (v. s.)

γ. *scabrida*, pube caulis et foliorum non molli et cinereâ, sed magis sparsâ subscabrâ. — in Senegaliâ (Perr.!) Sierrâ Leonâ (Smeathm!) Indiâ or. (Wall.!) Conyza cinerea Wall! cat. n. 118. H. (v. s.)

δ. *glabriuscula* (DC. l. c.), tota ferè glabra, caule striato, ramis elongatis, foliis ferè omnibus oblongis corymbis confertis. — in Indiâ apricis ad Madras et Megapatam legit cl. Wight. (v. s.). An species propria?

ε. *undulata* (DC. l. c.), tota cinereo-pubescenti, caulibus virgatis, foliis lanceolatis dentato-crispis seu undulatis acuminatis. — in Indiâ orient. Conyza cinerascens Wall! cat. n. 3062. comp. 172. Isomeria chinensis Wight. herb. Eandem ut videtur in Senegaliâ legit cl. Perrottet, sed corymbo conferto nec laxo. (v. s.)

ζ. *parviflora* (Reinw. in Blum! bijdr. ned. ind. p. 893), herbacea erecta, ramis fastigiato-paniculatis scabriusculis, foliis caulinis obovatis in petiolum angustatis, floralibus lanceolatis obtusis repandis parum pubescentibus, corymbo terminali dichotomo, squamis invol. apice cuspidatis. ① in Javæ graminosis et ruderatis, circa Buitenzorg (Blum!), et in insulâ Mauritiî (Bojer! Bontou!) Variat fl. violaceis et roseis et foliis infer. ovatis. Valdè affinis V. cineræ, differt tantum glabritie majore et capitulis dimidio brevioribus.

ribus. Pappus ext. brevissimus. Senecio indica foliis ternis Burm. f. zeyl. p. 211. t. 96. f. 1. optime, monente cl. L. Bouton, huc pertinet, sed folia non ternata; lobi laterales in icone design. sunt foliola novella ex axillaria! (v. s.)

53. *V. abbreviata*, radice suffruticulosa, caule à basi ramosissimo subdiffuso, foliis lanceolatis utrinque attenuatis subserratis utrinque ramisque cinereo-pubescentibus, corymbis oligocephalis, capitulis pedicellatis, invol. squamis lanceolato-linearibus acuminatis dorso puberis vix subtrinerviis. ? in Indiâ orient. *Conyza abbreviata* Wall! cat. n. 3048. comp. 158. nec fortè ab eâ diversæ *Conyza cinerea* B. Wall. n. 3008, et imò *Conyza incana* Wall! cat. n. 3046. Species *V. cinereæ* affinis, sed multò brevior, basi suffruticulosa et ut videtur perennis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
54. *V. vialis*, caule herbaceo erecto tereti ramoso minutissimè puberulo, foliis lineari-lanceolatis utrinque acuminatis breviter petiolatis, junioribus ramulisque villosis, adultis minutè pubescentibus margine denticulis remotis acutis instructis, corymbo laxo, capitulis longè pedicellatis, bracteolis subulatis paucis sub capitulo, invol. squamis dorso puberalis acuminatis, achæniis teretibus sparsè hirtellis, pappi serie externâ brevissimâ. ① in insulâ Madagascar secus vias et circa pagum Tamatave legit cl. Bojer. Flores rosei (v. s. comm. à cl. inv.)
55. *V. leptophylla* (DC. in Wight contrib. bot. ind. p. 2), caule herbaceo erecto ramoso tereti tenuiter pubescente, foliis oblongo-linearibus utrinque acuminatis integerrimis tenuissimè pulverulento-scrabris, paniculâ ramosissimâ laxâ divaricatâ nudâ, invol. squamis circ. 2-seriatis linearibus acuminatissimis vix puberulis. ① in Indiâ orient. *Conyza linifolia* Wall! cat. n. 2063. comp. 173. non Linn. *Calacia linifolia* Heyn! herb. Diversa videtur à Vern. *linifolia* sed non malè refert iconem Rumph. 6. t. 14. f. 2. et idè esset Vern. fasciculata Blum. bijdr. p. 893. in adn. non Michx. Achæn. villosiusculum teres (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et in h. Wight.)
56. *V. erigeroides*, glabra, caule diffuso ramoso striato, foliis lineari-lanceolatis obtusis submucronatis integris in petiolum attenuatis margine angustissimè subrevolutis, corymbo laxo subpaniculato, capitulis longè pedicellatis, invol. squamis mucrone acuminatis glabris disco brevioribus, achæniis glaberrimis. ①? in insulâ Timor et in Novâ Hollandiâ leger. collectores circumnav. Baudinianæ. *V. diffusa* Decaisn. novv. ann. mus. Par. 3 (1834) p. 408. non Less. (1829) *Cyanopsis*? *erigeroides* DC. in Wight. contr. bot. ind. p. 7. var. *Timorensis*. Folia 15-18 lin. longa, 2 $\frac{1}{2}$ lin. lata. (v. s.) Pappo ext. brevissimo ambigit inter *Vernoniam* et *Cyanopsidem*.
57. *V. laxiflora* (Less. in linnæâ 1831. p. 646), glabriuscula, caule erecto à basi ramoso, foliis infimis subrosulatis petiolatis obovatis obtusis dentatis, caulinis lanceolatis linearibusve distantibus, corymbo laxo, capitulis pedicellatis, invol. squamis vix pilosiusculis acuminatis subtrinerviis. ①? in Indiâ orient. ins. Timor (Gaudich.) et peninsulâ Malaicâ (Wight.). *Conyza bellidifolia* Wall! cat. n. 3047 comp. 157. Vix ab eâ differt *Conyza elegantula* Wall! cat. n. 3049 comp. 159. Invol. subrubens. Valdè accedit ad *V. cineream*. (v. s. comm. ab hon. cœt. merc. angl. Ind. or.)
58. *V. conyzoides* (DC. in Wight. bot. ind. 2. p. 6), caule herbaceo erecto striato brevissimè pubescente, foliis ovato- aut oblongo-lanceolatis acuminatis in petiolum brevem attenuatis serratis suprâ glabriusculis, subtus pubescenti-villosis, corymbo composito ramoso polyccephalo, invol. squamis lanceolato-linearibus pubescenti-villosis acuminatis disco brevioribus. — in Indiâ orient. prope Negapatam detexit cl. Wight. *Conyza ovata* Wall! cat. n. 3097. comp. 207. non Less. *Cineraria* glaberrima Spreng. in litt. ad Wall. Affinis *V. cinereæ*. Fol. inf. ignota, sup. 2. poll. longa, pollicem lata. (v. s. in herb. Wight et Wall.)
- β. *Leschenaultii*, foliis angustioribus lanceolatis minùs serratis. — in montibus Neelgerry legit cl. Leschenault. Vernac. dicitur *Kado*. (v. s. comm. ab am. Leschenault.)

59. *V. ALBICANS* (DC. in Wight. bot. ind. 2. p. 6), tota tomento adpresso sericeo villosa canescens, foliis ovato-rhombeis basi cuneatis integerrimis, supernè grossè dentatis acuminatis, corymbo subpaniculato nudo, invol. squamis linearibus villosis acuminatis, achænio villosa, pappo ext. quadruplò quàm int. breviorè. — in Indià orient. verisim. agro Maderaspatano. Conyza seu Chrysocoma sp. herb. Madr. Conyza albicans Wall! cat. n. 3072. comp. n. 182. Folia 15 lin. longa, 7-8 lata. Cor. lobi hispidi. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
60. *V. ? RUGOSA* (Blum! bijdr. ned. ind. p. 894), herbacea, ramis divaricatis pubescentibus, foliis oblongo-lanceolatis basi attenuatis apice cuspidatis remotè denticulatis suprà glabrinusculis, infrà rugosis et puberulis, corymbo terminali dichotomo divaricato laxifloro, squamis involucri acutiusculis. — in ins. Javà ad ortum fl. Tiburru in montanis Gede legit cl. Blum. Folia 3-4 poll. longa, 8-12 lin. lata majora subrugosa. Capitula parva. Pappus albus, ext. fortè nullus. (v. s. spec. incompl.)
61. *V. ? CYMOsa* (Blum! bijdr. ned. ind. p. 894), herbacea, ramis paniculatis pubescentibus, foliis lineari-lanceolatis remotè serrulatis scabris subtùs puberis, corymbo terminali dichotomo, pedicellis approximatis subumbellatis, squamis involucri acutiusculis. — in montanis ins. Javæ. Pedunculì compressi. Capitula parva. Folia 4-poll. longa, 6-8 lin. lata. Pappum ext. in specimine equidem manco non vidi et hic ex auct. cl. Blume et affin. cum *V. divergente* refero. (v. s. specim. incompl.)
62. *V. EUPATORIODES* (Blum! bijdr. ned. ind. p. 894), suffruticosa, ramis villosis foliis oblongo-lanceolatis argutè serratis rugosis suprà sparsi-setoso-scabris, subtùs villosa hirsutis, corymbo terminali divaricato, squamis involucri ext. brevibus acuminatis, intimis longioribus vix acutiusculis. — in ins. Javà ad cacumen montis Burangrang. Pappus albus ext. brevissimus vix ullus. (v. s. fruct.)
63. *V. DIVERSIFOLIA* (Bojer! in litt. 1835), caule herbaceo erecto tereti striatulo pubero paniculato-ramoso, foliis infer. elongatis caulinis ellipticis oblongisve integris aut sinuatis suprà scabridis, subtùs puberulo-hirsutulis, paniculà laxà apice corymbosà, invol. squamis ext. acutis, intimis lineari-oblongis apice subobtusis subcoloratis, achænio villosa, pappo exter. brevissimo. ① in fossis circa agros cultos prov. Emirnenis. ins. Madagascar legit cl. Bojer. Folia infer. 15 lin. longa, latitudine varia, caulina 6-9 lin. longa, omnia obscurè viridia. Cor. purpureæ. (v. s. comm. à cl. inv.)

Sect. VIII. LEPIDAPLOA. — Lepidaploæ et Vernoniæ sp. Cass. —
Vernoniæ propriæ pleræquæ Less.

Capitula plusquàm 4-flora, in paniculas dichotomas seu fasciculos seu cymas scorpioideas disposita, rariùs et verisim. abortu subsolitaria. Invol. squamæ apice non appendiculatæ.

§. 1. *Involucri squamis obtusis acutisve nec acuminatis, capitulis nequè solitariis neque in cymas scorpioideas dispositis, Less. in linnæ 1831. p. 635. — Species Americanæ, Asiaticæ, rariùs Africanæ.*

64. *V. ZEYLANICA* (Less. in linnæ 1829. p. 344. 1831. p. 636), suffruticosa, caule tereti tomentoso, foliis petiolatis membranaceis acutis crenatis suprà glabrinusculis, subtùs densè lanato-tomentosis, aliis ovatis basi cuneatis aliis panduratis, basi auriculato-cordatis, cymis subcorymbosis terminalibus, ramis elongatis apice ramosis, invol. ovatis squamis ovatis siccis vix acutis, achænio glabrinusculo, pappi serie ext. angustà paleaceà brevi. § in Zeylonà (Burm. Lesch!) in Madagascar (Bojer). Calalia Burm. Zeyl. 52. t. 21. Eupatorium Zeylanicum Lin. spec. 1172. Pappus pallidè rufescens. Cor. glabræ. Vernonia Wight! 416 sine loco design. (v. s. comm. à cl. Lambert et Leschenault.)

65. *V. APPENDICULATA* (Less. in *linnæa* 1831. p. 636), caule fruticoso ramosissimo, foliis infer. maximis basi cordatis inæqualiter serratis, caulinis lanceolatis petiolatis ad basin petioli auriculatis serrulatis supra glabris, subtus cum petiolis et ramis tomentosis, summis sessilibus sæpe cordato-auriculatis, corymbo fastigiato polyccephalo, capitulis 10-floris oblongis, invol. squamis imbricatis obtusis glaberrimis. 5 in Madagascar circa urbem Tananarivou, legit. cl. Bojer! Vernaculè dicitur *Ampiat* seu *Ampiadi*. Vern. macrophylla Bojer! in litt. 1833. Folia infer. 10 poll. longa margine plicata. Flores lilacini. Achæn. 3. lin. longa, subangulata sparsivillosa. Pappus albus pluriserialis, serie ext. breviori. Variat foliis omnibus aut plerisque exaristis. (v. s. comm. à cl. Bojer.)
66. *V. PANDURATA* (Link. enum. h. ber. 2. p. 276), herbacea, ramis teretibus subvillosis, foliis sessilibus amplexicaulis ovatis obtusis apiculatis basi auriculato-cordatis crenatis supra pilosis, subtus villosis, cymæ pluriès dichotomæ ramis 1-cephalis, capitulis multifloris campanulatis, invol. squamis siccis glabris 3-nerviis glandulosus ex rotundato-oblongis, achænio sparsè piloso multicostrato, pappi serie externâ minimâ. 2½ ad fluvium Congo, ex Less. in *linnæa* 1831. p. 637. Less. in *linnæa* 1829. p. 261. Cor. glabræ violacæ. An *V. pandurata* Jacq. in Spreng. syst. 3. p. 436. eadem?
67. *V. HYPOLEUCA*, caule fruticoso tereti erecto cum petiolis et paginâ fol. superiore aranceo-tomentoso, foliis oppositis petiolatis ovato-lanceolatis triplinerviis subtus cano-tomentosis trilobis, lobis acuminatis lanceolatis, supremis indivisis, paniculæ ramis ramulisque oppositis apice laxè corymbosis, capitulis 10-floris, invol. squamis 3-seriatis ovali-lanceolatis acutis substriatis, pappo biseriali ext. brevi latè paleaceo, int. multisetoso scabro. 5 in Mexico legit cl. Haenke. Cor. nec genitalia vidi. Species distinctissima. Petioli 8-10 lin. longi. Limb. fol. 2. poll. longus, sesquipollicem latus, pappus rufo-croceus, (v. s. in h. Haenke à cl. Sternb. comm.)
68. *V. NUDICAULIS* (Less. in *linnæa* 1831. p. 637), caule herbaceo simplici imâ basi tantum foliato pilis rigidis capitatis divergentibus villosis, ramis 1-cephalis, foliis sessilibus rosulatis obovatis rigidis subintegerrimis supra pilosis, subtus sericeo-lanatis, invol. campanulati squamis 3-seriatis linearibus acutis extus pilos dimorphos gerentibus, cor. glabris, achænio hirsuto-villoso, pappi plumoso serrato, seriebus confluentibus. (1) in Madagascar. Habitus *V. Sellowii* oligocephalæ.
69. *V. NUDIFLORA* (Less! in *linnæa* 1829. p. 258), basi suffruticosa, caulibus è rhizomate plurimis erectis striatis apice subvillosis, foliis sessilibus longè linearibus 1-nerviis angustissimis marginè revolutis supra scabridis, subtus adpressè villosis, corymbi subfastigiati ramis simplicibus bifidis vix folia superantibus, capitulis 15-16-floris ebracteatis pedicellatis, invol. turbinato-campanulati squamis ovatis subacutis glabris sæpius ciliatis, achænio oblongo piloso, pappi pluriserialis setis piliformibus ext. paucis brevibus cæt. inter. subæquantibus. 2½ in Brasiliâ meridionali. Cor. glabræ violacæ. Pappus albus. Folia 3-poll. longa, lineam lata. (v. s. comm. à Mus. reg. Berol.)
70. *V. SQUARROSA* (Less! in *linnæa* 1831. p. 678. non p. 627), canibus erectis teretiusculis basi suffruticulosus subvelutinis, foliis sessilibus elongato-linearibus margine subrevolutis, supra glabris rugosis, subtus albo-tomentosis, cymis scorpioideis summis subverticillatis, invol. campanulati squamis oblongis obtusis ciliatis apice squarrosis intimis subscariosis, achæniis multicostratis hirsutis, pappi serie ext. brevi paleacæ. 2½ in Brasiliâ. Vern. rubricaulis, squarrosa et plantaginoides Less! in *linnæa* 1829. p. 300. Cor. purpureo-violacæ. Differt à rubricauli, invol. squamis squarrosis nec adpressis.
- α, *angustifolia* (Less. ll. cc.), foliis confertis angustè linearibus uninerviis margine revolutis.
- β, *plantaginea* (Less. ll. cc.), foliis rarioribus flexuosis brevioribus penninerviis margine planis. (v. s. comm. à Mus. reg. Berol.)
71. *V. INTERMEDIA*, caule fruticuloso erecto tereti glabro, ramulis floridis pubescentibus foliosis 1-cephalis axillaribus racemoso-corymbosis, foliis sessilibus an-

gustè sublinearibus subacutis integerrimis margine revolutis suprâ glabris, subtus albidotomentosis nervo medio glabro, capitulis 30-35-floris, invol. squamis lanceolatis ciliatis dorso subvillosis mucrone brevi acuminatis, cor. glabrâ, achænio angulato parvè villoso, pappi ser. exter. paleaceâ brevi 2½ in prov. Rio-Grande Brasiliæ. Media inter V. nudicaulem et psilophyllum. Cor. purpureæ. Capitula subbracteata. (v. s. in h. Mus. reg. Par. et Mus. imp. Bras.)

β, *ramosior*, ramis floridis valdè elongatis scorpioideis sæpè bifidis, capitulis lateralibus subsessilibus aut breviter pedicellatis secus ramos 5-6 distantibus sæpiùs extrâ-axillaribus, invol. squamis paulò magis acuminatis. 2½ in prov. Rio-Grande Brasiliæ. Folia 3-6 poll. longa, lineam lata. Videtur eadem ac var. α in solo uberiore lecta. (v. s. in Mus. reg. Par. ex Mus. imp. Bras.)

72. V. *PSILOPHYLLA*, caule suffruticoso tereti erecto glabro tenui apice summo pubescente, foliis linearibus sessilibus acutis integerrimis 1-nerviis margine revolutis suprâ glabris, subtus albidopubescentibus, ramis foliosis apice pedunculosis 3-5 subcorymbosos 1-cephalos subnudos gerentibus, pedicellis axillaribus, capitulis axillaribus 15-20-floris ebracteatis, invol. squamis ovalilanceolatis glabrisculis ext. acuminatis int. mucronatis, corollâ glabrâ, achænio puberulo, pappi ser. ext. paleaceâ brevi acutâ. 2½ in prov. Rio-Grande Brasiliæ. Affinis habitu et inflorescentiâ V. nudifloræ, sed pappo faciliè diversa. Folia 3 poll. longa, semilineam lata. Cor. purpureæ. Pappus griseus. (v. s. in h. reg. Par. ex Mus. imp. Bras.)

β? *Paulina* (Mart! herb.), caulibus à collo plurimis, foliis subtus vix pubescentibus, cymæ extra axillaribus oligocephalis subscorpioideis pedunculis oppositifoliis, invol. campanulatis squamis oblongis dorso vix pubescentibus ext. subacutis, int. obtusis, cor. glabrâ, achænio sericeo-villoso. 2½ in editis campis prov. St-Pauli Brasiliæ legit cl. Martins. Herba pedalis. Pappus griseo-nigricans. (v. s. in h. Acad. reg. Monac.) An species propria? sed specimen mancum.

γ? *leiolepis*, corymbi polycephali pediculis inferioribus parvè ramosis, capitulis 10-15-floris, invol. squamis extimis secus pedicellum minimis acutis cæteris ovalibus obtusis lævigatis glabris purpureiscentibus, pappo albidiorè. 2½ in Rio-Grande. Habitus var. α; invol. squamæ obtusæ ut in β, valdè lævigatæ nec ut in α et β subpubescentes. (v. s. in h. Mus. reg. Par. ex Mus. imp. Bras.)

73. V. *LINEARIFOLIA*, basi suffruticulosa erecta glaberrima, caule subangulato apice ramoso, foliis linearibus utrinquè attenuatis subsessilibus rigidis penninerviis subtus glandulis elevatis punctatis, capitulis solitariis ramulos oppositifolios nudos aut ramos foliosos axillares terminantibus ovatis 25-floris, invol. squamis ext. ovato-oblongis obtusis adpressis, achænio sericeo-villoso, pappi serie externâ vix paleaceâ interiori triplò ferè breviorè. 3½ in Brasiliâ. Herba pedalis. Folia 1½ poll. longa, 2 lin. lata margine revoluta (an per siccationem tantum?). Cor. glabræ verisim. purpureæ. Pappus junior albus dein è griseo sordidè rufus. (v. s.)

74. V? *REVOLUTA* (Ham! tr. linn. soc. 14. p. 218), caule gracili ramoso striato, foliis linearibus subintegerrimis mucronulatis margine revolutis uninnerviis subtus glandulosis utrinquè ramulisque pubescentibus, capitulis terminalibus subsolitariis, invol. squamis linearibus acuminatis, exterioribus dorso carinatis, achænio villoso, pappi pilis æqualibus. 2½ in Indiâ orient. ad Sunnyasika et Monsargunny, ex Wall. cat. 3070. comp. 180 et in Camrupâ borealis et Cosalæ graminosis ex Hamilton. Folia pollicem longa, lineam lata, integerrima aut dente 1-2 medio notata. Pappus tennis argenteus. Cor. purpureæ. (v. s. comm. ab hon. cœtu merc. angl. Ind. or.)

75. V. *LUCIDA* (Less. in linnæâ 1829. p. 260. 1831. p. 640), suffruticulosa glaberrima, ramis virgatis angulatis striatis, foliis sessilibus oblongis basi obtusis apice acutis serratis glandulosis lucidis, cymâ paniculæformi, capitulis longè

pedunculatis subcylindraceis 10-12-floris. invol. squamis ovatis obtusissimis 3-nerviis, achænio piloso pappi seriebus fere longitudine aequalibus setiformibus. ♀ in Brasiliâ meridionali. Cor. violaceæ glabræ. Pappus argenteus. Capitula parva. (v. s. comm. à Mus. reg. Berol.)

76. *V. MITTIDULA* (Less. in Linnæâ 1829. p. 260. 1831. p. 649), caule fruticoso tereti ramosissimo glabro, ramulis angulatis, foliis sessilibus coriaceis oblongis uni-rarius-penninerviis margine subrevolutis integris aut denticulatis utrinque glabris glandulosis, cymis ad apices ramorum subpaniculatis, capitulis pedicellatis 7-8-floris, invol. turbinati foliolis oblongis subacutis adpressis, achænio piloso multicosato, pappi serie ext. brevi paleaceâ. ♀ in Brasiliâ campis ad Rio das Pedras, in prov. S^{ta}. Pauli, Rio-Grande, etc. Baccharis Montevidensis Spreng. syst. 3. p. 460. Pappus rufescens. Cor. violacea glabra. Involucra pilosiuscula seu glabra. Caulis albescens. (v. s. comm. à Mus. reg. Ber.)

77. *V?* *ANGULIFOLIA*, caule fruticoso? tereti striatulo foliisque pubè minimâ subpulverulento-velutinis, foliis petiolatis basi cuneatis cæterum ovatis acuminatis grosse inciso-dentatis basi 3-nerviis, nervis lateralibus bifidis, paniculâ terminali compositâ foliis intermixtâ, invol. squamis ovatis subacutis margine submembranaceis, capitulis 6-7-floris. ♀? in Africâ austr. orient. prope Omsamwubo legit cl. Drege, n. 5033! Flores in specim. verisim. masculi, ovario abortivo, stylo simplici à medio ad apicem hispidio. An Webbii spec.? sed habitus repugnat. Pappi setæ crassæ, exter. breviores. (v. s.)

78. *V. MESPILIFOLIA* (Less. in Linnæâ 1831. p. 641), fruticosa, ramis glabrisculis teretibus, foliis petiolatis longè obovato-cuneatis apice grosse 3-7-dentatis minutissimè (sub lente) puberulis, corymbo subnudo polycephalo, pedunculis pubescentibus, invol. squamis 7-8-subtriseriis obtusissimis dorso glabris margine ciliatis, floribus in cap. 4-5, pappi serie ext. brevi vix paleaceâ, int. multisetâ subpluriseriali. ♀ ad Cap. Bonæ Spei, in Uitenhagen (Burchell! n. 4500. Eckl! h. 1599. 1630. 1764. 1933), Cafferland, Swellendam, territorio cesso (Eckl! n. 1137. 199. 1375. 794), ad Omsamculo et Omtata. (Dreg! n. 5035.) Eupatorium Capense Spreng. fil. suppl. p. 22. Zeyh! fl. exs. n. 201. ex h. Moric. Achænio oblongum villosum. (v. s.)

♀. *subcanescens*, caule foliisque utrinque pubè brevi molli subcanescentibus. ♀ in Cafferland Dreg! n. 3687. (v. s.)

79. *V. CORYMBOSA* (Less. in Linnæâ p. 647), fruticosa, ramis striatis pubescentibus, foliis vix petiolulatis oblongo-cuneatis apice paucidentatis subtus tomentosis, suprâ glaberrimis, capitulis corymbosis, invol. squamis 12-15 obtusissimis dorso tomentosis, floribus in capit. 5, pappi serie ext. paucisetâ, inter. pluriseriali. ♀ ad Cap. Bonæ Spei. in coloniâ distr. oriental. (Burchell! n. 4343.) in collibus distr. Uitenhagen (Eckl! n. 144. 759. 400. 16. 1108. 1917. 1575.) ad Van Staedes rivier. (Dreg! n. 2118.) Stæhelia corymbosa Thunb. fl. cap. 628. ex Vahl! in h. Burm. Eupatorium Burm! herb. Capitula ad apices ramulorum 3-4 subsessilia subumbellata. Folia pollicem longa, 4-lin. lata. Pedunculi compressi folio florali fere breviores. Achæmium oblongum villosum. Pappus rufescens, involucri duplò longior. (v. s.)

♀. *mespiloides*, foliis majoribus acutè 7-9-dentatis subtus adpressè sericeo-canis. ♀ in Uitenhagen legit cl. Ecklon. n. 269. et ad Omtanda cl. Drege, n. 5036! Folia formâ et magnitudine *V. mespilifoliam* referentia; sed subtus canescentia. Capitula *V. corymbosæ*. An fortè utriusque hybrida? (v. s.)

80. *V. ASTRIFLORA* (Mart! herb.), fruticosa, ramis teretibus subtiliter striatis tomentoso-incanis, foliis sessilibus elliptico-oblongis utrinque vix acutis integerrimis suprâ adultis glabris, subtus tomentoso-canis penninerviis, capitulis 5-7 longè pedunculatis corymbosis ebracteatis circ. 40-floris, invol. squamis floribus multò brevioribus, ext. oblongis dorso hirsutis, intimis linearibus acuminatis glaberrimis, cor. glabrâ, achænio parçè villosò, pappi ser. ext. paleaceâ brevissimâ. ♀ in editis campis prov. Minarum gener. Brasiliæ. Folia 3 poll. longa, pollicem lata, pedunculi 1 $\frac{1}{2}$ -2 poll. longi, nudi. Cor. purpurea 5 lin longæ, pappus sordidus. (v. s. in h. Acad. reg. Monac.)

81. *V. ARIÆFOLIA* (Mart! herb.), caule suffruticoso simplici-erecto tereti adpressè sericeo-tomentoso, foliis sessilibus subrotundis obtusis subintegerrimis penninerviis rigidis suprà vix araneosis, subtùs adpressè sericeo-tomentosis, pedunculis 4-5 corymbosis folio paulò longioribus 1-cephalis sub capitulo 1-phyllis, capitulis circ. 70-floris, invol. campanulati squamis multiserialibus coriaceis ext. ovatis subacutis dorso araneosis intimis fere linearibus acuminatis glabris, cor. glabrà, achænio sericeo-villoso, pappi serie ext. angustè paleaceà acuminatà, inter. triplò breviorè. ζ in editis campis prov. Minarum gener. Brasilie. Caulis sesquipedalis. Folia 2 poll. longa, sesquipoll. lata. Cor. purpureæ. Pappus stramineus. Affinis videtur *V. argyrophyllæ*. (v. s. in h. Acad. reg. Monac.)
82. *V. PENTACANTHA*, caule fruticoso tereti supernè albo-lanuginoso, foliis petiolatis oblongis basi longè cuneatis apice obtusis integerrimis rigidis utrinquè reticulatis penninerviis suprà glabrinusculis, subtùs araneoso-tomentosis, capitulis ad apices ramorum subternis sessilibus aut brevè pedicellatis ebracteatis ovato-subglobosis 50-floris, inv. squamis coriaceis multiserialibus imbricatis, ext. ovatis mucrone brevi squarrosò, intimis longioribus acuminatis subecariosis, cor. glabrà, achænio villosò, pappi ser. ext. paleaceà acuminatà int. triplò breviorè. ζ Patr. ign. Valdè accedit ad hexacantham. Folia 3 poll. longa pollicem lata, petiolo 4-5 lin. Pappus subrufescens. (v. s. in herb. Acad. reg. Monac.)
83. *V. LAURIFOLIA*, caule fruticoso ramoso tereti ferrugineo-tomentoso, foliis brevissimè petiolulatis ovali-lanceolatis basi subobtusis apice acuminatis integerrimis rigidis penninerviis utrinquè nervis elevatis reticulatis suprà glaberrimis, subtùs secus nervum puberis, capitulis longè pedicellaribus axillaribus solitariis aut ad apicem laxè cymosis multifloris, invol. ovatis squamis rigidis glabrinusculis lanceolatis acuminatis, cor...., achænio sericeo-villoso, pappi ser. ext. brevè paleaceà. ζ Patr. ign. Folia 2-3 poll. longa, pollicem lata. Pedicelli circ. pollicares. Pappus rufo-griseus rigidulus. Recept. alveolatum. Flores in invol. circ. 25 sed in specim. vix ritè observandi. (v. s. in h. Acad. reg. Monac.)
84. *V. TOLUCCANA*, fruticosa glabra, ramis teretibus folisque oppositis petiolatis elliptico-oblongis integerrimis nitidis viscosis, pedunculis axillaribus oppositis corymbum subpanicalatum confertum gerentibus, capitulis pedicellatis circ. 20-floris, invol. squamis subacutis biserialibus disco dimidio brevioribus, achæniis angulatis pilosiusculis, pappi serie ext. brevè paucisetà. ζ in Mexico circa Toluccam legit cl. Andrieux, pl. exs. n. 273! Habitus Baccharidis seu Psiadiæ sed stigmata elongata hispida Vernoniæ. Fol. 1-2 poll. longa, 6 lin. lata. Cor. in sicco pallidæ. Achænia nigrescentia. Pappus albus. (v. s.)
85. *V. DOMINGENSIS*, fruticosa glabrinuscula, ramulis teretibus apice angulatis, foliis breviter petiolatis ovalibus obovatisve obtusis hinc indè dentatis penninerviis coriaceis suprà lucidis subtùs punctulatis, capitulis paucis subcorymbosis pedicellatis 10-floris, bracteis squamæformibus multiserialibus brevibus obtusis imbricatis sub invol., involucri squamis veris longioribus aicels vix acutis dorso ad apicem subpuberis, achænio 10-costato glabro, pappi ser. ext. paleaceà brevè. ζ in Sancto Domingo. *Lepidaploa buxifolia* Cass. dict. 26 (1823). p. 18 et idè Vern. *buxifolia* Less. in linnæa 1829. p. 313. *Pronotia Domingensis* Spreng! in herb. Balb. 1821. syst. 3. (1826) p. 502. Folia summa 6-7 lin. longa 4 lin. lata. Recept. nudum. Nomen Cass. etiamsi vetustius non admisi ne confundatur cum *V. buxoidi*. (v. s. ex itin. Berter.)
86. *V. ZANZIBARENSIS* (Less. in linnæa 1831. p. 638), caule fruticoso, ramis villosis, foliis ovalibus acutis subdentatis suprà sparse villosis subtùs petiolisque densè cano-tomentosis, corymbo laxo dichotomo subpanicalato, capitulis 12-floris, invol. squamis lanceolatis acuminatis dorso puberis, corollæ lobis intus ad apicem longè barbatis, achænio glabro, pappo biseriali ext. paleaceo brevè. ζ in insulâ Zanzibar locis à mare parùm dissitis legit cl. Bojer. Pappus intensè fulvus barbellatus rufescens. (v. s. comm. à cl. Bojer.)
87. *V. SOLIDAGINIFOLIA* (Bojer! in litt. 1835), fruticosa, ramis erectis apice angulatis pubero-scabridis, foliis subsessilibus lanceolatis acutis serratis suprà

glabris, subtus glancis secus nervos pubescentibus, cymâ corymbosâ, pedicellis valdè bracteolatis, invol. squamis exter. brevibus ferè subulatis, inter. lato-linearibus acutis, achæniis teretiusculis striatis confertè puberulis, pappi serie ext. setosâ brevi. $\frac{3}{4}$ in insulâ Zanzibar legit cl. Bojer. Frutex 8-10-pedalis. Folia omnino Solidaginis Canadensis. Flores flavi (Bojer). Folia super. $1\frac{1}{2}$ poll. longa, 4-5 lin. lata. Fl. in invol. circ. 12. Pappus pallidè cinnamomeus. (v. s. comm. à cl. Bojer.)

88. *V. SUAVEOLENS* (H. B. et Kunth nov. gen. 4. p. 30), fruticosa volubilis, ramis teretibus obscurè tomentosis, foliis breviter petiolatis oblongis penninerviis coriaceis subcrenulatis suprâ subpilosais demùm glabris, subtus petiolisque obscurè tomentosis, ramis floridis subverticillatis, capitulis 7-floris, invol. squamis 1-nerviis ex ovato-acutis ciliatis, achænio glaberrimo, pappi serie externâ brevi caducâ. $\frac{5}{8}$ in Novæ Granat. prov. Ayavaca et flum. Cutaco. Cor. albæ glabræ. Pappus ex albo flavescens. Less. in linnæâ 1829. p. 265. 1831. p. 647.
89. *V. PECTINATA*, fruticosa, ramis teretibus glabriusculis, foliis brevè petiolatis ovatis pectinato-acutè et elongato-serratis membranaceis utrinquè pilosiusculis, cymâ terminali corymbosâ nudâ, capitulis longè pedunculatis multifloris ovato-cylindraceis, invol. squamis siccis glabriusculis ex ovato-oblongis obtusissimis 3-nerviis, achænio glabro, pappi serie externâ brevissimâ. $\frac{3}{4}$ in Indiæ or. montibus Nilgherry legit cl. Noton. Eupatorium pectinatum Wall! cat. n. 3172. comp. 282. Species distinctissima nulli verè affinis. Cor. glabra. Pappus pallidè rufescens. Invol. squamæ apice obscuræ. Stylus et pappus Vernoniæ. (v. s. in herb. bon. cœtu merc. angl. Ind. or.)

90. *V. PECTINIFORMIS* (DC. in Wight, contrib. bot. ind. p. 2), fruticosa? ramis teretibus glabriusculis, junioribus angulatis pubescentibus, foliis breviter petiolatis lanceolatis acuminatis pectinato-acutè et elongatè serratis membranaceis suprâ glabris, subtus pilosiusculis, cymâ terminali corymbosâ nudâ, capitulis longè pedicellatis multifloris ovatis, invol. squamis siccis glabris ciliatis ovato-oblongis subacutis. $\frac{3}{4}$ $\frac{1}{2}$ in Indiæ orient. montibus Dendignul dictis legit cl. Wight. Valdè affinis *V. pectinatæ* et forsân mera varietas. Præter car. indicatos differt pappo nec rufo, involucrio evidenter longiore nec subæquali. Folia 3 pollices longa, vix pollicem lata. (v. s. in h. Wight.)

91. *V. ASPERA* (Ham. trans. lin. soc. 14. p. 219. non Less.), caule herbaceo? ramoso tereti subvillosa, foliis elliptico-oblongis subpetiolulatis basi attenuatis apice acuminatis serratis suprâ glanduloso-scabris, subtus reticulatis secus nervos rufo-villosis et inter eos glanduloso-impressis, paniculâ subcorymbosâ ramis inferioribus folio suo brevioribus, capitulis 30-floris, invol. squamis int. oblongis mucronatis, ext. minoribus valdè acuminatis, achænio oblongo-cylindrico angulato piloso. $\frac{1}{2}$ in Indiâ orient. ad Kodoya in Anggæ montibus, Bengalo et fortè agro Burmanico legit cl. Wall. et Ham. Eupatorium asperum Roxb. calc. 61. Vern. aspera Ham. trans. lin. soc. 14. p. 219. Wall! cat. et herb. n. 2922. comp. 32. non Less. Vernonia Roxburgi Less. in linnæâ 1831. p. 674. Conyza platylepis Spreng. in litt. ad Wall. Eupatorium pyramidale Don. prod. fl. nep. 170? — Pluk. Alm. t. 395. f. 7. malè. (v. s.)

92. *V. MULTIFLORA* (Less. in linnæâ 1831. p. 642), caule herbaceo erecto ramoso tereti tenuissimè puberulo, foliis breviter petiolatis lanceolatis utrinquè acuminatis à medio serratis suprâ asperulis, subtus tenuiter pubescentibus, paniculâ subcorymbosâ polycephalâ nudiusculâ, invol. squamis lanceolatis acutis pallidis glabriusculis, achænio glabro pappo subuniseriali. $\frac{1}{2}$ in Indiâ orient. ad Prome et Panlang. legit cl. Wallich. Conyza multiflora Wall! cat. n. 3032. comp. 142. An Eupatorium acuminatum Don. prod. fl. nep. 170? et idè Eupat. Nepalense Spreng. 3. p. 412? Conyza divergens E. Wall! n. 3007. et Eupatorium versicolor Wall! cat. n. 3167. Folia 3 poll. longa vix poll. lata. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

℞. *latifolia*, foliis ovali-lanceolatis suprâ magis asperis. $\frac{1}{2}$ in collibus Raymahal legit cl. Hamilton. *V. aspera* Ham! herb. ex Wall! cat. h. calc. Var. ℞. affinis *V. asperæ*, sed capitula dimidio minora ut in var. *f.* (v. s. comm. cum priorè.)

93. *V. NILGERRYENSIS*, herbacea? caule tereti subpuberulo, foliis brevè petiolatis ovatis acuminatis spinuloso- et acutè serratis suprâ scabridis, subtus folvo-subtomentosis, cymæ terminalis paniculæformis ramis ramosissimis polycephalis, capitulis ovatis confertis 2-5-floris, invol. squamis siccis ovali-oblongis acutis glabris apice pilosiusculis, achenio glabro, pappi serie externâ brevissimâ deciduâ. — in Indiâ orientali legit cl. Wight, et in montibus Nilgerry cl. Lesschenault. Eupatorium polyanthum Wall! cat. 3171. comp. 281. *V. aspera* Less. in Linnaea 1831. p. 643? Cor. glabra. Pappus albus. (v. s.)
94. *V. ACUMINATA*, caule fruticoso ramoso tereti, ramis junioribus villosis velutinis, foliis breviter petiolatis elliptico-obovatis longè cuneatis apice cuspidato-acuminatis subintegerrimis glabris nervis præcipuis subtus velutinis, paniculæ elongatæ nudæ ramis polycephalis, invol. squamis ovalibus obtusis dorso velutinis imbricatis. § in Indiâ orient. prov. Silhet, in montanis ad Panduah legit cl. Wallich. Conyza acuminata Wall! cat. n. 3034. comp. 144. Eupatorium acuminatum Wall! herb. Eupatorium cuspidatum Spreng. in litt. ad Wall. Folia 4-6 poll. longa 1 $\frac{1}{2}$ -2 lata, petiolo 2-5 lin. Fl. inv. 7-9 ex sicco purpurascens. Pappus albus pauciseriali, setis ext. irregularibus brevibus faciliè deciduis. Achen. glabrum basi attenuatum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
95. *V. PUNDUANA*, caule herbaceo? erecto tereti vix sub lente puberulo, foliis breviter petiolatis lanceolatis longissimis basi attenuatis integerrimis apice acuminatis grossè et rarer serratis glabris, nervo medio subtus tenuissimè puberulo, paniculæ elongatæ nudæ ramis apice confertè polycephalis, invol. squamis oblongis obtusis, intimis longioribus demum stellato-patulis. ¶ in Indiâ orient. prov. Silhet ad Panduah legit Fr. da Silva. Conyza Pundua Wall! cat. n. 3035. comp. 145. Folia 8 poll. longa, 2. poll. lata. Fl. in invol. 8-9. Recept. angustum. Achenia glabra basi attenuata. Pappus albus pauciserialis setis ext. brevibus irregularibus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
96. *V. VAGANS*, caule herbaceo tereti ramoso sarmentoso-scandente brevissimè puberulo, foliis breviter petiolatis ovato-lanceolatis acuminatis integerrimis glabris, ramis floridis axillaribus folio longioribus foliolosis polycephalis in paniculam amplam dispositis, invol. squamis lanceolatis acutis ciliolatis imbricatis. ¶ in Indiâ orient. prov. Silhet legit cl. Wallich. Conyza vagans Wall! cat. n. 3040. comp. 150. Eupatorium vagans Wall! herb. Flores in invol. 7-8 purpurei glabri. Pappus rufus, setis ext. paucis linearibus paleaceis ciliatis, cæteris pluriserialibus scabris. Achen. glabrum teretiusculum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
97. *V. SCANDENS*, caule scandente tereti, ramis junioribus velutinis adultis glabris, foliis breviter petiolatis ovalibus acuminatis integerrimis utrinquè glaberrimis, ramis floridis axillaribus foliosis subpaniculatis, capitulis pedicellatis ovatis, invol. squamis ciliatis dorso glabris ext. ovatis acutis, int. oblongis. § ¶ in Indiâ orient. prov. Silhet legit cl. Wallich. Conyza scandens Wall! cat. et herb. n. 3060. comp. 170. Fl. atropurpurei in invol. 9-11. Achenium junius teretiusculum glabrum. Pappus subrufescens multisetus æqualis, setulis ext. vix ullis deciduis. An fortè à genere removenda? (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
98. *V. BLANDA*, caule terecto tereti paniculato ramoso apice pubescente, foliis breviter petiolatis ellipticis basi attenuatis apice cuspidatis integerrimis glabris, ramis axillaribus foliosis paniculatis, capitulis pedicellatis, invol. squamis imbricatis dorso puberulis oblongo-linearibus mucronatis ext. brevissimis, int. sensim longioribus. § ¶ in Indiâ Birmanica propè Troglâ ad Saluen legit cl. Wallich. Conyza blanda Wall! cat. et herb. n. 3033. comp. 143. birm. n. 1926. Affinis *V. scandenti*. Achenium teres basi subattenuatum puberum. Pappus ext. vix ullus, inter. rufescens. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
99. *V. VOLKAMERTÆFOLIA*, fruticosa, caule tereti glabrisculo ad axillas villosis, foliis brevè petiolatis membranaceis amplissimis elliptico-obovatis acutis basi cuneatis subserratis suprâ glabris, subtus pubescentibus, utrinquè minutè glandulosis, cymæ terminalis paniculæformis rachi angulatâ, ramis tomentosis polycephalis ramosis, invol. squamis siccis extimis ovatis acutis villosis, intimis

- oblongis obtusis, achænio glanduloso, pappi serie externâ brevî. $\frac{3}{4}$ in Nepaliâ detexir indef. Wallich. *Conyza volkameriaefolia* Wall. ! cat. n. 3001. comp. 111. *Eupatorium volkameriaefolium* Wall. ! herb. Folia 9-10 poll. longa, 3 poll. lata. Cor. glabra. Pappus albus. (v. s. comm. ab hon. cœtu merc. angl. Ind. or.)
100. *V. EXTENSA*, suffruticosa glabra, ramis teretibus, junioribus subangulatis subpubescentibus, foliis brevè petiolatis oblongis basi attenuatis apice acumî-natis hinc indè serratis membranaceis, cymis terminalibus paniculæformibus ramosissimis aphyllis, capitalis pedicellatis 8-10-floris, involucri squamis oblongis obtusis, achænio pubescenti-glanduloso, pappi serie ext. interiori triplò breviorè. $\frac{3}{4}$ in Nepaliâ. *Conyza extensa* Wall. ! cat. n. 3016. comp. 126. *Eupatorium rufum* Spreng. in litt. ad Wall. Pappus pallidè aut intensè rufus. Folia 6-7 poll. longa, 1-2 poll. lata. Habitus ferè *V. Senegalensis* sed pappi series valdè inæquales. (v. s. comm. ab ampl. cœtu merc. angl. Ind. or.)
101. *V. MADAGASCARIENSIS* (Less. in linnæâ 1831. p. 644), caule fruticoso scandente, ramis striato-angulatis supernè adpressè pilosis, foliis petiolatis membranaceis oblongo-ovatis acuminatis basi acutis punctatis albo pilosiusculis, paniculis terminalibus dichotomis oligocephalis, capitulis pedicellatis multifloris, invol. campanulati squamis coriaceis glabris ciliatis ovatis, cor. et achænio glabris, pappi serie ext. setacèâ brevî. $\frac{3}{4}$ in insulâ Madagascar prope Maron-Voaî. Cor. in sicco flavidè? Recept. nudum. Pappus flavescens. Folia 2-3 poll. longa, 12-15 lin. lata.
102. *V. ATTENUATA*, caule herbaceo erecto striato infrâ hirsuto apice puberulo, foliis infrâ originem ramorum floralium confertis ellipticis basi cuneatis apice acuminatis grossè dentatis reticulato-nervosis suprâ glaberrimis, subtus in nervis puberulis, super. secus ramos florales distantibus linearibus acutis, dentatis, dentibus omnium mucronatis, corymbo laxo oligocephalo, capitulis longè pedunculatis. — in Indiâ orient. ad Moalmyne legit cl. Wallich. *Conyza attenuata* Wall. ! cat. n. 3020. comp. n. 130. Invol. squamæ glabriusculæ, ext. breves cuspidatæ, int. longiores acutæ. Achæn. teres villosum. Pappus ext. circ. lin. longus paucisetus. Folia 4 poll. longa, sesquipoll. lata. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
103. *V. LONGICAULIS*, herbacea, caule simplici tereti striato hirtello-pubescente, foliis brevissimè petiolatis oblongis ovalibusve basi cuneatis apice acuminatis serratis membranaceis glabris reticulatis venosis in medio areolarum suprâ elevato-glandulosis, cymæ terminalis paniculæformis ramis multi-capitatis subfoliiferis, invol. ovatis squamis siccis ovatis pungenti-mucronatis, achænio glabro, pappi serie externâ brevissimâ. $\frac{2}{3}$ in Indiâ orient. ad Pondua, Silhet, Nepaliâ legit cl. Wallich. *Eupatorium longicaule* Wall. cat. et herb. n. 2169. comp. 279, non 3073. *Eupatorium acuminatum* Spreng. in litt. ad Wall. 1830. Variat foliis latioribus et angustioribus. Habitu accedit ad *V. serratuloidem*. Cor. glabra. Pappus argenteus. (v. s. comm. ab hon. cœtu merc. angl. Ind. or.)
104. *V. SALIGNA*, caule herbaceo erecto tereti striato minutè puberulo paniculato-ramoso, foliis oblongo-lanceolatis utrinquè acuminatis hinc indè serratis, suprâ asperiusculis, subtus subpubescentibus, paniculæ polycephalæ ramis parcè foliosis elongatis 7-9-cephalis, invol. squamis submembranaceis oblongis acutis. — in Indiâ orient. prov. Silhet. legit. cl. Wallich. *Conyza saligna* Wall. cat. n. 3061, comp. 171. Folia 3-4 poll. longa, 10-12 lin. lata. Flores ex sicco vid. albi. Pappus albus, ext. circ. lineam longus. (v. s. comm. ab ill. Cur. merc. angl. Ind. or.)
105. *V. SERRATULOIDES* (H. B. et Kunth. nov. gen. 4. p. 26, t. 316), herbacea, caule simplici glabro tereti striato, foliis brevissimè petiolatis oblongo-lanceolatis acuminatis hinc indè serratis glabris, suprâ impresso-glandulosis, cymæ terminalis paniculæformis ramis apice 5-6-cephalis, capitulis 10-floris cylindraceis, invol. squamis ovatis pungenti-mucronatis adpressis ciliatis glabris, achænio velutino pubescente, pappi serie externâ vix paleacèâ brevissimâ. $\frac{2}{3}$ in Mexico temperato circa Valladolid de Mechoacan (H. et B.) et Leon (Mendez !) Less. in linnæâ 1829. p. 268. 1831. p. 645. *Serratula semiserrata* Willd. herb. ex

Less. Herba 2-3 ped. habitu serratulæ tinctoriæ. Caulis et ramî purpurascens. Cor. purpureæ glabræ. Pappus argenteus. (v. s. comm. à cl. Mendes et Alaman.)

106. *V. LIATROIDES*, caule herbaceo erecto tereti striato tenuissimè puberulo apice sulcato-angulato, foliis brevissimè petiolatis elliptico-oblongis basi acuminatis apice subacutis serratis penninerviis suprà glabris, subtus puberulis, ramis foliosis apice corymbiferis in paniculam dispositis, capitulis pedicellatis ebracteatis 10-12-floris, invol. squamis oblongis glabrisculis, ext. acuminatis, int. vix mucronatis, cor. glabrâ, recept. alveolato, achænio glanduloso subpubescente, pappi serie ext. brevissimâ angustè paleaceâ. $\frac{1}{2}$ in Mexico inter Tula et Tampico (Berl.!) Valdè affinis *V. serratuloides* sed capitula dimidio minora. Folia 2 poll. longa, 7-8 lin. lata. Pappus albidus. Invol. pallida 2. lin. longa. Variat nerviis subtus paulò prominulis aut reticulatis elevatis. (v. s.)

107. *V. MUCRONULATA* (Less. in linnæâ 1829. p. 266. 1831. p. 647), fruticosa, ramis teretibus striatis ferrugineo-tomentosis demùm glabris, foliis brevissimè petiolatis coriaceis ellipticis subcordatis retusis mucronatis integerrimis suprà glabris impresso-glandulosis, subtus fulvo-tomentosis, cymis terminalibus, capitulis brevè pedicellatis, invol. squamis coriaceis 1-nerviis ovatis ciliatis, cor. glabræ laciniis glandulosis, achænio piloso multicostato, pappi serie externâ paleaceâ brevi. $\frac{1}{2}$ in Brasiliâ tropicâ. Folia 11-15 lin. longa, 6 lin. lata. Pappus argenteus.

108. *V. OLIGACTOIDES* (Less. in linnæâ 1831. p. 648), caule fruticoso striato ramisque fulvo-tomentosis, foliis subsessilibus coriaceis oblongis basi obtusis (nec cordatis) penninerviis suprà lucidis scabriusculis, subtus fulvo-tomentosis, margine reflexis subserratis, paniculæ terminalis oblongæ ramis foliis parvis stipatis, capitulis 12-floris basi subsquamosis, invol. squamis siccis dorso fulvo-tomentosis, ext. subrotundis mucronatis, int. oblongis obtusis, cor. glabrâ, achænio hirsutiusculo, pappi ser. ext. paleaceâ brevi. $\frac{1}{2}$ in Brasiliâ. Folia 2 $\frac{1}{2}$ poll. longa, 1 $\frac{1}{2}$ lata. Cor. purpureæ. Pappus albidus. Facies *Liabi sessiliflori*.

109. *V. CORDATA* (H. B. et Kunth nov. gen. 4. p. 31), fruticosa, ramis subangulatis obscurè tomentosis, foliis petiolatis coriaceis ellipticis obtusis basi cordatis subdentulatis utrinquè obscurè tomentosis, cymæ terminalis ramis dichotomis, capitulis multifloris, invol. squamis ovato-oblongis plurinerviis puberulis, achænio pluricostato, costis hispidis, pappi serie externâ brevi. $\frac{1}{2}$ in montibus Mexicanis prope urbem Tasco. Less. in linnæâ 1829. p. 267. 1831. p. 648. *V. cordifolia* Willd. herb. ex Less. Pappus flavescens. Cor. ign.

110. *V. LEOCARPA*, caule fruticoso tereti apice velutino demùm glabro, foliis petiolatis ovato-lanceolatis integerrimis basi subobtusis apice acuminatis penninerviis suprà pubescentibus demùm glabris, subtus lanato-tomentosis, petiolo nervoque velutino subrufo, paniculæ terminalis compositæ ramis subcorymbiferis polycephalis, pedicellis aphyllis capitula 3-5 ovata congesta 5-flora gerentibus, inv. squamis ovalibus obtusis ciliatis sublanatis, cor. et achænio glabris, pappi ser. ext. setis paucis dimidiam int. longitud. æquantibus. $\frac{1}{2}$ in Mexico legit cl. Karwinski. Petioli 8-10 lin. longi. Limbus fol. 4 poll. longus, 1 $\frac{1}{2}$ poll. latus. (v. s. in h. Acad. reg. Monac.)

§: 2. *Involucris squamis obtusis aut acutis nec acuminatis, capitulis in cymis scorpioideis dispositis aut rarò solitariis, folia perfecta basi non gerentibus.* Less. in linnæâ 1831. p. 648. — *Species omnes Americanæ.*

111. *V. FERRUGINEA* (Less. in linnæâ 1829. p. 271. 1831. p. 649), fruticosa, ramis subangulatis fulvo-tomentosis, foliis petiolatis coriaceis ellipticis obtusis integerrimis margine undulatis suprà scaberrimis puberulis, subtus fulvo-tomentosis, cymis scorpioideis later. et termin., capitulis campanulatis sessilibus multifloris, invol. squamis coriaceis ciliatis apice pubescentibus obtusis mucronatis, achæni glandulosi costis hispidis, pappi serie ext. brevi. $\frac{1}{2}$ in Brasiliâ tropicâ. Cor. albæ? glabræ. Pappus flavescens seu ferrugineus.

112. *V. DISCOLOR* (Less! in *linnæa* 1829. p. 274. 1831. p. 649), fruticosa, ramis teretibus apice angulatis cum pedunculis petiolis et facie fol. inferâ pube brevissimâ adpressâ densâ canis, foliis petiolatis oblongo-lanceolatis utrinquâ acuminate integerrimis suprà glabris, cymis scorpioides divaricatis, capitulis secundis brevè pedicellatis nudis, invol. turbinati squamis ovatis acutis, achenio oblongo multicostrato pilosiusculo, pappi serie ext. subpaleaceâ brevissimâ. 3 in Brasiliâ tropicâ (Sellow!) ad Cercovado prov. Rio-Janeiro (Vanth.! n. 340). *Eupatorium discolor* Spreng. syst. 3. p. 472. Pappus substramineus. Cor. glabrâ purpureâ. (v. s.)
113. *V. DIFFUSA* (Less. in *linnæa* 1829. p. 272. 1831. p. 649), herbacea erecta ramosa, foliis longè petiolatis membranaceis angustè elliptico-lanceolatis integerrimis suprà puberulis, subtus ramisque obscurè tomentosis, cymæ scorpioides ramis divaricatis, capitulis pedicellatis secundis, invol. subturbinati squamis coriaceis 1-nerviis acutis extus lanato-tomentosis, achenio piloso, pappi serie externâ paleaceâ brevi. — in Brasiliâ tropicâ. *Conyza divaricata* Spreng. syst. 3. p. 508. Pappus argenteus. Petioli 1-2 poll. longi. Limbus fol. 4-10 poll. longus, $1\frac{1}{2}$ -5 lin. latus.
114. *V. OPPOSITIFOLIA* (Less. in *linnæa* 1829. p. 273. 1831. p. 649), herbacea, ramis oppositis teretibus subtomentosis suprà folior. orig. subincrassatis, foliis oppositis petiolatis membranaceis ellipticis acuminatis utrinquè glandulas sessiles gerentibus, suprà pubescentibus, cymæ scorpioides terminalis ramis aphyllis divaricatis, capitulis multifloris pedicellatis, invol. turbinati squamis 1-nerviis ovatis acutis extus pilosis demum stellato-patentibus, achenio piloso multicostrato, pappi serie ext. paleaceâ brevi. 5 in Brasiliâ tropicâ, circa Rio-Janeiro (Less.). *Conyza stellata* Spreng. neu. entd. 2. p. 141. syst. 3. p. 508. Cor. glabrâ purpureâ. Pappus niveus.
115. *V. VERBENACEA*, herbacea ad collum fruticosa, caule erecto tereti apice nudo undiquè pube brevi conferta hirtello, foliis caulinis alternis rameis sæpè oppositis lineari-oblongis obtusis sessilibus subdenticulatis suprà scabridis, subtus hirsutis, corymbo paniculato aphylo, invol. squamis lanceolatis acutis glabris, achenio hirtello striatulo, pappi serie ext. brevi. 4 5 in campis editis prov. Sancti Pauli Brasiliæ, legit cl. Lund. Flor. novemb. — Herba 1-1 $\frac{1}{2}$ ped. Flores albi. Fol. 2-3 poll. longa, 3-4 lin. lata. (v. s.)
- ℓ. latior*, foliis oblongo-lanceolatis basi attenuatis magis dentatis, invol. subpuberulo. 4 5 Cum var. et legit cl. Lund. Fl. albi in inv. 10-12. Achenium longius forsan magis etate provectum? Folia 3 poll. longa, pollicem lata. (v. s. comm. cum var. à cl. inv.)
116. *V. SERRATA* (Less. in *linnæa* 1829. p. 275. 1831. p. 649), herbacea, caule tereti striato glabro ramoso, ramis pubescentibus, foliis petiolatis membranaceis ellipticis utrinquè brevè acuminatis argutè duplò serratis suprà glabris, subtus petiolisque pilosiusculis, cymæ terminalis scorpioides ramis divaricatis, capitulis multifloris lateralibus subsessilibus, invol. campanulati squamis scariosis 1-nerviis ovato-oblongis mucronato-acutis, achenio piloso multicostrato, pappi serie ext. paleaceâ brevi. — in Brasiliæ nemorosis montibus Mandiocce. Cor. violaceâ? glabrâ. Pappus argenteus subplumosus. Petioli 6-18 lin. longi. Folia 2 $\frac{1}{2}$ -8 poll. longa, 1-4 poll. lata.
117. *V. PUBERULA* (Less. in *linnæa* 1831. p. 649), caule fruticoso, ramis teretibus sordidè puberulis apice striatis, foliis petiolulatis oblongis basi acuminatis apice acutiusculis subserrulatis, suprà glabris, subtus petiolisque puberulis, cymâ terminali decompositâ subaphyllâ, capitulis 8-9-floris, invol. turbinati squamis imbricatis obtusis, ext. brevissimis, cæteris gradatim longioribus, achenio oblongo hirsuto-canescente, pappi ser. ext. paleaceâ brevi. 3 in Brasiliâ legit cl. Sellow. Inv. squamæ siccæ adpressè pubescentes. Pappus albus. Fol. 2 poll. longa, 6 lin. lata, petiolo 2-lin. (v. s. comm. à Mus. reg. Berol.)
118. *V. WESTIMANA* (Less. in *linnæa* 1831. p. 650), caule herbaceo simpliciter hispidiusculo pubescente tereti, foliis petiolatis firmis oblongis obovatisve subintegerrimis penninerviis, suprà scabris hirsuto-pubescentibus, subtus tomen-

toxis, panicula terminalis ramis subverticillatis aphyllis, capitalis subsessilibus 12-floris, inv. campanulati squamis siccis ciliatis ellipticis obtusis, cor. glabrâ, achenio hirsuto, pappi ser. ext. brevi setaceâ. — in montosis Brasiliæ prope Villa-Ricca. Cor. purpureo-violaceâ. Pappus rufescens. Affinis V. Beyrichii dicitur.

a. angustifolia, foliis oblongo-ellipticis basi obtusis semipollicem latis.

β. cuneata, foliis obovatis ultrâ pollicem latis basi cuneatis.

119. V. BEYRICHII (Less. in Linnæa 1829. p. 275. 1831. p. 651), herbacea, caule tereti striato glabro, foliis confertis petiolatis lanceolatis rugulosis subdenticulatis, suprâ glaberrimis, infrâ minutè glandulosis ad nervos pilosiusculis, cymis terminalibus scorpioideis divaricatis, capitulis subsessilibus multifloris, invol. turbinati squamis siccis multinerviis ovatis acutis pubescentibus et ciliatis, achenio oblongo multicostato, pappi serie ext. brevi. — in humidis prov. Sanctæ-Annæ ad Rio-Janeiro. Cor. rosæ glabrâ. Pappus flavescens. Folia 3-3½ poll. longa, 10-11 lin. lata. (v. s. comm. à cl. Lund.)

120. V. INCANA (Less. in Linnæa 1829. p. 277), caulibus herbaceis teretibus simplicibus pube adpressâ subincanis apice angulatis ramoso-corymbosis, foliis sessilibus lineari-lanceolatis acuminatis subdenticulatis, suprâ glabriusculis, subtus pilis adpressis incanis, cymis scorpioideis in corymbum dispositis subaphyllis, capitulis multifloris sessilibus et pedunculatis, invol. subtrabinati squamis adpressis oblongis acutis extus piloso-subincanis, achenio oblongo multicostato piloso et glanduloso, pappi serie externâ internâ subæquantè. ♀ in Brasiliâ meridionali. Rhizoma multiceps lignosum. Cor. violaceâ glabrâ. Pappus argenteus. Herba fructifera refert Hieracium cymosum. (v. s. comm. à Mus. reg. Berol.)

121. V. ECHIOIDES (Less. in Linnæa 1829. p. 278. 1831. p. 651), caulibus è rhizomate lignoso pluribus teretibus simplicibus striato-sulcatis strigoso-pubescentibus, foliis sessilibus membranaceis utrinquè asperis elongato-lanceolatis undulato-crenatis integerrimis margine revolutis, cymâ divaricatè scorpioideâ ramis dichotomis, capitulis multifloris subsessilibus, invol. campanulati squamis oblongis 1-nerviis glabris, achenio multicostato piloso, pappi serie ext. brevi. ♀ in Brasiliâ meridionali. Habitus Hieracii echioidis. Cor. glabrâ violaceâ. Pappus niveus.

122. V. DENTICULATA, caule herbaceo erecto sulcato-angulato apice tenuissimè puberulo, foliis petiolatis lanceolatis utrinquè acuminatis serratulis paucis distantibus denticulatis, suprâ glabris, subtus pube tenuissimâ subpuberis, paniculâ laxâ polycephalâ aphyllâ, ramis infer. axillaribus hinc inde suboppositis folio longioribus, capitulis secus ramos sessilibus lateralibus 10-12-floris, inv. basi obtusis squamis glabris ext. brevioribus acutis, int. oblongis obtusis mucronulatis, cor. glabrâ, achenio brevissimo pubero striato, pappi ser. ext. setosâ brevi. — in Brasiliâ circa Rio-Janeiro. Folia 2-2½ poll. longa, 6 lin. lata, petiolo 2-3 lin. Pappus stramineo-subrufescens. Affinis videtur V. Beyrichii ex descr. Folia summa integra. (v. s. comm. à cl. Lund. ! Gaudich. ! Blanch.)

123. V. PITTAGORUM, arborea, ramis glabris, ramulis subhirtellis angulatis, foliis subsessilibus lineari-lanceolatis utrinquè acuminatis integerrimis, suprâ minutè scabridis, subtus hirsutis, cymâ ramosissimâ paniculæformi polycephalâ, invol. squamis ovatis glabriusculis, exter. mucronatis, inter. obtusiusculis, achenio pubero, pappi serie ext. setosâ brevi. ♀ in montosis circa Rio de Janeiro. Vulgatissima arbor, vulgo *capueiro* dicta. Herba tota et præsertim flores suaviter olentes. Flor. jul. aug. Trochili paniculas circumvolant. Pittacus vittatus (Kuhl.) semina avidè devorat (Lund.!). Folia 3-4 poll. longa, 6-8 lin. lata. Cortex lenticellis verruculosus. (v. s. comm. à cl. Lund.)

124. V. HERACLADA, fruticosa, ramis teretibus striatis pube brevissimâ rufâ puberulis, foliis petiolatis oblongo-lanceolatis utrinquè acuminatis integerrimis, suprâ aspero-scabridis, subtus hirsuto-tomentosis, cymæ polycephalæ ramis subscorpioideis, ramis inferioribus ex axillis fol. ortis, sup. nudis, invol. squamis

ovali-oblongis mucronatis, capit. aliis sessilibus aliis pedicellatis circ. 10-floris, achæniis striato-sulcatis puberis, pappi serie ext. brevi. $\frac{5}{8}$ in Brasiliæ prov. S^t Pauli frequens obs. cl. Lund. Fl. decemb. Valdè affinis *V. erioclada*, sed rami vix puberi, folia dimidio angustiora, panicula basi foliosa, et inv. squamæ mucronatæ nec acuminatæ. Pappus sordidè rufus. Cor. purpurea. Folia $\frac{3}{4}$ poll. longa, 5-6 lin. lata. (v. s. comm. à cl. Lund.)

125. *V. PETIOLARIS*, fruticosa, ramis teretibus velutino-tomentosis, foliis alternis longè petiolatis elliptico-oblongis utrinquè acuminatis integerrimis, suprâ sparsè puber. subtus molliter petiolisque hirsuto-tomentosis, paniculæ ampla ramis scorpioideis aphyllis, capitulis subsessilibus unilateralibus 30-floris, invol. turbinati squamis multiseriis semipatentibus dorso velutinis oblongis acutis, ext. brevibus secûs pedicellum imbricatis, cor. glabrâ, achænio villosò, pappi ser. ext. brevi vix paleacè, $\frac{5}{8}$ in Brasiliæ sylvis primævis leg. cl. Lhotsky. Petioli 9-10 lin. longi, limbi fol. 5-6 poll. longi, sesquipoll. lati. Inv. squamæ nec adpressæ nec squarrosæ, sed semipatentes! (v. s. comm. à cl. Lhotsky.)

β . *pauciflora*, foliis basi magis obtusis, capitulis paulò minoribus 10-12 floris. $\frac{5}{8}$ ad margines sylvarum circa Rio-Janeiro legit cl. Lhotsky. Arbor 25-30-pedalis. (v. s. comm. à cl. invent.)

126. *V. EUPATORIIFOLIA*, arborea, ramis teretibus densè pedunculis petiolisque hirsuto-tomentosis, foliis oppositis petiolatis ellipticis acuminatis basi obtusis integerrimis glandulas sessiles utrinquè gerentibus, suprâ pubescentibus nervo medio basi hirsuto, subtus hirsutis, paniculæ ramis ex axillis supr. aut ad apicem ortis subsorpioideis polycephalis, capitulis pedicellatis 30-floris, invol. turbinati squamis ovali-oblongis extus villosis demùm semipatulis, ext. brevioribus, cor. glandulosâ glabrâ, achænio puberulo, pappi serie ext. paleacè brev. $\frac{5}{8}$ in sylvis collium circa Rio-Janeiro legit cl. Lund. Affinis *V. petiolaris* sed oppositifolia! Differt à *V. oppositifolia* caule arboreo 8-12-pedali nec herbaceo suprâ fol. paria non dilatato, ramis folisque magis hirsutis. (v. s. comm. à cl. Lund.)

β . *pauciflora*, foliis basi subacutis parcius glandulosis, capitulis 10-12-floris, corollis subglandulosis, $\frac{5}{8}$ in fruticetis circa Rio-Janeiro legit cl. Lhotsky. Frutex 5-7-pedalis. (v. s. comm. à cl. invent.)

127. *V. HAVANENSIS*, caule fruticoso tereti ramoso glabro, ramulis striatis puberulis, foliis ovalibus utrinquè acuminatis basi cuneatis brevissimè petiolatis integerrimis penninerviis, suprâ muriculatis, subtus glandulosis et tenuiter pubescentibus, cymis axill. terminalibus pedunculatis laxis oligocephalis, capitulis pedicellatis aphyllis distantibus 25-floris, invol. turbinati squamis dorso glandulosis glabriusculis, ext. brevibus acutis, int. oblongis subcuneatis obtusiusculis, cor. et achænio glabris parè glandulosis, pappi ser. ext. brevi setiformi. $\frac{5}{8}$ in ins. Cubâ juxta Havanam legit cl. Ramon de la Sagra. Invol. subpurpurascens. Cor. purpurea. Pappus albus. Folia 4 poll. longa, sesquipollicem lata, juniora marginibus subrevoluta. (v. s. comm. à cl. R. de la Sagra.)

128. *V. LANCEOLARIS*, caule tereti striato pube ferrugineâ parè velutino, foliis petiolatis lanceolatis basi subobtusis apice acuminatis serrulatis, suprâ glabris, subtus petiolisque parè velutinis, paniculæ ramis axillaribus ebracteatis ramosis velutinis, capitulis sessilibus ovatis 20-floris lateralibus et terminalibus, invol. squamis omnibus mucronatis dorso subpuberis, cor. glabrâ, receptaculo alveolato, achænio puberulo, pappi ser. ext. brevi vix paleacè. $\frac{2}{3}$ in Mexico ex h. Haenke. Affinis *V. Haenkeana*. Pappus croceo-rufus rigidulus. Folia 5 poll. longa, pollicem lata, petiolo 4 lin. (v. s. in h. Haenke à cl. Sternberg missum.)

129. *V. HAENKEANA*, caule teretiusculo erecto ramis pedunculisque velutino-subhirsutis, foliis petiolatis ellipticis basi acutis apice acuminatis integerrimis penninerviis, suprâ demùm glabris, subtus velutino-subhirsutis, ramis axillaribus foliosis apice paniculatis, ramulis bi-trifidis, capitulis axillaribus latera-

libus sessilibus aut pedicellatis ebracteatis 20-floris, invol. squamis oblongis glabriusculis ext. mucronatis inter. obtusis, cor. glabrâ, achenio pilosiusculo, pappi ser. ext. brevissimâ vix paleaceâ. $\frac{1}{2}$? inter Peruanas Orinocoenses in herb. habet cl. Haenke. Folia 5-6 poll. longa, 2 poll. lata, petiolo 7-9 lin. longo velutino. Pappus croceo-rubens rigidulus. Pubes ramorum rufescens. (v. s. in h. Haenke à cl. Stenberg missum.)

130. *V. MICRADENTIA*, fruticosa, ramis teretibus densè et breviter sericeo-villosis foliis petiolatis ellipticis basi attenuatis apice acuminatis integerrimis penninerviis utrinquè secus nervos et in petiolo sericeo-pubescentibus suprâ sparse pilosis subtus sparè glandulosis, cymis scorpioideis, aliis axillaribus oligocephalis folio brevioribus aliis terminalibus stellato-divaricatis polycephalis, capitulis unilaterialibus ebracteatis 10-floris, invol. squamis dorso subvillosis 1-nerviis, ext. acuminatissimis, int. linearibus vix acutis, achenio subvillosio, pappi ser. ext. paleaceâ brevi. $\frac{1}{2}$ in Amer. calid. fortè Peruvia legit cl. Poeppig. (pl. exs. n. 1215.) Affinis ex descr. *V. frangulæfolia*. Folia caul. 6 poll. longa, 2-2 $\frac{1}{2}$ poll. lata, petiolo 8-9 lin. longa. (v. s. comm. à cl. Poeppig.)

131. *V. MENTHÆFOLIA* (Less. in linnæa 1829. p. 253. 1831. p. 645), fruticosa, ramis teretibus divaricatis apice obscure pubescentibus demum glabris, foliis petiolatis ovato-oblongis acutis basi attenuatis membranaceis utrinquè scabridis subtus pilosiusculis, cymis subfastigiatis dichotomis divaricatis, capitulis 15-floris, invol. campanulati squamis 1-nerviis cōriaceis ovatis acutis ciliatis, intimis obtusis glabris, cor. glabrâ, achenio glabriusculo, pappi serie ext. setiformi brevior. $\frac{1}{2}$ in fruticetis insulæ Cubæ. Eupatorium menthæfolium Poeppig. nus. ex Less. Cor. glabrâ. Pappus flavescens. Folia 1-3 poll. longa, 4-12 lin. lata, integra aut serrata. (v. s. comm. à cl. Ramon de la Sagra.)

§. 3. *Involucris squamis plerisque acuminatis, capitulis in cymis scorpioideas dispositis rarò subsolitariis, basi folia perfecta non gerentibus.* Less. in linnæa 1831. p. 651. — *Species omnes Americanae.*

132. *V. ODORATISSIMA* (H. B. et Kunth nov. gen. 4. p. 32), caule fruticoso tereti, ramis striatis velutino-pubescentibus, foliis brevè petiolatis rigidis obovatis basi cuneatis apice subdentatis, suprâ scabridis, subtus reticulatis hirsuto-pubescentibus, cymæ ramis plurimis ramosis in paniculam dispositis, capitulis innumeris pedicellatis later. et terminal. folio florali destitutis, invol. squamis pluriseriatis ovatis acuminatis, achenio vix pilosiusculo, pappi serie ext. setaceâ brevi. $\frac{1}{2}$ ad Caracas, Ibaque et Honda. Less. in linnæa 1829. p. 271. 1831. p. 631. Conyza clethræfolia et Eupatorium obovatum, Willd. herb. Cor. albæ glabræ. Pappus stramineus. Variat foliis serratis ex Kunth, crenatis in specim. meo Caracasano, et subintegerrimis ex Less. (v. s.)

133. *V. RUBRIRAMEA* (Mart. herb.), caule herbaceo quinquangulato glabriusculo erecto, foliis subpetiolatis oblongo-lanceolatis basi acuminatis apice subacutis subintegerrimis, suprâ glabris glabriusculisve, subtus pubero-subhirsutis, panicula ramis axillaribus polycephalis dichotomo-ramosis folio longioribus subaphyllis, capitulis lateralibus sessilibus terminalibus ebracteatis 20-floris, invol. squamis lineari-oblongis mucronatis glabris, cor. glabrâ, achenio striato pubero, pappi ser. ext. paleaceâ acuminatâ vix inter. triplò brevior. $\frac{1}{2}$? in campis prope Gongonhas de Sabara prov. Minas gener. Brasiliæ legit cl. Martius. Folia 4 poll. longa, 7-8 lin. lata. Rami purpurei. Corollæ albæ. Capit. vix 3-lin. longa. (v. s. in h. Acad. reg. Monac.)

134. *V. ASSANS* (Mart. herb.), caule herbaceo? erecto striato-angulato breviter velutino, foliis brevè petiolatis elliptico-oblongis utrinquè obtusis rigidis suprâ scabris, subtus velutino-puberis penninerviis nervis lateralibus reticulatis, ramis floridis bifidis subscorpioideis oligocephalis folio longioribus aphyllis, capitulis ovatis 30-floris sessilibus, invol. squamis imbricatis glabriusculis acuminatis apice non adpressis, cor. glabrâ, achenio pubero, pappi ser. ext. vix paleaceâ

brevissimâ panicetâ. $\frac{1}{2}$ $\frac{1}{2}$ in campis desertis Minarum gener. Brasiliæ collecti cl. Martius. Petioli 3 lin. Limbus fol. $2\frac{1}{2}$ -3 poll. longus, pollicem latus. Pappus rufus. Recept. areolatum. (v. s. in h. Acad. reg. Monac.)

135. *V. POLYANTHES* (Less. in linnæa 1831. p. 631), subarborea, ramis angulatis puberulis, foliis petiolatis lanceolatis utrinquè acuminatis subintegerrimis suprâ scabris, subtus puberis nervis elevatis subhirsutis, paniculâ ramosissimâ puberâ ramis scorpioideis subverticillatis divergentibus aphyllis, invol. 15-20-flori squamis oblongis subacuminatis glabriusculis, achænio puberulo punctato, pappi seriebus setiformibus subæqualibus. $\frac{1}{2}$ in montosis Brasiliæ circa Rio-Janeiro alt. 2-3000 ped., et in prov. Minar. gener. Eupatorium polyanthes Spreng. syst. 3. p. 435. Vern. patens Less. in linnæa 1829. p. 276. ex parte et excl. syn. Vern. Brasiliensis Mart. ! herb. Cor. glabræ albæ. Folia trita odorata (Lund.) 5-6 poll. longa, pollicem lata. Pappus flavescens. (v. s. comm. à cl. Lund.)

136. *V. GRANDIFOLIA*, arborea, ramis teretibus striatis apice subpubescentibus, foliis petiolatis ellipticis utrinquè acuminatis simpliciter et grossè serratis membranaceis glabriusculis secus nervos et venas minutè puberulis, paniculâ corymbosâ amplâ divaricatâ polycephalâ aphyllâ, capitulis pedicellatis 20-25-floris, invol. squamis glabris acuminatis margine membranaceis, cor. lobis puberulis, achænio tereti turbinato striato hispidovilloso, pappi ser. ext. paleaceâ brevi. $\frac{1}{2}$ in monte Cercovado Brasiliæ legit cl. Lund. Arbor 8-10 ped. Folia inf. 16 poll. longa, 8 poll. lata, petiolo 2 pollic. Pappus albus caducissimus. Recept. alveolare. Inv. demum patens. (v. s. comm. à cl. Lund.)

137. *V. PATENS* (H. B. et Kunth nov. gen. 4. p. 3a.), arborea, ramulis angulatis obscure pubescentibus, foliis breviter petiolatis angustè lanceolatis acuminatis basi angustatis subcrenato-undulatis suprâ rugosis scabriusculis, subtus hirtopubescentibus, cymæ scorpioideæ terminalis ramis patentibus, capitulis multifloris distantibus subsessilibus, invol. campanulati squamis siccis 1-nerviis ovatis acutis adpressis pilosiusculis, achænio multicostato piloso glanduloso, pappi ser. ext. paleaceâ brevi. $\frac{1}{2}$ in Amer. merid. (Humb.) V. patens Less. in linnæa 1831. p. 653. Baccharis rugosa Willd. herb. ex Less. Cor. glabræ albæ.

138. *V. SCARRA* (Pers. ench. 2. p. 404. Less. in linnæa 1831. p. 653), fruticosa, ramis angulatis divaricatis apice hirtello-puberulis, foliis brevissimè petiolatis ellipticis aut obovatis utrinquè obtusis subintegerrimis suprâ scabris, subtus hirsuto-tomentosis, paniculis terminalibus ramosissimis aphyllis, capitulis subsessilibus aut pedicellatis 30-40 floris, invol. turbinatis squamis glabriusculis oblongo-linearibus acuminatis, cor. glabrâ, achænio pubescente, pappi ser. ext. brevi setosâ. $\frac{1}{2}$ in Brasiliâ (ubi ex Marcgr. dicitur *Tremate*), in collibus juxta Bahiam (Salzm. ! Blanch. !), in montibus prov. Minarum gener. Mart. ! Baccharis Brasiliensis Linn. spec. 1205. — Marcgr. bres. 2. p. 81. fig. sup. Pappus subrufescens. Cor. purpurascens vanillam (Salzm. !) styracem (Marcgr.) olentes. Folia 12-15 lin. longa, 7-8 lin. lata. (v. s.)

139. *V. POLYCEPHALA*, caule fruticoso, ramis teretibus velutino-canescens, foliis brevè petiolatis ovalibus utrinquè obtusis margine subcrenulatis et revolutis suprâ subrugosis pubescentibus, subtus tomentoso-subincanis, cymæ ramis plurimis ramosis in paniculam dispositis, capitulis 20-floris axillaribus et lateralibus subsessilibus folio floralis destitutis, involucri obovati squamis pluriseriatis ovatis subacutis, achænio vix pubero, pappi serie ext. vix paleaceâ brevi. $\frac{1}{2}$ in collibus aridis circa Bahiam Brasiliæ legit cl. Salzmann. Caul. 4-pedalis. Rami cani. Cor. pallidæ. Pappus stramineo-subrufescens. (v. s.)

140. *V. MICROCEPHALA*, suffruticosa, ramis teretibus pube brevi subhirsutulis, foliis brevissimè petiolatis oblongis obtusis basi subcordatis integerrimis suprâ scabridis glabris subtus ferrugineo-villosis, cymis terminalibus subcorymbosis, ramis nudis lateralibus scorpioideis, capitulis 20-floris lateralibus subsessilibus obovatis, invol. squamis pluriserialibus imbricatis ext. brevissimis acutis, intimis longioribus, achænio glabriusculo, pappi serie ext. paleaceâ brevi. $\frac{1}{2}$ in Amer. calidior. legit cl. Née, sed locus proprius ignotus. Affinis *V. odora*

tissimæ et acabræ. Capitula minora. Folia 2-2 $\frac{1}{2}$ poll. longa, 6-9 lin. lata. (v. s. ex herb. Thibaud.)

141. *V. SORORIA*, suffruticosa, ramis teretibus hirsutulis, foliis breviter petiolatis ovatis acuminatis basi obtusis integerrimis suprâ villosis, subtus densè villosio-hirsutis, ramis floridis paniculatis elongatis bifidis hirsutulis, ramulis capitula ebracteata sessilia remota unilateralia gerentibus, invol. ovati hirsuti squamis omnibus acuminatis subpungentibus. $\frac{5}{8}$ in Brasiliâ circa Rio de Janeiro legit cl. Gandichaud. Valdè affinis *V. microcephalæ*, sed differt ramis verè hirsutis, foliis ovatis suprâ villosis, ramis cymæ magis elongatis, invol. hirsuto. Flores in specim. juniores et non satis vidi. (v. s. comm. à cl. inv.)
142. *V. MYRIOCEPHALA*, caule ramoso densè et adpressè velutino, foliis petiolatis ovali-lanceolatis utrinquè (apice præcipuè) acuminatis integerrimis penninerviis, subtus venoso-reticulatis suprâ glabris, subtus glandulosos et breviter pubescentibus, paniculæ amplæ ramosissimæ ramis axillaribus folio longioribus bifidis scorpioideis, capitulis lateralibus et terminalibus sessilibus solitariis ebracteatis 10-floris, invol. squamis ext. acuminatis ciliatis int. glabrisculis mucronatis, cor. glabrâ, achænio turbinato sericeo-villoso, pappi ser. ext. paleaceâ brevi. $\frac{2}{3}$? inter Peruanas Orinocenes herb. Haenkeani adest. Affinis *V. polyanthi*. Pubes caulis et pedunculorum subflavescentes velutina. Folia 4 poll. longa, 1 $\frac{1}{2}$ -2 poll. lata, petiolo semipollic. Invol. purpureascentia. Pappus albus. Cor. purpureæ. (v. s. in h. Haenk. à cl. de Sternberg missum.)
143. *V. DEPPHANA* (Less. in linnæâ 1831. p. 308 et 655), fruticosa, ramis teretibus obscure pubescentibus, foliis petiolatis membranaceis penninerviis ellipticis utrinquè subattenuatis apice mucronatis integerrimis suprâ scabris, subtus tomentoso-pubescentibus, cymis scorpioideis compositis in paniculam myriocephalam dispositis, capitulis secus ramos lateralibus sessilibus turbinatis 20-floris, invol. squamis rigidis mucronato-acuminatis, cor. glabrâ, achænio piloso, pappi ser. ext. paleaceâ brevi. $\frac{5}{8}$ in Mexico circa Misantla. Cor. albæ odoratissimæ. Folia 3-3 $\frac{1}{2}$ poll. longa, 12-15 lin. lata, petiolo 2-5 lin. (v. s. in h. Acad. reg. Monac.)
144. *V. ADPROXIMATA* (Less. in linnæâ 1831. p. 655), fruticosa, ramis teretibus striatis sordidè tomentosis, foliis petiolatis ellipticis basi cordatis subcrenatis suprâ scabrisculis, subtus molliter pubescenti-tomentosis, cymis scorpioideis carvatis aphyllis, capitulis 11-floris lateralibus pedicellatis, invol. turbinati squamis 8-seriatis siccis acuminatis muticis dorso tomentoso-incanis, ext. rotundia, cor. glabrâ, achænio villosa, pappi ser. ext. paleaceâ brevi. $\frac{5}{8}$ in Brasiliâ. Cor. purpureo-violaceæ. Fol. circ. semipedalia, 3-4 poll. lata. Similis *V. oppositifoliæ* et quinquefloræ dicitur.
145. *V. QUINQUEFLORA* (Less. in linnæâ 1831. p. 656), caule herbaceo tereti cum pedunculis petiolis folisque subtus hirsuto-pubescentibus, foliis petiolatis elliptico-oblongis utrinquè acuminatis rariter subserratis, suprâ scabris, cymâ terminali compositâ parçè ad basin pedunculorum bracteata, capitulis nudis pedicellatis ovali-oblongis 5-floris, invol. squamis siccis dorso villosio-pubescentibus imbricatis subacuminatis, achænio hirsuto, pappi ser. ext. brevi subpaleaceâ. $\frac{2}{3}$ in Brasiliâ legit cl. Sellow. Folia 5-6 poll. longa, 15 lin. lata. (v. s. comm. à Mus. reg. Berol.)
146. *V. FRANGULEFOLIA* (H. B. et Kunth nov. gen. 4. p. 37), fruticosa, ramis teretibus villosis, foliis brevè petiolatis oblongis acutis basi angustatis integerrimis rigidis suprâ pubescentibus nitidis, subtus glandulosos pubescenti-pilosis canescentibus, cymæ scorpioideæ terminalis ramis congestis fastigiatis, capitulis multifloris approximatis sessilibus, invol. hemisphæric squamis lanceolatis acutis 3-nerviis externis tomentoso-pubescentibus, achænio piloso, pappi serie ext. paleaceâ brevi. $\frac{5}{8}$ in calidis Novo-Granatensium prope Honda. Less. in linnæâ 1829. p. 276. 1831. p. 657. Cor. glabra violacea. Pappus argenteus. Fol. 1-2 $\frac{1}{2}$ poll. longa, 4-9 lin. lata, petiolo 2 lin.
147. *V. CANESCENS* (H. B. et Kunth nov. gen. 4. p. 35. t. 317. Less. in linnæâ 1829. p. 284. 1831. p. 657), caule basi suffruticosa, ramis teretibus pilis ad-

pressis canescentibus, foliis brevè petiolatis rigidis membranaceis oblongo-lanceolatis acuminatis basi angustatis integerrimis subrugosis suprà pubescentibus, subtùs piloso-canescentibus glandulosis, cymæ terminalis scorpioideæ ramis oligocephalis, capitulis distantibus sessilibus, involucri campanulati squamis siccis 1-nerviis acuminatis pilosis glandulosisque, achænio villosa, pappi serie ext. brevissima. 2/ 3 in Peruvia temperatâ prope Guamcabamba. *Lepidaploa canescens* Cass. dict. 26. p. 18. Vern. reticulata Willd. herb. ex Less. V. canescens Spreng. syst. 3. p. 437. excl. syn. Pers. et Sellow? V. geminata Kunth nov. gen. 4. p. 35? Cor. pallide violaceæ. Pappus argenteus. (v. s.)

β. *micrantha* (Less. in linneæ 1831. p. 637), suffruticosa? ramis subangulatis sericeo-pilosis, foliis petiolatis oblongis acuminatis basi rotundatis integerrimis suprà pubescentibus, subtùs argenteo-piloso-siccis, cymis terminalibus paniculaformibus ramosis, ramulis geminis dichotomis, capitulis solitariis sessilibus distantibus multifloris. 3/4 in prov. Comanensi? Vern. *micrantha* H. B. et Kunth nov. gen. 4. p. 37. V. *acuminata* Willd. herb. ex Less.

148. V. *SUBSQUARROSA*, ramis teretibus supernè villosa-hirsutis, foliis brevissimè petiolatis ovato-oblongis, basi obtusis apice acutis integerrimis suprà glabris rugosis, subtùs villosa-hirsutis subcanescentibus, cymis axillaribus terminalibusque bifidis divaricatis scorpioideis, capitulis ovatis 20-floris sessilibus, invol. squamis villosis lanceolatis externis subpungenti-acuminatis, acumine patente squarrosis, int. minutis acuminatis erectis, cor. glabris, achænio turbinato villosa, pappi serie ext. paleaceâ brevi. 2/4 in Brasiliâ ad margines sylvarum et aqueductus circa Rio-Janeiro (Mart.!) et Bahiam (Blanch.). Cor. verisim. purpureæ. (v. s.)

149. V. *MOLLIS* (H. B. et Kunth nov. gen. 4. p. 28), caule herbaceo tereti sericeo-villosa, foliis brevè petiolatis ovato-lanceolatis acuminatis basi angustatis remotè denticulatis integerrimisve suprà puberulis, subtùs argenteo-lauginosis, cymæ terminalis scorpioideæ ramis sæpè geminis elongatis, capitulis sessilibus ebracteatis 20-floris, inv. campanulati squamis 1-nerviis sericeo-pilosis acuminatis demùm patentibus, cor. parçè pilosis, achænio parçè sericeo-piloso, pappi serie ext. paleaceâ brevi. 2/4 in Peruvia? (herb. Dombey!) ad ripam fl. Catamayo prope Gonzanama Novo-Granatensium ex Humb. itin. et ad Orinocum ex herb. Hænk! Fol. 3-5 poll. longa, 1-2 poll. lata. Pappus argenteus. *Empatorium polystachyum* Willd. herb. ex Less. Vern. *mollis* Less. in linneæ 1823. p. 283. 1831. p. 657. (v. s. in h. Hænk. à cl. Sternb. missum.)

150. V. *FLAVESCENS* (Less. in linneæ 1831. p. 657), fruticosa, ramis teretibus striatis rufulo-tomentosis, foliis petiolatis ovatis subrotundisve suprà lævibus subtùs glandulosis flavido-villosis subintegerrimis, cymæ ramis scorpioideis patentibus, capitulis sessilibus lateralibus multifloris, recept. nudo, cor. glabris, achænio piloso, pappi ser. ext. paleis latissimis ovario brevioribus. 3/4 in Brasiliâ, ad Serra d'Estrella. Vern. scorpioideus Less. in linneæ 1829. p. 282. excl. syn. Folia basi obtusa aut acuta. Capitula approximata.

151. V. *TOURNEFORTIODES* (H. B. et Kunth nov. gen. 4. p. 34), herbacea, ramis teretibus striatisque flavido-pubescentibus, foliis breviter petiolatis ovato-lanceolatis subintegerrimis suprà scabris hirsutiusculis, subtùs pubescentibus punctatis, spicis scorpioideis geminatis, capitulis sessilibus 25-floris, recept. levissimè alveolato, corollis glabris, achænio glabrisculis, pappi serie ext. paleaceâ ovario subæquali. — in Americâ merid., sed locus propr. ignotus. V. *Tournefortioides* Less. in linneæ 1831. p. 657 non 1829. *Stæhelinia solidaginoides* Willd. herb. ex Less.

152. V. *SCORPIOIDES* (Pers. ench. 2. p. 404.), fruticosa, ramis teretibus puberulis, foliis breviter petiolatis ovali- aut ovato-lanceolatis acuminatis subdentatis aut subrepandis suprà glabriusculis, subtùs pubescentibus hirsutisve, cymæ ramis scorpioideis, capitulis secus ramos sessilibus 1-lateralibus ebracteatis circ. 20-floris, corollæ lobis pilosis, achænio turbinato piloso, recept. paleaceo fimbriifero, pappi ser. ext. vix paleaceâ achæni longitudine. 3/4 in

Brasiliâ tropicâ, Guianâ. *V. scorpioides* Less. in linnæâ 1831. p. 658. non alibi. *V. Tournefortioides* Less. in linnæâ 1829. p. 281. non Kunth *V. centriflora* Link et Otto abb. t. 55. *Lepidaploa scorpioides* Cass. dict. 26. p. 18. *Coryza scorpioides* Lam. dict. 2. p. 48.

a. *centriflora*, foliis ovalibus utrinquè acuminatis suprà glabris, subtùs pubescentibus. In Brasiliâ, Guianâ. Satis refert icon. Link. (v. s.)

β. *subrepanda*, foliis latè ovatis subrepandis apice acuminatis suprà glabris, subtùs pubescentibus. — in Guianâ gallicâ et Brasiliâ. Vern. *subrepanda* Pers. ench. 2. p. 404. (v. s.)

γ? *subtomentosa*, foliis latè ovatis subdentatis apice acuminatis suprà puberis, subtùs hirsutis. — ad Caracas legit cl. Vargas. (v. s.)

δ? *longifolia*, foliis elongato-lanceolatis acuminatis suprà glabris secus nervos puberis, subtùs præsertim in nervis puberis. — circa Bahiam in umbrosis legit cl. Salzmann. (v. s.)

ε? *longeracemosa*, foliis elongato-lanceolatis acuminatis suprà subscabris subtùs hirsuto-subtomentosis, spicis longissimis. — in Brasiliâ, Peruvîâ. Vern. *longeracemosa* Mart.! herb. — Vauth.! pl. exs. n. 262. Poepp.! n. 33. (v. s.)

153. *V. ignobilis* (Less. in linnæâ 1831. p. 658), fruticosa?, ramis teretibus hispidiusculo-pubescentibus, foliis petiolatis rigidis adproximatis oblongo-obovatis obtusis suprà asperimis, subtùs hirsutiusculo-pubescentibus, cymis scorpioideis aphyllis, pedunculis 2-3-cephalis, capitulis 10-floris, invol. semiglobosi squamis siccis glabrisculis r-nerviis acuminatis ext. rotundis, cor. glabrâ, achænio villosa, pappi seriebus confluentibus setaceis. ♂? in Brasiliâ. Cor. violacea. Fol. 3. poll. longa, pollicem lata.

154. *V. obovata* (Less. in linnæâ 1829. p. 279. 1831. p. 659), caule herbaceo simplici tereti striato villosa, foliis sessilibus membranaceis obovatis undulato-crenatis suprà scabris pilosis subtùs villosis, cymæ terminalis scorpioideæ ramis divaricatis dichotomis, capitulis adproximatis multifloris sessilibus, invol. campanulati squamis paucis siccis 3-nerviis adpressis acuminatis extùs pilosis, achænio subsericeo, pappi serie ext. palacæâ brevi. — in Brasiliâ tropicâ. Cor. glabrâ violacæâ. Pappus argenteus. Folia variant oblonga aut elliptica, obtusa aut acuta, 2 $\frac{1}{2}$ -4 poll. longa, 10-20 lin. lata.

155. *V. pellita* (H. B. et Kunth nov. gen. 4. p. 28. Less. in linnæâ 1829. p. 279), fruticosa?, ramis teretibus sordidè lanatis, foliis sessilibus lanceolatis coriaceis margine revolutis suprà rugosis scabro-pilosis, subtùs obscurè lanatis, cymæ terminalis scorpioideæ ramis dichotomis, capitulis adproximatis sessilibus circ. 20-floris, invol. campanulati squamis siccis adpressis acuminatis pungentibus extùs et apice lanatis, achænio oblongo densè sericeo-villosa, pappi serie ext. brevi vix palacæâ. ♂ in prov. Caracasanâ. Cor. ex Kunth glabrâ, ex Less. pilosâ, in nostro specim. lobis apice parçissimè pilosis. Pappus sordidus, ex Less. albus, in spec. Varg. rigidulus facilè deciduus. (v. s. comm. à cl. Vargas.)

156. *V. heptantha*, caule tereti erecto densè et obscurè velutino-hirsuto, foliis petiolatis elliptico-oblongis basi subacuminatis apice cuspidato-acuminatis subintegerrimis suprà brevissimè puberulis, subtùs petiolisquè densè hirsutis, paniculæ ramis axillaribus folio brevioribus simplicibus bifidis ebracteatis, capitulis sessilibus confertis spicatis 7-floris, inv. squamis lineari-lanceolatis subacuminatis dorso villosis, cor. lobis apice puberulis, achænio turbinato sericeo-villosa, pappi ser. ext. angustè palacæâ acuminatâ int. quadruplò breviorè. 2? ♂? inter Peruanas montanas Orinocences habet herb. Hænk. Fol. 3 poll. longa, 9 lin. lata, petiolo 3-4 lin. marginibus subrecurvatis. Cor. purpureâ. Tota planta in sicco obscura. (v. s. in h. Hænke à cl. Sternb. missum.)

157. *V. virgulata* (Mart.! herb.), caule herbaceo? erecto tenui virgato tereti striato glabro, foliis infer. . . . super. distantibus linearibus minimis integerrimis subtùs pubescentibus, capitulis 10-12 floris secus ramos spicatis sessilibus aphyllis, inv. squamis oblongis dorso adpressè pubescentibus, ext. brevibus mucronatis, intimis longioribus subacuminatis, cor. puberâ, achænio

sericeo-villoso; pappi ser. ext. angustè palacèà acuminatà brevi. — \mathcal{Z} ? in editis prov. Minarum gener. Brasiliæ. Folia sup. vix 2-3 lin. longa, inf. ignota. Pappus substramineus. Recept. nudum. (v. s. in h. Acad. reg. Monac.)

158. *V. DESKARTORUM* (Mart. herb.) glabriuscula, collo crasso, caulibus pluribus subsimplicibus sparsè foliosis apice hirsuto-villosis nudis 1-cephalis, foliis sessilibus linearibus acuminatis integerrimis planis nervoso-subreticulatis sparsè pilosis glabratiss, invol. 1-2-serialis squamis lanceolatis extùs villosis. — \mathcal{Z} in deserto trans flumen Sancti Francisci prov. Minarum gener. Brasiliæ legit cl. Martius. Herba 6 poll. longa. Affinis *V. campestri*. Folia 2. poll. longa vix 2 lin. lata. Cap. dimidio minus quàm *V. campestris*. Achænium sericeo-villosissimum. Pappi pallidè rufescentis palacè ext. breves lineares acuminatæ, int. elongatæ setiformes. Ad *V. simplicem* Less. valdè accedit. Flores in invol. circ. 30. An hujus sectionis? (v. s. in h. Acad. reg. Monac.)

159. *V. SIMPLEX* (Less. in linnæa 1829. p. 280. 1831. p. 659), caulibus è rhizomate lignoso-tuberoso pluribus simplicibus teretibus striatis villosis, foliis sessilibus oblongis acutis subintegerrimis suprà pilosis, subtùs villosis-lanatis, cymis scorpioides contractis oligocephalis ebracteatis, capitulis 25-floris, involucri campanulati squamis acuminatis subpungentibus 3-nerviis sericeo-pilosis, achænio brevi densè sericeo-piloso, pappi serie ext. internam subsequante. \mathcal{Z} in Brasiliâ tropicâ (Less.), in campis et paludosis prov. Minar. gener. (Mart.!) et prov. Sancti-Pauli (Lund.). Vern. erigerontis Mart. ! herb. Collum supernè tuberculosum villosum. Variat caulibus adpressè aut patentim pilosis, foliis margine planis aut revolutis uninerviis angustis aut penninerviis latioribus. Cor. ex sicco pallidæ glabræ. Pappus flavescens seu rufescens. (v. s.)

160. *V. CAMPESTRIS*, caule herbaceo erecto simplici confertè hispido, foliis sessilibus lineari-lanceolatis planis subintegerrimis utrinquè confertè villosis, capitulis 1-3 terminalibus, lateralibus breviter pedicellatis, terminali sessili, invol. squamis biserialibus villosis acuminatis, achænio hirtello, pappi serie ext. angustè palacèà brevi. — \mathcal{Z} in campis editis prov. Sancti-Pauli frequens, nov. flor. legit cl. Lund. Affinis *V. simplici*. Pubes confertissima in plantæ parte superiore, parca in inferiore. Cor. purpureæ glabræ. Pappus albus vix invol. longior. Folia $1\frac{1}{2}$ poll. longa, 3-3 $\frac{1}{2}$ lin. lata. (v. s. comm. à cl. Lund.)

161. *V. PSILOSTACHYA*, caule herbaceo erecto simplici villosa, foliis oblongo-lanceolatis utrinquè attenuatis integerrimis villosis, capitulis secus caulis apicem remotè spicatis subsessilibus ad axillas foliorum super. ortis et idèò cymam gracilem erectam simplicem constituentibus, invol. 2-3-serialis squamis dorso pubescentibus acuminatis, achænio hirtello, pappi serie ext. subpalacèà brevissimà acutà. \mathcal{Z} in campis editis prov. Sancti-Pauli sat. frequens, nov. fl. legit cl. Lund. Affinis *V. simplici* et *campestri* sed dispos. capitulorum faciliè diversa. Pappus albus, invo. vix superans. (v. s. comm. à cl. Lund.)

162. *V. DENSIVILLOSA* (Mart. ! herb.), caule herbaceo erecto simplici densè hirsuto tereti apice compresso, basi patentim folioso, apice ferè aphylo, foliis sessilibus obovatis basi cuneatis apice subobtusis acutis vix crenatis suprà adpressè sericeo-villosis, subtùs et margine densè sericeo-sublanatis, cymis corymbosis confertis aphyllis subscorpioides, capitulis 20-floris sessilibus, invol. squamis parcè villosis acuminatis, cor. glabrà, recep. nudo, achænio brevi villosa, pappi ser. ext. brevissimà palacèà acuminatà. — \mathcal{Z} in campis altis ad Serro Frio prov. Minar. general. legit cl. Martius, et in campis editis prov. Sancti-Pauli cl. Lund ! Folia 3-3 $\frac{1}{2}$ poll. longa, 18-20 lin. lata. Cor. violacæ. Pappus albus. Hirsuties tota sericeo-flavicans. (v. s. in h. Acad. reg. Monac.)

β . *angustior*, caule minùs hirsuto, foliis oblongis glanduloso-crenatis magis distantibus. \mathcal{Z} cum var. α . Folia sesquipollicem longa, 6. lin. lata, magis crenata. (v. s. in h. Acad. reg. Monac.)

163. *V. HOLOSERICA* (Mart. ! herb.), caule basi suffruticoso erecto simplici tereti pedunculisque villosis-velutinis, foliis sessilibus lanceolato-linearibus subacuminatis integerrimis utrinquè subtùs præsertim sericeo-villosis, paniculâ

terminalis ramis bi-trifidis aphyllis, capitulis pedicellatis 25-30-floris, invol. squamis glabris lanceolato-linearibus longè acuminatis, ext. brevioribus, cor. glabrâ, recept. subnudo, achænio densè sericeo-villoso, pappi ser. ext. angustâ paleaceâ acuminatâ brevi. ♀ in campis Sancti-Philippi versus Contendas prov. Minar. gener. Brasiliæ legit cl. Martius. Similis *V. schwenkiae* foliis, sed folia diversa 3 poll. longa, 4-lin. lata. Pappus uterque et achæniî pili crocei. (v. s. in h. Acad. reg. Monac.)

165. *V. SCHWENKIEFOLIA* (Mart. herb.) caule basi suffruticoso erecto simplici tereti densè pedunculisque velutino-villosis, foliis sessilibus ovatis basi obtusis apice mucrone calloso acutis utrinquè densè velutino-villosis subsericis penninerviis, paniculæ terminalis ramis lateralibus simplicibus aut bifidis aphyllis, capitulis pedicellatis 20-floris, invol. squamis glabriusculis longè acuminatis ext. brevioribus, cor. glabrâ, recept. nudo, achænio densè sericeo-villoso, pappi ser. ext. paleaceâ brevi. ♀ in Brasiliæ campis editis prov. Minar. gener. (Mart.), circa Tejuco (Vauth. pl. exs. n. 258). Pappus int. in sicco croceus, ext. albus. Folia 10-12 lin. longa, 5-6 lin. lata. Caulis sesquipedalis. (v. s.)

165. *V. TRIOLADA*, fruticosa, ramis teretibus hirsutis fuscâ densissimâ patulâ lanatis, foliis petiolatis elliptico-oblongis utrinquè acutis supra subasperis subrugulosis puberis, subtus reticulatis petiolisque fulvo-tomentosis, paniculæ terminalis ramis patentibus polycephalis bifidis subscorpioideis, capitulis sessilibus ovatis 10-12-floris, involucri squamis ovato-lanceolatis obtusiusculis aut mucronato-acuminatis, achænio villosio, pappi serie externâ setosâ, interior. multò brevior. ♀ in Brasiliæ prov. Sancti-Pauli. Pappus rufescens. Cor. glabrâ purpureâ. Affinis videtur *Vern. ferrugineâ*, sed inflorescentia diversa. Folia subtus non verè tomentosa, 2-2 $\frac{1}{2}$ poll. longa, pollicem lata. Inv. squamæ eciliatæ glabriusculæ. (v. s.)

166. *V. SESSILIFOLIA* (Less. in linnæâ 1831. p. 659), glaberrima, caule herbaceo erecto striato angulato apice nudiusculo, foliis lanceolato-linearibus elongatis sessilibus integerrimis penninerviis nervis utrinquè elevatis, subtus glanduloso-punctatis, capitulis circ. 40-floris longè pedunculatis aphyllis distanter racemosis, invol. campanulati squamis dorso pulverulento-puberulis et ciliatis lanceolatis acuminatis, cor. glabrâ, achænio teretiusculo sericeo-villosissimo, pappi ser. ext. paleaceâ acuminatâ achænio sextuplò ferè brevior. 2♀ in Brasiliâ. Simillima *V. grandiflorâ*, sed invol. et achænia diversa. Projecturæ 3 è foliis cujusque basi secus caulem decurrunt. (v. s. in h. Mus. reg. Paris acc. à Mus. imp. Bras. sub n. 441.)

167. *V. BREVIFOLIA* (Less. in linnæâ 1829. p. 285. 1831. p. 659), glabriuscula, caulibus imâ basi suffruticulosus erectis simplicibus aut à basi ramosis adpressè puberulis apice summo aphyllis sub-1-cephalis, foliis lineari-filiformibus sessilibus acutis sparse puberulis, invol. campanulati ebracteati squamis ovatis apiculato-cuspidatis extus adpressè villosis imbricatis. 2♀ in altis campis Brasiliæ (Sellow), inter Santo João d'Elkey et V^{na} Thitcam, prov. Minar. gener. legit cl. Martius. *V. filifolia* Mart. herb. Rhizoma lignosum. Folia 7-9 lin. longa. Herba in sicco nigricans. Cor. 15-20 purpureæ. Achænia turbinate sericeo-villosissima. Pappus albidus pluriserialis, setis ext. triplò ferè brevioribus paucioribus. An certè hujus sectionis? (v. s.)

§. 4. *Involucri squamis intimis obtusis, cymis scorpioideis, pedunculis ad basin folia perfecta gerentibus.* Less. in linnæâ 1831. p. 660. — *Species omnes Americanæ.*

168. *V. GRANDIFLORA* (Less. in linnæâ 1831. p. 660), glaberrima, caule herbaceo erecto simplici aut parcè ramoso teretiusculo, foliis linearibus elongatis utrinquè acuminatis sessilibus integerrimis penninerviis utrinquè nervoso-reticulatis, subtus glanduloso-punctatis, capitulis multifloris ad apicem caulis pedunculo-rumve-solitariis aphyllis, invol. campanulati squamis siccis imbricatis, ext. brevissimis acuminatis, mediis latè oblongis mucronatis, int. longioribus apice crosis obtusis, cor. glabrâ, achænio striato seriebus 10 pilosis notato, pappi ser.

ext. paleacea acuminatâ, achenio paulò brevior. $\frac{1}{2}$ in campis Brasiliæ (Sellow!), campis editis prov. Sancti Pauli (Lund!), ad Lorena prov. Sancti Pauli (Mart.). Vern. campestris Mart.! herb. Chrysocoma pumila Arrab. fl. flum. 8. t. 3a. Folia 5-6 poll. longa, 1-5 lin. lata. Capit. diam. pollicari. Cor. violaceæ. Achenium sesquilineare. Pappus albidus. Variat caule monocephalo aut racemoso, pedunculis longis distantibus 1-cephalis. (v. s.)

169. V. SELLOWII (Less. in linnæa 1829. p. 301. 1831. p. 661), caule tereti herbaceo simplici subaphyllo pubescente, foliis ad collum subsessilibus obovato-oblongis subacutis subserratis utrinquè subcabis penninerviis, pedunculis paucis ex axillis solitariis 1-cephalis, capitulis multifloris, invol. latè campanulati squamis oblongis acuminatis internis obtusis mucrone subpungentibus basi 3-nerviis, achenio oblongo multicostrato sericeo-piloso, pappi serie ext. paleacea brev. $\frac{1}{2}$ in Brasiliâ meridionali. Cor. albæ? glabræ. Pappus sordide flavescens. Variat pygmæa et bipedalis, involuero ferrugineo-villoso aut subglabro. (v. s. comm. à Mus. reg. Berol.)

170. V. OVATA (Less. in linnæa 1829. p. 294. 1831. p. 661), caule herbaceo tereti striatò simplici piloso, pilis sæpè reflexis, foliis brevè petiolatis membranaceis obovatis mucronulatis utrinquè pilosis, cymis scorpioides spiciformibus terminalibus, capitulis multifloris sessilibus, invol. campanulati squamis siccis 1-nerviis ciliatis, externis pungenti-acuminatis, internis obtusis, achenio piloso, pappi serie ext. brevi. $\frac{1}{2}$? in Brasiliâ ad Paraiba. Caulis pedalis. Cor. glabræ violaceæ. Pappus argenteus. Folia 15-18 lin. longa, 6-10 lata.

171. V. HYPOCHÆRIS, caule herbaceo è collo fruticoso erecto ferè aphylo tereti pubescente, foliis radicalibus patulis sessilibus obovatis obtusis margine subcrenatis penninerviis suprâ glabris subtus puberulis, caulinis oblongo-linearibus strictè adpressis crenulatis paucis, capitulis ad apicem geminis inferiore longius pedunculato ex axillâ folii supremi orto, invol. campanulati squamis ext. lanceolatis acuminatis pubescentibus, intimis oblongis subobtusis glabrisculis subcariosis, achenio sericeo-villoso, pappi ser. ext. paleacea brev. $\frac{1}{2}$ in Brasiliæ prov. Rio-Grande. Affinis V. Sellowii. Caulis pedalis. Fol. radie. 3 poll. longa, 2 poll. lata. Caulina pollicem longa, 1-2 lin. lata. Cor. purpureæ. Invol. et caulis purpurascens. (v. s. in h. Mus. reg. Paris. ex h. Mus. imp. Bras. n. 1025.)

172. V. GLABRATA (Less. in linnæa 1829. p. 294. 1831. p. 661), caule herbaceo simplici erecto supernè ramoso tereti striato subglabro, foliis brevissimè petiolatis rigidis lanceolatis elongatis basi subobtusis subdenticulatis reticulatis utrinquè glabris, cymis scorpioides terminalibus longis, capitulis sessilibus approximatis 20-25-floris, inv. campanulati squamis siccis 1-nerviis, exter. mucronato-acuminatis, intern. apice scariosis latè obtusis, achenio glabro, pappi serie ext. angustè paleacea brev. $\frac{1}{2}$ in Brasiliâ tropicâ. Caulis 2-4-pedalis. Cor. glabræ purpureæ. Pappus pallescens. (v. s. comm. à Mus. reg. Berol.)

173. V. BUDDLEIAEFOLIA (Mart.! herb.), fruticosa, ramis teretibus lanato-tomentosis, foliis brevissimè petiolatis elliptico-oblongis utrinquè subacuminatis crenatis, suprâ glabris venis impressis, subtus cano-tomentosis penninerviis, capitulis secus ramorum apicem sessilibus aut pedunculatis paucis extrâ-axillaribus ferè 60-floris, invol. subglobosi squamis coriaceis imbricatis adpressis obtusissimis dorso araneosis, ext. ovatis, int. oblongis, cor. glabrâ, achenio tereti villosò, pappi ser. ext. paleacea brev. $\frac{3}{5}$ in campis ad flum. Rio-Verde prov. Minarum general. Brasiliæ legit cl. Martins. Flores rosei. Folia 5-6 poll. longa, sesquipollicem lata, superiora, 9-10 lin. lata. (v. s. in h. Acad. reg. Monac.)

174. V. LIGULÆFOLIA (Mart.! herb.), fruticosa, ramis teretibus striatis glabrisculis, foliis sessilibus angustè ligulatis utrinquè subobtusis integerrimis suprâ glabris venoso-reticulatis, subtus cano-tomentosis penninerviis, capitulis ad apices ramorum spicatis in axillis foliorum sessilibus sæpius geminis folio triplò brevioribus 25-30-floris, invol. ovati squamis adpressis coriaceis glabris

- obtusis, ext. subtriangularibus brevibus, int. longis lato-linearibus, cor. glabrâ glandulosâ, achænio 10-costato basi sparse piloso apice glabro, pappi ser. ext. paleaceâ brevi. ζ in edijs campis prov. Minarum general. Cor. purpureæ. Folia 5-6 poll. longa, 9 lin. lata. (v. s. in h. Acad. reg. Monac.)
175. *V. CORIACEA* (Less. in linnæâ 1831. p. 661), caulibus è collo lignoso pluribus erectis teretibus simplicibus supernè albo-tomentosis, foliis brevissimè petiolatis coriaceis penninerviis suprâ nitidis, subtus tomentoso-incanis linearibus integerrimis, cymâ scorpioideâ spiciformi, capitulis multifloris folio flor. multò brevioribus, invol. semiglobosi squamis siccis vix subtomentosis ext. rotundis acutis, intim. linearibus obtusis, cor. glabrâ, achænio nitido sulcato, pappi ser. ext. paleaceâ brevi. — Patr. ignot. an Brasiliâ? Cor. purpureo-violaceâ. Capitula *V. macrocephalæ*. Folia 3-6 poll. longa, 4-6 lin. lata.
176. *V. RUFO-GRISÆA* (S. Hil. voy. diam. 2. p. 305 et 453), caule suffruticoso tereti, ramis supernè striato-angulatis rufo-griseove tomentosis, foliis petiolatis sublato-lanceolatis brevissimè mucronulatis basi acutis suprâ pubescenti-tomentosis subtus sericeis, cymis terminalibus bipartitis folio basilari brevioribus patentibus, capitulis secundis subsessilibus fol. flor. brevi brevioribus, invol. squamis pubescentibus acutis mucronulatis, achænio hirsuto villosa, pappi ser. ext. angustè paleaceâ brevi. ζ in Brasiliâ inter Almeida et Rio-Doce prope Quarta! do Riacho. Cor. dilutè purpureæ. Pappus subnrfescens.
177. *V. RUBRICAULIS* (H. et Bonpl. pl. equinox. 2. p. 66. t. 99), caule herbaceo sulcato-angulato pubescente simplici apice corymboso-paniculato, foliis sessilibus linearibus integerrimis margine revolutis suprâ scabris, subtus sericeo-tomentosis, capitulis secus ramos ad axillas sessilibus solitariis unilaterialibus 20-25-floris, invol. campanulati squamis imbricatis ovatis acutis dorso pubescentibus, achæniis glabris, pappi ser. ext. paleis lineari-subulatis brevibus. ζ in siccis montosis Novo-Granatensium prope Ybague H. B. et Kunth nov. gen. am. 4. p. 33. Less. in linnæâ 1831. p. 662. *V. rubricaulis* Bonplandiana, Less. in linnæâ 1829. p. 299, excl. var. γ et δ . Cor. purpureæ glabræ. Caulis ruber. — Eadem fortè ex Brasiliâ prov. Rio-Grande adest in h. Mus. reg. Par. ex Mus. imp. Bras. (ubi v. s.)
178. *V. OBTUSATA* (Less. in linnæâ 1831. p. 662), herbacea glaberrima erecta simplex, caule tereti striato, foliis petiolulatis rigidulis subintegerrimis oblongo-ovatis basi acuminatis lævibus penninerviis reticulatis, cymis scorpioideis, capitulis sessilibus ad folii valdè obtusati axillam sæpius geminis 10-floris, invol. cylindracei squamis siccis extus subpilosus nigro-ciliatis ovatis apiculatis, intimis obtusis, cor. glabrâ, achænio hirsuto, pappi ser. ext. paleaceâ brevi. — in Brasiliâ. Cor. purpureo-violaceâ. Pappus sordidus. Fol. 4 poll. longa, 15 lin. lata.
179. *V. SERICEA* (Rich. act. soc. h. nat. Par. 105), caule suffruticoso tereti striato tenuissimè puberulo, foliis brevissimè petiolatis lanceolatis acuminatis basi subobtusis subintegerrimis suprâ glabriusculis subtus adpressè pubescentibus, capitulis ad axillas sessilibus solitariis geminive circ. 50-floris, invol. campanulati squamis glabriusculis imbricatis, ext. acumine recurvo mucronatis, intimis longioribus obtusis, achænio sericeo-viloso, pappi ser. ext. angustè paleaceâ brevi. ζ in Brasiliâ, circa Rio-Janeiro (Gandich.! Lund.!), in monte Cercovado et in Serra de Estrella (Mart.!), et in Guiana Pers. ench. 2. p. 404. Ker bot. reg. t. 522. Less. in linnæâ 1829. p. 295. 1831. p. 663. Vern. imponens Mart. herb. Pappus uterque albus. Cor. glabriusculâ pallidè purpureæ aut ex Rich. albæ. Fol. 3 poll. longa, 9-10 lin. lata. Petioli sesquilineares! (v. s.)
180. *V. PERSICIFOLIA* (Desf. cat. h. Par. 1829. p. 400), caule fruticoso tereti striato flexuoso, foliis sessilibus lanceolatis longè acuminatis margine retrorsum asperis subintegerrimis deflexis nitidis, subtus subpubescentibus, capitulis axillaribus sessilibus solitariis circ. 30-floris, invol. teretis obovati squamis multiserialibus ovatis adpressis ciliatis, ext. brevibus mucrone rigido non adpresso recurvo superatis, intimis lineari-oblongis obtusis, pappi serie externâ paleaceâ brevi. ζ in Indiâ (verosim. occidentali). Cor. dilutè roseæ. Fol. 3-5 poll.

longa, 5-6 lin. basi lata. Pappus ext. fuscescens, int. setiformis albus. (v. s. in h. Desf.)

181. *V. ACUTIFOLIA* (Hook. bot. mag. t. 3062), caule herbaceo erecto glabro simplici tereti apice in ramos angulatos pubescentes diviso, foliis sessilibus linearibus acuminatis nervoso-reticulatis margine hinc inde subserratis suprâ glabris subtus obscure pilosis, capitulis 40-floris secus ramos subsessilibus lateralibus axillaribus, invol. hemisphærici squamis imbricatis glabrisculis, ext. brevibus acutis subsquarrosis, int. subscariosis obtusis, achænio pilos pluriseriales gerente, pappi ser. ext. brevi paleaceâ. $\frac{1}{2}$ in Brasiliâ ad Rio-Janeiro. — Differt à Vern. grandiflorâ, capitulis plurimis dimidio minoribus ebracteatis, foliis neo integerrimis nec omnino glabris; à *V. rubricauli*, capitulis 40 nec 20-25-floris, achænio pubigero, foliis subtus glabrisculis, etc.

β. ambigua, foliis suprâ junioribus scabridis demum glabris subtus glabrisculis punctatis, capitulis extrâ-axillaribus. $\frac{1}{2}$ in Brasiliâ. An Vern. rubricaulis latifolia Less. in linnæâ 1829. p. 299? (v. s. comm. sine nom. à Mns. reg. Berol.)

182. *V. SCHIEDANA* (Less. in linnæâ 1831. p. 399 et 663), fruticosa scandens, ramis teretibus pubescenti-canescensibus, foliis petiolulatis sublanceolato-ellipticis membranaceis penninerviis basi obtusis subintegerrimis suprâ glabris subtus pilosis, cymâ scorpioideâ divergente, capitulis sessilibus 25-floris folio stipatis, invol. campanulati squamis erectis ciliatis extus pilosis, ext. lanceolatis mucronatis, int. oblongis obtusis, cor. glabrâ, achænio sericeo-villoso, pappi ser. ext. paleaceâ brevi. $\frac{1}{2}$ in dumetis Mexici ad Papanila et Misantla. Folia 4-4 $\frac{1}{2}$ poll. longa, 1 $\frac{1}{2}$ -2 poll. lata; petiolo 3-4 lin. Affinis dicitur *V. sericea*.

183. *V. OBSCURA* (Less. in linnæâ 1829. p. 296. 1831. p. 663), caule fruticoso, ramis teretibus striatis fusco-villosis, foliis brevissimè petiolatis lanceolatis coriaceis subdenticulatis suprâ rugosis scaberrimis subtus flavescensibus villosotomentosis, cymis scorpioideis longis terminalibus, capitulis multifloris sessilibus approximatis folio florali multo brevioribus, invol. subcylindracei squamis r-nerviis ciliatis, intimis obtusissimis, achænio multicostato inter costas hirsuto, pappi serie ext. paleaceâ brevi. $\frac{1}{2}$ in Brasiliâ tropicâ. Cor. albâ? glabrâ. Pappus flavescens.

184. *V. AXILLIFLORA* (Mart. ! herb. et hort. Mon. ex Less. in linnæâ 1829. p. 297. 1831. p. 663), caule fruticoso, ramis teretibus villosis, foliis brevè petiolatis oblongo-ellipticis utrinquè acutis subintegerrimis suprâ scabris pilosis subtus villosotomentosis, cymis scorpioideis terminalibus longis, capitulis sessilibus folio florali multo brevioribus, invol. campanulati squamis r-nerviis obtusissimis subpungenti-mucronatis ciliatis, cor. lobis hirtis, achænio sericeo-piloso, pappi serie ext. brevi subpaleaceâ. $\frac{1}{2}$ in Brasiliâ ad Rio de S. Francisco prope Salgado prov. Minar. general. legit cl. Martius. Lodd. bot. cab. t. 1690. Cor. purpureo-violaceâ glabrâ. Pappus argenteus. (v. s. in h. Acad. reg. Monac.)

185. *V. COTONEASTER* (Less. in linnæâ 1829. p. 298. 1831. p. 663), caule basi suffruticuloso tereti striato apice laxè subtomentoso, foliis brevissimè petiolatis ovatis integerrimis suprâ scabrisculis demum glabris subtus villosotomentosis albis, floralibus ovato-cordatis, cymis longis simplicibus, capitulis ad axillas foliorum sessilibus solitariis rarius geminis, invol. squamis oblongis obtusis mucronatis laxè pilosis, achænio oblongo-villoso multicostato, pappi serie ext. paleis brevibus latioribus. $\frac{1}{2}$ ad Bahiam Brasiliæ. (Salzm. ! Blanch. !). *Conyza cotoneaster* Willd. in Spreng. syst. 3. p. 509. Invol. squamæ membran. sicce purpureæ. Cor. purpurea glabra. Pappus argenteus. (v. s. comm. à cl. Salzm. et Lhotsky.)

186. *V. ACUMINATA* (Less. in linnæâ 1831. p. 663), fruticosa, ramis divaricatis teretibus pubescentibus, foliis petiolulatis ovato-lanceolatis utrinquè acuminatis integerrimis scabris subtus punctatis, cymis scorpioideis brevibus polycephalis, capitulis 15-floris sessilibus approximatis folio flor. brevioribus, invol. squamis extus subpilosis, ext. minutissimis acuminatis, int. oblongis vix acutis, achænio multicostato piloso, pappi ser. ext. paleaceâ brevi. $\frac{1}{2}$ in insulis Cari-

- *hæsis?* Pappus flavescens. Recept. glabrum. Folia 1-2 $\frac{1}{2}$ poll. longa, 4-15 lin. lata. Petiolo 2 lin. Invol. in descr. nunc campanulatum nunc turbinatum dicitur.
187. *V. CHAMÆDRYS* (Less.! in linnæa 1829. p. 259. 1831. p. 641), caule fruticoso infernè glabro, ramis teretibus cano-tomentosis, foliis sessilibus oblongis subrotundisve margine subreflexo parcè serratis, suprâ glabris lucidis impressè glandulosis, subtus albo-tomentosis, cymæ ramis simplicibus ramosisve, capitulis sessilibus axillaribus et extrâ-axillaribus solitariis fasciculatisve cylindraceis 8-floris, invol. squamis 3-nerviis ovato-oblongis vix subtomentosis, achænio hiranto glanduloso, pappi serie externâ paleaceâ brevi. ζ in Brasiliâ tropicâ (Less.), ad Bahiam (Blanch.!). Cor. purpureo-violaceæ glabræ. (v. s.)
188. *V. AURIPODIARUM* (Mart.! herb.), suffruticosa?, caule erecto tereti striato pubescenti-villoso, foliis petiolatis elliptico-oblongis ntrinque acuminatis parcè dentatis, suprâ muriculatis secus nervos subvelutinis, subtus villosis-subhirsutis, capitulis ad axillas super. subsessilibus folio multò brevioribus solitariis 15-20-floris, invol. ovatis squamis adpressis ciliatis dorso glabrinaculis, ext. mucronatis, int. obtusis, corollâ glabrâ, achænio piloso striato, pappi ser. ext. brevi paleaceâ. ζ ? in campis prope Congonharde Sahara prov. Minar. gener. legit. cl. Martius. Aff. *V. lilacinea*. Petioli 8-9 lin. longi; fol. limbus 4-6 poll. longus, 18-20. lin. latus. Herba obscurè virens. (v. s. in h. Acad. reg. Monac.)
189. *V. LILACINA* (Mart.! herb.), fruticosa, ramis teretibus demùm glabris, ramulis subangulatis pubescenti-villosis, foliis breviter petiolatis ovalibus utrinquè subobtusis integerrimis suprâ tenuissimè pubescentibus, subtus pubescenti-villosis, ramis simplicibus bifidisve, capitulis ad dichotomias ram. aut. axillas foliorum sessilibus solitariis rarè geminis 10-12-floris, inv. ovatis squamis oblongis adpressis dorso apice puberis obtusis aut brevissimè mucronatis, achænio piloso turbinato, pappi ser. ext. brevi angustissimè paleaceâ. ζ in campis prope Tejucco prov. Minar. gener. Brasiliæ. Cor. lilacinæ apice parcè puberæ. Pappus rufescens vix invol. longior. Recept. alveolatum. Folia 2 poll. longa, 10 lin. lata, petiolo 1-2 lin. (v. s. in h. Acad. reg. Monac.)

§. 5. *Involucris squamis intimis acuminatis, cymis scorpioides folia perfecta ad basin pedunculorum gerentibus.* Less. in linnæa 1831. p. 664. — *Species omnes Americanæ.*

190. *V. ARBUSCULA* (Less. in linnæa 1831. p. 664), caule subarhoreo, ramis teretibus villo brevissimo incanis, foliis petiolulatis obovatis penninerviis suprâ pilosis, subtus villosis-incanis, cymis scorpioides, capitulis sessilibus foliis suis superatis, invol. turbinatis squamis acuminatis inermibus ciliatis dorso villosis, achænio villosis, pappi ser. ext. paleaceâ brevi. ζ in insulâ Mauritiî ex h. cl. Kunth; an fortè in horto bot. collecta? Affinis videtur *V. arborescenti*.
191. *V. ARBORESCENS* (Swartz fl. ind. occ. 2. p. 1320), fruticosa, ramis teretibus villosis, foliis brevè petiolatis oblongo-ovatis acuminatis sæpè deflexis subintegerrimis, suprâ pilosis subtus piloso-canescens, cymæ scorpioides terminalis ramis divaricatis, capitulis sessilibus axillaribus folio florali brevioribus, invol. campanulatis squamis acuminatis, achænio sericeo-piloso, pappi serie ext. brevi. ζ in montibus ins. Caribæarum Portorici, Sanctæ-Crucis. Less. in linnæa 1829. p. 302. 1831. 665. *Lepidaploa phyllostachya* Cass. dict. 26. p. 16. *Conyza arborescens* Linn. sp. 1209. — Plum. am. ed. Burm. t. 130. f. 2. Cor. glabrâ purpureo-cæruleæ nec flavæ ut vult Cass. Recept. subalveolatum. (v. s.)
- ρ . *ovatifolia*, foliis ovatis acuminatis, suprâ subrugosis, subtus pubescentibus paulò longius petiolatis. ζ in ins. Cubâ juxta Havanam et in Guadalupe. *Eupatorium secundiflorum* Bert.! herb. non Lag. An fortè *V. fruticosa* Lin.? sed folia acuminata nec obtusa. (v. s.)

192. *V. DIVARICATA* (Swartz fl. ind. occ. 3. p. 1319. non Less.), fruticosa,

ramis teretibus pubescentibus, foliis breviter petiolatis membranaceis ovali-oblongis utrinque acuminatis integerrimis supra et imprimis infra minutissimè pubescentibus, cymis axill. et terminalibus bifidis subscorpioideis, capitulis sessilibus nunc audis nunc bractea sua brevioribus 11-12-floris, invol. ovati squamis oblongis obtusis dorso parè puberis, achenio turbinato sericeo-villoso, pappi serie ext. brevi vix paleacea. ζ in ins. Caribæis. Conyza arboreascens Swartz! in h. L'Hér., sed antè fl. Ind. occ. script. V. acuminata Less. in linnæa 1831. p. 663? V. divaricata Spreng. syst. 3. p. 435 excl. syn. Cor. purpureæ glabræ. Pappus sordidè flavescenti-griseus. (v. s.)

193. V. ICOSANTHA, fruticosa, foliis petiolulatis oblongo-ovatis basi obtusis aut acutis membranaceis, subtus petiolisque villosis pellucido-glandulosis, cymis terminalibus scorpioideis oligocephalis, capitulis sessilibus alaribus aut extrà axillaribus folia sua superantibus 20-floris, invol. squamis oblongis adpressis mucronatis interioribus acuminatis, achenio turbinato densè piloso, pappi serie ext. paleacea brevi. ζ in Martinicâ et fortè in Brasiliâ. V. divaricata Less. in linnæa 1829. p. 306. 1831. p. 665. excl. syn. Pappus argenteus. Invol. à cl. Less. nunc cylindraceum nunc campanulatum dictum.

194. V. RIGIDA (Swartz! fl. Ind. occ. 1322), suffruticosa, ramis teretibus densè petiolisque pubescenti-velutinis, foliis petiolatis ovato-subrotundis utrinque obtusis integerrimis suprâ glabris sublucidis subtus glandulosis secus nervos velutinis, cymâ terminali subflexuosâ oligocephalâ, capitulis brevè pedunculatis folia floralia ovata subæquantibus, invol. squamis ext. carinatis subvillosis, intimis linearibus subacuminatis, achenio obconico hispidò, pappi serie latè paleacea brevi. ζ in montibus calcareis petrosis Jamaicæ borealis. Conyza rigida Sw. prod. 113. Cor. glabra purpurea. Pappus sordidè griseo-stramineus. An ipso suadente Swartzio à V. fruticosa satis differt? Folia rotundiora et cyma multò magis contracta. (v. s.)

195. V. VARIANA (Less. l. c. p. 306. 1831. p. 306), fruticosa, ramis teretibus striatis pube adpressâ sericeâ albidis, foliis brevè petiolatis membranaceis integerrimis ovato-oblongis utrinque minutè glandulosis et pubescentibus demum glabris, cymis scorpioideis brevibus oligocephalis, capitulis subsessilibus approximatis, invol. cylindracei squamis acuminatis 1-nerviis demum patentibus, achenio oblongo densè sericeo-piloso, pappi serie ext. paleacea brevi. ζ in insulâ Sanctæ-Crucis. Conyza glabra Willd. sp. 3. p. 1940. et Eupatorium obtusifolium Willd. sp. 3. p. 1768. ex Less. V. albicaulis Pers. syn. 2. p. 404. Cor. glabræ violaceæ. Pappus argenteus. (v. s. comm. à cl. Puerari, qui à cl. West receperat.)

196. V. LONGIFOLIA (Pers. ench. 2. p. 404), caule suffruticoso? tereti petiolis pedunculisque pubescentibus, foliis brevè petiolatis oblongo-lanceolatis basi brevè apice longè acuminatis membranaceis glabris integerrimis utrinque minutè glandulosis, cymis axillaribus et terminalibus folio brevioribus bifidis oligocephalis, capitulis folio oblongo-lineari brevioribus brevè pedicellatis, invol. turbinati squamis acuminatis, achenio sericeo-villoso, pappi serie ext. paleacea brevi. ζ in Guadalupâ. V. punctata Swartz in Wikstr. gnad. fl. 72. an ex ipso auctore Lepidaploa lanceolata Cass. dict. 26. p. 18, sed cor. certè in nostrâ non sunt flavæ! Folia 4-5 poll. longa, 9-14 lin. lata. Pappus argenteus. Cor. glabræ ex sicco purpureæ. (v. s. comm. à cl. Krauss.)

197. V. OBTUSIFOLIA (Less. l. c. p. 308), fruticosa, ramis teretiusculis glabris, junioribus pilosiusculis, foliis petiolatis carnosò-coriaceis ellipticis obtusis basi acutis integerrimis suprâ impresso-glandulosis utrinque pilosiusculis, cymis scorpioideis brevibus terminalibus oligocephalis, capitulis sessilibus multifloris subapproximatis folii floralis longitudine, invol. campanulati squamis oblongo-ovatis subacuminatis 3-nerviis, achenio sericeo-piloso, pappi serie ext. paleacea brevi. ζ in Brasiliâ tropicâ. Recept. paleaceo-fimbrilliferum. Pappus argenteus. Fol. 1-1 $\frac{1}{2}$ poll. longa, 4-6 lin. lata.

198. V. GRACILIS (H. B. et Kunth nov. gen. am. 4. p. 27), herbacea, ramis teretibus glabris aut subtomentosis, foliis brevissimè petiolatis oblonga acu-

minatis subintegerrimis puberulo-scabriusculis sæpè reflexis, cymis scorpioideis terminalibus longis divaricatis, capitulis sessilibus remotis in axillis foliorum solitariis conglomeratisve, invol. subturbinati squamis acuminatis pilosis demùm patentibus, achenio inter costas piloso, pappi ser. ext. brevi. $\frac{1}{2}$ in umbrosis prope Turbaco Novo-Granatensium et in Brasiliâ tropicâ. Less. in linnæâ 1829. p. 302. 1831. p. 667. V. elongata Willd. herb. ex Less. Cor. violaceæ glabræ.

ß. *villosa* (Less. l. c.) foliis subtus subtomentoso-villosis, capitulis magis conglomeratis.

199. V. STENOPHYLLA (Less. in linnæâ 1831. p. 667), caule è rhizomate lignoso erecto herbaceo tereti supernè albo-villosissimo simplici, foliis petiolulatis angustè linearibus aveniis acutis margine revolutis suprà pilosis, subtus sericeo-villosis, capitulis secus ramos paniculæ in axillis foliorum sessilibus et iis brevioribus, invol. campanulati squamis linearibus mucronatis extus villosis, cor. glabrâ, achenio villoso, pappi ser. ext. paleaceâ brevi. $\frac{1}{2}$ in Sancto-Domingo. Folia 1 $\frac{1}{2}$ - 2 $\frac{1}{2}$ poll. longa, lineam lata.
200. V. MACROPHYLLA (Less. in linnæâ 1831. p. 668), caule herbaceo altissimo tereti striato villoso-pubescente apice ramoso et foliato, foliis petiolatis membranaceis basi obtusissimis inf. ovatis acuminatis cæt. lanceolato-ellipticis penninerviis suprà hispidiusculis subtus molliter pubescentibus et punctatis, cymis scorpioideis divaricatis, capitulis sessilibus remotis solitariis plerisque folio suffultis, invol. latè campanulati squamis multinerviis, ext. ovatis acutis, med. obtusis, intimis acuminatis, cor. glabrâ, achenio villoso, pappi ser. ext. paleaceâ brevi. $\frac{1}{2}$ in Brasiliæ prov. Rio-Janeiro in Serra do Mar. Folia 17-18 poll. longa, 9-11 lata. Herba dendromorpha 20-pedalis. Cor. violaceæ. Pappus sordidè flavescens.
201. V. MULTIFLOSCULOSA (Mart. herb.) suffruticosa, ramis teretibus pedunculisque pube brevissimâ velutinis, foliis breviter petiolatis ovatis basi subcuneatis apice acuminatis callosò-serratis suprà glabriusculis subtus nascenti-subtomentosis, paniculæ ramis inf. axillaribus folio æqualibus, sup. brevioribus subscorpioideis, capitulis maximis 50-floris subsessilibus aliis ebracteatis aliis bractea oblongâ donatis, invol. squamis ovato-oblongis mucronatis adpressis ciliatis, intimis acuminatis, achenio villoso-sericeo, pappi ser. ext. brevi paleaceâ. $\frac{1}{2}$ $\frac{3}{4}$ in Brasiliæ umbrosis et sylvis cædvis prope Rio-Janeiro (Blanch.), ad montem Cercovado (Mart.), in siccis prope Rge. (Lhostky!), Chrysocoma pedunculata Arrab. fl. flum. 8. t. 37. Affinis V. macrocephalæ. Invol. squamæ dorso vix pubescentes. Pappus sordidè rufescens. Folia 8 poll. longa, 4-5 poll. lata. Variat foliis subtus subalbidis aut virescenti-pubescentibus. An eadem ac V. macrophylla? (v. s.)
202. V. HELOPHILA (Mart. herb.), caulibus basi suffruticulosus erectis simplicibus teretibus villoso-hirsutis apice glabriusculis, foliis brevissimè petiolatis ovatis basi obtusis apice acutis mucrone callosò integerrimis suprà sparsè pilosis demùm scabridis subtus glandulosus villosis, ramis axillaribus distantibus simplicibus bifidisve, capitulis sessilibus ad axillam folii ovato-lanceolati distantibus solitariis 20-floris, invol. squamis omnibus acuminatis, ext. brevioribus dorso pubescentibus, cor. glabris, achenio villoso, pappi ser. ext. paleaceâ brevi. 5 in paludosis montanis prov. Minarum General. prope Villa-Ricca leg. cl. Martins. Caules rubentes. Cor. albæ. Folia poll. longa, 6 lin. lata, petiolo lineam longo. (v. s.)
203. V. GEMINATA (Less. in linnæâ 1829. p. 303. excl. syn. ex linnæâ 1831. p. 669), herbacea volubilis, ramis teretibus obscurè villosis, foliis brevè petiolatis oblongo-ovatis acutis undulato-subcrenulatis suprà scabris rugosis subpilosis subtus obscurè villosis, cymæ scorpioideæ terminalis ramis longis, summis geminis, capitulis sæpius solitariis 20-floris approximatis sessilibus folia floralia subtus cordata ferè superantibus, invol. campanulati squamis in acumen setiforme patulum productis intimis linearibus rectis, achenio sericeo-piloso, pappi-serie ext. brevè paleaceâ. in Brasiliæ prov. Rio-Janeiro et in Serra d'Es-

trella. Recept. alveolare. Cor. albæ glabræ. Pappus albus. Spreng. syst. 3. p. 437. excl. syn. (v. s. comm. à Mus. reg. Berol.)

β. *aprica*, caule erecto ad apicem magis villosio, foliis oblongis magis acuminatis subtus densius et mollius villosis, cymæ ramis brevioribus, invol. squamis pallidis dorso omnibus molliter villosis. ½ in locis apricis vulgarissima circa Rio-Janeiro ex cl. Lund. Herba 1-2-pedalis. Cor. albæ. Capitula paulò minora quàm var. α. An species propria? (v. s. comm. à cl. Lund.)

204. V. *BARDANOIDES* (Less. in linnæa 1831. p. 669), caule herbaceo-erecto tereti striato apice hirsuto-viloso, foliis subsessilibus subrotundis ellipticisve basi subcordatis macronato-acuminatis reticulato-venosis asperimis suprà lucidis subtus pilosis, cymæ scorpioideæ pauciradiatâ polycephalâ, capitulis sessilibus foliis binis stipatis, invol. campanulati squamis acuminatis rigidis extus pilosis, corollâ glabrâ, achenio. . . . pappi ser. ext. paleaceâ brevi. — in Brasiliâ. Folia 2-3 poll. longa, ½-2 lata. Cor. purpureo-violaceæ. Pappus argenteus. Affinis dicitur V. geminata.

205. V. *CRAMISSONIS* (Less. in linn. 1829. p. 304. 1831. p. 670), herbacea, ramis teretibus albo-viloso-tomentosis, foliis breviter petiolatis rigidis ovato-lanceolatis acuminatis basi acutis integerrimis suprà rugosis pilosis subtus sericeo-incanis, cymis scorpioideis terminalibus longis, capitulis distantibus sessilibus folio florali sublongioribus multifloris, invol. campanulati squamis siccis 1-nerviis ciliatis pilosis longè acuminatis ext. subpatentibus subpungentibus, cor. glabrâ, achenio densè piloso, pappi seriebus inæqualibus. ½ in Brasiliâ ad fretum Sanctæ-Catharinæ. Cor. purpureo-violaceæ. Folia 2-3 poll. longa, 9-14 lin. lata.

206. V. *TOMENTOSA* (Nees amœn. 2. p. 11. t. 3. non Ell.), caule basi frutescente simpliciter virgato supernè villosio-tomentoso, foliis oblongo-lanceolatis brevè petiolatis integerrimis utrinquè molliter tomentoso-villosis, capitulis 12-floris corymboso-paniculatis, cymis subscorpioideis, invol. squamis lineari-lanceolatis acutis, interioribus apice coloratis, achenio obconico villosio, pappi serie ext. brevissimâ paleaceâ. ½ in Indiâ or. ex sem. hort. Less. in linn. 1829. p. 305. 1831. p. 670. Cor. glabræ purpurascens, in invol. teretiusculo. Pappus argenteus.

207. V. *MEGAPOTAMICA* (Spreng. syst. 3. p. 437), caule herbaceo simpliciter striato tereti glabriusculo, foliis sessilibus suprà glabris nitidis subtus albo-tomentosis oblongis inferioribus basi attenuatis superioribus basi dilatatis summis cordato-lanceolatis subangulatis, cymis scorpioideis in paniculam dispositis, capitulis cylindricis 6-floris, invol. foliolis ovato-oblongis mucronato-acutis erectis albidis-sericeo-pilosis, achenio obconico multicostato piloso, pappi serie externâ paleaceâ brevi. ½ in Brasiliâ meridionali ad Rio Grande. Less. in linnæa 1829. p. 308. 1831. p. 670. Cor. violaceæ glabræ. Pappus argenteus. Rhizoma repens. (v. s. comm. à Mus. reg. Berol.)

β. *brevifolia*, caule ad apicem foliis subtus et involucriis albo-lanatis, foliis ellipticis semi-pollicem circ. longis. — Cum var. α.

γ. *melanotrichia*, caule lanâ nigricante hirsuto, fol. nervo medio villis fuscis insignito, invol. squamis ciliis nigris hispidis, pappi interioris setis nigris. ½ in Brasiliæ prov. Sancti-Pauli in campis ad Lorena. Vern. Candolliana Mart. herb. Var. α. simillima, sed colore nigro pilorum ferè omnium etiam pappi inter. distincta. (v. s. in h. Acad. reg. Monac.)

208. V. *APICULATA* (Mart. herb.), caule herbaceo erecto tereti parçè ramoso pubescente subhirsutulo, foliis sessilibus erectis lanceolatis margine revolutis basi cordato-obtusis apice apiculatis suprà pilosis demum scabris subtus tomentosis, capitulis secus ramos sessilibus spicatis axillaribus folio florali paulò longioribus 4-floris, invol. cylindrici squamis dorso pubescentibus apice longè acuminatis, achenio adpressè pubero, pappi serie ext. setiformi paucisetâ int. triplò brevior. ☉ in campis Brasiliæ prope Barreiros Minarum Novarum legit cl. Martins. Herba sesquipedalis. Pili paginæ fol. sup. decidui basi tuberculo-formes. Accedere videtur ad V. Megapotamicam. (v. s. h. Acad. reg. Monac.)

209. V. *HEMOTIFLORA* (Rich. aest. soc. h. n. Par. 1792. p. 117), herbacea, caule

tereti adpressè pubescente, foliis brevissimè petiolatis ovali-lanceolatis membranaceis serratis basi cuneatis apice acutis suprâ pubescentibus, subtus sparè villosis, cymæ subpaniculatæ ramis subscorpioideis hinc inde foliiferis, capitulis 12-15-floris sessilibus distantibus, invol. ovati squamis sublinearibus acuminatis fructus superantibus extus villosis, achenio sericeo-villoso, pappi serie ext. paleaceâ brevissimâ. ☉ in Cayennâ et Brasiliâ. Less. in linnæâ 1831. p. 670. Pers. ench. 2. p. 404. V. sessiliflora Willd. herb. ex Less. in linnæâ 1829. p. 309. Habitus ferè V. canescens sed pubescens nec canescens, capitula angustiora longiora hinc inde è foliorum axillis orta. Pappus argenteus. (v. s.)

210. V. BERTERIANA, caule herbaceo erecto ramoso tereti villosa, foliis sessilibus oblongis utrinquè subattenuatis apice macronatis integerrimis suprâ setis brevissimis scabris, subtus villis adpressis subsericeis canescentibus, ramis scorpioideis, capitulis ad axillas foliorum sessilibus solitariis folio brevioribus 10-floris, invol. squamis lineari-lanceolatis acuminatis glabrisculis subnitidis fructus æquantibus, cor. glabrâ, achenio sericeo-subvillosa, pappi ser. ext. paleaceâ brevissimâ. ☉ in insulis Porto-Ricci et Sancti-Thomasii locis aridis legit cl. Bertero. Conyza Portoricensis Bert. herb. Pappo rufescente involucrum æquante et paginâ fol. breviter scabrâ differt à V. remotiflorâ. An varietas? (v. s.)

211. V. VERBASCIFOLIA (Less. in linnæâ 1829. p. 310. 1831. p. 670), caule herbaceo simplici tereti sulcato tomentoso-villoso, foliis sessilibus confertis rigidis utrinquè scabris et villosis obovatis subintegerrimis, cymis scorpioideis terminalibus subfastigiatis oligocephalis, capitulis multifloris sessilibus remotis folio florali ovato longioribus, invol. campanulati pauciserialis squamis acuminatis 3-nerviis, cor. glabrâ, achenio sericeo-villoso, pappi ser. ext. paleaceâ interiori parum brevior. 2/ in Brasiliâ tropicâ. Cor violaceâ. Pappus argenteus. Folia 2-5 poll. longa, 1-2 lata.

212. V. FLEXUOSA (Sims bot. mag. t. 2477), caulibus herbaceis è rhizomate lignoso teretiusculis striatis albo-pilosis, foliis sessilibus oblongis seu lineari-lanceolatis subintegerrimis utrinquè scabro-pilosis, cymis scorpioideis paucis, capitulis bracteâ oblongâ longioribus secus cymæ ramos ad dichotomias sessilibus multifloris, invol. campanulati foliolis 1-nerviis oblongis mucronato-acuminatis extus subvillosis, achenio turbinato undiquè villosa, pappi serie externâ paleaceâ acuminatâ interiore dimidio ferè brevior. 2/ in Brasiliâ meridionali. Lessing in linnæâ 1829. p. 311. 1831. p. 670. Lodd. bot. cat. t. 1880. Variat cor. purpureis et albis. Pappus argenteus valdè inequalis aut subæqualis. Rhizoma tuberosum. (v. s. comm. à Mus. reg. Berol.)

213. V. COGNATA (Less. in linnæâ 1831. p. 670), caule erecto striato villosa, foliis sessilibus penninerviis basi obtusiusculis, inf. obovatis acutis, sup. linearibus elongatis suprâ scabris hirsutiusculis subtus villosa-sericeis, cymæ ramis elongatis erecto-patentibus villosissimis polycephalis, capitulis 10-12-floris remotis folia sua superantibus sessilibus, cor. glabrâ, achenio piloso, pappi ser. ext. paleaceâ brevi. — in Brasiliâ. Affinis dicitur V. Platensis et V. obovatæ. Caulis totus foliatus. Villi in sicco flavicantes. Fol. infer. 4 poll. longa, sesquipollicem lata.

214. V. PLATENSIS (Less. in linnæâ 1829. p. 312. 1831. p. 671), caule herbaceo tereti striato apice subhirsuto, foliis sessilibus confertis elongato-lanceolatis integerrimis aut undulato-crenatis suprâ rugosis pubescentibus subtus villosa-canescantibus, cymis subscorpioideis axillaribus compositis in paniculam ferè dispositis, capitulis multifloris, invol. campanulati foliolis 1-nerviis oblongis mucronatis extus villosis, achenio piloso, pappi serie externâ brevissimâ. 2/ in Brasiliâ meridionali ad Rio de la Plata (Spreng.) et Rio-Grande (h. Mus. imp. Bras. n. 707). Conyza Platensis Spreng. 3. p. 509. Cor. purpureâ glabrâ. Flores in invol. 35-40. Pappus argenteus. (v. s.)

215. V. RADULA (Mart. herb.), caule herbaceo basi suffruticuloso erecto subangulato glabrisculo, foliis sessilibus oblongo-linearibus acutiusculis coriaceis utrinquè glabris asperulis venis elevatis et-reticulatis subintegerrimis, capitulis

ad apicem canlis ad axillas aut extrâ axillas folio flor. multò brevioribus sessilibus aut breviter pedicellatis 70-floris, invol. subglobosi squamis pluriserialibus ext. lanceolatis longè acuminatis semipatulis subaraneosis, intimis linearibus membranaceis subacuminatis, cor. glabrâ, achænio glabriusculo, pappi ser. ext. paleaceâ brevi, int. setis apice subclavellatis. $\frac{5}{2}$ in editis prope Sanctam Barbaram ad fl. Sabucahy prov. Minarum General. legit cl. Martius. Fol. 3-4 poll. longa, 4-6 lin. lata. Cor. purpureæ. Rhizoma lignosum. (v. s. in h. Acad. reg. Monac.)

β. latifolia, caule apice paulò magis villosiusculo, foliis oblongis vix acutis subtus puberulis perspicuè denticulatis, achænio longiore magis piloso. $\frac{5}{2}$ in Brasiliâ legit cl. Martius et cum var. α in herb. conjunxit: Folia 2-3 poll. longa, 9-12 lin. lata. An potiùs species propria? (v. s. in h. Acad. reg. Monac.)

216. *V. HECATANTRA*, caule suffruticuloso erecto ramoso tereti glabriusculo apice araneoso, foliis sessilibus linearibus utrinquè subattenuatis rigidis integerrimis suprâ glabris subtus subtomentoso-canescens, capitulis subcorymbosis terminalibus et panicis lateralibus folio florali vix brevioribus ovato-globosis 100-110-floris, invol. squamis multiserialibus adpressis, ext. latè ovatis vix submucronatis, inter. linearibus oblongis obtusis, cor. glabrâ glandulosâ, achænio 10-costato glabro, pappi ser. ext. paleaceâ brevi. $\frac{5}{2}$ patr. ign. Folia et invol. *V. ligulæfoliæ*, sed folia angustiora, ferè *V. radulæ* var. angustifoliæ similia, dispos. capit. diversa, flor. numerus multò major, achænia glabra, etc. (v. s. in h. Acad. reg. Monac.)

217. *V. ENSIFOLIA* (Mart. herb.), caule suffruticuloso tereti ramoso apice subvillosa, foliis sessilibus linearibus acutis subvillosis suprâ subasperis subtus penninerviis subreticulatis margine subintegerrimis, capitulis secus ramos elongatis sessilibus axillaribus aut extraaxillaribus folio brevioribus circ. 35-40-floris, invol. ovatis squamis pluriserialibus, ext. lanceolatis acuminatis subvillosis semipatulis, int. linearibus obtusis mucronatis, achænio glabro, pappi ser. ext. $\frac{5}{2}$ in subsylvaticis montanis prov. Minarum General. legit cl. Martius. Folia 5 poll. longa, 5 lin. lata. Affinis *V. radulæ* var. α . Flores in capit. junioris! (v. s. in h. Acad. reg. Monac.)

218. *V. SPIXIANA* (Mart. herb.), caule herbaceo ramoso erecto striato apice adpressè puberulo demum glabro, foliis subsessilibus linearibus utrinquè attenuatis integerrimis 1-nerviis suprâ et margine scabris subtus subhirsutis, paniculæ ramis bifidis folia ovali-lanceolata subsericea gerentibus subscorpioideis oligocephalis, capitulis confertis subsessilibus fol. flor. longioribus 7-floris, invol. oblongi squamis lanceolatis, exter. dorso villosis mucronatis, int. longioribus subglabris acuminatis, cor. subpilosâ, achænio sericeo-villoso, pappi ser. ext. paleaceâ brevi. $\frac{2}{2}$ in deserto Minarum Gener. Brasiliæ legit cl. Martius. Folia pollicem longa, 2 lin. lata. Pappus albens. Cor. pallidæ ex sicco. Invol. squamæ subrubentes. (v. s. in h. Acad. reg. Monac.)

219. *V. FRUTICULOSA* (Mart. herb.), caule fruticuloso ramoso tereti erecto apice pulveruleato-pubescente, foliis oblongis sessilibus utrinquè subacutis integerrimis aut panicis utrinquè minutè et adpressè pubescentibus cinereis 1-nerviis, cymâ scorpioideâ subaphyllâ patulâ, capitulis unilateralibus sessilibus oblongis 8-floris, invol. squamis oblongis, exter. submucronatis adpressis intimis subacuminatis, cor. hirtellâ, achænio sericeo-villoso, pappi ser. ext. angustè paleaceâ acuminatâ ferè triplò inter. brevior. $\frac{5}{2}$ in deserto ad flum. Sancti-Francisci prov. Minarum General. Brasiliæ legit cl. Martius. Folia 7-8 lin. longa, 2-3 lata. Capit. 3 lin. longum. (v. s. in h. Acad. reg. Monac.)

β? penninerviis, ramulis striatis, foliis duplò majoribus integerrimis subtus penninerviis, capitulis magis approximatis paulò majoribus 8-9-floris, invol. squamis mucrone subpatente brevi terminatis, pappi ser. ext. paleaceâ quintam seriei internæ longitudinem vix superante. $\frac{5}{2}$ cum priore juncta in herb. Mart. sed verisim. distinguenda. (v. s. in h. Acad. reg. Monac.)

220. *V. SENECIONIA* (Mart. ! herb.), caule herbaceo erecto tereti hirsuto, foliis sessilibus linearilanceolatis basi obtusis apice cuspidate acuminatis subintegerrimis penninerviis supra sparse pilosis subtus hirsutis, paniculae laxae ramis scorpioideis elongatis, capitulis ad axillas foliorum geminis sessilibus folio longioribus 10-floris, invol. squamis lanceolatis acuminatis dorso subhirsutis, cor. glabrâ, achænio puberulo parvisimè glanduloso, pappi ser. ext. paleaceâ brevi. 2 in campis editis prov. Sancti-Pauli Brasiliæ leg. cl. Martius et Lund. Herba bipedalis. Folia 2 poll. longa, 4-6 lin. lata. Pappus albidus. Cor. purpureæ. (v. s.)
- β. *adenocarpa*, foliis brevius cuspidatis, invol. squamis dorso magis hirsutis, foliis pube magis sericeâ hirsutis, achænio copiosissimè glanduloso. 2 in prov. Sancti-Pauli Bras. legit cl. Martins et cum var. α in herb. coadunavit. (v. s. in h. Acad. reg. Monac.)
221. *V. ARENARIA* (Mart. ! herb.), fruticosa tota tomento brevi cinatæscitovelutina, ramis teretibus, foliis petiolatis ellipticis basi acutis apice obtusis integerrimis penninerviis, nervis subtus prominulis, capitulis axillaribus solitariis sessilibus 30-40-floris, invol. hemisphærici squamis imbricatis acuminatis dorso puberulis, ext. acumine subrecurvo, int. rectis, achænio sericeo-villosissimo, pappi serie ext. paleaceâ brevi. 5 in rupibus arenariis prov. Piahiensis Brasiliæ legit cl. Martius. Cor. purpureo-violaceæ glandulosæ. Pappus nitens albidus. Fol. 1-2 poll. longa, 9-12 lin. lata, petiolo 2-3 lin. (v. s. in h. Acad. reg. Monac.)
222. *V. DEBILIS* (Mart. ! herb.), caule fruticoso ramosissimo, ramis teretibus debilibus apice velutino-tomentosis, foliis oblongo-lanceolatis vix petiolulatis utrinquè acuminatis integerrimis supra pubescenti-velutinis subtus cano-tomentosis, ramis flor. scorpioideis, capitulis ad axillas foliorum flor. solitariis geminisve sessilibus folio multò brevioribus 20-25-floris; invol. squamis oblongis dorso subvelutinis mucronato-acuminatis adpressis, corollâ extus hirtellâ, achænio tereti puberulo, pappi ser. ext. paleaceâ brevi. 5 in apricis aridis deserti versus flum. Sancti-Francisci prov. Minar. General. Brasiliæ legit cl. Martius. Folia 15-18 lin. longa, 4 lin. lata. Pappus argenteus. Cor. purpurascens. (v. s. in h. Acad. reg. Monac.)
223. *V. EREMOPHILA* (Mart. ! herb.), suffruticulosa tota pube brevi adpressâ subcanescens, caule erecto tereti ramoso, foliis elongatis lanceolato-linearibus utrinquè attenuatis integerrimis penninerviis, ramis scorpioideis oligocephalis sæpe bifidis, capitulis in dichotomiis ramorum et ad axillas fol. ram. sessilibus solitariis 10-floris, invol. ovatis squamis lanceolatis subacuminatis adpressis, cor. glabrâ glandulosâ, achænio turbinato sericeo-villoso, pappi ser. ext. paleaceâ brevi acuminatâ. 5 in campis arenosis versus Goyas prov. Minar. Gener. legit cl. Martius. Cor. albæ. Anthæræ roseæ. Pappus sordidè albescent. Folia 3-4 poll. longa, 3-4 lin. lata. (v. s. in h. Acad. reg. Monac.)
224. *V. CHALYBÆA* (Mart. ! herb.), suffruticulosa ? caule erecto tereti puberulo, foliis oblongo-linearibus basi acutis apice obtusis integerrimis subundulatis 1-nerviis utrinquè puberulis, ramis elongatis subscorpioideis, capitulis ad axillas foliorum sessilibus solitariis (raro geminis) 20-floris, invol. squamis omnibus acuminatissimis glabris nitidis subcoloratis, cor. glabrâ glandulosâ, achænio sericeo-villoso, pappi ser. ext. paleaceâ brevi. 5 in siccis prov. Bahiensis interioris legit cl. Martius. Herba ferè *Erigerontis canadensis*. Folia 6-9 lin. longa, sesquilineam lata. Pappus pallidus nitens. (v. s. in h. Acad. reg. Monac.)
225. *V. TRICHOLEPIS*, caule herbaceo erecto ramoso tereti glabro, ramulis subtiliter striatis et puberulis, foliis sessilibus membranaceis ovato-lanceolatis basi attenuatis apice acuminatis subintegerrimis, supra glabriusculis, subtus piloso-subhirsutis, capitulis secus ramos ad axillas foliorum sessilibus solitariis 15-floris folio flor. triplò brevioribus, invol. ovali-oblongi squamis glabris, ext. capillatis semipatulis, intimis lanceolato-linearibus acuminatis pappo et corollis longioribus, achænio teretiusculo adpressè piloso, pappi ser. ext. brevi angustè

paleaceâ. \mathcal{L} ? in siccis insulæ Cayennæ (Leprieur!) et in montosis prope Rio de Contas prov. Bahiensis Brasilæ! (Martius). Vern. patula Mart. ! herb. Folia 3 poll. longa, pollicem lata. Pappus griseus invol. brevior. (v. s.)

226. V. SALZMANNI, caule basi suffruticoso ramoso tereti pubescente, ramis elongatis, foliis lineari-lanceolatis utrinquè acuminatis vix petiolatis integerrimis membranaceis utrinquè subpubescentibus, capitulis 20-floris, secus ramos ad axillas foliorum sessilibus solitariis geminive ovatis, involucri squamis lineari-oblongis subacuminatis subvillosis, achænio piloso oblongo tereti, pappi serie externâ angustâ paleaceâ brevissimâ. \mathfrak{z} in umbrosis circa Bahiam Bras. semel legit cl. Salzmann. Caulis 4-pedalis. Folia 2-2 $\frac{1}{2}$ poll. longa, 5-6 lin. lata. Invol. rufescentia. Pappus pallidè rufescens. Cor. purpureæ, lobis apice subpuberis. (v. s.)

227. V. MUCRONIFOLIA, caule basi suffruticuloso tereti hirsuto-pubescente, foliis brevissimè petiolatis ovatis ovalibusve basi obtusis apice mucrone acuto apiculatis utrinquè minutissimè pubescentibus, cymæ ramis elongatis, capitulis ad axillas fol. sessilibus solitariis 10-12-floris, invol. squamis pilosiusculis acuminatis, achænio sericeo-villoso, pappi paleis externis brevibus oblongis paleaceis. \mathfrak{z} in sabulosis maritimis ad Bahiam Brasilæ leg. cl. Salzmann et Blanchet. Herba 4-pedalis. Aff. V. cotoneastri. Folia 15-18 lin. longa, 7-9 lata. Cor. purpureæ glabræ. Pappus argenteus. (v. s.)

228. V. POEPPIGIANA, caule herbaceo? tereti ramoso villosa, foliis sessilibus oblongo-lanceolatis utrinquè attenuatis subintegerrimis suprà sparsè pilosis subtus villosa-hirsutis, ramis scorpioides, capitulis ad axillas folior. sessilibus aut subgeminis 35-floris folio triplò brevioribus, invol. squamis lineari-lanceolatis omnibus acuminatis, ext. submucronatis dorso adpressè pubentibus, cor. glabrâ, achænio striato nervulis obtusis lævibus valleculis seriatis pilosis, pappi ser. ext. vix paleaceâ brevi. — in Americâ calidiore verisim. in Peruvia legit cl. Poeppig. (pl. exs. n. 1204). Affinis V. imponenti et salicifoliz: Cor. purpureæ. Folia 3-4 poll. longa, 5-6 lin. lata. Pappus argenteus. An satis ab obscurâ diversa? (v. s. comm. à cl. inv.)

229. V. MURICATA, herbacea?, caule tereti ramoso striato subglabrato, ramis pendunculisque densè hirsutis, foliis subsessilibus ovato-lanceolatis basi obtusis apice acuminatis subrenatis suprà junioribus puberis demùm muricatis subtus densè et molliter hirsutis, paniculæ terminalis ramosissimæ ramis scorpioides, capitulis sessilibus plerisque bractea foliaceâ suffultis 20-25-floris, invol. ovati squamis ext. hirsutissimis acumine rigido subpatente apiculatis, intimis glabrisculis acuminatis, achænio tereti villosa, pappi ser. ext. paleaceâ brevi. \mathcal{L} ? in Brasiliâ legit cl. Lhostky. Folia 3-4 poll. longa, 15-18 lin. lata. Pappus subrufescens. Cor. purpureæ? hirtæ? Tubercula paginæ fol. superæ sunt pilorum deciduorum bases. (v. s.)

230. V. SAGRANA, fruticosa ramosissima, ramis teretibus villosis, foliis brevissimè petiolatis ellipticis basi obtusis apice breviter acuminatis distanter serrulatis penninerviis suprà glabris subtus in nervis pubescentibus, ramis elongatis subscorpioides, capitulis unilateralibus ad foliorum axillas solitariis sessilibus folio multò brevioribus 30-floris, invol. ovati squamis ext. lanceolatis ciliatis, inter. longioribus glabris linearibus subacuminatis, cor. et achænio glabris, pappi ser. ext. angustè paleaceâ brevi. \mathfrak{z} in insulâ Cubâ circa Havanam legit cl. Ramon de la Sagra. Cor. purpureæ. Pappus albus. Achænium teres. Fol. 3 poll. longa, sesquipollicem lata, petiolo 1-1 $\frac{1}{2}$ lin. (v. s. comm. à cl. inv.)

231. V. ZUGCARINIANA (Mart. ! herb.), suffruticosa? caule tereti substriato erecto ramoso vix pubescente, foliis subsessilibus elliptico-oblongis utrinquè subobtusis hinc indè subserrulatis suprà muricatis subtus molliter hirsutis penninerviis, ramis scorpioides, capitulis 20-25-floris sessilibus distantibus solitariis geminive nunc ebracteatis nunc ad foliorum axillas sitis, invol. ovati squamis siccis imbricatis arcuè adpressis obtusis longè ciliatis dorso glabris, achænio glaberrimo, pappi serie ext. paleaceâ brevi. \mathcal{L} ? ad Minas Novas Brasilæ legit

cl. Martius. Fol. 3 poll. longa, sesquipollicem lata, Pappi setæ albæ, crassiusculæ. (v. s. in h. Acad. reg. Monac.)

232. *V. VARRONIAEFOLIA*, caule fruticoso ramoso tereti basi glabriusculo supernè pube pulverulentâ ferrugineo, foliis subsessilibus ovalibus utrinquè subobtusis subintegerrimis penninerviis reticulatis suprâ muriculatis bullulatis subtùs ferrugineo-velutinis, capitulis ad axillas foliorum secus ramos sessilibus solitariis aut geminis folio flor. triplò brevioribus oblongis 18-20-floris, invol. squamis ext. lanceolatis mucronato-acuminatis adpressis, inter. linearibus minùs acuminatis, cor. et achænio glabris, pappi serie ext. brevissimâ vix paleaceâ. 3 patr. ign. Cor. purpureæ 2 lin. longæ. Achænia striata. Fol. 2 poll. longa, 12-15 lin. lata. Capitula approximata secus ramos circ. 20-30. Pappus stramineus. Paulò accedit ad *V. Sagræanam* et ad *V. aurifodinarum*. (v. s. in h. Acad. reg. Monac.)

233. *V. LÆVIGATA* (Mart. ! herb.), glaberrima, caule herbaceo tereti, ramulis subangulatis, foliis lato-linearibus sessilibus integerrimis calloso-subacutis coriaceis utrinquè subreticulatis, ramis scorpioideis, capitulis ad axillam foliorum solitariis geminisve folio multò brevioribus 12-13-floris, invol. ovatis squamis oblongis adpressis subacutis, achænio villosa, pappi ser. ext. paleaceâ brevi. 2 in alpestribus prope Congonhas do campo prov. Minar. Gener. legit cl. Martius. Pappus stramineus scaber. Folia 4 poll. longa, 4 lin. lata. (v. s. in h. Acad. reg. Monac.)

♀. *bupleurifolia*, caule striato crassiori, foliis oblongis, super. ferè ellipticis, capitulis 17-18-floris, squamis invol. magis obtusis. 2 cum var. a? legit cl. Martius. Folia ferè Bupleuri fruticosi, 4 poll. longa, 8-10 lin. lata; summa poll. longa, 5 lin. lata. (v. s. in h. Acad. reg. Monac.)

234. *V. LACUNOSA* (Mart. ! herb.), caule herbaceo erecto tereti densè tomentoso, foliis subsessilibus ellipticis basi obtusis apice vix subacutis integerrimis suprâ pubero-velutinis subrugosis subtùs densè tomentosis penninerviis, nervis elevatis reticulatis, ramis floridis bifidis in paniculam ferè dispositis scorpioideis, capitulis sessilibus 10-floris ad dichotomias et in axillis foliorum ovatorum iis æqualium, invol. ovato-oblongi squamis erectis oblongis dorso lanatis intimis apice acuminatis, cor. lobis apice panicetosis, achænio sericeo-villoso, pappi ser. ext. paleaceâ acuminatâ. 2 in pratis montium prope Villa-Ricca Minar. Gener. Brasiliæ legit cl. Martius. Corollæ et stig. rosea. Herba 2-3-pedalis. Fol. 2 poll. longa, pollicem lata. Recept. nudum. (v. s. in h. Acad. reg. Monac.)

§. 6. *Involucris squamis intimis acuminatis, capitulis neque solitariis neque in cymas scorpioideas dispositis.* Less. l. c. p. 671. — *Species omnes Americanæ.*

235. *V. EXTRAXILLARIS*, caulibus è collo lignoso pluribus simplicibus subherbaceis teretibus tomentosis erectis apice arcuatis, foliis sessilibus linearibus acuminatis margine revolutis suprâ glabris nervo impresso subtùs tomentosis, cymâ terminali simplici, capitulis ebracteatis inter folia extra-axillaribus sessilibus solitariis ovatis 20-floris, invol. squamis rigidè acuminatis dorso araneosis, achænio teretisericeo-villoso, pappi aer. ext. angustè paleaceâ acuminatâ, inter. dimidio brevior. 2 in Brasiliæ prov. Minar. Gener. prope Tejuco unicâ vice legit cl. Vauthier. Caulis sesquipedalis gracilis. Folia sesquipoll. longa, 1-2 lin. lata, rigidula. Pappus stramineus. Capitula infima axillaria gemina, cætera extra-axillaria solitaria. (v. s. comm. à cl. Vauthier.)
236. *V. LINEARIS* (Spreng. syst. 3. p. 437), caulibus è rhizomate lignoso pluribus teretibus lignosis lanato-tomentosis simplicibus, foliis sessilibus linearibus acuminatis mucronatis 1-nerviis integerrimis margine revolutis suprâ glabris subtùs lanato-tomentosis, capitulis sparsis subsessilibus remotis folio stipatis, invol. subcylindracei squamis siccis pungenti-acuminatis arachnoideo-pilosis, cor. glabrâ, achænio densè piloso, pappi serie ext. paleaceâ brevi. 3 in Brasiliâ

- tropicâ. Less. in *linnæa* 1829. p. 285. 1831. p. 671. Cor. purpureæ. Pappus flavescent. Fol. 2 poll. longa, lineam lata.
237. *V. ROSMARTINIFOLIA* (Less. in *linnæa* 1829. p. 286. 1831. p. 672), caulibus lignosis ramosis teretibus lanato-tomentosis, foliis sessilibus 1-nerviis rigidis linearibus mucronato-acutis integerrimis margine revolutis suprâ glabris subtus lanato-tomentosis, capitulis approximatis multifloris subcylindraceis foliis plurimis stipatis, invol. squamis siccis pungenti-acuminatis ciliatis apice albo-pilosis, achenio densè piloso, pappi serie ext. brevi. $\frac{1}{2}$ in Brasiliâ tropicâ. Nervus folii pellucidus. Species valde affinis *V. lineari*.
238. *V. LINARIFOLIA* (Less. in *linnæa* 1829. p. 287. 1831. p. 672), caulibus è rhizomate lignoso plurimis teretibus subsimplicibus lanato-tomentosis, foliis sessilibus confertis linearibus 1-nerviis acuminatis coriaceis margine integerrimis et revolutis suprâ glabriusculis subtus lanato-tomentosis, capitulis campanulatis multifloris ramos nudos terminantibus, invol. squamis siccis 3-nerviis externis pungenti-acuminatis, achenio densè villosulo, pappi serie ext. brevi. $\frac{1}{2}$ $\frac{1}{2}$ in Brasiliâ tropicâ. Valde affinis *V. lineari*.
239. *V. ERYTHROPHILA*, caule herbaceo erecto simplici supernè angulato sparsè hispido apice nudo, foliis oblongo-linearibus sessilibus suprâ sparsè hirtis subtus nervo medio excepto glabris purpureis margine dentatis, dentibus acutis rigidè reflexis, corymbo 4-cephalo simplici aphylo, invol. squamis lanceolatis glabris, exter. obtusis, intimis acuminatis, achenio striato pilis rubris hirtello, pappi serie externâ squamellatâ brevi. $\frac{1}{2}$ in campis editis prov. Sancti-Pauli Brasil. ad Mogidas Cruzes legit cl. Lund. Nov. flor. Species dentibus retroflexis et pilis acheniî rubris spectabilis. Herba circ. pedalis. Folia 2 poll. longa, 4 lin. lata. (v. s. comm. à cl. inv.)
240. *V. CEPHALOTES*, caule herbaceo erecto simplici hirsuto, supernè nudo, infrâ medium folioso, foliis 4-5 approximato-subverticillatis sessilibus obovatis obtusis suprâ sparsè pilosiusculis, subtus canescenti-tomentosis, bracteolâ lineari in parte caulis nudâ, capitulis circ. 5-6 breviter pedicellatis ebracteatis in umbellam capitatum dispositis, invol. squamis lineari-lanceolatis pubescentibus apice acuminatis, acheniis villosis, pappi serie externâ paleaceâ brevi. $\frac{1}{2}$ in campis editis prov. Sancti-Pauli ad Mugi. Nov. flor. legit cl. Lund. An *Chrysocoma oligophylla* Arrab. fl. flum. 8. t. 2? Cor. purpureæ glabræ. Herba vix pedalis. Folia 3 poll. longa, sesquipollicem lata. (v. s. comm. à cl. Lund.)
241. *V. BARBATA* (Less. in *linnæa* 1829. p. 287. 1831. p. 672), fruticosa, ramis teretibus tomentosis, foliis sessilibus imbricatis cordato-ovatis acutis coriaceis integerrimis suprâ subpilosis infrâ villosis, cymæ terminalis ramis dichotomis, capitulis sessilibus folio stipatis, involucri turbinati pauciserialis squamis siccis 1-nerviis oblongo-acuminatis apice glabriusculis, achenio longè densèque sericeo, pappi serie ext. paleaceâ longiusculâ. $\frac{1}{2}$ in Brasiliâ tropicâ. Cor. lobis extus longè barbati. Pappus argenteus.
242. *V. VISCIDULA* (Less. in *linnæa* 1829. p. 289. 1831. p. 672), fruticosa, ramis teretibus viscido-pubescentibus, foliis sessilibus coriaceis ovato-oblongis acutis subdenticulatis suprâ lucidis subtus viscido-pubescentibus, cymis terminalibus ramosis, capitulis brevè pedicellatis multifloris, invol. turbinati squamis siccis glabris acuminatis, cor. glabrâ, achenio piloso, pappi serie paleaceâ brevi. $\frac{1}{2}$ in Brasiliâ tropicâ. Pappus argenteus.
243. *V. RUBROIDES* (Less. in *linnæa* 1829. p. 290. 1831. p. 672), caule suffruticoso simplici tereti supernè obscure pubescente, foliis brevè petiolatis confertis rigidis obovatis apice denticulatis suprâ glabris scabris subtus pubescentibus, cymis terminalibus oligocephalis, capitulis approximatis subsessilibus multifloris, invol. cylindracei squamis siccis 3-nerviis acuminatis glabris demum patentibus, achenio oblongo glabro basi et apice subpilosulo, pappi serie ext. paleaceâ brevi. $\frac{1}{2}$ in Brasiliâ tropicâ. Cor. ign. Pappus argenteus.
244. *V. SPECIOSA* (Less. in *linnæa* 1829. p. 290. 1831. p. 672), subarboorea, ramis teretibus villosulo-lanatis, foliis brevissimè petiolatis ovatis acutis coriaceis in-

tegerimiam, suprà glabris glandulosis subtus nervis villosis-lanatis, cymâ terminali spicæformi, capitulis sparsis in axillis fol. sup. sessilibus, invol. pauciserialis squamis siccis acuminatis subtomentosis, achænio densè piloso, pappi lati paleacei serie ext. longâ. 5 in Brasiliâ tropicâ. Folia $1\frac{1}{2}$ -2 $\frac{1}{2}$ poll. longa, 12-14 lin. lata.

245. *V. ARCTIOIDES* (Less. in linnæâ 1831. p. 400 et 672), fruticosa, ramis sulcatis angulatis tenerè tomentosis, foliis petiolulatis firmis obovato-lanceolatis cuneatis acuminatis inæqualiter serratis suprà lucidis glabris subtus arachnoideis, paniculâ terminali contractâ, capitulis multifloris, invol. squamis multinerviis ext. triangularibus rigidis, int. angustis acuminatis, corollâ et achænio glaberrimis, pappi seriebus latitudine non distinctis, ext. breviorè. 5 in Mexico. *Diazenxis serrata* Don trans. lin. soc. 16. p. 254. Folia 5-8 poll. longa, 2-3 $\frac{1}{2}$ poll. lata, petiolo 2-3 lin. Cor. purpureæ. Pappus argentens. Achænia 4-gona.

246. *V. ARGENTEA* (Less. in linnæâ 1831. p. 672), caule herbaceo simplici sulcato argenteo-tomentoso, foliis subsessilibus obovatis mucronatis coriaceis penninerviis inf. serratis suprà nitidis punctatis subtus argenteo-tomentosis, racemo terminali angusto, capitulis folio orbiculari basilari superatis, invol. campanulatis squamis laxè imbricatis linearibus pungentibus, cor. glabrâ, achænio hirsuto, pappi ser. ext. paleaceâ brevi. — in Brasiliâ. Fol. 4 poll. longa, 1 $\frac{1}{2}$ -2 lata. Racemus 2-8 poll. Cor. purpureæ.

247. *V. XANTHOPHYLLA* (Mart. ! herb.), caule fruticoso ramoso tereti tomento denso flavicante cum pedunculis et fol. paginâ infer. vestito, foliis subsessilibus ovalibus utrinquè obtusis subcrenatis coriaceis penninerviis suprà glabris, nervo medio basi tomentoso, capitulis 50-floris breviter pedunculatis in cymas contractas subpaniculatum dispositis, foliis floralibus ovatis sub ramulis, invol. squamis multiserialibus dorso tomentosis oblongis obtusis imbricatis, recept. nudo, achænio sericeo-villosissimo, pappi serie ext. ferè setiformi, int. triplò breviorè. 5 in campis arenosis in Chapada do Parana in confin. prov. Goyazæ legit cl. Martius. Cor. ign. Pappus rufo-flavescens. Folia sesquipollicem longa, pollicem lata. (v. s. in h. Acad. reg. Monac.)

248. *V. AUREA* (Mart. ! herb.), herbacea tota hirsuta, caule tereti subsimplici, foliis vix petiolulatis ovalibus utrinquè obtusis integerrimis, capitulis ad apices caulis et ramorum in axillis foliorum sessilibus solitariis folio paulò brevioribus circ. 40-floris, invol. campanulatis squamis lanceolatis, ext. in acumen gracile patulum productis, inter. acuminatis erectis, achænio sericeo-villoso, pappi ser. ext. angustè paleaceâ brevi. ○ in campis editis prov. Minarum Generalium Brasiliæ legit cl. Martius. Hirsuties partium juniorum flavescens. Fol. 15-16 lin. longa, 7-8 lata. Pappus albescens. Cor. glabræ ex sicco purpurascens. Radix descendens lateraliter fibrosa. (v. s. in h. Acad. reg. Monac.)

249. *V. CORDIGERA* (Mart. ! herb.), caulibus è collo suffruticoso pluribus herbaceis teretibus apice hirsutis, foliis sessilibus basi cordatis ovatis acutis rigidulis imbricatis integerrimis penninerviis reticulatis suprà glabris subtus parcè villosiusculis, ramulis foliosis simplicibus x-cephalis aut bifidis oligocephalis, capitulis subcorymbosis bracteatis 20-floris, invol. squamis oblongis parcè acuminatis basi glabris apice hirsutissimis, corollis achæniisque glabris, pappi ser. ext. angustè paleaceâ acuminatâ, inter. quadruplò breviorè. 2½ in editis prov. Minar. General. legit cl. Martius. Herba vix pedalis. Folia 15 lin. longa, 6-8 lata. Capit. 7-8 lin. longa. Cor. purpureæ. Pappus albidus. Achænia juniora hinc indè subpilosa. (v. s. in h. Acad. reg. Monac.)

250. *V. PYCNOSTACHYA*, fruticosa tota molliter hirsuto-tomentosa, caulibus teretibus ad apicem usque confertè foliosis, foliis ovali-aut obovato-oblongis acutis sessilibus planis integerrimis penninerviis, capitulis ad axillas supremas sessilibus folia sua non superantibus in spicam densam confertis circ. 40-floris, invol. ovato-campanulatis squamis coriaceis imbricatis pluriserialibus lanceolatis subacutis, junior. subvelutinis adultis glabris. 5 in Brasiliæ prov. Minar. Gener. ad Serro do Frio (Vauth. n. 267) in editioribus Minarum (Mart. !). Vern. decussata Mart. ! herb. sed folia non decussata, alterna. Flor. purpurei. Achænia.

sericeo-villosum. Pappi ser. ext. paleis oblongis apice incisis, int. setis filiformibus 2-3-serialibus crassiusculis. Recept. areolatum. (v. s.)

251. *V. TOMENTELLA* (Mart.! herb.), fruticosa, ramis teretibus junioribus tomentosis, foliis confertis sessilibus ovalibus subacutis integerrimis nervis utrinque prominulis supra glaberrimis subtus dense tomentosis, capitulis ad apices ramorum sessilibus apicatis extrâ-axillaribus folia sua non superantibus 20-floris, invol. ovati squamis imbricatis adpressis ciliatis acuminatis, cor. glabrâ, achenio sericeo-piloso, pappi ser. ext. paleaceâ brevi. $\frac{3}{4}$ in campis prov. Sancti-Pauli Brazilie legit cl. Martins. Simillima *V. pycnostachyæ* sed capitula 20 nec 40-flora, dimidio minora, folia paulò minora supra glaberrima. (v. s. in h. Acad. reg. Monac.)

252. *V. ROSA* (Mart.! herb.), fruticosa, ramis teretibus tomentosis, foliis sessilibus lanceolatis basi subacutis apice subacuminatis integerrimis supra glabrisculis subtus tomentosis penninerviis, ramis erectis, capitulis paucis sessilibus lateralibus et terminalibus folio flor. aequalibus 80-90-floris, invol. subglobosi squamis adpressis lanceolatis acuminatis dorso araneoso-sublanatis, cor. glabrâ, achenio sericeo-villoso, pappi ser. ext. angustè paleaceâ interiore circ. dimidio brevior. $\frac{3}{4}$ in campis prope Villa do Principe prov. Minarum General. Brazilie legit cl. Martins. Flores rosei. Folia 12-15 lin. longa, 3-5 lata. Capit. 7-8 lin. diam. Recept. nudum. (v. s. in h. Acad. reg. Monac.)

253. *V. VERTICILLATA* (Mart.! herb.), caule fruticoso ramoso, ramis teretibus sericeo-villosis ad apicem foliosis, foliis sessilibus ovalibus lanceolatisve planis subintegerrimis supra glabris subtus adpressè sericeo-villosis, capitulis ex axillis supremis pedunculatis subcorymbosis 20-25-floris, pedunculis nudis folio suo brevioribus, invol. ovati squamis imbricatis lanceolatis siccis dorso pubescentibus subacutis. $\frac{3}{4}$ in deserto trans Serram de Saint-Antonio et prope Caitete prov. Min. Gen. et Bahiam Brazilie legit cl. Martins. Recept. areolatum. Ach. sericeo-villosum. Pappi ser. ext. paleis oblongo-linearibus acutis, int. setis quadruplò longioribus scabris crassiusculis. Cor. purpureæ. (v. s. in h. Acad. reg. Monac.)

Variat α : foliis latioribus capitulis distinctè pedunculatis, et β : foliis angustioribus capitulis subsessilibus.

254. *V. SENSUENS* (Mart.! herb.), caule fruticoso tereti hirsuto-tomentoso, foliis sessilibus basi cordatis ovatis aut suborbiculatis apice calloso-subacutis integerrimis utrinque (subtus præcipuè) villosa-tomentosis, paniculæ ramis axillaribus brevibus oligocephalis, capitulis bracteâ foliaceâ longioribus 10-floris ad apices ramorum confertis subsessilibus, invol. squamis lanceolatis acuminatis extus velutino-villosis, cor. lobis extus ad apicem barbatis, achenio sericeo-villosissimo, pappi ser. ext. paleaceâ acuminatâ inter. triplò brevior. $\frac{3}{4}$ in campis editis prov. Minarum gener. et inter. Bahiensis legit cl. Martins. Variat foliis suborbiculatis 8-9 lin. longis latisque, et ovatis 2 poll. longis, pollicem latis. Invol. 4-6 lin. longum. Pappus albidus. (v. s. in h. Acad. reg. Monac.)

255. *V. FIRMULA* (Mart.! herb.), caule fruticoso tereti erecto ramulisque hirsutis, foliis sessilibus oblongo-lanceolatis rigidis acuminatis integerrimis supra glabrisculis subtus villis longis sparsis hirsutis, paniculæ elongatæ ramis oligocephalis foliosis brevibus, capitulis 40-floris subbracteatis, invol. squamis lineari-lanceolatis acuminatis ext. hirsutis intimis ferè glabris, cor. lobis sub apice barbatis, achenio sericeo-villoso, pappi ser. ext. paleaceâ brevi. $\frac{3}{4}$ in campis editis diserti Minarum General. legit cl. Martins. Valde affinis *V. dura* sed hirsutie faciliè distinctâ. Fol. 2 poll. longa, semi-poll. lata. (v. s. in h. Acad. reg. Monac.)

256. *V. DURA* (Mart.! herb.), suffrutescens? caule erecto tereti glabro, foliis sessilibus oblongo-lanceolatis rigidis subacuminatis integerrimis utrinque glabrisculis, ramis subscorpioides parcè hirsutulis, capitulis 40-floris pedicellatis bracteâ foliaceâ sæpius suffultis, invol. campanulati squamis acuminatis ext. dorso subhirsutis, corollæ lobis subbarbatis, achenio elongato sericeo-villo-

assimo, pappi ser. ext. angustè paleaceâ, int. ciliatâ externam duplò tantum superante. ? in campis Brasiliæ ad Contendas prov. Minar. General. legit cl. Martius. Folia 3 poll. longa, pollicem lata; axillæ subbarbatæ. Pappus rigidus stramineus. Recept. omninò nudum. (v. s. in h. Acad. reg. Monac.)

257. V. LESSINGIANA (Mart. ! herb.), caule suffruticoso tereti tomentoso-hirsuto, foliis sessilibus cordiformibus vix subacuminatis coriaceis integerrimis suprâ pube deciduâ subvillosis subtus subhirsutis, paniculæ ramis brevibus axillaribus bifidis oligocephalis, foliis subcapitulis ovatis utrinquè tomentosis capitulo brevioribus, capitulis confertis 3-5 sessilibus 10-floris, invol. ovalis squamis lanceolatis acuminatis dorso subvillosis, cor. lobis dorso hirsutissimis, achænio sericeo-villosissimo, pappi ser. ext. paleaceâ acuminatâ int. triplò breviora. ? in campis prope Formigas prov. Minar. Gener. Brasiliæ legit cl. Martius. Fol. 8-11 lin. longa, 4-6 lata. Pappus albus. (v. s. in h. Acad. reg. Monac.)

258. V. ECHITIFOLIA (Mart. ! herb.), caule herbaceo ? erecto simplici tereti striato subhirsuto apice subtomentoso, foliis breviter petiolatis ovatis ovalibusve utrinquè subacutis apice distanter et callosò denticulatis suprâ crebrè glandulosis puberulis subtus parçè glandulosis subhirsutis, paniculæ ramis brevibus subscorpioideis oligocephalis, capitulis confertis 10-12 sessilibus folio florali longioribus, invol. ovalis squamis lanceolatis dorso subtomentosis adpressis ext. mucronatis int. acuminatis, cor. glandulosâ apice setosâ, achænio glabro glanduloso?, pappi serie ext. paleaceâ brevissimâ. ? in montibus prov. Minar. Gener. Brasiliæ legit cl. Martius. Fol. 2 poll. longa, pollicem lata, petioli 2 lin. Nervus medius suprâ ad basin hirsuto-tomentosus. Pappus albus argenteus. Recept. nudum. (v. s. in h. Acad. reg. Monac.)

259. V. TRAGIÆFOLIA, caule basi suffruticuloso herbaceo confertè hispido usquè ad cymam folioso, foliis sessilibus oblongis acutis crenato-subdentatis suprâ confertè villosis subtus cano-tomentosis, capitulis secus ramulos cymæ sessilibus confertis hinc indè foliis stipatis in fasciculum densum glomeratis oblongis 10-floris, invol. squamis villosis pluri-serialibus acuminatis, apice uncinato-recurvus, achæniis pubescentibus, pappi serie ext. paleaceâ, paleis acutis ciliatis. ? in campis editis prov. Sancti-Pauli Nov. flor. legit cl. Lund. Species distinctissima vix scorpioidea. Cor. purpureæ glabræ. Folia pollicem longa, 4 lin. lata. (v. s. comm. à cl. Lund.)

Sect. IX. ASCARICIDA Less. in *linnæâ* 1831. p. 675. Ascaricida et Centrapalus Cass. dict. 3. suppl. p. 38 et 7. p. 383. — Baccharoides Lin. f. Zeyl. Moench meth.

Involucri campanulati squamæ foliæ aut apice appendiculatæ plûs minûs patentés et inter se dissimiles. Herbæ capitulis terminalibus solitariis aut paucis-corymbosis.

§. I. *Involucri squamis exterioribus latioribus.*

260. V? CALYCINA (Wall. ! cat. et herb.), caule tereti glabro apice minutissimè pubescente, foliis latè ovalibus obtusis penninerviis subsessilibus integerrimis glabrinusculis subtus tenuissimè glandulosis, capitulis brevè pedunculatis corymboso-adgregatis, involucri squamis ext. latè elliptico-oblongis obtusis submucronulatis pedunculisque pubescentibus intimis linearibus acutis minoribus achænio subvillosis, pappi pilis omnibus latitudine subæqualibus ext. dimidio brevioribus. ? in Ind. orient. regno burmannico ad Promè. Cor. extus glandulosa. Flor. in invol. circ. 40-45. Habitus Vernoniæ sed involucro et pappo fortè genericè distinguenda. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

261. V. MACROCEPHALA (Less. ! in *linnæâ* 1829. p. 298. 1831. p. 679), caulibus herbaceis basi suffruticosis teretibus simplicibus tomentosis, foliis brevè petiolatis oblongis utrinquè acuminatis integerrimis suprâ glabris subtus tomentosis penninerviis, cymis scorpioideis polycephalis, capitulis distantibus multi-

floris, involucri latè campanulati squamis oblongo-ellipticis obtusis extus tomentosis, achænio oblongo multicostato sericeo-piloso, pappi serie externâ brevi paleaceâ. ♀ in Brasiliâ meridionali. Pappus subflavidus. Cor. rosæ glabræ. Rhizoma lignosum repens. Folia 3-10 poll. longa, 8-30 lin. lata. Flores in capitulo 40-50. Capit. interdum proliferum vidi. Affinis videtur *V. ligulæfoliæ*. (v. s. comm. à Mus. reg. Berol.)

262. *V. MEXICANA* (Less. in *linnæa* 1831. p. 680), fruticosa, ramis teretibus cinereo-subtomentosis, foliis petiolatis obovatis acutis basi cuneatis denticulatis suprâ glabrinsculis scabridis subtus tenuiter tomentoso-incanis, paniculæ corymbosæ ramis aphyllis, invol. globosi squamis imbricatis glabris, ext. subrotundis acuminatis, med. ellipticis obtusis apice foliaceis, int. linearibus acutis, cor. puberis, achænio glaberrimo, pappo setaceo inæquali. ♂ in Mexico ad Cuesta-Grande de Chiconquiaco. Cor. cærulæ. Fol. 8 poll. longa, 6 poll. lata, petiolo 2 lin. Pappus flavescens. Habitus *V. arctioidis*.

263. *V. ALAMANI*, 'caule...., ramis apice striatis subpuberis, fol. inf...., summis brevissimè petiolatis elliptico-oblongis basi attenuatis apice acuminatis subintegerrimis suprâ muricato-scabris subglandulosis puberulis penninerviis, capitulis pedunculatis ex axillis et apice ramorum paucis nudis 40-45-floris, invol. subglobosi squamis coriaceo-scariosis multiseriatis glaberrimis ovalibus obtusis mucrone longo patente superatis, cor. glabrâ, achænio glanduloso striato, pappi serie ext. brevi paleaceâ. ♀ ? in Mexico. Frustula tantum sed distinctissima olim misit cl. Alaman. Capit. 8-10 lin. diam. Cor. purpureæ. Pappus albus, ext. subciliatus. (v. s.)

§. 2. *Involucri squamis exterioribus angustioribus.*

264. *V. BOJER* (Less. in *linnæa* 1831. p. 676), caule herbaceo erecto ramoso tereti glanduloso-hispido, foliis super. ovali-oblongis basi attenuatis apice subacuminatis integerrimis suprâ subpuberulis subtus cano-tomentosis reticulatis, capitulis circ. 30-floris, corymbo laxo divaricato, invol. campanulati squamis dorso glanduloso-puberis imbricatis oblongis productis in appendicem setiformem patulam, achæniis villosis. ♂ in campis et ad margines sylvarum insulæ Zanzibar legit cl. Bojer. Species distinctissima. Frutex 8-10-pedalis. Habitu *Decaneuri* sed achænium teretiusculum. Folia ferè *V. Zeylanicæ* sed indivisa. Pappus albus subplumosus, ser. ext. brevissima. (v. s. comm. à cl. Bojer.)

265. *V. ANTHELMINTHICA* (Willd. sp. 3. p. 1634), caule tereti hispidulo-pubescente superne ramoso, foliis ovato- aut ovali-oblongis utrinque acuminatis grossè serratis puberulis, capitulis corymbosis 40-45-floris, invol. squamis lanceolato-linearibus acutis, exterioribus subpatulis foliaceis obovato-linearibus, achæniis oblongis teretiusculis hispidulis, pappi serie ext. brevissima paleaceâ. ☉ in Indiâ orient. frequens. Ham. tr. lin. soc. 14. p. 217. Less. in *linnæa* 1829. p. 293. *Conyza anthelmintica* Lin. sp. 1297. *Serratula anthelmintica* Roxb. cat. calc. 60. *Baccharoides anthelmintica* Moench. 578. *Ascaricida Indica* Cass. dict. 3. suppl. p. 38. *Serratula elegans* herb. Maurit. — Burm. Zeyl. t. 95. Rheed. mal. 2. t. 24. Cor. purpurascens. Variat caule folisque plus minus hirsutis glabrativæ. (v. v. et s. sp. è plur. Indiæ locis.)

266. *V. PAUCIFLORA* (Less. in *linnæa* 1829. p. 292. 1831. p. 675), caule herbaceo ramoso teretiusculo pube brevi hirtello, foliis elliptico-lanceolatis utrinque acuminatis rariter serratis suprâ sparse hirtellis subtus pubè brevi præsertim secus nervos scabris et subglandulosis, capitulis 40-45-floris solitariis terminalibus, invol. squamis ext. subulatis laxis interiores lineares 3-nervias subæquantibus, achænio villosis, pappi setis barbellatis, serie externâ brevissima vix paleaceâ. ☉ in Senegalicæ plantie regionis Walo locis subhumidis legit cl. Perrottet. *Conyza pauciflora* Willd. sp. 3. p. 1927. Spreng. syst. 3. p. 514 excl. Link. syn. *Vernonia Senegalensis* Desf. cat. h. Par. 1829. p. 400. non Less. *Centrapalus Galamensis*, Cass. dict. 7. p. 383. *Vern. Galamensis* Less. in *linnæa*

1829. p. 314. *Vernonia cœlestina* Sohrad! hort. Gœtt. Cor. purpureo-cœruleæ. Affinis *V. anthelminticæ*. (v. v. c. et s. sp.)

β. *angustifolia*, foliis angustioribus oblongo-lanceolatis, summis ferè linearibus. — in sabulosis regionis Walo Senegalæ ubi à Nigritis dicitur *Soughan* ex Perrottet. (v. s.)

267. *V. BRACTEOLATA*, caule herbaceo? erecto tereti basi glabrinseulo apice adpressè villoso-hirsuto corymboso, foliis subsessilibus obovato-cuneatis breviter acuminatis distanter serratis utrinquè glabris, nervo medio subtus subvelutino, corymbi fastigiati pedunculis parcè ramosis sub capitulo longè bracteolis aristatis imbricatis onusto, invol. squamis multiserialibus aristatis subvillosis, intimis latoribus. ½ ex Nepaulià ad hortum Calcuttensem transl. et ibi à cl. Wallich observata; in Bengalià legit cl. Belanger. *Conyza bracteolata* Wall.! cat. n. 3036. comp. 146. Flor. in sicco albidæ, in invol. 50-60. Achæn. teres villosum. Pappus rufescens multiserialis, serie ext. brevi irregulari. Folia 7 poll. longa, 3-4 poll. lata. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
268. *V. SUBSESSILIS*, caule fruticoso erecto ramoso tereti, ramis apice pubescenti-villosis, foliis subsessilibus oblongis subintegerrimis basi subattenuatis apice longè cuspidato-acuminatis glabris subtus pallidis cum nervo medio pubescente, corymbo laxo, capitulis longè pedunculatis, pedunculis sub capitulo bracteolas subalatas paucis gerentibus, invol. multiser. squamis ext. brevioribus subulatis squarrosis, inter. longioribus planis lanceolato-linearibus acutis. ½ in Indiæ orient. prov. Silhet legit Franc. da Sylva. *Conyza divergens* Wall.! cat. n. 3000. comp. n. 110. Affinis *V. bracteolata*. Capit. 30-35-florum. Pappus rufus longus, ext. irregularis brevis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
269. *V. KARVINSKIANA*, caule herbaceo erecto basi terete glabro apice angulato puberulo, foliis petiolatis lanceolatis utrinquè acuminatis distanter serratis penninerviis suprà sparè scabridis subtus pubescenti-hirsutis, corymbo composito terminali fastigiato, pedicellis parcè bracteolatis capitulo 20-floro triplo longioribus, invol. squamis glabris, ext. lanceolato-acuminatis ferè subulatis patulis, mediis lanceolatis trinerviis, intimis lineari-lanceolatis subacutis subfoliaceis erectis, cor. et achænio glabris, pappi ser. ext. subnullà aut minimâ. ½ in Mexico legit cl. Karvinski. Habitus ferè *V. noveboracensis*. Caulis purpureus. Fol. 3 poll. longa, 7-8 lin. lata, petiolo 2-3 lin. Cor. purpureæ. Achænia angulata glandulosa. Recept. nudum. (v. s. in h. Acad. reg. Monac.)
270. *V. CORYMBIFORMIS*, caule herbaceo erecto tereti striato glabrinseulo apice angulato subpuberulo, foliis petiolatis ovali-lanceolatis utrinquè acuminatis rarissimè subserratis penninerviis suprà glabris, subtus tenuiter pubescentibus, corymbis ad apicem caulis et ramorum compositis, pedicellis parcè bracteolatis capitulo 20-floro quadruplo longioribus, invol. squamis ext. lineari-subulatis patulis, mediis lanceolatis, int. lineari-lanceolatis acutis submembranaceis, cor. et achænio glabris, pappi ser. ext. paleaceâ brevi distinctissimâ. ½ in Mexico legit cl. Karvinski. Affinis *V. Novæboracensi* et *Karvinskianæ*. Cor. purpureæ. Fol. 4-5 poll. longa, sesquipollicem lata, petiolo 2-3 lin. Pappus albus. Achænia striato-costata glandulosa. (v. s. in h. Acad. reg. Monac.)
271. *V. INULOIDES*, caule herbaceo erecto tereti ramoso pubescente, foliis petiolatis ovatis basi subacutis apice acuminatis subintegerrimis suprà pubescentibus subtus pubescenti-subhirsutis, corymbi compositi ramis sæpè trifidis, pedicellis vix bracteolatis capitulo 35-40-floro duplè ferè longioribus, invol. squamis ext. linearibus laxis subpatulis, inter. ovali-oblongis obtusis foliaceo-membranaceis, cor. et achænio glabris, pappi ser. ext. ferè setosâ brevi. ½ in Mexico legit cl. Karvinski. Achæniarum glandulosum. Pappus albus. Cor. purpureæ. Folia 3 poll. longa, sesquipollicem lata, petiolo 5 lin. Herba habitu ferè *Inulæ Conyzeæ*, affinis *Novæboracensi* et *anthelminticæ*. (v. s. in h. Acad. reg. Monac.)
272. *V. OLIGOPHYLLA* (Michx. fl. bor. am. 2. p. 94), caule herbaceo simplici glabro supernè ferè aphylo parcè ramoso, foliis dentatis acutis glabris, radicalibus

obovatis latis, caulinis lanceolatis angustis paucis, corymbo paniculato, involucri floribus brevioris squamis lanceolatis ciliatis acuminatis, ext. filiformibus, achæniis in costis pilosiusculis. \mathcal{U} in pinetis humidis et paludosis Carolinæ merid. Ell. sketch, 2. p. 286. Less. in linnæa 1831. p. 677. Chrysocoma acaulis Valt. car. 196. Cor. purpureæ. (v. s.)

a. verna, utroque flore pedunculato. (Michx. l. c.)

β. autumnalis, altero flore subsessili. (Michx. l. c.)

273. *V. NOVÆBORACENSIS* (Willd. sp. 3. p. 163a), caule herbaceo erecto striato-sulcato supernè ramoso glabriusculo, foliis breviter petiolatis elongato-lanceolatis suprâ glabris subtus subpubescentibus serrato-denticulatis, corymbo composito subscorpioideo fastigiato, invol. ovati squamis basi adpressis apice in setam filiformem patulam productis, achæniis glaberrimis, pappi ser. ext. paleis brevibus latiusculis. \mathcal{U} ad margines sylvarum et in pascuis à Carolinâ ad Canadâ. Serratula Novæboracensis Linn. sp. p. 1146. Vernonia præalta Willd. sp. 3. p. 163a. excl. syn. Less. in linnæa 1829. p. 264. — Dill. elth. 2. f. 342 et fere sinè dubio f. 341. Undè huc etiam sunt referendæ: Serratula glauca Linn. sp. 1146. Suprago glauca Gaert. fr. 2. t. 166. Vernonia glauca Willd. sp. 3. p. 1633. Huc etiam ex cl. Lessing verisim. pertinent Vernonia tomentosa Ell. sketch, 2 p. 288 non Nees (et ideo Chrysocoma tomentosa Valt. car. p. 196), nec non *V. Baldwini* James in ann. New-York 2 p. 211. Herba 5-pedalis. Caulis glaber pubescens aut apice subtomentosus. Cor. purpureæ. Achænia costata inter costas glandulosa. (v. v. c. et s. sp.)

274. *V. ANGUSTIFOLIA* (Michx. fl. bor. am. 2. p. 94), caule herbaceo simplici apice pilosiusculo, foliis crebris longè angustèque linearibus 1-nerviis subintegerrimis suprâ lævibus glabris secus nervum subtus pilosiusculis, corymbo fastigiato, ramis hinc inde scorpioideis, involucri squamis in acumen subfiliforme subpatulum desinentibus, achæniis hispidulis. \mathcal{U} in siccis et collibus arenosis Carolinæ. Ell. sketch, 2. p. 287. Less. in linnæa 1829. p. 262. Chrysocoma graminifolia Valt. Car. 196.

275. *V. SCABERRIMA* (Nutt. gen. am. 2. p. 134), caule herbaceo simplici basi piloso apice læviusculo, foliis lineari-lanceolatis acutis sessilibus remotè serrulatis suprâ scabris subtus pilosis, corymbo subumbellato panicapitulato, invol. squamis lanceolatis nudis in acumen filiforme desinentibus, achænio pilosiusculo. \mathcal{U} in Carolinâ australi et Floridâ, in pinetis. Ell. sketch, 2. p. 286. Less. in linnæa 1829. p. 263. 1831. p. 677. Serrulata pilosa Bosc! herb. Vern. angustifolia et scaberrima Ell. sketch, 2. p. 286. ex Less. Caulis 2-pedalis. Cor. lætè purpureæ. (v. s.)

§. 3. *Involucri squamis exterioribus brevissimis omnibus adpressis exappendiculatis. — Species in sect. dubiæ.*

276. *V. FASCICULATA* (Michx. fl. bor. am. 2. p. 94 non Blum.), caule herbaceo tereti glabriusculo, foliis longo-linearibus rariter serratis pube sparsè subscabridis margine subrevolutis, floribus corymbosis erecto-approximatis, involucri ovoidei lævis squamis adpressis acutis muticis. \mathcal{U} in pascuis reg. Illinoensibus et Virginæ. Capitula parva. Specimen meum (è Carol. merid. ortum fortè à sp. Michx. diversum?) differt à *V. scaberrimâ* exactè ut *V. præalta* à *Novæboracensi*.

277. *V. PRÆALTA*, caule herbaceo erecto striato elongato apice corymboso pubescente, foliis breviter petiolatis lanceolatis serratis suprâ glabris subtus puberulis, corymbo composito subscorpioideo irregulari, invol. ovati squamis ovatis vix acutis adpressis ciliatis, achæniis glabris, pappi ser. ext. paleaceâ brevi. \mathcal{U} in Amer. bor. à Georgiâ ad Kentucky, ad margines paludum. Serratula præalta Linn. sp. 1146. Chrysocoma gigantea Walt. car. 196. Vern. præalta et altissima Ell. sketch, 2. p. 289. Nutt. gen. am. 2. p. 236. Vern. altissima Less. in linnæa 1829. p. 265, 1831. p. 639. — Dill. elth. f. 343. Pluk. alm. t. 240. f. 6. Mill. ic. t. 234. Species Linnæana cum aliis sæpè confusa est

ex iconibus citatis facillè restituenda. Variat caule glabro aut pubero, foliis basi obtusis (Less.) aut sæpius acuminatis. (v. v. et s.)

β. *hebecarpa*, achæniis secus costas puberulis. 2. in Kentucky. Vern. altissima Rafin. in litt. 1831. Var. α simillima achæniis exceptis. (v. s. sp.)

† *Non satis notæ.*

278. V. ? *ACHYROCOMA* (Less. in Linnæa 1829. p. 313), caule herbaceo striato tomentoso ramoso, foliis subsessilibus oblongo-lanceolatis serratis suprà glabris subtus tomentosis, capitulis pedunculatis multifloris, invol. squamis coriaceis tomentosis apice glandulosis, ext. lanceolatis, int. oblongis obtusis, achænio villosa, pappi paleis pluriseriatis linearibus denticulatis exterioribus brevioribus, corollis apice glandulosis. — Patr. dub. forsan Amer. bor. Dispositio florum ignota. *Achyrocoma tomentosa* Cass. dict. 26. p. 21.
279. V. *AFFINIS* (H. B. et Kunth nov. gen. 4. p. 39), fruticosa, ramulis subangulatis puberulento-tomentosis, foliis petiolatis ovato-oblongis acuminatis basi rotundatis integerrimis coriaceis suprà glabris subtus hirtotomentosis, cymis terminalibus paniculæformibus foliatis, capitulis subsessilibus 5-6-floris, invol. squamis oblongis glabris, achænio glabro oblongo sulcato. 3. in Peruvia? cum V. floribunda à quâ vix differt nisi foliis basi magis rotundatis et achæniis minoribus.
280. V. *ANDROMACHIOIDES* (Less. in Linnæa 1831. p. 397 et 645), fruticosa, ramis diffusis teretibus striato-sulcatis supernè hispido-puberulis, foliis petiolatis decussatis membranaceis triplinerviis subrotundis mucronatis basi obtusis subintegerrimis suprà glabris subtus tenuiter villosotomentosis, paniculâ trichotomâ terminali ramis oppositis, capitulis 14-17-floris, invol. cylindraceis squamis glabris ciliatis ext. subrotundis acutis int. linearibus obtusis, recept. angusto fimbriifero, cor. glabrâ, achænio hirsuto, pappi ser. ext. brevi vix paleacea. 3. in sylvis Misanthæ Mexici. Cor. luteæ. An idèò *Vernonia* vera?
281. V. *ARISTATA* (Less. in Linnæa 1829. p. 313), fruticosa? ramis..., foliis petiolatis lanceolatis apiculatis subdentato-sinnatis, dentibus spinuliformibus, suprà sparè pilosis subtus villosis glandulosis, cymis scorpioideis, capitulis subaggregatis uniseriatis folio florali suffultis, invol. squamis exter. apice subulatis rigidis. — Patr. ign. *Lepidaploa aristata* Cass. dict. 26. p. 17.
282. V. *EMARGINATA* (Wikstr. quad. fl. 73), fruticulosa ramosa, ramis sursum villosis, foliis petiolatis ovalibus integerrimis apice emarginatis impresso-punctatis suprà glabris subtus pubescentibus, paniculâ multicapitatâ ramis secundifloris, invol. ovatis squamis lanceolatis acutis extus villosis. 3. in Guadalupe. Affinis dicitur V. albicauli Pers. sen Vahliaenæ Less. †.
283. V. *BACCHAROIDES* (H. B. et Kunth nov. gen. 4. p. 40), arborescens, ramulis angulatis hirtotomentosis, foliis petiolatis oblongis subacuminatis basi rotundatis remotè denticulatis suprà pubescentibus tomentoso-hirtis, cymis terminalibus paniculæformibus ramosissimis, capitulis 3-5 aggregatis multifloris, invol. hemisphærici squamis acutis glabris ciliatis. 3. in montosis Andium Novo-Granatensium. Cor. albæ glabræ. Pappus rufescens biserialis.
284. V. *CORDIFOLIA* (H. B. et Kunth nov. gen. 4. p. 38), suffruticosa?, ramulis subangulatis molliter tomentoso-pilosis, foliis petiolatis ellipticis breviter acuminatis basi rotundatis integerrimis rigidis suprà glabris nitidis subtus molliter piloso-pubescentibus, cymâ terminali paniculæformi patentissimâ, capitulis sessilibus, invol. semiglobosi squamis acutis subpungentibus. 3? in scopulis Novo-Granatensium.
285. V. *FRAGRANS* (Lallav. et Lex. nov. veg. descr. 1. p. 24), fruticoso-sarmentosa, foliis breviter petiolatis sublanceolatis integris margine revolutis scabriusculis, cymâ spicæformi subsimplici, capitulis sessilibus unilateralibus distantibus multifloris, involuero ventricosso. 3. ad vias in montosis prope S. Jose del Corral Mexicanorum. Flores albi vanillam olentes. Affinis videtur V. odoratissimæ aut Deppeanæ. †.

286. *V. FLORIBUNDA* (H. B. et Kunth nov. gen. 4. p. 38), fruticosa, ramulis angulatis hirta-tomentosis demum striatis glabris, foliis petiolatis oblongis acuminalatis basi obtusis subintegerrimis coriaceis supra glabris subtus canescentibus hirta-tomentosis, cymis paniculaeformibus axillaribus et terminalibus folio brevioribus, capitulis 4-5 floris subsolitaris, involucri oblongi squamis obovatis apice inciso-lacinatis?, achenio sulcato glabro. $\bar{\sigma}$ in Amer. merid. fortè in Peruvia. Cor. ign. Pappus duplex sed proportio serierum non indicata.
287. *V. FRUTICOSA* (Swartz fl. ind. occ. 1323), fruticosa, ramis teretibus..., foliis brevissimè petiolatis ovatis obtusis integris..., cymæ terminalis ramis elongatis flexuosis, capitulis sessilibus distantibus folio florali ovato brevioribus, $\bar{\sigma}$ in insul. Caribæis. *Conyza fruticosa* Lin. sp. 1209. — Plum. ed. Burm. t. 95. f. 1. †
288. *V. LINIFOLIA* Blum. bijdr. ned. ind. p. 893), herbacea, ramis paniculatis subtrigosis, foliis lineari-lanceolatis rariter serrulatis puberulis, corymbo terminali dichotomo divaricato laxifloro, squamis involucri cuspidatis. ☉? in montanis humidis ins. Javæ. Affinis dicitur *V. leptophyllæ*. An fortè eadem ac *V. erigeroides*?
289. *V. STELLARIS* (Lallav. et Lex. nov. veg. descr. 1. p. 23), arborescens, ramulis tomentosis, foliis brevè petiolatis lanceolatis supra pilosisculis subtus fuscescenti-tomentosis margine revolutis subintegerrimis, cymæ terminalis scorpioidæ ramis tomentosis flexuosis, capitulis sessilibus, invol. campanulati squamis acutis pilosis, corollæ tubo incurvato, lobis conniventibus. $\bar{\sigma}$ in pascuis ad José del Corral Mexicanorum. Flores albi vanillati. †.
290. *V. PROLIFERA* (Decaisn. n. ann. mus. h. n. Par. 3. p. 408), caule herbaceo striato puberulo, foliis subrotundo-ovatis obovatisve repando-subangulatis interdum oblongis obtusis subtus pallidioribus basi in petiolum attenuatis, panicula terminalibus ramosis, ramis proliferis, invol. floribus dimidio breviori. — in insulâ Timor. Cl. auctor hic citat *Conyzam proliferam* Lam. et *Sonchum volabilem* Rumph., sed ex descr. iconè et patriâ potiùs meo sensu ad *Microglossam volabilem* referendæ sunt.

†† Species vix notæ.

Obs. 1. Quid *Vernonia Missurica* et *Vernonia albiflora* (Raf. herb. p. 28)?

Obs. 2. Præter species hinc inde citatas ad *Vernoniam* referendæ videntur stirpes sequentes iconibus Floræ fluminensis incompletè cognitæ, nempe: *Chrysocoma arborea* S. t. 12, *C. repanda* t. 13, *C. paniculata* t. 14, *C. medio-serrata* t. 15, *C. cymosa* t. 16, *C. pilosa* t. 17, *C. sessilis* t. 20, *C. quadrangularis* t. 25, *C. linearis* t. 27, *C. horizontalis* t. 28, *C. herbacea* t. 29, *C. monantha* t. 33, *C. tomentosa* t. 38, *C. Precaa* t. 39, et *C. alterna* t. 40.

††† Species exclusæ.

- V. AMYGDALINA* Delil. = *Decaspermum amygdalinum*.
V. BRACTEATA Wall. = *Decaspermum Silhetense*.
V. BRASILIENSIS Less. = *Lychnophora brasiliensis*.
V. BRUNIIFORMIS Less. = *Lychnophora brunioides*.
V. CAPITATA Less. = *Albertinia capitata*.
V. CHINENSIS Less. = *Cyanopsis pubescens*.
V. ECHINOPS Mart. = *Echinopsidium*.
V. EXSICCATA Mart. = *Echinopsidium exsiccum*.
V. FIMBRILLIFERA Less. = *Decaspermum fimbriifolium*.
V. HALIMODENDRON Mart. = *Echinopsidium*.
V. HAKEFOLIA Less. = *Lychnophora hakeefolia*.
V. INCANA Less. = *Albertinia incana*.

- V. INCANESCENS Mart. = *Albertinia incanescens*.
 V. PINASTER Less. = *Lychnophora pinaster*.
 V. PINIFOLIA Less. = *Webbia pinifolia*.
 V. PLANTAGINIFOLIA Less. = *Pycnocephalum plantaginifolium*.
 V. POLYGALÆFOLIA Licht. = *Pegolettia polygalæfolia*.
 V. POPULIFOLIA Less. = *Distephanus populifolius*.
 V. PROTEÆFORMIS Less. = *Lychnophora proteæformis*.
 V. PYCNOCEPHALA Mart. = *Echinopsidium pycnocephalum*.
 V. SALICIFOLIA Less. = *Lychnophora salicifolia*.
 V. SCARIOSEA Poir. = *Liatris scariosa*.
 V. SCAPIGERA Less. = *Pycnocephalum scapigerum*.
 V. SENEGALENSIS Less. = *Decaneurum Senegalense*.
 V. SMITHIANA Less. = *Webbia Smithiana*.
 V. SPATHULÆFOLIA Mart. = *Pycnocephalum spathulæfolium*.
 V. STAAVIDOIDES Less. = *Lychnophora staauidoidea*.
 V. STACHYANTHUS Mart. = *Stachyanthus Martii*.
 V. TRICHOCARPHA Spreng. = *Piptocoma lychnophoroides*.
 V. VILLOSISSIMA Less. = *Lychnophora villosissima*.
 V. WIGHTIANA Wall. = *Blumea vernonioides*.
 X. DECANEURUM DC. in *ann. bot. Guill.* 1833. v. 2. p. 516. et in *Wight contrib. bot. ind.* p. 3.

Capitulum multiflorum homogamum; flor. tubulosis hermaphroditis 5-fidis. Recept. plantum alveolatum. Invol. imbricatum pluriseriale, nunc bracteis foliaceis cinctum, nunc nudum, squamis ext. brevioribus aristato-cuspidatis, intimis apice scariosis exaristatis. Stigma Vernoniæ. Achæmium nervis 10 prominulis angulatum sæpius glabrum. Pappus 1-serialis, setis crassissimis rigidis dense barbellatis, junioribus basi in annulum connatis, demum facile et segregatim deciduis. — Herbæ Indiæ aut suffrutices. Folia alterna petiolata dentata. Capitula sæpius ad apices ramorum solitaria, rarius corymbosa pedunculata. Flores ex sicco purpurei. — Genus affine Vernoniæ, Cyanopidi et Centrathero.

Sect. I. PHYLLOCEPHALUM Blum. *bijdr.* 888.

Capitula bracteis foliaceis amplis arcuè cincta. Achænia basi non sensim attenuata.

1. D. FRUTESCENS, caule suffruticoso erecto hirsuto, foliis breviter petiolatis oblongo-lanceolatis basi attenuatis apice cuspidatis serratis rugosis supra puberulo hirtis subtus pallidis pubescenti-tomentosis, capitulis paucis pedunculatis bracteatis. ♀ in uliginosis ins. Javæ, inter montes Burangrang et Tankawan Prah, prov. Krawang legit cl. Blume. *Phyllocephalum frutescens* Blum. l. *bijdr.* 888. Bractææ utrinquè hirsutæ subtus albidæ sessiles lanceolatæ, ext. capitulo longiores. Fol. 3-4 poll. longa, pollicem lata. (v. s. comm. à cl. Blume.)
2. D. RETICULATUM (DC. in *Wight contrib. bot. ind.* p. 3), caule suffruticoso? erecto ramoso setis undiquè scabro, foliis sessilibus ovatis mucronatis et mucronato-subdentatis supra asperis subtus dense albo-tomentosis, nervis venisque scabris reticulatis, pedunculis paucis axillaribus terminalibusve, capitulis bracteis pluribus foliaceis arcuè cinctis, invol. squamis internis chartaceo-scariosis glabris bractæas superantibus. ♀ in Ind. orient. ad montes Neelgerry legit cl. Wight. *Ampherephes Indica* Wall. l. cat. et herb. n. 3956. comp. n. 66. Jacea? vulgo Kadourou Lesch. l. herb. n. 290. Capitula multiflora. Achænia glabra 10-costata. Pappus 1-ser. deciduus. (v. s.)

3. *D. ALLE*, caule herbaceo erecto glabriusculo aut hinc inde setoso-scabro apice tomentoso, foliis in petiolum attenuatis ovato-lanceolatis acuminatis grossè et irregulariter serratis suprâ setoso-scabris glabrisve subtus albotomentosis, pedunculis axillaribus 1-cephalis, bracteis foliaceis capit. approximatis oblongis subtus tomentosis, invol. squamis internis chartaceo-scariosis glabris subacutis. — In Ind. orient.

a. epileium, foliis suprâ lævibus. — Ad Kortallum distr. Carnatici leg. cl. Hayne et Wight. Decan. epileium DC. in Wight contrib. bot. ind. p. 7. Ampherephis mollis Wall. ! cat. et herb. n. 3957. comp. 67. Wightia formosa Spreng. in litt. ad Wall. Folia 3-3½ poll. longa, sesquipoll. lata. (v. s.)

ß. scabridum, foliis suprâ scabris. — Wight cat. n. 1392. — *D. scabridum* DC. l. c. (v. s.)

Sect. II. GYMANTHEMUM. — Gymnanthemum spec. *Cass. bull. philom.* 1817. dict. 20. p. 108.

Capitula aut omnino ebracteata aut bracteolis paucissimis invol. squamas inferiores referentibus subinstructa. Achænia elongato-turbinata seu basi attenuata.

4. *D. GLABRUM*, caule suffruticuloso tereti glabro striato, foliis breviter petiolatis ovalibus basi cuneatis apice acuminatis distanter repando-serrulatis glabris cum nervis utrinquè tenuè puberulis, corymbis 2-5-cephalis, invol. squamis adpressis acutis (non aristatis) flosculis multò brevioribus. 5 in Ind. orient. Conyza Wall. ! cat. n. 3283. comp. n. 393. Eupatorium glabrum Heyn. ! herb. Folia 2-3 poll. longa, 8-10 lin. lata. Pappus rufus. Achænia teres 10-costatum basi attenuatum glabrum 2 lin. longum. (v. s. in h. Cur. Ind. or. nunc Soc. lin. Lond.)

5. *D. SILBERTÆÆ*, caule herbaceo erecto ramoso, foliis breviter petiolatis oblongo-lanceolatis utrinquè acuminatis remotè aristato-serratis suprâ ramisque glanduloso-scabridis subtus secus nervos scabridis, capitulis ad apices ramorum subsolitariis subcorymbosis, invol. squamis inter. oblongis mucronatis, exter. filiformi-subulatis subpatulis nonnullis secus pediculum ortis. 4 in Ind. or. prov. Silhet ad montes Pundoa legit Fr. de Sylva. Vernonia? bracteata Wall. ! cat. n. 2921. comp. n. 31. Achænia 10-costatum basi attenuatum glabriusculum 2 lin. longum. Pappus rufus 1-serialis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

6. *D. DENDRIGULÆÆ* (DC. in Wight contrib. bot. ind. p. 7), collo tuberoso suffruticuloso, caule herbaceo erecto parè ramoso tereti basi striato pubescente apice tomentoso, foliis subsessilibus latè ovatis subrotundisve coriaceis dentatis suprâ glabris asperulis subtus densè albo-tomentosis, summis ovalibus acutis parvis, corymbo fastigiato composito, invol. ovatis squamis ovalibus mucronatis imbricatis extus tomentoso-lanatis intus glaberrimis. 4 in Indiæ Peninsula montibus Dendigul legit cl. Wight. Conyza indica Wall. ! cat. n. 3042. comp. 152. non Blam. Folia inf. 4 poll. longa, 3 poll. lata. Achænia nervis 10 geminatis elevatis angulato-striatum glaberrimum inter nervos glandulosum. Pappus 1-serialis rigidus (v. s. in h. Wight. et comm. ab hon. Cur. merc. angl. Ind. or.)

7. *D. ? OBOVATUM*, caule suffruticoso scandente tereti, foliis petiolatis obovatis obtusis integerrimis suprâ glabris subtus ramisque pube minimâ subpubescentibus, paniculâ ovatâ, invol. squamis ovato-oblongis brevissimè mucronulatis. 5 in Rawak Moluccensium. Gymnanthemum obovatum Gaudich. ! voy. uran. bot. p. 471. Vern. cuneata Less. in linnæa 1831. p. 644. Folia in sicco margine subrevoluta 2-3 poll. longa, 12-15 lin. lata. Capitula 6-8-flora. Achænia glabrum versus apicem tumidum 10-costatum basi attenuatum, fortè in nostr. spec. sterile. Pappus rufus 1-serialis scaber. An omnino priorum congener. (v. s. comm. ab am. Gaudichaud.)

8. *D. GRANDE*, fruticoso glabrum, ramis teretibus apice summo pulverulento-puberulis, foliis longè petiolatis ovalibus utrinquè acuminatis integerrimis

sabitu secus nervos minutissimè puberulla, corymbo composito polycaphalo laxo fastigiato, invol. ovati squamis adpressis glabris, ext. acuminatis int. obtusis fere truncatis, acheniis glabris, 5 in insula Madagascar orâ occidentali circa pagum Majangay in sinu Bombatoc legit cl. Bojer! Vernonia grandis Bojer! in litt. 1835. Frutex 15-18-pedalis. Flores lilacini, in sicco albidii, in invol. 9-10. Pappus rutilans x-serialis crassiusculus. Achen. junctus teretiusculum. Valde affinis D. Senegalensi. (v. s. sp. comm. à cl. Bojer. et in ins. Maurit. ex sem. Bojerianis cult. comm. à cl. L. Bontou.)

9. *D. FIMBRILLIFERUM*, fruticosum, ramis tomentoso-pubescentibus, ultimis seu floralibus compressis, foliis obovato-lanceolatis breviter acuminatis in petiolum attenuatis vix apice subdenticulatis suprà glabris subtus nervum pubero-hirtellis, corymbo amplo composito aphylo, capitulis pedicellatis obovato-cylindraceis 3-4-floris, invol. pauciserialis squamis ovatis pubescentibus inter. longioribus, recept. breviter fimbriifero, acheniis subhispidulis, pappi seriebus satis inæqualibus. 5 in insula Borbonia sylvis frequens (Bontou!). *Gymnanthemum fimbriiferum* Cass. dict. 20. p. 109. *Vernonia fimbriifera* Less. in linnæa 1829. p. 313. *Eupatorium coriaceum* Vahl symb. 3. p. 95. non Spreng. Cl. Cassini dixit fl. flavos, sed aspitur dixit flores omnes decoloratos. Pappus intensè rufus. (v. s. sine cor. comm. à cl. Bontou.)

10. *D. SENEGALENSIS*, fruticosum, ramis subangulosis tomento brevissimo subvelutinis, foliis petiolatis ovato- aut oblongo-lanceolatis subintegerrimis suprà glabriusculis subtus villosopubescentibus, corymbo fastigiato nudo, invol. campanulati squamis ovatis ciliatis, ext. brevioribus macrone acutis, inter. longioribus obtusis demum stellatis, achenio maturo glabro 10-costato inter costas glanduloso. 5 in Senegaliâ (Roussillon! Sparman!) *Baccharis Senegalensis* Pers. ench. 2. p. 424. *Gymnanthemum cupulare* Cass. dict. 20. p. 109. *Eupatorium coloratum* Willd. sp. 3. p. 1768 ex Less. *Vernonia Senegalensis* Less. in linnæa 1829. p. 265. 1831. p. 641. non Desf. *Conyza rutilans* Poir. suppl. 2. p. 341? *Chrysocoma amara* Schum. in act. Dan. p. 157. ex Less. Cor. alba. Pappus rufo-aureus rigidus scaber. Huc referendæ mihi videntur: *Conyza radiata* in argillois circa Albreda ad fl. Gambie, C. aurea in insulâ prope Saint-Louis, et C. frutescens in peninsulâ Promontorii-Viridis prope Dakar, à cl. Perrottet collectæ et mecum commun. Variat squamis invol. plus minus ciliatis, imò subciliatis, foliis subrotundis, obovatis, obovato-oblongis aut oblongo-lanceolatis. (v. s.)

11. *D. AMYGDALINUM*, fruticosa, ramis teretibus apice pube brevi scabrinusculis, foliis oblongo-lanceolatis petiolatis utrinquè acuminatis, adultis glabris, corymbi paniculati ramis dichotomis, capitulis pedunculatis, involucri semiglobosi squamis obtusis, achenio villosopubido 10-nervio, 5 in Nubiâ ad Fazogl, ubi dicitur Kering et arabicè Kikir. *Vernonia amygdalina* Delil. in Cail. voy. et cent. n. 21. forsan in Sierra-Leonâ si specim. quoad patr. incertum huc ritè refero. Ex eo affinis videtur D. Senegalensi. (v. s.?)

12. *D. DIVERGENS* (DC. in Wight contrib. bot. ind. p. 8), caule herbaceo erecto tenuissimo velutino scabro paniculato ramoso, foliis brevissimè petiolatis ellipticis utrinquè acuminatis dentatis suprà glabris subtus puberulis reticulatis subrugosis, paniculæ ramis elongatis divergentibus foliosis ipsis paniculatis, capitulis ad apicem ramulorum confertis 7-8-floris, invol. squamis oblongis acutè mucronatis glabriusculis, achenio glabro glanduloso. 2 in Indiâ orient. verosim. Maderaspatanâ (h. Wall.!), in Pegu (Belanger!) *Conyza divergens* Wall. cat. n. 3027. comp. 137. excl. litt. E. *Vernonia* Wight herb. 2 Affinis D. aspero. Pappus rufus, ext. brevior paucisetus caducissimus. Folia 3 poll. longa, pollicem lata. An potius Vernoniæ sp. V. multifloræ proxima? (v. s.)

fl. lanceolata, foliis planis vix subrugosis capitulis paulò minoribus. — in regione Birmanorum ad Taong-Dong legit cl. Wallich. *Conyza lanceolata* Wall. cat. et herb. n. 3059. comp. 189. cat. Birm. n. 916: Panicula minus divaricatâ (v. s. comm. cum priore ab. hon. Cur. mere. angl. Ind. or.)

XX. CYANOPIIS Blum. in litt. 1831. — *Cyanthillium* Blum. (1826) *bijdr.* p. 889. — *Isonema* Cass. *bull. philom.* 1817. *dict.* 24 p. 25. non Brown. (1810). — *Centratheri* sect. Less. in *linnæâ* 1829. p. 320. — *Vernonia* spec. Less. *syn.* 146. — *Conyza* spec. Ait. *Lam.*

Capitulum multiflorum homogamum, fl. omnibus tubulosis inæqualiter 5-fidis hermaphr. Recept. alveolato-subfimbriiferum planum. Invol. hemisphærici squamæ pluriseriales imbricatæ lanceolatæ rigidularistatæ apice non adpressæ. Stigm. Vernoniæ. Achænium prismatocypentagonum glabrum, areolâ terminali latâ subcupulatâ, Pappus 1-serialis, setis rigidis imâ basi distinctissimis deciduis subscabris. — Herbæ Indicæ erectæ ramosæ verisim. annuæ. Folia petiolata ovata aut lanceolata alterna suprâ glabra subtus plûs minûs villosa. Capitula ad apices ramorum pauca pedicellata ebracteata corymbosa. Flores cœrulescentes, undè nomen ab ipso cl. auct. paulò mutatum ut habitum melius exprimat. — Genus Vernoniæ affine.

1. *C. VILLOSA* (Blum. *bijdr.* 889 sub *Cyanthillio*), foliis petiolatis ovato-oblongis infrâ caescenti-villosis margine subrepando-serrulatis, summis integerrimis, corymbis 3-4-cephalis, involucri floribus subæquali. ☉ in graminosis insulæ Javæ circa Baitenzorg. Proximè fortè nimis ad sequentem accedit. (v. s. comm. à cl. Blume.)
2. *C. PUBESCENS* (Blum. *bijdr.* 890 sub *Cyanthillio*), foliis petiolatis ovatis utrinquè acutis subtus pubescentibus margine repando-mucronulato-denticulatis, corymbis 3-4-cephalis, involucri floribus subæquali. ☉ in graminosis insulæ Javæ cum priori legit cl. Blume! in Manillâ et Guahan (Cham.), in Chinâ (Ait.), ins. Marianis (Hænk!), in Chinâ et Pegu (Less.), in Pegu ex cl. Belanger, ad Tavoy et Rangoon Ind. Birmanica ex cl. Wallich. *Conyza punctulata* Vahl! cat. et herb. n. 2995. comp. 105. *birn.* 1167. *Synantherea* Wall! cat. n. 7136. *Conyza patula* Ait. Kew, 3. p. 184. *Conyza Chinensis* Lam. *dict.* 2. p. 83. non Linn. nec Lour. *Isonema ovata* Cass. *dict.* 24. p. 25. *Serratula* Mill. ic. t. 247. *Centratherum Chinense* Less. in *linnæâ* 1829. p. 320. *Vernonia Chinensis* Less. in *linnæâ* 1831. p. 6-4. Achænia glabra obpyramidata 4-5-angulata. Variat invol. glabrato aut pubero, foliis subtus plûs minûs pubescenti-villosis. Specim. Indiæ contin. ferè media inter spec. hujus et prioris Blumeana et fortè in unam colligunt. (v. s.)

β. microcephala, capitulis dimidio minoribus. ☉ in insulâ Mauritiî introd. et quasi indigena facta ex Bojer in litt. An species propria? (v. s.)

3. *C. MOLUCCENSIS* (Blum. *bijdr.* 890 sub *Cyanthillio*), foliis breviter petiolatis lanceolatis utrinquè acuminatis subtus albido-villosis margine repando-mucronato-dentatis, capitulis geminis confertis, invol. squamis glaberrimis longè aristatis flosculos subæquantibus. ☉? in insulâ Moluccanis. Fol. 3 poll. longa, pollicem lata. (v. s. comm. à cl. Blume.)
4. *C. MADAGASCARIENSIS*, caule herbaceo erecto divaricato-ramoso tereti glabro, foliis ellipticis utrinquè acuminatis grossè serratis basi integerrimis cuneatis subpetiolatis sparse scabridis subtus punctulatis, capitulis pedicellatis ebracteatis, invol. squamis glaberrimis acuminatis, achæniis seriato-sulcatis glabris, pappo pauciseti caducissimo. — in insulâ Madagascar in pratis et ad margines sylvarum legit cl. Bojer. *Ethulia conyzoides* Bojer! in litt. 1835. Flores rosæ (v. s. comm. à cl. inv.)

† *Species exclusa.*

C. SALICOIDES DC. = *Vernonia erigeroides*.

XII. CENTRATHERUM Cass. *bull. philom. febr. 1817. dict. sc. nat. 7. p. 383. journ. phys. 89. p. 24.* — *Ampherephus H. B. et Kunth nov. gen. am. 4. (1818). p. 24. Less. syn. p. 148.* — *Spixia Schrank pl. rar. h. mon. 2. (1819). t. 80. non Leand.* — *Centratheri sect. 1. Less.* — *Amphirephidis sect. 1. Nees et Mart. act. soc. cur. nat. 12. p. 4.*

Capitulum multiflorum æqualiflorum. Involucrum imbricatum campanulatum quasi duplex nempe, foliis floralibus magnis cinctum, foliolis inv. veri siccis exter. brevioribus. Receptaculum planum nudum. Cor. regularis 5-fida, tubo gracili lobis acuminatis glabris longiore. Anthæræ inclusæ. Filam lævia. Styli rami semisubulati. Achænium oblongum subcylindraceum glandulosum glabrum, collo apicali parum prominulo. Pappus 1-serialis, setis serrulatis inæqualibus rigidis caducissimis! — Suffrutices aut herbæ. Folia alterna ovata aut oblonga serrata. Capitula ramosa terminantia, pedunculata. Flores purpureo-cœrulei. — Cl. Cassini in *journ. phys. 1. c.* asserit *Centratherum* idem esse ac *Ampherephum*, et in *tabl. comp. 1829.*, n. 36 et 37, genera duo admittit ex invol. squamis muticis aut spinoso-aristatis.

1. *C. INTERMEDIUM* (Less. in *linnæa 1829. p. 320*), caule herbaceo, involucri veri squamis integerrimis, exterioribus apice aristatis, interioribus ovato-oblongis obtusis. ☉ in *Brasiliâ* (Mart.), in cultis incultisque ad *Bahiam* (Salzm. ! Blanch. !). *Ampherephus intermedia* Link *abbild. 5. t. 29. Colla hort. rip. app. 3. p. 26. Spixia violacea* Schrank *pl. rar. h. mon. t. 80. Flores cœruleo-subviolacei. Cor. tubus puberulus. (v. v. cult. in h. Genev. et s. sp. è Bahiâ.)*
2. *C. PUNCTATUM* (Cass. *dict. 7. p. 384. Less. l. c.*), caule fruticoso, involucri veri squamis supernè eroso-denticulatis apice spinoso-dentatis. § in *ripâ flum. Orinoci ad pagum Maypures. Ampherephus aristata* H. B. et Kunth *nov. gen. 4. p. 25. t. 314. Cor. purpurea et ex cl. auct. glabra. Folia lanceolata. (v. s.)*
3. *C. MUTICUM* (Less. *l. c.*), caule fruticoso, invol. veri squamis denticulatis apice rotundatis mucronatis. § in *temperatis Novæ-Andalusie ad montem Cocollar. Ampherephus mutica* H. B. et Kunth *nov. gen. 4. p. 26. t. 315. Folia oblonga basi cuneata-angustata.*

† *Species exclusæ.*

C. CHINENSE Less. = *Cyanopsis pubescens.*

AMPHEREPHUS INDICA Wall. = *Decaneurum reticulatum.*

AMPH. MOLLIS Wall. = *Decaneurum molle.*

AMPH. PSILOCARPA N. et Mart. = *Oiospermum involueratum.*

AMPH. WIGHTIANA Wall. = *Oiospermum Wightianum.*

XIII. BECHIUM.

Capitula multiflora homogama, floribus tubulosis 5-fidis, ext. subpalmatis. Recept. alveolare. Invol. squamæ 2-3-seriales subimbricatæ longitudine subæquales, inter. angustiores glabræ apice coloratæ. Styli rami elongati hispidi. Achænium teretiusculum villosum. Pappus uniserialis pilosus scabridus. — Herbæ Madagascarienses. Folia alterna aut radicalia ovali-oblonga subsessilia subintegerrima suprâ piloso-scabra. Capitula terminalia corymbosa. Invol. squamæ apice subsca-riosæ purpureæ. Flores purpurei. — Nomen apud Plinium olim *Tussilagini* impositum admisi.

1. *B. RUBICAULE*, caule pubescente imâ basi hirtello ad medium folioso, foliis ovali-oblongis basi cuneatis subpetiolatis utrinquè puberulis, corymbo composito polycephalo, invol. squamis dorso villosio-hirtis. 2. in *insulæ Mada-*

gascar prov. Emirnensi legit cl. Bojer. Herba pedalis. Caulis purpureus. Folia 3 poll. longa $\frac{1}{2}$ poll. lata obscure virentia. Corymbi rami dense-villosi. (v. s. comm. à cl. inv.)

2. B. SCAPIFORME, caule nudo pube sublanatâ et pilis apice glandulosis in lanâ intermixtis hispido, foliis radicalibus ovali-oblongis basi cuneatis subsessilibus suprâ hirtis scabris subtus cania sericeo-tomentosis, capitulis circ. 3 terminalibus subsessilibus, invol. squamis dorso glanduloso-hirtis. $\frac{1}{2}$ in montibus nudis et siccis prov. Emirnensis Madagascariensium legit cl. Bojer. (v. s. comm. à cl. inv.)

XIV. STOKESIA Lher. sert. 27. DC. ann. mus. 16. p. 154. Cass. dict. sc. nat. 51. p. 64. Less. in linn. 1829. p. 321. syn. p. 148. — *Cartesia* Cass. bull. philom. 1816. p. 198. — *Carthami* sp. Hill.

Capitulum multiflorum radiatiforme. Involucrum subglobosum basi bracteatum pluriseriatum imbricatum, squamis adpressis, exterioribus ovalibus in append. foliaceam patulam ciliato-spinulosam desinentibus, interioribus oblongis ciliolatis. Recept. carnosum nudum. Cor. palmarum glandulis conspersæ, exteriores capituli majores intus profundius fissæ. Anth. inclusæ. Styli rami semi-subulati. Achæmium brevè tetra- (rarius tri-) gonum, areolâ terminali latâ 4-gonâ. Pappus 1-serialis 4-5-paleaceus, paleis deciduis longis planis scariosis integerrimis. — Herba erecta ramosa glaucescens glabriuscula, caule apice villosotomentoso. Folia alterna, infer. ovalia integerrima in petiolum longè attenuata, super. semi-amplexicauliâ lanceolata ad basin serrato-spinosa. Capitula ramos terminantia solitaria. Flores cœrulei.

1. S. CYANEA (Lher. sert. angl. 28). $\frac{1}{2}$ in Carolinâ australi. Videtur rara et à cl. Elliott non reperta. *Carthamus lævis* Hill. Kew. 57. t. 5. *Cartesia centauroides* Cass. bull. 1816. p. 198. *Carthamus Carolinianus* Michx. in h. Mus. Par. (v. s. ex herb. Lher.)

XV. PLATYCARPHA Less. in linnæâ 6. p. 688. syn. p. 148. — *Cynara Thunb.* fl. cap. 622. non Linn. — *Stobæa* sp. Spreng.

Capitulum homogamum multi- et æqualiflorum. Involucri squamæ pluriseriales lanceolatæ elongatæ submembranaceæ integræ acutæ, interiores angustiores recept. paleas ferè similes. Cor. tubo longissimo extus præsertim ad apicem hispidulo, lacinii 5-linearibus æqualibus. Stam. filam. glabra lævia. Anthere apice obtusæ et sub-æqualiter appendiculatæ basi ecaudatæ. Pollen globosum. Stigm. 2 longa apice divergentia teretiuscula undiquè ferè papillosa. Fructus glaber subcylindraceus. Pappus persistens paleis 7-9 scariosis albis linearibus acuminatis integerrimis constans. — Herba perennis. Folia radicalia longa petiolata pinnatipartita subtilis albo-tomentosa suprâ et in petiolo arachnoidea, lobis ovatis dentatis, inferioribus dissitis parvis, extremis majoribus confluentibus. Capitula plurima ad collum sessilia congesta. Cor. violacææ. — Genus distinctissimum!

1. P. GLOMERATA. $\frac{1}{2}$ ad Cap. Ronæ Spei (Th. Dreg!) in distr. Uitenhagen (Eckl. et Zeyh.) juxta Swartkopp' rivier. *Cynara glomerata* Thunb. prodr. cap. 141. flor. cap. 622. Willd. sp. 3. p. 1692. *Stobæa glomerata* Spreng. syst. 3. p. 394. (v. s.)

XVI. ODONTOCARPHA, — Gen. nov. Propp. n. 2.

Capitulum pluriflorum homogamum. Recept. angustum æbracteolatum. Invol. oblongo-cylindricum, squamis imbricatis adpressis. Cor. tubus pappo brevior, faux oblonga tubo latior, limbus 5-lobus acu-

finisculus. Stam. filam. glabra? brevior; antheræ ecaudatæ. Styli rami breviter exserti subulati hispiduli. Achænium villis sericeis adpressis brevibus munitum subcylindraceum striatum. Pappus 1-serialis persistens, paleis 7-8 lanceolatis acutis apice serrulatis. — Herba suffruticulosa gracilis Chilensis erecta ramosa, ramis apice 1-cephalis. Folia alterna angustè linearia acutiuscula enervia præsertim ad basin ciliolata. Capitula foliis parvis raris substipata oblonga. Cor. vix exsertæ ex sicco purpurascens. Invol. fortè viscosa. — Car. ferè Platycarpæ, sed habitus diversissimus, cor. tubus brevissimus, pappi palea denticulatæ, etc.

1. O. POEPPIGII. 3 in Chili legit cl. Poeppig. Folia pollicem longa semilineam lata. Capitula 3-4 lin. longa. (v. s.)

XVII. WEBBIA.

Capitula pluriflora discoidea dioica! Recept. alveolatum, alveolorum marginibus subexsertis erosis. Invol. imbricatum 3-5-seriale disco brevius. — ♂ Cor. tubulosæ breviter 5-dentatæ. Antheræ inclusæ lineares. Stylus indivisus aut apice subfissus. Ovarium extus villosum abortivum. Pappus pauciserialis, setis vix dentatis. — ♀ Cor. tubulosæ in lobos 5 elongatos angustos crassiusculos divisæ. Anth. 0, aut effatæ. Stylus è tubo exsertus, lobis corol. brevior, ramis 2 elongatis crassiusculis extus et styli partè indivisâ supremâ pilosiusculis. Achænium teres seu turbinatum pluricostatum erostre, costis villosissimis, sulcis glandulosus. Pappus pluriserialis, setis crassiusculis densè barbellatis exter. brevioribus. — Herbæ suffruticosæ Africanæ, caule erecto basi simplici apice corymboso. Folia sparsa linearia marginibus parçè revoluta suprâ glabriuscula subtus cum caule involucrique canescentia. Flores purpurei in capitulo 8-10. Pollen flavum. Stigmata purpurea. — Dixi in honorem cl. bot. angl. P. B. Webb, Orientis, Africæ borealis et insularum Canariensium strenuo observatori.

1. W. SERRATULOIDES, caule suffruticoso ramoso pubescente, capitulis ad apices ramorum solitariis hemisphæricis, invol. squamis adpressè imbricatis 5-seriatis, 3 in Senegambiæ locis arenosis ad Albreda secus flumen Gamble Martio ferè defloratam legit cl. Perrottet. Folia vix pollicem longa, semilineam lata. Capitulum 6 lin. diam. Serratulæ aut melius Jurinæ capitula referens. (v. s. specim. form. à cl. Perrottet comm.)
2. W. ? SMITHIANA, caule herbaceo striato sericeo-argenteo simplici dense foliato, foliis linearibus subtus sericeo-incanis, paniculâ subfastigiata, invol. campanulati squamis 2-3-seriatis 3-nerviis emucronatis, achænio pubescente, pappi ser. ext. dimidio interiore brevior. 2 ad fl. flavum Congo. Vernonia Smithiana Less. in linnæa 1831. p. 638. W. pinifolia dicitur proxima.
3. W. PINIFOLIA, caule herbaceo simplici striato subcanescente, foliis sessilibus confertis 1-nerviis linearibus margine revolutis suprâ glabris subtus sericeo-incanis, corymbo polyccephalo fastigiato, invol. squamis oblongo-ovatis subcanescentibus, achænio sericeo-piloso, pappi serie externâ dimidio interiore brevior. 2 ad Cap. B. Spei (Krebs!) in montibus Langekloof et Krumrivier (Th. Eckl. n. 352. et Zeyh.), in Coloniz distr. orientalibus (Burch.) cat. geog. n. 3840 et 7434, ad Willbergen, Zw. Key et Basche (Dreg.) in Cafferland et Ultenhagen (Eckl. et Zeyh.). Conyza canescens Thunb. fl. cap. 665. Conyza pinifolia Lam. dict. 2. p. 86. excl. syn. Seb. (1). Erigeron Capense Houtt.!

(1) Icon enim Sebana (mus. r. t. n. 3. f. 3) hic à Lamakio citata non malè habitum nostræ plantæ indicat, sed ex involucri ad Helichrysum intermedium referenda videtur.

- pf. syst. x. hl. 629. t. 69. f. 2. ex herb. Deless. *Vernonia pinifolia* Less. in Linnaea 1829. p. 257. 1831. p. 639. Caulis sesqui- aut bipedales. Fol. 2-3 poll. longa, 1-2 lin. lata, 1-nerviâ. Capitula 10-20 laxè corymbosa. (v. s.)
4. *W. ARISTATA*, caule herbaceo simplici striato subcanescente, foliis subsessilibus lanceolatis acuminatis basi attenuatis penninerviis integerrimis planis utrinquè aut subtus tantum pube confertâ sericâ adpressâ canis, corymbo polycephalo fastigiato, invol. squamis subcanescentibus lanceolatis in acumen aristæforme productis, achenio sericeo-villoso, pappi serie externâ triplò interiore brevior. \mathcal{L} ad Cap. Bonæ Spei ad Katrivier (Dreg. sp. exs.), ad Zw. Key et Basche (Dreg.!), in territorio cesso et Kafferland (Eckl. ! et Zeyh.). Affinis *W. pinifolia* sed folia ferè *Salicis* albæ nunc utrinquè nunc subtus tantum sericea. (v. s.)
5. *W. ELÆAGNOIDES*, caule herbaceo erecto striato subcanescente, foliis breviter petiolatis ellipticis acutis planis integris suprà glabris subtus pube sericâ adpressâ confertâ canis, corymbo polycephalo fastigiato, invol. squamis canescentibus oblongis mucronato-acutis, achenio villosio, pappi serie externâ triplò interiore breviori. \mathcal{L} in Africâ australi ad Zw. Omsamwubo et Omsanculo legit cl. Dregel! Folia 18-20 lin. longa, 9-10 lata. (v. s.)
6. *W. OLIGOCEPHALA*, caule herbaceo simplici striato glabriusculo, foliis breviter petiolatis ovalibus abruptè acuminatis integris planis penninerviis suprà glabris subtus pube sericâ confertâ adpressâ canis, corymbo 3-cephalo, invol. squamis subcanescentibus lanceolatis acutis, achenio tereti villosio, pappi serie externâ triplò interiore breviori. \mathcal{L} in Africâ australi ad Zw. Omsamwubo et Port Natal legit cl. Dregel! Priori affinis sed achenium duplò longius. Caulis multò tenuior. Folia pollicem longa, 8-10 lin. lata, etc. (v. s.)
7. *W. HIRUTA*, caule herbaceo simplici striato breviter hirsuto, foliis sessilibus oblongis basi cordatis apice mucronato-acuminatis subserratis suprà glabriusculis subtus hirsutis reticulato-nervosis, corymbo polycephalo conferto ramulis hirsuto-tomentosis, invol. squamis subcanescentibus lanceolatis acutis, acheniis villosis, pappi serie brevissimâ. \mathcal{L} in Africâ australi ad Tambukiland et Cafferland (Eckl. ! n. 725 et 1318), ad Katriviersberg et Port Natal (Dreg.!). Antheræ in fl. scem. effatæ. (v. s.)
8. *W. NUDICAULIS*, caule herbaceo subnudo striato glabriusculo, foliis radicalibus obovatis basi attenuatis mucronulatis glabris penninerviis planis subdenticulatis, canlinis remotis linearibus erectis margine revolutis integerrimis, corymbo laxo, invol. squamis canescentibus oblongis mucronatis, acheniis villosis, pappi serie externâ brevissimâ. \mathcal{L} in Africâ australi ad Zw. Key et Basche legit cl. Dregel! Folia radic. $1\frac{1}{2}$ poll. longa, 9-10 lin. lata; caulina ferè pollicem longa, vix lineam lata. Flores ferè *W. pinifolia*. (v. s.)

XVIII. HOPLOPHYLLUM. — *Pteronia* sp. *Lin. f. Thunb.*

Capitula 5-flora homogama. Invol. ovato-oblongi squamæ latæ imbricatæ obtusæ apice spinuloso-mucronatæ. Recept. angustum fimbriatiferum. Cor. glabræ tubulosæ profundè in lobos lineares æquales 5-fissæ. Antheræ ecaudatæ. Filamenta brevina glabra. Styli à medio setis erectis hispidi; ramis longissimis per totam longitudinem hispidis. Achenia crassa subtrigona villosissima apice glabra et sub pappo paululum contracta. Pappus multiserialis persistens, squamellis rigidulis exter. lineari-filiformibus, inter. basi membranaceis latioribus sensim acuminatis. — Suffrutex Capensis glaber rigidus, ramis teretibus striatis. Folia alterna rigida spinescentia integerrima aut hinc indè ad basin spinuloso-dentata crassiuscula patula utrinquè ut rami regulariter striata. Capitula in axillis supremis sessilia solitaria sed ob folia approximata 4-5 congesta capitata. Cor. pallida.

1. *H. spinosum*. ζ in Africâ Capensi in Carro (Thunb.), ad Olifanttrivier in regione Gariëpinâ (Dreg!), ad Clan William (Eckl. n. 1420. et Zeyh.). *Pteronia spinosa* Linn. f. suppl. 357. Thunb. fl. cap. 629. (v. s.)

XIX. PIPTOCOMA Cass. bull. philom. 1817. p. 10. 1818. p. 50. dict. sc. nat. 41. p. 111. Less. in linnæâ 1829. p. 315. syn. p. 147.

Capitulum pluriflorum æqualiflorum. Involucrum ovato-cylindricum imbricatum, foliolis coriaceis siccis ovato-aut obovato-oblongis. Recept. nudum angustum. Cor. regularis glabra, tubo longo, limbo 5-fido, lobis acuminatis glandulosis parte integra brevioribus. Anth. inclusæ. Achæ-nium obcompressum angulatum glabrum, disco epigyno magno, nectario styliformi. Pappus biserialis (ferè *Heterocomæ*), serie externâ brevissimâ cartilagineâ crenatâ, interiori ∞ -paleacea, paleis caducis longis planiusculis subserratis sæpè tortis. — Suffrutices Americani. Rami numerosi teretes tomentosi. Folia alterna integra 1-nervia suprâ glabra subtus tomentosa. Capitula plurima ad apices ramorum subsessilia aggregata in glomerulum subumbellatum. Flores purpurei aut rosei.

1. *P. ruvescens* (Cass. ll. cc.), foliis brevè petiolatis ovatis suprâ scabris subtus subreticulato-tomentosis, capitulis 12-floris. ζ in Sancto Domingo. *Achænia* 5-gona striatâ. Cor. purpureæ.

2. *P. lychnophoroides* (Less. l. c. f. 7. 8. 10. 12), foliis sessilibus linearibus suprâ lucidis subtus tomentosis margine revoluta, capitulis 3-4-floris. ζ in Brasiliâ tropicâ. *Vernonia trichocarpa* Spreng. syst. 3. p. 437. ex Less. Folia ferè Phyllicæ, 6 lin. longa lineam lata, coriacea. Glomeruli foliis cincti. Cor. roseæ. *Lychnophoræ* fortè nimis affinis. (v. s. comm. à Mus. reg. Berol.)

XX. DISTEPHANUS Cass. bull. philom. 1817. 76. dict. 13. p. 361. — *Vernoniæ* sp. Less. syn. 146.

Omnia *Vernoniæ*, sed involucri squamæ apice appendiculatæ. Invol. areolatum, areolis in dentes seu fimbrias breves exsertis. Pappi series utraq; constans paleis circ. 10-lamellatis coriaceis, exterioribus nunc interiores subæquantibus nunc dimidio brevioribus latioribus denticulatis, interioribus linearibus barbellatis. — Frutices Mauritanii seu Madagascarienses, ramulis foliisque tomentosis alternis integerrimis. Capitula densè corymbosa.

§. 1. *Pappi serie externâ dimidio quam interior breviori.* — Mauritaniani.

1. *D. populifolius* (Cass. l. c.), foliis petiolatis latè ovatis acuminatis, corymbo 7-8-cephalo, invol. squamis dorso præsertim ad margines lanatis, capitulis 30-40-floris. ζ in insulâ Mauritiî (Comm. ! Bory !), locis aridis elatis ventosis (Bouton !) ferè semper florens (Bojer !). *Conyza populifolia* Lam. ! dict. 2. p. 87. Willd. sp. 3. p. 1943. *Vernonia populifolia* Spreng. syst. 3. p. 434. Less. in linnæâ 1829. p. 256, et 1831. p. 673. Rami adulti atro-fusci teretes. Ramuli angulati densè albo-tomentosi. Folia ovato-cordata utrinquè tomentosa, 3 poll. longa (excl. petiol. semipollic.), 2 poll. lata. Capitula densè corymbosa. Involucri squamæ extus in parte liberâ lanato-villosæ. Cor. glabræ. Achæ-nium sericeo-villosum. (v. s.)

§. 2. *Pappi serie externâ interiorem subæquante.* — Madagascarienses.

2. *D. capitatus* (Bojer ! in litt. 1835), foliis subsessilibus lineari-lanceolatis basi attenuatis suprâ glabriusculis subtus adpressissimè cano-tomentosis, corymbo circ. 4-cephalo congesto, capitulis subsessilibus 20-25-floris, invol. squamis

acuminatis vix tomentosis. \S ad fossarum margines circa urbem Tananarivou insulæ Madagascar legit cl. Bojer. Frutex 2-3-pedalis. Rami virgati striati. Folia $1\frac{1}{2}$ poll. longa 3 lin. lata. Recept. eleganter areolatum, areolis inæqualiter dentatis medio punctigeris. Pappus rufus. (v. s. comm. à cl. Bojer).

3. D. TRINERVIS (Bojer! in litt. 1835), foliis breviter petiolatis ellipticis et super. oblongis utrinque tomentosis trinerviis, corymbo 5-7-cephalo conferto; capitulis pedicellatis panicifloris, invol. squamis acuminatis junioribus præsertim tomentosis. \S in insulâ Madagascar legit cl. Bojer. Frutex priori similis, sed paulò elatior. Rami angulati. Folia 9 lin. longa, 4 lin. lata. (v. s. comm. à cl. Bojer).

XXI. STROPHOPAPPUS. — Stilpnopappi sp. Mart.

Capitulum multiflorum. Invol. imbricatum, squamis ovali-oblongis obtusis extus apice subtomentosis. Recept. vix alveolatum. Pappus duplex, seriei utriusque ligulæ planæ acuminatæ nitentes elongatæ æquales. Achæmium villosissimum. — Frutex Brasiliensis, floribus non satis notis. Rami teretiusculi juniores tomentosi. Folia sessilia oblonga obtusa coriacea subintegerrima suprâ glabra subtus tomentosa. Capitula secus ramos ad axillas et apicem sessilia gemina folio breviora. †

1. S. BICOLOR. \S in adscensu montis Serra do Caraca prov. Minar. Gener. Brasiliæ legit cl. Martius. Stilpnopappus bicolor Mart.! herb. Folia 3 poll. longa, 8-9 lin. lata. Flores ign. Recept. vix alveolare ex statu imperfecto judicatum. (v. s. in h. Acad. reg. Monac.)

XXII. BLANCHETIA.

Capitulum pluri-(8-10) florum homogamum. Invol. ovati squamæ imbricatæ ovatæ mucrone acuminatæ. Recept. angustum epaleaceum. Cor. profundè et subæqualiter 5-fidæ. Anthæræ exsertæ basi obtusæ apice acutè appendiculatæ. Styli rami exserti hispidi acuminati. Achæmium glaberrimum obovato-oblongum subpentagonum striatulum. Pappi squamellæ circ. biseriales subæquales caducissimæ lineares rigidulæ subciliatæ. — Herba suffruticulosa Brasiliensis erecta alternè ramosissima. Caules rami et petioli tomento stellato brevi subrufescente et setis rigidis nigris patulis vestiti. Folia caulis alterna, ramorum steriliùm sæpius opposita petiolata ovali-oblonga basi longè cuneata subintegra aut hinc inde denticulata suprâ pube stellatâ parçâ notata subtus eodem tomento densè canescenti-rufa. Capitula in cymam umbellatam foliosam disposita subsessilia conferta ovata subaraneosa. Cor. ex sicco purpurascens. Pappus albus. — Genus habitu Stilpnopappo affine sed pappo diversum, dicavi cl. Blanchet, botanico Helvetico, qui, civitati Bahiæ dicatus, Brasiliæ res naturales strenuè collegit et plantam ipsam generis typum cum pluribus aliis detexit.

1. B. HETEROTRICHA. \S 2? in Brasiliâ ad Caxocira ab urbe Bahiâ lencis 15 dist. legit cl. Blanchet. (v. s.)

XXIII. STILPNOPAPPUS Mart. herb.

Capitula multiflora homogama. Invol. campanulatum squamæ circ. biseriales æquales acuminatæ. Recept. alveolatum fimbriiferum. Achæmium villosum. Pappus duplex uterque paleaceus planus nitidus, ext. paleis brevibus, int. paleis duplò longioribus vix exterioris numerum superantibus. — Herbæ annuæ Brasilienses. Ex pappo paulò accedit hinc ad Dialestam illinc ad Lychnophoram.

1. *S. TALCHOSPIROIDES* (Mart. ! herb.), caule herbaceo ramoso erecto puberulo, foliis oblongis basi attenuatis acutis subintegerrimis penninerviis planis supra glabris subtus tomentosis, pedunculis æcus ramos distantibus oppositifolius 1-cephalis nudis, capitulis 15-20-floris, invol. floribus brevioris biserialis squamis lanceolatis acuminatis foliaceis, ext. supra scabris subtus subtomentosis. — in Serra do Rio Contas prov. Bahiensis Brasiliz legit cl. Martins. Achænium villosum turbinatum 10-costatum. Pappus duplex persistens lacens albidus, ext. paleis 10 ovatis cuspidatis, int. triplo fere longior setis filiformibus scabris 15-20. Recept. fimbriatiferum. Cor. violaceæ. Capitulum circ. magnitud. Vern. anthelminthicæ. (v. s.)
2. *S. PRATENSIS* (Mart. ! herb.), caule herbaceo erecto tereti villosa, foliis ovali-oblongis acutis basi attenuatis integerrimis supra glabrisculis subtus cano-tomentosis, infer. paulo latioribus, cymæ ramis scorpioides oligocephalis capitulis 30-floris lateralibus solitariis sessilibus folio suo multo brevioribus, invol. squamis 2-3-seriatis acuminatis parce villosiusculis, recept. alveolato fimbriatifero, achænio brevi crasso sericeo-villosissimo, pappi ser. utraq. paleis 10 constante, ext. ovatis brevioribus, int. linearilongatis acuminatis. ① in campis apricis prov. Piautiensis et Fernambucanæ frequentissimam legit cl. Martins. Cor. purpureo-violaceæ glabræ. Herba pedalis. Radix fibrosa. Folia 1 $\frac{1}{2}$ pol. longa, 5-6 lin. lata. (v. s. in h. Acad. reg. Monac.)
β. crotonifolia (Mart. ! herb.), caule pubescente elatiore, foliis linearibus aut lanceolato-linearibus. ① in cultis apricis prov. Bahiensis Brasiliz. (v. s. in h. Acad. reg. Monac.)
3. *S. TOMENTOSUS* (Mart. ! herb.), caule herbaceo erecto tereti subamplici tomentoso basi folioso apice subaphyllo, foliis sessilibus linearibus integerrimis utrinque vix acutis 1-nerviis supra araneosis subtus tomentosis, capitulis 40-50-floris, ad apices aut æcus ramos sessilibus folio flor. paulo brevioribus, invol. campanulati squamis ovali-oblongis acuminatis dorso villosis adpressis, recept. alveolato longe fimbriatifero, achænio sericeo-villosissimo, pappi serie utraq. paleaceâ, ext. acuminatâ. ① in aridis prope Caitete prov. Bahiensis Brasiliz legit cl. Martins. Folia 1 $\frac{1}{2}$ -2 poll. longa, 2-3 lin. lata. Cor. purpureæ. (v. s. in h. Acad. reg. Monac.)
4. *S. SCARPEUS*, herbaceus collo frutescente, foliis radicalibus confertis linearibus integerrimis coriaceis supra glabris subtus cano-tomentosis 1-nerviis, scapo nudo tereti pubescente simplici aut bifido, capitulis remotis lateralibus sessilibus ebracteatis, invol. squamis dorso pubescentibus ovatis, achænio sericeo-villoso, pappi paleis ext. ovatis, inter. linearibus. ② in Brasiliâ circa Bahiam legit cl. Blanchet (scapo simplici n. 1874 ! scapo bifido n. 1887 !). Folia pollicem longa 1-2 lin. lata. Scapus 2-9 poll. longus. (v. s.)

† *Species exclusa.*

3. *BICOLOR* Mart. = *Strophopappus bicolor*.

XXIV. *DIALESTA* H. B. et Kunth nov. gen. am. 4. p. 35. t. 320. Cass. tabl. XL. n. 40. Less. in Linnaea 1829. p. 315. syn. p. 146.

Capitulum 2-florum æqualiflorum. Involucrum imbricatum cylindraceum fol. siccis 1-nerviis interioribus longioribus. Recept. minimum nudum. Cor. regularis glabra 5-fida, lobis acuminatis parte integrâ brevioribus. Fil. lævia. Achænium subcylindraceum multicostratum apice pilosiusculum, nectario styliformi. Pappus biserialis paleaceus serratus, paleis seriei ext. minimis multis inæqualibus, int. 1-4 tortis longis. — Arbor? ramis canescenti, seu fusciscenti-tomentosis. Folia alterna petiolata oblonga integerrima supra glabra subtus albedo-ferruginea. Capitula pedicellata solitaria in corymbum terminata approximata. Cor. albidæ.

2. *D. DISCOLOX* (H. B. et Kunth l. c.) 5 in calidis prope Honda Nove-Granatensium. Eupatorium cuspidatum Willd. herb. ex Less. (v. s. comm. à cl. Poeppig.)

XXV. MONOSIS DC. in ann. bot. Guill. 1833. v. 2. p. 515. et in Wight contrib. p. 1. — *Conyzæ* sp. Wall.

Capitula uniflora in glomerulum non concreta. Invol. oblongi squamæ imbricatæ obtusæ flore multò breviores. Cor. tubulosa 5-fida. Antheræ basi muticæ. Stigmata longè exserta Vernoniæ. Achænium glabrum teretiusculum. Pappus 2-3-serialis, setis rigidulis scabris æqualibus aut longitudine sed non latitudine discrepantibus. — Suffrutices. Folia alterna breviter petiolata obovato-cuneata acutiuscula vix subdentata suprâ glabra subtus cum petiolis et ramis tomentosa. Panicula nuda Rami apice capitula plurima subumbellata subsessilia conferta gerentes. — Affinis videtur Albertiniæ, sed capitula distiucta, Shawiæ, sed pappus crassus basi non pubescens et ovarium sæpius glabrum. — *Monos* græcè solitudo, ex flore in invol. unico.

Sect. I. EUMONOSIS.

Pappi series ext. ab interiore non discrepans. — Frutex Indicus.

1. *M. WIGHTIANA* (DC. in Wight contrib. p. 1), caule fruticoso?, ramis teretibus velutino-tomentosis, foliis petiolatis obovatis subacutis basi cuneatis et obtusis subintegerrimis penninerviis suprâ glabris et in nervis subvelutiniâ subtus velutino-hirsutis, paniculâ ramosissimâ capitulis ad apices ramulorum subcorymbosis subsessilibus, invol. squamis obtusis dorso tomentosis 5 in Indiæ orient. montibus Neelgerry leg. cl. Wight et Noton. *Conyza Wightiana* Wall. ! cat. n. 3028 (non 3092) comp. 138. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

Sect. II. EREMOSIS.

Pappus pluriserialis setaceus, ser. ext. brevi sed ab int. latitudine non discrepante. — Frutices Mexicana.

2. *M. SALICIFOLIA*, fruticosa glabrescens, ramis teretibus, foliis petiolulatis oblongo-lanceolatis utrinquè acuminatis subintegerrimis penninerviis sparsè glandulosi subtus glabris ad axillas venarum pilosis aut sparsè puberulis, corymbo composito terminali, capitulis pedicellatis, squamis infimis minimis, cæteris longis lanceolatis submucronatis subciliatis. 3 in Mexici montibus ad Guchinapa et Cuernavaca legit cl. Berlandier oct. florent. *Scirpium uniflorum* et *Eupatorium uniflorum* Moq. fl. mex. mss. Invol. squamæ corollæque subpurpurascens. Pappi setæ ext. paucæ vix inter. breviores albæ. Folia 2 poll. longa, 6 lin. lata. (v. s.)
3. *M. TARCHONANTHIFOLIA*, caule fruticoso tereti cano et adpressè tomentoso, foliis petiolatis oblongo-lanceolatis utrinquè attenuatis serraturas paucas callosas gerentibus suprâ glabris glandulosi subtus cano-tomentosis, ramis flor. ex axillis supr. et apice caulis nudis apice composito-corymbiferis, pedicellis ultimis confertis ternis brevibus, invol. oblongi 1-flori squamis glabris subcariosis imbricatis ext. brevibus submentosis int. longioribus subacutis glabris, achænio striato glabresculo, pappi setis ext. paucis brevibus et mediochr. ab int. latitudine non discrep. 3 in Mexico legit cl. Karwinski. Pappus albus. Fol. 4 poll. longa, ferè pollicem lata, petiolo 2-3 lin. Invol. sordidè fusciscentia. Cor. purpurascens, (v. s. in h. Acad. reg. Monac.)
4. *M. ? TOMENTOSA*, foliis ovatis integerrimis brevè petiolatis transversè venosis subtus ramisque tomentosis, capitulis in corymbum dispositis, invol. squamis ovatis acutis apice villosis. 3 in Mexico ad montem Guineo prope Vallabatum. Turpinia? *tomentosa* Lallav. et Lex descr. 1 p. 24. *Fulcaldea? tomentosa*

Less. in linnæa 1830. p. 248. Cor. purpureæ. Pappus pilosus et stigmata revoluta ex auct. undè potius huc quàm ad Fulcaldeam referenda videtur.

XXVI. SHAWIA Forst. nov. gen. p. 95. t. 48. Cass. dict. 34. p. 40. n. vi. Less. syn. p. 156.

Capitulum 1-florum. Invol. squamæ 5-6 regulariter imbricatæ adpressæ coriaceæ extus glandulosæ, ext. breves ovales, int. oblongo-lanceolatæ. Recept. punctiforme. Cor. infundibuliformis 5-fida. Styli rami graciles exserti divergentes. Achænium subcylindraceum villosa-subtomentosum. Pappus fructu longior, paleolis filiformibus inæqualibus numerosis basi subnudis apice barbellulatis. — Frutices. Folia alterna. Capitula in paniculam corymbumve disposita. — Genus à Cassinio olim ad Inuleas demùm ad Vernonia relatum. E Vernoniis excludit Lessing et cum dubio ad Eupatorieas refert, sed ex capitulis 1-floris potius ad Vernoniaceas fortè referendum.

1. S. PANICULATA (Forst. et Cass. ll. cc.), foliis ovali-oblongis integerrimis petiolatis subtus ramis pedunculisque tomentosis, capitulis in paniculam dispositis. 3 5 in Novâ Zeelandiâ. Rich. et Leason fl. nov. Zeel. p. 243.

Subdiv. 3. ALBERTINIÆ.

Pappo uni- bi- aut pluriseriali paleaceo aut setiformi, capitulis sæpius paucifloris in glomerulum concretis.

XXVII. HAPLOSTEPHIUM Mart. herb.

Capitula 1-flora in glomerulum globosum aggregata. Bracteæ paucæ foliaceæ sub glomerulo. Invol. partialia oblonga, squamis imbricatis ferè linearibus acutis coriaceo-scariosis glabris. Achænium obovatum glaberrimum nervis 10 costulatum, umbone prominulo. Pappus simplex, paleis caducissimis linearibus spiraliter tortis. — Frutex Brasiliensis Lychnophoræ ferè facie. Rami teretes subverticillati adpressè sericei demùm glabriusculi. Folia conferta sessilia patula subrecurva basi subdilatata ferè adnata lanceolato-linearia acuminata integerrima suprà glabra subtus sericeo-cana ferè ericæformia. — Genus affine Piptocomæ ex pappo caducissimo, Lychnocephalo ex pappi paleis tortis, sed præsertim Lychnophoræ ex habitu.

1. H. PASSERINA (Mart. herb.). 3 in summo monte Itambe de Villa prov. Minas Generalium Brasiliæ legit cl. Martins. Fol. 4 lin. longa, vix lineam lata. (v. s. in h. Acad. reg. Monac.)

XXVIII. LYCHNOPHORA Mart. in act. soc. sciab. v. 3. p. 2. p. 149 et diss. in-4°.

Capitula 1-4-flora in glomerulum subglobosum densè aggregata. Folia sæpius glomerulum bracteantia. Invol. oblongi squamæ 2-3-seriales lineari-oblongæ imbricatæ coriaceo-subscariosæ. Recept. nudum aut alveolare. Achænia glaberrima teretiuscula multicostata. Pappus biserialis, uterque latè paleaceus, serje exteriori multò breviori, interiori paleis tortis. — Fruticuli Brasilienses. Folia sessilia conferta uninnervia coriacea margine revoluta subtus sæpè cana. Glomeruli ramos terminantes. — Genus habitu distinctissimum ex cl. Martius et Saint-Hilaire.

Sect. I. EULYCHNOPHORA.

Capitula 1-flora. — *Lychnophora* genus Less. in *linnæa* 1829.

1. *L. ROSMARINIFOLIA* (Mart. diss. p. 10. t. 9), fruticosa, ramis teretibus crasso-tomentosis, foliis patentibus à basi dilatata cordata acuminato-subulatis acutis margine revolutis supra lineâ mediâ villosis subtus villo albo vestitis, capitulis adgregatis in glomerulos subglobosos interdum ipsos adgregatos subcylindricis 1-floris, invol. squamis obtusis. § in summis montibus distr. Adamantini Brasilie ad Tejuco et Milho-Verde. *Lychnophora rosmarinifolia* Less. in *linnæa* 1829. p. 316. Ob flores in invol. solitarios ab affinibus generice separat cl. Lessing, meo sensu nimis numero concedens. (v. s. in h. Acad. reg. Monac.)

Sect. II. Lychnophoroides.

Capitula 2-4-flora. — *Lychnophora* sect. *Vernonia* Less. in *linnæa* 1829.

2. *L. PHYLICÆFOLIA*, fruticosa demissa, ramis junioribus densè albo-tomentosis demum glabratis, foliis confertis erectis sessilibus lanceolato-linearibus mucrone cuspidatis margine latè revolutis supra glabris subtus albo-tomentosis, capitulis 2-floris in glomerulum foliis cinctum aggregatis, invol. squamis glaberrimis oblongis submembranaceis mucronato-acutis. § in altis ad Lincora prov. Bahiensis Brasilie legit cl. Martins. Suffrutex humilis. Pappus ext. brevissimus ferè coroniformis sed paleæ distinctæ erosæ obtusiusculæ, int. paleis elongatis apice serrulatis. (v. s. in h. Acad. reg. Monac.)
3. *L. STAAVIOIDES* (Mart. ! diss. p. 9. t. 8), fruticosa, ramis teretibus junioribus crasso-tomentosis, foliis patentibus è basi subcordata oblongis obtusis submucronulatis margine latè revolutis supra glabris subtus tomentosis, capitulis cylindricis 3-floris in glomerulum subglobosum foliis interpositis mixtum aggregatis, invol. squamis laxè imbricatis obtusis 3-nerviis. § in summis montibus distr. Adamantini Brasilie. *Vernonia staavioides* Less. in *linnæa* 1829. p. 249. 1831. p. 630. *Dialesta staavioides* Mart. ! herb. Invol. squamæ apice summo tomentosæ. Pappi paleæ ext. sub lente apice serratæ. Cor. rosæ. (v. s. in h. Acad. reg. Monac.)
4. *L. VILLOSISSIMA* (Mart. ! diss. p. 8. t. 7), fruticosa, ramis teretibus crasso et tessellato-tomentosis, foliis patentibus linearibus obtusis margine latè revolutis supra in basi longè lanuginosis subtus ubiquè tomentosis, capitulis 8-10 in glomerulum globosum lanatum adgregatis 4-floris cylindricis, invol. squamis laxè imbricatis subacutis. § in Brasilie distr. Adamantino in campis apricis prope Tejuco. *Vernonia villosissima* Less. in *linnæa* 1829. p. 249. 1831. p. 629. Cor. pallidæ lilacinæ.
5. *L. SALICIFOLIA* (Mart. ! diss. p. 12. t. 8), fruticosa, ramis teretibus crasso et pulvinato-tomentosis, foliis erecto-patentibus lineari-lanceolatis obtusiusculis mucronulatis margine vix revolutis subtus tomentosis, nervo dilatato glabro, capitulis ovatis 6-floris in glomerulos globosos adgregatis foliis floralibus intercinctis, invol. squamis arcuè adpressis lanceolatis. § in summo monte Itambe distr. Adamant. Brasilie. *Vernonia salicifolia* Less. in *linnæa* 1829. p. 249. 1831. p. 629. Cor. pallidæ rosæ. Invol. squamæ et cor. lobi apice barbati. Pappi paleæ sub lente serratæ. (v. s. in h. Acad. reg. Monac.)
6. *L. PINASTER* (Mart. ! diss. p. 7. t. 6), subarborea, ramis teretibus tessellato-tomentosis, foliis erecto-patentibus linearibus obtusis basi attenuatis in glomerulum subglobosum confertis 3-4-floris, invol. squamis glabris obtusiusculis. § in Brasilie montibus prov. Minas-Geraes. Cor. violaceæ. *Vernonia pinaster* Less. in *linnæa* 1829. p. 249. 1831. p. 629. (v. s. in h. Acad. reg. Monac.)
7. *L. HAKKÆFOLIA* (Mart. ! diss. p. 11. t. 10), fruticosa, ramis teretibus crasso et pulvinato-tomentosis, foliis erecto-patentibus æqualiter linearibus obtusis mucronulatis margine revolutis supra adultis glabris subtus tomentosis, nervo dilatato glabro, capitulis cylindricis 4-floris in glomerulos globosos congestis

- foliis floralibus instructis, invol. squamis laxè imbricatis linearibus obtusis. § in summo monte Itambe do Villa Brasiliæ. *Vernonia hakeæfolia* Less. l. c. p. 249. Cor. rosæ. (v. s. in h. Acad. reg. Monac.)
8. *L. PROTEÆFORMIS* (Mart. ! diss. p. 6. t. 5), fruticosa, ramis divaricatis basi nudis apice tessellato-tomentosis, foliis erecto-patentibus linearibus cum mucrone brevi margine revolutis suprâ glabris subtus albo-tomentosis, capitulis terminalibus sessilibus confertis ovato-cylindricis 3-4-floris, invol. squamis lineari-lanceolatis ciliatis. § in Brasiliæ monte Adamantino Serra de Santo Antonio prov. Minas-Geraes. *Vernonia proteæformis* Less. in linnæa 1829. p. 2419. 831. p. 629. Cor. lilacinæ.
9. *L. RACCORDIS* (Mart. ! diss. p. 6. t. 5), subarborea, ramis teretibus tessellato-tomentosis, foliis erecto-patentibus linearibus submucronulatis margine revolutis suprâ glabriusculis nervo medio impresso subtus tomentosis, capitulis oblongis 2-4-floris in glomerulum globosum foliis cinctum adgregatis, invol. squamis lineari-lanceolatis obtusis corollisque glabris. § in monte Adamantino Serra de Santo Antonio prov. Minas-Geraes Brasil. Fol. 15-18 lin. longa, lineam lata. Cor. pallidè lilacinæ. Pappi palæe subulatæ vix apice serrulatæ, exterior. interior. duplò brevior. (v. s. in h. Acad. reg. Monac.)
10. *L. BRUNIODES* (Mart. ! diss. t. 4. p. 4), fruticosa, ramis teretibus glabris nodulosis, foliis erecto-patentibus lineari-lanceolatis acutis basi attenuatis margine revolutis suprâ glabris subtus tomentosis, capitulis corymbosis in glomerulum terminalem congestis 3-4-floris cylindricis, invol. squamis oblongo-linearibus. § in Brasiliæ montanis siccis ad Itambe da Villa. *Vernonia brunioides* Less. in linnæa 1829. p. 248. 1831. p. 629. Rami tennes cicatrissati sæpè lichenibus onusti. Folia 5-6 lin. longa. Cor. pallidè rosæ in genere parvæ. Pappus ext. paleis brevibus erosis acutis, int. paleis elongatis purpurascens. (v. s. in h. Acad. reg. Monac.)
- §. *alpicola*, ramis junioribus pubescenti-subcanescentibus, foliis paulò minus confertis. — in Serro Frio ad Tejuco alt. 5000 ped. prov. Minarum General. Brasiliæ legit cl. Martins. *Vernonia alpicola* var. Mart. ! herb. Pappus ext. paulò adhuc brevior quam var. *α*, int. paleis linearibus purpurascens. (v. s. in h. Acad. reg. Monac.)

XXIX. ALBERTINIA Spreng. *neu entd.* 2 p. 133. *syst.* 3. p. 355 et 434 (excl. *car.*), Mart. *herb.*

Capitula 1-3-flora densissimè in glomerulos subglobosos collecta et sæpius involucrio universali brevi polyphylo cincta. Invol. partialis squamæ erectæ adpressæ sæpè pressione ad alveolorum rudimenta reductæ. Pappus bi- aut pluriserialis, setis omnibus filiformibus subscabris sæpius coloratis. — Frutices Brasilienses. Folia alterna petiolata elliptica utrinquè attenuata integerrima glabriuscula aut subtus albida.

Sect. I. ANISOTRICHIA.

Pappi series externa interiore duplò triplove brevior.

§. I. Capitulis 1-floris.

1. *A. BRASILIENSIS* (Spreng. *syst.* 3. p. 355 et 434), caule fruticoso ramoso tereti apice brevissimè hirsuto, foliis petiolatis ellipticis utrinquè acuminatis integerrimis membranaceis suprâ glabriusculis subtus petioloque subvelutinis, capitulis in racemum subcompositum ovatum dispositis pedunculatis ebracteatis, invol. campanulati floribus multò brevioris squamis velutinis concretis, recept. profundè alveolato. § in Brasiliæ prov. Rio de Janeiro ad Serra d'Estrella et Cercovado (Mart. ! Vauth. ! n. 341) in collibus circa Bahiam (Salzm. ! Blanch. !). *Albertinia Brasiliensis* Less. in linnæa 1829. p. 341. *Vern. Brasiliensis* Less. in

linnæ 1831. p. 682. Pappus rufus biserialis, exterior dimidio brevior. Flores carnei seu pallide violacei suaveolentes. (v. s.)

2. *A. RUFISETA*, caule fruticoso ramoso, ramis teretibus velutinis, foliis subsessilibus ellipticis obtusis integerrimis supra araneoso-velutinis obscure virescentibus subtus adpressè velutino-tomentosis canescentibus, summis pedunculatis suboppositis, capitulis 1-floris dense in glomerulum confertis, invol. squamis liberis lanceolatis, intimis valde acuminatis. $\frac{1}{2}$ in Brasiliæ siccis apricis montosis ad Serro Frio prov. Minarum General. legit cl. Martius. *Albertinia obtusata* Mart. ! herb. *Habitus* *Albertiniæ* et *Eremanthi*, à priori capitulis 1-floris et invol. squamis liberis, à posteriore pappi 2-3-seriali, serie extimâ dimidio breviori differt. Achenium sericeo-villosum turbinato-cylindraceum. Pappus rufus. (v. s. in h. Acad. reg. Monac.)
3. *A. PALLIDISETA*, caule fruticoso ramoso, ramis teretibus velutinis, foliis ellipticis basi cuneatis obtusis integerrimis supra glabriusculis obscure virentibus subtus adpressè velutino-tomentosis canis, pedunculis folia sua superantibus, capitulis 1-floris dense in glomerulum confertis, invol. squamis liberis subhirsutis, intimis mucronatis achenia non superantibus. $\frac{1}{2}$ in campis editis ad Serro Frio prov. Minar. Gener. Brasiliæ legit cl. Martius. *Albertinia obtusata* var. ? Mart. ! herb. A præcedenti tantum differt glomerulis longius pedunculatis, invol. squamis minus acuminatis et pappo paulò magis paleaceo stramineo nec raso. An var. ? (v. s. in h. Acad. reg. Monac.)

§. 2. Capitulis 3-floris.

4. *A. CROTOMOIDES*, ramis teretibus hirsuto-tomentosis super. subcorymbosis, foliis breviter petiolatis ellipticis obtusis basi subattenuatis integerrimis supra junioribus subvelutinis adultis glabris subtus petiolisque dense cano-tomentosis, capitulis 3-floris oblongis ad apices ramulorum conferto-umbellatis brevissimè pedicellatis, invol. squamis imbricatis lanceolatis acutis dorso tomentosis, achenio pilosiusculo. $\frac{1}{2}$ in montium sepibus prov. Minar. Gener. Brasiliæ legit cl. Martius. *Albertinia bardanoides* Mart. herb. Folia a poll. longa pollicem lata. Pappus duplex purpurascens, ser. ext. dimidio brevior magis paleacea. (v. s. in h. Acad. reg. Monac.)
5. *A. VERBASCIFOLIA* (Mart. ! herb.), ramis teretibus petiolis foliisque subtus densissimè hirsuto-tomentosis, foliis supra junioribus tomentosis adultis subvelutinis basi cordatis latè ovatis obtusis integerrimis lanugine crassis penninerviis, nervis later. utrinquè 15-16 subparallelis reticulato-ramosis, capitulis 3-floris oblongis ad apices ramulorum glomerulato-umbellatis ebracteatis, invol. squamis lanceolatis acutis dorso velutinis, achenio pilosiusculo. — in ferrugineis Serra de Ant. Pereira prov. Minar. General. legit cl. Martius. Pappus subrufescens biserialis, ext. dimidio brevior. Toto cælo diversa à *V. verbascifolia* Less. Fol. petiolus pollicaris, limbus 6 poll. longus, 4-5 poll. latus. (v. s. in h. Acad. reg. Monac.)
6. *A. ELÆAGNEUS* (Mart. ! herb.), caule fruticoso ramoso tereti subcanescente, ramulis ultimis angulatis, foliis petiolatis oblongis obtusis basi acuminatis integerrimis supra cinrescenti-subtus rufescenti-adpressè velutinis, capitulis 5-6 ad apices ramulorum umbellato-sessilibus 3-floris, invol. squamis subincanis lanceolatis acutis, achenio pilosiusculo. $\frac{1}{2}$ in altis lapideis Serro Frio ad Tejuco prov. Minar. Gener. Brasiliæ legit cl. Martius. Pappi pallide rufescentes series 2-3, ext. paucisetâ breviori. Folia utrinquè nervo medio saltem basi transversè ruguloso ! sen tuberculoso ; an casu ? (v. s. in h. Acad. reg. Monac.)
7. *A. STOLON*, fruticosa, ramis teretibus dense tomentosis, foliis petiolatis coriaceis ovali-oblongis basi attenuatis vix acutis integerrimis supra glabris subtus petiolisque cano-tomentosis, capitulis oblongis 2-3-floris in glomerulum globosum foliis paucis sessilibus sæpè cinctum dense aggregatis. $\frac{1}{2}$ in altis prov. Minarum Brasiliæ legit cl. Martius. *Lychnocephalus bicolor* Mart. ! herb. *Habitus* *Lychnophoræ*, capitulum *Echinopsidi*, bractææ fere *Hololepidis*, invol. squamæ apice obtusæ tomentosæ. Achenia glaberrima subangulata. Pappus

setosus pluriserialis, ser. ext. breviori. Folia 6-8 poll. longa, 1-2 poll. lata, nervo subtus valde prominente. (v. s. in h. Acad. reg. Monac.)

Sect. II. ISOTRICHIA.

Pappi series externa ab interiore nec longitudine nec latitudine discrepans.

§. 1. Capitulis 2-3-floris.

8. *A. CAPITATA*, fruticosa, ramis adultis teretibus junioribus pedunculisque angulatis canis, foliis petiolatis ellipticis utrinque acuminatis subintegerrimis suprâ glabris subtus petiolis ramulisque dense lepidoto-tomentosis albis, capitulis cylindraceis ad apices ramorum sessilibus dense corymbosis 3-floris, invol. squamis arcuè imbricatis ex ovato-oblongis, achænio teretiusculo multicostato glabro, pappi paleis seriebus 2 æqualibus. § in Brasiliâ tropicâ, ad Portam d'Estrellam (Less.!), in monte Cercovado (Lund! Lhostky!) Conyza capitata Spreng. syst. 3. p. 507. Less. in linnæâ 1829. p. 270. Vernonia incaulescens Mart. herb. Pappus atramineo-rufescens caducissimus. Achæniî callus basilaris vix perspicuus. (v. s.)
9. *A. RHYTHOPAPA*, caule fruticoso ramoso, ramis pedunculisque striato-angulatis adpressè incanis, foliis breviter petiolatis lanceolatis basi acutè apice subobtusè acuminatis integerrimis suprâ glabris subtus adpressimè canis, capitulis 2-3-floris dense in globulolum pedunculatum congestis, invol. squamis ext. subtomentosis truncatis, intimis paulò longioribus acutis. § in Brasiliâ prov. Minarum Generalium legit cl. Vauthier (coll. exs. n. 334). Pappus purpurascens, setis biserialibus æqualibus. Achæniolum glabriusculum. (v. s. comm. à cl. Vauth.)

§. 2. Capitulis unifloris. — *Eremanthus* Less. in linnæâ 1829. p. 317. 1831. p. 682. syn. 147. non Cass.

10. *A. GLOMERULATA*, ramis pedunculisque teretibus, foliis obtusis, floralibus sub glomerulo globoso eo longioribus reflexis, achænio pilosiusculo. § in Brasiliâ tropicâ (Less.) prov. Minar. General. (Gaudich.!) *Eremanthus glomerulatus* Less. II. cc. Folia et rami supremi suboppositi. Pappus rufus. (v. s. in h. Mus. reg. Paris.)
11. *A. INCANA* (Less. in linnæâ 1829. p. 342), ramis pedunculisque striatis compressis foliis lanceolatis utrinque acuminatis, floralibus capitulo brevioribus, achænio glabro. § in Brasiliâ tropicâ (Less.) prov. Minarum Generalium (Vauth. n. 355). Vernonia incana Less. in linnæâ 1831. p. 682. Invol. squamæ ext. breviter obtusæ subtomentosæ, intimæ abruptè duplò longiores glabriusculæ. Pappus albidus. (v. s.)
12. *A. POLYCEPHALA*, caule fruticoso striato corymboso ramoso, ramis angulatis adpressè subincanis, ultimis striato-sulcatis subcompressis, foliis lanceolatis suprâ glabris subtus adpressè canescentibus integerrimis basi acuminatis subsessilibus apice vix acutis, capitulis 1-cephalis umbellato-glomeratis glomerulis subbracteatis pedunculatis, invol. squamis ext. brevibus obtusis, intimis elongatis glabriusculis, achænio glabro. § in planitie altâ ad Pieda de Villam prov. Minar. Gener. Brasiliâ legit cl. Martius. Albertinia multiflora Mart. herb. Pappi series æquales subpurpurascens. (v. s. in h. Acad. reg. Monac.)
13. *A. SALICINA* (Mart. herb.) caule fruticoso tereti ramoso, ramulis ultimis angulatis subcompressis adpressè subincanis, foliis linearilanceolatis utrinque attenuatis subobtusis integerrimis suprâ glabris subtus adpressè camotomentosis, capitulis 1-floris in glomerulos bracteatos pedunculatos 8-9 umbellato-congestis, invol. squamis ext. brevibus subincanis, int. elongatis glabriusculis. § in editis sticciis rupestribus montium Serro Frio prov. Minar. General. Brasiliâ legit cl. Martius. Folia 2 poll. longa, 5 lin. lata. Pappus pallidus setis ex statu juniore æqualibus. (v. s. in h. Acad. reg. Monac.)

XXX. PYCNOCEPHALUM. — *Pycnocephalum* sect. *Vernonia* Less. in *linnæa* 1831. p. 630.

Capitula plurima 2-7-flora in glomerulum terminalem subglobosum collecta. Squamæ ovales multistriatæ paucæ sub glomerulo involucri locum gerentes. Invol. partialis squamæ oblongo-lanceolatæ 3-nerviæ subciliatæ. Cor. glabræ. Achænium sericeo-villosum. Pappus pluriserialis setaceus scaber, setis ext. paucis brevioribus sensim ad interiorum longitudinem accedentibus. — Herbæ perennes imâ basi foliatæ cæterum aphyllæ simplicicaules.

1. *P. SPATHULÆFOLIUM*, collo crasso subtomentoso, caulibus pluribus herbaceis basi foliatis cæterum aphyllis compressis densè sericeo-pubescentibus, foliis subradicalibus lineari-oblongis (spathulatisve ex Mart.) basi angustatis erectis integerrimis junioribus pubescentibus adultis glabris subnerviis, capitulis oblongis 5-7-floris in glomerulum globosum ebracteatum terminalem congestis, invol. squamis lanceolatis acuminatis basi 3-nerviis subciliatis, intimis glabris. ½ in aridis deserti versus Chapada de Parano prov. Minar. General. Brasiliæ legit cl. Martius. *Vernonia spathulæfolia* Mart. herb. Valdè affinis *P. scapigero*. Achænium sericeo-villosum. Pappus pluriserialis, setis scabris, ext. dimidio circ. brevioribus. Cor. purpureæ. (v. s. in h. Acad. reg. Monac.)
2. *P. SCAPIGERUM*, caule herbaceo simplicissimo aphylo tereti lanato-tomentoso, foliis radicalibus coriaceis brevè petiolatis spathulato-oblongis subintegerrimis glaberrimis junioribus subtomentosis, capitulis subcylindraceis 2-3-floris in glomerulum terminalem adgregatis, invol. squamis carinatis mucronato-acuminatis, achænio densè piloso. ½ in Brasiliâ tropicâ. *Vernonia scapigera* Less. in *linnæa* 1829. p. 250. Cor. purpureæ glabræ.
3. *P. PLANTAGINIFOLIUM*, caule herbaceo simplicissimo tereti supernè aphylo tomentoso, foliis subradicalibus membranaceis multiplinerviis lineari-lanceolatis integerrimis glabriusculis, junioribus tomentosulis, capitulis cylindraceis multifloris in glomerulum adgregatis, invol. squamis planiusculis 3-nerviis glabriusculis, achænio densè piloso. ½ in Brasiliâ tropicâ. Cor. purpureæ glabræ. *Vernonia plantaginifolia* Less. in *linnæa* 1829. p. 251.

XXXI. LYCHNOCEPHALUS Mart. herb.

Capitula plurima 8-flora ovali-oblonga densissimè conferta in glomerulum terminalem subrotundum involuacro generali destitutum. Bractæe foliaceæ sub capitulo quoque et ideò in medio glomeruli hinc inde perspicuæ. Involucrum parziale oblongo-cylindraceum in lanâ immersum, squamis plurimis glabriusculis obtusis. Achænium glabrum striato-sulcatum. Pappus duplex paleaceus, paleis ext. brevibus acutis persistentibus, int. elongatis spiraliter tortis caducissimis. — Frutex Brasiliensis. Rami teretes densè tomentosi. Folia petiolata ovali-oblonga obtusa basi cuneata coriacea integerrima suprâ glabrâ subtus petiolique densè tomentosa.

1. *L. TOMENTOSUS* (Mart. herb.). ¾ in campis sterilibus altis provinciæ Minas-Geraes Brasiliæ legit cl. Martius. Variat tomento albicante et flavicante. Invol. squamæ obtusæ glabriusculæ. Achænium glabrum striato-costatum. Pappus duplex paleaceus, ext. brevis acutus, int. elongatus spiraliter tortus caducissimus. (v. s. in h. Acad. reg. Monac.)

† *Species exclusa.*

L. BICOLOR Mart. = *Albertinia bicolor*.

XXXII. CHRONOPAPPUS. — *Heterocomæ* sp. DC.

Capitula pauca 8-10-flora in glomerulum axillarem conferta, bracteis foliaceis involucrium verum non constituentibus circumdata. Invol. partialis ovalis squamæ imbricatæ dorso tomentosæ acutæ lanceolato-lineares, int. angustissimæ subbreviares. Recept. nudum. Corollæ lobi dorso hirsuti. Achænia breviter angulata glaberrima. Pappus duplex uterque persistens, ext. paleis brevibus oblongis acutis, int. setis elongatis serratis. — Ab *Heterocomâ* differt bracteolis recept. nullis, corollâ hirsutâ, pappo ext. paleaceo nec coroniformi. Proximè ad *Lychnocephalum* accedit.

1. *C. bifrons*, fruticosa, ramis teretibus apice subcompressis valdè lanuginosis, foliis petiolatis ovatis basi obtusissimis apice subobtusis integerrimis suprâ adultis glabris eminentiis duris prismaticis eximie muricatis suprâ petiolisque cano-lanuginosis, glomerulis ad axillas superiores subsessilibus spicam compositam interruptam terminalem ferè constituentibus. ζ in Brasiliâ. *Heterocomâ* bifrons DC. ann. mus. 16. p. 191. t. 8. recueîl mém. 2. p. 34. t. 4. *Serratula bifrons* Pers. ench. 2. p. 391. (v. s.)

XXXIII. PITHECOSERIS Mart. herb.

Capitula 3-4-flora in glomerulum ovatum densè congesta sessilia subspicata ebracteata. Invol. oblongi squamæ erectæ glabriusculæ acuminatæ, ext. carinatæ, int. planæ lanceolato-lineares. Recept. nudum. Cor. tubus extus hirsutus, lobi glabri. Achænia in iisdem capitulis difformia, alia (juniora aut sterilia?) sericeo-villosa, pappo duplici ext. brevissimo paleaceo, int. pluriseriali setaceo: alia (matura aut fertilia?) glaberrima oblonga compressa, pappo setiformi caducissimo! — Herba. Folia ferè lyrata seu sinuato-pinnatifida.

2. *P. PACOURINOIDES* (Mart. ! herb.), glabriuscula, caule tereti, ramis apice fistulosis nudis, foliis sessilibus aurito-semiamplexicaulibus irreg. pinnati-lobatis lobis ovatis triangularibus dentatis subsinuatisve, terminali majore, glomerulo ovato glabro. \mathcal{U} in campis prov. Pernambucanæ legit cl. Martius. Herba grandis secus caulem et in foliorum paginâ infer. subtiliter pubescens. Folia membranacea pedem longa, 6 poll. lata. Capit. oblonga 3-5-flora. Invol. squamæ glabræ acuminatæ. Achænia juniora puberula adulta glaberrima. Pappus pluriserialis setosus, ser. ext. brevis, int. longa subfrescens, facillè deciduus in achæniis adultis. (v. s. in h. Acad. reg. Monac.)

XXXIV. STACHYANTHUS.

Capitula 12-13-flora secus rami apicem sessilia spicata subconferta, rachi ad eorum exortum scrobiculatâ, bracteis subcapitulis nullis. Invol. squamæ oblongæ adpressæ obtusæ, int. submucronatæ. Achænia sericeo-villosa. Pappus pluriserialis setaceus, serie ext. brevi basi flavidâ. — Folia sinuato-angulata. Affinis *Echinopsidio*.

3. *S. MARTII*, suffruticosa tota pube brevi adpressâ sericeo-canescens, caule tereti ramoso, foliis petiolatis ovato-rhombicis basi integris cuneatis à medio ad apicem dentato-angulatis, pedunculis longissimis nudis glomerulo ovato-globoso terminatis, capitulis secus axin spicatis 12-13-floris, invol. squamis oblongis submucronulatis. \mathcal{U} ζ in siccis sylvis æstu aphyllis prov. Bahiensis et Piahiensis Brasiliæ legit cl. Martius. *Vernonia Stachyanthus* Mart. ! herb. Achænium sericeo-villosum. Pappus pluriserialis, setis ext. brevibus, basi flavidis. Folia ferè *Atriplex* roseæ, sed limbo 3 poll. longo, 2 poll. lato, petiolo pollicari. (v. s. in h. Acad. reg. Monac.)

XXXV. CHRESTA *Fl. flumin. ic.*

Capitula 3-4-flora in glomerulos sphaericos terminales ebracteatos densissimè collecta. Invol. squamæ siccae acuminatæ paucæ. Achænium sericeo-villosissimum. Pappus multiserialis, serie ext. cæteris brevior. — Habitus distinctissimus hinc ad Elephantopum illinc ad Lychnocephalum accedens.

1. *C. SPHÆROCEPHALA*, tota canescens velutino-tomentosa, caule erecto tereti apice ramoso, foliis petiolatis ovatis penninerviis coriaceis dentato-subinnatis, ramis nudis capitulo globoso compacto terminatis. $\frac{1}{2}$ in saxosis calvis aridiusculis prov. Bahiensis legit cl. Martins, et in prov. Minas Geraes prope Tejuco cl. Vauthier (coll. exs. n. 294) Chresta cordata Arrab. fl. flum. 8. t. 150? In specim. Vauthieriano pedunculi apice subglomerulo fissi seu ramosi aut monstruositate aut compressione spectabiles, in Martiano indivisi. Invol. squamæ glaberrimæ. Cor. glabræ purpureæ. *Vernonia Echinops* Mart. ! herb. (v. s.)

2. *C. EXSUCCA*, caule erecto ramoso pubescenti-tomentoso, foliis sessilibus ellipticis acutiusculis integerrimis coriaceis supra glabris, subtus nervoso-reticulatis junioribus subvelutinis adultis glabris, ramis apice nudis glomerulum globosum gerentibus, invol. squamis lanceolatis cuspidatis. $\frac{1}{2}$ in montosis prov. Minar. Gener. Brasiliæ legit cl. Martins. *Vernonia exsucca* Mart. ! herb. Folia 12-15 lin. longa, 6-7 lata. Achænium sericeo-villosissimum. Pappus flavescens, setis ciliatis pluriserialibus ext. dimidiò triplòve brevioribus. Species affinis *C. pycnocephalæ*. (v. s. in h. Acad. reg. Monac.)

4. *C. PYCNOCEPHALA*, tota adpressè sericeo-pubescenti ferè lepidota, caule tereti apice ramoso, foliis sessilibus obovato-oblongis acutis subintegerrimis penninerviis, ramis nudis glomerulo globoso compacto terminatis. $\frac{1}{2}$ in campis deserti inter. Min. Nov. et S. Franciscum prov. Minar. General. Brasiliæ legit cl. Martins. *Vernonia pycnocephala* Mart. herb. Glomeruli dimidiò minores et folia diversissima à *C. sphærocephalâ* cui cæter. valdè affinis. Capitula 4-flora. Pedunculi folia quintoplò superantes. Invol. squamæ glabresculæ submucronatæ. (v. s. in h. Acad. reg. Monac.)

ß. Halimodendron, pedunculis folio triplò tantum circ. longioribus, invol. squamis extus puberulis. $\frac{1}{2}$ in campis siccis deserti prov. Minar. Gen. prope Reacho de Togo et ad fl. S. Francisci prope Salgado legit cl. Martins. *Vern. Halimodendron* Mart. ! herb. (v. s. in h. Acad. reg. Monac.)

Div. II. ELEPHANTOPÆ *Cass. tabl. xx. sect. 4. n. 111. excl. Dialista Less. in linnæâ 1830. p. 135. — Elephantopodæ Less. syn. 149.*

Capitula discoidea pauciflora. Invol. compressum pluriseriale, foliolis seriis cujusque 4, alternis conduplicatis, alternis planis conduplicatis distichis. Recept. nudum aut alveolatum. Cor. palmata. Antheræ ecaudatæ. Styli rami semisubulati. — Folia alterna.

XXXVI. ELEPHANTOPUS *Cass. dict. 14. p. 341. Less. in linnæâ 1829. p. 323. syn. 149. — Elephantopi sp. Linn. Gartn.*

Capitula pluriflora (3-5 sæpius 4-flora) aequaliflora in glomerulum foliis cinctum densè adgregata. Involucrum compressum biseriale, foliolis siccis oblongis alternè planis et conduplicatis interioribus sæpius 3-nerviis. Rhachis nuda. Cor. palmata, limbo 5-fido, laciniis acuminatis, sinu uno cæteris profundiore. Stam. filam. lævia. Styli rami semisubulati. Achænium compressiusculum multicostatum oblongum pilosum. Pappus 1-serialis, paleis plurimis rectis basi dilatatis cæterum

angustissimis acuminatis æqualibus serratis. — Herbæ perennes pilosæ erectæ. Folia alterna sessilia penninervia. Glomeruli ramos terminantes subcorymbosi. Cor. violacæ rarius albæ.

§. 1. *Corollâ glabrâ.*

1. *E. SCABER* (Linn. sp. 1313. excl. syn. Gron. et Brown.), caule dichotomè ramoso strigoso-villoso, foliis scabris radicalibus crenatis cuneatis basi longè attenuatis, caulinis lanceolatis, floralibus latè cordato-ovatis acuminatis canescentibus. \mathcal{U} in Ind. orient. frequens in Timor, Javâ, Luconia, Silhet, Nepaliâ, agro Burmannico, etc. Less. in linnæâ 1829. p. 325. Wall. ! herb. comp. n. 89. Wight ! herb. — Dill. elth. 126. t. 106. Breyn. ic. 32. t. 34. (v. s.)
2. *E. CAROLINIANUS* (Willd. sp. 3. p. 2390 excl. Br. syn.), caule piloso apice corymboso-ramoso, foliis scabris, radicalibus crenatis cuneatis basi attenuatis, caulinis oblongis basi angustis floralibus ovato-oblongis scabridis. \mathcal{U} in siccis substerilibus Carolinæ, agri Kentuckensis, Floridæ, et ex Willd. in Jamaicâ, ex Sieber in Martinicâ, ex variis in Brasiliâ, Mexico, etc.; sed fortè cum sequentibus sæpè confusus. Ell. sketch, 2. p. 480. Less. in linnæâ 1829. p. 325. *E. scaber* Swartz obs. 316? Michx. fl. bor. am. 2. p. 148. Glomeruli bractæe latitudine duplè et ultrâ longiores nec ut in priore latitudini ferè longitudine æquales. (v. s.)
3. *E. NUDICAULIS* (Ell. sketch, 2. p. 481), caule subhirsuto subnudo parcè ramoso, foliis radicalibus ovali-sublanceolatis crenato-serratis suprâ scabrinisculis subtus villosis, floralibus latè ovato-cordatis villosissimis. \mathcal{U} in siccis substerilibus Carolinæ Poir. suppl. 2. p. 543. *E. Carolinianus* var. *simplex* Nutt. gen. 2 p. 187. *E. tomentosus* Linn. sp. 1314? excl. syn. auct. Carib. Pursh. fl. 2. p. 582? Less. in linnæâ 1829. p. 326? an *E. Caroliniani* var.? (v. s.)
4. *E. MOLLIS* (H. B. et Kunth nov. gen. am. 4. p. 26), caule strigoso-piloso supernè dichotomo-ramoso, foliis ovatis inferioribus basi attenuatis crenato-serratis suprâ glabrinisculis subtus molliissimè canescenti-villosis, caulinis oblongis acuminatis serratis, floralibus ternis latè ovatis acuminatis serratis. \mathcal{U} in Cumanâ et Caracas ex H. B. et K., Havanâ, Portorico, Peruvîâ (Poepp.), Brasiliæ prov. S. Pauli (Lund). Videtur *E. tomentosus* auct. in Caribæis indicatus. *E. Carolinensis* albiflorus Meyer esseq. 256? Sloan. hist. t. 156. f. 1. 2? Cor. alba ex Kunth et Meyer. (v. s.)
5. *E. MARTII* (Graham list. pl. rar. mart. 1830. p. 2), caule ramoso piloso folioso, foliis subsessilibus undulatis rugosis supernè hispidis subtus tomentosovillosis serrato-crenatis denticulatis, inferioribus spatulato-oblongis breviter attenuatis, floralibus latè ovato-cordatis acutis villosis. \mathcal{U} ? in Brasiliâ ad Rio-Janeiro (Lund!), Sanctam Catharinam (Bacle!), Bahiam in incultis (Salzman!) El. cervinus Arrab. fl. flum. 8. t. 148. Vulgatissimus in pratis apricis et dumetis circa habitationes usque ad altitud. 3000 ped. (herb. Acad. Monac.) Ab indigenis *Yerva grossa* dictus. Infusio theæformis sudorifica. Cor. pallida. (v. s.)

§. 2. *Corollæ lobis strigoso-pilosis.*

6. *E. HIRTIFLORUS*, caule hirtello sulcato subdichotomo, foliis oblongis sessilibus acutis villosissimis, inferioribus serratis, cæteris integris, floralibus oblongo-lanceolatis acuminatis involucrique foliolis villosis. — in sabniosis maritimis circa Bahiam legit cl. Salzman. Cor. ex eo pallidè cœrulescens, lobis ad apicem strigis hispidis. (v. s.)

† *Species non satis notæ.*

7. *E. VIRGATUS* (Desv. in Ham. fl. ind. occ. 52.) caule virgato hirsuto-piloso, foliis subsessilibus trinerviis lanceolato-linearibus remotè serratis utrinquè strigoso-pilosis, pedunculis longissimis paniculatim dispositis. — in Guianâ. Nullam novi *E. speciem* foliis trinerviis.

8. *E. FLEXUOSUS* (Raf. fl. Ind. 73), caule ramoso flexuoso, foliis radicalibus caulinisque sessilibus ovatis denticulatis acutis subtus tomentosis margine revolutis. — in Louisiana ex Robin, voy. p. 448. An fortè *E. mollis*, sed corollacina dicitur.

†† *Species exclusæ.*

- E. ANGUSTIFOLIUS* Sw. = *Elephantosis angustifolia*.
E. NUDIFLORUS Spr. = *Elephantosis angustifolia*.
E. NUDIFLORUS Willd. = *Distreptus nudiflorus*.
E. SPICATUS Juss. et Lam. = *Distreptus spicatus*.
E. STRIGOSUS Willd. = *Distreptus spicatus*.

XXXVII. *ELEPHANTOSIS* Less. in linn. 1829. p. 322. f. 14-17 syn. 149.

Omnia Elephantopi sed pappus biserialis, paleis angustissimis rectis. — Capitula seu glomeruli spicatum secus caulis apicem dispositi.

1. *E. NIFLORA* (Less. l. c.), pappi serie exteriori brevi ab interiori longitudine distinctâ, capitula biflora. 2. in Brasiliâ tropicâ. Pappi series interna setis deciduis tortis constans.
2. *E. QUADRIFLORA* (Less. l. c.), pappi seriebus longitudine non distinctis, capitula 4-floris, glomerulis inferioribus subpedunculatis, foliis utrinquè villosis, remotè dentatis. — in Brasiliâ tropicâ juxta Novum Friburgum (Sell!), in prov. Sancti Pauli (Lund!). Folia etiam in specim. herb. Mus. Bras. video remotè denticulata! An ideò *El. liatroides* cat. h. Petropol. certè distincta? (v. s.)
3. *E. ANGUSTIFOLIA*, pappi seriebus longitudine non distinctis, capitulis 4-floris, glomerulis sessilibus, foliis glabresculis. 2. in locis montosis Jamaicae, Sancti Domingi (Bertero!) et Brasiliâ circa Bahiam (Salzman!). *Elephantopus angustifolius* Swartz fl. ind. 1383. Less. in linnæa 1829. p. 327. *Distreptus angustifolius* Cass. *Eleph. nudiflorus* Spreng. syst. 4. p. 454. non Willd. — Sloan. hist. 5. t. 148: f. 4. (v. s.)

XXXVIII. *DISTREPTUS* Cass. bull. philom. apr. 1817. dict. sc. nat. 13. p. 366. Less. in linnæa 1829. p. 327. f. 19. 33. 34. syn. 149. — *Pseudo-Elephantopus Rohr* in Skrivt. af. n. h. Selsk. Kiob. 11. p. 2. p. 213. — *Matamoria Lallav. et Lex. nov. veg. deser.* 1. p. 8.

Omnia Elephantopi sed pappi paleæ duæ cæteris longiores apice deorsum deinde sursum plicatæ (vid. Gærtn. t. 165. ic. *Eleph. spicati*). — Capitula secus axin ad foliorum axillas sessilia et ideò spicam interruptam ferè formantia.

1. *D. SPICATUS* (Cass. et Less. ll. cc.), foliis floralibus saltem inferioribus capitula superantibus. 2. in cultis ruderalisque Jamaicae, Hispaniolæ, Porto-Ricci, Martinicæ, Cubæ, Trinilitis, Mexici ad Veracruz et Tampico, prov. Caracasæ Mexici, etiam in Philippinis verisim. introductus. *Elephantopus spicatus* B. Juss. in Aubl. guian. 868. Swartz! fl. ind. occ. 1380. Poepp.! per. exs. n. 38. *Matamoria spicata* Lallav. et Lex. nov. veg. descr. 1. p. 8. — Sloan. hist. t. 150. f. 3. 4. Flores albi. Variat foliis latioribus et angustioribus, superficie glabrata aut villosa, ramis erectis aut divaricatis. Huc *E. strigosus* Willd. herb. adductus ut mera varietas à cl. Lessing. (v. s.)
2. *D. CRISPUS* (Cass. dict. 60. (1830.) p. 601. excl. syn.), foliis hirtis, floralibus saltem inferioribus densè ciliatis capitula superantibus, spicis acuminatis, achæniis glaberrimis, pappi setis spiraliter tortis. — in Jamaica. *D. spiralis* Less. in linnæa 1831. p. 690. syn. 149.
3. *D. NUDIFLORUS* (Less. l. c. p. 329), foliis floralibus capitulis dimidio brevioribus. — in Sancto Domingo. *Elephantopus nudiflorus* Willd. sp. 31 p. 2390

non Spreng. — Sub hoc nomine habeo ex Tabago? stirpem priori similem, sed spica magis continua, capitula minora ad axillas fol. flor. geminata terna quaternave 4-flora. Pappi paleæ 4-5 apice crispato-undulatæ nec regulariter biflexæ. An eadæm ac stirps Willd. et Less.?

†. *Species exclusa.*

D. ANGUSTIFOLIUS Cass. = Elephantosis angustifolius.

DIV. III. ROLANDRÆ Cass. tabl. xx. sect. v. (excl. gen.) Less. syn. 150.

Capitula 1-flora. Cor. 5-fida regularis. Invol. 1-seriale ad summum 4-5-phyl-lum sæpè gamophyllum aut rarissimè 2-phyl-lum basi breviter bracteolatum. Pappus brevis sæpissimè coroniformis. — Septibus forsan artificialis et typo colligens plurimos valdè diversos ob numerum specierum paucissimum nondum separatos.

XXXIX. GUNDELIA Tourn. itin. 2. p. 251. ic. cor. 51. inst. t. 486. Gaert. fr. 2. p. 386. t. 163. Cass. dict. 20. p. 93. Less. in *linnaë* 1829. p. 334. f. 43. 47. 68. 74. 76. 79-83. syn. p. 151. — Hacub Vaill. act. sc. par. 17. 18. p. 150. f. 16. — Gundelheimeria Cass. tabl. xx. n. 52.

Capitula ad bractearum axillas 5-7 uniflora inter se ferè ad apicem concreta, centrali fertili, cæteris sterilibus. Involucra 1-serialia gamophylla flore breviora apice spinosa cum adstantibus demum connata. Cor. regularis glabra, limbo tubo longiore 5-fido. Antheræ inclusæ. Styli rami semisubulati extus pilis erectis hispidi intus plani glabri. Achænium glabrum compresso-subtetragonum, nectario alveolari basi pyramidato, disco epigyno magno. Pappus 1-serialis coroniformis, ore brevi spinuloso. — Herba lactescens Dipsaci seu Eryngii facie. Folia alterna sessilia semiamplexicaulia pinnatilobata, lobis dentibusque spinosis ferè ut in Acantho spinoso. Flores purpurascens. Capitula suprâ descripta in capitulum generale ovatum basi exinvolucratum adgregata.

1. G. TOURNEFORTII (Lin. sp. 1315). ½ in Persiâ, Armeniâ, Syriâ, agro Aleppensi, locis aridis et lapidosis. Tourn. II. cc. Silybum seu Hacub, etc. Rauw. itin. 74. ic. — Lam. dict. 3. p. 60. Mill. ic. t. 287.

a. *glabra*, capitulis glabris. — G. glabra Mill. dict. n. 2 — Tourn. cor. 51. n. 1. (v. v.)

β. *araneosa*, capitulis araneosâ lanugine obsitis. — G. Tournefortii Mill. dict. n. 1. — Tourn. cor. 51. n. 2. (v. s.)

XL. CORYMBIUM Linn. gen. n. 1004. Lam. ill. t. 723. Gaert. fr. 2. p. 42. t. 86. DC. *recueil mém.* 1. p. 4. Cass. dict. 10. p. 580. Lessing in *linnaë* 1829. p. 330. f. 13. 20-25. et 1831. p. 691. syn. p. 150. — Contarena Adams. fam. 2. p. 120.

Capitulum 1-florum. Involucrum cylindraceum 2-phyl-lum, squamis canaliculatis liberis, basi foliolis exiguis paucissimis calyculatum. Cor. carnulosa regularis, limbo 5-partito. Antheræ inclusæ, in sicco subliberatæ. Styli rami semicylindracei. Achænium teres oblongum basi longè attenuatum extus undiquè pilis longissimis densis obsitum. Pappus 1-serialis brevis membranaceus, in lacinias piliformes fissus. — Herbæ Capenses, radice crassâ perenni, collo lanuginoso, foliis radicalibus linearibus parallelè plurinerviis limbo nullo ad petiolum reductis. Caulis subnudus foliis parvis semiamplexicaulibus apice ramoso-corym-

posus. — Falsò dixerunt Gærtner semen albuminosum, Willdenow antheras verè liberas, Cassini involucri squamas cum ovario et inter se concretas.

1. *C. nervosum* (Thunb. fl. cap. 729), totum glaberrimum, caule subangulato lævi, foliis latè lanceolato-linearibus, nervis subtùs valdè prominulis, marginibus crassis callosis, corymbo laxo, invol. squamis lævibus. \mathcal{U} ad Cap. B. Spei, circa urbem (Th. Gaud. ! Burch. ! n. 6771), ad montem Tabularem et Kleindrakenstein (Dreg. !), in distr. Uitenhagen (Eckl. et Zeyh. !). *C. glabrum* Lin. f. suppl. p. 392. Lam. ill. t. 273. f. 2. Less. l. c. non Thunb. Pluk. t. 272. f. 4? Axillæ et bases foliorum nunc glabræ nunc subbarbatæ. Folia 3-6 lin. lata, nervis 5-7 subtùs quasi quadratis. (v. s.)
2. *C. glabrum* (Thunb. fl. cap. 729), totum glaberrimum, caule subangulato lævi, foliis linearibus seu angustè lanceolatis, nervis subtùs prominulis, marginibus non incrassatis, corymbo laxo, invol. squamis lævibus. \mathcal{U} ad Cap. Bonæ Spei (Th.), ad Drakeenstenberg et Onder-Bokkewald (Dreg. !), in distr. Worcester (Eckl. et Zeyh. !). *C. gramineum* Lam. dict. 2. p. 129? ill. t. 723. f. 3. Affine priori sed folia longiora, 2 lin. tantùm lata, nervis multò minùs prominulis, margine non incrassato. (v. s.)
3. *C. cymosum* (E. Mey. ! in Dreg. coll. n. 1525), totum glaberrimum, caule teretiusculo lævi, foliis linearibus ad marginem non incrassatis, nervis vix subprominulis, corymbo conferto, invol. squamis lævibus. \mathcal{U} ad Cap. Bonæ Spei, in parte Coloniæ orientali (Burch. ! n. 6661), ad Paarlberg (Dreg. !), in Stellenbosch, Swellendam et circa urbem (Eckl. et Zeyh. !). Priori affine sed corymbus confertus nec laxus. An *C. gramineum* Lam. hùc potiùs referendum? (v. s.)
4. *C. scabrum* (Linn. f. suppl. 392), caule tereti muricato, foliis elongato-linearibus glabris planiusculis, invol. squamis extùs muricatis. \mathcal{U} ad Cap. B. Spei in lateribus montium prope urbem (Th.), ad Paarlberg (Dreg. !), distr. Worcester (Eckl. et Zeyh. !). Thunb. fl. cap. 729. Lam. ill. t. 723. f. 1. Less. l. c. — Burm. afr. t. 70. f. 1. Pluk. alm. t. 272. f. 5. mal. et ideò *C. Africanum* Lin. sp. 1317. Burm. ! prod. 25. sed *C. glabrum* eodem nom. insignitum adest. Folia 3-5 lin. lata, basi canaliculata. (v. s.)
5. *C. villosæ* (Lin. f. suppl. 392), caule tereti muricato, foliis convolato-semiteretibus glabris, involucri cylindrici squamis extùs muricatis. \mathcal{U} ad Cap. B. Spei in monte Tabulari et aliis circa urbem locis (Burch. ! n. 6663. Dreg. ! Eckl. et Zeyh. !). Less. l. c. Thunb. in fl. cap. fortè non immeritò cum priora conjunxit. *C. gramineum* Burm. ! prod. fl. cap. 26. (v. s.)
6. *C. luteum* (E. Mey. ! in Dreg. coll. n. 1527), caule tereti muricato subætuloso, foliis lineari-filiformibus glabris, invol. oblongi subtrunati squamis vix subscabris. \mathcal{U} ad Cap. Bonæ Spei circa Paarlberg legit cl. Dregel Corollæ luteæ. Habitus *C. filiformis* sed invol. squamæ latiores magis foliaceæ vix ac ne vix scabræ. (v. s.)
7. *C. villosum* (Less. in linnæa 1831. p. 691), caule hirsuto, foliis planis utrinquè muricatis et hirsutis, invol. squamis muricatis subvillosis. \mathcal{U} ad Cap. Bonæ Spei. Burch. ! n. 6760. Dreg. ! Eckl. et Zeyh. ! *C. villosum* et *C. hirsutum* Thunb. fl. cap. 730 et 729. *C. hirsutum* Ecklon ! pl. exs. cap. in h. Durrant. Variat plùs minùs hirsutum et æqm pili pauciores magis muricatum evadit. (v. s.)
8. *C. congestum* (E. Mey. ! in Dreg. coll.), caule hispido, foliis elliptico-lanceolatis utrinquè attenuatis submembranaceis utrinquè hirtellis, corymbo congesto, involucri squamis hirtellis. \mathcal{U} in Cap. Bonæ Spei ad Drakeenstenberg alt. 2-300 ped. legit cl. Dregel ! et in distr. Worcester cl. Ecklon et Zeyher. Folia infer. pollicem, imò sesquipollicem lata. (v. s.)

XLI. ROLANDRA *Rottb. coll. hafn. 2. p. 256. Swartz fl. ind. occ. 1388. t. 17. Cass. dict. 46. p. 170. Less. in linn. 1829. p. 332. f. 44. 75. syn. p. 150. — Echinopsis sp. Lin.*

Capitula 1-flora super receptaculum generale piloso-fimbrilliferum globosum adgregata. Invol. 2-valve compressum, valvis membranaceis carinatis aristatis, exteriore longiori. Cor. regularis glabra 3-5-fida, sæpius 4-fida, limbo subinflato. Stam. filamenta lævia. Styli rami semi-subulati. Achænium oblongo-teretiusculum 4-costatum glabrum glandulosum. Pappus 1-serialis coroniformis. brevis dentatus, dentibus ciliatis. — Suffrutex. Folia alterna breve petiolata oblonga utrinquè attenuata suprâ glabra muriculata subtus araneoso-villosa alba. Capitula generalia globosa ramulos terminantia sæpè ramulis abbreviatis falso-axillaria. Cor. alba.

1. R. ARGENTEA (Rottb. l. c.). $\frac{3}{4}$ in subhumidis Jamaicae, Guadalupe, Surinam, Cajennæ, Bahiæ, etc. Sloan. hist. jam. 1. p. 43. t. 7. f. 3. Plum. ed. Barm. t. 123. f. 1. Echinops fruticosus Lin. Echinops nodiflorus Lam. dict. p. 337. Varias foliis oblongis ovalibusve semper acuminatis, subtus plus minus sericeo-argenteis. Variat adhuc cor. lobis 3, 5 sæpius 4; involucri valvulâ minore nunc longiusculè nunc brevè aristatâ, nunc muticâ, in speciminibus cæterum simillimis, undè *R. monocantha* et *R. diacantha* Cass. dict. 46. p. 171. vix ut varietates habendæ. (v. s.)

† *Species exclusæ.*

R. REPTANS Willd. = *Trichospira menthoides*.

R. TERMINALIS Spreng. = *Bailliera commelinoides*.

XLII. SPIRACANTHA *H. B. et Kunth nov. gen. am. 4. p. 28. t. 313. Less. syn. 150. — Acosta DC. mss. 1810.*

Capitula 7-11 uniflora adgregata in glomerulum ovatum bracteatum et ad bractearum axillas solitaria sessilia. Bracteæ subdisticho-imbricatæ basi concavæ in spinam desinentes. Involucrum 4-5-phyllum, foliolis lanceolatis diaphano-membranaceis subæqualibus. Cor. regularis, tubo gracili, fauce infundibuliformi; lobis 5 lineari-lanceolatis. Stylus apice bifidus? Achænium obovato-cuneatum læve. Pappus squamellulis pili-formibus brevibus integris scabridis constans. — Suffruticulus ramosissimus. Ramuli subpubescentes filiformes. Folia alterna brevè petiolata oblonga utrinquè attenuata integerrima suprâ glabra subtus pube adpressâ albâ. Glomeruli ad apices ramulorum axillarium et ad apicem ramorum. Cor. purpureæ. — Genus affine *Rolandræ*.

1. S. CORNIFOLIA (H. B. et Kunth l. c.). $\frac{3}{4}$ in umbris humidis Novæ Granatæ ad portum Sapote prope Rio Sinu. Acosta *Rolandræ* DC. mss. (v. s. comm. à cl. Bonpland.)

XLIII. TRICHOSPIRA *H. B. et Kunth nov. gen. am. 4. p. 23. t. 312. Less. in linnæa 1829. p. 342. syn. p. 150.*

Capitulum multiflorum seu glomerulus polycephalus, bracteis è rachide ortis margine scariosis apice fimbriatis basi 1-nerviis apice 3-nerviis, in axillâ florem unicum foveantibus sed involucrum verum subnullum. Cor. regularis glabra, tubo tereti, fauce campanulatâ, lobis 4 obtusis. Antheræ è fauce semiexsertæ. Styli rami exserti semisubulati. Achænium compressum oblongo-obovatum basi acuminatum puberu-

lum in medio utrinquē 3-costatum. Pappus 1-serialis, aristis nunc 2 divaricatis nunc insuper 4-5 prioribus adjectis multò minoribus. — Herbæ humiles Americanæ Filaginis facie. Radicellæ pubescentes! Folia alterna aut opposita subtus lanâ albâ tomentosa sessilia integra penninervia superiora opposita. Capitula axillaria et oppositifolia sessilia. — Genus affine Rolandrae et Spiracanthæ, Lessingii subtribus.

* *Pappo pluriaristato.*

1. *T. MENTHOIDES* (H. B. et Kunth l. c.), pappo pluriaristato, caule repente procumbente, foliis fere omnibus auriculato-semiamplexicaulibus. ② in arenosis fluvii Apures prov. Varinensis et in Cubâ. Less. in linnæâ 1829. p. 342. f. 63. Rolandra reptans Willd. herb. ex Less.
2. *T. PULEGIUM* (Mart. ! herb.), pappo pluriaristato, caulibus à collo pluribus diffusis non radicanibus, foliis fere omnibus auriculato-semiamplexicaulibus. ② in herbidiis prov. Rio-Negro Brasiliæ fluv. Japura conterminis legit cl. Martius. Valdè accedit ad *T. menthoidem* sed ex omni parte major, multicaulis, non repens et aristæ fructus paulò longiores, Squamæ paucissimæ involucri loco. (v. s.)

** *Pappo biaristato.*

3. *T. BIARISTATA* (Less. ! l. c. f. 57. 58. 65. 66. 70), pappo biaristato, caule erecto aut vix adscendente, foliis alternis subtus tomentosis, superioribus tantum auriculatis. (①) ② ? in Brasiliâ tropicâ. Priori paulò minor. (v. s. comm. à Mus. reg. Berol.)
4. *T. PRIEURÆ*, pappo biaristato, caule erecto, foliis sinuato-dentatis suprâ glabris subtus canescenti-tomentosis, infer. alternis obovatis in petiolum attenuatis, super. oppositis basi dilatato-auriculatis. (⊙) in Guianâ Gallicâ ad ripas flum. Oucassa, in solo argillaceo paludoso legit cl. Le Prieur, qui ex naufragio miserrimo hanc et paucas alias salvavit! Herba pedalis. (v. s. comm. à cl. inv.)

XLIV. *LAGASCEA* H. B. et Kunth nov. gen. am. 4. p. 24. Less. in linnæâ 1829. p. 333. f. 36-38. 45. 46. 48. 78. — *Noccæa* Cass. dict. 25. p. 102. non Mœnch.

Capitula 1-flora in glomerulum generale aggregata. Involucrum (proprium) 1-seriale gamophyllum 4-5-dentatum tubulosum. Cor. regularis extus pubescens, tubo gracillimo, fauce dilatata cylindracea, lobis 5 acuminatis. Stam. filam. papillosa. Styli rami semisubulati. Achænium obcompressio-tetragonum pluricostatum. Pappus 1-serialis coroniformis membranaceus brevissimus nunc subinteger nunc dentatus. — Herbæ aut suffrutices Americanæ. Folia opposita sæpius triplinervia. Capitula generalia terminalia bracteis plurimis (exterioribus oppositis) obvallata. Flores albi, flavidi aut rubri. — *Noccæa* nomen quidem vetustius (1794), sed Kunth, qui primus genera duo conjunxit, *LAGASCEA* nomen admisit ex usu hortulanorum. Cur autem cl. Lessing qui genera distinxit eorum nomina transposuit, ignoro.

Sect. I. *LAGASCA* Cav. anal. sc. nat. 6. (1800) p. 333. t. 44. et omn. fere auct. excl. Less. — *Noccæa* Jacq. fragm. (1805) p. 58. t. 85. f. 1. Less. syn. 151. non Mœnch.

Pappus coroniformis minimus subinteger ciliatus. — Herbæ annuæ. Glomerall ad apicem pedunculi elongati nudi solitarii. Folia petiolata. Cor. albæ.

1. *L. MOLLIS* (Cav. anal. sc. nat. 6. p. 333. t. 44), herbacea, foliis petiolatis ovatis acuminatis subcrenatis molliter sericeo-tomentosis, capitulis generalibus solitariis longè pedunculatis. (①) in collibus insulæ Cubæ et ad Santa-Fé Mexicæ.

norum, Desv. Journ. bot. 1. p. 28. t. 2. Noccæa mollis Jacq. frag. t. 13. Flores albi. Involuelli foliola concreta 3-nervia. Capitula et bractæ sericeo-villosæ. (v. v.)

Sect. II. NOCCÆA. — *Nocca Cav. ic. 3. (1794) p. 12. t. 224. non Less.*
— *Lagascea Less. syn. 131. non Cav.*

Pappus paleis brevibus subdentatis. — Suffrutices aut herbæ perennes à Mexico. Glomeruli adgregati foliis bracteati. Folia sessilia aut vix petiolata. Cor. albidæ ochroleucæ aut purpurascences.

2. *L. RUBRA* (H. B. et Kunth nov. gen. am. p. 24. t. 311), suffruticosa, foliis breviter petiolatis ellipticis ovatisve subacutis dentatis coriaceo-rigidis scabris, capitulis glomerato-corymbosis. $\frac{1}{2}$ in temperatis Mexici ad montes Guchilapa, etc. *Nocceæ rubra* Cass. l. c. *Nocca rigida* Cav. ic. 3. p. 12. t. 224. Sweet! br. fl. gard. ser. 2. t. 26. Flores pallidè rubentes. (v. s.)
3. *L. MOQUINIANA* (DC. in Moq. ic. ined. mexic. 1814), suffruticosa, foliis petiolatis ovatis ovalibusve integerrimis suprâ piloso-scabridis subtus villosis, paniculæ ramis distantibus oppositis, invol. generalis squamis 4-6 lineari-lanceolatis flores superantibus. $\frac{3}{4}$ in Mexico observ. cl. Moquinno et Hænke. Huc referenda ex cl. Moquinno Caçanaca Hern. mex. 184. ic. sed foliis sessilibus et glomerulorum dispositione differt et potius *Lagasceæ* species hodiernis ignota! In nostrâ ex ic. floræ Mexic. flores purpurei. Caulis rami et involucri apice glanduloso-villosa; invol. sericeo-villosissima. Folia 2-2 $\frac{1}{2}$ poll. longa, pollicem lata. (v. ic. pict. Moq. et sp. sicc. in h. Hænke.)
4. *L. LATIFOLIA*, fruticosa, foliis sessilibus amplexicanlibus oblongis acuminatis serratis integerrimisve suprâ scabris subtus molliter pilosis reticulatis, capitulis corymboso-paniculatis. $\frac{3}{4}$ in Mexico. *Nocca latifolia* Lallav. et Lex. nov. veg. descr. 1. p. 31. Sweet br. fl. gard. t. 215. Flores suaveolentes ochroleuci aut albidî. Cor. ferè pollicem longæ. Styli longè exserti. (v. s. ex hort. Mexic. ubi Cervantez eam descr.)
5. *L. HELIANTHIFOLIA* (H. B. et Kunth l. c.), suffruticosa, foliis sessilibus amplexicanlibus oblongis subacuminatis cordatis serratis coriaceo-rigidis suprâ scaberrimis subtus scabrinsculo-hirtellis, capitulis subcorymbosis. $\frac{3}{4}$ juxta Acapulco Mexic. *Nocceæ helianthifolia* Cass. l. c. *Nocceæ macrophylla* Zucc. in h. Acad. reg. Monac. Cor. albidæ? pollicem ferè longæ. Folia 7-8 poll. longa, 2 lata. Involucellum externè sericeo-pilosum. (v. s. comm. à cl. Mairet.)
6. *L. SUAVEOLENS* (H. B. et Kunth l. c.), suffruticosa, foliis sessilibus amplexicanlibus oblongis acuminatis supernè serratis coriaceo-rigidis suprâ scaberrimis subtus secus nervos valdè prominentes molliusculè puberulis inter eos aparsè subscabridis, capitulis corymboso-paniculatis. $\frac{3}{4}$ in declivitate orientali montium Mexicanorum. *Nocceæ suaveolens* Cass. l. c. Cor. alba. Folia 4 poll. longa, 15 lin. lata more *Borraginearum* demum suprâ albo-punctata. (v. s. in h. Hænk. à cl. Sternb. comm.)
7. *L. ANGUSTIFOLIA*, suffruticosa, foliis brevè petiolatis oblongo-linearibus integerrimis utrinquè attenuatis villosis-scabris, capitulis villosissimis subfasciculatis. $\frac{3}{4}$ in Mexico circa Leon legit cl. Mendez. Cor. ex statâ immaturo 3 lin. longæ albidæ. Involucellum villosissimum 5-dentatum, dentibus subulatis. Folia 3-4 poll. longa semipollicem lata non triplinervia. (v. s.)

Div. IV. BOJERIEÆ.

Capitula pluriflora homogama. Anthæræ basi bicandatæ. Pappus setis 1-seriatis. — Arbores frutices aut herbæ Madagascarienses, omnes à cl. Bojer detectæ.

XLV. SYNCHODENDRON Bojer in litt. 1833.

Capitulum pluriflorum homogamum (ex Bojer), recept. angusto nudo, invol. obconici squamis multiseriatis imbricatis coriaceis siccis gla-

bris, exter. brevissimis, intimis longioribus linearibus. Cor. tubulosæ æquales 5-fidæ. Anthæræ longè caudatæ exsertæ. Styli rami breves glabriusculi. Achænia teretiuscula subvillosa. Pappus uniserialis, setis rigidulis integris sublaminatis cor. longitudine æquantibus. — Arbor Madagascariensis 50-pedalis Compositarum omnium maxima! Rami juniores petioli et imò folia pube brevi velutino-tomentosâ oblecta. Folia alterna petiolata integra orbiculata aut obovata, demùm supernè glabra. Capitula ante folia secus ramos juniores et crassos sessilia fasciculata, 2 lin. longa. (Bojer.)

1. *S. RAMIFLORUM* (Bojer! in litt. 1833 et 1835). § in insulæ Madagascar prov. Emirnensi in vallibus inter montes excelsos et circa pagos cultum. Florescentia hujus arboris Madagascariensibus tempus oryzae culturæ proprium indicat (Bojer). (v. s. comm. à cl. inv.)

XLVI. CENTAUROPSIS Bojer in litt. 1833.

Capitulum pluri- (12-20-) florum homogamum. Recept. angustum areolatum paleis linearibus paucis deciduis onustum. Invol. oblongum arcuè imbricatum, squamis coriaceis obtusis, ext. brevissimis, inter. elongatis ferè linearibus. Cor. tubulosæ, fauce vix dilatata, limbo 5-dentato, exter. incurvæ majores. Anth. loculi basi producti in appendices polliniferas? Stigm. 2 longa exserta divergentia recurvato-reflexa setis hispida. Achænium teretiusculum gracile striatum glaberrimum erostre. Pappus 1-serialis è margine calyculi coroniformis achænium coronantis ortus rigidus, setis inæqualibus achænio duplò triplòve brevioribus, in margine capituli brevioribus aut subnullis. — Frutices Madagascarienses sylvicolæ. Folia alterna breviter petiolata subintegerrima obovata aut ovalia basi cuneata penninervia. Capitula corymbosa aut solitaria ovato-oblonga. Flores ex sicco purpurascens. — Genus affine Stokesiæ, Platycarpæ et Distephano.

1. *C. LAMUGINOSA* (Bojer! in litt.), foliis latè obovatis obtusissimis suprà adultis glabriusculis, subtus ramulis pedunculisque densè velutino-tomentosis, capitulis laxè corymbosis. §? §? in insulæ Madagascar sylvis vastis loco Bé-Fou-roun dicto detexit cl. Bojer! Folia 6 poll. longa, 4 poll. lata, basi distinctè cuneata, suprà jun. furfuraceo-tomentosa, adulta punctulis elevatis subscabrida. Invol. squamæ exteriores dorso velutinæ. Flores in invol. 12-13. Cor. extus incurvæ. (v. s. comm. à cl. inv.)
2. *C. FRUTICOSA* (Bojer! in litt.), ramulis junioribus vix pubescentibus, foliis ovalibus involucriisque glaberrimis, pedunculis solitariis angulatis. § ad margines sylvarum in prov. Emirnensi inter. ins. Madagascari detex. cl. Bojer. Priori multò minor. Folia pollicem longa, 5-6 lin. lata. Pedunculi folii circ. longitud. æquales 1-cephali sub capitulo bracteam foliaceam ovalem sessilem gerentes. Fructus exactè prioris sed pappus paulò longior et magis caducus. Flores vix vidi, ex fruct. circ. 20 in invol. (v. s. comm. à cl. Bojer.)

XLVII. TECMARSIS.

Capitulum circ. 6-florum homogamum. Recept. angustum nudum. Invol. oblongum ante explicationem acutum, squamis arcuè imbricatis, exter. brevibus obtusis, inter. linearibus subacutis. Cor. tubulosæ filiformes inconspicuæ.... Anth..... Stigm..... Achænia gracilia subangulata striata glabrâ erostria. Pappus 1-serialis, setis basi in tubum concretis subinæqualibus erectis subrigidulis achæniū longitudine æqualibus invol. ferè superantibus. — Frutex Madagascariensis debilis,

ramis teretibus procumbentibus apice setoso-ciliatis villosisque. Folia alterna fasciculata glabra obovato-oblonga acuta basi cuneata paucidentata, petiolis ciliato-villosis brevibus. Capitula corymbosa, pedunculis ramosis nudis. Invol. glabra. — Genus ex floribus ignotis dubium, et ideo Tecmarsis (græcè conjectura) dictum, fortè ex invol. et pappo ad Centauropsidem et Synchodendron accedens.

1. T. BOJERII. ♂ ad margines sylvarum in locis umbrosis insulæ Madagascar legit cl. Bojer, qui mecum sine nomine commun. Rami et folia in sicco non malè Bucidam referunt. (v. s. comm. à cl. invent. sed capitula alia juniora minima alia nimis etate proventa!)

XLVIII. BOJERIA.

Capitulum multiflorum homogamum, fl. tubulosis 5-fidis hermaphroditis. Invol. campanulatum polyphyllum, squamis multiseriis longitudine subæqualibus acutis, ext. lineari-lanceolatis foliaceis dorso villosis, int. linearibus glabriusculis. Recept. planum subalveolatum. Anth. basi bicaudatæ. Stylus basi bulbosus, ramis exsertitis teretibus pube minimè scabridis. Achænia striato-subangulata glabra. Pappus 1-serialis, setis longis subscabris basi inter se variè et brevè subconcretis. — Herba perennis elata Madagascariensis. Caulis teres ramosus præsertim ad apices dense villosus. Folia caulina alterna amplexicaulia ovato-lanceolata integra suprè sparsè puberula subtus subtomentosa. Capitula ad apices ramorum solitaria. Flores numerosissimi. Cor. purpureæ. Calyx in achænio facilè à pericarpio separatur et pappum secum trahit! Pericarpium teres apice cupulam exiguum styli basi per anthesin foveentem gerit. — Genus spectabile Eupatoriaceis stylo subaffine, sed habitu et antheris Vernoniaceis referendum, dicavi cl. Guill. Bojer, hist. nat. in insulâ Mauritiî professori, plantæ detectori et qui insulas Mauritanas, Madagascar et alias Australi-Africanas laboribus indefessis egrediè explorat.

1. B. SPECIOSA. in locis humidis insulæ Madagascar circa montes Angavon dictos detexit cl. Bojer. Herba interdum 10-pedalis. Capitula diam. pollicari. (v. s. comm. sine nomine à cl. invent.)

Subtrib. II. PECTIDÆ Less. in linnæd 1830. p. 134.

Capitula radiata nec discoidea. Styli flor. herm. rami apice obtusi nec acuminati. Folia sæpius opposita. Herbæ aut frutices. Species omnes Americanæ. Flores sæpè flavi.

DIV. 1. LIABÆ Cass. tabl. p. 28. Less. in linnæd 1831. p. 696. syn. p. 151. — Vernonicarum gen. Cass. Less. in linnæd 1829. p. 318.

Cor. disci regulariter 5-fidæ. Stylus fl. herm. ramis longiusculis plerumque obtusis. — Herbæ seu frutices. Folia sæpius opposita eglandulosa basi ciliis destituta, subtus tomentosa.

XLIX. XANTHISMA.

Capitulum multiflorum radiatum, fl. radii ligulatis neutris? seu femineis? integris, disci hermaphroditis 5-fidis lobis erectis. Recept. fimbriiferum, fimbriis laceris achænio ferè longioribus. Invol. hemisphæricum, squamis imbricatis coriaceis ferè ovalibus obtusissimis

adpressis. Anthæræ (disci) ecaudatæ. Stylus radii brevis simplex intra partem ligulæ tubulosam inclusus, disci apice bifidus, lobis linearibus inclusis obtusis breviter hispidulis. Achænium obovatum vix subangulatum pubescens. Pappus paleis elongatis acuminatis inæqualibus, ext. paulò brevioribus et angustioribus. — Herba Mexicana monocarpica erecta glabra simplex apice laxè ramosa subcorymbosa, ramis erectis foliosis. Folia caulina alterna sessilia oblonga aut linearia uninervia rigidula, pleraque integerrima, pauca apice acutè serrata. Flores lutei.

1. *X. TEXANUM*. ① ② in sylvis Mexicanæ provinciæ Texas legit cl. Berlandier (pl. exs. n. 2639). Herba habitu ad *Centauream cerinthefoliam* ferè accedens. (v. s.)

L. HECTORÆA.

Capitulum multiflorum radiatum, ligulis fœm. 12-15 acutè 3-dentatis. Invol. pluriseriale, squamis acuminatis subimbricatis. Recept. alveolatum. Cor. disci glabræ 5-fidæ. Stylus teres nec basi bulbosus, ramis terebribus hispidulis. Achænium teres villosus. Pappus biserialis, serie ext. paleaceâ brevi, inter. elongatâ setiformi scabrâ. — Herba Mexicana basi suffruticosa simplex erecta tota cano-villosa. Caulis teres ad apicem usquè foliosus. Folia alterna conferta erecta oblongo-lanceolata integra mucrone calloso superata. Capitula ad apicem caulis breviter pedicellata subcorymbosa flava. Pappi series ext. alba, int. subrufa. — Cor. *Andromachiæ* sed habitus diversissimus, stylus non bulbosus, etc. Nomen ad *Hectoris* et *Andromachiæ* affinitatem alludit.

1. *H. VILLOSISSIMA*. $\frac{1}{2}$ $\frac{3}{4}$ in Mexico prope Real del Monte legit cl. Hænke. Caulis pedalis. Folia 10-12 lin. longa, 2-3 lin. lata.

LI. *ANDROMACHIA* *H. et Bonpl. pl. eq. 2. p. 104. t. 112. Less. in linnæâ 1829. p. 318. non Cass.* — *Andromachiæ* sect. 2 et 3. *H. B. et Kunth nov. gen. 4. p. 99.* — *Oligactis Cass. dict. 36. p. 16.* — *Diplostephii sp. Spreng.* — *Viviania Willd. herb. non Cav. nec Colla.*

Capitulum radiatum, disco regulari multifloro, radio uniseriali ligulato fœmineo. Invol. squamæ imbricatæ siccæ acuminatæ. Recept. breviter paleaceo-fimbriliferum. Cor. glabræ, disci 5-fidæ, lobis acuminatis revolutis tubo paulò brevioribus, radii ligulatæ, tubo longo, ligulâ oblongâ. Filam. lævia. Styli basi bulbosi rami semicylindracei. Achænium cylindraceum multicostatum, valliculis hirsutis, nectario styliformi. Pappus 2-serialis paleaceus serratus, paleis seriei ext. multò brevioribus et sublterioribus quam inter. — Herbæ aut frutices. Folia opposita petiolata subtus tomentosa dentata. Capitula corymbosa.

Sect. I. PLEIONACTIS.

Capitulum multi (20-30)-florum. Ligulæ radii crebræ. — Herbæ, caulibus foliisque subtus cano-tomentosis. Flores ligulati flavi. — *Andromachiæ* veræ *Kunth.*

1. *A. GRANDIFLORA* (*H. B. et Kunth l. c. p. 99*), herbacea ramosa, foliis ovatis acutis leviter cordatis suprâ glabris rugoso-bullatis subtus albo-lanatis, petiolis alatis basi auriculatis et connatis, capitulis 3is-4is-ve corymbosis. $\frac{1}{2}$ in aridis prope urbem Alaui Quitensium. *Diplostephium grandiflorum* *Spreng. syst. 3. p. 544.*
2. *A. IGNIARIA* (*H. et Bonpl. pl. eq. 2. p. 104. t. 112*), herbacea ramosa, foliis ovatis acutis basi obtusis suprâ glabris subtus albo-lanatis, petiolis basi auricu-

latis et connatis, capitulis fasciculatis in corymbos terminales trifidos dispositis. $\frac{1}{2}$ in altâ planitie Quitensi. H. B. et Kunth nov. gen. 4. p. 100. Less. in linnæâ 1829. p. 318. Amellus floribundus Willd. herb. ex Less. *Diplostephium igniarum* Spreng. l. c. *Liabum Bonplandii* Cass. dict. 26. p. 206. *Tomentum Agarici igniarum* succedaneum.

3. *A. SOLIDAGINEA* (H. B. et Kunth l. c. p. 100), herbacea ramosa, foliis ovato-oblongis acuminatis basi rotundatis et in petiolum basi nudum decurrentibus suprâ glabris subtus albo-tomentosis, corymbis terminalibus ramosis, ramulis fasciculatis. $\frac{1}{2}$ in Andibus Peruvianorum inter Ayavaca et fl. Cutaco. *Diplostephium solidagineum* Spreng. syst. 3. p. 544. (v. s. in h. reg. Acad. Monac. specimen Peruv. à descr. Kunth discrepans tantum tomento paginæ inf. fol. in sicco rufo nec albo.)
4. *A. VERBASCIFOLIA* (H. B. et Kunth l. c. p. 100), herbacea ramosa, foliis oblongis acutis basi in petiolum nudum angustatis suprâ rugoso-bullatis glabris subtus lacunosus cano-lanatis, corymbis terminalibus ramosissimis, capitulis fasciculatis. $\frac{1}{2}$ in montibus Quitensium cum Cinchonâ inter Malacates et Gonzanama. *Diplostephium verbascifolium* Spreng. l. c.
5. *A. MELASTOMOIDES* (H. B. et Kunth l. c. p. 101. t. 337), herbacea ramosissima, foliis oblongo-ellipticis acuminatis basi acutis coriaceis suprâ scabris subtus cano-tomentosis, petiolis nudis, pedunculis axillaribus et terminalibus polycephalis. $\frac{1}{2}$ in temperatis prope Ibague Novo-Granatensium. Less. in linnæâ 1829. p. 318. f. 1-3. 6. *Diplostephium melastomoides* Spreng. l. c. *Viviania bicolor* Willd. herb. ex Less.

Sect. II. OLIGACTIS H. B. et Kunth nov. gen. 4. p. 101.

Capitulum pauciflorum. Ligulæ radii 3-7. — Frutices, ramis glabris. Flores ligulati pallidi fortè albi?

6. *A. NUBIGENA* (H. B. et Kunth nov. gen. 4. p. 102), fruticosa, foliis lanceolato-oblongis acutis basi obtusis membranaceis suprâ petiolisque glabris subtus nervo medio excepto albo-tomentosis, corymbis terminalibus ramosis, capitulis pedicellatis. $\frac{1}{2}$ in excelsis montis Chimborazo. *Diplostephium nubigenum* Spreng. syst. 3. p. 543. *Oligactis nubigena* Cass. dict. 36. p. 17.
7. *A. SESSILIFOLIA* (H. B. et Kunth l. c.), fruticosa, foliis oblongis acutis basi rotundatis membranaceis suprâ glabris subtus petiolisque albo-tomentosis, paniculis axillaribus et terminalibus, capitulis conglomeratis sessilibus. $\frac{1}{2}$ in Peruvia? *Diplostephium sessiliflorum* Spreng. l. c. *Oligactis sessiliflora* Cass. l. c.
8. *A. VOLUBILIS* (H. B. et Kunth l. c.), volubilis, foliis lanceolato-linearibus acutis brevè petiolatis suprâ glabris subtus cano-tomentosis, paniculis terminalibus simplicibus, capitulis solitariis pedicellatis. $\frac{1}{2}$ in monte Antisanâ Quitensium. *Diplostephium volubile* Spreng. l. c. *Oligactis volubilis* Cass. l. c.

† Species exclusæ.

- A. ACAULIS* Kunth. = *Liabum acaule*.
A. HIERACIOIDES Kunth. = *Liabum hieracioides*.
A. JUSSIÆI Cass. = *Liabum Jussæi*.
A. POTRAVI Cass. = *Liabum Brownei*.

LII. *LIABUM* Adans. fam. 2. p. 131. excl. sp. Cass. dict. 26. p. 203. excl. sp. Less. in linnæâ 1829. p. 319. — *Andromachia* sp. Kunth. — *Andromachia* Cass. bull. philom. 1817. p. 184. non Bonpl. — *Amelli* sp. Linn. — *Starkea* Willd. sp. 3. p. 2216. — *Starckia* Juss. dict. 50. p. 403.

Capitulum radiatum, disco regulari multifloro, radio uniseriali liga-

lato fœmineo. Invol. squamæ imbricatæ subulatæ. Recept. fimbriilis membranaceis subulatis hispidum. Cor. glabræ, disci 5-fidæ regulares, lobis 5 linearibus tubo brevioribus, radii ligulæformes longæ lineares apice integræ. Styli (disci) rami semiteretes graciles parçè hispidi. Achænium cylindraceum striatum, callo basilari instr. Pappus 1-serialis setis filiformibus barbellatis subinæqualibus. — Herbæ Americane sæpè cano-lanatæ, caulescentes aut acaules. Folia opposita triplinerviâ subdentata. Flores flavi.

§. 1. *Capitulis plurimis corymbosis.* — *Starkea Willd. l. c.*

1. L. BROWNEI (Cass. dict. 26. p. 203), caule erecto infernè folioso tomentoso, foliis ovatis subserratis in petiolum attenuatis suprâ lanuginosis demùm glabris subtùs tomentosis, cymâ umbellatâ terminali, pedunculis elongatis aliis monogaliis oligocephalis. 2/ in Jamaicâ et Santo-Domingo. Solidago P. Brown. jam. 320. t. 33. f. 2. Amellus umbellatus Linn. sp. 1276. Swartz! obs. 310. *Starkea umbellata* Willd. sp. 3. p. 2216. *Andromachia Poiteavi* Cass. bull. philom. 1817. p. 184. Variat foliis obtusis et acutis. (v. s.)
2. L. JUSSIEI (Cass. dict. 26. p. 205), caule erecto glabrinsculo subanguloso, foliis petiolatis ovalibus dentatis basi stipulato-auriculatis suprâ glabris subtùs tomentosis, capitulis in cymam paniculato-corymbosam dispositis. 2/ in Peruvîâ. *Andromachia Jussiei* Cass. bull. philom. 1817. p. 184. *Conyza stipulata* Vahl in herb. Juss.

§. 2. *Capitulis solitariis.* — *Chrysactinium H. B. et Kunth l. c.*

3. L. HIERACIOIDES, caule procumbente cano-lanato, foliis sessilibus connatis lanceolatis acutis subdentatis suprâ pilosis subtùs cano-lanatis, pedunculis terminalibus 1-cephalis. 2/ in siccis temperatis prope Loxam Quitensium. *Andromachia hieracioides* H. B. et Kunth nov. gen. am. 4. p. 98. Solidago hieracioides Spreng. syst. 3. p. 543.
4. L. ACAULE, caule subnullo, foliis radicalibus sessilibus lanceolato-oblongis basi angustatis remotè dentatis suprâ glabris subtùs densè candido-lanatis, scapo erecto longissimo 1-cephalo lanato. 2/ in montibus Quitensibus. *Andromachia acaulis* H. B. et Kunth nov. gen. 4. p. 98. t. 336. Solidago acaulis Spreng. syst. 3. p. 543.

+ *Species exclusa.*

L. BONPLANDII Cass. = *Andromachia igniaria*.

LIII. ALIBUM Less. syn. 152.

Omnia Liabi sed pappus disci biserialis, serie exteriori coroniformi dentatâ, interiori multisetacæ: radii coroniformis dentatus, 1-serialis. — Herba trichotoma erecta Australi-Americana. Rami teretes hirsuti. Folia opposita obsolete auriculata et connata ovato-elliptica integrâ seu basi pinnatifida suprâ hirsutiuscula subtùs tenuiter albo-tomentosa. Capitula solitaria nutantia. Invol. squamæ biseriales æquales oblongo-ellipticæ mucronatæ plurinerviæ. Recept. planum nudum. Achænia turbinata hirsuta.

1. A. LIABOIDES (Less. l. c.). — in Pomelacte Americæ æquinoctialis.

LIV. CACOSMIA H. B. et Kunth nov. gen. 4. p. 289. t. 404. Cass. tabl. xx. n. 4. Less. in Linnæâ 1829. p. 338. f. 31. 53-55. syn. 152. — *Xantholepis Willd. herb.*

Capitulum radiatum, floribus disci plurimis hermaphr., radii ligulatis fœmineis. Invol. teres imbricatum, squamis siccis plurinerviis

interioribus longioribus. Recept. nudum. Cor. glabræ, disci regulares 5-fidæ, lobis acuminatîs revolutis parte integrâ brevioribus, radii ligulâ latè oblongo-ellipticâ apice 3-crenulatâ tubo duplò longiore. Filam. lævis. Styli rami semicylindræci. Achænium obpyramidatum 4-quetrum truncatum glaberrimum, nectario styliformi. Pappus o. — Herba suffruticosa graveolens. Rami tenuiter lanati. Folia opposita subpetiolata connata ovato-oblonga remotè dentata trinervia suprâ rugosa glabra subtus pubescenti-cana. Corymbi ad apices ramorum axillarium. Cor. flavæ.

1. *C. RUGOSA* (H. B. et Kunth, l. c.). 2½ in Andibus Peruviz. Xantholepis tomentosa Willd. herb. ex. Less. l. c.

Div. II. EUPECTIDÆ. — Pectidæ Less. in *linnæâ* 1831. p. 706. *syn.* p. 152. — Tagetineæ Pectidæ pleræque Cass. *tabl.*

Corolla disci subbilabata aut rarissimè regulariter 5-dentata. Styli disci rami semicylindræci breves. — Herbæ glabriusculæ pleræque annuæ Americanæ. Folia opposita basi lineâ transversali subconnata pellucido-punctata uninervia basi denticulis setigeris ciliata, apice variis subserrulata. Invol. 1-seriale. Recept. nudum. Cor. flavæ glabræ nunc regulares 5-dentatæ nunc lobo unico profundius fissæ subbilabiata, lobo exteriore indiviso, inter. 4-dentato. Antheræ ecadatæ. Stylus radii glaber. Achænia angulata basi attenuata, callo basilari. Pappus flavescens. — Pectidæ à cl. Cassini ad Tagetineas relatæ, à cl. Kunth inter Tagetem et Steviam mediæ, à cl. Lessing juxta Liabeas in Veronicearum tribu admisæ.

LV. PECTIDOPSIS.

Cor. disci regulariter 5-dentatæ. Pappus paleaceus brevis 5-dentatus. — Herba boreali-americana basi ramosa diffusa lævis. Folia angustolinearia basi dentato-ciliata. Capitula solitaria brevissimè pedicellata. Invol. 8-phyllum. Radii ligulæ 7-8. — Habitus Pectidis sed pappo eodem jure ac Pectidium et Lorentea separanda.

1. *P. ANGUSTIFOLIA*. (1) in Americ. bor. montibus dictis Rocky-Mountains. Pectis angustifolia Torr. acc. rock. mount. in ann. Lyc. N.-York. 2. p. 214.

LVI. PECTIDIUM Less. in *linnæâ* 1831. p. 706. *syn.* p. 152. — Pectis Cass. *dict.* 38. p. 202. non Less. — Pectidis sp. Linn.

Cor. disci regulariter 5-dentatæ. Pappi setæ subulatæ rigidæ corneæ læves nudæ demum divergentes. — Herba annua glabra erecta di- aut trichotoma. Folia linearia cilia utrinquè 1-2 basi gerentia glandulis grossis pellucidis ad utrumque nervi medii lateris ornata. Capitula longè pedicellata solitaria. Invol. 5-phyllum. Ligulæ breves.

1. *P. PUNCTATUM* (Less. in *linnæâ* 1831. p. 707). (1) in insulis Caribæis nempe Santo-Domingo (Bertero!), Santo-Thomasio (Ehr.), Sancta-Martha (West!), Sanctæ-Crucis (Jacq.), Porto-Ricco (Ledra!). Pectis punctata Jacq. amer. 216. t. 128. Linn. sp. 1230. Swartz obs. p. 308. Linn. fil. herb. Pectis linifolia Gærtn. fr. 2. t. 171 (excl. syn.). Lam. ill. t. 684. — Plum. et Burm. t. 86. f. 1? Sloan. jam. hist. 1. t. 149. f. 3? sed icon utraque pessima et fortè removenda. Conf. Pectidem linifoliam et Swartzianam. (v. s.)

LVII. PECTIS Less. in *linnæâ* 1831. p. 708. *syn.* 153. — Lorentea Lag. nov. gen. (1816). p. 28. Cass. *dict.* 27. p. 202. nec Less. nec Ort. — Chthonia Cass. *bull. philom.* 1817. p. 33. — Pectidis sp. pleræque Linn.

Cor. disci obscurè bilabiata. Pappus disci et radii uniserialis, paleis basi scariosis latis, apice setiformibus serrulatis, sæpius inæqualibus. —

Herbæ annuæ aut rariùs perennantes. Folia glabra margine angustissimè cartilaginea. Capitula pedunculata, ramis nempe floridis nudis aut bracteolatis 1-cephalis elongatis, aut subsessilibus inter folia plùs minùs occultata.

* *Capitulis pedicellatis, pedicellis nudis capitulo longioribus.*

1. *P. LINIFOLIA* (Linn. sp. 1250. amœn. 5. p. 107), caule suberecto trichotomo tetragono ad angulos puberulo, foliis linearibus mucronatis margine scabris grossè glandulosis basi cilia setosa gerentibus, capitulis pedicellatis 4-18-floris, invol. squamis 4-6 elliptico-lanceolatis conduplicatis, pappo latè paleaceo, setis disci 5 longis, radii 2 oppositis. — in Jamaica et Sanct. Thomas. (Less.). — Sloan. jam. hist. 1. t. 149. f. 3. ex Less. in linnæa 1831. p. 709, sed icon à descr. Less. in plurimis recedit nec malè Pectidis convenit.
2. *P. SWARTZIANA* (Less. in linnæa 1831. p. 711), erecta, ramis patulis glabris subtrigonis, foliis oblongo-linearibus acutis basi cilia 2 utrinquè gerentibus grossè pellucido-glandulosis, pedicellis ex dichotomiis et apice ramorum aphyllis elongatis 1-cephalis, invol. squamis 5-6 linearibus obtusiusculis subconvolutis, floribus 10, acheniis glabriusculis, pappo 1-seriali brevi dentato-eroso radii brevi inæquali interdum uniaristato. (1) in Sancti-Domingi aridis. Pectis patula Linn. fil. herb. *P. punctata* Swartz in herb. L'Hér. *P. ciliaris* Swartz obs. bot. 309. non Linn. *P. Swartziana*? *Chthonia leptocephala* Cass. dict. 27. p. 206? — Icon Plum. ed. Burm. t. 86. f. 1. ad *P. punctatam* à Linnæo et Lessingio relata huc meo sensu pertinet, sed pessima! Icon Plum. t. 151. f. 2. à Linnæo ad ciliarem suam relata toto cœlo differt, nec descr. amœn. acad. convenit. (v. s.)
3. *P. CAPILLARIS*, caule erecto diffusè ramoso glaberrimo, foliis linearibus basi setigero-ciliatis subtus glandulas grossas ad utramque nervi mediè latus gerentibus, pedicellis elongatis capillaribus, invol. squamis 5 oblongo-linearibus acutis margine scariosis, capitulis 12-15-floris, ligulis oblongis minimis, pappo 5-aristato. (2) in Mexici ditione Oaxacana prope las Bacas septemb. flor. Caules pedicellique purpurascens tenues. Herba 5-6 poll. (v. s. comm. à cl. Mairet.)
4. *P. TENELLA*, caule herbaceo erecto glabriusculo tetragono oppositè ramosissimo, foliis linearibus submucronatis marginibus subrevolutis minutè et parcè glandulosis cilia ad basin utrinquè sæpius solitaria gerentibus, pedicellis 1-cephalis parcè bracteolatis capitulo paulò longioribus, invol. squamis 8 linearibus obtusius convolutis, floribus 13-15, pappo radii 3-4, disci 5-6-setoso. (1) in Mexico prope Laredo ad Tamaulipas legit cl. Berlandier flor. mense julio. Herba 3 pollic. glabra. Radix exilis simplex. Fol. 9-12 lin. longa. Flores flavi. (v. s.)
5. *P. ELONGATA* (H. B. et Kunth nov. gen. 4. p. 262. t. 392), caule erecto subtereti glabro virgato, foliis elongato-linearibus acutis basi dentibus aristatis utrinquè 3-8-ciliatis utrinquè subscabris, capitulis subcorymbosis ad apices ramulorum pedicellatis, involucri squamis 5 lineari-oblongis acuminatis, pappi paleis in disco et radio 10-20 setiformibus scabris inæqualibus. (1) circa Papayan? (Kunth), ad Hondam (Willd.), ad Panama (h. Hænke!). Herba pedalis basi simplex. Folia 15-18 lin. longa, 1-1 1/2 lata. (v. s. in h. Hænke. à cl. Sternh. comm.)
6. *P. BONPLANDIANA* (H. B. et Kunth nov. gen. 4. p. 263), erecta oppositè aut dichotomè ramosa, foliis lanceolato-linearibus acuto-mucronatis grossè pellucido-glandulosis basi dentibus aristatis 1-2 utrinquè ciliatis, capitulis terminalibus et alaribus longè pedunculatis solitariis, involucri sub-10-phyllis, pappo disci 5-setoso, radii pauciseti, utroque inæquali setis 3 elongatis scabris 2 brevibus membranaceis denticulato-erosis. (1) in Mexico prope Gueretaro et Tampico de Tamaulipas. Less. in linnæa 1831. p. 712. *P. linearis* Willd. herb. ex Less. Achenia pilosiuscula. Pedunculi sesquipollicares. (v. s.)
7. *P. UNIARISTATA*, caule herbaceo erecto tereti glabro oppositè ramosissimo, foliis linearibus mucronato-aristatis cilia utrinquè 3-4 ad basin et pappo

pellucidas ad utrumque nervi medii latus 1-seriatis gerentibus, pedicellis elongatis bracteolatis 1-cephalis, invol. squamis 5 linearibus acuminatis, floribus circ. 10, pappo radii et disci coroniformi hinc in setam unicam aristatam producto. ① in Mexico. Herba pedalis elegans tota glabra. Cor. flavæ. Folia 6-10 lin. longa, vix ultra lineam lata. (v. s. comm. à cl. Alaman.)

8. *P. PYGMÆA* (H. B. et Kunth nov. gen. 4. p. 262), caule erecto basi ramoso, ramis teretibus puberulis fastigiatis, foliis lanceolato-linearibus acuminato-piliferis per totam longitudinem dentibus aristatis ciliatis, capitulis terminalibus subcorymbosis brevissimè pedicellatis inter folia suprema absconditis, capitulis 12-floris, invol. 5-phyllo tubuloso, pappi paleis 3-10 subscariosis inæqualibus scabris. ① in calidis prope Honda ad fl. Magdalensæ. Less. in linnæa 1831. p. 713. *P. minuta* Willd. herb. ex Less. non Lin. Proxima dicitur *P. serpyllifolia* et ancipitis.
9. *P. BERLANDIERI*, caule herbaceo erecto di-trichotomè ramoso, ramis angulatis secus angulos hinc inde subscabris, foliis linearibus mucronatis subglandulosis supernè et margine plus minùs scabridis basi cilia 2-4 utrinquè gerentibus, pedicellis 1-cephalis subbracteolatis involucri paulò longioribus, invol. squamis linearibus acuminatis convolutis, pappi setis 1-serialibus vix basi dilatatis disci 5-6, radii 2-3. ① in Mexico ad Tancoyuca legit cl. Berlandier. Radix exilis ramoso-fibrosa. Herba 5-6 pollicaris. Habitus ferè Lorentæ Hænkeanæ, sed pedicellis brevior, caulis ferè omninò glaber, involucri acuminata et pappus diversus. (v. s.)

** *Capitulis sessilibus aut subsessilibus.*

10. *P. ANCEPS* (Less. in linnæa 1831. p. 715), caule ascendente trichotomo ancipiti biseriali puberulo, foliis obovato-linearibus muticis subtùs glandulosis margine scabris basi setigeris, capitulis subsessilibus circ. 13-floris, ligulis brevissimis, invol. squamis 5 obtusissimis, pappi paleis latiusculis in disco 4-5, in radio 3 haud unilateralibus. ① Patr. non satis nota. An ins. Caribæ? Achænia hirsutiuscula.
11. *P. COSTATA* (Ser. et Merc. mss. in h. DC.), prostrata ramosissima, foliis oblongis acuto-mucronatis basi dentibus aristatis utrinquè 2-5-ciliatis, capitulis inter folia suprema congesta subsessilibus, involucri 5-angulato 5-phyllo, pappi paleis 3-5 membranaceis erectis basi latius apice aristatis subserratis inæqualibus. ① in insulâ Cubâ legit cl. Ramon de la Sagra. Lorentea prostrata Lag. nov. gen. et sp. 28. n. 359. (v. s.)
12. *P. PROSTRATA* (Cav. ic. 4. p. 12. t. 324), prostrata ramosissima, foliis oblongis acuto-mucronatis basi dentibus aristatis 5-7 utrinquè ciliatis, capitulis ad ramorum dichotomias et inter folia suprema congesta sessilibus, involucri 5-phyllo ovato-cylindraceo, pappi paleis membranaceis erectis basi latius apice subaristatis in disco 5, in radio 2 unilateralibus. ① in Mexico, in montibus Sancti Angeli (Moq. ic. ined.), ad occid. urbis Guanaxuato (Alam.), circa Leon et alibi. Less. in linn. 1831. p. 714. Chthonia prostrata Cass. dict. 9. p. 173. *P. fasciculata* Poir. dict. 5. p. 120. Achænium hirsutiusculum. (v. s.)
13. *P. CILIARIS* (Lin. sp. 1050. amen. acad. 5. p. 497), erecta dichotomè ramosa, caule tetragono sub foliis bifariam puberulo, foliis linearibus amplexicaulibus acutis basi ciliis utrinquè 3-5-ciliatis, capitulis intrâ folia superiora congesta subsessilibus, invol. squamis 5, pappi paleis membranaceis lanceolatis acuminatis serratis radii 3 inæqualibus, disci 5. ① in Santo-Domingo. *P. ciliaris* Swartz! in herb. L'Hér. (sed descr. suæ non congr.) Less. in linnæa 1831. p. 708. An fortè Chthonia glaucescens Cass. dict. 9. p. 173? (v. s.)
14. *P. FASCICULIFLORA*, erecta glabra, caule tereti subsimplici, foliis lato-linearibus subobtusis, infer. oppositis eciliatis, summis alternis ad ramos floridos congestis pauci-ciliatis, omnibus crebrè et minutè pellucido-punctatis, capitulis ad axillas et apicem inter folia floralia congestis, invol. oblongis squamis 5 acutiusculis, ligulis parvis oblongis, achæniis 1-2-aristatis. ① in Mexico legit cl. Hænke. Affinis *P. carthusianorum* sed duplò major. Capitula ferè ut in Campanulâ glomeratâ disposita. (v. s. in h. Hænk. à cl. de Sternberg misso.)

15. *P. CARTHUSIANORUM* (Less. in *linnæâ* 1831. p. 712), caule erecto tereti, sub foliis obscure biserialium pubescente, foliis lato-linearibus acutiusculis ciliis 5-9 utrinquè ad medium usque instructis minutè et crebrè pellucido-glandulosis, ramis floridis brevissimis apice corymboso-congestis, capitulis inter folia occultatis sessilibus, invol. squamis 6 oblongis obtusis conduplicatis, flor. cir. 12, acheniis hirsutiusculis, pappi paleis latis in setam acuminatis, radii sæpius 3, disci 5. (1) in Santo-Domingo legit cl. Bertero, qui fortè non immeritò sub nomine *P. ciliaris* comm. Huc enim apprimè convenit icon *Plum.* ed. Burm. t. 151. f. 2. à Linnæo et aliis ad ciliarem relata. (v. s.)
16. *P. SERPYLLIFOLIA* (Less. in *linnæâ* 1831. p. 715), caule suffruticuloso ramossissimo procumbente basi repente, foliis oblongis obovatisve obtusis cilia utrinquè 3-4 ad basin conferta gerentibus glandulosis internodio longioribus, capitulis paucis ad apices ramorum sessilibus solitariis, invol. squamis 5 obtusis, floribus 15-30, pappi paleis basi dilatatis in radio 2-3, in disco 5-6, achenio hirsuto. 2/3 in insulis Caribæis (Sw.), Portorico (Wydł.), Sanct.-Thomas. (Less.), Chthonia repens Cass. dict. 27. p. 205. *P. humifusa* Wydł. pl. exs. n. 208. Icon *Plum.* ed. Burm. t. 95. f. 2. ad Lorenteam *serpyllifoliam* ab auct. relata huc fortè ex capitulis sessilibus potiùs pertinet. Valdè accedit ad *P. humifusam* Sw. (Lorenteam Less.) sed differt radice perenni, caule basi radicante capitulis sessilibus et pappo disci uniseriali. (v. s.)
17. *P. SIBIRICI* (Less. in *linnæâ* 1831. p. 717), caule adscendente basi repente tetragono subdichotomo, foliis oblongo-obovatis muticis internodio sæpius brevioribus basi cilia 4-6 gerentibus, capitulis inter folia subsessilibus sparsis, invol. squamis obtusis, pappo latè paleaceo disci paleis circ. 8, radic. paucioribus. 2/3 in Martinicâ (Sieb.) et Guadalupâ (Perr.). Pectis prostrata Sieb. fl. Mart. exs. suppl. n. 24. Pappi radii palear. plùs quàm 2. Cor. luteæ. Affinis *P. serpyllifoliæ*. (v. s.)

† *Species exclusæ.*

- P. ANGUSTIFOLIA* Torr. = *Pectidopsis angustifolia*.
P. CANESCENS Kunth. = *Lorentea canescens*.
P. CANESCENS Less. = *Lorentea auricularia*.
P. DISCOIDEA Horn. = *Leyssera capillacea*.
P. HUMIFUSA Sw. = *Lorentea humifusa*.
P. MEIOPOLIA Wall. = *Glossocardia Boswellæ*.
P. MINUTA Linn. = *Bellium minutum*.
P. PINNATA Lam. = *Schkuhria abrotanoides*.

LVIII. *LORENTEA* Less. in *linnæâ* 1831. p. 717. syn. 153. non Ort. (1) nec Lag. (2). — *Cryptopetalum* Cass. dict. 27. p. 206. non Hook. — *Pectidis* sp. Linn.

Corollæ disci obscure bilabiatae. Pappus disci biserialis inæqualis serrulatus. — Herbæ interdum basi suffruticulosæ. Capitula pedicellata in omnibus, unicâ (*L. sessiliflora*) exceptâ.

Sect. I. *CRYPTOPETALUM* Cass. dict. 12. p. 123. 27. p. 206.
 non Hook.

Pappus radii paleis plurimis acuminato-setiformibus pappo disci brevioribus constans.

1. *L. CRYPTOPETALA*, caule diffuso bifariam piloso, foliis lineari-lanceolatis,

(1) *LORENTEA* Ort. = *Sanvitalia*.

(2) *LORENTEA* Lag. et Cass. dict. 27. p. 202. = *Pectidis* sp. Less. seu *Chthoniæ* Cass.

caulis punctatis dentibus aristatis utrinque 7-8-ciliatis, capitulis terminalibus solitariis pedunculatis, invol. cylindraceo 5-phylo, ligulis brevissimis, pappi paleis inaequalibus pluriseriatis rigidis scabris. ① in Peruvia ad Guamanga legit Jos. de Jussieu. Cryptopetalum ciliare Cass. l. c. (v. s. in h. Juss.)

2. *L. HÆNKEANA*, caule herbaceo erecto tereti plus minus hirsuto di-trichotomè ramoso, foliis linearibus aut sublanceolato-linearibus mucronatis plus minus utrinque hirsuto-scabris cilia utrinque 2-3 basi gerentibus, pedicellis 1-cephalis bracteolatis capitulo quintuplò longioribus, invol. squamis 5 oblongo-linearibus subobtusis achænia vix superantibus, pappo disci biseriali setis 5 elongatis cæteris brevioribus tenuioribus, radii paucioribus brevioribus. ② in Mexico legit cl. Hænke.

a. *hirsuta*, ramis folisque hirsuto-scabris, ciliis foliorum breviusculis. (v. s. in h. Hænke à cl. Sternberg comm.)

β. *leviuscula*, ramis folisque glabrisculis, setis foliorum longioribus. — Etiam in Mexico legit cl. Hænke. Habitus omnino *Pectidis Berlandieri*, sed caulis basi durior et verisim. suffruticulosus, pappus et involucria valde diversa etc. (v. s. cum var. α.)

3. *L. HUMIFUSA* (Less. in linnæâ 1831. p. 719), prostrata ramosissima, foliis obovato-oblongis obtusis sparsè et minutè glandulosis basi dentibus aristatis utrinque 2-4-ciliatis, capitulis ex axillis foliorum superiorum confertorum brevè pedicellatis, involucriis ovatis 5-phyllis, pappo disci biseriali, paleis majoribus 5 membranaceis basi latis apice angustè acuminatis serratis disci brevioribus. ① in Antillarum sterilibus apricis, Guadalupâ, Sancta-Cruce. *Pectis humifusa* Swartz fl. ind. occ. p. 1362. — Plum. ed. Burm. t. 95. f. 2? (v. s.)
4. *L. MULTIFLOSCULOSA*, caule suffruticuloso diffuso tereti glabro, foliis lato-linearibus seu lingulatis obtusis mucronatis basi cilia utrinque 3-4 et in paginâ inf. glandulas plurimas parvas gerentibus, pedunculis elongatis bracteolatis 1-cephalis, invol. squamis 8 oblongis obtusis, floribus 30-60, achæniis glaberrimis, pappo disci biseriali aristato serrato inæquali, radii 1-seriali breviori. ② in Peruvia montibus Guanacentibus (sic script. in herb. an Guanacensisibus ex Leon de Guanuco?) legit cl. Hænke. Folia 9-12 lin. longa, 2 lin. lata. Invol. 4 lin. longa, floribus breviora. Ligulæ disco longiores. Pappus etiam disci achænio brevior. Variat ramis foliis pedunculisque plus minus elongatis. Specim. longifolia optimè referunt ic. Banks. rel. houst. t. 19. à Lessingio ad Lorenteam saturoioidem relata, sed in nostrâ achænia radii papposa nec calva. (v. s. in h. Hænke à cl. Sternberg comm.)
5. *L. SESSILIFLORA* (Less. in linnæâ 1831. p. 720), caule herbaceo, ramis ad apicem usque foliatis bifariam sub foliis puberulis, foliis linearibus mucronatis internodio brevioribus basi ciliigeris pellucido-glandulosis, capitulis inter folia absconditis, invol. 5-phylli squamis obtusis, pappo in radio et disco conformi — Patr. ign.

SECT. II. STAMMARIUM Willd. herb. — Lorentea Less. in linnæâ 1830. p. 135.

Pappus radii nullus aut dimidiatus auriculæformis brevis.

6. *L. AURICULARIS*, canescenti-hirtella, caulibus erectis aut procumbentibus ramosis, foliis linearibus acutis basi ciliis utrinque 2-3 instructis, capitulis pedicellatis circ. 40-floris, pappo radii auriculæformi. ① in temperatis prope Chilpancingo Mexicanorum ex Less. *Pectis canescens* Less. in linnæâ 1830. p. 135. t. 2. f. 7-12. et 1831. p. 718. non Kunth. *Stammarium hyssopifolium* Willd. herb. ex Less. (v. s. in h. Hænke à cl. de Sternb. comm.)

β. *Acapulcensis*, caulibus diffusis, foliis latius linearibus, invol. disco dimidio breviori. ② circa Acapulco. An species propria? (v. s. in h. Hænke.)

7. *L. CANESCENS*, canescenti-subhirtella, caulibus basi suffruticulosus diffuso-

ramosis, foliis linearibus acutis basi ciliis 2-3 utrinquè instructis, capitulis pedicellatis 40-floris, pappo radii nullo aut brevi paleaceo setigero. \mathcal{L} \mathfrak{z} in Mexico ad Tehuentepec et Chiguitan. Pectis canescens H. B. et Kunth nov. gen. am. 4. p. 263. t. 393. non Less. Cinerariæ sp. nov. Alam. herb. Affinis *L. auriculari*, sed pappus radii nunc omnino nullus, nunc setis paucis inæqualibus constans, nunquam nec coroniformis nec auriculæformis. (v. a. comm. à Alaman et Mairët.)

3. *L. SATUREIODES* (Less. in LINNÆ 1830. p. 135. t. 2. f. 7-12. et 1831. p. 718), caulibus suffrutescentibus erectis, foliis oblongo-linearibus mucronatis glabris basi ciliis utrinquè circiter 8-10 instructis, capitulis multifloris longè pedicellatis, pappo radii nullo. \mathcal{L} \mathfrak{z} in collibus arenosis prope Vera-Cruz. Aster satereiæ foliis Banks in Houst. rel. t. 19. Inula satereioides Mill. dict. Inula Satereiæ Spreng. syst. 3. p. 520.

9. *L. TENUESIPOLIA*, glabra, caule basi suffrutescente erecto-subdiffuso ramosissimo, foliis linearibus acuminatis grosse glandulosis ciliis utrinquè 3-4 basè gerentibus, capitulis longè pedicellatis circ. 10-floris, pappo radii nullo. \mathfrak{z} \mathcal{L} in insulâ Albemarle à numero archip. Gallapagos sub æquatore non longè à littore Peruviano jacentis, legit cl. Macrae. Rami tennes teretes. Fol. 8-10 lin. longa, semilineam lata. Involucra oblongo-cylindracea. Achæniâ elongata, radii glabra, disci paberula. (v. a. comm. ab hon. Soc. hort. Londin.)

Trib. II. *EUPATORIACEÆ* Less. syn. 154. — Eupatoriæ, Adenostylæ et Tussilaginéæ Cass. tabl. XIX et XX.

Stylus fl. hermaphr. supernè cylindraceus, ramis longis teretibus seu clavatis exsertis extûs supernè papillois rariùs puberulis, fl. masc. integer aut breviter bifidus clavulatus. Stigmatis series angustæ parùm prominulæ antè medios styli ramos desinentes, rariùs apicem attingentes, rarissimè confluentes. Corolla staminigera regularis 5-dentata, dentibus sæpiùs erectis. Anthæræ ecaudatæ. Pollen globosum echinulatum. — Capitula pauci- aut multiflora nunquam 1-flora, homogama fl. tubulosis, rariùs heterogama fl. radii tubulosis ligulatisve. Flores (paucissimis exceptis) cyanici. Herbarum aut frutices. Folia sæpiùs opposita. — Tribus accedens hinc ad Vernoniaceas et ad Senecionideas oppositifolias.

CONSPECTUS TRIBUS.

Subtrib. I. *EUPATORIÆÆ*. Capitula homogama. (Gen. LIX-LXXXIX.)

Div. I. *ALOMIÆÆ*. Pappus nullus. (Gen. LIX-LXIV.)

Div. II. *AGERATEÆ*. Pappus paleaceus aut partim squamellatus. (Gen. LXV-LXXIII.)

Div. III. *ADENOSTYLÆÆ*. Pappus setosus scaber aut plumosus. (Gen. LXXIV-LXXXIX.)

Subtrib. II. *TUSSILAGINÆÆ*. Capitula heterogama aut dioica. (Gen. XC-XCVII.)

Div. I. *PETASITEÆ*. Flores feminei tubulosi. (Gen. XC-XCIII.)

Div. II. *EUTUSSILAGINÆÆ*. Flores feminei ligulati. (Gen. XCIV-XCVII.)

Subtrib. I. *EUPATORIÆ*. — Eupatoriæ et Adenostylæ Cass. —
Alomiæ Ageratæ, Eupatoriæ Less.

Capitula homogama. — Flores nunquam stricto sensu flavi.

Div. I. *ALOMIÆ* Less. syn. 154.

Pappus nullus. — Divisio nimis artificialis ex affinitate Alepidis cum Agerato, Astemmati cum Adenostemmate et Orsinia cum Eupatorio, imò Isocarphæ cum Carphophoro.

LIX. *ORSINIA* Bertol. in litt. 1834. non in ann. st. nat. — Eupatorii sp. Balb.

Capitulum 6-7-florum. Recept. nudum. Invol. ovati squamæ 6-7 ovales erectæ æquales. Cor. 3-5-fidæ, hirtellæ, lobis ciliatis. Styli rami exserti muriculati obtusi. Antheræ inclusæ. Achænium obcompressum apice villosum. Pappus nullus. — Suffrutex aut Herba suffruticans Brasiliensis. Caulis teres pubescenti-velutinus apice ramoso-corymbosus. Folia opposita et alterna breviter petiolata ovata rigida acutiuscula acutè serrata penninervia reticulata suprâ scabrida subtus secus nervos nervulosque densè hirta. Corymbi terminales, ramis hirsutis. Capitula subsessilia. Invol. squamæ pallidæ dorso puberæ latè ovales subconcavæ. — Cum primum genus cl. Orsini à cl. Bertoloni dicatum in Jasoniam ceciderit hoc novum consecravit cl. auctor et mecum caract. ex specim. authentico desumptos communicavit.

1. O. *EUPATORIA*. \tilde{J} in Brasiliæ prov. Minarum Generalium ad Mariannam legit cl. Vauthier. Eupatorium Armani Balb. pl. rar. h. taur. 1810. p. 27. t. 6. sine loci indic. Flores albi. Folia 2 poll. longa, 15 lin. lata, petiolo 3 lin. (v. s. comm. à cl. Vauthier sub n. 263.)

LX. *PIQUERIA* Cav. ic. 3. p. 19. t. 235. Willd. sp. p. 1748. Cass. dict. 41. p. 115. Less. syn. 154. — Flaveriæ sp. Juss.

Capitulum 4-5-florum. Invol. 4-5-phyllum 1-seriale oblongum. Recept. nudum planum parvum. Cor. tubus brevis sub dilatatus sæpius hirsutus. Antherarum appendix terminalis subnulla. Styli rami obtusi. Achænia juniora compressa adulta 5-angulata glabra stipiti brevi articulata. — Herbæ aut suffrutices Americanæ. Folia opposita sæpius trinervia. Capitula corymbosa aut paniculata parva. Corollæ albæ etiamsi ex siccis flavas dixerit Cassini, qui sæpiissime album videbat flavum.

Sect. I. *EUPIQUERIA*.

Capitula 4-flora. Cor. tubus hirsuto-barbatus. Invol. squamæ obtusæ mucronatæ. — Caules herbacei.

1. P. *TRINERVIA* (Cav. ic. 3. p. 19. t. 235), herbacea glabra, foliis ovato-ant oblongo-lanceolatis subserratis trinerviis, paniculâ corymbosâ laxâ polycephalâ. \mathcal{L} in Mexico locis rupestribus prope S. Rosa de la Sierra (H. B.), Jalapam (Schied. et Depp.), etc. in omnibus Mexici circuitibus ubi *Xoxonitztal* et *Yoloxitlic* nuncupatur (Moq. fl. mex.) Ageratum febrifugum Less. fl. mex. ined. Stevia febrifuga Moq. fl. mex. ic. ined. — Hern. mex. p. 353. fl. 1. ex Moq. et fortè p. 437. f. 2. ex DC. — P. trinervia Jacq. f. eclog. 1. p. 70. t. 48. Variat ramis omnino glabris aut sub foliorum origine bifariam pilosis. (v. v. c. et s. sp.)
2. P. *PILOSA* (H. B. et Kunth nov. gen. am. 4. p. 153), herbacea piloso-glandulosa, foliis latè ovatis aut ovato-oblongis trinerviis grossè serratis, paniculâ corymbosâ laxâ. \mathcal{L} in temperatis Mexici inter Capula et Pascuaro (H. B.). Folia 2 $\frac{1}{2}$ poll. longa, 1 $\frac{1}{2}$ ferè lata. (v. s. comm. à cl. Alaman.)

Sect. II. PIQUERIOIDES.

Capitula 5-flora. Invol. squamæ striatæ in appendicem subulatam desinentes.
— Caules fruticosi. Folia triloba.

3. *P. PUBESCENS* (Smith in Rees cycl. 27. n. 2), fruticosa, caule pube densâ brevi velutinâ, foliis trilobis acutis glabriusculis, paniculâ densè corymbosâ villosâ, capitulis 5-floris. ♀ in Peruvîâ ex herb. Linn. fil.
4. *P. QUINQUEFLORA* (Cass. dict. 41. p. 116), fruticosa, caule basî glabro apice rufo-tomentoso, foliis trilobis glabriusculis pellucido-glandulosis, lobis acutis dentatis, paniculâ densè corymbosâ subviscosâ, capitulis 5-floris. ♀ in Peruvîâ legit Dombey. Corollæ à Cassinio dicuntur flavæ, sed frequentissimè ex sicco flores albos dixit flavos. An certè à priore diversa?

Sect. III. ARTEMISIOIDES.

Capitula 3-4-flora. Cor. glabræ, tubo longiusculo basi suprâ ovarium subdilatato. Invol. squamæ acuminatæ aut obtusæ. — Caules fruticosi glabri apice summo pubescentes aut glutinosi. Folia glabra petiolata. — An genus proprium?

5. *P. FLORIBUNDA*, fruticosa glaberrima apice glutinosa, foliis oppositis petiolatis ovato sen rhombeo-lanceolatis grossè dentatis, paniculæ ramosissimæ ramis corymbiferis, capitulis 4-5-floris, invol. squamis acuminatis. ♀ in Peruvîâ montanis Oronocensibus (et fortè in Chili) legit cl. Hænke. Inv. 3 lin. longa. Folia ferè Sedi populifolii sed membranacea. Specimina Chilensia in herb. Hænkeano (fortè schedulæ transpositione) dicta Peruvianis simillima in ditissimis Chilensibus herbariis nunquàm vidi et mihi quoad patriam valdè dubia! (v. s. in herb. Hænke à cl. Sternb. comm.)
6. *P. ARTEMISIOIDES* (H. B. et Kunth nov. gen. am. 4. p. 153), glabriuscula, foliis caulinis oppositis petiolatis lanceolatis acuminatis basi cuneatis grossè dentatis summis lineari-lanceolatis integerrimis sparsis, invol. squamis obtusis. ♀ in Peruvîâ (Domb.), in frondosis Quitepsium (H. et B.), in vallibus Cordillieris (Hænke). Flaveria Peruviana Juss. gen. 187. Flaveria spicata Smith in Rees cycl. v. 14. n. 2. Herba sicca subflavicans. Rami et folia novella subpubescentia, cæteræ partes glabræ. (v. s.)
7. *P. CALIOIDES*, fruticosa glaberrima, foliis linearibus integerrimis subnerviis, paniculæ ramosissimæ ramis racemiferis, capitulis 3-floris, involucri squamis subobtusis. ♀ in Peruvîâ vallibus Cordillieris legit cl. Hænke. Folia 8-10 lin. longa, vix lineam lata. Invol. 2 lin. longa. (v. s. in h. Hænke à cl. Sternb. comm.)

LXI. *ALOMIA* H. B. et Kunth nov. gen. am. 4. p. 151 et 312. t. 354.
Less. syn. 154.

Capitulum multiflorum. Invol. campanulatum imbricatum, squamis angustis acutis. Recept. convexum nudum. Cor. tubo gracili glabro. Antheræ appendicibus ovato-oblongis obtusis terminatæ. Styli lobi elongati obtusi. Achænia 5?-angularia apice calva. — Herba Mexicana ramosissima glabriuscula albiflora. Piquerias (nec Ethulias) caracteribus et habitu omninò referens.

1. *A. AGERATOIDES* (H. B. et K. l. c.), in Mexicis calidis prope Mescala. Ethulia ageratoides Spreng. syst. 3. p. 458.

LXII. PHALACRÆA.

Capitulum multiflorum. Invol. squamæ subbiseriales æquales 3-nerviæ. Recept. nudum. Corol. tubus basi piloso-hispidus, faux ampliata, limbus 5-fidus. Antheræ vix apice appendiculatæ. Styli rami exserti

apice clavati. Achænium compresso-angulatum calvum. — Herba Peruviana breviter puberula fortè subviscida. Folia opposita petiolata latè ovata basi truncata et ad petiolum subcuneata grossè et interdum duplicato-dentata. Pedunculi nudiusculi apice corymbosi. Capitula pedicellata. Flores purpurascens. — Genus ex pappo nullo affine Alomiæ sed diversum cor. tubo hispido, invol. non imbricato, etc. Aff. etiam Astemmati, sed differt cor. fauce ampliata nec cylindraca; tubo hispido nec glabro. — Nomen ex *φάλακρος*, calvus.

1. *P. LATIFOLIA*. \mathcal{L} in Peruvia circa Limam legit cl. Née. *Stevia* seu *Ageratum* Née! herb. *Ageratum latifolium* Cav. ic. 4. p. 33. t. 35? Planta mea (etiam in herb. Hænkano obvia) ut Cavanillesiana à Neesano itinere proveniens apprimè, ex habitu toto, iconi et desc. respondet, sed Cav. vidit pappum 5-paleaceum et in meâ pappus nullas. An diversa? An à Cavan. malè descr.? Ex Cav. vulgò *Teatina* nuncupatur. Petioli 6 lin. longi pedunculique caulesque pube brevi velutini. Limbus 2 poll. ferè longus sesquipollic. latus. (v. s.)
 β . *glabra*, caule glabro, pedunculis vix apice puberulis, petioli 15 lin. longis, fol. limbo 3 poll. longo, 2 poll. lato, capitulis paulò majoribus.
 \mathcal{L} Etiam habeo ex itin. Neesano sed locum propr. ignoro. (v. s.)

LXIII. GYMNOCORONIS.

Capitulum multiflorum homogamum. Invol. squamæ 1-seriales acuminatæ subnerviæ. Recept. nudum. Corolla tubulosa glabra, fauce non dilatata, lobis brevibus conniventibus. Styli rami exserti clavati. Achænium compressum dorso nervoso-subangulatum calvum. — Herbæ Brasilienses erectæ glabræ. Caules infernè sulcati. Folia opposita petiolata oblongo-lanceolata acuminata dentata. Pedunculi corymbosi. Capitula pedicellata subpuberula. Flores albi. — Genus valdè Adenostemmati affine sed achænia calva! Nomen ex *γυμνος* nudus, et *κορυς*, apex, terminus. An stirpes sequentes sint species aut varietates dubius hæreo.

1. *G. ATTENUATA*, foliis oblongato-lanceolatis basi cuneatis secus petiolum decurrentibus. \mathcal{L} in Brasiliæ prov. Rio-Grande. (v. s. in h. Mus. reg. Par. à Mus. imp. Bræ. sub n. 782 miss.)
2. *G. SUBCORDATA*, foliis oblongato-lanceolatis basi truncatis subcordatis in petiolum non decurrentibus. \mathcal{L} in Bras. prov. Rio-Grande. (v. s. cum priori sub n. 886 miss.)

LXIV. ISOCARPHA *R. Brown. obs. comp. 110 excl. syn. Less. in linnæâ 1830. p. 141. 1831. p. 405. syn. p. 154.*

Capitulum multiflorum. Invol. squamis circ. biseriatis conduplicatis. Recept. conicum, paleis acuminatis squamas referentibus. Corolla tubulosa 5-dentata. Styli rami elongati extus convexi hispiduli acuti. Achænium prismaticum epapposum. — Herbæ Americæ æquinoctialis erectæ oppositifoliæ. Folia integra rarius dentata. Capitula terminalia.

1. *I. ATRIPLICIFOLIA* (Brown. l. c.), erecta glabra, caule tereti, foliis oppositis auriculato-connatis aut alternis stipulato-auriculatis in petiolum alatum basi attenuatis supernè deltoideo-ovatis acuminatis grossè et obtusè dentatis triplicinerviis membranaceis, capitulis paucis pedicellatis subpaniculatis. — in Amer. merid. (Mill.) ad flumen Magdalenzæ (Bertero!), *Calidemos atriplicifolia* Spreng.! syst. 3. p. 457. *Bidens atriplicifolia* Linn. sp. 2. p. 1168. *Spilanthes atriplicifolia* Linn. syst. veg. 610. Ad aliam stirpem alludere videtur Necker cum recept. nudum dixit, achænia angulata nuda, juniora interdum setas 1-2-gerentia. (v. s.)

2. *I. OPPOSITIFOLIA* (R. Brown. l. c.), caule dichotomo, ramis longissimis erectis tripartitis simplicibusque, foliis lanceolatis, capitulis subternis sessilibus. — in Jamaica montibus. *Calea oppositifolia* Linn. sp. 1179. amœn. acad. 5. p. 404. Swartz obs. 302. — *Santolina* Brown. jam. 315. Gærtner fr. 2. p. 391. Invol. squamæ binerviæ laxæ muticæ rectilineo-acuminatæ ex Less.
3. *I. BILLEBRØGIANA* (Less. in linnæâ 1831. p. 405. in adn.), ramis flexuosis, foliis sessilibus penninerviis basi auriculatis spathulato-rhomboidis, squamis invol. penninerviis, achæniis pubescentibus. — in Tierra Bomba prope Carthagenam.
4. *I. ECHIOIDES* (Less. in linnæâ 1830. p. 141. t. 2. f. 14-16), caule infernè dense foliato supernè nudo, foliis sessilibus lineari-lanceolatis triplinerviis scabris et hispidis, paniculâ corymbosâ confertâ, corollis apice pilosisculis, invol. squamis linearibus canaliculatis ciliato-hirsutis, achæniis glaberrimis. ♀? in graminosis Mexici prope Hacienda de la Laguna. — Paniculæ rami foliis suis breviores. Invol. floribus æquale, squamis ciliatis, inter. latius scariosis.

† *Species exclusa.*

I. BEATERII Balb. = *Critonia Dalea*.

Div. II. AGERATEÆ Less. syn. 155.

Pappus paleaceus, squamellis membranaceis rigidisve distinctis aut in coronam plus minus concretis.

LXV. *CORLESTINA* Cass. dict. 6. suppl. p. 8. et v. 26. p. 227. Kunth nov. gen. am. 4. p. 150. Less. syn. 155. — *Agerati* sp. auct.

Capitulum multiflorum. Invol. cylindraceo-hemisphæricum polyphyllum subimbricatum. Recept. convexum paleaceum aut nudum. Achænia pentagona. Pappus coroniformis membranaceus dentatus, paleis nempe concretis apice subinæqualibus. — Herbæ annuæ erectæ ramosæ. Caules teretes. Folia opposita breviter petiolata dentata. Capitula corymbosa approximata pedicellata. Corollæ cæruleæ aut purpureæ. — Genus vix ab *Agerato* distinctum et cum eo fortè (eodem jure ac sectiones *Stevie*) coadunandum?

Sect. I. ISOCARPHOIDES.

Recept. paleis linearibus onustum.

1. *C. ISOCARPHOIDES*, caule suffruticoso tereti hispidulo scabro, foliis lineari-lanceolatis acuminatis subsessilibus triplinerviis integerrimis suprâ scabris subtus velutino-villois, invol. squamis lineari-subulatis scabridis, capitulis circ. 25-floris, pappo brevi truncato. ♀ in Mexico legit cl. Hænke. Proximè accedit ad *Isocarpham* echioideam ex descr. Less., sed habitus ab *Hieracio* echioidei diversus. Caulis per ferè totam longitudinem foliosus. Pappus brevis equidem non verò nullus. (v. s. in h. Hænke. à cl. Sternb. comm.)

♀. *dentata*, foliis majoribus hinc inde dentatis. (v. s. cum var. α.)

2. *C. PALMACEA* (Gray! in Andr. pl. mex. exs. n. 287), caule herbaceo erecto tereti molliter ac brevissimè velutino, foliis petiolulatis ovatis seu ovato-lanceolatis tripli- aut quintuplinerviis subdentatis suprâ scabridis subtus velutino-villois, invol. squamis lineari-subulatis scabridis, capitulis 25-floris, pappo coroniformi dentato hinc inde in setam expanso. ♀? in Mexico circa Oaxacam legit cl. Andrieux mense julio. Flores violacei. (v. s.)
3. *C. ALBIDA*, caule herbaceo erecto puberulo, foliis ovalibus utrinquè attenuatis vix dentatis quintuplinerviis suprâ hispido-scabris subtus tomentoso-hirsutis, invol. squamis glabresculis linearibus acutis, capitulis circ. 60-floris, pappo coroniformi dentato. ♀? in Mexico inter Oaxacam et Mitla julio ineunte legit cl. G. Andrieux. Cor. albida. (v. s. comm. à cl. inv. snb n. 548.)

Sect. II. AGERATOIDES.

Receptaculum epaleaceum.

4. *C. AGERATOIDES* (H. B. et Kunth nov. gen. am. 4. p. 151), caule pulverulento scabrido, foliis ovato-oblongis acuminatis utrinque scabris, corollæ tubo extus pilifero, pappo cupulæformi obtusè dentato. ① in Mexici montanis ut videtur sat. frequens. *Ageratum cœlestinum* Sims bot. Mag. t. 1730. *Sparganophorus ageratoides* Lag. elench. h. madr. 1815. p. 25. ex Spreng. *Ageratum corymbosum* Balb. 1 hort. taur. 1806 et 1808. Pers. ench. n. 4. (v. s. sp.)

β. *latifolia*, foliis latè ovatis. — in Mexico. (v. s.)

5. *C. CORYMBOSA*, caule glabriusculo, foliis ovatis acuminatis utrinque subcabridis, corollæ tubo extus glandulifero, pappo cupulæformi inæqualiter dentato, dentibus 2 longioribus acutis. — in Mexico? *Ageratum corymbosum* Zucc. obs. n. 85. *Cœlestina cærulea* Cass. dict. 6. suppl. p. 8. An prioris varietas? (v. v. c. in hort. Florent. à Zuccagnii tempore seminibus servatam.)

6. *C. PARVIFOLIA*, caule vix puberulo, foliis ovatis crenato-dentatis brevissimè petiolatis, pedunculis racemoso-corymbosis, corollæ tubo glabro, pappo coriiformi obtusè dentato. — in Amer. calid. (fortè circa Limam) legit cl. Née. Petioli lineam vix longi. Limbus folii 6 lin. longus, 4. lin. latus. (v. s. ex itin. Née.)

LXVI. *AGERATUM* Linn. gen. n. 936. Gærtn. fr. 2. p. 398. t. 165. — *Carelia Pont. diss. Adans. fam. 123. non Less.*

Capitulum multiflorum subglobosum. Invol. polyphyllum imbricatum, squamis linearibus acuminatis. Recept. nudum. Cor. 5-fidæ. Styli rami elongati cylindracei obtusiusculi. Achænium subpentagonum basi attenuatum. Pappus paleaceus, paleis 5-10 liberis aristato-acuminatis aut pectinatis obtusis. — Herbæ pleræque annuæ erectæ Americanæ aut rarius Indicæ. Folia opposita petiolata ovata dentata. Capitula corymbosa. Flores cærulei aut albi. — Sectiones non magis inter se differre videntur quàm sectio *Steviæ*.

Sect. I. EUAGERATUM. — *Ageratum* Cass. dict. Less. syn. 155.

Pappi paleæ 5 ad basin distinctæ subserratæ apice acuminatæ aristatæ.

1. *A. CONYZOIDES* (Linn. sp. 1175), caule ramoso, foliis ovatis rhombeis cordatisve longiusculè petiolatis, pappi paleis basi dilatatis serrulatis apice longè aristatis corollam subæquantibus. ① frequens in Americâ et Africâ æquinoctiali, in Indiâ orientali, insulis Mauritanis (introd.?) et Maderæ secus vias (Low!), in ins. Sandwich Owyhee (Macrae!). — Schkuhr handb. t. 238. Hook. exot. flor. t. 15. Herba valde varians, caule nunc hispidò nunc ferè glabro, foliis nunc ovatis nunc obcordatis, acutis aut obtusis, plus minùs dentatis pilosis, petiolis hispidissimis aut glabriusculis, corollis cæruleis (*Ag. cæruleum* Desf. cat. h. par.) aut albis (*A. album* Stend. nom. ph. p. 13), undè plurimæ species ab auctoribus admixtæ hic ut varietates indicatæ, vix inter se nunc satis distinctæ sed tamen recognoscendæ.

a. *obtusifolium*, caule glabriusculo, foliis ovatis obtusis, floribus pallidè cæruleis. *A. obtusifolium* Lam. dict. 1. p. 54. ill. t. 672 f. 1. — *Sparganophorus obtusifolius* Lag. elench. h. madr. 1815. p. 25? (v. s.)

β. *hirtum*, caule hirtò, foliis ovatis acuminatis. *A. hirtum* Lam. l. c. Poir. suppl. 1. p. 241. — Lam. ill. t. 672. f. 2. (v. v. c.)

γ. *Mexicanum*, caule hirtello, foliis ovatis subobtusis, floribus cæruleis. *A. Mexicanum*, bot. mag. t. 2524. Sweet br. fl. gard. t. 89. (v. s.)

δ. *cordifolium*, caule hispidò, petiolis hispidissimis, foliis ovatis acuminatis. *A. cordifolium* Roxb. hort. calc. Wall. herb. et list, n. 237. (v. s.)

Sect. II. PECTINELLUM.

Pappi paleæ circ. 10 distinctæ per totam ferè longitudinem pectinato-ciliatæ.

2. *A. MELISSAFOLIUM*, caule ramoso glabro, foliis petiolatis glabriusculis ovatis crenato-dentatis truncatis aut subcordatis apice subacutis, pedunculis corymbosis, corollâ glabrâ brevi, pappi paleis oblongis pectinato-ciliatis subacutis. — in Brasiliâ circa Bahiam leg. cl. Salzmänn (n. 431) et Blanchet (n. 136 et 1974!). An *Chrysocoma pauciflora* Arrab. fl. flum. v. 8, t. 3? Petioli 6-7 lin. longi; limbi fol. pollicem longi, 6 lin. lati. Achænia angulata basi attenuata glabra. (v. s.)
3. *A. MATRICARIOIDES* (Less. syn. 155), caule ramoso, foliis petiolatis rhombeo-ovatis obtusis crenatis glabriusculis, pedunculis corymbosis, pappi paleis lanceolatis inciso-laceris pubescentibus. — in insulâ Cubâ. *Hymenopappus matricarioides* Spreng. syst. 3. p. 450. *Anthodium patena* (Spr.) Corolla Piqueriæ (Less.). Cæt. ign.
4. *A. DOMINGENSE* (Spreng. syst. 3. p. 446), caule gracillimo vix puberulo, foliis petiolatis orbiculatis cordatis puberulis crenato-dentatis, pedunculis 1-cephalis solitariis, pappi paleis oblongis serratis breviter acuminatis corollâ dimidio brevioribus. ① in Santo-Domingo legit cl. Bertero! Herbula fortè decumbens. Petioli 4 lin. longi. Folia 4 lin. diam. Invol. squamæ paucæ. (v. s.)

† *Species dubiæ.*

5. *A. CILIARÆ* (Linn. sp. 1175), caule glabro, foliis petiolatis ovatis crenatis obtusis. — in Ind. orient. prope Bisnagas. Pluk. alm. t. 81. f. 4. Spec. omnino obscura †.
6. *A. ? MARITIMUM* (H. B. et Kunth nov. gen. am. 4. p. 150), caule herbaceo ramoso glabriusculo, foliis petiolatis ovatis leviter cordatis obtusiusculis crenatis glabris, capitulis corymbosis pedicellatis, circ. 70-floris, corollâ glabrâ, pappi paleis crebris subulatis. ① in littore insulæ Cubæ prope Havanam. Corollæ violacæ. An *Eupatorii* species? An genus proprium?

† *Species exclusæ.*

- A. ANGUSTIFOLIUM* Sp. = *Eriophyllum angustifolium*.
A. ARBUTIFOLIUM Kunth. = *Phania arbutifolia*.
A. CŒLESTINUM Sims. = *Cœlestina ageratoides*.
A. CORYMBOSUM Balb. = *Cœlestina ageratoides*.
A. CORYMBOSUM Zucc. = *Cœlestina corymbosa*.
A. FRERIFUGUM fl. mex. = *Piqueria trinervia*.
A. GUIANENSE Aubl. = *Habeclinium macrophyllum*.
A. LATIFOLIUM Cav. = *Phalacræa latifolia*.
A. LINEARE Cav. = *Palafoxia linearis*.
A. PEDATUM Ort. = *Stevia pedata*.
A. PUNCTATUM Jacq. = *Stevia serrata*.
A. PUNCTATUM Ort. = *Stevia Eupatoria*.
A. VISCOSUM Cav. = *Stevia salicifolia*.

LXVII. ANISOCHÆTA.

Capitulum multiflorum (fortè dioicum?). Invol. ovatum imbricatum, squamis lineari-lanceolatis adpressis disco brevioribus. Recept. nudum. Cor. tubulosæ profundè 5-fidæ glabræ exter. ad peripheriam dejectæ. Anthæræ ecaudatæ. Styl. in plerisque inclusus subabortivus, in paucis ramis exsertis cylindraceis subclavatis vix papillosis. Achænia teretiuscula vix subpuberula erostria (immutura aut abortiva?)

Pappus paleis sæpius 3 rarius 4-5 erectis squamellosis angustis acuminatissimis inaequalibus. — Frutex Australi-Africanus subscondens. Caulis teres striatus pubescens, ramis patulis. Folia alterna petiolata latè ovata basi subtruncata 3-5-nervia grossè inciso-dentata glabra, petiolo nervisque pubescentibus. Paniculæ ramos terminantes subundæ laxæ. Bracteolæ minimæ secus pedicellos. Cor. ex sicco albidæ.

1. A. MINAHOIDES. ♂ in Africâ austr.-orient. ad Zw. Omsamcaba et Omsamwabo legit cl. Drege! Petiolus semipollic. Limbus folii 2 poll. longus et latus. (v. s. comm. à cl. inv.)

LXVIII. ADENOSTEMMA Forst. nov. gen. (1776) p. et t. 45. Cass. dict. 25. p. 360. Less. syn. 156. — Lavenia Swartz Prod. (1788) 112. fl. ind. occ. 1329. — Verbesinæ et Cotulæ sp. Linn.

Capitulum multiflorum homogamum. Recept. planum nudum foveolatum. Invol. campanulatum floribus ferè brevius post anthesin reflexum, squamis 1-seriatis foliaceis oblongis. Corolla tubulosa 5-dentata infrà lobos subvillosa, fauce vix ampliata. Stigmata longè exserta apice dilatata colorata. Achænium obovato-oblongum subangulatum superatum, aristis 3-5 brevibus rigidis apice glandulâ globosâ aut clavatâ terminatis. — Herbæ æquinoctiales glanduloso-pilosæ seu glabriusculæ. Folia opposita petiola ovata seu rhomboidea triplinervia dentata. Capitula pedunculata corymboso-paniculata. Flores albi. — Nomen Swartzii à Rajo mutuatum erat ritè conditum (planta enim erat Verbesina Lavenia Linn.), sed nomen Forteri vetustius et characterem indicat. — Species inter se valdè affines examen ulterius requirunt.

* *Caule basi radicante adscendente.*

1. A. SWARTZII (Cass. dict. 25. p. 362), caule basi radicante adscendente simplici apice minutissimè puberulo, foliis petiolatis ovato-subcordatis basi cuneatis grossè dentato-crenatis subacutis glabris, pedunculis 3-5 corymbosis elongatis, invol. squamis obtusis dorso glabris apice ciliolatis, achæniis distinctè stipitatis scabridis. (1) in subhumidis sylvaticis umbrosisque Jamaicæ. Lavenia decumbens Swartz! fl. ind. occ. 1331. Cotula Verbesina Linn. au. ac. 5. p. 407. — Sloan. jam. 126. p. 262. t. 155. f. 2. Caulis non decumbens ut vult Swartz sed adscendens. Cor. 15-20 albidæ. Achæni puncta parva albidâ pulveris moleculam referunt. Pappi setæ 3. (v. s.)
2. A. BERTERII, caule basi adscendente glanduloso-puberulo, foliis petiolatis ovato rhombeis basi cuneatis apice acutis hinc inde dentato-serratis glabris, corymbo 5-7-cephalo, involucri squamis ovalibus dorso puberulis subacutis, achæniis distinctè stipitatis muricatis. (1) in Santo-Domingo legit amic. Bertero. Lavenia decumbens ex Hispaniolâ Spreng. syst. 3. p. 445. A. Swartzii simillima, sed differt notis indicatis, consist. foliorum magis papyraceâ, capitulis duplò ferè majoribus, achæniis multò magis muricatis, setis pappi ferè ubiq̃ue 2 nec 3. (v. s. comm. à cl. Balbis.)
3. A. PERROTTETII, caule basi radicante adscendente glanduloso-subpuberulo, foliis petiolatis ovatis crenatis obtusis basi cuneatis glabris, paniculâ polycéphalâ laxâ, invol. squamis oblongis acutiusculis dorso puberulis, achæniis distinctè stipitatis muricatis. (1) ? in paludibus turfosis peninsulæ Promontorii Viridis Africanæ legit cl. Perrottet. Similis A. Swartzii, sed folia rigidiora obtusiora. Capitula 11-15. Achænia punctis glandulosis rubris muricata. Caules rubescentes. (v. s. comm. à cl. inv.)
4. A. MAURITIANUM, caule basi radicante adscendente simplici vix apice puberulo, foliis petiolatis ovato-lanceolatis basi cuneatis grossè dentatis glabris, pedunculis elongatis 3-7 in corymbum dispositis, invol. squamis obtusis

- glabris, achæniis lævibus. — in insulæ Mauritiæ sylvis montosis (Comm., Sieb. Bontou! etc.). Ad. viscosa var. Commersonii Cass. dict. 25. p. 362. *Lavenia erecta* Sieber! fl. maur. exs. 2. n. 119 et 364. Non malè refert t. 42. Burm. Zeyl. Folia papyracea cum petiolo 3-5 pollic. longa. (v. s.)
5. *A. DREGEI*, caule basi radicante adscendenti-erecto glabro vix hinc inde asperulo, foliis petiolatis ovatis basi longè cuneato-attenuatis apice acutis dentatis membranaceis suprà pube minimâ sparsâ puberulis subtus secus nervos glanduloso-scabridis, corymbo laxo, pedicellis elongatis, invol. squamis elliptico-oblongis pubescentibus, achæniis valdè muricatis. — in Africâ australi ad Zw. Omsamwubo et Omsamculo legit cl. Drege! Habitus *A. Mauritiani*, sed achæniis muricatis et aliis notis faciliè differt. (v. s.)
6. *A. MICROCEPHALUM*, caule basi radicante adscendente minutissimè glanduloso-pubero, foliis breviter petiolatis ovatis denticulatis obtusis glabris, paniculæ laxiusculæ capitulis parvis, invol. squamis oblongis subacutis glabrisculis, achæniis vix muricatis. — in Indiâ orient. ad Gualpara et Silhet legit cl. Hamilton. *Lavenia viscida* Ham. ex Wall.! cat. n. 332. Affine *A. Perrottetii*, sed capitula dimidio minoribus sanè primâ fronte diversum. (v. s. comm. ab ampl. Cur. merc. angl. Ind. or.)
7. *A. TINCTORIUM* (Cass. dict. 25. p. 364), caule subprocumbente diffuso, radice repente, foliis oppositis lanceolatis inæqualiter serratis glabris nerviis, invol. squamis obtusis. — in Cochinchinâ et Chinâ ad colorem cœruleum foliis contusis eliciendum culta et *cham-lonla* nuncupata. *Spilanthus tinctorius* Lour. coch. ed. Willd. 2. p. 590. Species ex pappi ariatis 3-capitatis ad *Adenostemma* ritè videtur relata, sed fruct. non descr.
- ** *Caule erecto, achæniis muricatis.*
8. *A. GLUTINOSUM*, caule erecto ramoso villosoglutinoso, foliis ovatis acutis remotè dentato-serratis ciliatis basi cuneatis utrinquè glutinoso-hirtellis, corymbo paniculato laxo, pedicellis capitulo triplo longioribus, achæniis papuloso-muricatis. — in insulis Sandwich ad alt. 450-500. hexap. *Lavenia glutinosa* Gaudich.! bot. voy. Freyc. p. 471. Ach. basi attenuata sessilia glabra, sed evidentissimè tuberculis minimis muricata. (v. s. comm. à cl. invent.)
9. *A. VISCOSUM* (Forst.! nov. gen. n. 15), caule erecto pubescente aut glabro, foliis petiolatis oblongo-lanceolatis basi cuneatis grossè serratis, capitulis laxè paniculatis, pedicellis capitulo vix duplo longioribus, achæniis papuloso-scabris. ☉? in ins. Moluccanis, Rawak (Cham.!) Timor (Gaudich.!), et in ins. Societatis (Forst.) *Lavenia erecta* Gaudich. in bot. voy. Freyc. 470. excl. syn. — Putumba Rheed mal. 10. t. 63 in arenosis Malabarix crescens, huc ex forma foliorum fortè referenda sed capitulis majoribus rufescentibus differt. (v. s.)
10. *A. PARVIFLORUM* (Blum. bijdr. p. 906. sub *Laveniâ*), caule erecto simplici glabriusculo, foliis petiolatis rhombeo-ellipticis obtusiusculis basi cuneatis repando-obtusè-serratis, corymbo terminali divaricato, ramis puberulis, invol. squamis dorso puberis, stigmatibus vix exsertis, achæniis muricato-tuberculatis. ☉ in umbrosis ins. Javæ circa Buitenzorg. (v. s. in h. Blum.)
- a. divaricatum*, paniculâ laxiore valdè divaricatâ, invol. minùs puberulis. — in ins. Javâ. *Lavenia erecta* Blum.! bijdr. 906. non alior. (v. s. comm. à cl. Blum.)
11. *A. FASTIGIATUM* (Blum.! bijdr. p. 905. sub *Laveniâ*), caule erecto subramoso glabro, foliis in petiolum attenuatis ovato-lanceolatis obtusè acuminatis grossè et inæqualiter dentato-serratis glabris, paniculâ terminali fastigiatâ, ramis involucriisque glanduloso-pilosiusculis, achæniis pube minimâ et tuberculis densè scabridis. — in ruderatis et humidis ins. Javæ (Blum.!), ins. Marianarum (Hænke!), in Indiâ orient. ad Bologâ. *Lavenia divaricata* Ham. herb. L. erecta var. β . Wall.! cat. n. 328. Folia papyracea 5-6 poll. longa, $1\frac{1}{2}$ -2 poll. lata. (v. s.)

12. *A. ROYLEI*, caule erecto teretiusculo ramoso, foliis (saltem superioribus) alternis ovalibus basi cuneatis subsessilibus apice acutis grossè dentatis secus nervos ramis caulibusque brevissimè pulverulento-puberulis, corymbo laxo paniculato, invol. squamis ovali-oblongis obtusis, achæniis muriculatis. — in Ind. orient. regionibus ad radices Hymalayæ sitis legit cl. Royle. Flores albi. Affinis *A. elato* et hirsuto. (v. s. comm. à cl. inv.)
13. *A. ELATUM* (Don prod. fl. nap. 181), caule erecto obtusè tetragono glabro simplici, foliis ovato-oblongis subsessilibus basi attenuatis crenato-dentatis subobtusis glabris, paniculâ subconfertâ, invol. squamis lineari-oblongis obtusis glabris, achæniis sparsè muricatis. — in Nepaliâ. *Lavenia carnea* Wall. ! cat. et herb. n. 329 (exclusis specim. nonnullis casu mixtis). *Lavenia elata* Spreng. syst. 3. p. 445 ex ipso in litt. ad Wall. Folia summa interdum alterna. (v. s.)
- β. laxum*, caule ramoso, paniculâ polycephalâ divaricatâ. — in Nepauliâ et Silhet. *Lavenia alba* (seu alta) Wall. ! cat. et herb. n. 330. exclusâ var. *C. Lavenia nudiflora* hort. bot. calc. à var. *α* nullo modo nisi car. indic. differre videtur. (v. s. comm. cum var. *α* ab hon. aulâ merc. angl. Ind. orient.)
14. *A. LATIFOLIUM* (Don prod. 181), caule erecto ramoso apice præsertim puberulo hirsuto, foliis petiolatis basi cuneato-acuminatis latè ovato-rhombéis aut subcordatis vix acutis grossè serratis secus nervulos puberulis, paniculâ corymbosâ hirsutâ polycephalâ, invol. squamis subacutis dorso scabridis, achæniis muricato-tuberculatis. (I) in Nepauliâ. *Lavenia latifolia* Spreng. syst. 3. p. 445. Habeo ex pl. Wallichianis, sed ex confus. quadam in numer. serie locus proprius incertus. Specim. habeo à cl. Leschenault in montibus Nelligherryanis lectum. *Ageratum strictum* Sims bot. mag. t. 2410 huc ex icon. pertinere videtur, sed descr. desideratur. (v. s. comm. ab ampl. aulâ merc. angl. Ind. or.)
15. *A. RUGOSUM* (Wight in Wall. cat. n. 331. sub *Laveniâ*), caule erecto basi subtetragono ramoso puberulo seu scabrido, foliis petiolatis latè ovatis subcordatis irregulariter subtrilobis grossè dentatis nervis subtus prominulis reticulato-rugosis rigidis suprà glabrisculis subtus puberulo-scabridis, paniculâ laxâ polycephalâ, invol. squamis oblongis dorso scabridis, achæniis muricato-tuberculatis. — in Peninsulâ Indicâ legit cl. Wight. DC. in Wight. contrib. bot. ind. p. 8. *Aden. aquaticum* Don prod. fl. nepal. p. 81. ipso mon. Lindl. Valdè affiné *A. reticulat.* sed achænia muriculata nec lævia? An character varians in hoc genere? (v. s. in h. hon. Cur. Ind. or. nunc Soc. Lin. Lond.)
16. *A. CAFFRUM*, caule erecto basi glabro apice pedicellisque pube brevi molli subvelutinis, foliis petiolatis ovato-rhombéis longè acuminatis basi cuneatis grossè villosò-dentatis suprà glabris subtus secus nervos junioribus tenuissimè subvelutinis, corymbo trichotomo laxo, invol. squamis lineari-oblongis obtusis basi velutinis, achæniis glanduloso-muricatis. (I) in Africæ austr. dist. Cafferland legit cl. Drege! Affine *A. elato* et latifolio, sed capitula duplò ferè majora. Cor. albæ. Folia 5 poll. longa, 2 poll. lata. (v. s.)
17. *A. NATALENSE*, caule erecto hinc indè (præsertim sub foliorum jugis) petiolisque muricato-echinulatis, foliis petiolatis ovato-subcordatis basi cuneatis apice acutis dentatis suprà puberulis subtus hirtellis, corymbo laxo subpaniculato, pedicellis involucrisque puberulis, achæniis muricatis. — in Africâ australi-orient. ad Port Natal legit cl. Drege! Caulis è nodis infimis subradicans, circ. bipedalis. Petiol. pollicaris. Limbus folii 3 poll. longus, 2 poll. latus. Capitula florida parva. (v. s.)
18. *A. BRASILIANUM* (Cass. dict. 25. p. 363), caule erecto scabrido, foliis petiolatis latè cordato-ovatis obtusis secus petiolum subcuneatis grossè et irregulariter dentatis glabrisculis, corymbi polycephali ramis puberulis, invol. squamis glabris ovali-oblongis obtusis, corollâ sub lobis hirsutâ, achæniis muricatis. — in Brasiliâ (Vand. ex Cass.), ad Sanctam-Catharinam (Bacle). Verbesina

Brasiliæ Pers. ench. 2. p. 472. Petiol. 12-15 lin. Folia 2 poll. longa, 2 $\frac{1}{2}$ poll. lata. (v. s. comm. à cl. Bacle.)

20. *A. TRIANGULARE*, caule herbaceo erecto petiolisque subtrigosis foliis petiolatis basi truncatis aut subcordatis idè fere triangularibus dentatis glabriusculis, corymbo laxo paniculato, capitulis longè pedunculatis, invol. campanulato truncato, squamis subconcretis corollà sub lobis hirsutà, achæniis subnuberculatis, pappi setis 2-3. 2/ in montosis ad margines rivulorum circa Rio-Janeiro legit cl. Lond. Flores albi. Rhizoma obliquum. Caulis pedalis. Petiol. pollicaris. Limbus fol. 2 poll. longus, 1 $\frac{1}{2}$ latus. (v. s. comm. à Mus. reg. Ber.)

*** *Caule erecto, achæniis lævibus.*

20. *A. RECTUM*, caule erecto simplici glabro, foliis petiolatis ovatis basi acuminatis grossè dentatis glabris, paniculà divaricatà laxà, ramis glanduloso-puberulis, invol. squamis glabris obtusis, achæniis lævibus. — in Zeylonà et in Ind. orientali (Wall.!) comp. n. 328 et 329, sed in n. 327 est *Myriactis* spec. (forte num. transp.). Burm. Zeyl. p. 95. t. 42. *Lavenia* erecta Willd. sp. 3. p. 1725? excl. Rheed. et Forst. syn. *Verbesina Lavenia* Linn. fl. Zeyl. 310. excl. syn. Plures species sub hoc nomine in herbariis et in libris meo sensu remanent confusæ! (v. s.)

21. *A. HIRSUTUM* (Blum. ! bijdr. p. 905. sub *Lavenià*), caule erecto subramoso glanduloso-pilosiusculo, foliis petiolatis infer. ovato-lanceolatis utrinquè acuminatis à medio ad apicem argutè serratis, paniculà terminali virgatà depauperatà, ramis glanduloso-puberis, invol. squamis subobtusis dorso puberulis, achæniis subscabridis. — in humidis insulæ Javæ ad montem Gede. (v. s. comm. à cl. Blum.)

22. *A. MADURENSE* (DC. in Wight contrib. bot. ind. p. 9), caule erecto juniore cum foliis brevissimè puberulo adulto glabrato, foliis petiolatis latè ovato-rhombis utrinquè acuminatis à basi grossè et acutè dentatis, corymbo suprà ultima folia breviter exserto ramis puberulis, invol. squamis glabris subscutis, achæniis lævisculis. — in Indiæ orient. peninsulà Maderaspatanà prope Madura legit cl. Wight. *Lavenia macrophylla* Wight! herb. non malè refert ic. Barm. Zeyl. t. 42. sed corymbus minùs exsertus et folia acutè nec obtusè dentata. (v. s. in h. Wight.)

23. *A. RETICULATUM* (DC. in Wight contrib. bot. ind. p. 8), caule erecto subtetragono glabro secùs paniculæ divaricatisimæ ramos parçè glanduloso-puberulo, foliis ovatis grossè dentatis rigidis, nervis subtùs prominulis reticulatis puberulis, invol. squamis lineari-oblongis obtusis dorso subpubescentibus, achæniis lævibus. — in Ind. orient. Maderasp. australi circa Dendigul legit cl. Wight. Folia infer. non vidi. Species distinctissima nervatione foliorum! (v. s. in h. Wight.)

24. *A. LEOCARPUM* (DC. in Wight contrib. bot. ind. p. 9), caule erecto tetragono glabro secùs paniculæ ramos glanduloso-puberulo, foliis ovali-oblongis subobtusis basi attenuatis subpetiolatis grossè dentatis glabris, paniculà laxà polycéphalà, invol. squamis lineari-oblongis obtusis dorso vix puberulis, achæniis sublævibus. — in Ind. or. *Lavenia alta* Wall. ! cat. et herb. n. 3220. comp. n. 330. *Ageratum aquaticum* Roxb. ! herb. sed diversum ab *Aden. aquatico* Don. Herba fere *A. elati*, fructus *A. Mauritanii*. (v. s.)

† *Species recognoscendæ.*

25. *A. MACROPHYLLUM* (Blum. ! bijdr. 905 sub *Lavenià*) caule erecto simplici glabro, foliis infer. longè petiolatis subrhombico-ovatis utrinquè acuminatis à medio ad apicem distanter serratis, paniculæ laxæ ramis vix puberulis, invol. squamis linearibus obtusis ciliolatis, achæniis.... — in locis humidis montis Gede in ins. Java. (v. s.)

! ? *repens*, caule subramoso infernè repente, foliis angustè serratis, ramis paniculæ subaltratis — ad cataractas fervidas montis Gede. Huc relata ex auctoritate cl. Blume l. c.

26. *A. AQUATICUM* (Don fl. nep. prod. p. 81), caule glabro, foliis cordatis serratis acutis glabris. ☉ in Nepaliæ inundatis. *Ethulia aquatica* Roxb. mss.
27. *A. PLATYPHYLLUM* (Cass. dict. 25. p. 363), caule erecto, pilis glandulosis sparsis, foliis oppositis petiolatis glabris ovato-deltoides secus petiolum cuneato-decurrentibus grossè crenato-dentatis, paniculâ corymbiformi laxissimâ. — in Peruvia legit Jos. de Jussieu. Fruct. indescr.
28. *A. BIFLORUM* (Less. syn. 136). — Huc cl. Lessing. refert sine descript. Verbesinam bifloram Linn. sp. 1272. sed descr. Linn. ab *Adenostemmati* caract. differt floribus luteis, pedunculis axillaribus bifloris et achæniis muticis. Icon Rheed. mal. 10. t. 40. à Willd. sub *Verbesinâ biflorâ* citatâ ab *Adenost.* omninò diversa. Species idèò omninò dubia.

† *Species exclusæ.*

LAVENIA DENTATA Wall. = *Myriactis Wallichii*.

LAVENIA RIGIDA Wall. = *Leptocoma racemosa*.

LAVENIA SPHERANTHA Wall. = *Myriactis Nepalensis*.

LXIX. *SCLEROLEPIS* Cass. dict. 25 (1816). p. 365. 26. p. 227 et 233. 48. p. 155. Less. syn. 136. non Monn. — *Sparganophorus Michx.* Nutt. Ell. non Gærtn. — *Ethulia* sp. Walt.

Capitulum multiflorum. Invol. squamæ biseriales lineares æquales. Recept. nudum. Corolla tubulosa glabra 5-dentata, fauce vix à tubo distinctâ. Styli rami exserti cylindræo-clavati. Achænium 5-angulare. Pappi squamæ 5 uniseriales æquales subcornæ breves subovales obtusiusculæ. — Herba Boreali-americana glabra simplex. Folia verticillata linearia. Capitula sæpius solitaria, rariùs 2-3, ramos supernè aphyllis breves terminantia. Cor. pallidè purpureæ.

1. *S. VERTICILLATA* (Cass. dict. 48. p. 155). ? in aquosis humidisve pinetis Carolinæ mediæ (Walt. Michx.!), et Novæ-Cæsariæ (Greene! Torr.). *Ethulia uniflora* Walt. car. 195. *Sparganophorus verticillatus* Michx. fl. am. 2. p. 95. Nutt. gen. 2. p. 139. Ell. sketch 2. p. 312. (v. s.)

LXX. *PHANIA*. — *Stevia* sp. Linn. — *Agerati* sp. Kunth.

Capitulum 15-25-florum. Invol. squamæ 8-12 subuni- aut biseriales non imbricatæ. Corolla tubulosa glabra 5-fida, lobis acuminatis. Achænium angulatum glabrum. Pappi paleæ 5-8 oblongo-lineares ciliatæ. — Frutices Americani glabriusculi. Rami tenues. Folia opposita ovalia integerrima aut subdentata basi subattenuata. Invol. puberula. Capitula pauca pedicellata. Corollæ albæ. — Genus à *Stevia* ferè ut *Alomia* à *Piqueria* differt, habitu etiam sat distinctum, sed fortè dividendum. — Nomen à *Phanio* vetustissimo botanico Theophrasti cœvo.

Sect. I. *EUPHANIA*.

Corolla pappi longitudine æqualis, tubo vix nullo, fauce obovatâ, lobis brevibus subacutis. — Suffrutex multicanlis. Folia trifida aut trisecta.

1. *P. MULTICAULIS*, glaber, caulibus pluribus erectis, foliis petiolatis trilobis (nunc trifidis nunc trisectis) lobis obovatis obtusis subdentatis, capitulis 2-5 subcorymbulosis axillaribus 20-floris, invol. squamis 10-12 acutiusculis. ? in America calidiore, Santo-Domingo? seu Brasiliâ? Species distinctissima. Capitula parva albida. (v. s.)

Sect. II. OXYLOBUS. — *Oxylobus Moc. fl. mex. ic. ined.*

Corolla pappo triplō circiter longior, lobis acuminatis. — Frutices Mexicani. Folia integerrima aut subdentata.

2. *P. ARBUTIFOLIA*, ramulis hirtis, foliis apice subcrenatis, inv. 12-phyllo, capitulis 20-25-floris. ♂ in Mexici montibus Nauhcampatepetl vel Coñre del Perote. *Ageratum arbutifolium* H. B. et Kunth nov. gen. et sp. 4. p. 149. *Stevia arbutifolia* Willd. herb. ex Less. in linnæa 1830. p. 140.
3. *P. TRINERVIA*, ramulis glabris, foliis integerrimis, invol. 8-phyllo, capitulis 15-16-floris. ♂ in Mexico. *Oxylobus trinervius* Moc. ic. fl. mex. Pappus dicitur 8-paleaceus et achænium tetragonum. (v. ic. pict.)

LXXI. *STEVIA* Cav. icon. 4 (1797). p. 33. Lag. nov. gen. et sp. (1816). p. 26. H. B. et Kunth nov. gen. et sp. 4 (1820). p. 139. Cass. dict. 26. p. 227. Less. syn. 135.

Capitulum 5-florum. Invol. cylindricum squamis 5-6 acutis acuminate subæqualibus. Recept. nudum. Achænium striato-nervosum angulatumve elongatum. Pappus paleis 1-serialibus nunc in iisdem aut sæpius in diversis individuis omnibus scariosis planis parvis, nunc aliis scariosis, aliis (pler. alternis) in aristas 1-5 rigidas margine scabras abeuntibus, nunc omnibus (pler. 5-20) in aristas mutatis. — Suffrutices aut herbæ omnes ex Americâ præsertim è Mexico, rariùs ex Amer. merid. Folia infer. opposita, super. opposita aut alterna, in unicâ ternato-verticillata, linearia lanceolata aut ovata, sæpius triplinervia integerrima serrata aut in unicâ trifida. Capitula in corymbum densum laxumve disposita, rariùs subsparsa longius pedicellata. Cor. albæ, rosææ aut purpureæ, interdum subochroleucæ.

* *Exaristatæ, nempe pappo exaristato paleaceo scarioso, paleis 5-6 nunc liberis parvis nunc in coronulam dentatam concretis.*

a. Herbaceæ perennes.

1. *S. TRIFLORA*, caule herbaceo erecto tereti oppositè ramoso tenuissimè præsert. ad apicem pulverulento-velutino, foliis oppositis lanceolatis in petiolum subattenuatis trinerviis apice serratis suprâ glabris subtus pulverulento-velutinis, corymbo laxo composito fastigiato myriocephalo, invol. squamis acutis glabriusculis, floribus sæpius in invol. 3, pappo coroniformi dentato brevi. ♀ in Mexico legit cl. Karwinski. Cor. extus subglandulosæ ochroleucæ. (v. s. in h. Acad. reg. Monac.)
2. *S. CORONIFERA*, herbacea glabra erecta, caule tereti oppositè ramoso, foliis oppositis petiolatis ovali-lanceolatis utrinquè acuminatis subdentatis, corymbo laxè composito, capitulis ad apices ramulorum confertis, invol. squamis acuminatis, floribus circ. 5, pappo coroniformi brevi subintegro. — è Mexico misit cl. Mairet. Cor. albæ glandulosæ. (v. s. comm. à cl. inv.)
3. *S. TRACHELIODES*, caule herbaceo erecto densè pubescenti-velutino, ramis oppositis alternisque subcorymbosis, foliis pl. oppositis basi caucatis sessilibus integerrimis cæterum latè ovatis acutis triplinerviis grossè dentatis utrinquè minutè pubescentibus subtus pallidis subvillosis, corymbis confertis polycephalis, inv. squamis puberis mucronato-acuminatis, corollâ puberulo-hirtellâ, pappo coroniformi brevissimo. ♀ in Mexici valle Tolucana legit cl. Berlandier. Species distinctissima. Pappo perexiguo ferè transit in Aloniâ seu Piqueriam sed evid. *Stevia* congener. Cor. purpurascens. Folia 3 poll. longa, $1\frac{1}{2}$ -2 lata. (v. s. comm. à cl. Berl. sub n. 1164.)
4. *S. MONARDEZOLIA* (H. B. et Kunth nov. gen. am. 4. p. 147), caule herbaceo apice ramoso hispidò-hirto, ramis fastigiatis polycephalis folisque suboppositis

petiolatis ovatis acutis duplicato-serratis basi angustatis triplinerviis involucri- que hirtis, capitulis acutè congestis, pappo paleaceo brevi. $\frac{1}{2}$ in Mexico versus Valladolid. Cor. violaceæ. (v. s.)

5. *S. CLINOPODIA*, caule herbaceo erecto puberulo, foliis oppositis in petiolum brevem attenuatis ovato-lanceolatis parvè dentatis utrinquè puberulis tenuiter nervosis, ramis oppositis paniculatis apice corymbiferis, corymbis confertis, invol. puberuli squamis mucronatis, pappo brevi paleaceo subacuto. $\frac{1}{2}$ in Mexico leg. cl. Alaman et Mairet. Corollæ purpureæ involucri paulò longiores extus glabriusculæ. Internodia distantia. Caulis fuscus. Variat foliis ovatis et lanceolatis. (v. s.)

6. *S. JORULLENSIS* (H. B. et Kunth nov. gen. et sp. p. 144), caule herbaceo apice ramoso pubescente, ramis sparsis abbreviatis corymbosis subfastigiatis polycephalis, foliis oppositis petiolatis ovatis obtusis crenatis basi cuneatis triplinerviis involucrique puberulis, pappo paleaceo brevi. $\frac{1}{2}$ in monte Jorullo Mexici. Cor. purpurea.

7. *S. ELONGATA* (H. B. et Kunth nov. gen. am. 4. p. 144), caule herbaceo erecto apice ramoso piloso-hirto, ramis alternis elongatis polycephalis, foliis oppositis sessilibus obovatis basi cuneatis serratis triplinerviis piloso-hirtis, super. rhomboideo-ovatis acutis, capitulis longè pedunculatis, involucri glanduloso-hirto, pappo scarioso brevi irregulariter fisso. $\frac{1}{2}$ prope Santa-Fe de Bogota. Affinis St. Jorullensi et St. ovata. Cor. glabra.

8. *S. CONFERTA*, caule herbaceo erecto piloso aut pubescente paniculato, foliis pl. alternis adproximatis sublinearibus integerrimis aut subdentatis puberulis subglandulosis, corymbis confertis polycephalis, invol. squamis glabriusculis apice acuminatis, pappi paleis 5-6 approximatis exaristatis. $\frac{1}{2}$ in Mexico prope Guanaxuato (Née) et Mexico (Berl.). Cor. purpureæ involucri apice purpurascente paulò longiores. Achænia glabra.

a. pilosa, caule pilis articulatis subhispido. — St. pilosa Lag. et Kl. b. madr. 1805. nov. gen. et spec. 26. n. 341. Pappi paleæ obtusæ. Invol. apice vix patentia. (v. s.)

β. puberula, caule pube brevi confertâ velutino-pubescente. — Pappi paleæ acutiores. Invol. apice subpatula. (v. s.)

β. Fruticosa.

9. *S. CORIACEA* (Lag. nov. gen. et sp. p. 28. n. 355), fruticosa, foliis oppositis ternisve subcoriaceis latè lanceolatis argutè serratis summis linearibus acutissimis integerrimis} corymbo umbellato polycephalo compacto, pappo paleaceo exaristato. $\frac{1}{2}$ in Peruvîâ.

10. *S. SUBPUBESCENS* (Lag. nov. gen. et sp. p. 28. n. 352), fruticosa minutè pubescens, ramis teretibus, foliis oppositis petiolatis ovato-lanceolatis acutis hinc indè serratis subtus pubescentibus, corymbo fastigiato composito, invol. puberulis, pappi subcoroniformis paleis brevibus exaristatis. $\frac{1}{2}$ in Mexico (Lag.) St. lanceolata Moc. fl. mex. ic. ined.? Cor. ex albo carneæ. Specimen habeo à cl. Berlandier in montibus Guchilacensis lectum huic descr. simile, sed foliis suprâ glandulosis. (v. s.)

11. *S. GLUTINOSA* (H. B. et Kunth nov. gen. am. 4. p. 148. t. 353), fruticosa glutinosa glabra, ramulis alternis corymboso-fastigiatis, foliis oppositis longè petiolatis ovato-oblongis utrinquè acutis argutè serratis, corymbis capitato-congestis polycephalis subsessilibus, achæniis. . . . pappo paleaceo brevi exaristato. $\frac{1}{2}$ in plantis Novo-Granatensium prope Santa-Fe de Bogota et Suacha. Cor. albæ? Petioli 9-10 lin. longi, quâ notâ differt ab omnibus et specialiter à St. lucidâ ad quam cæterum accedit. Achænia ex icone puberulo-hirta sed de hoc caract. descriptio silet.

β? Oaxacana, achæniis glaberrimis. $\frac{1}{2}$ in Mexici ditione Oaxaca legit cl. Andrieux (pl. exs. mex. n. 285). Optimè refert iconem et descr. var. α sed folia longiora ($3\frac{1}{2}$ poll.) et ferè angustiora (9-12 lin.) minùs serrata.

Cor. ferè purpurascens et certè achænia glabra. An species propria?, suadente patriâ sat distante. (v. s.)

12. *S. LUCIDA* (Lag. nov. gen. am. 28), suffruticosa glabra glutinosa, foliis oppositis petiolatis lato-lanceolatis dentatis suprâ lucidis penninerviis, ramulis oppositis, corymbo conferto polycephalo, pappo coroniformi paleaceo subintegerrimo. ζ in Mexico ad Ixmiquilpan et Cimnapan legit cl. Née. St. hyemænoppa Moc. fl. mex. ic. ined. Achænia angulata glabra. Cor. glabræ ex sicco albide? Petioli 3 lin. longi, limbus 2 poll. longus 6-8 lin. latus. Affinis videtur *S. glutinosa*: (v. s. ex h. Née sine nom.)

13. *S. ANGUSTIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 149. excl. var. α ?), suffruticosa, caule supernè puberulo-hirto, ramis fastigiato-corymbosis oppositis rariùs alternis, foliis oppositis breviter petiolatis lineari-lanceolatis utrinquè angustatis glabris subtùs involucrioque subglutinosi, capitulis dense conglomeratis, pappi subcoroniformis paleis ovatis acutis brevissimis. ζ in Mexico. Cor. albæ. Achænia angulata vix subscabrida. Pappum in floribus 1-2 inter sexcentas vidi 1-aristatum. St. salicifoliæ affinis.

α . *integrifolia*, foliis integerrimis. — Folia 3 poll. longa 5-6 lin. vix lata. Videtur diversa à var. α Kunthianâ, fortè ipso auct. momento à specie separandâ. (v. s.)

β . *subserrata*, foliis remotè serratis. — St. angustifolia β Kunth.

- ** Paleaceo-aristatæ, nempe pappo paleis paucis scariosis brevibus et aristis 1-5 rigidulis scabrido-serrulatis constante.

α . Fruticosæ.

14. *S. OLIGOCEPHALA*, caule gracili imâ basi fruticuloso et ramoso infernè hispidulo, foliis oppositis lineari-lanceolatis petiolulatis tenuiter trinerviis à mediò remotè serratis glabris, paniculâ laxâ oligocephalâ subnudâ, capitulis longè pedicellatis, invol. squamis glabriusculis acuminatis, pappo brevissimè paleaceo et 1-3-aristato. ζ in Brasiliæ prov. Minarum Gener. ad Serro Frio legit cl. Vauthier (pl. exs. n. 309) Species elegans distinctissima. Cor. tubus purpureus, limbus albidus. (v. s. comm. à cl. Vauthier.)

15. *S. HÆRVOSA*, fruticosa, ramis breviter et dense velutinis, foliis oppositis alternis ellipticis basi attenuatis apice subacutis subserratis utrinquè tenuiter pubescentibus penninerviis, nervis subtùs prominulis, ramulis summis alternis involucriisque subhirtellis, corymbis confertis, achæniis glabris, pappo paleaceo et 1-3-aristato. ζ in Mexico prope Villalpando ad orient. urbis Guanajuato legit cl. Mendez. Corollæ pubescentes albæ. Achænia nigra. Folia 12-15 lin. longa 5-6 lata minutè subtùs glandulosa, nervis elevatis albidis distincta. (v. s.)

16. *S. SALICIFOLIA* (Cav. ic. 4. p. 32. t. 354), fruticosa glabra, foliis oppositis angustè lanceolatis brevissimè petiolatis et subconnatis integerrimis aut mediò subserratis utrinquè attenuatis, corymbo patente, pappi paleis nunc omnibus brevibus scariosis nunc 1, 2 aut 3 in setas aristatas exsertis. ζ in Mexici montanis (Berl. ex. n. 868. 675. 825. 1202.), juxta Ixmiquilpan (Lag.) et alibi. Lag. nov. gen. p. 28. Ageratum viscosum Ort. dec. p. 36. ex Lag. St. semperflorens Ten. in litt. Rami et folia subtùs sæpè subviscosa. Pedunculi pube brevi villosiusculi. Cor. albæ. (v. s.)

17. *S. CONNATA* (Lag. elench. h. matr. 1815. nov. gen. et sp. p. 27. n. 351), fruticosa stricta glaberrima, foliis sessilibus oblongo-lanceolatis serratis acutissimis, infer. oppositis connatis, super. subparais, corymbo fastigiato, invol. 5-7-floris, pappo paleaceo et aristato. ζ in Mexico. St. glabra Moc. fl. mex. ic. ined. Corollæ albæ. An fortè eadem ac *S. viminea*?

β . Herbacææ.

18. *S. VIMEINA* (Schrad. hort. Goett. 1832), herbacea glabra erecta, caule simpliciter apice corymboso, foliis oppositis alternisve basi semiamplexicaulis elongato-linearibus acutis apice subserratis, corymbo umbellato composito.

polyccephalo laxiusculo, inv. squamis acutis, pappo paleis brevibus 2-4 et aristis 1-3 constante. \mathcal{L} in Mexico. St. longifolia Moq. fl. mex. ic. ined. St. umbellata Alam. in litt. 1831. Valdè accedit ad Scannatam Lag. sed herbacea. Cor. albæ invol. ferè duplò longiores. (v. s.)

19. *S. serrata* (Cav. ic. 4. p. 33. t. 355), caule herbaceo erecto ramoso pubescente, foliis alternis subfasciculatis lineari-lanceolatis glabrisculis serratis basi integerrimis in petiolum attenuatis, corymbis fastigiatis, pappo paleaceo et bi- sæpius triaristato. \mathcal{L} in Mexici montanis, circa urbem, in valle Toluca (Berl.) ad Temascatio et Guanaxuato (H. et B.), etc. Lag. nov. gen. et sp. n. 336. H. B. et Kunth nov. gen. am. 4. p. 141. Moq. fl. mex. ic. ined. Ageratum punctatum Jacq. h. Schœnb. 3. p. 28. t. 300 non Ort. An St. virgata H. B. et Kunth nov. gen. am. 4. p. 142. satis differt? Variat caule erecto aut demisso, foliis lanceolatis aut ferè linearibus sæpè subsecundis, corollis albis aut ex albo roseis, pubescentiâ parvâ aut subhirsutâ (v. v. c. et s. sp.)

20. *S. ivæfolia* (Willd. mag. ges. naturf. Berl. 1807. p. 137. enum. 855), caule herbaceo erecto hirsuto-villoso apice ramoso-corymboso, foliis lanceolatis in petiolum attenuatis triplinerviis apice serratis, corymbis fastigiatis, invol. et corollâ glandulosi et puberis, pappo brevi paleaceo et hinc indè 1-2-aristato. \mathcal{L} in Mexico. Affinis St. serratæ sed caulibus villosis-hirsutis, folia sæpius latiora plana. (v. v. c. in hort. Par.)

ß? *Bogotensis*, foliis minoribus non triplinerviis suprâ glabris et viscosis, corymbo coerctato, pappi aristis ternis. \mathcal{L} in altâ plantie Bogotensi inter Santa Fe et Chipo. St. ivæfolia H. B. et Kunth nov. gen. am. 4. p. 142. sed à verâ ivæfoliâ patriâ et caract. videtur aliena.

21. *S. eupatoria* (Willd. sp. 3. p. 1775. enum. 854. mag. naturf. Berl. 1807. p. 138), herbacea subdiffusa glabriscula, foliis lanceolatis in petiolum subattenuatis trinerviis supernè obsolete serratis, corymbis fastigiatis laxiusculis, pappo paleaceo et sæpius 2-4-aristato. \mathcal{L} in Mexico. St. hyssopifolia Cav. præl. 465. Lag. nov. gen. et sp. 26. n. 340. non Kunth nec Sims. Ageratum punctatum Ort. dec. 3. p. 37. (ex Lag.) non Jacq. St. punctata Pers. ench. 2. p. 403. Mustelia eupatoria Spreng. trans. Linn. soc. Lond. 6 p. 152. t. 13. St. eupatoria Sims bot. mag. t. 1849. Corollæ involuero duplò longiores, tubo incarnato, limbo intus albo. Achæn. glabra. (v. v. et s.)

22. *S. purpurea* (Pers. ench. 2. p. 403 excl. syn.), caule herbaceo erecto pubescenti-velutino ramosissimo, foliis lanceolatis alternis, inferioribus obovatis canaliculatis in petiolum angustatis triplinerviis apice serrulatis, corymbis subcoerctato, pappo paleaceo et triaristato. \mathcal{L} in Mexico (Pers. Moq.), ad radices montis Orizabæ (Sch. et Dep.). Willd. enum. 855. Moq. fl. mex. ic. ined. Less. in linnæa 1830. p. 139. Jacq. fragm. 80. t. 127. f. 2. St. eupatoria bot. reg. t. 93. Corollæ purpureæ extâs puberulæ. Invol. vix puberum pallidè virescens. An satis à St. eupatoriâ diversa? An St. stricta Horn. hort. hafn. 2. p. 799. huc revoc. ? (v. v.)

23. *S. microphylla* (H. B. et Kunth nov. gen. am. 4. p. 140), viscoso-hirto, caule herbaceo paniculato-ramoso, ramis floriferis dichotomis polyccephalis foliisque oppositis subsessilibus lineari-lanceolatis obsolete crenatis, invol. squamis acuminatis, pappo breviter paleaceo et triaristato. \mathcal{L} in uliginosis inter Tasco et Thuilopec Mexicanorum. Cor. glabra alba aut rosea.

24. *S. origanoides* (H. B. et Kunth nov. gen. am. 4. p. 147), caule herbaceo oppositè ramosissimo hirto pubescente, foliis oppositis sessilibus lanceolatis angustato-acuminatis medio serratis triplinerviis utrinquè tenuissimè hirtis, capitulis fasciculatis congestis, involucri squamis pubescentibus vix acutis, pappo breviter paleaceo et triaristato, aristis cor. puberâ brevioribus. \mathcal{L} in Mexico. Habitus Origani. Cor. albæ. Achænia glabra. (v. s.)

25. *S. microcephala*, caule herbaceo erecto simplici hirsuto-villoso, foliis alternis subsessilibus ovalibus obovatisve basi attenuatis integerrimis apice serratis dentatis subtripplinerviis utrinquè puberulo-hirtis, summis lanceolatis

linearibusve acutis, corymbo conferto alternè ramosissimo polycephalo, capitulis pedicello proprio multò longioribus, achenio glabro, pappo brevi paleaceo et 3-5-aristato, aristis cor. extus puberulâ paulò brevioribus. $\frac{1}{2}$ in Mexico legit cl. Hænke. Capitula 3 lin. longa. Cor. ex sicco pallidè purpurascens. Folia 9-10 lin. longa, 4-5 lata. Au St. incanescens Lag. nov. gen. et sp. n. 348? an fortè eadem ac. St. pubescens Lag.? (v. s.)

26. *S. LANCEOLATA* (Lag. nov. gen. et sp. 26. n. 338), caule herbaceo subhirsuto supernè ramoso-corymboso, foliis lanceolatis acutis planiusculis in petiolum attenuatis subtùs scabris, summis lineari-lanceolatis integerrimis, corymbo fastigiato laxiusculo, pappo paleaceo et triaristato. $\frac{1}{2}$ in Mexico circa Pachua. Var. duas dist. cl. Lagasca.

a. floribus albis, foliis subtùs glabris.

β. floribus purpureis, foliis subtùs subhirsutis.

27. *S. CANESCENS* (H. B. et Kunth nov. gen. am. 4. p. 143), caule herbaceo apice corymboso-ramosissimo polycephalo villosio-tomentoso, ramis subfastigiatis foliisque sparsis subsessilibus lanceolatis basi angustatis serratis hirsuto-pubescentibus, involucri pilosiusculo, pappo paleaceo et triaristato. $\frac{1}{2}$ in Mexico inter Guanaxuato et montem Serena. Cor. alba.

28. *S. PUBESCENS* (Lag. nov. gen. et sp. 26. n. 337), caule herbaceo simpliciterectiusculò pubescente, foliis infer. oppositis subspathulatis apice dentatis in petiolum attenuatis, super. sparsis linearibus subintegerrimis, corymbo fastigiato densiusculo, invol. pubescentibus, pappo paleaceo et triaristato. $\frac{1}{2}$ in Mexico circa Guanaxuato et Chalma. Cor. purpureo-sanguinea. — St. pubescens H. B. et Kunth nov. gen. am. 4. p. 143. Videtur diversa foliis lanceolatis, corollis albis, etc.

29. *S. CARACASANA*, caule herbaceo erecto tenuiter puberulo, foliis oppositis in petiolum attenuatis ovatis acutis basi cuneatis dentato-serratis subtriplinerviis subtùs glandulosis puberulisque, ramis sup. oppositis paniculatis apice corymbos polycephalos confertos gerentibus, invol. puberuli squamis apice mucronatis, pappo breviter paleaceo et 3-aristato corollam æquante. $\frac{1}{2}$ circa Caracas detexit cl. Vargas. Glandulæ fol. in sicco nigricantes. Invol. apice purpurascens. Cor. albidæ, tubo ochroleuco. (v. s. comm. à cl. invent.)

30. *S. TERNIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 146), caule herbaceo obsoletè trigono ramosissimo subpubescente, ramis fastigiatis polycephalis foliisque ternis sessilibus oblongis acutis crenato-serratis triplinerviis involucrique glabris, capitulis pedicellatis fasciculatis, pappo paleaceo et subbiaristato. $\frac{1}{2}$ in provinciâ Mechoacanensi Mexici. Cor. alba. Species foliis ternato-verticillatis-facilè distincta.

31. *S. NEPTEFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 145), caule herbaceo erecto paniculato-ramoso tomentoso-pubescente, ramis oppositis fastigiatis apice corymbiferis, foliis oppositis ovato-lanceolatis basi longè cuneatis et subsessilibus supernè triplinerviis grossè serrato-crenatis suprâ tenuissimè pubescentibus subtùs pallidis pubero-subtomentosis, corymbis confertis, invol. pubescenti-hirtellis, pappo paleaceo et 3-aristato. $\frac{1}{2}$ in planitie Mexicanâ (H. et B.) inter Vittoria et Tula (Berl.!) circa Guanaxuato (Lag.?) St. suaveolens Lag. nov. gen. et sp. 27. n. 342? St. pubescens Moc. fl. mex. ic. ined. Cor. albæ aut fortè rosæ extus puberulæ. Aristæ cor. æquales. Achen. glabra. (v. s. comm. à cl. Berlandier.)

32. *S. RHOMBIFOLIA* (H. B. et Kunth nov. gen. am. 4 p. 143), pubescens, caule herbaceo apice subramoso polycephalo, foliis oppositis petiolatis rhomboidè-ovatis serrato-crenatis triplinerviis, capitulis fasciculatis breviter pedicellatis, invol. squamis acutis, pappo triaristato tubum cor. basi pubescentem æquante. $\frac{1}{2}$ in collibus montis Sanctæ Guadalupæ prope Mexico. Cor. alba.

33. *S. FASTIGIATA* (H. B. et Kunth nov. gen. am. 4. p. 148) glutinosa, caule herbaceo ramoso, ramis superioribus corymboso-fastigiatis polycephalis foliisque oppositis breviter petiolatis ovato-oblongis, super. oblongo-lanceolatis

acutis grossè serratis basi cuneatis glabris, capitulis congestis, pappo brevi paleaceo rariùs 1-aristato. $\frac{1}{2}$ locis scopulosis Mexici. Affinis *St. glutinosæ* dicitur.

34. *S. UNIARISTATA*, caule herbaceo erecto pubescente subhirtello, foliis oppositis ovato-lanceolatis acuminatis in petiolum brevem attenuatis dentatis subtriplinerviis utrinquè puberulis, ramis oppositis apice corymbiferis in paniculam dispositis, corymbis confertis, involucri puberuli squamis mucronatis, achæniis capituli circ. 4, pappo paleaceo brevi obtuso, uno aristam unicam gerente. $\frac{1}{2}$ in montibus Leoninis ad occid. urbis Guanajuato legit cl. Mendez. *St. incanescens* Lag. nov. gen. et sp. 27. p. 348. var. β ? et γ ? Folia subtus nervosa. Corollæ in sicco flavidæ involucri paulò longiores. Folia summa oblonga lanceolata utrinquè acuminata integerrima. (v. s.)
35. *S. FASCICULARIS* (Less. in linnæa 1830. p. 140), caule herbaceo, foliis oppositis rhomboideo-lanceolatis profundè et argutè serratis vix pilosiusculis, super. sessilibus, ramis inflorescentiæ fastigiatis aut subfastigiatis plerisque oppositis, capitulis fasciculatim congestis, corollis glabris, pappo florum cujusque capituli sæpiùs 2 brevi paleaceo, aliorum 3-aristato. — in Mexico prope Jalapam et Hacienda de la Laguna.
36. *S. TOMENTOSA* (H. B. et Kunth nov. gen. am. 4. p. 145. t. 352), caule herbaceo supernè fastigiato-ramoso pubescente, foliis alternis rariùs oppositis basi cuneatis et petiolatis ovatis dentato-serratis triplinerviis suprâ tenuissimè pubescentibus subtus albo- et densè tomentosis, corymbis polycephalis confertis, invol. squamis puberulis, pappo nunc brevi paleaceo nunc 1-3-aristato. $\frac{1}{2}$ in Mexici et Tolucæ vallibus (Berl. l. exs. n. 1100. et 1196), prope Valladolid (H. B.) Cor. ex sicco incarnatæ. (v. s. comm. à cl. Alam. Mairet, etc.)
37. *S. MOLLIS* (Schrad. l. in litt. 1832), caule herbaceo pubescente supernè fastigiato-ramoso, foliis infer. oppositis petiolatis latè ovatis grossè dentato-serratis basi subcuneatis suprâ minutissimè pubescentibus subtus cano-tomentosis, super. oblongis sessilibus integerrimis, corymbis laxis oligocephalis, invol. squamis dorso puberulis, pappo nunc brevi paleaceo nunc 1-2-aristato $\frac{1}{2}$ patr. ign. Cor. ex sicco albæ. An fortè var. *St. tomentosæ*? (v. s. comm. à cl. Schrad.)
38. *S. ADENOPHORA* (Lag. nov. gen. et sp. 27. n. 347), caule herbaceo glanduloso-villoso, foliis oppositis sessilibus ovatis obtusis serratis triplinerviis, basi subdeltoideæ integerrimæ, paniculâ trichotomâ laxè ramosâ, pedicellis 1-cephalis villosis-glanduliferis, pappo 3-aristato, rariùs 2 seu 4-aristato. $\frac{1}{2}$ in Chili. Cor. purpureæ. Folia summa minora integerrima.
39. *S. PANICULATA* (Lag. elench. h. madr. 1805. nov. gen. et sp. 27. p. 343), caule herbaceo erecto breviter pubescente paniculato, foliis infer. oppositis ovatis triplinerviis, super. alternis ovato-oblongis serratis basi cuneatis integerrimis, summis linearilanceolatis, pedunculis parçè ramosis 3-4-cephalis laxè corymbosis, pappo nunc paleaceo brevi obtuso, nunc in iisdem capitulis 1-2-aristato. $\frac{1}{2}$ in Mexico (Lag.) *St. hyssopifolia* Sims bot. mag. t. 1861. non Cav. nec Kunth. Corollæ paulò involucri longiores albæ, tubo purpurascante. Achænia glabra 2-3 exaristata, 3-2 aristam 1-2 gerentia. (v. v. c.)
40. *S. OVATA* (Lag. elench. h. madr. 1805. nov. gen. et sp. 27. n. 344), caule herbaceo erecto glabriusculo paniculato, foliis ovatis triplinerviis serratis basi cuneatis integerrimis, super. oblongis subintegris, corymbis fastigiatis subcompactis, pappo plerumquè brevi paleaceo obtuso hinc indè uni- rariùs biaristato. $\frac{1}{2}$ in Mexico. Willd. enum. 855. Dum. Cours. bot. cult. 3. p. 93. Similis *St. paniculatæ* nec fortè satis distincta. (v. v. c.)
41. *S. ELATION* (H. B. et Kunth nov. gen. am. 4. p. 144), caule herbaceo erecto, ramis oppositis elongatis piloso-hirtis, foliis oppositis, infer. petiolatis subrotundo-ovatis obtusis grossè crenatis basi cuneatis pilosiusculis, super. subsessilibus oblongo-ovatis, capitulis pedicellatis, invol. glanduloso-hirto, pappo

brevi paleaceo et triaristato. $\frac{1}{2}$ prope Santa-Fé de Bogota in umbris humidis. Corolla alba extus tenuissimè glanduloso-hirta. Aff. St. elongata.

42. *S. QUITTENSIS* (H. B. et Kunth nov. gen. am. 4. p. 145), caule herbaceo erecto supernè ramoso viscoso-villoso, ramis subfastigiatis polycephalis foliisque oppositis petiolatis ovato-oblongis acutis grossè crenato-serratis triplinerviis subulis hispidulis, capitulis pedicellatis confertis, involucrio glabriusculo, pappo paleis parvis 3-1 et aristis 3-5 alternantibus. $\frac{1}{2}$ in apricis inter urbem Quito et montem Burro Potrero. Cor. alba glabriuscula.
43. *S. PODOCEPHALA*, caule herbaceo erecto villosa-subhirsuto in paniculam laxam divaricatam ramosissimam supernè diviso, foliis caulinis oppositis sessilibus basi cuneatis ovato-rhomboidèis grossè dentatis triplinerviis utrinquè pubescenti-hirtis, ramis alternis ovatis subintegris, ramis pedicellis pilosis glandulosis, invol. pedicello brevioribus puberulis, pappo 3-4-paleaceo et 3-2-aristato, aristis rigidis demum divergentibus. $\frac{1}{2}$ in Mexico leg. cl. Née, Alaman, Berlandier, Mendez. Flores invol. duplò longiores albidì extus parè puberuli. Achænia glabra. Pili caulis articulati. An ideò St. enarthotricha Lag. huc revocanda? sed in nostrà invol. puberum nec glabrum, folia (saltem mihi obvia) sessilia nec petiolata, corolla alba nec purpurea. (v. s.)
44. *S. HIRSUTA*, caule herbaceo erecto apice subnudo foliisque utrinquè pube articulata longà confertè subdecidua hirsutis, foliis in petiolum attenuatis basi cuneatis ovato-lanceolatis dentatis, capitulis fasciculatis sessilibus, fasciculis in corymbum trichotomum dispositis, invol. glabriusculis, pappo paleaceo et triaristato, aristis divergentibus corollà puberulà paulò longioribus. $\frac{1}{2}$ in Mexico ad Villalpando legit cl. Mendez. Pappus St. podocephala. Cor. ferè St. tenellæ. Hirsuties St. enarthotrichæ. Cor. pallidè purpurea ex sicco. Inv. purascens. (v. s. comm. à cl. Mendez.)
45. *S. MULTIFIDA* (fl. mex. ic. ined.), caule herbaceo filiformi piloso-glanduloso, foliis imis oblongis dentatis, summis alternis linearibus integerrimis, cæteris omnibus oppositis trifidis, lobis elongatis aliis dentatis aliis pinnato-incisais, corymbo laxo paucifloro, invol. squamis acuminatis dorso puberulis, achænio tenui glabro, pappo vix paleaceo et 3-aristato. ①? in Mexico, verisim. circa urbem. Species distinctissima. (v. s. in h. Hænk. ab ill. Sternberg miss.)
- ß. trifida*, foliis trifidis, lobo medio sæpè dente utrinquè exciso. — in Mexico. St. trifida Lag. nov. gen. et sp. 27. n. 350.
46. *S. MICRANTHA* (Lag. elench. h. madr. 1805. nov. gen. et sp. 27. n. 349), caule herbaceo sparè pubero paniculato, foliis oppositis alternisque ovatis serratis, inf. subcoarctatis, capitulis pedicellatis laxè subcorymbulosis, pappi paleis minimis, aristis 3 corollà pubescente tubulosà longioribus. ① in Mexico prope Leon ad occid. urbis Guanaxuato legit cl. Mendez. St. tenella Moq. fl. mex. ic. ined. Species distinctissima. Flores albidì minimi. (v. s. sp.)
47. *S. DISSESTIVOLIA*, herbacea glabriuscula erecta, ramis oppositis distantibus divaricatis, ramulis alternis, foliis oppos. alternisque dissitis sessilibus ellipticis integerrimis, sup. ovali-oblongis, capitulis pedicellatis ad apices ramulorum paucis, invol. squamis acuminatis, pappo breviter paleaceo et aristis 3-6 corollam vix puberulam paulò superantibus. $\frac{1}{2}$ in Mexico legit cl. Hæнке. Habitus ferè Hyperici pulchri. Internodia ferè 3 pollic. Folia 4-5 lin. longa 3-4 lata. Cor. parvæ albidæ. Ambigit inter secundam tertiamque divisionem. (v. s. in h. Hæнке à cl. Sternberg comm.)
48. *S. LEUCONEURA*, caule basi fruticuloso ramoso, ramis elongatis infernè pilis articulatis demum deciduis hispidis cæterum pedicellis involucriisque brevissimè puberulis, foliis oppositis et alternis lanceolatis utrinquè acuminatis serratis et super. integerrimis, nervis albis elevatis, foliis infer. basi penninerviis, super. uninerviis, paniculà laxà diffusâ, capitulis pedicellatis, pappo 4-5 aristato, aristis cor. æqualibus. $\frac{3}{5}$ $\frac{1}{2}$ in Mexico ad Leon ad occid. urbis Guanaxuato legit cl. Mendez. St. longiseta Moc. fl. mex. ic. ined.? Affinis St. laxifloræ. Invol. et cor. extus purpurascens. Achænia glabra aristis breviora (v. s. sp.)

49. *S. LAXIFLORA* (DC. h. monsp. 1813 et in Moç. fl. mex. ic. ined.), caule herbaceo pubescente ramosissimo, foliis linearî-lanceolatis plerisque alternis canaliculatis apice serratis, corymbo laxo, capitulis pedicellatis, pappo paleis parvis et aristis 5 alternantibus. $\frac{1}{2}$ in Mexico. St. purpurea (quinquearistata) Lag. nov. gen. et sp. 26. n. 339. non Pers. St. hyssopifolia H. B. et Kunth nov. gen. am. 4 p. 140. non Cav. nec Sims. Affinis St. purpureæ et cum eâ in hortis sæpe ob flores purpureos confusa. Differt staturâ majore, caule infernè pilis raris subhispidò, ramulis pube tenuissimâ subincanis, corymbo laxiore, corollis involucri duplò longioribus et aristis pappi 5 cum paleis parvis alternantibus. (v. v. c.)

50. *S. VISCIDA* (H. B. et Kunth nov. gen. am. 4. p. 140. t. 351), caule herbaceo viscoso-hirto supernè paniculato, foliis alternis sessilibus linearibus apice subcrenulatis uninerviis pubescentibus et glandulosis, corymbis oligocephalis in paniculam subcorymbosam dispositis, involucri squamis hirto-viscosis acuminatis, pappo 5-aristato epalaceo. $\frac{1}{2}$ in umbrosis circa Mexico (Humb. et Bonpl., Berl.!) Ageratum purpureum Sesse et St. villosa Moç. fl. mex. ic. ined. ? ex quo fortè *Pipitzahuatl* Mexicanorum. Affinis laxifloræ. Cor. purpureæ invol. duplò longiores extûs puberæ. An potiùs ad sect. 3^{am}? (v. s.)

*** Multiaristatæ, pappo paleis nullis aut subnullis sed aristis 6-20 constante.

51. *S. HEPTACHÆTA*, caule herbaceo ? gracili vix sparè pilosulo, foliis oppositis linearibus obtusis integerrimis vix pilosiusculis, infer. apice sublterioribus et hinc indè dentatis, paniculâ laxâ nudiusculâ, capitulis longè pedicellatis, pappo 6-8 aristato. $\frac{1}{2}$? in Brasiliæ prov. Minarum General. ad Sabarra legit cl. Vauthier. Caulis fortè imâ basi fruticulosus ad apicem foliis distantibus minoribusque subnudus. Invol. et pedicelli extûs puberuli. Aristæ æchanio puberulo longiores corollâ æquales, hinc indè in æchanio quodam capituli subnullæ. (v. s. comm. à cl. Vauthier sine nom. n. 310.)

52. *S. LUNDIANA*, subherbacea simplex erecta, caule glabriusculo tereti, foliis oppositis sessilibus oblongis erectis dentatis trinerviis subtusque subreticulatis glabris, paniculâ corymbosâ laxâ, invol. squamis glandulosis pubescentibusque subacutis, pappo 12-13-aristato. $\frac{1}{2}$ in campis editis prov. Sancti Pauli nov. flor. legit cl. Lund. Herba circ. pedalis. Cor. rubentes. Pappus subnigrescens. Habitu affinis St. heptachætæ. (v. s. comm. à cl. inv.)

53. *S. SUBOCTOARISTATA* (Lag. nov. gen. et sp. 27. n. 345), herbacea glabriuscula, foliis petiolatis obovatis venosis subserratis, corymbo fastigiato (Spr. syst. 3. p. 448), pappo 7-9-aristato (Lag.). $\frac{1}{2}$ in Peruvîâ. Habitus St. ovatæ. Cæst. ign. †.

54. *S. HÆNKEANA*, caule basi suffruticuloso erecto oppositè ramoso apice tenuiter velutino, foliis oppositis petiolatis ellipticis utrinquè attenuatis dentatis utrinquè punctulato-subscabris, capitulis in corymbos plurimos ramosos confertis, invol. squamis acuminatis, pappo 10-12-seto corollâ vix puberulâ evidenter breviorè æchanii longit. æquali. $\frac{1}{2}$ in Mexico legit cl. Hænke. Habitus ferè Origanî. Capitula breviter pedicellata. Achænia scabrinuscula. Cor. ex sicco purpurascens. Petioli 3 lin. longi. Limbus fol. 12-14 lin. longus 6 lin. latus. (v. s. in h. Hænke à cl. Sternberg comm.)

55. *S. DODECACHÆTA*, caule herbaceo adscendente supernè villosò confertè folioso, foliis oppositis sessilibus ovalibus grossè dentatis triplinerviis subpuberulis, corymbis confertissimis ad apices ramorum subsessilibus polycephalis, pappo constante aristis circ. 12 æchanio longioribus corollam æquantibus. $\frac{1}{2}$ in Peruvîâ circa Lima lectam misit cl. Abadia. Habitus ferè *Pycnanthemî*. Cor. tubus purpureus, limbus albidus. Invol. squamæ vix puberulæ breviter acuminatæ. (v. s.)

56. *S. HISPIDULA*, caule herbaceo erecto apice tenuissimè glanduloso-velutino, foliis oppositis oblongo-lanceolatis utrinquè attenuatis apice serratis margine et subtus in nervo sparè setoso-hispidis, corymbo terminali conferto, capitulis pedicellatis, invol. squamis lanceolatis acuminatis dorso puberulis æchanio

parcè longioribus, pappi setis 20-25 scabris corollæ intus ad faucem et extus subvillosæ subæqualibus. \mathcal{U} in Mexico verisim. legit cl. Née, saltem ex ejus herb. habui commixtam cum *St. vigintisetâ*. (v. s.)

57. *S. PUBERULA* (Hook. misc. bot. 2. p. 225), herbacea erecta pubera apice corymbosa, foliis alternis sessilibus ovalibus basi apiceque acutis obtusè serratis triplinerviis, corymbo denso, invol. glanduloso, pappi aristis subnatis scabris. — in Peruviae valle Santa ad Obrajillo. Cor. albæ aut carneæ.
58. *S. DIVARICATA*, caule herbaceo erecto pubescente, ramis alternis divaricatis, foliis alternis sessilibus lanceolatis integerrimis aut subdentatis utrinquè attenuatis parcè pubescentibus subtriplinerviis, capitulis in fasciculos 2-3-cephalos pedicellatos aggregatis laxè paniculatis, invol. brevishortus, pappo 15-seto. \mathcal{U} in Mexico? Cor. tubus purpurascens puberulus, limbus albidus pappi longitudinem vix superans. Achænium in capitulo sæpè adest unicam exaristatum. (v. s. in h. Mart.)
59. *S. POLYPHYLLA*, caule basi suffruticuloso erecto apice corymboso et pubescente, foliis oppositis in petiolum attenuatis linearibus subintegerrimis subnerviis pubescentibus, ramis ex axillis ortis subconfertis, fasciculis oligocephalis in corymbum dispositis, invol. squamis 5-6 pappo dimidio brevioribus vix acutis, pappo 12-16-seto, corollæ puberæ æquali, achæniis puberulis. \mathcal{J} inter Peruvianas et Panamenses herb. Hænk. recenset. Caulis pedalis strictus. Rami steriles frequentes foliosi breves. Cor. purpurascens. Habitus ferè *Silenes polyphylla*, (v. s. in h. Hænk. comm. à cl. Sternberg.)
60. *S. VERONICÆ*, caule herbaceo basi suffruticuloso erecto simplici glanduloso-piloso, foliis oppositis oblongis basi cuneatis integris apice serratis subacutis minutè et sparse puberulis, corymbo fastigiato, capitulis pedicellatis, invol. squamis lanceolatis pubescentibus corollas ferè adæquantibus, pappo 15-16-seto stramineo scabrido corollam glabram subæquante. \mathcal{U} in Brasiliæ prov. Minas Generalium ad Serro do Frio legit cl. Vauthier (n. 325!). Cor. vid. roseæ ex sicco. Achænium glabrum. (v. s. comm. à cl. inv.)
- β . *erythrochata*, pappi setis rubescentibus. — in prov. Rio-Grande. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 943 miss.)
- γ . *glabrata*, invol. squamis glabris. — in prov. Rio-Grande (v. β . in eod. herb. n. 925.)
61. *S. LINARIFOLIA*, caule herbaceo erecto ramosissimo parcè piloso, foliis alternis linearibus integerrimis vix uninerviis utrinquè pubescenti-subhirtellis, pedicellis elongatis 1-cephalis paucis ad apices ramorum, invol. squamis acutiusculis, pappo 12-14-seto corollæ puberulæ longitudinem subæquante. \mathcal{U} in Brasiliæ prov. Rio-Grande. Folia 12-15 lin. longa, sesquilineam lata. Cor. ex sicco albidæ. Achænium subscabridum. (v. s. in h. Mus. reg. Par. sub n. 852 à Mus. imp. Bras. acc.)
62. *S. MEGAPOTAMICA*, caule basi suffruticuloso ramoso, ramis sparse hispidulis teretibus, foliis oppositis linearibus glabriusculis hinc indè subserratis, pedicellis elongatis 1-cephalis in corymbum laxum dispositis, invol. 5-6-phyllis, squamis acutis glabriusculis, achænio glabro striato maturo invol. vix breviorè, pappo 20-seto scabro cot. æquali. \mathcal{J} \mathcal{U} in Brasiliæ prov. Rio-Grande. Herba 6-8 poll. longa. Folia pollicem longa, lineam lata. Cor. purpurascens. Pappus subflesceus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 928 miss.)
63. *S. OXYLANA*, caule imò basi suffruticuloso tereti glabriusculo, foliis oppositis oblongo-lanceolatis sessilibus utrinquè attenuatis dentatis hinc indè subtiùs et ad margines sparse setosis, capitulis ad apices ramorum paucis brevè pedicellatis, invol. squamis 5-6 lanceolatis acuminatis dorso setosis, achænio glabro, pappi setis circ. 20 scabris. \mathcal{J} \mathcal{U} in Brasiliæ prov. Rio-Grande. Herba 7-8 pollic. Cor. apice puberæ. Pappus flavescens. (v. s. in h. Mus. reg. Par. à Mus. Bras. sub n. 998 miss.)
64. *S. VIGINTISETA*, caule herbaceo glabriusculo erecto alternè paniculato-ramoso, ramis apice corymbiferis, foliis alternis brevissimè petiolatis lanceolatis

acuminatis subintegerrimis triplinerviis parvè subscabridis, capitulis breviter pedicellatis confertiusculè corymbosis, invol. squamis subacutis achænia vix superantibus, achæniis multicostatis piloso-hirtellis, pappi aristis 20-24 scaberrimis corollâ paulò brevioribus. — in Mexico legit cl. Karvinski. Cor. purpureæ glabræ. Pappus sordidè purpurascens. Styli rami valdè elongati ferè omninò epapuloi. (v. s. in h. Acad. reg. Monac.)

65. *S. RARUNCULOIDES*, caule herbaceo glabriusculo apice pubero erecto alternè paniculato, foliis alternis brevissimè petiolatis elongato-lanceolatis utrinquè attenuatis triplinerviis crenato-dentatis utrinquè scabridis, capitulis ad apices ramorum paucis pedicellatis, invol. squamis linearibus acuminatis glabris achænia superantibus, achænio angulato sulcato pilosiusculo, pappi aristis 25-30 scaberrimis achænio longioribus. $\frac{1}{2}$ in Mexico ad occid. urbis Guanaxuato legit cl. Mendez. Inv. purpurascens. Pappus flavescens. Capitula majora quàm in cæteris Steviis. (v. s. comm. à cl. inv.)

66. *S. CALYCINA*, caule suffruticoso erecto tereti basi glabro apice pulverulento-subvelutino, foliis linearibus utrinquè attenuatis integerrimis glabriusculis axillis foliiferis, corymbo laxo paniculato, invol. duplici, ext. squamis bracteivæ 5-6 linearibus, inter. squamis 5 lineari-lanceolatis margine scariosis pappum maturum subequante, achænio nitido glabro, pappi setis circ. 50 scabris. $\frac{1}{2}$ in Brasiliæ prov. Rio-Grande. Habitus ferè *St. polyphyllæ*, sed invol. achæn. et pappus valdè differunt. Fruticulus sesquipedalis. Folia pollicem longa, $\frac{1}{2}$ lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 905 miss.)

67. *S. MULTIARISTATA* (Spreng. syst. 3. p. 449), herbacea glanduloso-pubescent, foliis alternis linearibus canaliculatis subintegerrimis subitè canescentibus, pedunculis 1-cephalis subracemosis, pappo multiaristato corollam longè superante. — in Monte-Video.

† *Non satis nota.*

68. *S. URINCEFOLIA* (Thunb. ex florâ 1821. p. 331), herbacea, foliis ovatis acutis serratis capitulis paniculatis. — in Brasiliâ. Cæt. ex operè Thunb. mihi ignoto invitus omitto.

†† *Species exclusæ.*

S. CALLOSA Nutt. = *Florestina callosa*.

S. FEBRIFUGA Fl. mex. = *Piqueria trinervia*.

S. LINEARIS Cav. = *Palafoxia linearis*.

S. PEDATA Cav. = *Florestina pedata*.

LXXII. *PALAFOXIA* Lag. nov. gen. h. matr. (1815. et april. 1816) p. 26. *Sims bot. mag. t. 2132. Spreng. syst. 3. p. 449.* — *Paleolaria* Cass. bull. philom. decemb. 1816. mart. 1818. dict. suppl. 1. p. 59. 37. p. 256. Less. syn. 155. — *Polypteris* Nutt. gen. am. (1818). 2. p. 139. non Less. — *Stevia* et *Agerati* sp. auct.

Capitula pluriflora. Invol. oblongum campanulatumve vix subimbriatum, squamis 8-14 in fructu maturo stellato-patentibus. Recept. nudum. Achænia angulata, exter. squamis involuta. Styli rami elongati obtusi semiteretes dorso papillis elongatis subhispidi. Pappus paleis 8-12 scariosis 1-nerviis pinnato-striatis siccis 1-serialibus alternis brevioribus aut biseriatis exterioribus brevioribus constans. — Frutices aut herbæ Americanæ habitu Stevias referentes. Folia inf. opposita et sup. alterna indivisa. Capitula laxè corymbosa. Cor. albæ carneæ aut purpureæ. — Genus habitu valdè accedens ad *Florestinam Helianthis* adjunctam, et styl. habitu ferè *Vernoniaceas* referens.

1. *P. LINEARIS* (Lag. nov. gen. et sp. h. matr. 1815. n. 335), fruticosa cinereo-

- pubescens, foliis linearibus integerrimis obtusis, involacro oblongo 10-12-floro, pappi paleis 8 inæqualibus acuminatis longioribus achænio puberulo elongato brevioribus. ♀ in Mexico. Bot. mag. t. 2132. *Stevia linearis* Cav. l. præl. n. 464. *Ageratum lineare* Cav. icon. 3. p. 3. t. 205. *Paleolaria carnea* Cass. ll. cc. *Stevia linearis* et *lavandulifolia* Willd. Corollæ carneæ. Achænia exteriora interdum pappo minimo subabortivo tantum donata. (v. v. et s.)
2. *P. FASTIGIATA* (Less. syn. 156), caule herbaceo subscabro, ramis fastigiato-corymbosis, foliis lineari-lanceolatis integerrimis utrinque scabris, invol. squamis 12-15-biserialibus, ext. brevioribus, pappi paleis 12-14 lineari-lanceolatis, achæniis subscabri longitudine. 2♂ in Amer. bor. ad ripas flum. Altamshah. *Polypteris integrifolia* Nutt. gen. am. 2. p. 139. ex Less. Cor. tubus pubescens. Achænium subquadrangulare obconicum. Pappi paleis interdum lacera.
3. *P. TEXANA*, caule herbaceo puberulo, ramis glanduloso-puberis, foliis lanceolato-linearibus acuminatis trinerviis integerrimis puberulis, capitulis circ. 30-floris, invol. squamis 10-12 dorso glanduloso-puberis, pappo fl. ext. brevi vix acuto, fl. inter. aristato-acuminato achænio aequali. — in Mexici provinciâ Texas legit cl. Berlandier (pl. exs. n. 2014). Folia petiolo 4 lin., limbo bipoll. Cor. ex sicco incarnatæ. Achænia tetragona puberula basi attenuata. (v. s.)
4. *P. LATIFOLIA*, caule herbaceo erecto basi pubescente apice pubescenti-glanduloso, foliis oppositis petiolatis cordatis suborbiculatis grossè dentatis utrinque pubescenti-villosis basi trinerviis, paniculâ laxâ subnudâ, invol. squamis obovatis obtusis, pappi paleis 8 alternis elongatis acuminatis, alternis brevioribus obovatis. — in Mexico circa Tehuantepec mens. august. legit cl. G. Andrieux (pl. exs. mex. n. 286). Species distinctissima! Cor. purpureæ in invol. circ. 20. (v. s. comm. à cl. inv.)

• LXXXIII. *CARELIA* Less. syn. 156. non *Adans.* nec *Ponted.* nec *Rieux.*

Capitulum multiflorum. Recept. convexum pilosiusculum. Invol. campanulati squamæ ovali-oblongæ 2-3-seriales margine membranaceæ. Cor. tubulosæ 5-dentatæ. Achænia 5-gona. Pappus biserialis æqualis, paleis brevibus obtusis enerviis apice serrulatis. — Fruticulus *Brasiliensis* villosio-canescens. Folia opposita oblongo-ovata petiolata basi obtusa. Capitula dichotomè corymbosa.

1. *C. CISTIFOLIA* (Less. l. c.). ♀ in Brasiliâ (Less.), in prov. Rio-Grande. Hoc refero specimen descr. equidem brevissimæ Lessingii optimè conveniens, sed fortè specificè distinguendum. Petioli 4 lin., limbus fol. 15-16 lin. 5-6 lin. latus. Capitula pedicellata 50-flora. (v. s. in h. Mus. reg. Paris. à Mus. imp. Bras. sub n. 907 miss.)

LXXXIV. *AGRIANTHUS* Mart. l. in herb. Acad. Monac.

Capitulum circ. 20-florum. Invol. squamæ 1-2-seriales lineares acuminatæ subæquales. Recept. nudum. Cor. cylindraceæ obtusè 5-dentatæ. Styli rami longè exserti cylindrici obtusi. Achænia acutè pentagona secus angulos scabra, areolâ basilari prominulâ. Pappus corollâ brevior 1-serialis, squamellis lineari-lanceolatis acuminatis rigidulis inæqualibus densè barbellato-ciliatis. — Frutices *Brasilienses* ramosi humiles glabriusculi. Rami teretes. Folia conferta subimbricata alterna rigida lineari-lanceolata aut acerosa. Capitula in axillis supremis subsessilia aut breviter pedicellata in corymbum densum capitatum adgregata. Cor. ex sicco pallidè purpurascentes. — Genus ambiguum inter *Agerateas* et *Adenostyleas*.

1. *A. CAMPESTRIS* (Mart. l. c.), foliis lato-linearibus planiusculis minutissimè pubescentibus subtus reticulatis, pappo corollâ triplo fere breviori. ♀ in Brasiliæ prov. Minarum Generalium, in altis graminosis deserti inter Minas Novas et

flum. Sancti-Francisci, legit cl. Martius. Folia 10-12 lin. longa, $1\frac{1}{2}$ -2 lata. (v. s. in h. Acad. reg. Monac.)

2. *A. xiphioides* (Mart. l. c.), foliis acerosis subtriquetris glaberrimis, pappo corollâ paululum brevior. $\frac{5}{8}$ in Brasiliæ prov. Bahiensi interiori in montibus ad Rio de Contas Vales legit cl. Martius. Flor. octob. Rami divaricati breves. Folia 2-3 lin. longa, lineam lata. (v. s. in h. Acad. reg. Monac.)

Div. III. ADENOSTYLEÆ. — Eupatorieæ *Less. syn.* 156. — Eupatorieæ prototypæ et Adenostyleæ *Cass. tabl.*

Pappus pilosus barbellatus plumosusve uni- aut pluriserialis. — Adenostyleæ non divellendæ, nec ex recept. paleaceo aut nudo, nam Chromolœna et Decachæta ad Eupatorium nimis accedunt; nec ex pappo piloso aut plumoso, nam Liatrides plurimæ transitum exhibent.

LXXV. KUHNIA *Linn. sp. pl. ed. 2 (1763). p. 1662. Vent. h. cels. n. 91. Cass. dict. 24. p. 515. Less. syn. 157. Rafin. herb. p. 29. — Critonia Gærtn. fr. 2. p. 411. t. 174. non R. Brown. — Eupatorii sp. Ort.*

Capitulum multiflorum. Invol. squamis pauciserialibus. Recept. nudum. Cor. tubus ad faucem non ampliatus. Anthæræ interdum liberæ (nunc sæpius abortivæ). Styli rami exserti cylindracei obtusi. Achæ-nium elongatum sessile aut stipitatum. Pappus 1-serialis plumosus aut varius barbellatus. — Herbæ aut suffrutices Americani. Folia alterna 1-nervia rarius triplinervia linearia aut lanceolata. Capitula paniculato-corymbosa pedicellata. Cor. purpurascens aut albæ.

Sect. I. STRIGIA.

Achænia teretiuscula multistriata. Pappus plumosus. Invol. squamæ imbricatæ, exter. breviores.

1. *K. eupatorioides* (Linn. sp. 1662. Linn. f. decad. 2. p. 21. t. 11), caule herbaceo, foliis lato-lanceolatis serratis, corymbis paniculatis terminalibus. $\frac{2}{3}$ in Pennsylvania (Linn.), in Sibiria? (Ard. sed verisim. falso). Vent. h. Cels. n. 91. in adn. Ell. sketch 2. p. 291. Eupatorium alternifolium Ard. spec. 2. p. 40. t. 20. Flores albi aut albi. Folia ex Ell. subtus pubescentia glanduloso-punctata. Capitulum 10-florum.
2. *K. paniculata* (Cass. dict. 24. p. 516), caule herbaceo, foliis linearibus aut linearilanceolatis integerrimis junioribus margine revolutis, ramis apice corymbosis in paniculam laxam dispositis. $\frac{2}{3}$ in siccis Carolinæ (Ell. Bosc.), in montanis Pennsylvania, Virginie (Pursh) et Alabama (Torr.). Critonia Kuhnia Michx. fl. bor. am. 2. p. 101. Kuhnia Critonia Willd. sp. 3. p. 1773. Ell. sketch 2. p. 291. Flores albi. Folia subtus punctata. (v. s.)
3. *K. rosmarinifolia* (Vent. h. Cels. n. et t. 91), caule suffruticuloso, foliis linearilanceolatis integerrimis junioribus margine revolutis, pedunculis paucis subcorymbosis. $\frac{5}{8}$ in insulâ Cubâ. Eupatorium canescens Ort. dec. p. 34. *K. frutescens* Horn. h. hafn. 2. p. 791. Eupatorium Cubense Pers. ench. 2. p. 402. Flores purpurei. Folia subtus glabra punctata. (v. v. c. in h. Cels. et s. ex h. Madr.)

Sect. II. TRICHOGONIA.

Achænia pentagona, angulis piloso-scabris. Pappus plumosus. Invol. squamæ squales nec imbricatæ. — An genus proprium?

4. *K. arguta* (H. B. et Kunth nov. gen. am. 4. p. 105. t. 393), caule herbaceo subviscoso-hirto, foliis angustè lanceolatis argutè serratis tenuissimè hirtis, corymbis terminalibus. $\frac{2}{3}$ in montanis siccis prope Popayan Novo-Granat. Flores rosei.

5. *K. VILLOSA* (Spreng. syst. 3. p. 439), caule fruticuloso ramoso apice densè piloso-glanduloso, foliis linearibus integerrimis aut remotè denticulatis margine revolutis densè piloso-glandulosis, pedicellis paucis 1-cephalis in corymbum subdigestis. $\frac{5}{2}$ in Brasiliæ prov. Minarum General. ad Serra do Frio legit cl. Vauthier (pl. exs. n. 313). Folia secus ramos conferta, pollicem longa vix lineam lata. Invol. oligophyllum dorso pubescens. Corollæ subhispidulæ. (v. s. comm. à cl. Vauthier.)
6. *K. HIRTIFLORA*, fruticulosa, ramulis striatis hirtis, foliis brevissimè petiolatis ovatis subcordatis acutiusculis dentatis basi trinerviis suprâ sparsè puberulis subtus pubescenti-subhirsutis, corymbo terminali subsimplici oligocephalo conferto, capitulis breviter pedicellatis circ. 40-floris, invol. squamis 1-2-seriatis aequalibus oblongis dorso puberis obtusiusculis, cor. fauce dilatata extus hirtâ. $\frac{5}{2}$ in Brasiliæ prov. Minarum General. legit cl. Gaudichaud. Achænia longa 5-costata basi attenuata secus angulos hirta. Pappus plumosus. Folia 6 lin. longa 5 lin. lata. (v. s. in h. Mus. reg. Par.)
7. *K. GLABRATA*, caule erecto simplici basi suffruticoso glabro apice nudo ad summitatem pubescente, foliis glabris sessilibus obovato-cuneatis triplinerviis, corymbo composito, invol. squamis circ. biserialibus ovali-oblongis obtusissimis. $\frac{5}{2}$ in Brasiliæ prov. Rio-Grande. Habitus *K. glutinosæ*. Flor. in invol. 10. Achænia pentagona obpyramidata parcè villosa. Pappi setæ crassæ barbellatæ.
- a. foliis subintegerrimis. (v. s. in h. Mus. Par. à Mus. imp. Bras. sub n. 1055 miss.)
- b. foliis serratis. (v. s. in eod. herb. sub n. 784.)

Sect. III. LEOGONIA.

Achænia pentagona omninò lævia. Pappus barbellatus. Invol. squamæ paucæ subimbricatæ. — An potius *Eupatorii* species?

8. *K. PODOCARPA*, fruticulosa erecta ramosa, ramis teretibus minutissimè hirtellis, foliis alternis petiolatis ovatis acutis dentatis suprâ glabriusculis subtus petiolisque puberulo-velutinis, capitulis ad apices ramorum paucis pedicellatis, invol. squamis oblongis biserialibus, ext. foliaceis dorso puberulis, achæniis 5-gonis glabris longiusculè stipitatis. $\frac{5}{2}$ in Brasiliæ arenosis maritimis Restingas dictis. Fl. nov. et dec. Tota planta odorata (Lund in litt.). Folia fere formâ *Melissæ*. Cor. in sicco pallidæ. Habitu accedit ad sect. secundam, sed achænia lævia. (v. s.)
9. *K? GLUTINOSA* (Ell. sketch 2. p. 292), glutinoso-pubescent, caule herbaceo, foliis lanceolatis basi attenuatis triplinerviis integerrimis apice acuminatis rarer et grossè dentatis, capitulis corymboso-paniculatis. $\frac{4}{2}$ in pratis Alabamæ Amer. bor. Capitulum 8-10-florum. Cor. alba. Styli rami breviter exserti. Achænium acutè pentagonum, angulis lateribusque glaberrimis. Pappus albidus 1-serialis multisetus, setis scabris nec verè plumosis. (v. s. comm. à cl. Elliott.)
10. *K. BACCHAROIDES*, fruticosa glabra ad summitates viscosa, foliis obovatis basi cuneatis in petiolum attenuatis apice obtusis dentatis coriaceis reticulato-venosis glabris, corymbo composito nudo, invol. squamis linearibus paucis seriatis. $\frac{5}{2}$ in Brasiliâ circa Bahiam legit cl. Blanchet (pl. exs. n. 1826). Folia sesquipollic. longa 8-9 lin. lata. Cor. ex sicco pallidè flavæ. Achænium acutè pentagonum læve. Pappus uniserialis setis breviter et densè plumosis. (v. s.)

LXXVI. CLAVIGERA.

Capitulum 5-20-florum. Invol. pluriserialia imbricatum, squamis longitudinaliter striatis, ext. brevissimis intimis elongatis linearibus. Recept. angustum epaleaceum. Cor. tubulosæ cylindricæ brevissimè 5-dentatæ, dentibus extus glanduliferis. Anthæræ apicè appendiculatæ. Styli rami teretes nunc inclusi nunc parcè exserti. Achænium subcy-

lindraceum glabrum. Pappus 1-serialis, setis breviter barbellato-plumosis. — Suffrutices Mexicani erecti ramosissimi. Folia alterna integerrima aut dentata, linearia aut oblonga, uninervia impunctata. Capitula in paniculam spicæformem aut corymbosam disposita. Flores albidii. — Genus medium inter Kuhniam et Liatridem, ob specierum in Mexico omnium habitantium, dicavi cl. Franc. Xav. Clavijero qui historiam Mexici non tantum civilem, sed etiam naturalem scripsit. (1)

1. *C. CORYMBOSA*, glabra, caule basi suffruticoso, foliis linearibus integerrimis, capitulis circ. 20-floris corymbosis. $\frac{5}{2}$ in Mexici provinciâ Leonensi legit cl. Mendez. Invol. squamæ intense striatæ. (v. s.)
2. *C. SCOPARIA*, glabra, caule basi suffruticuloso, foliis linearibus integerrimis, capitulis circ. 10-floris in paniculam angustam spiciformem longè digestis. $\frac{2}{5}$ in Mexici prov. Leonensi legit cl. Mendez, in Cordillerâ Guchilapanâ Berlandier, et ad sinum Mulgrave ex herb. Hænke. Invol. squamæ striatæ. Rami paululum ad Liatridem mucronatam accedunt sed caract. diversissimi. (v. s.)
3. *C. DENTATA*, pubescens cinerea fruticosa, foliis ovali-oblongis dentatis hinc inde apice sublobatis, ramulis foliolis oligo- aut monocephalis in paniculam angustam dispositis, capitulis 12-floris, invol. squamis lineari-lanceolatis acuminatis ciliatis apice subscariosis, parce striatis. $\frac{5}{2}$ in Mexicanæ prov. Texas distr. (Commancheries) orientalibus necnon circa Bejar legit cl. Berlandier, pl. exs. n. 1897. 2079 et 2081! (v. s.)

LXXXVII. *LIATRIS* Schreb. gen. (1791). n. 1263. Willd. sp. 3. p. 1634. Nutt. gen. am. 2. p. 131. Less. syn. 157. non Don. — Anonymos Walt. fl. car. (1788). p. 196. — Suprago Gärtn. fr. (1791). 2. p. 402. — Liatris Suprago et Trilisa Cass. — Serratulæ sp. Linn. — Vernoniæ sp. Poir. — Kuhniz sp. Juss.

Capitulum multi-(5-30) florum. Invol. pauciseriale imbricatum. Recept. nudum. Cor. tubulosa ad faucem ampliata, lobis elongatis. Styli rami longè exserti cylindracei. Achænium circ. 10-costatum subcylindraceum. Pappus 1-3-serialis, setis plumosis barbellatisve. — Herbæ rarè frutices. Species omnes genuinæ ex Amer. bor. ortæ. Caulis sæpius simplices. Folia alterna integerrima aut in paucissimis dentata. Capitula spicata racemosa paniculata aut corymbosa. Cor. purpureæ roseæ rarius variantes albæ. — Interdum in plantis cultis vidi styli ramos in ligulas petaloideas eglandulosas mutatos. Radices tuberosæ resinosæ terebinthinam spirant undè à Canadensibus dicuntur *Pinetta de Prairie* et hinc inde ut diureticæ imò antisiphiliticæ admisse.

Sect. 1. *EULIATRIS*. — *Liatris* Cass. dict. 26. p. 235. non alior.

Pappus 1-serialis evidenter et longè plumosus. — Herbæ radice globoso-tuberosâ vix deorsum subfibrillosâ. Caulis simplices. Capitula spicata aut simpliciter racemosa.

1. *L. TENUIFOLIA* (Nutt. gen. am. 2. p. 131), radice tuberosâ, caule gracili glabro, foliis filiformi-linearibus longissimis infernè confertis summis brevibus setaceis infernè subpilosisculis, capitulis pedicello ramoso insidentibus in racemum longissimum digestis oblongis 5-floris, invol. squamis oblongis submucronatis, achænio villosa, pappo plumoso. $\frac{2}{5}$ in pinetis arenosis Carolinæ

(1) Opus anno 1780 italicè, et curante C. Cullenio anno 1787, anglicè editum. (Conf. Bibl. voy)

- utrinque. Ell. sketch 2: p. 275. *L. graminifolia* Willd. sp. 3. p. 1636? Folia ferè Pini palustris. Racemus 1-2-pedalis. Pedicelli circ. pollicares. Flores purpurei.
2. *L. mucronata*, radice....., caule simplici glabro, foliis angustè linearibus acutissimis calloso-mucronatis glaberrimis erectis punctatis, capitulis circ. 5-floris ad axillam bracteam sessilibus approximatis in spicam digestis, invol. squamis ciliatis obtusis abruptè mucronatis, achanio pubescente, pappo plumoso. $\frac{1}{2}$ in Mexicanæ provinciæ Texas distr. (Cammancheries) orientalibus legit cl. Berlandier. Fructif. mens. nov. et dec. Affinis videtur *L. tenuifolia* sed capitula sessilia. (v. s.)
3. *L. elegans* (Willd. sp. 3. p. 1635), radice tuberosâ, caule simplici pubescenti-villoso, foliis linearibus mucronatis scabrido-punctatis, capitulis breviter pedicellatis adproximatis 3-5-floris, racemo spiciformi elongato, invol. squamis acuminatis, interioribus coloratis, achanio villosa, pappi setis plumosâ. $\frac{1}{2}$ in Virginîæ (Ker), Carolinæ? (Walt. Fras. etc.), Floridæ (Michx.) et Georgiæ (Dob.) siccis. Bot. reg. t. 267. Ell. sketch 2. p. 279. *Stehelina elegans* Walt. fl. car. 202. *Serratula elegans* Ait. Kew. ed. 1. v. 3. p. 138. *Eupatorium speciosum* Vent. h. Cels. n. et t. 79. Folia sæpè falcata patentia seu reflexa. (v. s.)
4. *L. squarrosa* (Willd. sp. 3. p. 1634), radice tuberosâ, caule simplici pubescente, foliis longissimè linearibus nervosis margine scabriusculis, capitulis paucis in racemum digestis ad apicem pedicelli foliosi solitariis, invol. cylindrici squamis supernè foliaceis lanceolatis rigidis patentibus, achanio glabrescens, pappi setis plumosis. $\frac{1}{2}$ in arenosis campis et sylvis Canadæ (Hook. fl. bor. 1. p. 306), Virginîæ, Carolinæ et agri Kentuckensis. Pursh fl. 2. p. 509. Cass. dict. 26. p. 235. Sweet br. fl. gard. t. 44? malè. Ell. sketch 2. p. 282. *Serratula squarrosa* Linn. sp. 1147. *Pteronia caroliniana* Walt. fl. car. p. 292. ex Pursh. — Dill. elth. 83. t. 71. f. 82. Moris. ox. s. 3. t. 27. f. 1. Cór. latè purpurea, exter. demùm lateraliter rejectæ. Pappus 1-serialis rufo-purpureus. Ach. striatum. Var. foliis et invol. glabrescens aut pubescentibus. (v. s.)
- β. intermedia*, caule humiliore foliisque magis pilosis, invol. squamis ext. elongatis acuminatis erectis. — in Canada. *L. intermedia* bot. reg. t. 948. (v. s.)
5. *L. punctata* (Hook. fl. bor. am. 1. p. 306. t. 105), radice....., caule erecto simplici piloso, foliis linearibus marginatis glabris ciliatis utrinquè impresso-punctatis, spicis elongatis densis non rarè foliosis, invol. subcylindracei circ. 5-flori squamis imbricatis subequarosis, ext. rigidioribus, omnibus impresso-punctatis pungenti-acuminatis margine ciliato-lanatis, achanio piloso, pappo valdè plumoso. — in Amer. bor. occid. planitiebus ad Saskatchewan Red-deer et Eagl. hills. Cor. rosea.
6. *L. resinosa* (Nutt. gen. am. 2. p. 131), glabra, radice tuberosâ, caule erecto simplici aut apice vix ramoso, foliis linearibus rigidulis integerrimis utrinquè impresso-punctatis, spicis elongatis densis sæpius foliosis, invol. cylindracei 5-6-flori squamis imbricatis erectis longè acuminatis punctulatis, achanio villosa, pappo densè plumoso. — in Carolinæ utriusque pinetis (Nutt.), in territorio Arkansasano (Pourt.). Flores purpurei elegantes. (v. v. c. in h. Genev. ex sem. à cl. de Pourtales lectis.)

Sect. II. SUPRAGO. — *Suprago* Cass. dict. 51. p. 385. — *Supraginis* sp. Gært.

Pappus 1-3-serialis, setis densè et breviter barbellatis. — Herba, radice globoso-tuberosâ vix deorsum subfibrillosâ. Caules simplices. Capitula spicata aut racemosa. — Radix et inflorescentia sect. 1^{ma}. Pappus inter 1^{am} et 3^{am} medius.

7. *L. scariosa* (Willd. sp. 3. p. 1635), radice tuberosâ, caule simplici pubescenti-villoso, foliis lanceolatis pubescentibus margine scabris, infer. longissimis basi attenuatis, capitalis breviter pedicellatis racemoso-spicatis 15-30-floris, invol.

squamis obovatis obtusis margine scariosis, exter. patentibus acutis, achænio villosio, pappo subplumoso achænio subbreuiore. $\frac{1}{2}$ à Virginîâ ad Carolinam in montanis pratis (Pursh), in siccis (Ell. sketch 2. p. 280 cum var. plur.). bot. reg. t. 590. non Curt. Anonymos racemosus Walt. car. 198. ex Ell. *Liatris squarrosa* Michx. fl. bor. am. 2. p. 92. *Serratula scariosa* Linn. sp. 1147. excl. Pluk. syn. *Vernonia scariosa* Poir. dict. 8. p. 502. — Pluk. alm. t. 177. f. 4. Variat capitulis 15-20-25-36-floris, foliis glebruiculis pubescentibus aut ferè villosis, floribus involucri brevioribus aut longioribus, apicâ compactâ aut dissitiflorâ. (v. s. comm. à cl. Ell.)

8. *L. SPHÆROIDRA* (Michx. fl. bor. am. 2. p. 92), radice tuberosâ, caule pubescente, foliis lævibus infer. lato-lanceolatis super. lanceolato-linearibus, capitulis pedunculatis majusculis distantibus subglobosis 30-40-floris, invol. squamis latè ovalibus obtusis adpressis margine scariosis, achænio multicostato piloso scabro, pappi setis barbellatis achænio longioribus. $\frac{1}{2}$ in Carolinæ distr. superioribus. Ell. sketch 2. p. 281. Sw. br. fl. gard. t. 87. Suprago sphærocephala Cass. dict. 61. p. 386. *Liatris scariosa* bot. mag. t. 1709. excl. syn. (v. s.)

9. *L. SPICATA* (Willd. sp. 3. p. 1636), radice tuberosâ, caule simplici glabro, foliis linearibus glabris basi ciliatis nervosis et punctatis, capitulis longè apicatis ad axillam bractearum sessilibus 8-floris, invol. squamis oblongis adpressis obtusis sæpè mucronatis, achænio piloso, pappi setis densè barbellatis. $\frac{1}{2}$ in pratis et pinetis Lemmisi à Pensylvaniâ ad Georgiam. Sims bot. mag. t. 1411. *Serratula spicata* Linn. sp. 1147. And. rep. t. 401. Suprago spicata Gærtn. fr. 2. p. 402. t. 167. Cass. dict. 51. p. 385. *Liatris picta* Bart. ex Green. herb. L. gracilis Lodd. bot. cab. t. 1909. non Nutt. Anonymos ciliatus Walt. car. p. 197. An L. *Linaria* Raf. fl. lud. 61 satis differt? Flores purpurei. (v. s.)

β . *macrostachya*, apicâ densiore, capitulis brevissimè pedicellatis. L. macrostachya Michx. fl. 2. p. 91. ex Ell. sketch 2. p. 273. L. spicata Sweet br. fl. gard. 1. t. 49. L. pumila Lodd. cab. t. 147? — Dill. elth. t. 72. f. 83. (v. s.)

γ . *racemosa*, capitulis distinctè pedicellatis. *Liatris pilosa* bot. reg. t. 395. Lodd. bot. cab. 4. t. 356. non alior. An L. *turbinata* Sweet in Loud. hort. brit. 383. sine descr.? Differt à verâ L. pilosâ caule glabro nec pubescente, foliis ciliatis ut in var. α nec pilosis, floribus in invol. 8 nec 10-14. (v. s.)

10. *L. PYCNOSTACHYA* (Michx. fl. 2. p. 91), radice tuberosâ, caule simplici hirsuto, foliis strictis angusto-linearibus pubescentibus, capitulis in spicam longam digestis confertim sessilibus, invol. squamis adpressis supernè squarrosis. $\frac{1}{2}$ in pratis Illinoensibus (Michx.), montanis Virginie et Carolinæ (Pursh.) Huc cl. Pursh refert ic. Dill. t. 72. f. 83. à cæteris ad spicatam pertin. — Anonymos pilosus Walt. car. 197. Capitula minora quàm L. spicata. Cæt. ign.

11. *L. GRAMINIFOLIA* (Pursh fl. bor. am. 2. p. 508), radice tuberosâ, caule simplici glabro, foliis linearibus longissimis nervosis eciliatis margine scabrinis costâ mediâ hispidis, capitulis remotiusculis ad axillam bractearum sessilibus in spicam digestis, invol. squamis oblongis obtusis submucronatis ciliatis adpressis, inter. coloratis, achænio villosulo, pappi setis longiusculè barbellatis. $\frac{1}{2}$ in pinetis humidis Lemmisi Carolinæ Ell. sketch 2. p. 274. Anonymos graminifolius Walt. car. 197. — Pluk. alm. t. 424. f. 6? In plantâ cultâ capitula majora et magis adproximata. (v. v. c. et s. sp.)

12. *L. CYLINDRACEA* (Mich. fl. bor. am. 2. p. 93), radice tuberosâ, tota hirsuta, caule gracili, foliis gramineis, capitulis cylindraceis paucifloris sessilibus distanter spicatis, squamis apice rotundatis abruptè mucronatis. $\frac{1}{2}$ in sylvis pratisque Illinoensibus. Cor. lobi intus hirsuti (Nutt.). — Species homonyma à cl. Pursh in Carolinâ indicata et à cl. Elliot sketch 2. p. 276. descr. videtur diversa sed mihi non satis nota.

13. *L. ASPERA* (Michx. fl. bor. am. 2. p. 92), radice....., caule subramoso scabro-pubescente, foliis lineari-lanceolatis asperissimis, capitulis brevibus sessilibus apicatis distinctè alternis solitariis, invol. squamis rotundato-obtusis conniventibus. $\frac{1}{2}$ in pratis Illinoensium. Cæt. car. ign. — Anonymos ramosa

Walt. fl. car. 198 huc à cl. Pursh revocata videtur diversa et species in Carolinâ non indigena ex Ell. sketch 2. p. 276.

14. *L. HETEROPHYLLA* (Ait. kew. ed. 2. v. 4. p. 502), radice...., caule simplici glabro, foliis lanceolatis glabris lævibus, superioribus lineari-lanceolatis multoties minoribus, capitulis brevissimè pedunculatis spicato-racemosis, invol. subquattroai squamis lanceolatis acutis nudis. 2/ in Americâ bor. (Ait.), in Carolinâ australi et Georgiâ (Ell. sketch 2. p. 277). *L. varia* Banks ined. Cæst. ign.
15. *L. PILOSA* (Willd. sp. 3. p. 1636), radice...., caule simplici pubescente subpiloso, foliis linearibus pilosis ciliatis, capitulis longè pedicellatis in racemum laxum dispositis 10-14-floris, invol. squamis oblongis obtusis adpressis villosis, achenio multicostato puberulo, pappi setis densè barbellatis. 2/ in Amer. bor. à Georgiâ ad Nov. Cæsaream. Pursh fl. 2. p. 508. Ell. sketch 2. p. 277. Cor. lætè purpureæ. Species ab hortulanis sub hoc nomine admixta est *L. spicata* var. *γ*.
β. levicaulis, caule glabro, foliis basi remotè ciliatis, invol. squamis oblongis, capitalis 7-8-floris. 2/ in Novâ Cæsareâ (Torr.!) *L. pilosa* Torr. herb. 1835. *L. gracilis* Pursh fl. bor. am. 2. p. 508. Var. aut spec. recognoscenda (v. s.)
16. *L. GRACILIS* (Ell. sketch 2. p. 278), radice...., caule gracili pubescente subpiloso, foliis linearibus glabris basi subciliatis punctatis, capitulis longè pedicellatis laxè racemosis sub-7-floris, invol. squamis obovatis ciliatis adpressis, acheniis villosis, pappi setis barbellatis. 2/ in pinetis sicciæ Carolinæ (Ell.) et Georgiæ (h. Daby!). Cl. Nuttall ut mera var. *L. pilosæ* hanc fortè non immeritò habet. *L. gracilis* Pursh et Lodd. bot. cab. t. 1909. fortè diversa? (v. s.)
17. *L. SECUNDA* (Ell. sketch 2. p. 278), radice tuberosâ, caule reclinato pubescente, foliis linearibus glabris basi parcè ciliatis, capitulis breviter pedicellatis racemosis secundis, invol. squamis lanceolatis acuminatis adpressis dorso punctatis. 2/ in sicciæ et arenosis collibus Carolinæ, vulgaris prope Columbia (Ell.!). Cor. pallidè purpureæ in invol. 6-8. Achenium villosum. Pappi setes 2-3-seriales breviter et densè barbellatæ. (v. s.)

Sect. III. TRILISA. — *Trilisa* Cass. dict. 55. p. 310.

Pappi setæ 1-3-seriales breviter et distanter barbellatæ. — Herbæ aut suffrutescentes. Radices à trunco subtuberoso fibrosæ. Capitula in corymbum paniculamve disposita.

* *Herbaceæ.*

18. *L. PAUCIFLORA* (Pursh fl. bor. am. 2. p. 510), radice...., caule simplici glabro, foliis linearibus, paniculâ virgatâ foliosâ, ramis brevibus oligocephalis, capitulis subsessilibus secundis 3-5-floris, inv. squamis erectis lanceolatis acutis glabris. — in Georgiâ. Capitula magnit. *L. heterophyllæ*. Cæst. ign.
19. *L. ODORATISSIMA* (Willd. sp. 3. p. 1637), radice crassâ fibrosâ?, caule simplici folisque glaberrimis ovatis lanceolatisque nervosis subdenticulatis subglaucis, caulibus semi-amplexicaulibus, paniculâ polycephalâ subcorymbosâ, capitulis pedicellatis 7-8-floris, invol. squamis oblongo-obovatis obtusis, achenio multicostato vix puberulo, pappi setis parcè et breviter barbellatis. 2/ in paludosis apricis et pinetis demissis Virginie (Nutt.) Carolinæ et Floridæ. Michx. fl. 2. p. 93. Pursh fl. 2. p. 510. Nutt. gen. am. 2. p. 132. Andr. bot. rep. t. 633. *Trilisa odoratissima* Cass. dict. 55. p. 310. Anonymos odoratissimus Walt. car. 198. Cor. tubulosa, lobis brevibus. Pappus biserialis. Cor. lætè purpureæ. Folia trita vanillam ferè spirant. (v. s.)
20. *L. PANICULATA* (Willd. sp. 3. p. 1637), radice fibrosâ, caule simplici piloso-viscoso, foliis lanceolatis nervosis glabriusculis, paniculæ coarctatæ ramulis brevibus oligocephalis, capitulis sub-5-floris, invol. lanceolatis puberis erectis, achenio puberulo, pappi setis densè barbellatis. 2/ in pinetis Virginie (Pursh fl. 510), Carolinæ (Fras. Bosc.!), Floridæ (Pursh), Georgiæ (Daby!), Ell.

sketch 2. p. 283. *Anonymos paniculatus* Walt. car. 198. Variat cor. purpureis et albis. Capitula generis minora. Folia radic. multò majora 3-5-nervia basi attenuata. (v. s.)

21. *L. OPPOSITIFOLIA* (Nutt. in Sillim. journ. 5. 1822. p. 299), pubescens, caule ramoso, foliis lanceolatis suboppositis nervosis, ramulis paniculato-corymbosis, capitulis cylindraceo-ovatis 15-20-floris, invol. squamis oblongis obtusis, intercoloratis, pappi setis scabruiculis. ♀ in Louisiana circa Novam Aureliam. Capitula parva *L. paniculata* sed numerosiora.
22. *L. COMYMBOSEA* (Nutt. gen. am. 132), radice fibrosâ, caule solitario simplici hirsuto, foliis lanceolato-oblongis obtusis lævisculis, super. brevioribus, corymbo partim simplici circ. 20-cephalo, pedicellis hirsutis subundis 1-5-cephalis, capitulis 20-floris, invol. squamis oblongo-ovalibus obtusis lævibus. ♀ in Carolinæ bor. (Nutt.) et Georgiæ (Ell.) humidis ad margines paludum. *L. tomentosa* Michx. fl. bor. 2. p. 93 ? Ell. sketch am. 2. p. 284. Cor. pallide purpureæ.
23. *L. WALTERI* (Ell. sketch 2. p. 285), radice fibrosâ, caule simplici supernè piloso, foliis lanceolatis acutis glabris punctatis basi attenuatis, capitulis paucis corymbosis multifloris, invol. squamis acutis tomentosis, achænio parvè piloso, pappi setis subplumosis. ♀ in Carolinâ ad St. John's Berkeley (Ell.). *Anonymos uniflora* Walt. car. 198. Cor. intensè purpureæ.
24. *L. ABLUDIFOLIA* (Nutt. gen. am. 2. p. 133), demissa laxiuscula, radice tuberosâ fibrosâ, caulibus adgregatis, foliis radicalibus spatulato-lanceolatis obtusis 3-nerviis, caulinis sublinearibus, corymbi ramis subfastigiatis 1-5-cephalis, capitulis circ. 20-floris, invol. squamis oblongis obtusis, infer. subpatulis, pappi setis vix plumosis. ♀ in collibus arenosis Carolinæ borealis. Cor. purpureæ.

** *Fruticose.*

25. *L. FRUTICOSA* (Nutt. in Sillim. am. journ. 5. 1822. p. 299), glabra fruticosa, ramis corymbosis supernè nudis, foliis cuneato-obovatis punctatis, capitulis sub-5-floris, invol. squamis lineari-lanceolatis acuminatis. ♀ in Floridâ orientali. Drumetum ramosum. Folia ferè *Portulacæ* oleracæ. Invol. basi attenuatum atomis resinosis punctatdm. Cor. purpureæ. Pappus subconcolor.

† *Species exclusæ.*

L. BAICALENSIS Adams. = *Carphephorus Baicalensis*.

L. LATIFOLIA Don. = *Ainslæa pteropoda*.

LXXXVIII. *CARPHEPHORUS* Cass. bull: philom. 1816. dict. 7. p. 149. Less. syn. 158.

Capitulum multiflorum. Receptaculum paleaceum!, paleis inv. squamas referentibus corollâ brevioribus deciduis. Invol. squamæ 3-seriales imbricatæ adpressæ. Achæmium cylindraceum seu 10-costatum basi attenuatum. Pappus uni- aut pluriserialis, setis crassis plumosis aut grossè et irregulariter barbellatis. — Herbæ perennes, aut Sibiricæ aut originis dubiæ, habitu *Liatridis*. Caulis simplex. Folia alterna linearia aut lanceolata integra aut dentata. Capitula corymbosa aut racemosa. Cor. purpureæ. — Species recognoscendæ.

1. *C. PSEUDO-LIATRIS* (Cass. l. c.), caule pubescente, foliis linearibus acutis semi-amplexicaulibus integerrimis? adpressis, capitulis densè corymbosis, invol. squamis oblongis, achæniis 10-costatis hirtellis. ♀ patr. ign. Cor. et styli purpurei.
2. *C. BAICALENSIS*, caule infernè glabro supernè subvillosa, foliis lanceolatis in petiolum attenuatis dentatis, racemo terminali folioso, pedunculis 1-cephalis nudis, invol. squamis ovato-lanceolatis acuminatis. ♀ in regione alpinâ ad meridiem lacus Baikal. *Liatris Baicalensis* Adams act. soc. h. n. Mosc. 5. (1817).

p. 115. Cor. violaceæ. Pappus plumosus 20-setisq., setis basi subconcretis. Recept. paleaceum.

3. *C. REVOLUTIFOLIUS*, caule simplici suffruticoso basi glabro apice cum pedicellis puberulo, foliis linearibus glabresculis margine revolutis, ramulis axillaribus confertè foliosis, capitulis pedicellatis in racemum terminalem foliosum elongatam dispositis, achæniis teretibus puberulis. ♀ in Mexico legit cl. Hænke. Folia pollicem longa, explicata 2 lin. lata. (v. s. in h. Hænke ab ill. de Sternberg comm.)

LXXIX. DECACHÆTA.

Capitulum multi- (20-) florum. Invol. ovatum imbricatum, squamis 3-4-seriatis substriatis caducis ext. ovatis, inter. oblongis obtusiusculis. Recept. subconvexum, paleis linearibus caducissimis inter flores onustum. Cor. glabræ. Antheræ inclusæ. Styli rami longè exserti. Achænium acutè pentagonum glabrum, areolâ terminali convexâ. Pappus 1-serialis, setis 10 persistentibus basi vix scabris apice subclavellatis. — Suffruticulus erectus glaber Mexicanus. Rami teretes. Folia alterna subpetiolulata oblongo-lanceolata utrinquè acuminata serrata penninervia tenuiter reticulata. Thyrsus terminalis. Rami axillares apice capitula racemoso-congesta sessilia gerentes.

1. *D. HÆNKEANA*. ♀ in Mexico legit cl. Hænke. Folia 3-6 poll. longa, 10-12 lin. lata. (v. s. in h. Hænke ab ill. com. de Sternberg miss.)

LXXX. CHROMOLÆNA.

Capitulum multiflorum. Recept. paleis paucis linearibus apice coloratis inter flores hinc inde onustum. Invol. squamæ innumeræ erectæ sed laxiusculæ (Lupuli more) imbricatæ oblongæ obtusæ subscariosæ apice coloratæ, venis subelevatis parallelis substriatis. Cor. tubulosæ ad faucem non ampliata. Antheræ appendice ovatâ subpetaloideâ coloratâ donatæ. Styli rami crassi elongati glandulosi. Achænium 5-gonum basi attenuatum glaberrimum. Pappus 1-serialis setosus scaber. — Herba perennis Brasiliensis tota capitulis exceptis hirsuto-tomentosa. Radices fibrosæ. Caules teretes. Folia opposita lanceolata breviter petiolata basi integrâ à medio ad apicem dentato-serrata. Capitula cymosa, ad apicem caulium, medio subsessili precociori, ramis ex hujus basi ortis mono- aut oligocephalis. Invol. (et corollæ?) latè roseo-violaceæ, formâ Lupulina capitula, colore Salviæ Hormini bracteas, referentia. — Differt ab Eupatorio ut Carphæphorus à Liatride.

1. *C. NORMINOIDES*. ♀ in Brasiliæ prov. Minarum Generalium legit cl. Vauthier. Herba sequepedalis basi simplex ramis paucis oppositis. (v. s. comm. à cl. Vauthier sub n. 296.)

LXXXI. OOCLINIUM. — *Præxelis Cass. dict. 10. p. 261?* — Eupatoriæ sp. auct.

Capitulum multiflorum. Invol. squamæ biseriales aut sæpius pluries imbricatæ deciduæ. Recept. nudum post invol. collapsum ovatum! Achænium angulatum. Pappus 1-serialis setosus scaber. — Herba sæpius suffruticosa Americana. Folia opposita dentata aut integrâ. Pedunculi sæpius elongati 1-cephali. Cor. purpureæ. — Genus affine Eupatorio, sed receptaculo oviforme et etiam habitu diversum.

§. 1. *Species fruticosæ aut perennes, foliis integris aut dentatis, corollis æqualibus.*

1. *O. GRANDIFLORUM*, collo suffruticuloso, caulibus herbaceis teretibus sparse hispidulo-setosis, foliis oppositis sessilibus lanceolatis integerrimis acuminatis hispidulo-setosis, ramis elongatis subnudis 1-cephalis, capitulis oboeonicis circ. 40-floris, invol. squamis imbricatis acuminatis glabris. $\frac{5}{2}$ in Brasiliæ prov. Sancti-Pauli. Herba vix pedalis. Folia pollicem longa, 2-3 lin. lata. Capitula 6 lin. longa. Flosculi 5 lin. longi duplò nempe triplòve sequentibus majores, invol. æquales, ex sicco purpurei. Achænia exter. ad angulos scabra obcompressa. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 434 missa.)
2. *O. FAUCIDENTATUM*, collo suffruticoso, caule erecto basi sparse hispido apice glabro simplici aut trichotomo, ramis aphyllis apice 1-cephalis, foliis oppositis sessilibus lanceolatis ciliatis utrinquè glabris dentibus paucis acutis remotis subserratis, invol. ovali-subcylindracei glabri squamis adpressis ovalibus acutè mucronatis acuminatisve, exter. striatis, recept. ovato. $\frac{2}{2}$ in Peruvia legit olim cl. Hænke ex ill. de Bausa (sed in herb. Hænk. non reperire quivi). Medium inter grandiflorum et pedunculare. (v. s. in h. Acad. reg. Monac.)
3. *O. PEDUNCULARE*, suffruticulosum gracile diffusum adscendensve, caule tereti hispidulo, foliis oppositis linearibus utrinquè acutis subdentatis hispidulis, ramis elongatis subnudis apice 1-cephalis, capitulis 35-40-floris obovato-cylindraceis, invol. squamis imbricatis, ext. striatis acuminatissimis, inter. submembranaceis vix acutis, recept. ovato. $\frac{3}{2}$ in Brasiliæ prov. Minarum Generalium ad Sabara legit cl. Vauthier et hab. etiam in prov. Sancti-Pauli (h. Mus. Paris!). Folia 1-1 $\frac{1}{2}$ poll. longa, 1-2 lin. lata. Rami floridi 3-4 poll. Invol. apice purpurascens. Achænia compresso-tetragona hispidiuscula. Cor. et styli purpurei. (v. s.)
4. *O. SIDERITIS*, caule herbaceo adscendente tereti sparse hispido oppositè ramoso, foliis oppositis distantibus ovato-lanceolatis sparse hispidulis basi cuneatis integerrimis trinerviis apice acuminatis dentatis, ramis elongatis apice 1-3-cephalis nudis, capitulis obovato-cylindraceis circ. 30-35-floris, invol. squamis 2-seriatis erectis glabris, exter. acutis, int. obtusis apice subscariosis, recept. ovato. $\frac{2}{2}$? \odot ? circa Bahiam semel legit cl. Saltzman. Affine E. pedunculari. Habitus ferè Stachydis Sideritis. Radix fibrosa. Cor. ex sicco purpurascens. Capit. 4 lin. longa. (v. s. comm. à cl. Saltzman.)
5. *O. RIGIDUM*, suffruticulosum, caule erecto tereti striato glabro, foliis oppositis breviter petiolatis ovatis acuminatis remotè dentato-serratis trinerviis supra glabris subtus crebrè glanduloso-punctatis secus margines et nervos villosis, corymbo oligocephalo, capitulis longè pedunculatis 20-30-floris, invol. squamis acutè imbricatis deciduis striatis ext. obtusis, recept. ovato. $\frac{3}{2}$ in calcareis montibus Jamaicae borealis. Eupatorium rigidum Swartz! prod. III. flor. 1373. Caulis subherbaceus. Folia pollicem longa, 6-7 lin. lata. Flores cærulei. Odor plantæ suaveolens, sapor terebinthinaceus. (v. s. ex h. Swartz.)

§. 2. *Species herbacea annua, foliis tripartitis, coroll. marginalium lobis exter. majoribus.*

6. *O. CAPILLARE*, herbaceum erectum glabrum tenne, foliis oppositis usquè ad basin in segmenta 3 lineari-filiformia partitis et inde falso-verticillatis, pedunculis elongatis laxis 1-cephalis, invol. oblongi squamis acutis, corollis radii palmatifidis majoribus, achæniis angulatis glabris. $\textcircled{1}$ in Brasiliæ prov. Minarum Generalium ad Serra do Frio legit cl. Vauthier (pl. exa. n. 3111). Herba ferè Galii facie ramosissima gracillima. Cor. latè violaceæ. (v. s.)

† *Species dubia.*

7. *O. VILLOSUM*, herbaceum erectum, caule pilis articulatis sublanato, foliis oppositis petiolatis ovalibus serratis utrinquè petioloque villosissimis, capitulis

paucis subpaniculatis, invol. squamis glabris deciduis, recept. ovali seu conoideo, achæniis hispidulis. — in Guianâ Gallicâ legit cl. Poiteau. *Praxelis villosa* Cass. dict. 10. p. 262. An *Ooclinii*, an *Conoclinii* spec.?

LXXXII. CONOCLINIUM. — *Eupatorii* sp. auct.

Capitulum multiflorum. Invol. campanulati squamæ 2-3-seriales lineares acutæ subæquales. Recept. nudum conicum! Achænium angulatum. Pappus 1-serialis pilosus scaber. — Herbæ Americane aut suffrutices. Caules teretes striati. Folia opposita petiolata dentata. Corymbi conferti terminales. Cor. cœruleæ aut purpureæ: — Genus affine *Eupatorio*, sed recept. conico nec plano diversum sat videtur.

* *Species herbaceæ.*

1. *C. CŒLESTINUM*, herbaceum, caule tereti pubescente, foliis oppositis petiolatis ovatis basi truncatis aut subcordatis apice acutis obtusè dentatis triplinerviis subscabris, corymbis confertis, invol. squamis circ. 30, 2-3-serialibus ferè subulatis, floribus in invol. circ. 40, recept. conico, achænio angulato glabro. 2½ in pinguibus umbrosis montanis Virginie (Pursh), Carolinæ (Fras.! Boec.!), ad ripas fl. Ohio (Raf.!). *Eupatorium cœlestinum* Linn. sp. 1174. Ell. sketch 2. p. 306. Steud. nom. phan. 322. excl. Cass. syn. E. deltofolium Raf. in litt. 1831. — Dill. elth. 140. t. 114. f. 139. Pluk. t. 394. f. 1. Cor. lætè cœruleæ parvæ odoratæ. Styli lobi longè exserti. (v. s.)
2. *C. BETONICUM*, herbaceum, caule tereti striatulo glabro apice pubescenti-velutino, foliis oppositis sinu rotundato cordatis oblongis crenatis vix subacutis suprâ scabridis subtus petiolisque pubescenti-villosis, capitalis in corymbos terminales simplices confertis 60-70-floris, invol. squamis circ. 25 biserialibus linearibus acuminatis scabridis, recept. conico, achænio angulato glabro. 2½ in Mexico ad Tampico de Tamaulipas legit cl. Berlandier. Affine *E. cœlestino*. Cor. purpureæ. Folia exactè *Betonicæ*. (v. s.)
3. *C. BETONICÆFORME*, herbaceum?, caule tereti cum petiolis pediculisque densè breviterque velutino, foliis oppositis petiolatis sinu rotundato cordatis oblongis crenatis acutis suprâ pulverulento-scabris subtus velutinis, capitalis in corymbos terminales confertos digestis pedicellatis circ. 40-floris, invol. squamis circ. 16 biserialibus linearibus acutis dorso subpuberis, recept. conico, achænio angulato glabro globulos resinosos gerente. — in Brasiliâ circa Rio-Janeiro legit cl. Lund. *E. betonica* simillimum. Petioli 4 nec 6-8 lin. longi. Superficies tota magis velutina. (v. s. comm. à cl. Lund.)

** *Species basi suffruticosa.*

4. *C. FRAXIFOLIUM*, basi suffruticosum, caulibus erectis teretibus hispidis, foliis alternis rariis oppositis petiolatis ovatis basi cordatis truncatisve grossè crenato-dentatis suprâ glabriusculis subtus pallidis villosis, capitalis in corymbum terminalem densè congestis pedicellatis circ. 20-floris, invol. squamis 10-12 subbiserialibus ovali-oblongis mucronatis dorso puberis, recept. elevato subconico, achænio angulato pilosiusculo. 3 2½ in Brasiliâ circa Bahiam leger. cl. Blanchet, Salzmann et Lhotsky, et circa Caracas cl. Vargas. *E. nepetoides* Lindl. herb. Petioli 3-9 lin. longi; limbus fol. 9-18 lin. longus, 6-9 lin. latus. Folia subtus nunc ferè cana nunc glabriuscula, foliis *E. chamædrifolii* sat similia, sed fructif. valdè diversâ. Flores rosei odorati. (v. s.)
5. *C. MICROPHYLLUM* (Linn. suppl. 355 sub *Eupatorio*), suffruticosum adscendens, ramis teretibus subhirsutis, foliis oppositis petiolatis triangulari-ovatis obtusis crenatis margine revolutis suprâ glabris subtus albo-lanatis, corymbo terminali conglomerato, capitalis pedicellatis multifloris, invol. squamis subimbricatis lanceolatis ciliatis, recept. conico, achænio glabro. 3 in Novâ Granatâ. Crenæ fol. utrinquè 3-4. Petiolus ferè limbo æqualis nempe 3-4 lin. longus. Cor. purpureæ ex icone fortè 10-12, lobis puberis.

LXXXIII. HEBECLINIUM. — Eupatorii sp. auct. — Coleosanthi sp. Cass.

Capitulum multiflorum. Invol. campanulati squamæ pluriseriales laxè subimbricatæ sæpè in appendicem subcoloratam productæ. Recept. elevatum supernè plano-convexum pube brevi confertâ hirsutum et, ideo ferè piloso-frimbilliferum. Achænia angulata. Pappus 1-serialis pilosus scaber. — Herba australi-americanæ pubescentes. Caules teretes. Folia opposita petiolata cordata acuminata dentata. Corymbi terminales compositi conferti. Cor. albæ aut roseæ. — Genus affine Eupario, sed differt recept. villosa et invol. squamis sæpius appendiculatis.

* *Involucri squamis exappendiculatis.*

1. *H. MACROPHYLLUM*, basi suffrutescens, caule tereti pubescente, ramis patulis, foliis oppositis longè petiolatis lato-cordatis acuminatis crenato-serratis trinerviis reticulatis suprâ glabris subtus molliter pubescentibus, corymbo terminali composito polyccephalo, capitulis ad apices ramulorum confertis ovatis 30-floris, invol. squamis laxè imbricatis linearibus obtusis, achænio angulato glabro. $\frac{3}{4}$ in rudertis montium Jamaicæ (Sw.), Guadalupæ (Kraus!), Porto-Ricci (Wydler!), Cûbæ (Ossa), Guianæ (Aubl.), Brasilis (Lund!), agri Guyaquilensis (Hænke!), Peruvias (Poepp.), Novæ-Andalusis (H. B. K.). Eupatorium macrophyllum Linn. sp. 1175. Vahl symb. 3. p. 96. E. molle Swartz prod. xii. E. macrophyllum Sw. flor. 1303. E. molle et macrophyllum Willd. Ageratum guianense Aubl. guian. 800. Coleosanthus tiliaefolius Cass. dict. 24. p. 519. — Plum. et Burm. t. 129. Herba 4-5-pedalis. Flor. albi. (v. s.)

β. rivale, frutescens, caule tereti glabro ramis divaricatis, foliis oppositis longè petiolatis lato-cordatis acuminatis dentato-serratis trinerviis reticulatis subtus vix pubescentibus, corymbo terminali conferto parvo, capitulis pedicellatis circ. 20-floris, invol. squamis oblongis glabrisusculis ferè obtusis, achænio angulato glabro. $\frac{3}{4}$ in montosis juxta rivos Brasilis ad Rio-Janeiro legit cl. Lund. (v. s. in h. Mus. reg. Berol.)

* * *Involucri squamis omnibus aut plerisque in appendicem subcoloratam productis.*
— Urolepis.

2. *H. HECATANTHUM*, caule herbaceo erecto subsimplici tereti striato supernè præsertim cum petiolis et pediculis pube glandulosâ velutinis, foliis oppositis longè petiolatis cordatis brevè acuminatis acutè dentatis suprâ sparse subtus secus nervos puberulis, corymbo terminali composito conferto, capitulis ad apices ramulorum plurimis sessilibus circ. 100-floris, invol. squamis pluri-seriatis striatis, intimis productis in appendicem linearem villosam subcorollinam albidam, achænio glabro. ☉ in Brasilis prov. Rio-Grande. Herba pedalis sesquipedalive. Petioli ferè 2 poll.; limbus 2 $\frac{1}{2}$ poll. longus, 2 poll. latus. Cor. albæ. Recept. hemisphericum villosissimum. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 787 miss.)
3. *H. UROXANA*, caule herbaceo erecto tereti, petiolis pedunculisque pube brevi glandulosâ confertâ velutinis, foliis oppositis petiolatis cordatis acutis dentatis utrinquè glanduloso-puberulis, corymbo terminali oligocephalo, capitulis pedicellatis circ. 60-floris, invol. squamis 2-3-seriatis ferè omnino linearibus striatis puberulis productis in appendicem longam puberam roseo-coloratam, recept. hemispherico villosa, achænio glabro. — in Brasilis prov. Rio-Grande. Cor. roseæ. Valdè accedit ad *H. hecatanthum*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 807 et 803 miss.)

LXXXIV. CAMPULOCINIUM. — Eupatorii sp. auct.

Capitulum multiflorum. Invol. campanulati squamæ 2-3-seriales inæquales, ext. ovales, inter. angustiores. Recept. convexum hemisphæ-

ricum nudum. Achæniû angulatum. Pappus 1-serialis pilosus scaber. — Herbæ rarius suffrutescentes Americanæ pilosæ. Folia opposita petiolata dentata aut integerrima. Capitula pedicellata, in corymbum panicul. disposita, aut solitaria. Cor. purpurascens aut albidæ. — Genus Eupatorio affine, sed receptaculo hemisphærico et habitu quodam sat diversum videtur.

* *Ramis apice elongatis nudis 1-cephalis.*

2. *C. MACROCEPHALUM* (Less. in Linnæâ 1830. p. 136 sub Eupat.), caule herbaceo! erecto tereti hirsuto apice oligophyllo, foliis infer. oppositis? sup. alternis subsessilibus basi attenuato-cuneatis oblongis aut ovali-lanceolatis integris apice dentatis utrinquè hirtis, capitulis corymbosis paucis longè pedunculatis erectis circ. 100-floris campanulatis, invol. squamis 3-4-seriatis longit. subæqualibus, ext. ovalibus foliaceis hirtis, intimis linearibus scariosis serrulatis. 2? in Brasiliâ (Less.), in Bandâ orientali (Bacle!), prope Rio-Janeiro (Lund!) et Bahiam (Blanch!), et in Mexico prope Hacienda de la Laguna (Less.). Recept. convexum. Cor. glabra. Achænia 5-costata basi attenuata glabra. Pappus scaber. (v. s.)

β. *strigosum*, caule foliisque inferioribus multò magis strigosis, sup. linearibus, — in Brasiliæ prov. Rio-Grande. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 908. miss.)

2. *C. KLEINIOIDES* (H. B. et Kunth nov. gen. am. 4. p. 120. sub Eupatorio), herbaceum hispidum, ramis alternis teretibus striatis elongatis 1-cephalis, foliis oppositis brevissimè petiolatis lanceolatis acutiusculis obsolete serratis subtriplinerviis, capitulis solitariis longè pedunculatis multifloris, invol. campanulatis squamis subimbricatis glabriusculis, int. linearibus obtusis, ext. lanceolatis acuminatis brevioribus, recept. subgloboso, achænio hispido. ☉ in aridis montium Novæ Andalusie juxta Cocollar et Tomiriquiri. Føl. 6-9 lin. longa, 1-2 lata. Cor. dentes purpurascens. Habitus Porophylli olim Kleinie suffruticosi.

** *Capitulis corymboso-paniculatis.*

3. *C. PALUSTRE*, caule herbaceo erecto tereti subpuberulo apice nudo, foliis oppositis breviter petiolatis oblongis crenato-dentatis glabriusculis, capitulis 8-10 paniculato-corymbosis subglobosis, invol. squamis lineari-subulatis dorso-puberis, floribus circ. 70, recept. hemisphærico, achænio glabriusculo. 2 ♂ in palustribus prov. Sancti-Pauli Brasiliæ legit cl. Lund. Flos colore albo et violaceo variat. (v. s.)
4. *C. SUBPANICULATUM*, fruticosum, ramis teretibus apice summo puberulo-velutinis, foliis oppositis petiolatis elliptico-lanceolatis basi brevi cuneatis apice acuminatis subtriplinerviis integerrimis suprâ glabris subtus rufescenti-puberulis, paniculæ ramis oppositis oligocephalis, infer. folio suo brevioribus, capitulis pedicellatis 45-50-floris, invol. squamis ovali-oblongis 2-seriatis obtusiusculis dorso-puberis, int. paulò longioribus acutis, recept. convexo subalveolato, achænio glabro. ♂ in Brasiliæ prov. Sancti-Pauli. Petioli 5 lin. longi. Føl. limbus 2 poll. et paulò ultrà longi, 9-10 lin. lati. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 457. miss.)
5. *C. URTICÆFOLIUM* (Linn. f. suppl. 354. sub Eupatorio), caule herbaceo erecto tereti striato piloso, foliis oppositis petiolatis cordatis obtusiusculis irregulartè et grossè serratis trinerviis utrinquè piloso-hispidis paniculâ terminali subcymosâ, capitulis pedicellatis multifloris, invol. squamis 2-3-seriatis subpatulis lanceolatis apice subulato-mucronatis pubescentibus, cor. lobis villosis obtusis, recept. hemisphærico nudo, achænio glabro. 2? in Novâ Granatâ. Smith. id. pl. ined. 3. t. 68. Tratt. thes. t. 15. Petioli ferè longit. limbi; limbus basi subcuneatus, nervis 2 lateralibus margines constituentibus.

LXXXV. BULBOSTYLIS. — *Coleosanthi* sp. Cass. — *Eupatorii* sp. auct.

Capitulum 10-25-florum. Invol. squamæ lineari-lanceolatæ acuminatæ striatæ laxè imbricatæ 2-3-seriales. Recept. angustum epaleaceum. Cor. tubulosæ basi dilatatæ brevissimè 5-dentatæ. Stylus basi bulbosus. Achænium teres striatum sessile. Pappus 1-serialis setosus scaber. — Suffrutites Mexicani aut rariùs austr.-Americani. Rami teretes. Folia opposita aut superiora alterna, petiolata, lanceolata, aut ovata serrata, pubera aut glabra, subtùs sæpè glandulosa nunquàm pellucido-punctata. Panicula subthyrsoidea foliosa. Capitula oblonga.

1. *B. CAVANILLESII*, suffruticosa erecta, caule tereti villosa-velutino, foliis oppositis (summis alternis) petiolatis ovato-lanceolatis subacutis crenato-dentatis suprà sparsè scabridis subtùs trinerviis reticulatis glandulosis nervis petiolisque pubescentibus ramulis axillaribus sæpiùs 1-3-cephalis in paniculam elongatam foliosam dispositis, capitulis 20-25-floris, invol. squamis lanceolatis acuminatis striatis laxè imbricatis erectiusculis (non verè squarrosis), achænio tereti striato pubero. $\frac{5}{4}$ in Mexico ut vid. non infrequens et in Californiâ (Dougl.!). *Eupatorium axillare* fl. mex. ic. ined. *Eupatorium squarrosum* Cav. ic. 1. p. 66. t. 98. *Coleosanthus Cavanillesii* Cass. dict. 10. p. 37. Specim. habeo ex herb. Cavan. proveniens sed à descr. cl. auct. subrecedens foliis basi truncatis nec cordatis, floribus imò fructibus vix involuero longioribus, etc. Cor. ochroleucæ basi tumidæ apice brevissimè 5-dentatæ. Stylus basi dilatatus et hispidus. (v. s.)
2. *B. SECUNDIFLORA*, suffruticosa erecta, caule tereti pubescenti-viloso, foliis alternis et oppositis petiolatis subcordato-ovatis acuminatis dentatis trinerviis utrinquè parcè pubescenti-hirtellis subtùs glandulosis, ramis floridis racemosis in paniculam foliosam secundam dispositis, pedicellis glanduloso-villosis, capitulis nutantibus circ. 20-floris, invol. squamis imbricatis lanceolatis striatis acuminatis, achænio teretiusculo striato puberulo. $\frac{5}{4}$ in Mexico (Lag.) inter Tula et Tampico (Berlandier!) *Eup. nutans* H. B. et Kunth nov. gen. am. 4. p. 105? *Eup. secundiflora* Lag. nov. gen. 1814. p. 25. n. 334. Cor. ochroleucæ. Species media inter priorem et sequentem. (v. s.)
3. *B. PENDULA*, suffrutescens erecta, caule tereti glabriusculo, foliis oppositis (vix summis alternis) petiolatis ovato-lanceolatis basi cuneatis apice acuminatis dentatis triplinerviis tenniter pubescenti-hirtellis subtùs crebrè glandulosis, paniculâ racemosâ secundâ, pedicellis glanduloso-pilosis, capitulis pendulis 11-15-floris, invol. squamis imbricatis lanceolatis acuminatis striatis, achænio tereti striato puberulo. $\frac{5}{4}$ in Mexico. *Eupat. pendulum* Schrad! cat. sem. h. Gott. 1830. pl. rar. descr. et ic. ined. 1832. *E. scariosum* Wallr. in litt. 1834. Differt à duobus præced. florum in invol. numero minore, foliis tripli- nec trinerviis subtùs pilos pauciores glandulas numerosiores gerentibus et basi cuneatis. Cor. flavo-virides basi vix dilatatæ. An hæ tres species in unicam fortè colligendæ? (v. s. cult. h. Götting. et à cl. Schrad. comm. cum ic. exim.)
4. *B. REBERCARPA*, suffruticosa ramosissima, caulibus ramis pedunculis petiolisque pubescenti-hirtellis, foliis oppositis breviter petiolatis ovatis utrinquè subacutis dentato-serratis suprà scabridis subtùs trinerviis reticulatis villosis-hirtellis, ramulis axillaribus 1-cephalis, capitulis circ. 10-floris, invol. squamis laxè imbricatis subacutis striatis pubescentibus, ext. ovatis, inter. oblongo-linearibus; achænio teretiusculo striato pubescenti-viloso. $\frac{5}{4}$ in Mexico legit cl. Hænke. Cor. tubulosa brevissimè 5-dentata. Stylus basi bulbosus et villosus. Folia (incl. petiolo 2 lin.) 15 lin. longa, 6 lin. lata. (v. s. in h. Hænke. ex Mexico et comm. à cl. Née sine loci design.)
5. *B. PEDUNCULOSA*, fruticosa ramosissima subcæspitosa, ramulis teretibus pulverulento-velutinis apice longè, in pedicellum 1-cephalum dentatis, foliis oppositis (et summis alternis) breviter petiolatis ovatis dentibus paucis crenatis pube minimâ utrinquè cinereis subtùs glandulosis, capitulis circ. 25-floris, invol. squamis laxè imbricatis striatis mucronato-acuminatis, achænio tere-

tiusculo striato glabro. ζ in Mexico circa Villapando legit cl. Mendez. Suffrutex vix pedalis tortuosus. Cor. ex sicco albæ basi tumidulæ. Pedunculus a poll., quâ notâ faciliè ab *E. veronicæfolio* differt. Fol. cum petiolo 1 lin., 4-5 lineas longa. (v. s.)

6. *B. VERONICÆFOLIA*, fruticosa ramosissima, ramulis tenuissimè pulverulento-hirtis, foliis infer. ternis, cæteris oppositis, brevissimè petiolatis ovato-subrotundis obtusis subcordato-reniformibus crenatis tenuissimè pubescenti-canescenscentibus, capitulis ad apices ramorum solitariis breviter supra ultimum jugum exsertis circ. 15-20-floris, invol. squamis laxè imbricatis striatis mucronato-subacuminatis, achænio glabro. ζ in collibus Mexici prope sacellum Guadalupe. Eup. glechomæfolium Moc. fl. mex. ic. ined. Eupat. veronicæfolium H. B. et Kunth nov. gen. am. 4. p. 112. t. 441. Cor. alba. Suffrutex pedalis ad B. pedunculosam accedens, sed pedunculus vix invol. longitudine.
7. *B. ? NEPETÆFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 112), fruticosa, ramis sulcato-striatis hispido-hirtis, foliis alternis petiolatis triangulari-ovatis acutis subcordatis crenatis trinerviis rigidiusculis canescentibus supra scabris subtus tenuissimè tomentosis, capitulis axillaribus et terminalibus subspicatis subsecundis 20-25-floris, invol. squamis imbricatis striatis lineari-lanceolatis, ext. ovatis. ζ in exsiccatis montis Sanctæ Rosæ Mexici. Eupat. nepetæfolium H. B. et Kunth nov. gen. am. 4. p. 112. Cor. alba. Petiol. 2 lin., limbus pollicem longus, 8-9 lin. latus. An fortè *B. Cavanillesii* affine? sed de stylo et corollâ ailet cl. auct.
8. *B. OLIGANTHES*, herbacea erecta, foliis oppositis alternisve breviter petiolatis supra scabriusculis subtus pubescentibus, capitulis solitariis paucisve cynosis terminalibus multifloris, invol. turbinati squamis siccis multinerviis ext. subrotundis ciliatis, intus linearibus elongatis. — in collibus apricis Mexici prope Hacienda de la Laguna. Eupat. oliganthes Less. in linnæa 1830. p. 137. t. 2. f. 13. Cor. ut in *B. nepetæfoliâ* cylindricæ à typo generis recedunt, sed cor. basis non inflata ex icon. cit. Cæt. tamen ad *Bulbostylium* accedunt. Forma fol. ab auct. omisa.
9. *B. GLABRA*, herbacea glabra, ramis teretibus, foliis oppositis petiolatis ovali-lanceolatis acuminatissimis remotè serratis triplinerviis, paniculâ laxâ subcorymbosâ, capitulis longè pedicellatis erectis 16-floris, invol. squamis laxè imbricatis lanceolatis acuminatis striatis, ext. multò brevioribus, achænio angulato parçè puberulo. — Patr. ign. Petioli 8-9 lin. longi; fol. limbus 5-6 poll. longus, 1 $\frac{1}{2}$ -2 latus. Capit. 5-6 lin. longa. (v. s. comm. à cl. Gaudichaud.)
10. *B. SPINACLÆFOLIA*, fruticosa glabra, ramis teretibus, foliis oppositis longè petiolatis hastato-triangularibus lobis apiceque acuminatis ad margines grossè et acutè dentatis triplinerviis punctulatis, paniculis foliosis, pedunculis trifidis gracilibus folii longitudine, capitulis pedicellatis 10-12-floris, invol. squamis laxè imbricatis oblongis obtusis substriatis, achænio nervis exsertis 5-costato, nervis lateraliter hispidis. ζ in Mexico inter Victoria et Tula legit cl. Berlandier. Petioli 2 poll. longi. Limbus basi breviter cuneatus 3 poll. longus, 2 poll. latus. Cor. teres albidæ. (v. s.)
11. *B. ? PAUCIFLORA*, herbacea erecta hispido-pilosa, ramis teretiusculis, foliis oppositis petiolatis ovatis acutis serratis basi cuneatis et integerrimis reticulato-triplinerviis, capitulis axill. et termin. solitariis longè pedunculatis multifloris, invol. campanulatis squamis subimbricatis lanceolatis acuminatis striatis subæqualibus, achænio ad angulos scabro. \mathcal{Y} in Novâ Granatâ? Eupat. pauciflorum H. B. et Kunth nov. gen. am. 4. p. 120. Cor. violacæ? Recept. indescrib. Achænia compresso-quadrangularia ex cl. auct. Petioli 2 lin. longi. Limbus pollicaris, 7 lin. latus.

LXXXVI. CRITONIA *P. Brown. jam. (1756) p. 490, non Gærtn nec Cass. (1).*
 — *Dalea P. Brown. p. 314, non Linn., — Wikstrœmia Spreng. syst. 3. (1826) non Schrad. nec Endl. (2).* — *Eupatorii sp. auct.*

Capitulum 3-5-florum. Invol. oblongi squamæ 2-3-seriales imbricatæ oblongæ striatæ obtusæ. Recept. angustum nudum. Stylus basi non bulbosus. Achænia acutè pentagona basi attenuato-stipitata inter costas hispidula aut glandulifera. Pappus 1-serialis setosus scaber. — Frutices Caribæi et Mexicani glabri elegantes. Rami teretes striati. Folia opposita petiolata ovalia utrinquè acuminata subserrata membranacea punctis glanduloso-pellucidis ut in Samydâ sæpius oblongis perforata trita suaveolentia. Corymbi compositi. Capitula straminea ad apices ramulorum corymbi congesta. — Genus characteribus levibus sed habitu distinctissimo insignitum.

1. *C. DALEA*, fruticosa glaberrima, foliis oppositis petiolatis lanceolatis utrinquè sensim acuminatis subserratis penninerviis punctis liturisque pellucidis notatis, paniculâ subcorymbosâ, capitulis ad apices ramulorum ternis fasciculatis oblongis 4- (rariùs 3-) floris, invol. squamis 7-8 oblongis imbricatis substriatis, achænio acutè 5-gono hispidulo et glandulifero. $\frac{3}{4}$ in frutetis Jamaicæ (nec alibi ritè cognitum!) *Eupatorium* Dalea Linn. sp. 171. Swartz obs. 298. Jacq. ! h. Schœn. 2. p. 10. t. 146. *Isocarpha* Berterii Balb. ! herb. et ex eo *Calydermos Jamaicensis* Spreng. syst. 3. p. 457. excl. syn. — *Dalea* sen *Critonia* *P. Brown. jam. p. 314 et 490. t. 34. f. 1.* Odor totius fruticis gratissimus etiam in statu secco diù permanens. Petioli 4 lin. longi. Limbus fol. 3 poll. longus, pollicem latus. (v. s.)
2. *C. PSEUDO-DALEA*, fruticosa glaberrima, foliis petiolatis lanceolatis utrinquè sensim acuminatis serratis penninerviis punctis liturisque pellucidis notatis, paniculâ subcorymbosâ, capitulis ad apices ramorum quinis fasciculatis oblongis 5-floris, invol. squamis 7-8 oblongis imbricatis substriatis, achænio acutè pentagono hispidulo eglanduloso. $\frac{3}{4}$ in frutetis insulæ Cubæ circa Havanam. *Eup. Dalea* H. B. et Kunth nov. gen. am. 4. p. 106. excl. syn. Valde accedit ad *C. Daleam* sed præter notas ind. differt paniculâ magis confertâ. De odore fruticis silent autoptæ. Petiolus 3-4 lin. limbus 4 poll. longus, 12-15 lin. latus. (v. s. comm. à cl. Ramon de la Sagra.)
3. *C. MACROPODA*, fruticosa glaberrima, foliis oppositis longè petiolatis ovato-lanceolatis basi breviter attenuatis apice subabruptè et longè acuminatis penninerviis serratis punctis liturisque pellucidis notatis, paniculâ corymbosâ, capitulis ad apices ramulorum 3-5 breviter pedicellatis oblongis 5-floris, invol. squamis imbricatis oblongis acutiusculis, achænio acutè pentagono hispidulo eglanduloso. $\frac{3}{4}$ in insulâ Trinitatis Sieb. fl. trinit. exs. n. 184. Affine *C. Daleæ* et præsertim pseudo-daleæ, sed ab utraq. differt petiolo pollicari et ideo triplò longiori, foliis 5-7 poll. longis, 2 poll. latis magis ovatis et abruptius acuminatis, capitulis pedicellulatis, etc. (v. s. comm. à cl. Sieb.)
4. *C. PARVIFLORA*, fruticosum glabrum, ramulis nervisque fol. subtus subpubescentibus, foliis oppositis longè petiolatis ovato-lanceolatis acuminatis serratis penninerviis punctis liturisque notatis, paniculâ corymbosâ, capitulis ad apices ramulorum confertis sessilibus 3-floris, invol. squamis 6-8, inter. obtusis, achænio angulato. $\frac{3}{4}$ inter fruteta in montibus excelsis Jamaicæ australis. *Eupator. parviflorum* Swartz prod. p. 111. fl. ind. occ. 3. p. 1299. non Aubl.

(1) *Critonia* Gærtn. (1791) est *Kuhniz* sp. *Critonia* Cass. dict. v. 12 (1818) p. 1. est *Decaneuri* sp.

(2) *Wikstrœmia* Schrad. (1821) est genus *Tiliacæis* affine. Vid. prod. 1. p. 517. *Wikstrœmia* Endl. (1833) est genus *Thymelæarum*.

Sæpè cum *C. Dalea* mixta, sed præter notas ind. differt odore ferè nullo, corymbo magis divaricato, foliis non nitidis, etc.

5. *C. DALCROIDES*, fruticosa, ramulis pedunculis foliorumque petiolis nervisque utrinquè villis rufis velutino-subhirtis, foliis oppositis breviter petiolatis elliptico-oblongis utrinquè acuminatis penninerviis basi integris cæterum serratis punctis pellucidis crebris liturisque paucissimis pertusis, paniculâ divaricatâ, pedunculis apice capitula 3-5 glabra straminea 5-flora gerentibus, invol. oblongi squamis imbricatis 3-4 seriatis vix acutis. $\frac{5}{2}$ ad Tampico de Tamaulipas in Mexico legit cl. Berlandier. Affinis *C. Dalea* sed faciliè distincta. (v. s.)
6. *C. HÆBETRYA*, fruticosa, ramis subangulatis glabris, ramulis paniculâ velutino-villosis, foliis oppositis petiolatis ellipticis acuminatis minutè et rariter serrulatis penninerviis punctis pellucidis rotundis et paucis oblongis pertusis, paniculâ ramulosissimâ, capitulis ad apices ramulorum confertis breviter pedicellatis 5-floris, invol. oblongi squamis stramineis obtusiusculis striatis, achenio 5-costato inter costas hispidulo. $\frac{5}{2}$ in Mexico legit cl. Hænke. Petioli pollicares. Limbus fol. 5 poll. longus, 2-2 $\frac{1}{2}$ latus. (v. s. in h. Hænke. à cl. Sternb. comm.)

LXXXVII. EUPATORIUM *Tourn. inst. 455. t. 259. Linh. gen. n. 935. Gærtn. fr. 2. p. 401. t. 166. Adans. fam. 2. p. 123. Cass. dict. 16. p. 2. 25. p. 432. 26. p. 228. Less. syn. 157. excl. hinc inde sp.*

Capitulum multi (3-100)-florum. Recept. planum nudum. Invol. squamæ uni-bi- aut pluriseriales æquales aut inæquales laxè aut strictè imbricatæ. Cor. faux vix dilatata. Antheræ inclusæ. Styli rami exserti cylindracei obtusi. Achenium angulatum striatumve. Pappus 1-serialis pilosus scaber. — Herbarum aut suffrutices. Species paucæ gerontogæ, innumeræ, Americanæ. Folia sæpius opposita, rarè alterna aut verticillata. Capitula sæpius corymbosa aut paniculata. Cor. purpureæ cœruleæ, albæ aut albidæ uno verbo cyanicæ, nunquam flavæ seu xanthinæ. — Genus vastissimum, etiam post priorum et sequentis separationem species ferè innumerosas continens, fortè quoad specierum in sectionibus seriem recognoscendum.

Ser. I. IMBRICATA. *Nempè capitula squamis pluribus serialibus imbricatis, ext. brevibus, inter. sensum elongatis constantia.*

§. 1. CYLINDROCEPHALA.

Capitula cylindrica 5-60-flora. Invol. squamæ pluriseriales arcuè imbricatæ obtusæ intensè striatæ.

* Capitulis 20-60-floris, foliis oppositis.

1. *E. MULTIFLOSCULOSUM*, fruticosum, ramis teretibus hirsutis, foliis oppositis brevissimè petiolatis ovatis subcrenatis basi subtrinerviis subtus nervulosis utrinquè glabrinusculis subtus multiglandulosis, capitulis ad apices ramulorum intrâ duo ultima folia subsolitariis sessilibus ovato-cylindraceis 60-65-floris, invol. squamis arcuè imbricatis obtusis ciliatis substriatis. $\frac{5}{2}$ in Brasiliâ. Affine *E. squalido* sed faciliè ab eo et ab omnibus hujus sect. differt flosculis in invol. numerosissimis, capitulis solitariis, etc. (v. s.)
2. *E. MACRANTHUM* (Swartz fl. ind. oco. 1315), fruticosum glaberrimum, ramis angulato-sulcatis, foliis oppositis petiolatis ovatis acuminatis cartilagineo-serratis trinerviis subtus granulis pellucidis adspersis, corymbo terminali decomposito, pedicellis angulatis, capitulis cylindricis 30-40-floris, invol. squamis arcuè imbricatis obtusis. $\frac{5}{2}$ in Martinicâ. Spreng. syst. 3. p. 410. exclud. syn. Petioli semunciales. Capitula semipoll. longa. Cor. albidæ.

3. *E. SQUALIDUM*, fruticosum, ramis teretibus hirsutis, foliis oppositis breviter petiolatis ovato-rotundis crenato-dentatis trinerviis suprâ sparsè puberulis subtus nervoso-subreticulatis villosis-hirsutis, corymbis terminalibus oligocephalis suprâ folia vix exsertis, capitulis brevè pedicellatis cylindricis circ. 30-floris, invol. squamis arcè imbricatis obtusissimis dorso puberulis vix striatis. ♀ in Brasiliæ prov. Minarum Gener. ad Mariannam legit cl. Vauthier. Folia obscurè virentia vix cum petiolo lineam longo pollicaria. Cor. ex sicco purpureæ. (v. s. comm. à cl. Vauthier sub n. 279.)
4. *E. LIATRIDEUM*, caulibus basi suffruticulosus adscendenti-erectis teretibus simplicibus molliter villosis, foliis oppositis subpetiolulatis internodio brevioribus lanceolatis dentato-serratis basi trinerviis suprâ villosis-scabris subtus villosis subhirsutis, corymbis confertis oligocephalis, capitulis brevè pedicellatis circ. 30-floris, invol. cylindrici squamis circ. 30 imbricatis ovatis basi adpressis glabris striatis apice foliaceis subpatulis villosiusculis, achænio glabro. ♀ in Brasiliæ prov. Rio-Grande. Caules pedales aut paulò ultrâ. Folia pollicem longa, 5-6 lin. lata. Internodia 3 pollic. Capit. 4 lin. longa. Cor. purpureæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 756 miss.)

ℳ. *subcanum*, foliis petiolo magis distincto donatis utrinquè magis villosis-hirsutis, capituli paulò magis crassi squamis in appendicem foliaceam paulò longiorem productis. ♀ (v. s. cum α sub n. 751.)
5. *E. ORIGANOIDES* (H. B. et Kunth nov. gen. am. 4. p. 114), fruticosum, ramis teretibus glabris, ramulis angulato-striatis tenuissimè albidis-tomentosis, foliis oppositis petiolatis ovato-oblongis obtusis crenatis trinerviis suprâ rugosis tenuissimè hirtis subtus albo-tomentosis, corymbis terminalibus ramosis, capitulis pedicellatis multifloris, invol. oblongo-cylindracei squamis imbricatis trinerviis obtusis mucronulatis villosis-pubescentibus. ♀ in Novâ-Granatâ? Cor. alba. Petiol. 5 lin. Limbus pollicem longus, 6 lin. latus.
6. *E. STACHADIFOLIUM* (Linn. f. suppl. 355), fruticosum, ramis teretibus albolanatis, foliis oppositis petiolatis lanceolatis crenatis suprâ viridibus subtus cano-tomentosis, corymbis terminalibus conglomeratis, ramulis alternis albolanatis, capitulis multifloris, invol. squamis imbricatis acutis lanceolatis deorsum lanatis int. apice ciliatis, achænio glabro. ♀ in Novâ-Granatâ. Smith ic. pl. ined. 3. t. 69. H. B. et Kunth gen. nov. am. 4. p. 116. t. 343. An stirps à Kunthio descr. omninò eadem ac Linnæana?
7. *E. OASCURUM*, suffruticosum, caule tereti hispidulo, foliis oppositis subsessilibus linearibus trinerviis utrinquè attenuatis rariter serratis suprâ scabridis subtus secus nervos scabris inter nervos glanduloso-punctatis, corymbo composito trichotomo subpaniculato, pedunculis apice tricephalis, capitulis oblongo-cylindraceis circ. 30-floris, invol. squamis imbricatis striatis apice obtusis subfoliaceis inadpressis. ♀ in Brasiliæ prov. Rio-Grande. Folia sesquipollicem longa, ferè 2 lin. lata. Capit. 3 lin. longa, (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sine num. miss.)
8. *E. HYPERICIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 118), fruticosum, ramis sulcatis glabris, ramulis pubescenti-hirtis, foliis oppositis breviter petiolatis ovatis acutis integerrimis subtriplinerviis subrigidis suprâ glabrinusculis subtus molliter tomentoso-hirtis, corymbis trifidis oligocephalis, capitulis longè pedicellatis circ. 25-floris, invol. cylindracei squamis arcè imbricatis obtusis striatis glabris, achænio glabro. ♀ in Novâ-Granatâ? Cor. violacea. Petiol. sesquillin. Limbus fol. 9-10 lin. longus, 6 lin. latus.
9. *E. SERICEUM* (H. B. et Kunth nov. gen. am. 4. p. 110), caule herbaceo volubili?, ramis petiolisque albo-sericeis, foliis oppositis petiolatis ovatis acuminatis cordatis inæqualiter serratis suprâ hispido-scabris subtus sericeo-tomentosis, paniculæ terminales subcorymbosæ ramis subumbellatis, capitulis scasilibus ternis quaternisve circ. 20-floris, invol. oblongi squamis imbricatis striatis glabris, extèr. obtusis, inter. linearibus, achænio glabro. ♀ in Novâ-Granatâ. Cor. violacea? Petioli pollicares. Limbus fol. 4 pollic. longus, 2 ½ latus.

10. *E. FLORIBUNDUM* (H. B. et Kunth nov. gen. am. 4. p. 118. t. 344), fruticosum, ramis sexangularibus sulcato-striatis glabriusculis, foliis oppositis petiolatis ovatis angustato-acuminatis leviter serratis supra glabriusculis subtus molliter tomentoso-pubescentibus, corymbis ramosissimis diffusis, capitulis longè pedicellatis circ. 25-floris, invol. cylindracei squamis arcè imbricatis striatis obtusis glabris, achenio glabro. ζ in temperatis versus Loxam Peruvianorum. Cor. violaceæ. Petioli 3-6 lin. Limbus fol. 3 poll. longus, a $\frac{1}{2}$ poll. latus.
11. *E. SERRATULOIDES* (H. B. et Kunth nov. gen. am. 4. p. 117), caule herbaceo glabriusculo oppositè ramoso, ramis apice furcatis oligocephalis, foliis oppositis petiolatis ovatis acutis serratis triplinerviis supra scabriusculis subtus glabris margine ciliato-scabris, capitulis solitariis pedunculatis multifloris, invol. anbcylindracei squamis arcè imbricatis obtusis striatis ciliatis, achenio glabro. \mathcal{Z} in præruptis juxta Guancabamba Peruvianorum. Cor. violaceæ. Petioli 3 lin. Fol. limbus 13-14 lin. longus, 7 lin. latus.
12. *E. MAXIMILIANI* (Schrad. in litt. 1832), fruticosum, ramis virgatis teretibus apice tenuissimè pubescentibus, foliis oppositis petiolatis ovato-rhombeis acuminatis per totam ferè longitudinem grossè serratis trinerviis utrinquè glabrisculis, corymbo composito terminali, capitulis longè pedicellatis breviter cylindraceis 35-40-floris, invol. squamis arcè imbricatis obtusis. ζ in Brasilià circa Rio-Janeiro inter sepes primus detexit Princ. Maxim. Neovidenis. Subscandens. Rami oppositi patentes. Flores pallidissimè violacei aut nivei. Achenia glabra. (v. s. comm. à cl. Schrad. ex hort. Gett. et sp. à cl. Lund.)
 β . hispidulum, foliis utrinquè pilis sparsis hispidulis. ζ in Brasilià ad Rio-Janeiro et Bahià (Blanchet!), in Guianà Gallicà (Perr.). Flores in invol. 36 reperi. (v. s.)
13. *E. ODORATUM* (Linn. sp. 1174. amæn. 5. p. 405), fruticosum, ramis angulo recto divaricatis pubescenti-villosis, foliis oppositis petiolatis triangulari-ovatis apice integris acuminatis basi grossè dentato-serratis triplinerviis subtus densè pubescentibus punctatis corymbis ramos terminantibus, capitulis pedicellatis cylindricis 20-floris, invol. squamis arcè imbricatis plerisque acuminatis. ζ in ruderalis Jamaicæ. (Br. Linn. Sw.), Sancti Domingi (h. Lher.), Guadalupæ ubi vulgò dicitur *langue-de-chat* (Bad. ! Krauss !) Martinicæ (h. Duby!), Guianæ (h. Lindl.). Swartz obs. 300. — Pluk. alm. t. 177. f. 3? Flores albi sævè odorati! Floret decemb. Fortè à plurimis cum sequentibus confusum (v. s.)
 β . Cubense, invol. squamis striatis plerisque obtusis. — in insulà Cubà prope Havanam leg. cl. de la Ossa et R. de la Sagra. Fl. in invol. circ. 18. (v. s.)
 γ . *brachiatum*, invol. squamis plerisque acuminatis, foliis subtus parciis pubescentibus. ζ in Guadalupà (Badier) *E. atriplicifolium* Hort. Par. Lam. dict. 2: p. 407. excl. Plum. syn. *E. brachiatum* Wikstr. fl. guad. p. 71. (v. s. ex hort. Par. tempore Lam. lectum et sp. sicc. à cl. Badier comm.)
14. *E. COMPTROIDES* (Vahl symb. 3. p. 96), suffruticosum, ramis angulo ferè recto divaricatis pubescentibus, foliis oppositis petiolatis latè ovatis acuminatis basi grossè dentato-serratis triplinerviis apice integris supra subpuberulis subtus densè pubescentibus et punctulatis, corymbis terminalibus compositis, capitulis pedicellatis cylindricis circ. 20-floris. invol. squamis arcè imbricatis striatis obtusis. ζ in umbrosis Brasiliæ circ. Bahiam et in prov. Sancti-Pauli, in ins. Cubà, et in Mexico inter Santander et San. Fernando. Schrank pl. rar. h. mon. t. 85. Affine odorato sed certè dist. Cor. pallidè cœruleæ ex Berland. purpurascens ex Saltzm. sed cæt. nullum discrimen video inter specim. Mexicana et Brasiliensis. Florum vix fragrantium numerus in invol. variat à 15 ad 27. (v. s.)
15. *E. DIVERGENS* (Less. in linnæa 1830. p. 138), herbaceum, ramis divergentibus, foliis oppositis breviter petiolatis ovato-lanceolatis longè acuminatis triplinerviis subcrenato-serratis sparsè pilosiusculis, paniculis parvis oligoce-

- phalis fastigiatis, capitulis 20-25-floris, invol. cylindracei squamis erectis obtusissimis glabris striatis apice subciliatis. — in Mexico prope Hacienda de la Laguna. Cor. lilacinæ. Affine *E. conyzoidi*.
16. *E. CLEMATITIS*, fruticosum, ramis teretibus striatis glabriusculis, ramulis angulo recto divergentibus, foliis oppositis petiolatis triangularibus basi truncatis apice acuminatis subintegerrimis subtrinerviis glabris, corymbo paniculato laxo oligocephalo, capitulis longè pedicellatis cylindricis 25-floris, invol. squamis arcuè imbricatis obtusis. $\frac{5}{3}$ in Peruvia? legit cl. Pœppig. Petioli 8-10 lin. longi. Limbi vix prope basin 1-dentatis. Capit. semipollic. (v. s. comm. à cl. inv. sub n. 3108.)
17. *E. SMILACINUM* (H. B. et Kunth nov. gen. am. 4. p. 110), caule herbaceo volubili sulcato-angulato, ramulis piloso-hispidis, foliis oppositis petiolatis ovatis acuminatis grosse crenato-serratis trinerviis supra hispidulis subtus hispidis, corymbo terminali subtrifido divaricato, capitulis sessilibus congestis, invol. oblongi squamis imbricatis obtusis striatis ciliatis. $\frac{2}{4}$ in Andibus Quinduenais Novæ Gran. ? Flosc. et fruct. ign.
18. *E. INTEGRIFOLIUM* (Bert. ! herb. Spreng. syst. 3. p. 410), fruticosum, ramis petiolisque minutè pubescentibus, foliis oppositis petiolatis cordatis ovatis obtusis subintegerrimis supra glabris subtus glabresculis dense punctatis, corymbo terminali subsessili, capitulis cylindricis circ. 20-floris, invol. squamis arcuè imbricatis obtusis. $\frac{5}{3}$ in Guadalupâ legit cl. Bertero. Capit. 4 lin. longa. Petioli 2-4 lin. longi. Folia subtus pallidiora. (v. s. comm. à cl. amic. inv.)
19. *E. ATRIPPLICIFOLIUM* (Vahl symb. 3. p. 96. non Lam.), fruticosum, ramis petiolisque velutinis, foliis oppositis petiolatis triangulari-ovatis dentatis acutis supra glabris subtus glabresculis dense punctatis, corymbo terminali subsessili, capitulis cylindraceutis circ. 20-floris, invol. squamis arcuè imbricatis obtusis. $\frac{5}{3}$ in ins. Sanctæ Crucis. *E. triangulare* Poir. suppl. 2. p. 604. Fortè nimis ad *E. repandum* accedit? — Stirps Guadalupensis huc à Willd. relata videtur *E. odorati* aut punctati var. (v. s. sp. sine fl. cum herb. Vahl. coll. in h. Puer.)
20. *E. OSSÆANUM*, fruticosum glabrum, ramis teretibus vix cum foliis novellis subpuberulis, foliis oppositis petiolatis ovato-lanceolatis basi obtusis aut subattenuatis apice acuminatis grosse dentatis rigidis basi 3-5-nerviis supra lucidis subtus tenuiter multipunctatis, corymbo trichotomo, capitulis ad apices ramulorum 3 sessilibus cylindricis circ. 20-floris, invol. squamis arcuè imbricatis vix striatis obtusiusculis, achænio angulato glabro. $\frac{5}{3}$ in insulâ Cubâ legit cl. de la Ossa. Petioli 6-12 lin. Limbus fol. 4 poll. longus, ferè 2 poll. latus. (v. s. comm. à cl. inv.)
21. *E. ALTERNANS*, suffruticosum glabrum, caule tereti striatulo, ramis alternis, foliis lanceolatis subpetiolatis basi cuneatis apice acuminatis grosse serratis triplinerviis subreticulatis subtus punctatis, corymbo composito, ramis ramulisque alternis, capitulis pedicellatis 20-25-floris cylindraceutis, invol. squamis arcuè imbricatis striatis obtusis. $\frac{5}{3}$ in Brasiliæ prov. Sancti-Pauli. Valdè accedit ad *E. propinquum*. Folia 2 poll. longa, 9 lin. lata. Capitula 3 lin. longa. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 815 miss.)
22. *E. GLABERRIMUM*, fruticosum undiquè glabrum, ramis teretibus, foliis oppositis subsessilibus ovalibus acuminatis basi obtusis serratis penninerviis reticulato-nervulosis, corymbo composito fastigiato polycephalo, capitulis pedicellatis oblongo-cylindraceutis circ. 20-floris, invol. squamis adpressis tenuiter striatis. $\frac{5}{3}$ in Mexico legit cl. Hænke. Petioli semilineares primâ fronte nulli. Folia 6-7 poll. longa, 2 poll. lata. Capitula 3-4 lin. longa. Styli rami refracti unde in sicco primo intuitu non apparent. (v. s. in h. Hænke. à cl. Sternb. comm.)
23. *E. PSIADIEFOLIUM*, fruticosum glaberrimum subviscosum, caule tereti, ramis subhexagonis, foliis oppositis breviter petiolatis oblongo-lanceolatis utrinquè breviter acuminatis trinerviis basi integris apice grosse et pauci-serra-

tis, corymbo trichotomo laxo, capitulis pedicellatis cylindricis 15-18-floris. invol. squamis arcuè adpressis obtusis. ζ in Brasiliâ (h. L'Hér.), in siccis ad Cercovado (Lhostky!). (v. s.)

β . *latifolium*, ramis subhexagonis, foliis ovatis acuminatis. ζ in Cajennâ legit cl. Patris! (v. s.)

γ . *tereticaule*, ramis teretibus, foliis ovatis acuminatis grossius dentatis, capitulis gracilioribus. ζ in Mexico legit cl. Hænke! (v. s. in ejus herb. à cl. Sternb. comm.) An species propria?

**** Capitulis 10-20-floris, foliis oppositis.**

24. *E. MACRONON*, fruticosum glaberrimum, ramis teretibus, foliis oppositis petiolatis ovatis acuminatis grosse et obtusè cartilagineo-serratis subtrinerviis subtus eglandulosis, corymbo terminali decomposito, pedicellis compresso-angulatis, capitulis 16-floris, invol. squamis arcuè imbricatis obtusis. ζ ad vertex montium Dominicæ locis humidis. An var. *E. macranthi*?, sed differt præter notâ indicatas, petiolo 3 lin., capitulis 4 lin. longis. (v. s. ex h. L'Hér.)

25. *E. REPANDUM* (Willd. sp. 3. p. 1767. excl. Lam. syn.) fruticosum, ramis petiolis pedunculisque pubescenti-tomentosis, foliis oppositis petiolatis ovatis acuminatis sinuato-dentatis supra junioribus velutinis adultis glabris subtus velutino-tomentosis parçè glandulosis, corymbi ramis vix folia superantibus, capitulis pedicellatis cylindricis circ. 15-floris, invol. squamis arcuè adpressis obtusis. ζ in Sancto-Domingo (Plum. Lam.) et Cuba (R. de la Sagra!). Plum. amer. ed. Burm. t. 130. f. 1. (nec 128. f. 1 in quâ corollæ involucreo duplò longiores et folia obtusa). (v. s.)

26. *E. SINUATUM* (Lam. dict. 2. p. 407. ill. t. 672. f. 5), suffruticosum, ramis teretibus apice pubescentibus gracilibus, foliis oppositis petiolatis ovatis basi truncatis subtrinerviis apice subobtusis margine dentibus utrinquè 4-5 grossis obtusis lobato-crenatis supra scabridis subtus velutino-pubescentibus et glandulosis, pedunculis axillaribus divaricatis folio aequalibus et terminalibus parçè corymbosis, capitulis pedicellatis oblongis 8-10-floris, invol. squamis imbricatis ext. dorso puberis int. glabris substriatis, achænio glabro. ζ in Sancto-Domingo legit cl. Bertero. Petioli 2 lin. Limb. fol. pollicem longus, 8-9 lin. latus. Icon Plum. t. 128. f. 1. à Lamarckio citata verisim. excludenda. (v. s. comm. à cl. Bert.)

27. *E. GRACILIFLORUM*, fruticosum glabrum, ramis teretibus angulo recto divergentibus, foliis oppositis petiolatis ovato-lanceolatis acuminatis basi trinerviis dentatis apice integerrimis, corymbo oligocephalo laxo, capitulis pedicellatis teretibus gracilibus circ. 12-floris, invol. squamis arcuè imbricatis obtusis. ζ in Mexico leg. cl. Hænke, circa Acapulco (Karvinski). Affine *E. clematiti*. Petioli 5 lin. longi. Fol. limbi 2 poll. longi, 7-9 lin. lati. Capitula 4-5 lin. longa sesquilineam lata. Corymbi 5-15-cephali. (v. s. in h. Hænke à cl. Sternb. comm. et h. Acad. reg. Monac.)

28. *E. PUNCTATUM* (Lam. dict. 2. p. 408. non Willd.), fruticosum glaberrimum, ramis oblique divergentibus, foliis oppositis petiolatis ovatis utrinquè acuminatis grosse serratis subtus punctatis, corymbis ramos terminantibus, capitulis pedicellatis cylindricis 15-16-floris, invol. squamis arcuè imbricatis obtusis. ζ in ins. Caribæis (Sur. Vaill. ex Lam.), Guadalupâ (Bad.) Trinitatis (Sieb.) *E. odorato* affine Sieb.! fl. trin. exs. n. 75. — Syn. Pluk. cum signo? à Lam. citatum est certè exclud. Herba ex sicco videtur subviscosa. (v. s.)

29. *E. OXYLEPSIS*, fruticosum, ramis teretibus dense hirsuto-velutinis, foliis oppositis brevissimis petiolatis ellipticis subacutis integerrimis aut subnidentatis supra subscabridis subtus dense velutino-hirsutis, paniculæ oligocephalæ ramulis oppositis circ. 3-cephalis, capitulis cylindricis glabris 15-16-floris, invol. squamis arcuè imbricatis vix striatulis, ext. brevibus ovatis acutis, int. linearibus acuminatis. ζ in Brasiliæ prov. Sancti-Pauli. Fol. pollicem longa 6 lin. lata. Petioli 1-2 lin. Capitula 5 lin. longa ad *E. leptocephalum* accedentia. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 468.)

30. *E. SALICINUM* (Lam. dict. 2. p. 409), fruticosum, caule subangulato pubescente ramis teretibus substriatis glabriusculis, foliis oppositis subpetiolatis oblongo-lanceolatis acuminatis supernè rariter dentatis suprâ glabris et rugosis subtus cano-tomentosis, corymbo glomerato convexo, capitulis 10-12-floris, invol. ovalis squamis imbricatis pubescentibus, intimis longioribus obtusis. ♀ in Peruvîa ex h. Juss. ad radices montis ignivomi Tunguraguz Quitensium. Spreng. syst. 3. p. 412. excl. syn. Folia 6-7 poll. longa.
31. *E. PROPINQUUM*, fruticosum glabrum, caule tereti, ramis oppositis, foliis oppositis subpetiolatis lato-lanceolatis basi cuneato-attenuatis apice acuminatis grossè serratis triplinerviis subreticulatis subtus subpunctatis, corymbo composito trichotomo polyccephalo, capitulis pedicellatis cylindricis 16-floris, invol. squamis arcè adpressis striatis obtusis. ♀ in Brasiliæ prov. Rio-Grande. Folia (cum petiolo 2-3 lin.) 3 poll. longa, ferè pollicem lata. Capitula 4 lin. lata. Accedit ad *E. ivæfolium* et *E. alternans*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 1064.)
32. *E. IVÆFOLIUM* (Linn. amœn. ac. 5. p. 405. sp. 1175), herbaceum, caule tereti hispidulo, foliis oppositis angusto-lanceolatis utriusque attenuatis subpetiolatis trinerviis subserratis glabris, corymbo trichotomo laxo, capitulis oblongis pedicellatis circ. 15-20-floris, invol. squamis paucis erectis striatis obtusis. ♀ in cultis Jamaicae (Swartz! obs. 301), Cuzcæ (Spr.) et in Mexici ditione Oaxaca (Andr.! pl. exs. n. 283). (v. s.)
 β. foliis medio serratis, capitulis 20-25-floris. — in Guadalupe (Bert!). (v. s.)
33. *E. CHAMÆDRYFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 113), fruticosum, ramis teretibus pubescenti-velutinis, foliis oppositis breviter petiolatis ovatis obtusiusculis crenatis trinerviis suprâ scabris subtus pubescenti-hirtis subtomentosis, junioribus canescentibus, corymbis compositis confertis breviter ramosis, capitulis sessilibus 10-12-floris subcylindraceis, invol. squamis adpressis obtusis substriatis dorso subpuberis, inf. ciliatis. ♀ in Andibus Peruvianorum frequens (H. et B.), in vallibus (Hænke!). Cor. violaceæ. (v. s. in h. Hænke à cl. Sternb. comm.)
34. *E. FURCATUM* (Lam. dict. 2. p. 407), caule dichotomo glabro tereti, foliis oppositis, infer. ovatis, sup. lineari-lanceolatis, angustis integerrimis internodio longioribus, paniculis terminalibus, capitulis 10-12 floris, invol. ovali-oblongi imbricati squamis striatis obtusis, achenio angulato. — in ips. Caribæis ex h. Juss.
35. *E. RHINANTHACRUM*, suffruticosum glabrum, caulibus à collo plurimis teretibus simplicibus, foliis oppositis (supremis subalternis) subsessilibus lanceolatis basi attenuatis trinerviis parè dentatis impunctatis, corymbo pedunculato subgloboso conferto, capitulis oblongis pedicellatis 10-12-floris, invol. squamis imbricatis oblongo-linearibus substriatis acutiusculis. ♀ in Brasiliæ provinciâ Sancti-Pauli. Caulis pedalis. Folia sesquipollicem longa, 6 lin. lata. Accedit ad *E. ivæfolium* sed caule glabro, corymbo coarctato et invol. squamis acutis facillè differt. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 500 miss.)
36. *E. VERBENACEUM*, suffruticosum, caule tereti aparsè hispido, foliis oppositis petiolatis lanceolatis basi cuneatis apice acuminatis grossè dentatis margine scabris secus nervum subtus hispidis cæterum glabriusculis, corymbo fastigiato, ramis oppositis, capitulis longè pedicellatis cylindricis 10-12-floris, invol. squamis arcè imbricatis striatis glabris obtusis. ♀ in Brasiliæ prov. Rio-Grande. Folia internodiis dimidio ferè breviora, caulina bipollicaria, ramea pollicaria, Verbenæ officinalis folia referentia. Corymbi rami glabri. Capit. 4 lin. longa. Cor. ex sicco purpureæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 760 miss.)
37. *E. TOZZIÆFOLIUM*, caulibus ad collum suffruticulosos simplicibus teretibus apice præsertim puberulo-velutinis, foliis oppositis brevissimè petiolatis ovatis obtusè dentatis trinerviis suprâ pubescentibus subtus subtomentosis, summis etiam dentatis, corymbi ramis oppositis folio longioribus, capitulis ad apices

- ramulorum subsessilibus confertis 10-floris, invol. squamis imbricatis 4-5-seriatis striatis obtusis, int. apice villosiusculis. $\frac{5}{2}$ in Brasiliæ prov. Sancti-Pauli. Valde accedit ad *E. bartisiæ* folium. Achenia 5-gona ad angulos subvillosa. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 504 miss.)
38. *E. bartisiæ* folium, suffruticosum, caule simplici tereti supernè pubescente, foliis confertis, infer. oppositis subsessilibus ovato-lanceolatis, sup. alternis exactè sessilibus angustius lanceolatis basi tantum dentatis apice acuminatis integris, summis integerrimis angustis, sessilibus basi trinerviis grossè incisedentatis utrinquè pubescenti-scabris subtus glandulosis, corymbo composito fastigiato, ramis alternis erectis, capitulis pedicellatis cylindricis 11-floris, invol. squamis imbricatis striatis obtusis dorso puberulo et glanduloso, intimis mucrone subpatulo brevi apiculatis. $\frac{5}{2}$ in Brasiliæ prov. Rio-Grande (h. Mus. imp. bras. n. 822.) et in Bandâ orientali (Bacle!). Folia pollicem longa. Rami floridi foliis multo longiores. Capit. 4 lin. Cor. et styli purpurei (v. s.)
39. *E. phlogifolium*, fruticosum, caule tereti glabro, ramis hispidis, foliis oppositis sessilibus angusto-lanceolatis acuminatis hinc inde suberratis glabris subtus punctulatis, corymbo composito fastigiato polycephalo, capitulis oblongis 5-floris. invol. squamis 9-10 glabris striatis obtusis. $\frac{5}{2}$ in Brasiliæ prov. Rio-Grande. Habitus ferè *Eup. ivæ* folii sed corymbo multo magis amplo, ramis magis hispidis et capitulis 5-floris faciliè distinctum: (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 799 miss.)
40. *E. serrulatum*, fruticosum, ramis teretibus apice sex-sulcatis puberulo-subhispidis, foliis oppositis breviter petiolatis linguato-lanceolatis utrinquè subattenuatis regulariter serrulatis penninerviis utrinquè sparè glandulosis subtus secus nervos puberulis, petiolo ciliato, corymbo oligocephalo, capitulis oblongis pedicellatis 11-floris, invol. squamis imbricatis glabris oblongis striatis obtusiusculis. $\frac{5}{2}$ in Brasiliæ prov. Rio-Grande. Folia (incl. petiolo 2-3 lin.) 2 $\frac{1}{2}$ poll. longa, 4-6 lin. lata. Capit. 3 lin. longa pallida. Cor. ex sicco ochroleucæ. Affinis *E. comptoniæ* folio. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 790 miss.)
41. *E. tachyophorum*, canibus herbaceis erectis simplicibus setoso-hispidis, foliis oppositis et sup. alternis petiolo brevissimo hispidissimo donatis lanceolatis basi subtruncatis apice acuminatis grossè et obtusè dentatis utrinquè hispidis, corymbo terminali composito conferto polycephalo, capitulis breviter pedicellatis 10-floris, invol. squamis circ. 3-seriatis oblongis basi glabris striatis apice submembranaceis obtusis villosis. — in Brasiliæ prov. Sancti-Pauli. Setæ articulatæ. Folia sesquipollicem longa, 7 lin. lata: Achenia glabriuscula. Affine *E. bartisiæ* folio. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 497 miss.)
- *** Capitulis 10-15-floris, foliis alternis.
42. *E. fulgellum* (H. B. et Kunth nov. gen. am. 4. p. 119. t. 345), fruticosum, ramis teretibus pubescentibus, foliis alternis oblongo-lanceolatis utrinquè attenuatis dentatis suprà scabriusculis subtus tenuissimè pubescentibus penninerviis, corymbo composito terminali conferto polycephalo, capitulis 8-13-floris cylindricis pedicellatis, invol. squamis arcè imbricatis obtusis ciliatis. $\frac{5}{2}$ in convalli Mexicanâ prope lacus Texcenses. Cor. supernè violaceæ. Styli longè exserti. (v. s. comm. à cl. Alaman, Mendez, Mairret.)
- **** Capitulis 5-9-floris, foliis oppositis.
43. *E. punctulatum*, fruticosum, caule tereti glabriusculo, foliis oppositis petiolatis ellipticis basi attenuatis apice acuminatis rariter dentatis aut integerrimis triplinerviis glabriusculis subtus punctulatis, corymbo fastigiato composito, pediculis pubescentibus apice capitula 3-5 sessilia fasciculata cylindrica 8-9-flora gerentibus, invol. squamis arcè imbricatis obtusis substriatis. $\frac{5}{2}$ in Amer. calid. sed locus proprius ignotus. Valde accedit ad *E. punctatum*. Petioli 6-8 lin. longi. Folia 4 poll. longa, 12-14 lin. lata. Capit. 4 lin. longa. Achenia glabra. (v. s. in h. Martius.)

44. *E. SCABRUM* (Linn. f. suppl. 354), suffruticosum? erectum, ramis striatis villosis-scabris, foliis oppositis breviter petiolatis ovalis acuminatis subserratis variis integris subtrinerviis rugosis subtus hirtis supra scaberrimis, paniculis coarctatis terminalibus bracteatis, capitulis pedicellatis cylindricis 7-8-floris, invol. squamis acutè imbricatis glabris oblongis obtusis trinerviis apice callosis, achen. glabro. ζ in Novâ-Granatâ (Mut. in Linn.) juxta Santa-Fe (H. B. et Kunth). Smith pl. ined. 3. t. 67. *E. scabrum* H. B. et Kunth gen. nov. an. 4. p. 119 à Mutisio ut specim. Linn. communic. ab icone et descr. Smith. multum recedit ex cl. auct. Planta nigrescens. Flores ex Sm. flavescences, potius verisim. albid.
45. *E. LEPTOCEPHALUM*, fruticosum glabrinsculum, ramis teretiusculis, foliis oppositis breviter petiolatis ovali-lanceolatis acuminatis serratis triplinerviis subtus crebrè glanduloso-punctatis, corymbo composito fastigiato polycephalo, pedicellis 3-cephalis, capitulis cylindricis 7-8-floris gracilibus, invol. squamis acutè adpressis substriatis ext. obtusiusculis intimis subacutis. ζ in Andibus Peruvianorum legit indef. Hænke. Corymbus ferè Odontolomæ. Rami subnubentes. Capitula glabra 4 lin. longa. Petioli sesquilineares. Fol. 2 poll. longa, 7-9 lin. lata. (v. s. in h. Hænke à cl. Sternb. comm.)
46. *E. HEXANTHUM*, suffruticosum, caule simplici tereti pubescente subhirtello, foliis sessilibus infer. oppositis cordatis breviter acuminatis grossè dentato-incisis subtus reticulato-nervosis supra scabris subtus secus nervos puberis, super. sensim alternis vix cordatis longius acuminatis subintegris, corymbi compositi ramis alternis paulò folia superantibus capitula 3-7 fasciculata cylindracea 6-flora gerentibus, invol. squamis imbricatis dorso puberulis et subglandulosis, int. acutè mucronatis. ζ in Brasiliæ prov. Rio-Grande. Pedale. Affine *E. hartszefolium*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 829. miss.)
47. *E. GAUDICHAUDIANUM*, caule herbaceo? erecto tereti densè et sordidè cum petioli pedicellisq. velutino-hirtis, foliis oppositis petiolatis ellipticis obtusis basi subcuneatis crenato-dentatis penninerviis supra sparse scabridis, subtus pubescenti-hirtellis, corymbo terminali composito polycephalo densè conferto, capitulis congestis subsessilibus 5-floris, invol. ovalis squamis 10 biseriatis, ext. ferè subulatis, inter. ovali-oblongis obtusis apice ciliatis, achenio angulato glabro. \mathcal{L} ? circa Rio-Janeiro Brasiliæ in apricis legit cl. Gaudichaud. Petioli 2-3 lin. longi. Limb. fol. 15-16 lin. longus, 7-8 latus. Folia suprema sæpè alterna. (v. s. comm. à cl. invent.)
48. *E. INTERMEDIUM*, fruticosum, ramis teretibus apice præsertim velutino-hirsutis, foliis oppositis basi brevissimè cuneatis et petiolulatis elongato-lanceolatis serratis penninerviis supra puberis subtus velutino-hirsutis, corymbi compositi terminalis ramis oppositis polycephalis, capitulis ad apices ramulorum confertis subsessilibus 5-floris, invol. squamis laxè imbricatis, ext. minimis acutis, inter. oblongis basi striatis glabris apice submembranaceis obtusis ciliatis dorso villosulis, achenio glabro. ζ in Brasiliæ prov. Rio-Grande. Exactè medium inter *E. Gaudichaudianum* et *E. comptoniaefolium*. Folia 2-2 $\frac{1}{2}$ poll. longa, 6-9 lin. lata. (v. s. in h. Mus. Par. à Mus. imp. Bras. sub n. 844 miss.)
49. *E. COMPTONIAEFOLIUM*, fruticosum, caule tereti glabro, ramulis brevissimè velutinis, foliis oppositis brevissimè petiolatis lineari-sublanceolatis acutis regulariter serratis 1-nerviis supra glabris subtus brevissimè velutinis, corymbo composito fastigiato denso, capitulis oblongis 4-5-floris vix pedicellatis, invol. squamis 8-9 imbricatis subacutis striatulis int. apice ad dorsum tomentosis. ζ in Brasiliæ prov. Rio-Grande. Cor. ex sicco subochroleuce breves. Achenia glabra. Fql. 1 $\frac{1}{2}$ -2 poll. longa, 3-4 lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 76 miss.)
50. *E. ASCLETADEUM*, suffruticosum, caule tereti apice villosa-hispido, foliis oppositis sessilibus aut summis alternis cordatis lanceolatis acuminatis integerrimis glabris subtus crebrè punctatis, corymbi compositi ramis oppositis villosa-hispidis, ramulis apice capitula 3-5 subsessilia oblonga 4-5-flora gerentibus,

invol. squamis imbricatis striatis achæniisque glabris. $\frac{5}{2}$ in Brasiliæ prov. Rio-Grande. Folia conferta more Gentianæ asclepiadæ decussata sesquipollicem longa 5 lin. lata. Invol. 2-3 lin. longa pallida subpurpurascens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 473, 774 et 84a.)

51. *E. CUNEATUM*, fruticosum erectum ramosum glaberrimum viscosum, ramis teretibus striatis, foliis breviter petiolatis oppositis obovatis basi cuneatis et plurinerviis apice obtusis grosse dentatis coriaceis, corymbo composito fastigiato, invol. squamis 3-4 seriatis lanceolatis acutis ciliolatis pallidis, flor. 5, achæniis obovato-teretiusculis minutissimè scabridis. $\frac{5}{2}$ in Brasiliæ prov. Sancti-Pauli. Folia sesquipollicem longa, pollicem lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 503 miss.)

52. *E. POLYSTACHYUM* (DC. non Willd.), fruticosum glaberrimum, ramis teretiusculis striatis, foliis oppositis petiolatis ellipticis acutis serratis trinerviis subtus punctatis suprâ glutine viscoso flinitis, paniculæ oblongæ ramis subracemosis, capitulis oblongis 5-floris pedicellatis, invol. squamis 8-10 imbricatis obtusis substriatis. $\frac{5}{2}$ in Brasiliæ prov. Sancti-Pauli. Petioli 3 lin. longi. Folia sesquipollicem longa, 8-9 lin. lata. Capit. 2-3 lin. longa. Achænia glabra angulata. Styli longè exserti. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 488 miss.)

ℓ. angustius, foliis longius petiolatis oblongis suprâ vix viscosis. $\frac{5}{2}$ in prov. Rio-Grande (h. Mus. imp. Bras. n. 976!). Petioli 6 lin. longi. Fol. limb. 2 poll. longus, 8-9 lin. latus. (v. s. in h. Mus. reg. Par.)

53. *E. XYPHYLOIDES*, fruticosum glaberrimum, foliis oppositis longissimis sublinearibus utrinquè attenuatis subpetiolatis denticulatis coriaceis reticulato-venosis, paniculæ subcorymbosæ ramis oppositis, capitulis oblongis pedicellatis 4-floris, invol. squamis 8-10 imbricatis brevissimis subrotundis int. longioribus obtusiusculis substriatis. $\frac{5}{2}$ in Brasiliæ prov. Rio-Grande et Sancti-Pauli (h. Mus. imp. Bras. n. 997 et 513), prov. Minarum General. (Gaud. in h. Mus. Par.), circa Tejuco (Vauthier! pl. exs. n. 277). Species distinctissima, foliis coriaceis subfalcatis Xylophyllas angustifolias subreferens. Folia 4-7 poll. longa, 1 $\frac{1}{2}$ -3 lin. lata. Capit. vix 3 lin. longa. Achænia angulata glabra. Receptac. elevatum minimum. (v. s.)

54. *E. PINNATIFIDUM*, fruticosum erectum ramosum glabrum, ramulis angulatis, adultis teretibus, foliis oppositis profundè pinnatifidis, lobis 1-2-jugis linearilanceolatis integerrimis, capitulis racemosis subpaniculatis 5-floris, invol. squamis 1-3-seriatis imbricatis oblongis acutis scarioso-rufis, achænio glabro. $\frac{5}{2}$ in Americâ austr. legit cl. Née, sed locus propr. mihi ignotus. Affine *E. xylophylloidi* ex inflorescentiâ, sed foliis faciliè diversum (v. s.)

***** Capitulis 5-floris, foliis alternis.

55. *E. RUPELURI-FOLIUM*, fruticosum, caule tereti glabro, paniculæ ramis puberulis, foliis alternis sessilibus lato-linearibus acuminatis integerrimis trinerviis nervis lateralibus tenuissimis subtus minutè glandulosi utrinquè subviscosi, paniculâ corymbosâ compositâ polycephalâ, capitalis oblongis 5-floris brevè pedicellatis, invol. squamis viscosis glabris imbricatis obtusis striatis, pappo involucri non superante. $\frac{5}{2}$ in Brasiliæ prov. Sancti-Pauli. Affine *E. linifolio* sed ex omni parte majus. Folia 4 poll. longa, 3-4 lin. lata. Corymbus major. Invol. fusco-purpurascens. Folia summa basi dilatata more Bupleuri. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 464 et 465 miss.)

56. *E. LINIFOLIUM*, suffruticosum, caule tereti breviter pubescenti-hispidulo, foliis alternis subsessilibus erectis linearibus acuminatis integerrimis 1-nerviis margine subrevolutis subtus crebrè glandulosi, corymbo composito fastigiato, capitalis oblongis brevè pedicellatis 5-floris, invol. squamis erectis, ext. brevibus puberis ciliatis, int. longioribus obtusiusculis pallidis striis tribus notatis, pappo involucri longiore. $\frac{5}{2}$ in Brasiliâ ad Sanctam-Catharinam legit cl. Gaudichaud. Folia 2 poll. longa, 2 lin. lata. Cor. ex sicco purpurascens. Styli rami longissimi. (v. s. comm. à cl. invent.)

57. *E. ericoides*, fruticulosum erectum, caule tereti infernè glabro superne pulverulento-velutino, foliis alternis sessilibus linearibus basi subattenuatis integerrimis apice callosis et obtusis erectis confertis, corymbo composito terminali fastigiato, capitulis 5-floris pedicellatis, invol. squamis circ. 10 dorso velutinis margine ad apicem ciliatis demum patulis. $\frac{5}{8}$ in Bras. prov. Rio-Grande. Specimen quoad flores manentem sed evident. *E. ericoidi* affine et foliis facile distinctum. Folia 6-7 lin. longa, lineam lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 834 miss.)
58. *E. halimifolium*, fruticulosum erectum, caule tereti glabro, ramis ad apicem velutino-canis, foliis alternis cuneatis obtusis basi attenuatis integerrimis erectis confertis utrinque pube sericeâ glandulis intermixtâ cano-sericeis, corymbo composito terminali, capitulis pedicellatis 5-floris, invol. squamis laxè imbricatis dorso subvillosis ext. acutis int. obtusis ciliatis, acheniis 5-gonis glabris. $\frac{5}{8}$ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 766) et in prov. Minarum Generalium in monte Serra do Caraca (Mart.!) *Vernonia halimifolia* Mart. herb. Invol. subpurpurea. Cor. purpurea. Pappus sordidus. Folia 6-9 lin. longa, $1\frac{1}{2}$ lin. lata. (v. s. in h. Mus. reg. Par. et Acad. reg. Monac.)
59. *E. ericoides*, fruticulosum glabrum erectum, caulis teretibus, foliis alternis sessilibus lineari-subulatis acutis erectis confertis integerrimis rigidulis, corymbis terminalibus oligocephalis, capitulis pedicellatis 5-floris, invol. squamis 10 laxè imbricatis, ext. parvis acutis, int. longioribus ovalibus obtusis margine membranaceis ciliatis flosculis dimidio ferè brevioribus, achenio pentagono glabro. $\frac{5}{8}$ in Brasiliæ prov. Rio-Grande. Fruticulus pedalis. Folia 3-4 lin. longa. Styli exserti obtusi. Cor. ex sicco albidæ. Pappus achenio longior, setis rigidis basi ferè in anulum concretis crebre barbellatis. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 843 et 775 miss.)
60. *E. stramineum*, caule suffruticoso erecto ramosissimo tereti, ramulis foliisque minutè puberulis, foliis alternis linearibus sublanco-latis unijnerviis integerrimis, capitulis 5-floris ad axillas supremas breviter pedicellatis in racemose foliosos subdispositis, invol. squamis imbricatis lanceolatis acuminatis stramineis glabris striatis, achenio 5-gono ad angulos subscabro. $\frac{5}{8}$ in Peruvia legit cl. Hænke. (v. s. in h. Hænke. à cl. de Sternberg. comm.)

§. 2. *Pteropoda*.

Capitula 8-10 flora. Invol. oblongi squamæ imbricatæ oblongæ pallidæ. — Caules herbacei. Folia in petiolum alatum angustata glabra impunctata.

61. *E. quadrangulare*, herbaceum? undiquè glabrum, caule acutè tetragono, foliis sessilibus subconnatis lanceolato-ovatis apice acuminatis basi attenuatis penninerviis serratis membranaceis, paniculæ amplæ ramis oppositis striato-angulatis corymboso-paniculatis polycephalis, capitulis oblongis 8-10-floris: invol. squamis adpressis oblongis vix obtusis striatis. — in collibus Mexici ad Tantoyuca legit cl. Berlandier (n. 2151), ad Acapulco (seu fortè ad montes Oro-noenses) legit cl. Hænke. Folia 6 poll. longa, 2 poll. lata. Panicula pedalis. Capit. 3 lin. longa pallida. Achenia angulata puberula. (v. s. in h. Hænke.)
62. *E. aromatisans*, herbaceum? glaberrimum, caulibus striato-sulcatis subangulatis, foliis oppositis latè ovatis acuminatis basi longè cuneato-angustatis in parte petioliiformi subintegris cæterum grossè serratis, paniculæ amplæ ramis oppositis polycephalis, capitulis oblongis 10-12-floris pedicellatis, invol. squamis striatulis subciliatis obtusiusculis adpressis. — in insulâ Cubâ juxta Havannam, ubi ex cl. Ramon de la Sagra vulgò adhibetur ad odore aromatico impregnandas præparationes tabacinas vulgò *agarrés* dictas. Affine *E. quadrangulare*! Fol. pars petioliiformis 3 poll. longa, limbus 6-7 poll. longus, 5 poll. latus. Styli rami longissimè exserti! (v. s. comm. à cl. de la Sagra.)
63. *E. thyrsoides* (Moc. fl. mex. ic. ined.), herbaceum? glaberrimum, caulibus teretibus striatis, foliis oppositis cordatis acuminatis serratis, petiolis latè marginalis integerrimis basi subdilatis et ferè connatis, paniculæ amplæ

thyrsoides ramis oppositis, capitulis pedicellatis oblongis 8-9-floris, invol. squamis imbricatis acutis. — in Mexico. Affine *E. aromatisanti*, sed folia cordata serraturis minoribus donata et rigidiora, petioli subconnati basi latiores nec angustiores et capitula pauciflora. (v. ic. pict. et spec. sicc. in h. Hænk. à cl. Sternberg comm.)

β. *puberum*, foliis secus nervos velutino-puberis, canle apice subpuhero: — in Mexico legit cl. Hænke. Stirps coll. Andrieux exs. mex. n. 282. foliis subtus ubique sed tenuissimè pubescentibus huc fortè etiam referenda? (v. s. in ejus herb. à cl. Sternb. comm.)

§. 3. *Sessilifolia*.

Capitula 5-12-flora. Invol. oblongi squamæ oblongæ imbricatæ. — Folia opposita sessilia. Caules herbacei.

64. *E. sessilifolium* (Linn. sp. 1172), canle subtereti glabrinusculo, foliis sessilibus semimplexicaulibus distinctis ovato-lanceolatis acuminatis basi rotundatis serratis glabris, corymbo composito, invol. squamis paucis oblongo-linearibus obtusis imbricatis, floribus in invol. 5-12. ♀ in montanis rupestribus præc. calcareis in Alleghany (Pursh) occidentalibus (Michx.), in sylvis ad Roxburg prope Bostoniam (Big. fl. ed. 2. p. 293). Flores albi. Affine *E. perfoliato* sed glaber et folia non connata (v. v. c. et sp.)

65. *E. truncatum* (Willd. sp. 3. p. 1751), canle tereti striato villosulo-hispidulo, foliis oppositis amplexicaulibus distinctis basi obtusis lanceolatis acuminatis rugosis dentato-serratis subtus pubescenti-villosis, corymbo composito conferto, capitulis 5-10-floris, invol. squamis 12-15 imbricatis linearibus obtusis. ♀ in Marylandiâ (Sims) Pennsylvania (Willd.). Enp. salviaefolium Sims bot. mag. 1. 2010. Flores albi. Omnino medium inter *E. perfoliatum* et *E. sessilifolium*. (v. s.)

66. *E. perfoliatum* (Linn. sp. 1174), canle tereti striato villosulo-hirsuto, foliis oppositis connato-perfoliatis lanceolatis acuminatis dentatis rugosis suprâ glabris pubescentibus subtus tomentosis aut subhirsutis, summis distinctis, corymbo conferto, invol. squamis 12-15 imbricatis lineari-lanceolatis, floribus in inv. 8-10, achænio glabro. ♀ in pratis humidis Amer. bor. à Canadâ ad Floridam. Pursh fl. bor. am. 2. p. 516. Ell. sketch 2. p. 302. Big. fl. bost. ed. 2. p. 297 et med. bot. 1. p. 38. t. 2. Raf. fl. med. 1. n. 36. ic. *E. connatum* Michx. fl. bor. am. 2. p. 99. Flores albi. Variat foliis super. alternis! Herba (vulgò *Boneset*) amaræ decoctum emeticum sudorificum febrifugum in Amer. popolare. (v. s.)

§. 4. *Verticillata*.

Capitula 5-10-flora. Invol. oblonga squamis oblongis imbricatis. Folia 3^{im} verticillata. Caules herbacei.

67. *E. ternifolium* (Ell. sketch 2. p. 306), canle tereti striato pubescente solido, foliis 3-4 verticillatis ovato-lanceolatis acuminatis dentatis suprâ sparsè subpuberis subtus pubescentibus glandulosis, corymbo conferto, invol. squamis imbricatis, ext. brevibus valde puberis, int. lineari-oblongis striatis subpubescentibus, floribus in capitulo paucis. — in pinguibus et demissis Carolinæ (Fras. ! Ell.). Capitula parva pallida. Achænia angulata. Cor. albidæ vix purpureo tinctæ. *E. trifoliatum* Linn. sp. 1173 ex syn. Gron. virg. 119, sed species cum aliis verticillatis sæpè confusa. (v. s.)

β. foliis majoribus basi longius cuneato-acuminatis. *E. verticillatum* var. Torr. ! in litt. 1820. Ell. l. c. in adn. (v. s.)

68. *E. purpureum* (Linn. sp. 1173), canle tereti lævi glaucescente fistuloso, foliis 4-5^{is} verticillatis brevissimè petiolatis ovato-lanceolatis dentato-serratis subrugoso-venosis scabrinusculis, capitulis corymbosis 5-floris, invol. squamis interioribus longioribus subscariosis lineari-oblongis glabris, achænio pentagono vix pubescente. ♀ in humidis pinguibus Amer. bor. à Canadâ ad Caro-

linam. Pursh fl. bor. am. 2. p. 515. Ell. sketch 2. p. 307. — Corn. cad. cap. 72. p. 191. ic. Moris. ox. 3. s. 7. t. 13. f. 4. male. Dod. mem. t. 217. An E. levigatum Torr. cat. pl. New-York. p. 92. non Lam. huc revocandum? E. purpureum Lour. videtur omnino diversum! Herba obscure virescens sæpe purpureascent. Cor. pallide purpureæ. (v. s.)

69. E. MACULATUM (Linn. amœn. ac. 4. p. 288), caule solido sulcato superne pubescente sæpe lineis purpureis maculato, foliis 4-5^{te} verticillatis petiolatis ovato-lanceolatis inæqualiter serratis subtus pubescentibus, corymbo polycephalo, capitulis pedicellatis 5-8-floris, invol. squamis imbricatis lævibus ext. subrotundis int. linearibus elongatis, achenio angulato glabro glanduloso. 2/ in humidis Amer. bor. à Canadâ ad Carolinam. Pursh fl. 2. p. 1760. Ell. sketch 2. p. 308. — Herm. parad. 158. t. 159. Moris. ox. s. 7. t. 18. f. 3. An confirmante cl. Hooker fl. bor. 1. p. 305, ab E. purpureo satis differt? (v. s.)
70. E. VERTICILLATUM (Muhl. in Willd. sp. 3. p. 1760), caule tereti lævi solido sæpe lineis purpureis maculato, foliis 3-4^{te} verticillatis basi cuneato-attenuatis ovato-lanceolatis inæqualiter serratis glabrisculis, corymbo composito, capitulis pedicellatis 5-10^{te}-floris, invol. squamis imbricatis lævibus. 2/ in pinguius Carolinæ super. (Bosc!) Georgiæ (Ell. sketch 2. p. 308) agri Bostoniensis (Big. ! fl. hort. ed. 2. p. 298). E. purpureum Michx. fl. 2. p. 99. non alior. E. fusco-rubrum Walt. car. 199? Cor. purpureæ. Affine E. maculato et sæpe cum eo confusum, sed foliis non verè petiolatis facillè differt. (v. s.)
71. E. DUBIUM (Poir. suppl. 2. p. 606), caule tereti striato solido pubescente, foliis 3^{is} verticillatis petiolatis ovatis vix acuminatis subserratis suprâ sparse puberulis subtus pubescentibus glandulosisque, corymbo longè pedunculato, capitulis 3-floris, invol. squamis imbricatis lævibus, int. longioribus acuminatis lineatis, achenio angulato glanduloso. 2/ in Pennsylvania (Muhl. !) novâ Cæsareâ (Pursh.) Carolinâ merid. (Fraser. !). E. punctatum Willd. enum. 853. non Lam. Affine videtur E. ternifolium sed capitula pauciflora, invol. glabra, folia ovata, etc. (v. s.)

Ser. II. SUBIMBRICATA. *Nempè involucris aut laxè aut paucis seriebus imbricatis ovatis campanulatisve.*

* *Capitulis 3-5-floris, foliis alternis.*

72. E. TRIFLORUM (Aubl. guian. 2. p. 795. t. 314), fruticosum, ramis teretibus sarmentosis albo-velutinis, foliis alternis breviter petiolatis ovato-lanceolatis acutis subcordatis integerrimis suprâ scabris subtus albidovelutinis, pedunculis axill. et termin. racemoso-paniculatis, capitulis glomeratis 3-floris, invol. oblongi squamis imbricatis oblongis acutis concavis. 3 in Guianâ Gallicâ ad ripam fl. Aroura. Lam. dict. 2. p. 404. Willd. sp. 3. p. 1749. Cor. alba. Petiol. 4 lin. Limb. fol. 5 poll. longus, 2 poll. latus.
73. E. CAMPESTRE, herbaceum erectum glabrum, caulibus simplicibus striatulis, foliis alternis oblongis basi subtrinerviis cuneato-subpetiolulatis integris, à medio ad apicem dentatis, corymbo composito fastigiato, capitulis breviter pedicellatis 5-floris, invol. squamis subimbricatis acutiusculis, corollis acheniisque puberulis. 2/ in campis editis prov. Sancti-Pauli novæ flor. legit cl. Lund (n. 852!). Herba pedalis stricta. Folia pollicem longa, 4 lin. lata. Cor. dilatè violaceæ. (v. s.)
74. E. SUBALTERNIFOLIUM, caule herbaceo erecto gracili tereti pubescenti-velutino, foliis inf. oppositis cæteris subalternis breviter petiolatis ellipticis seu obovato-oblongis utrinquè attenuatis rigidulis subdentatis suprâ glabris subtus trinerviis et conspicuè reticulatis canescenti-puberulis, corymbo terminali pedunculato conferto, capitulis ad apices ramulorum plurimis sessilibus 5-floris, invol. squamis 2-seriatis subimbricatis, ext. obtusis dorso puberulis, inter. glabris apice serrulatis, achenio glabro. 2/ in Brasiliâ prov. Minas General. ad Serra do Frio (Vauth. ! n. 335), in campis editis prov. Sancti-Pauli (Lund herb. n. 851!). Herba circ. pedalis, rhizom. verisim. lignoso. Folia pseudo-

opposita cum petiolo 10-12 lin. longa, 4 lin. lata. Cor. ex sicco purpurascens. Styli longè exserti. Pappus longit. corollæ. (v. s.)

75. *E. DICTIOHYLLUM*, caule suffruticoso erecto tereti pulverulento-velutino, foliis alternis coriaceis ovatis basi in petiolulum cuneato-attenuatis supernè remotè et calloso-serratis utrinquè glandulosis et puberulis subtus reticulo nervulorum elevatorum eleganter areolato, corymbo paniculato polycephalo, invol. squamis omnibus linearibus acuminatis exter. dorso puberulis cæter. glabris, flor. 5, acheniis villosis. $\frac{3}{4}$ in campis editis prov. Sancti-Pauli nov. flor. legit cl. Land (n. 853!). Flores albi. Folia 3 poll. longa, $1\frac{1}{2}$ lata. (v. s.)

** Capitulis 5-10-floris, foliis oppositis.

76. *E. IRESINOIDES* (H. B. et Kunth nov. gen. am. 4. p. 106. t. 340), caule herbaceo vulnibili striato villosa-pubescente, ramis divaricatis, foliis oppositis petiolatis deltoideo-ovatis acutis serrato-dentatis triplinerviis suprâ pubescenti-hirtis subtus molliter canescenti-pubescentibus, paniculâ terminali trichotomâ diffusâ, capitulis pedicellatis subfastigiatis sub-5-floris, invol. squamis circ. 16 subimbricatis oblongis glabris, ext. mucronatis, int. acuminatis. $\frac{2}{3}$ ad radices Andium Novo-Granatensium prope Ybague. Cor. alba. Habitus Iresinis. Petiol. 4 lin. limbus 20-21 lin. longus, 13-14 latus.

77. *E. PYRIFOLIUM*, fruticosum glabrum, ramis teretibus, foliis oppositis petiolatis ovatis ovalibusve utrinquè vix acutis rigidulis integerrimis penninerviis, paniculis axillarihus oppositis terminalibusque polycephalis, capitulis pedicellatis 5-floris, invol. squamis 4-seriatis imbricatis ferè 5-fariis striatis ext. brevibus ovatis acutis, int. oblongis longioribus, achenio puberulo. $\frac{3}{4}$ in fruticetis ad Cercovado Brasilæ legit cl. Lhostky. Petioli 6-9 lin. longi. Limb. fol. 3 poll. longus. 18-21 lin. latus. (v. s. comm. à cl. inv.)

78. *E. DENDROIDES* (Spreng. syst. 3. p. 415), arboreum glabrum, ramis sexangularibus, foliis oppositis petiolatis ovatis acutis basi cuneatis crenato-dentatis coriaceis reticulato-venosis, corymbis terminalibus ramosissimis, capitulis geminis ternisve subsessilibus sub-5-floris, invol. oblongi squamis 7-8 imbricatis linearibus obtusis viscosis, achenio glabro. $\frac{5}{8}$ prope Loxa Novo-Granatens. Mikania arborea H. B. et Kunth nov. gen. am. 4. p. 139. Petioli 7-8 lin. longi. Limbus fol. sub 4-pollicaris, 21-22 lin. latus.

79. *E. BACCHAROIDES* (H. B. et Kunth nov. gen. am. 4. p. 132), fruticosum glabrum, ramis sulcato-striatis, foliis oppositis petiolatis lanceolato-oblongis utrinquè acutis remotè serratis coriaceis, corymbis terminalibus, ramulis diffusis viscosis, capitulis pedicellatis 6-floris, invol. cylindraceo-campanulatis squamis imbricatis oblongis acutiusculis subciliatis, achenio ad angulos scabro. $\frac{3}{4}$ in umbrosis juxta Alagner Novo-Granatensium. Petiol. 3 lin. Limb. fol. 2 $\frac{1}{2}$ poll. longus, 10-11 lin. latus. Cor. alba.

80. *E. ANGUSTIFOLIUM* (Spreng. syst. 3. p. 415), fruticosum, ramis teretibus albedo-tomentosis, foliis oppositis petiolatis oblongo-lanceolatis acutiusculis basi rotundatis integerrimis reticulato-venosis coriaceis suprâ glabris subtus albedo-tomentosis, corymbo paniculatis, capitulis subgeminis sessilibus 6-floris, invol. oblongi squamis subdenis imbricatis oblongo-linearibus apice fimbriatis, acheniis glabrisculis. $\frac{3}{4}$ in umbrosis Nova-Granatæ inter Ybague et Cnestade Tolima. Cor. pilosiuscula. Anthæræ subexsertæ. Mikania angustifolia H. B. et Kunth nov. gen. am. 4. p. 138. Petioli semipoll. Limb. fol. 4 $\frac{1}{2}$ poll. longus, 14 lin. latus.

81. *E. ACUMINATUM* (H. B. et Kunth nov. gen. am. 4. p. 107), caule herbaceo-sexangulari glabro, foliis oppositis petiolatis subrotundo-ovatis læviter cordatis valde acuminatis dentato-serratis 5-nerviis glabris subtus in venis nervisque pubescentibus, corymbo terminali conglomerato, capitulis sessilibus sub 7-floris, invol. cylindræci squamis imbricatis lineari-oblongis striatis glabris, exter. minimis, achenio glabro. $\frac{2}{3}$ in Novâ-Granatâ. Petiol. 15 lin. longus. Limb. fol. 4 poll. longus, 3 poll. latus. Cor. alba? glabra.

82. *E. tenuifolium* (H. B. et Kunth nov. gen. am. 4. p. 107), caule herbaceo erecto subangulati glabro, foliis oppositis petiolatis ovatis acutis grosse dentato-serratis trinerviis glabris subtus in venis nervisque pubescentibus, corymbo terminali sessili oppositè ramosissimo, capitulis sessilibus conglomeratis sub 7-floris, invol. cylindracei squamis imbricatis, ext. ovatis brevibus ciliatis, int. oblongis obtusis glabriusculis. $\frac{1}{2}$ in humidis Amer. austr. inter Cumana et Bosdones. Petiol. 10 lin. Fol. limbus 2 $\frac{1}{2}$ poll. longus, 20 lin. latus.
83. *E. micranthum* (Less. in linnæa 1830. p. 138), fruticosum glabrum, foliis oppositis petiolatis papyraceis penninerviis ovato-lanceolatis acutiusculis remotè serratis subtus dense glandulosis, paniculis trichotomis subfastigiatis, capitulis 7-8-floris, invol. turbinate-cylindracei squamis siccis erecto-patentibus apice ciliatis. $\frac{3}{4}$ in sylvis Xalapæ Mexic. Cor. purpureæ. An *E. micranthum* Lag. nov. gen. et sp. 25 n. 333 omninò idem? Ex Lessingio aff. *E. fastigiato* Kunth! ex Lag. Conoclinio celestino!
84. *E. pallescens*, suffruticosum?, ramis teretibus tenuissimè et molliter pubescentibus, ramulis divaricatis, foliis oppositis petiolatis ovato-lanceolatis basi breviter cuneatis integris triplinerviis apice longè acuminatis dentato-serratis suprâ glabriusculis subtus pubescentibus, corymbis ad apices ramorum confertis polycephalis, capitulis breviter pedicellatis 7-9-floris, invol. squamis ovali-oblongis obtusis striatis 2-3-seriatis laxè imbricatis glabriusculis. $\frac{3}{4}$ in Brasiliæ prov. Minarum General. ad Sahara legit cl. Vauthier. Invol. straminea, corollas subæquantia. Achaenium glabrum. Petiol. semipollic. Limb. fol. 4 poll. longus, pollicem circ. latus. (v. s. comm. à cl. Vauthier n. 273.)
- E. hirsutum*, fruticosum, ramis velutino-hirsutis, foliis suprâ puberulis subtus velutino-hirsutis. $\frac{3}{4}$ in Brasiliæ prov. Rio-Grande, (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 789 miss.)
85. *E. glomeratum*, caule suffruticuloso erecto cum petiolis et pedunculis hirsutulo, foliis oppositis petiolatis basi cordatis ovato-lanceolatis acuminatis dentato-crenatis suprâ minutè pubescentibus subtus molliter subtomentosis, ramis floridis oppositis divaricatis paniculam confertissimam gerentibus, ramulis apice 3-cephalis, capitulis 7-8-floris, invol. squamis glabris striatulis parçè imbricatis, acheniis pentagonis hispidulis. $\frac{3}{4}$ inter Chilenses herb. Hænkeani adnumeratur. Folia ferè Betonicæ. Styli longè exserti. Capitula adeò aggregata ut primâ fronte multiflora videantur. (v. s. in h. Hænke.)
86. *E. leucophyllum* (H. B. et Kunth nov. gen. am. 4. p. 115), fruticosum, ramis teretibus albis tomentosis, foliis oppositis petiolatis ovatis apice angustatis acutis leviter cordatis subtrinerviis suprâ pubescentibus subtus albotomentosis, paniculis terminalibus ramosissimis foliatis diffusis, capitulis pedicellatis 9-10-floris, invol. oblongo-campanulati squamis imbricatis obtusis pubescentibus, ext. brevibus acutiusculis, achenio glabriusculo. $\frac{3}{4}$ in temperatis Andium Peruvianum. Cor. albæ? Petiol. subsemipollic. Limb. fol. 2 poll. longus, basi pollicem latus.
87. *E. lavandulæfolium*, fruticosum, ramis teretibus junioribus cano-tomentosis, foliis oppositis subsessilibus linearibus integerrimis obtusiusculis margine revolutis suprâ molliter villosis demùm glabris subtus tomentosis, paniculâ ovatâ brevi confertâ, capitulis ad apices ramorum congestis sessilibus 9-10-floris, invol. squamis oblongis obtusis laxè imbricatis, ext. dorso villosis, inter. glabriusculis subscaiosis. $\frac{3}{4}$ in Amer. merid. legit Dombey, sed locus prop. ign. in vallibus Andium Peruviae (h. Hænk!). Accedit ad *E. stachadifolium* sed folia revoluta nec plana. Recept. planum nec semiglobosum. Invol. squamæ obtusæ nec acutæ etc. (v. s. comm. à Mus. reg. Paris.)

*** Capitulis 10-20-floris, foliis oppositis.

88. *E. decemflorum*, suffruticosum, ramis teretibus apice præsertim pubescentibus, foliis oppositis brevissimè petiolatis elliptico-oblongis utrinquè acuminatis suprâ basin triplinerviis serrulatis suprâ aparsè puberulis subtus pubescentibus et glandulosis, corymbi terminalis congesti ramis oppositis, capitulis

subsessilibus 10-floris, invol. squamis 2-3-seriatis subimbricatis obtusis, exter. brevibus pubescentibus, inter. longioribus membranaceis glabris, achenio glabro. $\frac{5}{8}$ in Peruvia (Poeppig!), in vallibus Andium (h. Hænk.). Folia 3 poll. longa, pollicem lata. Capitula straminea. Invol. corollas aequantia. (v. s. comm. à cl. Poeppig. sub n. 30.)

89. *E. PELUCIDUM* (H. B. et Kunth nov. gen. am. 4. p. 108), caule herbaceo erecto sexangulati glabro, foliis oppositis petiolatis subrotundo-ovatis leviter cordatis acuminatis dentato-serratis 5-nerviis glabris subtus in venis nervisque pubescentibus, venulis pellucidis, corymbo terminali conglomerato, capitulis sessilibus 10-floris, invol. cylindracei squamis imbricatis lineari-oblongis glabris nitidis albidis, achenio secus angulos scabriusculo. $\frac{7}{8}$ in Novâ-Granatâ. Simile *E. acuminato* sed præter notas ind. differt ex cl. auct. Invol. 24-27 nec 15-phyllis.

90. *E. RELIANTHIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 127), fruticosum, ramis teretibus hispidis, foliis oppositis petiolatis ovato-oblongis acuminatis serratis supra scabriusculis subtus glanduloso-punctulatis et pilosisculis, corymbis terminalibus trifidis, capitulis pedicellatis 10-floris, invol. campanulato-cylindracei squamis imbricatis oblongo-linearibus obtusis glabriusculis. $\frac{5}{8}$ prope pagum Ayavaca Peruvianorum. Petiol. 4 lin. Limb. fol. sub 5-pollicaris, 22 lin. latus.

91. *E. CANESCENS* (Vahl! symb. 3. p. 95. t. 73), fruticosum, ramis teretibus apice pulverulento-velutinis, foliis oppositis breviter petiolatis ovatis dentibus paucis obtusis sublobatis integerrimisve obtusis supra glabriusculis subtus canescenti-velutinis, corymbo simplici terminali, capitulis pedicellatis 10-floris ovatis, invol. squamis 3-serialibus laxè imbricatis obtusis, inter. striatis. achenio pube minimâ subvillosa $\frac{5}{8}$ in ins. Sanctæ-Crucis (Vest in h. Puer.!) et Sanct.-Thomas. (Willd.). Folia petiolo 1-2 lin., limbo 9-12 lin. longo, 5-6 lato. (v. s.)

β. adenocarpum, foliis paulò minoribus, achenio glabro globulis glandulisve consperso. $\frac{5}{8}$ in Sancto-Domingo legit cl. Bertero. (v. s.)

92. *E. VARGASIANUM*, fruticosum, ramis teretibus petiolis pedunculisque densè breviterque hirsutis, foliis oppositis petiolatis latè ovatis acutis basi subcordatis 5-nerviis dentatis supra sparse scabridis subtus pubescenti-hirsutis, corymbo amplo conferto polycephalo oppositè ramoso, capitulis pedicellatis 10-floris, invol. squamis ovali-oblongis vix acutis 2-3-seriatis ciliatis dorso glabris, achenio puberulo. $\frac{5}{8}$ circa Caraccas legit cl. Jos. Vargas. Petiol. ferè sesquipollic. Fol. limbus 3-4 poll. longus, 2-2 $\frac{1}{2}$ poll. latus. Invol. straminea. (v. s. comm. à cl. Vargas.)

93. *E. NIVRUM* (H. B. et Kunth nov. gen. am. 4. p. 115. t. 342), fruticosum, ramis sulcatis tenuissimè petiolis pediculisque albo-lanatis, foliis oppositis petiolatis triangulari-ovatis acutis leviter cordatis crenatis trinerviis supra glabris nitidis subtus niveo-tomentosis, paniculis subcorymbosis ramosissimis foliatis diffusis, capitulis pedicellatis 10-11-floris, invol. oblongi squamis imbricatis striatis villosis-pubescentibus, ext. ovatis, achenio glabro. $\frac{3}{8}$ juxta Popayan Novo-Granatensium? Cor. alba? Petiol. 5-6 lin. Limb. fol. 15 lin. longus, 11 lin. latus. Species elegans!

94. *E. POPULIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 111), arborescens glabrum, ramis sulcatis, foliis oppositis petiolatis ovatis acutis obtusè grossè et inæqualiter dentatis subcoriaceis, paniculis terminalibus ramosissimis patentibus, capitulis sessilibus subquaternis approximato-congestis 10-12-floris, invol. oblongo-cylindraceo imbricato, squamis obtusiusculis trinerviis. $\frac{5}{8}$ in aridis calidissimis juxta pagum Mazatlan Mexicanorum. Petiol. 2 poll. Limb. fol. 7 poll. longus, 5 poll. latus.

95. *E. SOLIDAGINOIDES* (H. B. et Kunth nov. gen. am. 4. p. 126), fruticosum, ramis sexangularibus glabris, foliis oppositis petiolatis ovatis acuminatis serratis subtrinerviis supra glabris subtus punctulatis et pilosisculis, paniculis terminalibus ramosissimis foliatis divaricatis, capitulis pedicellatis 10-13-floris.

invol. campanulati squamis imbricatis lanceolatis acuto-subulatis subciliatis, achænio ad angulos hispidio. $\frac{3}{4}$ in temperatis Quitensium inter Tiscan et Alausi. Cor. albæ? E. celtidifolio et E. cœlestino (mihi diversissimis) dicitur affine.

96. *E. ELLIPTICUM*, scandens, ramis elongatis teretibus striatis pubescentibus, foliis alternis petiolatis ellipticis acuminatis glabris integerrimis, capitulis ad apices ramorum paniculatis pedicellatis 12-15-floris, invol. squamis 10-12 glabris subciliatis acutis, int. longioribus sed disco dimidio brevioribus. $\frac{3}{4}$? (in Indiâ or. ad Penang legit cl. Vallich. Commixtum errore quodam cum Mikaniâ volubili à Wall. cat. n. 3174 sed evidenter diversissimum. Flores violaceo-purpurei, fauce campanulatâ. Pappus rufus copiosus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
97. *E. FASTIGIATUM* (H. B. et Kunth nov. gen. am. 4. p. 125. t. 347), fruticosum glabrum, ramis sulcatis, foliis oppositis petiolatis oblongis acutis basi angustatis crenato-serratis coriaceis subviscosis, corymbis terminalibus ramosissimis fastigiatis, capitulis pedicellatis circ. 12-floris, invol. subcampanulati squamis imbricatis lineari-oblongis obtusis subciliatis. $\frac{3}{4}$ in frigidis montium Novo-Granatensium. Cor. alba. Petiol. 7-8 lin. Limb. fol. sesquipollicem longus, 13-14 lin. latus.
98. *E. PRUNIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 132. t. 349), arboreum, ramis teretiusculis, ramulis piloso-pubescentibus, foliis oppositis sparsive petiolatis oblongis acutis basi rotundatis subintegerrimis coriaceis suprà glabris subtus hirtis pubescentibus, corymbis terminalibus oppositis ramosissimis, capitulis pedicellatis 12-floris, invol. oblongi squamis laxè imbricatis acutis glutinosi glabris, achænio ad angulos scabriusculo. $\frac{5}{8}$ in convalli Tarquiensi juxta urbem Quito. Petiol. 2-3 lin. Limb. fol. 16 lin. longus, 8 lin. latus. Cor. alba.
99. *E. INULÆFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 109), caule herbaceo erecto sexangulari tomentoso-pubescente, foliis oppositis breviter petiolatis ovato-lanceolatis longissimè acuminatis basi in petiolum angustatis remotè serratis suprà pubescentibus subtus tomentoso-pubescentibus et canescentibus, paniculâ corymbosâ diffusâ, capitulis subsessilibus congestis circ. 13-floris, invol. oblongi squamis laxè imbricatis lineari-oblongis obtusis striatis ciliatis, achænio glabro. $\frac{1}{2}$ in dumetis circa urbem Mariquitæ Novo-Granatensium.
100. *E. MOLLE* (H. B. et Kunth nov. gen. am. 4. p. 109), caule herbaceo erecto sexangulari cano-tomentoso, foliis oppositis petiolatis ovato-oblongis longissimè acuminatis basi attenuatis obsolete crenatis suprà pubescentibus subtus molliter et tenuissimè tomentosis canescentibus, paniculâ subcorymbosâ patente, capitulis subsessilibus conglomerato-congestis circ. 13-floris, invol. oblongi squamis laxè imbricatis lineari-oblongis obtusis ciliatis, achænio glabro. $\frac{1}{2}$ juxta Mariquitæ Novo-Granatensium cum *E. inulæfolio* à quo vix satis differt.
101. *E. SUAVEOLENS* (H. B. et Kunth nov. gen. am. 4. p. 109), caule herbaceo erecto teretiusculo molliter pubescente, foliis oppositis petiolatis ovato-oblongis angustato-acuminatis basi angustatis obtusè serratis subtriplinerviis suprà scabris subtus hirtis pubescentibus subrugosis, paniculâ corymbosâ subfastigiata, capitulis subsessilibus conglomeratis circ. 13-floris, invol. oblongi squamis laxè imbricatis lineari-oblongis obtusis ciliatis, achænio glabro. $\frac{1}{2}$ in temperatis montium Novæ-Granatæ. Flores suaveolentes. Valdè affine *E. inulæfolio* et *E. molli*. — *E. circa Montevideo à Sellowio lectum et huc à Spreng. relatum est verisim. diversum.*
102. *E. CONCINNUM*, fruticosum, caule tereti glabro, ramis præ. ad apicem hirsutis, foliis oppositis breviter petiolatis ovali-lanceolatis acuminatis serratis penninerviis suprà glabriusculis subtus cano-tomentosis, corymbo laxo fastigiato, capitulis pedicellatis 15-floris, invol. squamis 3-seriatis laxè subimbricatis oblongo-linearibus acutiusculis ciliatis dorso puberulis, achænio angulato glabro. $\frac{3}{4}$ in Brasiiliæ prov. Rio-Grande. Folia 2 poll. longa, 7 lin. lata.

Capit. 6-7 lin. longa. Achænium bilineare. Cor. ex sicco pallidæ. Styli valdè exserti. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 827 miss.)

103. *E. PENTLANDIANUM*, fruticosum glabrum, ramis teretibus, foliis oppositis breviter petiolatis ovato-lanceolatis subacuminatis serratis basi trinerviis subtus pallidis, corymbis terminalibus fastigiatis, capitulis pedicellatis 12-15-floris, invol. squamis oblongis obtusiusculis 2-3-seriatis laxè imbricatis puberulis corollis dimidio brevioribus, acheniis glandulosis glabris. $\frac{5}{8}$ in Amer. merid. parte centrali nunc Bolivia dicta legit cl. Pentland. Frutex ramosissimus obscure virens. Folia conferta. Cor. purpureæ. Pappus sordidus. Achænia albidæ. (v. s. comm. à cl. inv.)

104. *E. MONTANUM* (Swartz! prodr. III. flor. 1311), fruticosum, ramis teretibus hirsutiusculis, foliis oppositis breviter petiolatis cordiformibus acutis denticulatis suprâ sparsè hispido-scabris subtus secus nervos hirsutis granuliferis, corymbo pedunculato, ramis trifidis polycephalis, capit. circ. 15-floris, invol. squamis imbricatis conniventibus acutis hirtis. $\frac{5}{8}$ in montibus altis Jamaicae. Petioli 2 lin. hirsuti. Limbi fol. 2 $\frac{1}{2}$ poll. longi, 1 $\frac{1}{2}$ lati, lobis baseos conniventibus. Cor. exalbidæ. Pubes ferruginea. Affine *E. cordifolio*. (v. s. ex h. Swartz.)

105. *E. GLABRATUM* (H. B. et Kunth nov. gen. am. 4. p. 127), fruticosum glabrum, ramis teretibus striatis, foliis oppositis petiolatis ovato-oblongis acutis dentato-serratis subcarnosis, pedunculis axillaribus terminalibusque subtricephalis, capitulis circ. 16-floris, invol. campanulati squamis subimbricatis lanceolatis obtusiusculis externè subviscosis, achænio 5-angulato ad angulos scabro. $\frac{5}{8}$ in locis altis montium Mexicanorum inter Omiltau et Real del monte. Cor. albæ glabræ invol. duplò longiores. Petiol. $\frac{1}{2}$ poll. Limb. fol. 2 $\frac{1}{2}$ poll. longus, pollicem et paulò ultra latus.

106. *E. RUPICAULE* (H. B. et Kunth nov. gen. am. 4. p. 124), herbaceum glabrum, caule simplicis, foliis oppositis petiolatis ovato-deltoides acuminatis grossè dentato-serratis trinerviis, venulis pellucidis, corymbi terminalis ramosis ramulis pubescentibus, capitulis pedicellatis circ. 16-floris, invol. subcampanulati squamis subimbricatis lanceolatis acutis subciliatis. $\frac{7}{8}$ in Mexici montanis inter Puerto de Varietos et Santa-Rosa de la Sierra. Cor. albæ. Caulis petiolique purpurei. Petiol. 7-8 lin. Limb. 3 poll. longus, 2 poll. latus.

107. *E. GLUTINOSUM* (Lam. dict. 2. p. 408), fruticosum, ramis teretibus sulcatorugosis glabris apice glutinosis, foliis oppositis petiolatis cordato-lanceolatis acuminatis serratis suprâ bullato-rugosis glabris subtus tomentosis, corymbo oblongo paniculato terminali, capitulis 15-18-floris, invol. ovali-oblongi squamis imbricatis pubescentibus acutis. $\frac{5}{8}$ in Peruvia ex h. Juss. inter Mulato et El Tambillo Quitensium (H. B. et Kunth 4. p. 131). Fol. 5-6 poll. longa, 1 $\frac{1}{2}$ lata. Cor. purpureæ. Arbor 2-3-org. cortice cinereo.

108. *E. SALVIA* (Colla pl. chil. f. 2. p. 8. n. 53 act. tur. 38. p. 8), fruticosum, ramis teretibus striatis, junioribus foliisque novellis viscosis, foliis oppositis petiolatis lanceolatis acutis basi truncatis aut subattenuatis grossè crenato-dentatis rigidulis suprâ glabris bullulatis subtus nervis elevatis reticulatis pubescentibus, corymbo terminali composito conferto, capitulis pedicellatis 15-20-floris, invol. squamis 2-3-seriatis imbricatis lanceolatis acuminatis, ext. brevioribus pubescenti-viscosis, intimis lævibus coloratis. $\frac{5}{8}$ in Chili circa Valparaíso, Quillota, Rancagua, inter fruticeta et sylvas (Bert. ! Gay ! etc.). An ex loco natali *E. reticulatum* Hook. et Arn. bot. 1. p. 29? non Desv. Cor. violaceæ. Achæn. 5-costatum. Vulgò à Chilenis dic. *Salvia Macho* ex Bertero. (v. s.)

109. *E. BACLEANUM*, suffruticosum pube brevi confertâ pubescenti-hirtellum, caulibus teretibus apice subaphyllis, foliis oppositis et hinc inde alternis in petiolum subattenuatis oblongis ovalibusve dentatis aut profundè incisis, paniculæ subcorymbosæ ramis alternis, capitulis pedicellatis circ. 16-20-floris, invol. squamis 2-3-seriatis laxè imbricatis lineari-oblongis obtusissimis apice membranaceis subciliatis, achænio angulato villosa. $\frac{5}{8}$ in Brasiliæ merid. ad

oram dictam Banda orientale legit cl. Baile, et paulò post naufragium miserabile collectiones suas ferè totas amisit! undè levis solatii modò hujus nomen huic stirpi miraculè salvatè dedi. Etiam vidì inter Pernambuco (h. Hænk.!) et sine loco (h. Née!). Species ex habitu cinerascens, foliis variè incisis facilè distincta, Pappus sordidus. (v. s.)

110. *E. KRÖDIFOLIUM*, herbaceum? caule erecto tereti infernè hispido folioso supernè nudo pulverulento-pubescente, foliis oppositis dorso et margine hispidis in petiolum longè attenuatis supernè pinnatipartitis, lobis pinnatifidis lobulis oblongo-linearibus subacutis, corymbi terminalis conferti ramis alternis approximatis, capitulis 18-20-floris pedicellatis, invol. squamis circ. 2-seriatis, ext. paucis parvis acutis, int. oblongis basi striatis apice submembranaceis obtusis puberis. ? in Brasiliæ prov. Rio-Grande. Affine *E. Baileano* sed dist. caule foliisque setoso-hispidis, capitulis dimidio minoribus, etc. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 808 miss.)

111. *E. STEVIAEFOLIUM*, herbaceum, caule erecto tereti hirsuto oppositè ramoso, foliis oppositis in petiolum attenuatis lanceolatis acuminatis serratis trinerviis suprà sparè puberis subtus pubescenti-hirsutis, corymbis axillaribus parvis et terminalibus fastigiatis, capitulis breviter pedicellatis 15-20-floris, invol. biserialis squamis ext. parvis paucis acutis, int. oblongis pubescentibus apice membranaceis obtusis. ? in Brasiliæ prov. Sancti-Pauli. Invol. apice subcolorata. Folia 15-18 lin. longa, 5 lin. lata. Cor. parvæ invol. ferè breviores. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 496 miss.)

β. *angustius*, caule magis hirsuto, foliis angustioribus et longioribus (2 circ. pollices longis, 3 lin. latis). — in prov. Rio-Grande (h. Hænk. imp. Bras. n. 806). An species diversa. (v. s. in h. Mus. reg. Par.)

112. *E. PAULINUM*, herbaceum, ramis divaricatis teretibus hispidis, foliis oppositis et alternis lanceolatis basi in petiolum attenuatis triplinerviis integrè supernè acuminatis dentatis, paniculæ corymbosæ ramis alternis velutinis, capitulis ad apices ramulorum confertis pedicellatis 15-20-floris, invol. squamis biserialis, ext. paucis parvis acutis, int. oblongis basi striatis apice obtusis pubescentibus ciliolatisque submembranaceis, achænio glanduloso. — in Brasiliæ prov. Sancti-Pauli. Folia sesquipollicem longa, 4-5 lin. lata, summa subintegerrima. Species affinis *E. pubescenti* et *E. decemfloro*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 498 miss.)

*** Capitulis 20-floris.

113. *E. HÆNKEANUM*, fruticosum, ramis teretibus apice pubescenti-velutinis, foliis oppositis petiolatis latè ovato-rhombicis basi cuneatis trinerviis apice acuminatis integerrimis glabresculis, corymbo laxo, pedicellis longè bracteolatis, capitulis circ. 20-floris, invol. ovalis squamis imbricatis acutis vix pubescentibus, achæniis glabris. ? in Mexico legit cl. Hænke. Petiol. 9-12 lin. longi. Limb. fol. 3 poll. longus sesquipollicem latus. Folia referunt *Hederæ helicis* var. integrifoliam. (v. s. in h. Hænke à cl. Sternb. comm.)

β? *velutinum*, ramis petiolis folisque pubescenti-velutinis, capitalis circ. 30-floris, achæniis brevioribus glaberrimis. ? ad Acapulco legit cl. Hænke. An fortè species propria inter *E. Neesianum* et *E. Hænkeanum* media? (v. s. in h. Hænke à cl. Sternb. comm.)

114. *E. HEMISPHERICUM*, fruticosum erectum, ramis 3-4-angulatis petiolis pedicellisque pubescenti-hirtellis, foliis oppositis petiolatis ovatis acutis trinerviis dentatis glabresculis aut sparè puberulis subtus crebrè glanduloso-punctatis, paniculâ laxâ subirregulari, capitulis longè pedicellatis 20-floris ferè hemisphæricis, invol. squamis 3-seriatis laxè imbricatis striatis ext. brevibus acutis int. longioribus ciliatis apice ad dorsum hirsutis corollas æquantibus, achænio glabro. ? verè in Brasiliâ legit cl. Lund! Habitus paulò *E. glechonophylli* sed capitula diversissima. Petiol. 7-8 lin. longi. Limb. fol. pollicem longus, 8-10 lin. latus. (v. s.)

β. *molliusculum*, foliis duplò triplòve majoribus vix glandulosis, ramis te-

retiusculis. $\frac{3}{4}$ circa Rio-Janeiro (Lund!). Capitula exactè var. α . Prioris videtur var. solo pinguiori et magis umbroso lecta. (v. s. cum. α .)

115. *E. CACALIOIDES* (H. B. et Kunth nov. gen. am. 4. p. 130), fruticosum, ramis tetragonis striatis glabris viscosis, foliis oppositis petiolatis oblongo-lanceolatis acuminatis basi acutis profundè crenatis subcoriaceis suprâ glabris subtus tomentoso-pubescentibus, corymbis terminalibus trifidis ramosissimis, capitulis pedicellatis 20-floris, invol. campanulato-oblongi squamis laxè imbricatis striatis acutis ciliatis viscosis. $\frac{3}{4}$ in agro Quitensi? Cor. violacea? Petiol. 7-8 lin. Limb. fol. 3 poll. longus, pollicem latus.

116. *E. XALAPPENSE* (H. B. et Kunth nov. gen. am. 4. p. 128), fruticosum glabrum, ramis sexangularibus striatis, foliis oppositis petiolatis ovato-ellipticis acutis serratis subtriplinerviis subcarnosis, paniculis terminalibus ramosissimis foliatis, ramis subgeminis oppositisve, capitulis pedicellatis ternis circ. 20-floris, invol. campanulati squamis oblongo-lanceolatis acutis imbricatis subviscosis. $\frac{3}{4}$ prope urbem Xalappam Mexicanorum. Cor. alba. Petiol. $\frac{1}{2}$ poll. Limb. fol. 20 lin. longus, 10 lin. latus. Affine *E. glabrato*.

117. *E. RUGOSUM* (H. B. et Kunth nov. gen. am. 4. p. 114. non Schrad.), fruticosum, ramis teretibus molliter pubescentibus, foliis oppositis subsessilibus triangulari-ovatis acutis grossè crenatis coriaceis suprâ bullato-rugosis scabris subtus hispido-tomentosis et canescentibus, corymbis terminalibus ramosis densis, capitulis circ. 20-floris, invol. campanulato-cylindracei squamis imbricatis, ext. subrotundis, int. oblongis obtusis striatis ciliatis, achænio ad angulos scabriusculo. $\frac{3}{4}$ in collibus apricis aridis inter urbem Alausi et Cerro de Sitzan prov. Quitensis. Cor. pallidè purpureæ. Folia 14-15 lin. longa, 13 lin. lata, petiol. 2 lin.

118. *E. SCHIEDRANUM* (Schrad.! ind. sem. h. Goett. 1832. p. 3), suffruticosum, caulibus teretibus puberulis, foliis oppositis petiolatis ovato-lanceolatis acuminatis crenato-serratis basi obtusis truncatisve suprâ sparse subtus secus nervos scabrisculis, corymbis pedunculatis oligocephalis axillaribus terminalibusque, capitulis breviter pedicellatis circ. 20-floris, invol. squamis laxè imbricatis oblongo-linearibus apice scariosis glabris ciliatis obtusis mucronatisve, achænio acutè angulato secus nervos scabro. $\frac{2}{3}$ $\frac{3}{4}$ in Mexico (Schrad.) circâ urbem et ad Tampico de Tamaulipas (Berl.). Cor. lilacinæ. Habitus ferè *E. iresinoidis*. (v. s.)

β . *virgatum*, foliis deltoideis imâ basi cuneatis. — in Americâ meridionali. *E. virgatum* Schrad.! ind. sem. h. Goett. 1829 ex litt. 1832. Caulis 3-4-pedalis, cortice sanguineo-castaneo. (v. s. ex h. Goett. comm. à cl. Schrad.)

γ . *diversifolium*, foliis deltoideis longè acuminatis, ramorum juniorum subcordatis, summis subrotundo-ellipticis. — in Americâ merid. sed ut var. β locus proprius incertus. *E. diversifolium* Schrad.! ind. sem. h. Goett. 1829. ex litt. 1832. Var. β et γ vix ab α distinctæ ex specim. Schraderianis. Nomen var. α selegi etiam paulò recentius, ob patriam certam. (v. s. ex h. Goett.)

119. *E. SALZMANNIANUM*, caule herbaceo tereti apice glabro cæterum sparse pubero, foliis oppositis petiolatis elliptico-oblongis obtusis basi attenuatis rigidulis integerrimis penninerviis suprâ scabris subtus glabriusculis, paniculæ corymbosæ ramis oppositis elongatis subfastigiatis, capitulis breviter pedicellatis 20-floris, invol. squamis 3-4-seriatis imbricatis linearibus obtusiusculis dorso puberis, achænio minutè pubescente. $\frac{2}{3}$ circa Bahiam Brasiliæ in collibus aridis legit cl. Salzmann. Caulis sesquipedalis. Invol. purpurascens. Cor. purpurea ex Salz. sed ex sicco aurea videtur. Pappus albus. Folia 3 poll. longa, 10 lin. lata, petiolo 4 lin. (v. s.)

120. *E. VAUTHIERIANUM*, caule herbaceo erecto tereti oppositè ramoso tenuissimè pulverulento-velutino, foliis oppositis breviter petiolatis oblongo-lanceolatis utrinquè attenuatis subtriplinerviis distanter et acutè serratis suprâ scabridis subtus pubescentibus, paniculâ compositâ amplâ polycephalâ, capitulis pedicellatis 20-floris, invol. squamis 3-4-seriatis laxè imbricatis striatis pube-

rulis et apice subciliatis, intimis acutis. $\frac{1}{2}$? in Brasiliæ prov. Minarum Gener. prope Mariannam legit cl. Vauthier. Achænium glanduloso-puberulum. Folia 5-6 poll. longa, 1-2 poll. lata. (v. s. comm. à cl. Vauthier sub n. 274.)

***** *Capitulis 10-25-floris, foliis alternis.*

121. *E. VINDEX*, rhizomate crasso lignoso, caulibus pluribus herbaceis simplicibus erectis teretibus foliisque puberulis, foliis paucis remotis parvis alternis ovali-lanceolatis subsessilibus integris, capitulis 7-8 subracemosis, invol. ovatis squamis subimbricatis striatis ovali-oblongis acutis disco brevioribus, achæniis pentagonis subhirtellis. $\frac{3}{4}$ in Brasiliæ prov. Minarum Gener. in altis deserti Minar. Novarum legit cl. de Martius. E campis adustis invicta quotannis assurgit. Vernonia vindex Mart. ! Herb. Habitus omnino Vernoniæ, sed pappus et stigmata Eupatorii. Caules pedales. Folia 4-5 lin. longa. Cor. et styli purpurascens. Flores in inv. 20-25. (v. s. in h. Acad. reg. Monac.)

***** *Capitulis 25-30-floris, foliis oppositis.*

122. *E. NEEANUM*, fruticosum, ramis teretibus apice pubescenti-velutinis, foliis oppositis petiolatis latè ovatis acuminatis ferè triangularibus angulis infer. obtusis subcrenatis trinerviis anprà glabriusculis subtus pubescentibus, paniculæ subcorymbosæ ramis oppositis, capitulis pedicellatis obovatis circ. 25-floris, invol. squamis imbricatis pubescentibus, ext. acutis intimis obtusis, achænio angulato secus angulos scabro. $\frac{3}{4}$ circa Acapulco legit cl. Née. Petioli 9-10 lin. longi. Limb. 2 poll. longus, 15 lin. latus. Cor. invol. longiores. (v. s. ex herb. Thibaud.)
123. *E. TINIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 133), fruticosum glabrum viscosum, ramis subaulcatis, foliis oppositis petiolatis ovato-ellipticis acutis basi rotundatis crenato-serratis coriaceis nitidis, corymbis terminalibus oppositè ramosis, capitulis breviter pedicellatis 20-25-floris, invol. campanulati squamis imbricatis lanceolatis scutiusculis striatis, achænio glabriusculo. $\frac{3}{4}$ in Novâ-Granatâ? Petiol. semipoll. Limbus fol. $3\frac{1}{2}$ poll. longus, 2 poll. latus. Cor. alba.
124. *E. STILLINGIAEFOLIUM*, fruticosum glabrum, ramis teretibus, foliis oppositis petiolatis latè ovato-lanceolatis acuminatis basi trinerviis obtusis serratis apice integerrimis subtus crebrè et minutè glandulosis, paniculâ compositâ corymbosâ, capitulis pedicellatis ovatis circ. 20-25-floris, invol. squamis pauciserialis imbricatis linearibus acutis pedicellatisque minutè pubescentibus, achænio angulato secus angulos scabrido. $\frac{3}{4}$ in Mexico ad Tamaulipas legit cl. Berlandier. Cor. rosæ. Petiol. 9-12 lin. longi. Fol. limbi 3 poll. longi, 2 poll. lati. Affinis *E. collino*. (v. s. comm. à cl. Berland.)
125. *E. MENDEZII*, fruticosum, ramis teretibus apice breviter velutino-hispidulis, foliis oppositis petiolatis ovato-lanceolatis acuminatis basi et apice integris medio serratis trinerviis anprà subpuberulis subtus petiolisque velutino-hispidulis, paniculâ amplâ trichotomâ ramosissimâ, capitulis pedicellatis oblongo-obovatis circ. 25-floris, invol. squamis imbricatis acuminatis dorso puberulis, achænio glabro. $\frac{3}{4}$ in Leon Mexic. ad occid. urbis Guanajuato legit cl. Mendez. Petiol. 6 lin. longi. Limbus 30-33 lin. longus, sesquipollicem latus. Frondescentia cinerea. (v. s.)
126. *E. MALVÆFOLIUM*, fruticosum?, ramis teretibus tenuiter cum pedunculis petiolis nervisque foliorum pulverulento-velutinis, foliis oppositis longè petiolatis latè cordiformibus breviter acuminatis grossè dentatis trinerviis hinc indè subtrilobis utrinquè glabriusculis, corymbis terminalibus compositis foliis intermixtis, capitulis pedicellatis circ. 25-floris, invol. squamis laxè imbricatis striatulis subsentis glabris, achænio glabriusculo, pappo tenui caducissimo. $\frac{3}{4}$? in Mexico inter Vittoria et Tula legit cl. Berlandier. Petiol. 3-5 pollic. Limbi fol. 5-6 poll. longi, 4-5 lat. Cor. teretes brevissimè 5-dent. Styli longissimi. (v. s.)
127. *E. FULGIDIOSUM* (H. B. et Kunth nov. gen. am. 4. p. 110), caule suffruti-

coso volubili, ramis ferrugineo-tomentosis, foliis oppositis petiolatis oblongis acuminatis basi angustatis remotè denticulatis suprâ scabris subtus tenuissimè tomentoso-hirtis foliigineo-fuscescentibus, paniculis terminalibus et axillaribus pedunculatis divaricatis, capitulis subsessilibus approximatis circ. 25-floris, invol. semiglobosi squamis imbricatis oblongis obtusis ciliatis, recept. convexo, achænio ad angulos scabro. 5 \mathcal{L} in Andibus Quindensibus Novo-Granat. Cor. albæ? Petioli pollicares. Fol. limbus 6 poll. longus, 2-2 $\frac{1}{2}$ latus.

128. *E. ARBORUM* (H. B. et Kunth nov. gen. am. 4. p. 131), arboreum, ramis sexangularibus glabris, foliis oppositis petiolatis oblongo-lanceolatis basi rotundatis crenato-denticulatis coriaceis suprâ glabris subtus tenuissimè pubescentibus, paniculis terminalibus subcorymbosis, capitulis pedicellatis sessilibusve circ. 25-floris, invol. cylindraceo-campanulati squamis imbricatis striatis acutis subciliatis subviscosis, achænio ad angulos scabro. 5 inter Pomalacta et Alausi Quitensium. Petiol. 9 lin. long. Limb. fol. 3 poll. longus, 10-11 lin. latus. Cor. lilacina. Affine *E. cacalioidi*.

129. *E. DODONÆFOLIUM*, fruticosum glaberrimum, ramis teretibus, foliis oppositis subpetiolatis basi cuneato-attenuatis oblongis obtusis integerrimis aut summis callosio-denticulatis rigidis reticulatis subtus punctulatis, paniculæ ramis oppositis apice subcorymbosis, capitulis pedicellatis circ. 25-floris, invol. squamis 3-4-seriatis imbricatis linearibus striatulis apice obtusis ciliatis dorso pubescentibus. 5 in Peruvîâ legit cl. Poeppig. Affine *E. Blanchetiano*. Folia ramea internodio multò breviora, 2-2 $\frac{1}{2}$ poll. longa, 7-9 lin. lata. Achænia minima glabra. Pappus tenuis albus. (v. s. comm. à cl. Poeppig sub n. 18.)

130. *E. DISCOLOR*, fruticosum, ramis sexangularibus glabris vix apice summo subpuberulis, foliis oppositis petiolatis oblongo-lanceolatis basi attenuatis apice acuminatis suprâ glabris subrugosis subtus densè cano-tomentosis margine junioribus angustè revolutis vix crenulatis, corymbo trichotomo conferto terminali, capitulis ad apices ramulorum congestis subsessilibus 25-floris, invol. squamis laxè imbricatis pluri-seriatis acutis ciliatis, achæniis ad angulos subscabridis. 5 in Amer. calid. (Née!), in Peruvîâ montanis et in Huanacera circa urbem (Hænk!). An *E. persicæfolium* H. B. et Kunth sed folia basi non rotundata! An *E. salicinum* Lam. sed capitula non 10-12 sed 25-flora? Cæt. à foliis Persicæ et Salicis differt lanugine canâ paginæ infer. *E. discolor* Spreng. est *Vernonia*. (v. s.)

131. *E. SALVIEFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 131 non Sims), fruticosum volubile, ramis teretibus sulcatis glutinosis, foliis oppositis petiolatis oblongo-lanceolatis acuminatis basi rotundatis argutè serrato-dentatis coriaceis suprâ bullato-rugosis glabris subtus cano-tomentosis, corymbis terminalibus, capitulis pedicellatis et sessilibus 20-30-floris, invol. cylindraceo-campanulati squamis imbricatis striatis acuminatis glabris acutis, achænio ad angulos scabro. 5 in frigidis ad declivitatem montis Chimborazo. Cor. violacea. *E. glutinoso* affine.

132. *E. GLABRIUSCULUM*, herbaceum erectum glabriusculum, ramis oppositis cum nervis fol. subtus pedicellisque minutissimè pubescentibus, foliis oppositis petiolatis ovato-lanceolatis longè acuminatis grossè et acutè serratis basi cuneatis triplinerviis membranaceis, paniculæ amplæ ramis ramulisque oppositis, capitulis longè pedicellatis circ. 25-30-floris, invol. squamis 3-4-seriatis laxè imbricatis dorso minutè puberulis vix substriatis, intimis longioribus acutis, achænio glanduloso-pubero. 2 \mathcal{L} in Brasiliâ. Nimis affine *E. Vauthieriano* sed glabritie, petiolis triplò longioribus, foliis longius cuneatis regul. serratis, etc. differre videtur. (v. s.)

133. *E. SUBORTUsum*, fruticosum glabrum subviscosum, ramis teretibus, foliis oppositis breviter petiolatis oblongis obtusis basi attenuatis integerrimis rigidulis penninerviis, paniculæ ramis oppositis elongatis apice trifidis 15-20-cephalis, capitulis pedicellatis circ. 30-floris, invol. squamis 2-3-seriatis linearibus imbricatis dorso subpuberulis, intimis apice obtusis araneoso-ciliatis. 5 in Brasiliâ circa Bahiam legit cl. Blanchet (n. 1409!) et in siculis ad montem Sinet

prope Cayennam cl. Leprieur! Cor. purpureæ. Pappus albissimus. Styli longè exserti. Achænium secus costas scabridum. Folia 3 poll. et ultrà longa, pollicem circ. lata. (v. s.)

134. *E. OXYCHLÆNUM*, suffruticosum, ramis teretibus hispidis, foliis oppositis integerrimis aut apice subrepando-dentatis obtusis rigidulis suprâ scabris subtus pubescenti-hirtis penninerviis, inf. ellipticis in petiolum attenuatis, super. oblongis basi cuneatis subsessilibus, paniculæ ramis oppositis apice 5-6-cephalis, pedicellis glanduloso-pilosis, capitulis pedicellatis circ. 30-floris, invol. squamis 2-3-seriatis imbricatis linearibus acuminatissimis corollas paulò superantibus dorso glanduloso-puberis. ♀ in Bras. prov. Minas Gener. prope Mariannam legit cl. Vauthier (n. 2801). Affine priori. Cor. purpureæ. Rami rubentes. (v. s.)

135. *E. PERSICÆVOLIVUM* (H. B. et Kunth nov. gen. am. 4. p. 130), fruticosum, ramis sexangularibus glabriusculis, foliis oppositis petiolatis oblongo-lanceolatis acuminatis basi rotundatis crenulatis subcoriaceis suprâ glabris subtus cano-tomentosis, corymbis terminalibus, capitulis pedicellatis sessilibusve multi-? floris, invol. cylindraceo-campanulati squamis laxè imbricatis striatis acutis ciliatis. ♀ ad radices montis ignivomi Côtapaxi Quitensium prope Mulalo. Valdè affine *E. cacalioidi* dicitur.

136. *E. CORDIFOLIUM* (Swartz prod. III. flor. 1309), fruticosum, ramis teretibus glabris, ramulis hirsutis, foliis oppositis brevissimè petiolatis cordiformibus acuminatis serratis suprâ sparè hispidis subtus præsertim secus nervos majores hirsutis et granulis conspersis, corymbo terminali sessili, capitulis ad apices ramulorum confertis subsessilibus multifloris, invol. squamis imbricatis linearibus acutis semireflexis hirtis, achænio lineari. ♀ in rupibus calcareis et præcipitiis montium Jamaicae. Hirsuties ferruginea. Folia 2 poll. longa, $\frac{1}{2}$ poll. lata. Recept. convexum. (v. s. sine fl. ex h. Swartz.)

137. *E. GRACILE* (H. B. et Kunth nov. gen. am. 4. p. 124), fruticosum procumbens repens, ramis adscendentibus hirtis, foliis oppositis breviter petiolatis ovatis acutiusculis crenato-serratis glabris subtrinerviis, pedunculis axillaribus et terminal. 3-cephalis, capitulis pedicellatis multifloris, invol. campanulati squamis imbricatis lanceolatis acutis ciliatis glabris, achænio hispidulo. ♀ in temperatis juxta Guaduas Novo-Granat. Cor. alba? Petiolus vix lin. longus. Folia semipollicaria.

138. *E. EPHLOBIOIDES* (H. B. et Kunth nov. gen. am. 4. p. 125), fruticosum erectum glabrum, ramis teretibus lævibus, foliis oppositis subsessilibus ovato-oblongis acutiusculis remotè crenulatis trinerviis, pedunculis axill. et termin. 3-cephalis, capitulis pedicellatis multifloris, invol. campanulati squamis imbricatis oblongis acutis ciliatis, achænio ad angulos scabro. ♀ in temperatis Novæ-Granatæ? Affine *E. gracili* dicitur.

139. *E. ELEGANS* (H. B. et Kunth nov. gen. am. 4. p. 133 non Vent.), fruticosum, ramis teretibus piloso-tomentosis, foliis oppositis petiolatis elliptico-ovatis obtusis basi rotundatis crenatis coriaceis utrinquè scabriusculis subtus in nervis venisque pubescentibus; corymbis terminalibus congestis, capitulis subsessilibus multifloris, invol. campanulati squamis imbricatis lanceolatis obtusiusculis pilosis, achænio glabro. ♀ circa urbem Quito? Petiol. 2-3 lin. Limbus fol. 15 lin. longus, 8 lin. latus. Cor. alba?

140. *E. ARGUTUM* (H. B. et Kunth nov. gen. am. 4. p. 121), caule herbaceo striato hirtello, foliis suboppositis petiolatis lanceolatis acutis argutè serratis utrinquè tenuissimè glanduloso-hirtis, capitulis 1-2 longè pedunculatis terminalibus semiglobosis multifloris, invol. squamis subæqualibus 2-3-seriatis imbricatis hirtellis linearibus apice subulatis, achænio glabro. ♀ ad radice montis Chapoltepec Mexicanorum. Cor. violacea supernè hirta. Petiol. semipoll. Limb. fol. sesquipollicem longus, $\frac{1}{4}$ lin. latus.

141. *E. SORDIDUM* (Less. in linnæa 1831. p. 403), suffruticosum, ramis teretibus sordidè lanuginoso-tomentosis, foliis oppositis petiolatis membranaceis suprâ

basin quintuplinervio-penninerviis ellipticis utrinque obtusis supra hispidiusculis subtus tomentoso-pubescentibus, paniculis subcorymbosis trichotomis, capitulis pedicellatis multifloris, invol. campanulati squamis 3-seriatis linearibus acuminatis villosa-pubescentibus. $\frac{3}{4}$ in Mexici Cuesta-Grande de Chicon-quaco et in sylvis Jalapensisibus. Cor. purpureo-violaceæ. Fol. 2-5 poll. longa, 2 $\frac{1}{2}$ -3 $\frac{1}{4}$ lata.

***** *Capitulis 30-100-floris, foliis oppositis alternisve.*

142. *E. KARVINSKIANUM*, fruticosum, ramis teretibus vix apice puberulis, pediculis et præsertim petiolis hispidis, foliis oppositis petiolatis ovato-lanceolatis acuminatis argutè et parvè serratis basi ciliatis glabrusculis, corymbis ramos terminantibus 8-12-cephalis, capitulis longè pedicellatis 35-40-floris, invol. campanulati laxè imbricati squamis oblongo-linearibus obtusis ciliatis, achænio glabro. $\frac{5}{8}$ in Mexico legit cl. Karvinski. Affine *E. adenophoro*. Pili petiolorum articulati. Petiol. 4 lin. longi. Limb. fol. pollicem longus, 5-6 lin. latus. Styli longè exserti. (v. s. in h. Acad. reg. Monac.)
143. *E. AMPHIDYCTIUM*, caule è collo lignoso simplici herbaceo tereti confertè hirtello à medio ad apicem aphylo, foliis alternis sessilibus obovato-oblongis basi cuneatis obtusis integris apice serratis acutis utrinque puberulis et æqualiter nervis elevatis reticulatis, corymbo 7-8-cephalo conferto terminali, invol. squamis omnibus acuminatis ext. vix puberulis, flor. 30-35, achæniis minutissimè scabridis. $\frac{2}{3}$ in campis editis prov. Sancti-Pauli nov. flor. legit cl. Lund. Herba pedalis. Affinis *E. subalternifolio* et *dyciophyllo*, sed florum numero (30 nec 5) evidenter diversum. (v. s.)
144. *E. BALLOTÆFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 121), caule herbaceo erecto striato pubescenti-hirto, foliis alternis petiolatis ovato-triangularibus cordatis acutis inciso-crenatis subtrinerviis supra hispidis subtus hispidopubescentibus, corymbis terminalibus ramosissimis, capitulis pedicellatis 36-40-floris, invol. cylindræo-campanulati squamis imbricatis linearibus acuminatis glabris, ext. brevibus pubescentibus, achænio glabro. $\frac{2}{3}$ in Novâ-Granatâ? Cor. violacea. Petiol. 7 lin. longi. Fol. subpollic.
145. *E. LAMIIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 126 non Link.), caule herbaceo? volubili? striato hirto, foliis oppositis petiolatis ovatis cordatis angustato-acuminatis grossè crenatis rigidis supra bullato-rugosis et scabriusculis subtus hirtis, corymbis terminalibus, capitulis pedicellatis circ. 40-floris, invol. cylindræo-campanulati squamis imbricatis striatis ciliato-pubescentibus, ext. acutis, int. lanceolatis, achænio glabro. $\frac{2}{3}$? in agro Quitensi? Cor. violacea?
146. *E. VISCOSUM* (H. B. et Kunth nov. gen. am. 4. p. 129), fruticosum glabrum viscosum, ramis sexangularibus, foliis oppositis petiolatis obovato-oblongis acutis crenato-serratis coriaceis glanduloso-punctatis, corymbis terminalibus ramosissimis diffusis-subfastigiatis, capitulis pedicellatis circ. 40-floris, invol. campanulati squamis laxè imbricatis lanceolatis acutis, achænio ad angulos hispido. $\frac{3}{4}$ in agro Quitensi? Cor. violacea. Petiol. 5 lin. Limbus fol. 3-3 $\frac{1}{4}$ poll. longus, 16 lin. latus.
147. *E. FRATERNUM*, fruticosum glabrum, caule tereti, foliis oppositis vix petiolatis ovalibus integerrimis mucrone calloso subacutis rigidis penninerviis reticulatis, sup. internodio brevioribus, paniculæ ramis oppositis polycephalis, sup. corymbosis, capitulis pedicellatis 40-floris, invol. squamis 2-3-seriatis subimbricatis linearibus dorso et ad apicem margine subvillosis, intim. mucronatis, ext. subobtusis. $\frac{5}{8}$ in Amer. merid. (fortè Peruvia?) legit cl. Poeppig. Proximè accedit ad *E. Blanchetianum*, sed folia magis distantia, 3 poll. longa, 1 $\frac{1}{2}$ lata, capit. 40-flora, achænia tenuia puberula. (v. s. comm. à cl. inv. sub n. 2075.)
148. *E. VITALBÆ*, fruticosum scandens glabrum, ramis teretibus junioribus pubescentibus, foliis oppositis petiolatis ellipticis acutis seu plerisque ovato-lanceolatis acuminatis subdentatis quintuplinerviis, paniculæ ramis oppositis

- oligocephalis, capitulis laxis longè pedicellatis circ. 45-floris, invol. squamis glabris 3-4-seriatis laxè imbricatis striatis, ext. latè ovatis, int. linearibus acuminatis, achenio glabro. ζ in Brasiliâ prope Rio-Janeiro legit cl. Lund. Habitus ferè Clematidis. Petiol. 4 lin. Limb. 3 poll. longi, 10-15 lin. lati. Invol. foliola subscariosa, ext. latiora quàm in plerisque. (v. s. comm. à cl. Lund.)
149. *E. COLLINUM*, fruticosum glabrum, ramis teretibus apice summo subpulverulentis, foliis oppositis petiolatis ovato-lanceolatis acuminatis medio subserratis basi breviter cuneatis trinerviis subtùs parè glandulosis, paniculâ corymbosâ fastigiatâ, capitulis pedicellatis oblongis circ. 50-floris, invol. squamis imbricatis adpressis lanceolatis acuminatis, achenio secus angulos subscabro. ζ in collibus Mexici ad Tantoyuca legit cl. Berlandier. Frutex 6-pedalis. Petioli 9-10 lin. longi. Fol. limb. 3 poll. longus, ferè 2 poll. latus. Pedicelli bracteolati. (v. s. comm. à cl. Berlandier.)
150. *E. THESPISELEFOLIUM*, herbaceum?, caule tereti, pedunculis petiolis nervisque fol. pulverulento-velutinis, foliis oppositis petiolatis latè cordatis subrotundis cum acumine tenuissimè serrulatis præter nervos glabriusculos penninerviis, nerv. jugis 3 infer. approximatis, corymbo amplo composito fastigiato, capitulis pedicellatis campanulatis 80-100-floris, invol. squamis laxè imbricatis lanceolatis-linearibus acutis, achenio glabro. — in Mexico legit cl. Karvinski. Petioli 3 poll. longi. Limb. fol. 6-7 poll. longus totidemque latus. Cor. ex sicco purpurascens. Pappus albus. Folia sub paniculæ ramis ferè linearia. (v. s. in h. Acad. reg. Monac.)
- Ser. III. EXIMBRICATA. *Involucra nempe squamis 1-2-serialibus subæqualibus nec imbricatis constantia.*
- * *Capitulis 20-70-floris.*
151. *E. INNUMEROSUM*, herbaceum, caule angulato pube tenui brevi glandulosâ præsertim ad apicem hirtello, foliis oppositis et summis alternis breviter petiolatis cordatis acuminatis crenatis trinerviis suprâ sparsè scabridis subtùs pubescenti-tomentosis, corymbo composito terminali alternè ramoso, capitulis pedicellatis circ. 70-floris, invol. campanulatis squamis biseriatis linearibus acuminatis glabriusculis, achenio sparsè puberulo. ζ in Amer. calid.? legit cl. Née, sed locus proprius in herb. omissus. Petiol. fol. sup. 2 lin. Limb. 3 poll. longi 2 poll. lati. (v. s.)
152. *E. HERACLADUM*, caule herbaceo? tereti pube brevi glandulosâ confertâ cum petiolis et præsertim pediculis velutino, foliis oppositis petiolatis ovatis basi truncatis apice acuminatis grossè dentatis suprâ scabridis subtùs trinerviis velutino-subpubescentibus, corymbis plurimis alternè ramosis glanduloso-velutinis confertiusculis, capitulis pedicellatis 35-50-floris, invol. squamis linearibus subniseriatis dorso puberulis disco paulò brevioribus, achenio glanduloso. ζ in Brasiliæ prov. Rio-Grande. Folia summa alterna. Petiol. 3 lin. longi. Limb. fol. 15 lin. longi, 7-8 lin. latus. Recept. subconvexum. Cor. ex sicco incarnatæ. Styli teretes longè exserti. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 786 et 780 miss.)
153. *E. ADENANTHUM*, suffruticosum?, caule tereti alternè ramoso puberulo subviscidulo, foliis (sult. sup.) alternis basi longè cuneatis et petioliformibus medio ferè ovatis acuminatis dentatis subtriplinerviis suprâ glabris subtùs pubescenti-hirtellis glandulosis, corymbis ad apices ramorum 3-7-cephalis, capitulis pedicellatis 40-50-floris, corollis glanduliferis, invol. squamis 1-2-seriatis æqualibus acuminatis dorso puberulis demùm reflexis. ζ in Brasiliâ, sed locus proprius (verisim. Rio-Janeiro) incertus. Folia 3 poll. longa, pollicem lata. Achenia secus angulos scabra. (v. s.)
154. *E. ADENOPHORUM* (Spreng. syst. 3. p. 420), caule herbaceo subangulato ramis pedunculisque glanduloso-hirtis, foliis oppositis petiolatis ovato-triangularibus subacuminatis basi cuneatis grossè serratis suprâ glabris subtùs pilo-

- sinuosis, corymbis terminalibus trichotomis, capitulis pedicellatis 35-40-floris, invol. campanulati squamis 1-2-seriatis aequalibus acutis subciliatis, ext. pubescentibus, achænio 4-5-angulato glabro. $\frac{1}{2}$ in altâ planitie Mexicanâ inter Carpio et Gasave. E. glandulosum H. B. et Kunth nov. gen. am. 4. p. 122. t. 346. Lindl. bot. reg. t. 1723. non Michx. Cor. alba. Petiol. 8 lin. Limbus sesquipedalis longus, 14-15 lin. latus. (v. v. c.)
155. E. PRICHINCHENSE (H. B. et Kunth nov. gen. am. 4. p. 122), caule herbaceo volubili sexangulari villosa-hispido, foliis oppositis petiolatis subrotundo-ovatis acuminatis cordatis subduplicato-crenato-serratis trinerviis utrinquè pilosis-culis, corymbis terminalibus trichotomis, capitulis pedicellatis circ. 40-floris, invol. campanulati squamis subimbricatis lineari-lanceolatis acutis subciliatis, ext. pubescentibus, achænio ad angulos scabro. $\frac{1}{2}$ in declivitate montis Pichinchæ Quitensium. Cor. alba. Petiol. 10 lin. Limbus fol. 27 lin. longus, 18 latus.
156. E. CADUCISETUM, herbaceum glabrum, caule tereti oppositè et parcè ramoso, foliis oppositis breviter petiolatis ovatis acutis mediò parcè dentatis, super. valdè distantibus, corymbo trichotomo laxo, capitulis pedicellatis circ. 40-floris, invol. squamis biseriatis linearibus, ext. substriatis, int. apice submembranaceis acutis, achænio glabro, pappo parco caducissimo. $\frac{1}{2}$ circa Santa-Fé de Bogota. Herba circ. pedalis. Folia incl. petiolo lineam longo circ. 8-9 lin. longa, 3-4 lata. (v. s. comm. ab amic. Delessert.)
157. E. GLECHONOPHYLLUM (Less. in linnæâ 1831. p. 105), caule herbaceo erecto apice cum petiolis pedunculisque minutissimè pulverulento-puberulis, foliis oppositis petiolatis ovato-lanceolatis acuminatis dentatis membranaceis trinerviis glabrisculis, capitulis corymbosis paucis pedicellatis 30-floris, invol. squamis 1-2-seriatis linearibus acutiusculis 3-nerviis vix puberulis, achænio angulato subscabrido. $\frac{1}{2}$ in Chili (Cham.), prope Quillota in sylvaticis collium (Bert.), Valparaiso (Gaudich.), Conceptionem (Macrae!), E. chilense (Bert. non Mol.), Vulgò *Barba de Viajo*. Cor. roseæ. Petioli 4-12 lin. longi. Herba basi suffruticulosa. Affine E. urticæfolio (ex ic. Smith) sed jam glabritie ferè totâ et præsertim corollæ diversum. (v. s.)
158. E. CONGLOBATUM, suffruticosum, ramis teretibus striatulis apice petiolisque pubescenti-velutinis, foliis oppositis petiolatis ovatis subcordatis acuminatis dentatis trinerviis utrinquè subpuberulis subtus crebrè glandulosis, ramis floridis oppositis corymbosis apice capitula densè congesta subsessilis circ. 30-flora gerentibus, invol. squamis circ. biseriatis acuminatis dorso puberulis, corollâ glabrâ. $\frac{1}{2}$ circa Bahiam Brasiliæ in collibus umbrosis leger. cl. Salzmann et Blanchet. Cor. albidæ. Achænia subpubera. Frutex 5-6-pedalis. Simile E. urticæfolio, sed capitula conferta corollæ glabræ dimidio breviores, etc. (v. s.)
159. E. MYRTILLOIDES, suffruticulosum, ramis virgatis teretibus pulverulento-velutinis, foliis alternis breviter petiolatis ovatis basi obtusis apice acutis, dentatis suprâ glabris subtus tenuissimè pubescentibus glandulosis et trinerviis, corymbo terminali 5-6-cephalo conferto, capitulis brevè pedicellatis circ. 30-floris, invol. squamis 2-3-seriatis aequalibus linearibus acuminatis dorso pubescentibus, achænio glanduloso. $\frac{1}{2}$ in Brasiliæ prov. Minarum General. legit cl. Vauthier. Folia 7 lin. longa, 4-5 lin. lata. Petiol. 1 lin. Styli longè exserti. (v. s. comm. à cl. Vauthier.)
160. E. REMOTIFOLIUM, caule suffruticoso diffuso subvolubili tereti vix puberulo, foliis oppositis internodio multò brevioribus petiolatis ovatis basi obtusis apice acuminatis dentatis utrinquè petiolisque puberulo-hirtellis, ramis divaricatis in paniculas laxas subtrichotomas productis, capitulis pedicellatis circ. 25-floris, invol. squamis 2-3-seriatis glabris acutis, inter. longioribus, achænio glaberrimo. $\frac{1}{2}$ in Brasiliæ prov. Minarum Generalium prope Mariannam legit cl. Vauthier (n. 259!) et in apricis montosis prov. Rio-Janeiro cl. Lund! (Mus. Ber.). Herba Clematidea gracilis. Internodia 6-12 poll. longa. Flores albi odore Anthemidis Cotonæ (Lund). Petiol. 9 lin. Fol. limbus 15 lin. longus, 9 lin. latus. Pappus albiissimus. Achæniû nigrum. (v. s.)

161. *E. DELTOIDEUM* (Jacq. h. schoenbr. 3. p. 63. t. 369), herbaceum erectum, caule teretiusculo pulverulento velutino, foliis oppositis petiolatis hastato-triangularibus lobis apiceque acuminatis trinerviis inæqualiter serratis subtus pubescentibus, paniculâ thyrsoidæ apice subcorymbosâ, capitulis pedicellatis circ. 40-floris, invol. squamis linearibus acutis subniseriatis, achænio puberulo. $\frac{2}{3}$ in Mexico ad Villalpando (Mend.), in valle Toluca (Berl.), in Xxtapalapanæ montibus (Moq.). Bonpl. nav. 1, p. 38. t. 14. *E. triangulare* Moq. fl. mex. ic. Mexicanis dicitur *Xolochichitl* ex Moq. Cor. pallidè purpureæ. (v. v. et s.)
162. *E. PETIOLARE* (Moq. fl. mex. ic. ined.), fruticosum, ramis teretibus pubescenti-velutinis, foliis oppositis longè petiolatis cordiformibus acuminatis crenatis basi trinerviis suprâ sparse puberulis subtus pubescenti-velutinis, paniculâ ramis axillaribus et terminalibus folio æqualibus polycephalâ, capitulis pedicellatis circ. 35-40-floris, invol. squamis linearibus acuminatis subniseriatis pubescentibus, achænio gracili pubero. $\frac{5}{8}$ in Mexico. Bustamenta cordata Alam! in litt. 1831. Petiol. sesquipollic. Limbus fol. 3 poll. longus, 2-2 $\frac{1}{2}$ latus. Cor. albæ basi attenuatæ. (v. s. comm. à cl. Alaman et Mairet et in Andr. pl. mex. ined. n. 279.)
163. *E. LONICEROIDES* (H. B. et Kunth nov. gen. am. 4. p. 116), fruticosum, ramis subviscosis, foliis oppositis subsessilibus elliptico-oblongis obtusis subintegerrimis subcoriaceis glabris, paniculis terminalibus patulis, capitulis breviter pedicellatis 30-40-floris, invol. campanulati squamis lanceolato-linearibus acutis subviscosis ciliato-pubescentibus, achænio ad angulos hispidulo. $\frac{5}{8}$ in temperatis Andium Novo-Granatensium. Cor. purpureæ. Fol. 2 poll. longa, pollicem lata. Affine *E. amygdalino*.
164. *E. CONSANGUINEUM*, suffruticosum, caule erecto tereti cum petiolis pedunculisque tenuissimè pubescente, foliis oppositis petiolatis latè ovatis acuminatis vix obtusè acuminatis grossè crenatis basi truncatis trinerviis membranaceis suprâ secus nervos subtus ubique pube minimâ velutinis, ramis floridis axillaribus oligocephalis folio brevioribus, capitulis pedicellatis circ. 30-floris, invol. squamis 1-2-seriatis linearibus acutis dorso subpuberulis achænio angulato glabriusculo. $\frac{5}{8}$ in Brasiliâ circa Rio-Janeiro legit cl. Gaudichaud. Petioli pollicares. Limb. fol. 3 poll. longus, 2 poll. latus. Affine *E. sordido* sed multò minùs velutinum, folia longius petiolatum, folia latiora magis deltoidea, rami floridi breviores, etc. (v. s. comm. à cl. Gaudichaud.)
165. *E. TRISTE*, fruticosum, ramis teretibus cum petiolis pedunculisque pube ferrugineâ densâ hirsuto-velutinis, foliis oppositis petiolatis latè ovatis acuminatis basi obtusissimis crenato-dentatis suprâ scabridis subtus velutino-hirsutis, corymbi ramis primariis oppositis, capitulis longè pedicellatis 25-floris, invol. vix biserialis squamis linearibus acutis dorso velutinis, achæniis angulatis glandulosis. $\frac{5}{8}$ in Jamaicâ legit cl. Bertero. Eup. ceanothifolium Spreng. Syst. 3. p. 418. excl. syn. et patr. primâ. Petioli semipollicares. Limb. fol. 2 poll. longus, 1 $\frac{1}{2}$ latus. Cor. faux magis inflata quam in cæteris. Anthæræ odore perspicuæ. (v. s. comm. à cl. Bertero et Balbis.)
166. *E. ARALIEFOLIUM* (Less. in linnæâ 1831. p. 403), fruticosum scandens, ramis glabris junioribus subancipitibus, foliis oppositis petiolatis oblongo-ellipticis obovatis integerrimis penninerviis carnosius nitidis, paniculis suprâ decompositis trichotomis, capitulis 23-28-floris, invol. squamis pauciseriatis glabris, intim. linearibus acuminatis. $\frac{5}{8}$ in sylvis Misantlæ Mexicanorum.
167. *E. ORIGOCERHÆUM*, caulibus basi fruticulosius à rhizomate lignoso plurimis erectis teretibus pulverulento-subvelutinis, foliis oppositis petiolatis ovato-lanceolatis acutiusculis serrulatis glabriusculis, capitulis paucis corymbosis longè pedicellatis circ. 25-floris, invol. squamis 1-2-seriatis oblongis acuminatis glabris, achænio angulato glabriusculo, pappo pauciseto caducissimo. $\frac{2}{3}$ $\frac{5}{8}$ in Mexico legit cl. Hænke. Herba suffruticulosa vix pedalis. Radix teres lignosa simplex. Folia internodiis breviora, cum petiolo bilineari 9 lin. longa, 3 lin. lata. (v. s. in h. Hænke à cl. Sternb. comm.)

168. *E. MAIRETIANUM*, fruticosum, ramis teretibus glabriusculis, foliis oppositis petiolatis ovato-lanceolatis acuminatis minute serrulatis triplinerviis glabris subtus albidis, paniculæ ramis oppositis apice corymbosis, capitulis pedicellatis campanulatis circ. 25-floris, invol. squamis linearibus acutis 1-2-seriatis, achæniis angulatis glabris. $\frac{3}{4}$ in Mexico. Bustamentæ sp. Mair. ! Folia ferè Populi. Petiolus ferè pollicaris; limbus 3 poll. longus, $1\frac{1}{2}$ latus. (v. s. comm. cum pl. aliis Mexic. à cl. Mairet.)
169. *E. BERLANDIERI*, fruticosum glabrum, ramis teretibus striatis, foliis oppositis petiolatis ovatis acuminatis basi et apice integris medio crenas paucas gerentibus utrinquè eglandulosis, paniculæ ramis oppositis apice corymbosis, capitulis pedicellatis circ. 25-floris, invol. squamis subbiseriatis linearibus acuminatis, achænio angulato ad angulos vix scabro, pappo corollâ ferè duplò breviorè. $\frac{3}{4}$ in Mexico inter Tala et Tampico legit cl. Berlandier. Affine *E. ageratifolio* var. β fortè ab α distincta. Petiol. 4 lin. longi; limbi fol. sesquipollic. longi, 9 lin. lati. Cor. albæ. (v. s. comm. à cl. Berland.)
170. *E. GRANDIDENTATUM*, caule herbaceo tenuissimè pulverulento-velutino, ramis oppositis erectis elongatis, foliis oppositis petiolatis ovato-lanceolatis acuminatis basi obtusis subquintuplinerviis grossè dentato-serratis glabriusculis internodio brevioribus, corymbis pluribus fastigiatis trichotomis, capitulis pedicellatis 25-floris, invol. squamis subuniseriatis linearibus acutis, achænio 5-gono vix scabrido. $\frac{2}{3}$ in Mexico legit cl. Luc. Alaman. Petioli 9-12 lin. longi. Limbus fol. 3 poll. longus, $1\frac{1}{2}$ latus. Juxta foliorum distantia. Cor. ex sicco albidæ. Achænia parva nigra. Affinè ex descr. *E. Pazcuarensi*. (v. s.)
171. *E. STERNBERGIANUM*, caule herbaceo ? erecto tereti glabro apice minutè puberulo, foliis oppositis petiolatis latè ovatis vix acuminatis grossè dentatis suprâ glabriusculis subtus petiolisque subpuberulis, corymbis compositis confertis, capitulis pedicellatis 25-30-floris, invol. squamis biseriatis linearibus glabriusculis, achænio pentagono ad angulos scabro. — in Peruvîæ collibus Cordillerianis legit cl. Hænke. Petiol. 5-6 lin. longus, limbus 1-1 $\frac{1}{2}$ poll. longus, 10-18 lin. latus, imâ basi trinerviis, supernè penninerviis. Affine *E. Dombeyano*. (v. s. in h. Hænke. ab ill. com. de Sternberg comm.)
172. *E. DOMBEYANUM*, fruticosum glaberrimum, ramis teretibus, foliis oppositis petiolatis ovatis acuminatis serratis longè suprâ basin triplinerviis, paniculâ tripartitâ, ramis divergentibus ramosis laxis, capitulis longè pedicellatis circ. 25-floris, invol. squamis biseriatis sublinearibus acuminatis vix subciliatis apice ferè scariosis, achænio glabro angulato. $\frac{5}{8}$ in Amer. merid. legit cl. Dombey, sed locus propr. in ejus herb. non indic. Petioli 4 lin. Limb. fol. 2 $\frac{1}{2}$ poll. longus, pollicem latus. Cor. ex sicco purpurascens. (v. s. comm. à Mns. reg. Par.)
173. *E. PANICULATUM* (Schrad. ! ind. sem. h. Gætt. 1832. p. 3. diss. Blum. p. 44). caule herbaceo tereti glabriusculo, foliis inf. et ramis oppositis super. et ramis alternis petiolatis ovato-triangularibus basi truncatis aut breviter cuneatis grossè et obtusè dentatis membranaceis glabriusculis, corymbo laxo foliis parvis intermixto, capitulis pedicellatis circ. 25-floris, invol. squamis oblongo-linearibus mucronatis glabris 1-2-seriatis subæqualibus, achænio angulato glabriusculo. ☉ in Brasiliâ ad sepes circa Bahiam legit cl. Salzmann, etiam vidi ex insulâ Trinitatis ortum. Habitus in sicco *E. Schiedeani* sed ex cl. invent. annum herbaceum. Invol. albida. Cor. minimæ pallidè rubentes. *E. microstemon* Cass. dict. 25. p. 432. Habui etiam sub nomine *E. paniculati* Bernh., sed descr. editam non reperio. (v. s.)
174. *E. SORDESCENS*, fruticulosum, caule erecto tereti cum petiolis pedunculisque velutino-hirtello, foliis oppositis petiolatis ovato-lanceolatis acuminatis dentatis trinerviis suprâ sparse scabris subtus pubescenti-velutinis pallidioribus, paniculæ ramis folio paulò longioribus ultimis corymbosis, capitulis pedicellatis 25-floris, invol. squamis 1-2-seriatis linearibus acutis dorso puberis, achænio secus costas scabrido. $\frac{3}{4}$ in Brasiliâ circa Rio-Janeiro legit cl. Lund. An *E. lauiifolium* Link. enum. h. ber. 2. p. 306 (non Kunth) huc refer. ? Petiol.

9-12 lin. longi; limb. fol. 3 poll. longus, pollicem latus. (v. s. comm à cl. Lund.)

175. *E. RUFESCENS* (Lund in herb. Mus. reg. Berol.), fruticosum, ramis teretibus glabris apice rufo-velutino-hirsutis, foliis oppositis petiolatis ovalibus basi subattenuatis apice longè et angustè acuminatis subintegerrimis penninerviis supra glabris subtus rufis velutino-hirsutis, paniculæ amplæ ramis oppositis polycephalis, capitulis pedicellatis 20-25-floris, invol. squamis circ. biseriatis linearibus acutis dorso puberulis corollis multò brevioribus, achænio glabro. $\frac{5}{8}$ in montosis prov. Rio-Janeiro Brasiliæ legit cl. Lund. Petiol. 1 $\frac{1}{2}$ -2 poll. longi. Limb. fol. 6-7 poll. longus, 2 $\frac{1}{2}$ -3 poll. latus. Cor. ex sicco purpurascens. Pappus sordidus. (v. s. in h. cit.)

176. *E. BREVIPEXA*, fruticosum, ramis teretibus velutino-hispidis, foliis oppositis breviter petiolatis ovali-oblongis acuminatis triplinerviis à medio ad apicem crenatis supra scabris subtus reticulatis cinereo-pubescentibus, nervis petioloque villosis, corymbo terminali conferto vix pedunculo composito, capitulis 20-25-floris pedicellatis, invol. squamis linearibus 1-2-seriatis acutis demum patulis, achænio angulato vix pubero. $\frac{5}{8}$ in Mexico ad Villalpando legit cl. Mendez. Fol. incl. petiolo 1-2 lin., sesquipollicem longa, 5-6 lin. lata. Cor. ex sicco pallidè purpurascens. (v. s. comm. à cl. Mendez.)

177. *E. AZURÆUM*, suffruticosum, ramis teretibus densè villosis-velutinis, foliis oppositis petiolatis ovato-lanceolatis acuminatis excepto apice grossè et inæqualiter dentatis supra glabriusculis subtus petiolisque velutinis subvillosis, corymbo terminali conferto, capitulis pedicellatis 20-25-floris, invol. squamis biseriatis linearibus acuminatis, achænio costato glabriusculo. $\frac{5}{8}$ in Mexico ad Monterey in Nova Leonis prov. legit cl. Berlandier. Cor. cœruleæ. Petiol. 6 lin. longi. Limbus fol. pollicem longus, 8 lin. latus. (v. s.)

178. *E. BUSTAMENTA*, suffruticosum?, ramis teretiusculis apice pilosis, foliis oppositis petiolatis ovatis basi cuneatis apice acuminatis medio dentatis trinerviis supra sparsè puberulis subtus secus nervos villosis, corymbo trichotomo, capitulis ad apices ramorum confertis brevè pedicellatis circ. 20-floris, invol. squamis linearibus subniseriatis mucronulatis margine scariosis, achænio angulato puberulo. — in Mexico. Bustamenta ovata Mairet! in litt. Cor. albidæ. Petioli pollicares. Limbus 28-30 lin. longus, 14-15 lin. latus. (v. s.)

179. *E. PRUNELLÆFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 123), caule herbaceo substriato hirsuto subglutinoso, foliis oppositis petiolatis ovatis acutis serratis utrinquè pilosisculis, capitulis longè pedunculatis terminalibus subcorymbosis multifloris, invol. campanulatis squamis lanceolatis apice subulatis ciliatis, exter. pubescenti-hirtis. $\frac{2}{3}$ in Mexico juxta pagum Ario, in decliv. montium Agnarsacensium et in monte Orizaba. Cor. violacæa? Petiol. 4 lin. Limb. fol. 11 lin. longus, 5-6 latus.

180. *E. PAZCUARENSE* (H. B. et Kunth nov. gen. am. 4. p. 123), caule herbaceo erecto pubescenti-hirsuto, foliis oppositis petiolatis subrotundo-ovatis breviter acuminatis leviter cordatis grossè subduplicato-serratis subtriplinerviis supra scabriusculis subtus in venis pubescentibus, corymbis terminalibus fastigiatis, capitulis longè pedicellatis multifloris, invol. subcampanulatis squamis acutis subciliatis, exter. pubescentibus, achænio ad angulos hispido. $\frac{2}{3}$ in temperatis montanis Mexici ad lacum Pazcuarensem. Cor. alba? Affine *E. ageratoidi* dicitur.

181. *E. VALLINCOLA*, caule suffruticoso erecto tereti, ramulis petiolisque pilos patulos articulatos gerentibus, foliis oppositis longè petiolatis subcordatis acutis ferè triangularibus (angulis inf. obtusis) basi trinerviis grossè dentatis utrinquè sparsè hispidulis, capitulis 5-13 in corymbum compositum fastigiatum dispositis pedicellatis 20-25-floris, invol. squamis linearibus 1-2-seriatis disco brevioribus, achæniis pentagonis ad angulos pubero-scabris. $\frac{5}{8}$ in vallibus Andium Peruvix legit cl. Hænke. Petioli circ. 10-11 lin. longi. Limbus fol. pollicem longus, 10-12 lin. latus. Cor. ex stæce purpurascens. (v. s. in h. Hænke à cl. de Sternberg comm.)

182. *E. PAUPERATUM* (H. B. et Kunth nov. gen. am. 4. p. 121), caule suffruticoso? tereti piloso, foliis oppositis petiolatis triangularibus acutis basi truncatis grossè crenato-serratis subtrinerviis suprâ pilosisculis subtus molliter piloso-pubescentibus, corymbo terminali simplici, capitulis pedicellatis multifloris, invol. campanulati squamis lanceolatis apice subulatis striatis ciliatis, ext. paulò brevioribus, achænio glabro. $\frac{3}{4}$ in Novâ Granatâ? Fortè idem ac *E. microphyllum* L.

** *Capitulis 20-floris.*

183. *E. NERVOSUM* (Swartz! prod. iii. flor. 1307), fruticosum, ramis teretibus erectis, ramulis pubescentibus, foliis oppositis petiolatis ovatis acuminatis grossè et acutè dentatis triplinerviis rugoso-nervosis glabris subtus globuliferis, corymbo composito polycephalo fastigiato, capitulis pedicellatis 20-floris, invol. squamis linearibus acuminatis 8-10 subuniseriatis. $\frac{3}{4}$ in altis montibus Jamaicæ (Sw.), Sancti Domingi (Mack. in h. Lindl.!) Ramuli rubentes. Petioli puberi 3 lin. longi. Fol. 2 poll. longa, 10-12 lin. lata, infrâ nervos 2 laterales, nervulos 2 minores è nervo medio emittentia. Capitula parva. Cor. albida. Achænia minima. (v. s. ex h. Swartz.)
184. *E. PAPANTLENSIS* (Less. in linnæa 1831. p. 403), fruticosum, ramis teretibus junioribus pubescentibus, foliis oppositis membranaceis triplinerviis ovatis basi cordatis acuminatis serrato-crenatis non glabris, paniculis fastigiatis myriocephalis, capitulis pedicellatis 20-floris, invol. campanulati squamis bisseriatis linearibus acuminatis glabris, corollis tubulosis. $\frac{3}{4}$ in Mexico prope Papantla. *E. leiophyllo* proximum. Petioli, pubes foliorum, achænia indescr. $\frac{1}{2}$.
185. *E. AREOLARE*, fruticosum, ramis teretibus petiolis pedunculis nervisque foliorum subtus pube rufâ brevi confertâ velutinis, foliis oppositis petiolatis ovato-lanceolatis acuminatissimis rigidis basi obtusis 3-nerviis suprâ glabris minutissimè areolatis subtus puberis reticulatis, pedunculis axillaribus oppositis et subterminalibus trichotomis folio brevioribus, capitulis brevissimè pedicellatis 20-floris, invol. squamis 1-2-seriatis linearibus acutis, achænio pubescente. $\frac{3}{4}$ in Mexici montibus Guachilacensibus specimina paucâ legit cl. Berlandier. Petioli 3 lin. longi. Fol. limb. 30 lin. longus 12-14 latus. Cor. ex sicco albida. (v. s.)
186. *E. BREVISSETUM*, fruticosum, ramis teretibus densè petiolisque velutino-hispidis, foliis oppositis petiolatis ovato-lanceolatis acuminatis serratis trinerviis suprâ sparse scabris subtus pubescenti-subvelutinis glandulosisque, paniculis axillaribus terminalibusque folio brevioribus oppositè ramosis, capitulis pedicellatis circ. 20-floris, invol. squamis linearibus acutis 1-seriatis dorso puberis, pappi setis corolla dimidiò brevioribus, achænio glabro. $\frac{3}{4}$ in Mexico. Genus novum Alam! in litt. 1831. Mair! exs. 1833. Cor. albæ invol. duplò longiores. Petioli 4 lin. longi; limbus 3 poll. longus, sesquipollicem latus. (v. s.)
- β . foliis suprâ confertè scabris subtus velutino-tomentosis. — in Mexico. (v. s.)
187. *E. LÆVE*, fruticosum glaberrimum, ramis teretibus substriatis, foliis oppositis petiolatis ovato-lanceolatis acuminatis apice obtusè serratis triplinerviis utrinquè lævigatis, paniculis ramosissimis ramos terminantibus, capitulis pedicellatis 20-floris, invol. squamis 1-2-seriatis linearibus acuminatis. $\frac{3}{4}$ inter frutices circa Rio-Janeiro leg. cl. Lund Gandichaud et Lhostky. Frutex 10-12 pedalis. Rami junior. subrubentes, adulti albidii. Petiol. 3-5 lin. longi. Folia 3-5 poll. longa, 9-15 lin. lata. Pappus stramineus corollæ longit. Achæn. angulatum glabrum. An fortè *E. Freyresii* Thunb. pl. bras. dec. 2. n. 19. ex florâ 1821. p. 382? sed descr. non suff. (v. s.)
188. *E. AYAPANA* (Vent. ! malin. t. 3. mag. enc. an VIII. v. 3. p. 76), caule basi suffruticoso ascendente ramoso glabro, foliis subsessilibus oppositis lanceolatis triplinerviis acuminatis subintegerrimis glabris, corymbo laxo oligocephalo, capitulis pedicellatis circ. 20-floris, invol. subuniseriatis squamis

- linearibus acuminatis inæqualibus dorso puberulis. $\frac{5}{8}$ in America merid. ad ripam fl. Amazonum dexteram (Vent.), undè migravit in ins. Sanctæ Crucis (Vahl), ins. Mauritanias (Petit-Th.), Javam (Blam.) etc. Petit-Th. obs. in dict. Lam. p. 9. cum ic. Tratt. thes. t. 16. Eup. triplinerve Vahl! symb. 3. p. 97. Foliorum infusio est eximium sudorificum et ideò alexipharmacum in utrâque Indiâ sub Ayapanæ nomine celeberrimum, undè nomen Vent. etiamsi recentius Vahliano plerisque Eupatoriis convenienti selegi. (v. s. et olim v. v.)
189. *E. LUNDIANUM*, suffruticosum, ramis teretibus subtiliter striatis puberis apice hirsutis, foliis oppositis petiolatis ellipticis utrinquè acuminatis medio grossè crenatis basi et apice integris penninerviis suprâ scabriusculis subtus hirsutis, paniculæ ramis oppositis hirsutis vix folio longioribus apice confertè subcorymbosis, capitulis vix pedicellatis 20-floris, invol. squamis 2-3-seriatis dorso puberis, intim. obtusissimis corollis dimidio ferè brevioribus, achænio angulato glabriusculo. $\frac{5}{8}$ in Brasilæ collibus apricis circa Rio-Janeiro mense august. legit cl. Lund. Petiol. 4 lin. Fol. limbus 2-2 $\frac{1}{2}$ poll. longus, pollicem circ. latus. (v. s. comm. à cl. Lund.)
190. *E. HIRSUTUM*, suffruticosum, ramis teretibus petiolis pedunculis folisque subtus densè hirsutis, foliis imis oppositis super. alternis confertis breviter petiolatis oblongo-lanceolatis utrinquè acutiusculis crenato-subdentatis suprâ sparsè puberulis penninerviis, corymbo terminali composito conferto, capitulis pedicellatis circ. 20-floris, invol. squamis subuniseriatis glabriusculis linearibus subacutis, achænio 5-gono secus angulos scabro. $\frac{5}{8}$ in Mexico legit cl. Hænke. Folia (incl. petiolo 1-2 lin.) ferè 2 poll. longa, 6 lin. lata (v. s. in h. Hænke à cl. Sternb. comm.)
191. *E. PYCNOCEPHALUM* (Less. in. linnæa 1831. p. 404), caule herbaceo tereti erecto, foliis... petiolatis membranaceis triplinerviis ovatis s. rhomboideo-ovatis utrinquè acuminatis serrato-crenatis glabris, paniculis trichotomis laxis, capitulis circ. 20-floris congestis brevissimè pedicellatis, invol. campanulati squamis glabris obtusissimis, ext. minoribus mucronulatis. — ad Papantla Mexicanorum. Affine *E. prunellæfolio*.
192. *E. CILIATUM* (Less. in linnæa 1831. p. 404), caule herbaceo erecto, foliis oppositis petiolatis membranaceis triplinerviis ovatis acutis serrato-crenatis subsetuloso-ciliatis sparsè et rigidè pilosis, paniculæ fastigiatæ ramis hispidiusculis, capitulis 18-22-floris pedicellatis, invol. campanulati squamis subbiseriatis linearibus subscariosis apice ciliatis, achænio ad angulos hirsuto. — in Mexici regione subfrigida inter San Miguel del Soldado et la Joya. Fol. circ. pollicaria.
193. *E. BERTERIANUM* (Colla! act. Acad. Taur. 33. p. 130. t. 8. app. hort. rip. 3. p. 18. t. 9), herbaceum glabrum, caule erecto tereti ramoso, foliis alternis petiolatis ovatis acutis crenatis, paniculis axillaribus terminalibusque diffusis, invol. squamis linearibus acutiusculis circ. 15 subbiseriatis, achænio angulato glabro. ① in Guadalupâ legit cl. Bertero. Mikania Berteriana Spreng! syst. 3. p. 423. Folia infima fortè opposita. Herba pedalis. (v. s.)
194. *E. GUADALUPENSE* (Spreng. syst. 3. p. 414), herbaceum glabriusculum erectum, caule tereti apice alternè ramoso, foliis infer. oppositis super. alternis petiolatis rhombeo-ovatis trinerviis crenatis acuminatis suprâ sparsè subtus in nervis subpuberulis, paniculâ laxâ, capitulis pedicellatis circ. 20-floris, invol. squamis biseriatis ext. brevibus, inter. longioribus submembranaceis obtusis submucronatis, achænio glabro. ① in Guadalupâ legit cl. Krauss. E. violaceum Bert. ex Spreng. Petioli pollicem longi. Fol. limb. 2 poll. longus, sesquipollicem latus. Cor. violacæ. (v. s.)
- β. Wydlerianum*, foliis minoribus acutius dentatis. ① in Porto-Ricco legit cl. Wydler. (v. s.)
- *** Capitulis 10-20-floris.
195. *E. SOPHIOIDES*, herbaceum, caule adscendente simplici basi folioso hirsuto apice subnudo glabriusculo, foliis oppositis petiolatis pubescentibus bipinna-

tisectis, lobis inaequalibus obovatis oblongisve, ultimis longioribus sinuato-dentatis, corymbo simplici oligocephalo, capitulis pedicellatis 15-20-floris, invol. squamis linearibus obtusis 10-12-uniseriatis persistentibus demum reflexis, achænio angulato puberulo. ① in Sancto-Domingo legit cl. Bertero. *Mikania sophiæfolia* Spreng. syst. 3, p. 423. *Eupat. sophiæfolium* auct. sed differt ab *icone Plumieri* (speciei archetypo) involucri nullo modo imbricatis et foliorum divisuris. (v. s.)

196. *E. GONOCADUM*, fruticosum glabrum, ramulis sexangulatis, foliis oppositis petiolatis rhombeo-lanceolatis basi cuneatis integerrimis apice acuminatis serratis rigidis supra basin triplinerviis, pedunculis axillaribus et terminalibus folio brevioribus capitula plurima pedicellata corymbosa 15-20-flora gerentibus, invol. squamis circ. 2-seriatis ext. angustis acutis, int. oblongis obtusis longioribus sed flosculis dimidiò brevioribus, achænio ad angulos scabro. 3 circa Mexico (h. Lindl.). Petioli 3-4 lin. longi. Limbus fol. sesquipoll. longus, 7-8 lin. latus. Affine *E. ligustrino* et glabrato. (v. s. in h. Lindl.)

197. *E. ERIGEROIDES*, fruticosum, ramis oppositis et alternis teretibus junioribus tenuissimè puberulis, foliis oppositis et alternis petiolatis subnerviis lanceolato-linearibus acuminatis integerrimis subpuberulis, paniculis terminalibus laxis, capitulis pedicellatis 15-floris, invol. squamis circ. 10 subbiseriatis dorso puberis, inter. longioribus oblongis apice submembranaceis obtusis ciliatis, achænio subscabrido angulato. 3 in Brasiliæ prov. Rio-Grande. Petioli 3-4 lin. longi. Limb. fol. pollicem longus, 2-3 lin. latus. Cor. ex sicco pallidæ. Affine *E. steviæfolio*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 809 miss.)

ℓ. *ramulosum*, ramis pluribus brevioribus, foliis linearibus hinc indè subdentatis, capitulis 15-20-floris. 3 in Brasiliæ prov. Rio-Grande. Folia 1 $\frac{1}{2}$ lin. lata. (v. s. cum var. α sub n. 779.)

198. *E. VIBURNOIDES*, fruticosum, ramis teretibus junioribus pedunculis petiolis nervoque medio fol. araneoso-subtomentosis, foliis oppositis breviter petiolatis latè ovalibus utrinquè subacutis à mediò ad apicem argutè et rariter serratis supra glabris subtis villosis hirtellis, corymbo trichotomo terminali vix exserto, capitulis pedicellatis 15-floris, invol. squamis biseriatis subæqualibus linearibus obtusis, achænio pentagono hispidulo. 3 in Mexico inter Victoria et Tula legit cl. Berlandier. Petioli 3 lin. longi. Limbus fol. 3-3 $\frac{1}{4}$ poll. longus, 2-2 $\frac{1}{2}$ latus. Invol. et coroll. purpurascens. (v. s.)

199. *E. CALAMINTEÆFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 129. t. 343), fruticosum, ramis teretibus pulcherrime pubescentibus, foliis brevissimè petiolatis ovato-subrotundis obtusis dentibus 2-3 vix utrinquè notatis aut subintegerrimis supra glabriusculis subtis puberulis punctulatis subtrinerviis, ramis lateralibus apice corymbulosis in paniculam foliosam dispositis, capitulis pedicellatis 14-15-floris, invol. squamis linearibus acuminatis apice subscariosis, achænio angulato pilosulo. 3 in Mexici montibus Guanajuatensibus. Cor. alba. Invol. squamæ paulò longiores et magis acuminatæ quàm in *icone citatâ*. (v. s. in h. Acad. reg. Monac.)

ℓ. *pluridentatum*, foliis latius ovatis dentibus utrinquè 7-8 notatis, petiolo paulò longiore. paniculâ confertiore. 3 in Mexico legit cl. Alaman. An species propria? (v. s.)

200. *E. TETRAGONUM* (Schrad. cat. sem. h. Goett. 1830 et in litt. 1832), caule suffruticoso debili erectiusculo glabro, ramis oppositis tetragonis vix apice pubescentibus, foliis oppositis petiolatis ovato-lanceolatis acuminatis dentato-serratis trinerviis glabriusculis, paniculâ thyrsoidæ, capitulis pedicellatis 12-15-floris, invol. squamis subuniseriatis linearibus acuminatis dorso pubescentibus demum patulis, achænio angulato puberulo, pappo corollâ ferè dimidiò brevior. 2ℓ. 3 in Mexico (Schrad.). Flores albi. Petioli pollicares. Limb. fol. 3-4 poll. longi, 1 $\frac{1}{4}$ -2 $\frac{1}{2}$ lati. Caulis in solo fertiliori usquè ad 10-12 pedes longus, digitum crassus. (v. s. cult. in hort. Goett. à cl. Schrader. comm.)

201. *E. villosum* (Swartz! prod. III. flor. 1305), fruticosum, ramis teretibus ferrugineo-velutinis, foliis oppositis petiolatis ovatis acutis subserratis trinerviis supra scabridis subtus ferrugineo-hirsuto-subtomentosis, corymbo trichotomo fastigiato polycephalo, capitulis pedicellatis 8-16-floris, invol. squamis 8-10 uni- seu biseriatis linearibus subpatulis pubescentibus, achænio angulato puberulo. ζ in frutetis montosis Jamaicae (Sw.!), Sancti-Domingi? (Willd. sp. 3. 1763? excl. syn. Sloan.). Hirsuties tota ferruginea. Petioli 2-3 lin. longi hirsuto-velutini; limb. fol. 12-15 lin. longi, 6-7 lati. Capitula parva. Cor. albidæ. (v. s. ex h. Swartz.)
202. *E. orbiculatum*, caulibus fruticulosius erectis simplicibus teretibus glabris, foliis oppositis brevissimè petiolatis orbiculatis dentatis basi trinerviis internodiis longitudini plerisque subæqualibus utrinquè glabriusculis, corymbi fastigiati ramis oppositis oligocephalis, capitulis pedicellatis 12-floris, invol. squamis 1-2-seriatis acutis oblongo-linearibus dorso subpuberulis. ζ \mathcal{L} in Brasiliæ prov. Sancti-Pauli. Petioli lineam longus. Limb. fol. diam. 9-10 lin. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 507 miss.)
203. *E. puberulum*, suffruticosum, ramis teretibus tenuissimè puberulis, foliis oppositis petiolatis ovato-lanceolatis subacuminatis grossè dentatis apice integris basi trinerviis supra glabriusculis subtus petiolisque tennè velutino-puberulis, paniculæ ramis apice ramulosis, capitulis ad-apices ramulorum confertis sessilibus circ. 15-floris, invol. squamis linearibus acutis 1-2-seriatis dorso subpuberulis, achænio breviter hispidulo. ζ in Sancto-Domingo legit cl. Bertero. B. molle Bert.! herb. non Sw. nec Kunth. *E. salicinum* ex Hispaniâ Spreng. syst. 3. p. 412? excl. syn. Petioli 6 lin. longi, limb. fol. 2-2 $\frac{1}{2}$ poll. longus, pollicem latus. Cor. ex sicco albidæ. (v. s. comm. à cl. Bertero.)
204. *E. rhomboideum* (H. B. et Kunth nov. gen. am. 4. p. 127), fruticosum, ramis striatis glabris ramulis pubescenti-hirtis, foliis oppositis petiolatis rhomboideo-ovatis obtusiusculis crenatis subtriplinerviis supra pilosiusculis subtus glabris, corymbis terminalibus, capitulis pedicellatis, invol. campanulatis squamis oblongo-lanceolatis acutis hirtio-pubescentibus. ζ in frigidis Mexici inter Toluca et Tianguillo. Petioli 4 lin. Limb. fol. 15 lin. longus, 10 latus.
205. *E. triangulatum* (Alam. in litt. 1831), caule herbaceo erecto simplici tereti purpureo glabro, foliis oppositis petiolatis ovato-triangularibus apice acuminatis basi truncatis integris, angulis lateralibus rotundatis, lateribus dentato-serratis rigidis trinerviis reticulatis glabriusculis, paniculæ subcorymbosâ, pediculis pubescentibus, capitulis breviter pedicellatis 12-floris, invol. squamis subuniseriatis linearibus acutis, pappo corollâ dimidiò breviorè. \mathcal{L} à Mexico misit cl. Luc. Alaman cum plur. aliis rarissimis. Ex deser. valdè accedit ad *E. rubricaulæ*, sed caulis teres nec hexagonus, fol. venæ obscuræ nec pellucidæ, folia rigidula nec membranacea, invol. non imbricata, etc. Cor. albæ invol. duplò longiores. (v. s.)
206. *E. cubense*, fruticosum, ramis teretibus apice pubescenti-velutinis, foliis oppositis petiolatis ovato-lanceolatis basi subtruncatis apice acuminatis subdentatis trinerviis supra glabris subtus pube subvelutinâ canescentibus glandulosisque, corymbo trichotomo fastigiato, capitulis 10-12-floris, invol. squamis 8-10 linearibus subpatulis pubescentibus, achænio angulato puberulo. ζ in ins. Cubâ prope Havanam legit cl. Ramon de la Sagra. Petioli hirsutè brevi canescentes, 2-3 lin. longi. Limb. fol. 15-16 lin. longi, 7-8 lati. Inflor. rescentia omninò *E. villosi*. (v. s. comm. à cl. de la Sagra.)
207. *E. cynanchifolium*, fruticosum, ramis teretibus apice tenuissimè pulverulento-velutinis, foliis oppositis breviter petiolatis ovatis subcordatis obtusis trinerviis reticulato-venosis supra glabriusculis subtus pulverulento-velutinis, corymbo trichotomo fastigiato, capitulis pedicellatis 10-12-floris, invol. squamis 8-10 subbiseriatis pubescentibus subacutis, achænio subangulato vix puberulo. ζ in ins. Cubâ juxta Havanam legit cl. Ramon de la Sagra. Folia cinereo-subcanescentia petiolo 1-2 lin., limbo pollicem longo, 8 lin. lato. *Inflor.* *E. cubensis*. Cor. albæ. (v. s. comm. à cl. de la Sagra.)

208. *E. LRIOPHYLLUM* (Less. in Linnæa 1831. p. 402), fruticosum, ramis teretibus junioribus pubescentibus, foliis oppositis membranaceis triplinerviis ovatis acuminatis basi obtusissimis profunde serrato-crenatis glaberrimis, paniculis subfastigiatis myriocephalis, capitulis pedicellatis 12-floris, invol. campanulati squamis biseriatis linearibus acuminatis glabris. ♂ in Mexico prope Papanla. Folia 12-21 lin. longa, 9-15 lin. lata, an sessilia? an petiolata? Affine *E. micrantho* dicitur, sed paniculæ ramî primaria folia sua superantes, corollæ limbo amplo tubum superante.

209. *E. LUCIDUM* (Orc. dec. 35), fruticosum, ramis teretibus, junioribus breviter glanduloso-hirsutis, adultis glabris, foliis oppositis petiolatis ovatis utrinque breviter acuminatis integerrimis aut hinc inde serratis, supra sparse subscabridis, subtus præsertim secus nervos et petiolum hirtellis, corymbo terminali conferto, capitulis pedicellatis 10-12-floris, invol. squamis biseriatis linearibus subobtusis extus subhirtellis, achænio angulato glabro glanduloso. ♂ in Mexico (Orc.). Specimen habeo sub Ortegiano nomine à cl. Néé lecto, sed locus non indic. Folia supra vix in sicco lucentia. Affine *E. calaminthæfolio*. (v. s.)

210. *E. OBTUSISSIMUM*, fruticosum, ramis teretibus hirsuto-villosis, foliis oppositis breviter petiolatis subcordatis ovato-rotundis obtusis supra minutè velutinis subtus velutino-tomentosis subtriplinerviis reticulatis, corymbo terminali subcesili conferto, capitulis breviter petiolatis 10-12-floris, invol. squamis linearibus subacutis puberis 2-seriatis subæqualibus, achænio angulato glabro. ♂ in Sancto-Domingo legit cl. Bertero. *E. villosum* Bert. herb. non Willd. *E. domingense* Spreng. syst. 3. p. 412? (etiãmsi folia alterna dicat). Petioli 2-3 lin. longi. Limb. fol. sesquipoll. longus latusque (v. s. comm. à cl. Balbis.)

211. *E. INCOMPTUM*, caule herbaceo erecto teretiusculo villoso-subtomentoso, foliis alternis fere sessilibus basi longè cuneatis ovatis acutis triplinerviis supra scabridis subtus villoso-tomentosis, paniculæ elongatæ ramis alternis apice racemoso-paniculatis, capitulis ad apices ramulorum confertis brevè pedicellatis 10-12-floris, invol. squamis 1-2-seriatis linearibus obtusis pubescentibus, achænio angulato glabriusculo. — in Mexico legit cl. Hænke. Fol. infer. ignota, an opposita? media 3 poll. longa, sesquipollicem lata; super. longè acuminata. (v. s. in h. Hænke à cl. Sternberg comm.)

**** *Capitulis 6-10-floris.*

212. *E. AGERATIFOLIUM*, fruticosum glabrum, ramis teretibus, foliis oppositis petiolatis latè ovatis basi subtruncatis apice attenuatis obtusis grossè dentatis trinerviis eglandulosis, corymbis terminalibus trichotomis, pedicellis subviscosis vix puberulis, capitulis circ. 10-floris, invol. squamis linearibus biseriatis acutiusculis apice subciliatis, demùm patulis, achænio puberulo ad angulos subscabrido. ♂ in insulâ Cubâ circa Havanam legit cl. R. de la Sagra. *E. Havanense* H. B. et Kunth nov. gen. am. 4. p. 128? sed pedicelli non verè glabri, pedunculi trichotomi nec trifidi, corollæ pallidæ nec violacæ, involutum vix imbricatum, etc. — Petioli 9 lin. longi; limb. fol. 24-30 lin. longus, 20-22 latus. (v. s. comm. à cl. de la Sagra.)

♂? *Mexicanum*, foliis magis ovatis basi subobtusis medio crenato-dentatis, pedicellis subpubescentibus, capitulis 12-13-floris. ♂ in Mexico inter Victoria et Tula legit cl. Berlandier. *E. Berlandieri* formâ foliorum fere accedit, sed pappus corollam adæquans.

♀? *Domingense*, foliis magis ovatis basi obtusis crenato-dentatis, pedicellis subpubescentibus, capitulis 18-20-floris. ♂ in Sancto-Domingo legit Bertero. *E. ageratoides* Bert. herb. non Linn. Foliorum limbus sesquipollicem longus poll. latus. Invol. squamæ acutiores. An species propria? (v. s. comm. à cl. Balbis.)

213. *E. CELTIDIFOLIUM* (Lam. dict. 2. p. 406), fruticosum glabrum, ramis teretibus striatis, foliis oppositis petiolatis ovato-lanceolatis basi obtusis apice acu-

- minatissimis subserratis basi subquintuplinerviis, paniculæ amplæ ramis oppositis polycephalis, capitulis pedicellatis 10-floris, invol. squamis 7-9 subuniseriatis linearibus acuminatis apice serraturis 1-2 notatis demum patulis, achenio subpuberulo. 5 in ins. Caribæis (Lam.), in montibus Guadalupæ (Perrottet!) et in Cubâ circa Havanam (La Ossa!) Mikania verrucosa Spreng. syst. 3. p. 423? Verrucæ sunt lenticellæ omnibus Eupat. lignosis solemnes. Fol. 4-5 lin. longa, $1\frac{1}{2}$ lata, petiolo 4 lin. longo. Cor. parvæ albidæ. (v. s.)
214. E. SCHRADERI, frutescens, ramis teretibus striatis pubescenti-hirtellis, foliis infer. oppositis rameis alternis brevissimè petiolatis oblongo-lanceolatis basi cuneatis integerrimis à medio dentatis suprâ sparsè scabridis subtus pubescenti-hirtellis triplinerviis, corymbis terminalibus laxis, capitulis pedicellatis circ. 10-floris, invol. squamis subuniseriatis erectis lanceolatis acuminatis, achenio angulato puberulo. 5 in Americâ merid. sed locus proprius ign. E. rugosum Schrad. l. pl. rar. h. Gœtt. ined. ex litt. 1832. non Kunth nec Sprengel. Folia subrugosa 3 poll. longa, 6-9 lin. lata. Cor. albæ. (v. s. cult. ex h. Gœtt. à cl. Schrad. comm.)
215. E. VOLIOLOSUM, caule suffruticoso tereti vix tenuissimè pulverulento-velutino, foliis oppositis (alternisque?) petiolatis ellipticis crenatis subtus reticulatis utrinquè cinereo-subvelutinis, paniculæ elongatæ ramis brevibus oligocephalis, capitulis pedicellatis 9-10-floris, invol. squamis linearibus obtusiusculis subuniseriatis pubescentibus, achenio angulato scabrido. 5 in Chili circa Coquimbo leger. cl. Gandichaud et Macrae. Petioli 2-3 lin. longi. Limb. fol. 3 lin. longus, circ. 2 lin. latus. Folia ramea ad axillas conferta. Ramo alterni. (v. s.)
216. E. DIFFUSUM (Vahl symb. 3. p. 94?), caule herbaceo erecto tereti vix pubescente, foliis oppositis petiolatis basi cuneatâ cordato-ovatis acuminatis obtusè et sæpè duplicato-serratis membranaceis basi trinerviis glabris subtus crebrè glandulosis, paniculæ diffusæ ramosissimæ ramis inf. oppositis sup. alternis, pedicellis longis capillaribus, capitulis 7-8-floris, invol. squamis 1-2-seriatis linearibus acuminatis nitidis subscariosis demum reflexis, achenio teretiusculo pubescente. — in Americâ merid. (Vahl), prov. Santa-Marta (Bert.), ad flav. Magdalene (Spreng. syst. 3. p. 419), in sylvis Jamaicæ interioris (Swartz flor. Ind. occ. 1301), in Hispaniolâ (Ham.). E. capillare Desv. in Ham. flor. Ind. occ. p. 51. E. flaccidum Spreng. in litt. ad Balbis. Species habitu distinctissima sed corollæ ignotæ. An fortè huc species duæ confusæ? (v. s. comm. à cl. Bert. et Balb.)
217. E. POLYBOTRYUM, caule fruticuloso tereti ramosissimo tenuissimè cum pedunculis petiolis nervisque fol. pulverulento-velutinis, foliis oppositis rarius ternis et super. alternis petiolatis ovatis acuminatis subdentatis penninerviis glabriusculis, ramis axillaribus racemoso-paniculatis in paniculam longam foliosam dispositis, capitulis pedicellatis 6-8-floris, invol. squamis paucis 1-2-seriatis oblongo-ovalibus acutiusculis, achenio vix pubente. 5 in Mexico legit cl. Haenke. Petioli 4-6 lin. longi. Limb. fol. 30 lin. longus, circ. 20 lin. latus. Capitula parva innumerosa. Cor. ex sicco albidæ. Styli valdè elongati. Achenium exangulatum. (v. s. in herb. Haenke à cl. Sternb. comm.)
218. E. AURICULATUM (Vahl symb. 3. p. 95. t. 72. non Lam.), fruticosum, ramis teretibus striatis tomentoso-canis demum calvis, foliis alternis petiolatis ad basin petioli utrinquè rotundè auriculatis ovatis acutis denticulatis suprâ glabris subtus cano-tomentosis, spicis plurimis paniculatis elongatis, capitulis sessilibus distantibus 7-floris, invol. squamis 8 fusco-ferrugineis basi squamulis parvis calyculato. 5 in Brasiliâ.
219. E. ORGYALE, herbaceum altissimum, ramis hexagonis vix apice summo puberulis, foliis oppositis petiolatis latè ovato-lanceolatis basi cuneatis apice acuminatis grossè dentatis penninerviis reticulatis suprâ sparsè scabridis subtus pubescenti-hirtellis et glandulosis, corymbo trichotomo terminali confertissimo, capitulis ad apices ramulorum densè adgregatis sessilibus 6-floris, invol. squamis 2-ser. oblongis obtusis ext. brevioribus subpubescentibus, int.

longioribus glabris. $\frac{1}{2}$ circa Bahiam (Blanch. ! n. 1923), in montosis circa Rio-Janeiro frequens ex cl. Lund! Herba 6-7-pedalis. Rami oppositi. Flor. albi odoratissimi insecta odore allicientes. Invol. albida. (v. s.)

220. E. PIQUERIODES, caule herbaceo tereti glabro, ramulis oppositis apice petiolisque puberulis, foliis oppositis petiolatis lanceolatis acuminatis trinerviis remotè serratis secus nervos subtus puberulis cæterum glabris, paniculis laxis oppositirameis ramos terminantibus, invol. 5-phyllis, floribus circ. 6-7, styli ramis exsertis apice crassè clavellatis. — in Peruviae montibus Guanoccensibus legit cl. Haenke. Simillimum Piqueriae trinerviae sed folia subtus secus nervos hirtella et car. gener. diversi. (v. s. in h. Haenke ab ill. de Sternberg comm.)

**** Capitulis 8-20-floris. — *Herbæ Boreali-Americæ, foliis petiolatis oppositis aut in unicâ specie alternis.*

221. E. CORDATUM (Walt. car. 199), caule tereti tenui-pubescente, foliis oppositis breviter petiolatis ovatis subcordatis grossè et obtusè dentatis triplinerviis supra sparsè et subtus secus nervos scabridis, corymbo composito, capitulis 12-20-floris, invol. squamis circ. 10 lanceolatis subæqualibus vix pubescentibus, achænio subangulato glabro. $\frac{1}{2}$ in siccis Carolinæ (Michx. ! Bosc !), Novæ-Cæsareæ (Green. !). Eop. aromaticum Ell. sketch 2. p. 304. non Lin. E. cordiforme Poir. suppl. 2. p. 600. E. melissoides Willd. spec. 3. p. 1754? sed capit. dicuntur 5-6-flor. Cor. albæ odoratæ. Petioli 2 lin. longi. (v. s.)

ß. Fraseri, foliis ovatis paulò longioribus nec basi subcordatis. $\frac{1}{2}$ in Carolina. E. Fraseri Poir. suppl. 2. p. 600. Lam. ill. t. 672. f. 4. (v. s. ex herb. Fraser ex quo etiam Lam. et Poir. habuerunt.)

222. E. AROMATICUM (Lin. sp. 1175), caule tereti pubescente, foliis oppositis petiolatis ovatis acuminatis trinerviis grossè et inæqualiter serratis scabriusculis, corymbo amplo subpaniculato, capitulis circ. 20-floris, invol. squamis 10-12 linearibus acutis æqualibus, achænio angulato glabro. $\frac{1}{2}$ in Virginia (Lin.), Novâ-Cæsareâ (Torr. !), à Pensylvaniâ ad Floridam (Pursh) Willd. sp. 3. p. 1765. — Pluk. alm. t. 88. f. 3. Cor. albæ. Styli breviter exserti. (v. s.)

223. E. CRANTHIFOLIUM (Muhl. in Willd. sp. 3. p. 1755), caule tereti glabriusculo, foliis oppositis petiolatis ovatis acuminatis dentatis triplinerviis subglabris, corymbo subconferto, capitulis 5-10-floris, invol. squamis 10 linearibus subæqualibus, achænio angulato glabro. $\frac{1}{2}$ in pinguibus Carolinæ ad Paris island prope Beaufort (Ell. sketch 2. p. 303), à Novæboraco ad Virginiam in rupibus (Pursh fl. 2. p. 514). An stirps Purshiana eadem ac Elliotiana? (v. s.)

224. E. AGERATOIDES (Lin. f. suppl. 355), caule tereti glabro, foliis oppositis longè petiolatis latè ovatis acuminatis trinerviis inæqualiter et grossè serratis glabris, corymbo composito, capitulis circ. 20-floris, invol. squamis linearibus acutis æqualibus, achænio angulato glabro. $\frac{1}{2}$ in Amer. bor. Canadâ (Michx.) Virginiâ (Willd.) Carolinâ (Ell.) Pensylvaniâ et montibus Alleghany (Pursh). E. urticæfolium Reich. syst. 3. p. 719. Michx. fl. bor. 2. p. 100. E. altissimum Lin. syst. veg. 614. Ageratum altissimum. Lin. sp. 1176. — Corn. can. t. 21. Mor. ox. s. 7. t. 18. f. 11. Cor. albæ. Styli longè exserti. Petiol. 1-2 poll. longi. Fol. interdum subcordata. (v. s.)

225. E. SEROTINUM (Michx. fl. bor. am. 2. p. 100), caule tereti pulverulento-pubescente, foliis oppositis petiolatis ovato-lanceolatis superne attenuatis acutis grossè et acutè serratis triplinerviis pubescentibus, corymbo composito, capitulis 10-12 floris, invol. squamis 10 linearibus puberis subimbricatis, achænio angulato. $\frac{1}{2}$ in vallibus Carolinæ maritimæ (Michx.). Ell. sketch 2. p. 304. Folia non rariter serrata ut vult Michaux, sed serraturis grossis plurimis utrinquè notata. Cor. albæ. (v. s. c. ex sem. Michaux in h. L'Hér.)

226. E. INCARNATUM (Walt. flor. car. 200. Ell. ! sketch 2. p. 305), caule tereti pubescente, foliis oppositis longè petiolatis ovatis basi truncatis subcordatisve

acuminatis grossè et obtusè dentatis triplinerviis glabriusculis, corymbi laxi ramis subracemosis, capitulis 20-floris, invol. squamis circ. 15-20 linearibus acutis biserialibus subæqualibus, achænio angulato glabro. ♀ in pinguibus Carolinæ. Cor. purpureæ. Petioli 1-2 poll. longi. Affin. E. cœlestino sed certè distinctum. An fortè E. violaceum Raf. fl. Ind. 62? (v. s.)

227. E. RESINOSUM (Torr. in litt. 1835), caule tereti erecto brevissimè velutino, foliis oppositis sessilibus elongato-lineari-lanceolatis acuminatis serratis basi obtusis supernè glabriusculis subtus velutino-tomentosis, corymbo composito fastigiato polycephalo, invol. squamis ovalibus obtusis dorso velutino-villosis, flor. 10-12, achæniis resinoso-muricatis. ♀ in Novâ-Casariâ (Torr. Green.). Habitus potiùs sect. sequentis sed ex numero florum huc referendus. Flores albi. Styl. modicè exserti. (v. s.)

228. E. OCCIDENTALE (Hook. fl. bor. am. r. p. 305), glabriusculum, caule striato, foliis alternis brevè petiolatis ovatis acuminatis grossè inæqualiter serratis apice integerrimis, summis lanceolatis integerrimis, capitulis paniculato-corymbosis, invol. pluriflori squamis 1-serialibus acutis laxis, achæniis linearibus angulatis. — in collibus lapidosis inter flum. Lewis et Clark Amer. bor. occid. Flores albi.

ℓ. *subroseum*, caule tereti, corollis apice stylisque roseo-purpurascens. — ad flum. Columbia (Douglas). Flor. in invol. 15-20. (v. s. comm. ab hom. Soc. hortie. Londin.)

***** Capitulis 5-floris. — *Herbæ Boreali-Americane, foliis indivisis pinnatifidisve alternis aut oppositis, rarissimè verticillatis.*

229. E. LEPTOPHYLLUM, caule glaberrimo tereti erecto apice paniculato-ramoso, foliis alternis filiformibus tenuibus glabris integerrimis, racemis spicatis elongatis secundis ramos ramulosque terminantibus in paniculam dispositis, capitulis 3-5-floris breviter pedicellatis, invol. squamis lanceolatis margine albo angusto cinctis. ♀ in Georgiâ circa Savannah à cl. Herbmout lectam habui. Habitus inflorescentiæ Solidaginum, cæterum E. fœniculaceo affine. (v. s. comm. à cl. Duby.)

230. E. FœNICULACEUM (Willd. sp. 3. p. 1750), caule striato pubescente paniculato ramosissimo, foliis alternis glabris, infer. pinnatifidis, super. fasciculatis, omnibus linearifiliformibus, capitulis innumeris parvis 3-5-floris, invol. 10-phylli squamis ext. 5 brevioribus pubescentibus, achænio cylindrico glabro. ♀ in pascuis pinguibus præ. maritimis à Virginiâ ad Floridam et Georgiam. Pursh fl. 2. p. 512. Ell. sketch 2 p. 294. E. fœniculoides Walt. car. 199. Chrysocoma capillacea Michx. fl. bor. am. 2. p. 101. Cor. ochroleucæ interdum purpureo tinctæ (Ell.). Habitus Artemisiæ ex Pursh. (v. s.)

ℓ. *traganthes*, paniculæ strictæ angustæ ramulis rachi approximatis, invol. 6-phylo 6-floro. — in Amer. bor. (ex L'Hér. mss. qui asserit plantam in horto Par. natam in terrâ in quâ Dionæa muscipula adfecta fuerat) nec in Chinâ. Artemisia capillifolia Lam. dict. r. p. 266. Art. tenuifolia Willd. sp. 3. p. 1819. non Mœnch nec Adams. Traganthes Wallr. sched. crit. 456. Mikania artemisioides Cass. dict. 55. p. 130. — Dill. elth. 37. t. 33. Specimen habeo ex cl. Jan lectum Venetiis. An spontè? an in horto lectum? (v. s.)

231. E. CORONOPHYLLUM (Willd. sp. 3. p. 1750), caule tereti pubescente diffusopaniculato, foliis subpubescentibus alternis, infer. pinnatifidis lobis linearibus, reliquis fasciculatis angustè linearibus integerrimis, capitulis 5-floris, invol. 10-phylo pubescente basi imbricato, achænio angulato glabro. ♀ in siccis sterilibus Carolinæ. E. compositifolium Walt. car. 199. Chrysocoma coronopifolia Michx. fl. am. 2. p. 102. Cor. albæ. Affine E. fœniculaceo. (v. s.)

232. E. PINNATIFIDUM (Ell. sketch 2. p. 295), caule striato pubescente apice corymboso-ramoso, foliis pubescentibus pinnatifidis, infer. 4^{te} verticillatis, super. alternis, capitulis 5-floris, invol. squamis 8-10 oblongo-lanceolatis pubescentibus, achænio oblongo sulcato. ♀ in humidis Carolinæ mediæ. Cor. albæ. Species media inter. E. fœniculaceum coronopifoliumve et reliquas generis stirpes.

233. *E. LINEARIFOLIUM* (Walt. car. 199), caule tereti pubescente apice ramoso-corymboso, foliis infer. oppositis, sup. alternis, omnibus pubescentibus linearibus aut sublancoelatis 1-nerviis integerrimis aut hinc inde subdentatis, capitulis 5-floris, invol. squamis 10 oblongo-linearibus imbricatis dorso puberis et glandulosis, achænio sulcato-angulato glabro glanduloso. $\frac{1}{2}$ in Carolinâ frequens (Fras.!), circa Charleston (Nois.), à Novâ-Cæsareâ (Greene), ad Carolinam (Pursh fl. 2. p. 513), in Novo-Eboraco (Torr.). Michx. fl. 2. p. 97. Willd. sp. 3. p. 1750. Pluk. alm. t. 88. f. 2? — An *E. hyssopifolium* Ell. sketch 2. p. 297? *E. crassifolium* Rafin. fl. lud. 62? — Styli plerumque corollâ longiores. Flores albidî. (v. s.)
234. *E. HYSSOPIFOLIUM* (Lin. sp. 1171), caule tereti pubescente apice laxè corymboso-ramoso, foliis oppositis summisve alternis lineari-lanceolatis trinerviis pubescentibus punctatisque integerrimis aut inferioribus dentatis, capitulis 5-floris, invol. squamis 10 imbricatis dorso puberis glandulosisque. $\frac{1}{2}$ in pinetis et arenosis à Carolinâ (Fras.!) Bosc!), ad Novam-Cæsaream (Pursh fl. bor. am. 2. p. 512). — Dill. elth. 141. t. 115. f. 140. Flores albi. An *E. linearifolium* Ell. sketch 2. p. 296 huc referendum? sed caulis dicitur subprocumbens, atque styli corollam subæquantes! undè suspicor speciem Elliottii diversam et fortè novam sed mihi ignotam. (v. s.)
235. *E. GLAUDESCENS* (Ell. sketch 2. f. 297), caule pubescente, foliis plerisque oppositis subsessilibus lanceolatis obtusè pauci-serratis basi integerrimis triplinerviis subglanciis pubescentibus, capitulis corymbosis 5-floris, invol. squamis 8-10 lanceolatis acutis dorso glandulosis, stylo corollâ longiore $\frac{1}{2}$. In pin-guibz umbris Carolinæ. Cor. albæ. Serraturæ 3-4 utrinque à medio ad apicem folii.
- ß. leucolepis*, capitulis 5-7-floris, invol. squamis lanceolatis acuminatis apice albido-scariosis. $\frac{1}{2}$ in Novâ-Cæsareâ (Torr. ! Greene!) Eup. linearifolium Michx? ex Torr. in litt. sed ab *E. linearifolio* differt foliis evidenter à medio ad apicem serratis, invol. acuminatis nec subacutis. (v. s.)
236. *E. ALTISSIMUM* (Lin. sp. 1171), caule tereti pubescente apice ramoso-corymboso, foliis oppositis subsessilibus lanceolatis trinerviis utrinque attenuatis pubescentibus, inf. medio aut à medio ad apicem serratis, sup. integerrimis, capitulis 5-floris, invol. squamis 10 oblongo-linearibus imbricatis obtusiusculis pubescentibus, achænio glabro glanduloso teretiusculo. $\frac{1}{2}$ in arenosis sylvaticis demissis Pennsylvaniae, Virginiae et ad ripas fl. Mississipi, Missourii. Pursh fl. am. bor. 2. p. 514. Jacq. hort. Vind. t. 164. Willd. sp. 3. p. 1754. Fl. albi. Caulis 3-4-pedalis. (v. s. sp. et v. c.)
237. *E. SEMISERRATUM*, velutino-pubescentibus subvillosis, caule tereti, foliis 3^{is} verticillatis, ramis summisve oppositis lanceolatis utrinque longè acuminatis triplinerviis infernè integerrimis subpetiolatis, supernè serraturis obtusis 9-10-profundè notatis, corymbo composito fastigiato, capitulis 5-floris, invol. squamis 10, int. linearibus obtusis, ext. brevissimis, achænio angulato glabro. $\frac{1}{2}$ in Georgiâ circa Savannah legit cl. Herbemont. Cor. albæ parvæ. Affine *E. cuneifolio*. (v. s. comm. à cl. Duby.)
238. *E. CUNEIFOLIUM* (Willd. sp. 3. p. 1753), caule tereti pubescente, foliis oppositis alternisve angusto-lanceolatis basi cuneato-atteuatis subpetiolatis et integerrimis cæterum dentatis triplinerviis utrinque velutino-pubescentibus, corymbo composito, capitulis 5-floris, invol. squamis 10 linearibus obtusis, ext. brevissimis, achænio angulato parè glanduloso. $\frac{1}{2}$ in Carolinâ (Bosc.!) ad St. Thomas (Ell.). *E. parviflorum* Ell. sketch 2. p. 299. non Sw. Cor. albæ parùm involucri longiores. *E. Marrubium* Walt. huc à Willd. revocatum videtur omninò diversum. (v. s. comm. à cl. Bosc.)
239. *E. SCABRIDUM* (Ell. sketch 2. p. 299), caule pubescente, foliis oppositis sessilibus ovato-lanceolatis acutè serratis basi integris pubescentibus subacabris subtus subglaucis, capitulis corymbosis 5-floris, invol. squamis 10 acutissimis puberis glandulosis, achænio angulato. $\frac{1}{2}$ in siccis Carolinæ. Cor. albæ invol. longiores. Rami inf. brachiati, sup. alterni.

240. *E. PUBESCENS* (Muhl. in Willd. sp. 3. p. 1753), caule tereti pubescente, foliis oppositis sessilibus ovatis acuminatis utrinque sparsè pubescentibus et punctato-glandulosis, infer. duplicato-serratis, sup. subserratis, corymbo composito fastigiato, capitulis 5-floris, invol. squamis 10 lineari-lanceolatis acutis puberis, achænio angulato. \mathcal{U} in sylvis arenosis Pennsylvaniae (Muhl. Ell.) à Novâ-Cæsareâ ad Carolinam? ex Pursh fl. 514. excl. syn. Michx. Pers. ench. 2. p. 400. excl. syn. Michx. Fl. albi. An *Critonia elliptica* Raf.! in Kentucky orientali lecta ab hoc diversa? (v. s.)

241. *E. AMORNUM* (Pursh fl. bor. am. 2. p. 514), caule solido lævi, foliis oppositis ternisque brevè petiolatis lanceolato-oblongis utrinque acutis serratis glabrisculis subrugosis subtus reticulato-venosis, paniculæ corymboso-fasciculatæ confertæ pedunculis tomentosis, capitulis 5-floris, invol. squamis oblongis acutis coloratis. \mathcal{U} in montanis Novæ-Cæsareæ. Fl. carnei.

242. *E. FALCATUM* (Michx. fl. bor. am. 2. p. 99), glabellum, foliis 4^{te} verticillatis ovali-lanceolatis utrinque acuminatis subfalcatis rariter serratis, paniculâ brachiâtâ multicorymbosâ, capitulis 5-floris, invol. gradatim imbricatis, squamis extimis brevibus ovalibus. \mathcal{U} in Amer. bor. ad ripas fl. Ohio et Scioto. Affine dicitur *E. albo*.

243. *E. ALBUM* (Linn. mant. III), caule tereti pubero apice ramoso-corymboso, foliis oppositis subsessilibus lato-lanceolatis basi attenuatis apice grossè paucidentatis penninerviis utrinque scabriusculis subtus punctatis, capitulis 5-floris, invol. squamis 10 elongato-lanceolatis acuminatis corollas ferè superantibus dorso glandulosis, achænio angulato glabro. \mathcal{U} in Pennsylvania (Linn.), Novâ-Cæsareâ (Torr.), Georgiâ prope Savannah (herb. Duby!), Willd. sp. 3. p. 1752 excl. syn. Kæmpf. Icon Banks. Kæmpf. t. 26. ad *E. Finlaysonianum* relata, hanc speciem refert sed in pl. Japonicâ corollæ involucri longiores et circ. æquales in Americanâ. Cor. albæ. Invol. squamæ apice albidæ. (v. s.)

β . *glandulosum*, caule magis hispidulo, foliis basi minùs attenuatis per totam ferè longitudinem dentatis subtus involucrique multò magis punctato-glandulosis. \mathcal{U} in aridis sylvarum Carolinæ. *E. glandulosum* Michx. fl. bor. am. 2. p. 98. *E. album* Ell. sketch 2. p. 298. (v. s.)

244. *E. TRICRIFOLIUM* (Willd. sp. 3. p. 1753. hort. ber. t. 32), caule tereti pubescente, foliis oppositis sessilibus distinctis ovatis subacuminatis scabris dentato-serratis basi profundius dentatis, corymbo composito subpaniculato, capitulis 5-6-floris, invol. squamis dorso puberis apice acutis, achænio angulato glanduloso. \mathcal{U} in Amer. bor. à Novâ-Angliâ ad Carolinam. Fl. albi. (v. s.)

β . *verbenæfolium*, foliis oblongo-lanceolatis profundius incis. \mathcal{U} à Bostoniâ ad Carolinam. Eup. pilosum Walt. car. 199. Eup. verbenæfolium Michx. fl. bor. am. 2. p. 98. An fortè *E. lanceolatum* Willd. sp. 3. p. 1752? (v. s.)

245. *E. ROTUNDIFOLIUM* (Linn. sp. 1733), caule tereti pubescenti-velutino, foliis oppositis sessilibus distinctis subrotundo- aut ovato-cordatis obtusis dentatis venosis utrinque pubescentibus subtus punctato-glandulosis, corymbo fastigiato, capitulis 5-floris, invol. squamis 10 acuminatis, achænio angulato glanduloso. \mathcal{U} in pinetis et sylvis à Canadâ ad Carolinam (Pursh), in Novâ-Cæsareâ (Raf.), Georgiâ (Herbem.), Pursh fl. 2. p. 514. Ell. sketch 2. p. 300. — Pluk. alm. t. 88. f. 4. — *E. Marrubium* Walt. car. 199? ex Ell. — Flor. albi. Herbæ decoctio tónica febrifuga. (v. s.)

β ? *ovatum* (Torr. in litt. 1835), capitulis 7-8-floris, caule magis hirsuto. \mathcal{U} in demissis circa Bostoniam. Eup. ovatum Bigel. ex Torr. an spec. propria? (v. s.)

***** Capitulis 5-8-floris. — Species *Gerontogea*, foliis indivisis.

246. *E. LONGICAULE*, caule herbaceo erecto tereti ramoso hirsuto-velutino, foliis petiolatis ovato-lanceolatis acuminatis serratis basi tripinnerviis suprâ glabris

subtus pubescenti-velutinis, corymbo composito fastigiato amplo, capitulis 8-floris, invol. squamis circ. 10 oblongis acutis glabrescentibus. 2/ in Nepaliâ legit cl. Wallich, etiam recepi à cl. Royle. *Conyza longicaulis* Wall.! cat. n. 3073. comp. 183 non n. 2169 ad *Vernoniâ longicaulem* referend. *Mikania?* longicaulis Wall.! herb. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

247. *E. WALLICHII*, caule erecto herbaceo ramoso tereti striato pubescenti-velutino, foliis vix petiolatis ovali-lanceolatis acuminatis serratis supra glabrescentibus, subtus pubescentibus, capitulis in corymbum subglobosum densum terminalem congestis subsessilibus paucifloris, invol. squamis paucis obtusis, exter. brevioribus subacutis. 2/ in Nepaliâ legit cl. Wallich. *E. viscosum* Wall.! cat. n. 3287. comp. 397 non Kunth. Ex statu florum juniore nec pumera nec formam ritè vidi sed ex habitu sine dubio huc refero. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

248. *E. BIRMANICUM*, caule suffruticose erecto ramoso apice subpuberulo, foliis breviter petiolatis lanceolatis utrinquè acuminatis parè serratis glabrescentibus, corymbo oligocephalo, capitulis parvis paucifloris, pedicellis canescenti-villosis, invol. glabro. 3/ in Indiâ or. regione Birmanorum ad Segain legit cl. Wallich. *E. suaveolens* Wall.! cat. n. 3290. comp. n. 400 non Kunth. Folia fere *E. triplinervii* sed penninervia. Capitula in specim. obvio nimis juniora! (v. s. in h. Cur. Ind. or. nunc Soc. Lin. Lond.)

249. *E. FINLAYSONIANUM* (Wall.! cat. n. 7133), caule herbaceo ramoso erecto? striato tereti apice puberulo, foliis lanceolatis in petiolum attenuatis acuminatis grossè et irregulariter inciso-serratis glabris, corymbo composito fastigiato, pedunculis subbracteolatis, capitulis 5-floris, invol. squamis circ. 10 inæqualibus oblongo-linearibus glabris, ext. acuminatis. 2/ in Indiâ or. (ex herb. Finl.) et Japoniâ si ut autumnò *E. album* Thunb. fl. jap. 308. Banks ic. Kæmpf. t. 26 non Linn. huc pertinet. (v. s. comm. ab ill. Cur. merc. angl. Ind. or.)

250. *E. PUNDUANUM* (Wall.! cat. n. 3170. comp. n. 280), caule herbaceo erecto ramoso tereti pubescenti-velutino, foliis vix petiolatis lanceolatis acuminatis à medio acutè serratis utrinquè puberis subtrinerviis, ramis in paniculam subdispositis nudiusculis apice confertè corymbosis, capitulis 4-floris, invol. squamis glabris 5-7 oblongo-linearibus obtusis. 2/ ad Pundua Ind. or. legit cl. Wallich. Achænia subangulata glabra basi attenuata subpuberula. Pappus albidus cor. æqualis. Styli rami exserti longissimi! (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

β. *Lambertianum* (Wall.! cat. n. 3285. comp. 280), caule paulò magis hirsuto. — cum priori. (v. s. comm. cum α.)

251. *E. REEVESII* (Wall.! cat. n. 3168. comp. 278), caule herbaceo erecto tereti glabrescentulo, foliis subsessilibus ovatis acutis grossè et obtusè dentatis glabris nervis subtus et interdum supra puberulis, pedunculis axill. nudis apice congestè corymbosis, supremis in corymbum compositum confertis, capitulis subsessilibus 5-floris, invol. squamis circ. 10 inæqualibus glabris oblongo-linearibus obtusis. 2/ in Chinâ legit cl. Reeves. Fl. ex sicco vid. albi. Pappus albidus. Achænia glabra striato-angulata. (v. s. in h. Calc. conum. ab ill. Cur. Ind. or. et in h. Lindl. à cl. Parkes circ. Canton lect.)

252. *E. NODIFLORUM* (Wall.! cat. n. 3166. comp. 276), caule herbaceo erecto subsimplici tereti hirsuto, foliis sessilibus oblongo-lanceolatis mucrone acutis grossè serratis sparse puberis subtus in nervis hispidis, corymbo composito conferto, capitulis ad apices ramorum congestis 4-floris, invol. squamis 5-6 inæqualibus dorso pubescentibus. 2/ in Indiâ or. montibus Silhetensibus. Strictum 1-2-pedale. Folia vere penninervia sed nervis infimis 2 validioribus. Achænia striato-angulata glabra glandulosa. Pappus sordide albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

253. *E. CINEREA* (Linn. sp. 1172), herbaceum, caule tereti striatulo tenuè

puberulo oppositè ramoso, foliis oppositis breviter petiolatis ovato-lanceolatis acutis serratis utrinquè tenuissimè puberulis, corymbis terminalibus compositis subfastigiatis, capitulis pedicellulatis 5-floris, invol. squamis 5-7 oblongo-linearibus acutis dorso puberulis, achænio teretiusculo striato glabro. \mathcal{L} in Chinâ (Linn. h. Lindl.), Japoniâ? (Thunb. fl. jap. 308). An stirps Japonica de quâ Thunb. caulem et folia glabra et Willdenowiana (sp. pl. 1753) cui auctor folia obovatâ glabra tribuit, sunt omninò similes? (v. s. in h. Lindl.)

254. E. LINDLEYANUM, caule herbaceo erecto simplici tereti striato puberulo apice nudiusculo, foliis oppositis subsessilibus lineari-lanceolatis remotè serratis glabriusculis basi trinerviis, corymbo composito terminali, pedicellis sub capitulo minutissimè bracteolatis, invol. squamis 5 oblongo-lanceolatis acutis, floribus 5, achæniis angulatis minutè glandulosis. \mathcal{L} in Chinæ collibus. Radices è collo fasciculatæ teretes. Folia sesquipoll. longa, 2-3 lin. lata. (v. s. in h. Lindley.)

***** Cannabina. — Capitula 5-6-flora. Invol. squamæ 7-9 circ. biseriatae oblongo-lineares non verè imbricatæ. — Folia opposita aut vericillata, indivisa aut trifida. Caulis herbacei.

255. E. CANNABINUM (Linn. sp. 1173), caule herbaceo erecto striato scabro, foliis plerisque trisectis, segmentis lanceolatis acuminatis basi attenuatis regulariter serratis pubescentibus, corymbo fastigiato, capitulis 5-6-floris, invol. cylindrico glabro circ. 10-phylo, squamis ext. 5 obtusis brevissimis. \mathcal{L} in fossis, aquosis, inundatis totius Europæ, Persiæ (Szw.!), Tauriæ (Bieb.). OED. fl. dan. t. 745. Sow. engl. bot. t. 478. E. trifoliatum Habl. non Linn. Flores pallidè purpurascens et var. albi, odore melleo insignes. Achænia subpentagona glandulosa. (v. v.)

β . *indivisum* (DC. fl. fr. n. 3107), foliis omnibus aut plerisque indivisis. Ray. syn. 180. E. Soleirolii Lois. nouv. not. 36. fl. gall. ed. 2. v. 2. p. 223 huc relat. ex Petit ann. sc. obs. 1829. p. 463 et ex specim. Soleir. (v. s.)

256. E. SYRIACUM (Jacq. ic. rar. 1. t. 170. misc. 2. p. 349), caule herbaceo erecto subpubescente, foliis plerisque trisectis, segmentis ovato-lanceolatis acuminatis basi breviter attenuatis subpubescentibus grossè et inæqualiter dentatis, corymbo fastigiato, capitulis 5-6 floris, invol. cylindrico glabro circ. 10-phylo, squamis ext. 5 obtusis brevissimis. \mathcal{L} in Syriâ (Jacq.) et Persiâ (Szw.) Willd. sp. 3. p. 1757. Fortè mera E. cannabini varietas? (v. v. c.)

β . *indivisum*, foliis plerisque indivisis. \mathcal{L} in Persiâ boreali. (Stev. l Szw.!). (v. s.)

257. E. HETEROPHYLLUM, caule herbaceo tereti striato puberulo, ramis gracilibus, foliis oppositis petiolatis suprâ sparè subtùs secus nervos puberulis, caulinis trisectis, segmentis ovali-lanceolatis utrinquè acuminatis medio grossè serratis, ramis indivisis ovali-lanceolatis subintegerrimis, corymbis terminalibus confertis, capitulis pedicellatis 5-floris, invol. glabro 7-8-phylo, squamis 3-4 exter. inæqualibus brevibus, 4 oblongis obtusis subæqualibus, achænio glabro. — Patr. ign. Speciem distinctissimam vidi in tepidario univ. Vindob. anno 1827 à cl. et amic. Jacquin commun. Stylus basi filiformis. Cor. basi inflato-globosa teretiuscula intùs glabra. (v. s.)

258. E. JAPONICUM (Thunb. fl. jap. 308), caule herbaceo erecto striato scabro, foliis longiusculè petiolatis, inf. trisectis, segmentis foliisque anper. ovatis acutis sæpius serratis subtùs pallidè scabridis, paniculis terminalibus ovatis, pedunculis brevissimis, capitulis 5-floris. \mathcal{L} in Japoniâ. An ex Willd. ab E. cannabinò (seu syriaco) satis differt? Petioli dicuntur unguiculares nec 2-4 lin. longi. †

***** Capitulis 4-5-floris. — Species ex Americâ calidiore foliis indivisis.

259. E. NITIDUM, fruticosum glaberrimum, caule tereti, foliis oppositis bre-

vissimè petiolatis approximatis ellipticis basi et apice attenuatis integerrimis triplinerviis utrinque lucidis, corymbi compositi fastigiat ramis inf. oppositis, summis alternis compressis, capitulis pedicellatis 5-floris, invol. squamis 1-2-seriatis subæqualibus mucronato-acuminatis, achænio puberulo. $\frac{3}{4}$ in sabulosis aridis circa Bahiam Brasilis leg. cl. Salzmann et Blanchet. Frutex 5-7-ped. Folia 2 poll. longa, pollicem lata, petiolo bilineari. Cor. alba. Antheræ ferè cor. orem æquant. (v. s.)

260. *E. BOGOTENSIS*, fruticosum glabrum, ramis teretibus, foliis breviter petiolatis oblongo-lanceolatis utrinque subacutis serratis coriaceis penninerviis, paniculâ terminali cernuâ viscosâ, capitulis longè pedicellatis 5-floris, invol. squamis 7-8 biserialibus inæqualibus obtusiusculis, achænio angulato glabro. $\frac{3}{4}$ circa Santa-Fe de Bogota. Species paniculâ cernuâ et quasi reflexâ (ex n. specim.) distinctissima. Petiol. 2 lin. Limb. fol. 2 poll. longus, 7 lin. latus. (v. s. comm. ab amic. Delessert.)

261. *E. SERRATIFOLIUM*, fruticosa erecta glabra, ramis sexangularibus, foliis petiolatis ovato-oblongis acuminatis basi cuneatis dentato-serratis reticulato-trinerviis, corymbis terminalibus ramosissimis, capitulis solitariis pedicellatis 5-floris, invol. squamis 5 oblongo-linearibus obtusis umbellatis viscosis, achænio angulis pilosisculis. $\frac{3}{4}$ in Novâ Granatâ. Petiol. 3-4 lin. Limb. fol. 15-16 lin. longus. Mikania serratifolia H. B. et Kunth nov. gen. am. 4. p. 138.

262. *E. SIBERIANUM*, herbaceum pubescens, caule tereti striatulo erecto, foliis petiolatis ovato-lanceolatis basi cuneatis apice acuminatis repando-dentatis suprâ sparè subpuberulis subtis in nervis petiolisque puberis, paniculâ laxâ subtrichotomâ, ramis apice capitula oblonga glabra 5-flora fasciculato-sensilia gerentibus, invol. squamis 6-7 subbiserialis oblongis mucronatis, achænio angulato puberulo. — in insulâ Trinitatis legit cl. Sieber. Mikania serratifolia Sieb. f. fl. trin. exs. n. 72 non Kunth. Petiol. 6 lin. longi. Limbus 3 poll. ferè longus, 12-15 lin. latus. Capitula pallida 2 lin. longa. (v. s.)

263. *E. LIGUSTRINUM*, fruticosum, ramis teretibus glabris, ramulis subangulatis pulverulento-velutinis, foliis oppositis petiolatis oblongis aut subellipticis obtusiusculis basi attenuatis et subrecurvis ad apicem dentibus paucissimis notatis glabris subtis crebrè punctatis, corymbis trichotomis polycephalis, pedicellis glanduloso-puberulis ferrugineis, capitulis 4-5-floris, invol. squamis 5-6 linearibus obtusiusculis sub 1-seriatis, ext. paucis parvis acutis, achænio angulato subacabrido. $\frac{3}{4}$ in Mexico ad Tamaulipas legit cl. Berlandier. Cor. alba, tubo brevi, fauce obconicâ. Folia circ. Ligustri. (v. s.)

264. *E. TRIFIDUM* (Vahl symb. 3. p. 94), scandens, caule striato vix pubescente, foliis alternis petiolatis tripartitis, lacinii linearibus extus serratis, summis indivisis lanceolatis, corymbis longè pedunculatis compositis, capitulis 5-floris, invol. squamis glabris 10-phyllis biserialis, extimis brevissimis. — in Caribæis.

265. *E. GLUMACEUM*, caule herbaceo erecto tereti pubescenti-piloso, foliis oppositis petiolatis ovato-lanceolatis acuminatis hinc indè subserratis suprâ pubescenti velutinis subtis velutino-tomentosis trinerviis, paniculâ laxâ diffusâ trichotomâ, ramulis ad apices capitula plurima oblonga 4-flora breviter pedicellata gerentibus, invol. squamis stramineis glaberrimis acuminatis biserialis, ext. brevioribus. — in insulâ Cubâ legit cl. de la Ossa. Folia internodiis multò breviora, adjecto petiolo 4-lin., 3 poll. ferè longa, ferè pollicem lata. Invol. glumacea subnitentia. Achæniû angulatum puberulum. (v. s.)

266. *E. CHILCA* (H. B. et Kunth nov. gen. am. 4. p. 126), fruticosum glabrum, ramis striatis, foliis oppositis petiolatis ovato-oblongis acuminatis serratis basi angustatis triplinerviis, paniculis terminalibus corymbosis foliatis, capitulis pedicellatis et sessilibus sub-4-floris, invol. squamis 4-6 subimbricatis oblongo-lanceolatis acutis externè glutinosas, achænio glabro $\frac{3}{4}$ ad radicea monda Sanctæ-Polonis juxta Caxamarca Peruvianorum. Cor. alba. Petiol. lineam longus. Fol. limbus sesquipollicem longus, 4 lin. latus. Frutex aromaticus Chilca vern. dictus. An Mikaniæ spec.?

267. *E. AFFINE* (H. B. et Kunth nov. gen. am. 4. p. 126), fruticosum glabrum, ramis striatis, foliis oppositis petiolatis ovato-oblongis acuminatis serratis basi cuneatis triplinerviis, paniculis terminalibus corymbosis, capitulis pedicellatis et sessilibus 3-6-floris, invol. squamis 4-6 subimbricatis oblongo-lanceolatis acutis externè glutinosis, achænio ad angulos scabriusculo. $\frac{5}{2}$ cum *E. Chilcæ*? à quo ipso docente Kunthio vix differt, nisi foliis latioribus longius petiolatis, capitulisque majoribus.

***** *Multiloba*. — *Capitula 5-flora. Invol. squamæ 5 æquales subuniseriales.* — *Folia multifida, lobis linearibus. Caules fruticulosi.*

268. *E. ANETHIFOLIUM*, suffruticulosum? ramosissimum adscendens glabrum, ramis teretibus junior. striatis, foliis oppositis petiolatis multipartitis, lobis oppositis linearibus integerrimis ultimis trifidis, corymbi terminalis ramis ramulisque hirtellis, bracteis setaceo-subulatis, capitulis pedicellatis 5-floris, invol. squamis 5 glabrinusculis oblongo-linearibus subobtusis. $\frac{2}{2}$ in Brasiliæ prov. Rio-Grande. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 913 et 929 miss.)

269. *E. MULTIFIDUM*, fruticosum ramosissimum adscendens, ramis teretibus junioribus setoso-hispidis et pube minimâ glandulosâ subvelutinis, adultis glabris, foliis oppositis subsessilibus multipartitis, lobis oppositis apice trifidis linearibus oblongis integerrimis glabris rarius dorso subsetosis, corymbo composito pedunculato conferto, ramis alternis brevibus, capitulis subsessilibus circ. 5-6-floris, invol. biseriali, ext. foliolis setaceis, inter. squamis 5-6 oblongis obtusis apice ciliatis. $\frac{5}{2}$ in Brasiliæ prov. Sancti-Pauli. Species distinctissima vix pedalis aut sesquipedalis. Folia 6-8 lin. diam. cum invol. subviscosa. Flor. ex sicco albi. Achæniolum glabrum angulatum. Pappus corollâ dimid. ferè brevior. Habitus à genere alienus! (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 508 et 509 miss.)

270. *E. BRASILIENSE* (Spreng. syst. 3. p. 417), caule herbaceo simpliciusculo, foliis subsessilibus confertis 3-5-partitis linearifiliformibus glabris, corymbis lateralibus, capitulis sub-3-floris, invol. glabro subimbricato aut simplici. — in Brasiliâ. Affine fortè *E. multifido*. Cæt. ign. †.

† *Species non satis notæ.*

271. *E. AMYGDALINUM* (Lam. dict. 2. p. 408), caule... tereti simplici glabro, foliis oppositis sessilibus oblongo-lanceolatis subdentatis glabris, paniculâ laxâ subnudâ, capitulis 12-15-floris, invol. ovalis squamis oblongis inæqualibus margine subvillosis. — in Peruviâ ex h. Juss. Folia inf. 3 poll. longa, summa subalterna.

272. *E. AUSTRALE* (Thunb. in fl. 1821. p. 332. n. 20), fruticosum, ramis..... foliis..... triplinerviis ellipticis serratis, paniculâ fastigiatâ, capitulis..... $\frac{5}{2}$ in Brasiliâ. †.

273. *E. BRICKELLIA*, caule herbaceo tenuissimè pubescente apice subtomentoso, foliis oppositis cordatis acuminatis dentatis triplinerviis tenuè pubescentibus, summis subalternis basi obtusis, corymbo paniculato, invol. squamis ext. setaceis, int. linearibus lanceolatis, achænio subglabro ro-striato. — ad latera collium in Georgia occidentali, Brickellia cordifolia Ell. sketch 2. p. 290. Jam suadente cl. Lessing hanc stirpem ab Eupatorio non separare queo, sed ex numero florum potius ad Eupatorii species Mexicanas quàm boreali-Americanas accedere videtur.

274. *E. CHILENSE* (Molin. chil. ed. gall. p. 335), foliis oppositis amplexicaulibus lanceolatis denticulatis, invol. 5-floris. — in Chili. Cæt. ign.

275. *E. COTINIFOLIUM* (Willd. phytogr. II. t. 7. f. 1. ex sp. pl. 3. p. 1769), fruticosum, foliis alternis petiolatis obovatis mucronatis integerrimis triner-viis, corymbo terminali. $\frac{5}{2}$ in Martinicâ. Capitula multiflora, involacro non imbricato ex Spreng. †.

276. *E. FREYREYII* (Thunb. in florâ 1821. p. 332. n. 19), fruticosum scandens, ramis... foliis... lanceolatis triplinerviis glabris serratis, paniculis patentibus, capitulis... 5 in Brasiliâ. †.
277. *E. HISPIDUM* (Pers. ench. 2. p. 402. n. 66), caule suffruticoso hispido erecto, foliis alternis ovatis sinuatis denticulatis pilosis, paniculis terminalibus capitulis sub-8-floris, invol. oblongo imbricato. 5 in insulâ Afric. Mozambico. *E. sinuatum* Lour. fl. coch. 2. p. 595. non Lam. Flor. purpurascens. Styli longi semibifidi. Recept. concavum. An. *Vernoniæ* potius species? †.
278. *E. LUDOVICIANUM*. — in Louisianâ. Cactis n. 1. Rob. louis. 432. Cact. pubescens Raf. fl. lud. 61.
279. *E. MEGAPOTAMICUM* (Spreng. syst. 3. p. 420), herbaceum, foliis alternis petiolatis ovatis mucronato-dentatis ramisque villosis, paniculâ terminali, capitulis paucifloris, invol. glabro subimbricato aut subsimplici. — in Brasiliæ prov. Rio-Grande. Cact. ign. †.
280. *E. MENTHAFOLIUM* (Poepp. ex Spreng. syst. 3. p. 412), caule fruticoso, foliis alternis petiolatis oblongis utrinquè attenuatis medio serratis pubescentibus, paniculis axillaribus terminalibusque patentibus, capitulo multifloro; invol. imbricato. 5 in Cubâ. Cact. ign.
281. *E. MONTEVIDEENSE* (Spreng. syst. 3. p. 417), caule herbaceo ramoso, foliis sparsis subsessilibus linearibus margine subserrulato-scabris, capitulis racemosis paucifloris, pappo rufo. — in Montevideo. Cact. ign. †.
282. *E. MYOSOTIFOLIUM* (Jacq. ic. rar. 3. t. 282. coll. 2. p. 341), herbaceum villosulum, caule tereti striato, foliis alternis in petiolum attenuatis obovato-cuneatis obtusis integerrimis, paniculâ diffusâ laxâ, capitulis longè pedicellatis cylindricis multifloris, invol. squamis acutis erectis. ② in Amer. calid. Cor. purpureâ. Cact. ign. †.
283. *E. FRAXINIANUM* (Spreng. syst. 3. p. 412), fruticosum, foliis alternis petiolatis oblongis utrinquè attenuatis obtusiusculis repandis suprâ opacis glabris subtus fusco-tomentosis, paniculâ terminali foliosâ, capitulis multifloris, invol. imbricato oblongo-cylindraceo. 5 in Amer. austr.? Cact. ign. †.
284. *E. POEPPIGII* (Spreng. syst. 3. p. 415), fruticosum, ramis... foliis oppositis petiolatis ovato-lanceolatis subintegerrimis subtus subtomentosis, paniculâ corymbosâ terminali, capitulis paucifloris, invol. pauciseriato. — in insulâ Cubâ. Eup. deltoideum Poepp. exs. non Jacq. Cact. ign. An fortè idem ac *E. cynanchifolium*, sed descr. Spreng. non sufficit.
285. *E. QUATERNUM*, caule suffruticoso subsimplici striato, foliis subquaternis congestis subsessilibus ovato-lanceolatis subserratis rugoso-asperis, corymbo fastigiato, capitulis sub-8-floris, invol. oblongo-cylindrico. 5 in agris Cochinchinæ. E. purpureum Lour. fl. coch. 2. p. 595. non Linn. Suffrutex 3-pedalis. Flores purpurei. Recept. subnudum. †.
286. *E. RETICULATUM* (Desv. in Ham. fl. ind. occ. p. 51. non Hook.) glaberrimum, ramis pentagonis, foliis oppositis obovato-oblongis retusis subundulatis consistentibus subtus reticulatis, paniculâ contractâ apice subumbellatâ, capitulis multifloris. — in Jamaicâ. Cact. ign.
287. *E. ROSALESIA*, caule fruticoso, ramis virgatis apice involucriisque glanduloso-viscosis, foliis oppositis petiolatis ovato-subcordatis crenato-subseratis suprâ asperis subtus subtomentosis, capitulis corymbosis, pedunculis capitula 2-5 gerentibus, invol. imbricato, squamis apice revolutis. 3 in Mexici pascuis ad S. José del Corral. *Rosalesia glandulosa* Lallav. et Lex. nov. veg. 1. p. 9. Frutex biorgyalis. Cor. pallidè flavæ, limbo brevissimè 5-dentato. Styli exserti insigniter clavati. Achænia cylindrica striata villosa.
288. *E. SERRATUM* (Spreng. syst. 3. p. 415), fruticosum, ramis... foliis oppositis subpetiolatis lineari-lanceolatis argutè serrulatis subtus tomentosis, paniculæ terminalis ramis divaricatis, capitulis paucifloris, invol. paucis seriebus imbricato aut subsimplici. 5 in Monte-Video. Cact. ign.

289. *E. SOPHIAEFOLIUM* (Linn. sp. 1175), herbaceum, caule adscendente à basi multiplici, foliis oppositis bipinnatisectis, lobis sinuatis oblongis obtusis, corymbo paniculato, capitulis pedicellatis, invol. cylindrici squamis arcuè imbricatis multiseriatis. — in insulis Caribæis. Eup. sophiaefolio flore purpureo Plum. amer. ed Burm. t. 128. f. 2. †.
290. *E. SPICATUM* (Lam. dict. 2. p. 409), caule herbaceo? simplici striato subangulato, foliis oppositis lineari-spathulatis superne rariter dentatis subtilissimè tomentosis, capitulis in spicam terminalem basi interruptam dispositis sessilibus subglomeratis multifloris, invol. ovoidei squamis oblongis margine scariosis. 2L (Pers.) 5 (ex Lam.) — in Monte-Video ex Comm. in h. Juss. Folia 2 poll. longa.
291. *E. SPRENGELIANUM*, fruticosum, ramis teretibus pubescentibus, junioribus albo-tomentosis, foliis alternis brevissimè petiolatis oblongis obtusis repandentatis coriaceis rugosis subtus tomentosis, racemis terminalibus oligocephalis. 5 in insulis Indiæ occid. E rugosum Spreng. neu entdt. 3. p. 39. non Kunth nec Schrad. Species valde dubia ab auct. ipso in systemate omissa. †.
292. *E. SQAMOSUM* (Don prod. fl. nep. 170), caule fruticoso diffuso, ramis teretibus, foliis ovatis acutis petiolatis crenatis 3-nerviis utrinquè tomentosis, corymbo paniculato, capitulis. . . . , involucri oblongis pedicellisq. scabris, squamis ext. ovatis acutis, intimis oblongis obtusis. 5 in Nepaliâ. Flores albi.
293. *E. STACHYOPHYLLUM* (Spreng. syst. 3. p. 420), caule herbaceo ramoso hirsuto, foliis oppositis alternisque subsessilibus lanceolatis subserratis rugosis 3-nerviis pubescentibus, corymbis terminalibus oligocephalis, capitulis paucifloris, invol. squamis obtusis glabris. — in Brasiliâ. Cœt. ign.
294. *E. TOMENTELLUM* (Schrad. ind. sem. h. Gœtt. 1833. p. 3), frutescens, foliis cordatis acuminatis subtus cano-pubescentibus integris, ramorum juniorum 3-5-lobis villosis, fol. radic. ligulis obovato-ellipticis. 2L 5 in Mexico.

N. B. Ad Eupatorii genus pertinere videntur stirpes floræ fluminensis vol. 8 sequentes, nempe *Chrysocoma maculata* t. 6, *Chrysocoma alata* t. 48 (affinis *E. quadrangulata* sed canes teretes), *Chrysocoma punctulata* t. 49, *Cacalia emarginata* t. 52, *Cacalia punctulata* t. 66, *Cacalia cor Jesu* t. 71?, *Cacalia triphylla* t. 74, et *Cacalia lata* t. 76, sed descriptionibus deficientibus non satis notæ.

† Species exclusæ.

- E. ACUMINATUM* Wall. = *Vernonia acuminata*.
E. ACUMINATUM Don. =? *Vernonia multiflora*.
E. ALTERNIFOLIUM Ard. = *Kuhnia eupatorioides*.
E. AMARUM Steud. = *Mikania amara*.
E. AREOLATUM Pers. = *Ethulia conyzoides* var. *l.*
E. ARMANI Balb. = *Orsinia eupatorioides*.
E. ASPERUM Roxb. = *Vernonia aspera*.
E. AURICULATUM Lam. = *Senecio deltoides*.
E. AXILLARE fl. mex. = *Bulbostylis Cavanillesii*.
E. CAPENSE Spr. = *Vernonia mespilifolia*.
E. BICOLOR Lam. herb. Juss. = *Gynoxis buxifolia*.
E. CELEBICUM Blum. = *Vernonia Celebica*.
E. CINEREUM L. f. = *Pteronia incana*.
E. CÆLESTINUM L. = *Conoclinium cælestinum*.
E. COLORATUM Willd. = *Decaneurum Senegalense*.
E. CONDENSATUM Reichenb. = *Adenocyclum condensatus*.
E. CORDATUM L. in. non Walt. = *Mikania volubilis*.

- E. CORTICACRUM** Vahl non Spr. = *Vernonia fimbrillifera*.
E. CORNIFOLIUM Willd. = *Odontoloma acuminatum*.
E. CUBENSE Pers. non DC. = *Kuhnia rosmarinifolia*.
E. CUSPIDATUM Willd. = *Dialesta bicolor*.
E. CYDONIÆFOLIUM Willd. = *Vernonia dichocarpa*.
E. DALEA Linn. = *Critonia Dalea*.
E. DALEA H. B. K. = *Critonia pseudodalea*.
E. DELTIFOLIUM Raf. = *Conoclinium coelestinum*.
E. DENTICULATUM St. = *Mikania denticulata*.
E. DIVARICATUM Thunb. = *Pteronia divaricata*.
E. DISCOLOR Spr. = *Vernonia discolor*.
E. ELEGANS Vent. = *Liatris elegans*.
E. ELÆAGNIFOLIUM Hort. Calc. = *Vernonia elæagnifolia*.
E. FERRUGINEUM Labill. = *Ozothamnus ferrugineus*.
E. FLEXUOSUM Lam. = *Senecio flexuosus*.
E. GLABRUM Heyn. = *Decaneurum glabrum*.
E. HOUTONIS Linn. = *Mikania Houstonia*.
E. JAVANICUM Blum. = *Vernonia Javanica*.
E. KLEINIODES H. B. K. = *Campuloclinium kleiniioides*.
E. LONGICAULE Wall. n. 2109. = *Vernonia longicaulis*.
E. MACROCEPHALUM Less. = *Campuloclinium macrocephalum*.
E. MACROPHYLLUM Vahl. = *Hebeclinium macrophyllum*.
E. MELISSÆFOLIUM Lam. = *Nothites melissæfolia*.
E. MICROPHYLLUM L. = *Conoclinium microphyllum*.
E. MOLLE Swartz = *Hebeclinium macrophyllum*.
E. NEPTEÆFOLIUM H. B. K. = *Bulbostylis nepetæfolia*?
E. NERVOSUM Sieb. non Sw. = *Baccharis nervosa*.
E. NUTANS H. B. K. = *Bulbostylis secundiflora*?
E. OBTUSIFOLIUM Willd. = *Vernonia Vahlia*.
E. OBOVATUM Willd. = *Vernonia odoratissima*.
E. OLIGANTHES Less. = *Bulbostylis oliganthes*.
E. PARVIFLORUM Aubl. = *Mikania amara*.
E. PARVIFLORUM Swartz. = *Critonia parviflora*.
E. PAUCIFLORUM H. B. et K. = *Bulbostylis*? *pauciflora*.
E. PECTINATA Wall. = *Vernonia pectinata*.
E. PENDULUM Schrad. = *Bulbostylis pendula*.
E. POLYANTHES Spr. = *Vernonia polyanthes*.
E. POLYANTHUM Wall. = *Vernonia Neelgherryana*.
E. POLYSTACHYUM Willd. = *Vernonia mollis*.
E. PYRAMIDALE Don = ? *Vernonia aspera*.
E. RETROFRACTUM Thunb. = *Pegoletia polygalæfolia*.
E. RIGIDUM Swartz = *Ooclinium rigidum*.
E. ROSMARINIFOLIUM Labill. = *Ozothamnus rosmarinifolius*.
E. SATURIAÆFOLIUM Lam. = *Nothites satureiæfolia*.
E. SCANDENS Linn. = *Mikania scandens*.
E. SCANDENS Link = *Senecio deltoides*.
E. SCANDENS Thunb. = *Mikania auriculata*.

- E. SCANDENS Wall. = *Mikania pubescens*.
 E. SCARIOSUM Wall. = *Bulbostylis pendula*.
 E. SECUNDIFLORUM Bert. = *Vernonia arborescens* β.
 E. SECUNDIFLORUM Lag. = *Bulbostylis secundiflora*.
 E. SPECIOSUM Vent. = *Liatris elegans*.
 E. STIPULACRUM Steud. = *Mikania stipulacea*.
 E. SQUARROSUM Cav. = *Bulbostylis Cavanillesii*.
 E. TOMENTOSUM Lam. = *Senecio penicillatus*.
 E. UNIFLORUM Moq. = *Monosis salicifolia*.
 E. URTICIFOLIUM L. f. = *Campuloclinium urticifolium*.
 E. VAGANS Wall. = *Vernonia vagans*.
 E. VERONICÆFOLIUM H. B. K. = *Bulbostylis veronicæfolia*.
 E. VESTITUM Spreng. = *Salmea hirsuta*.
 E. VINCEFOLIUM Lam. = *Mikania amara*.
 E. VOLKAMERIEFOLIUM Wall. = *Vernonia volkameriaefolia*.
 E. VOLUBILE Vahl. = *Mikania volubilis*.
 E. ZEYLANICUM L. = *Vernonia Zeylanica*.

LXXXVIII. NOTHITES Cass. dict. 35 (1825) p. 163. — *Eupatorii* sp. Lam. Willd. Less.

Capitulum oblongum 5-florum. Recept. nudum angustum. Invol. 5-phyllum, squamis 1-seriatis erectis oblongo-lanceolatis. Cor. tubo brevi, fauce elongatâ intus villosâ. Antheræ subinclusæ, appendicibus obtusissimis. Achænia subpentagona hispidiuscula. Pappi setæ 10-20-uniseriales filiformes apice barbellatæ basi membranâ angustissimâ marginatæ. — Herbæ ex Amer. merid. ortæ. Folia opposita rarius alternata, plerumquæ sessilia, dentata. Corymbi terminales. — Genus mihi ignotum et hic ex cl. Cassini omninò expositum, restitutis tamen nominibus arbitrariò mutatis.

* *Foliis omnibus aut plerisque oppositis.*

1. N. MELISSÆFOLIA, erecta pubescens, foliis oppositis subsessilibus ellipticis inæqualiter serratis utrinquæ punctato-resinosi triplinerviis reticulatis, corymbo terminali glanduloso-pubescente, pappi paleis 10-12. ½ in Peruvia. *Eupatorium melissæfolium* Lam. dict. 2. p. 411. *Mikania melissæfolia* Willd. sp. 3. p. 1747. *Nothites latifolia* Cass. dict. 35. p. 163. Cor. purpureæ. Folia a poll. longa, sesquipollicem lata.
2. N. SATUREIÆFOLIA, erecta suffruticulosa pubescens, foliis suboppositis sessilibus linearibus integerrimis obsolete lanuginosis, corymbis terminalibus. ½ 5 in Monte-Video. *Eupatorium satureiæfolium* Lam. dict. 2. p. 411. *Mikania satureiæfolia* Willd. sp. 3. p. 1747. *Nothites angustifolia* Cass. dict. 35. p. 164. Folia pollicem longa, lineam sesquilineamve lata.
3. N. BREVIFLORA (Cass. dict. 35. p. 165), erecta pubescens, foliis oppositis subsessilibus oblongo-lanceolatis basi angustatis et integris supernè serratis, corymbo terminali glanduloso-pilosiusculo, pappi paleis circ. 17. ½ in Brasiliâ. Folia sesquipoll. longa, semipoll. lata.
4. N. PETIOLATA (Cass. dict. 35. p. 166), ramosissima glabriuscula, foliis oppositis alternisque petiolatis ovalibus obtusè serratis subpubescentibus glanduloso-punctatis, corymbo terminali pubescente, pappo 12-15-setoso, et in unico cujusque capituli flore coroniformi dentato. — in Peruvia circa Limam.

** *Foliis alternis.*

5. *N. BACCHARIDEA*, fruticosa virgata glabra, foliis alternis subpetiolatis linearilanceolatis utrinque acuminatis, aliis medio grossè inciso-serratis, aliis rameis seu ad axillas fasciculatis integerrimis, paniculâ elongatâ parçè foliaceâ polycephalâ, pappi setis circ. 20 scaberrimis. ζ in fruticetis apricis collium circa Valparaiso et Quillota Chilensium legit cl. Bertero. Decemb. flor. Baccharis? Bert. ! herb. n. 837. Invol. 1-ser. 5-phylum. Flores 5. Achænia glabra, matura vix involucri breviora. Styli longè exserti obtusi subclavati. Cor. ex sicco ochroleucæ. (v. s.)
6. *N. OVATIFOLIA*, fruticulosa minutissimè subvelutina, foliis petiolatis ovatis acuminatis hinc indè sublobatis denticulatis penninerviis, paniculâ polythryâ, ramis floridis axillaribus, capitulis brevè pedicellatis confertis, pappi setis circ. 12-15 scabris. ζ in Mexico legit cl. Hænke. Petioli 6 lin. longi. Limb. fol. 27-30 lin. longus, 20-24 latus. Capitula ferè Mikaniæ sed 5-flora. (v. s. in h. Hænk. à cl. de Sternberg.)

LXXXXIX. *MIKANIA Willd. sp. 3. p. 145a. Kunth nov. gen. am. 4. p. 134. Less. syn. 157. non Neck. — Eupatorii sp. Linn. Less. in linnæa 1830.*

Capitulum 4-florum. Recept. nudum angustum. Invol. 4-phylum, adjectâ aut ad basin aut infra basin bracteolâ. Cor. tubus brevis, faux dilatata subcampanulata. Antheræ subexsertæ. Achænium angulatum. Pappus 1-serialis pilosus scabridus. — Frutices aut herbæ, sæpissimè scandentes. Folia constanter opposita, sæpius petiolata, projecturâ transversali inter se ligata, raro ad basin petioli auriculata, limbo sæpius cordato aut ovato sed in genere multiformi. Capitula spicata, racemosa, corymbosa, aut paniculata. Corollæ albidæ aut pallidè flavescentes. — Caract. gener. levis equidem, sed genus adeo naturale ut ordine subartificiali species hinc subungere tantum potui.

§. 1. *Erectæ, caulibus nempè non scandentibus.*

1. *M. LINEARIFOLIA*, herbacea?, erecta parçè ramosa glabra basi radicans, caule tereti, ramis sulcato-angulatis, foliis distantibus sessilibus linearibus acutis integerrimis, corymbo terminali composito, capitulis pedicellatis, bracteolâ adpressâ acutâ minutâ, invol. squamis oblongis mucronatis, achæniis subglandulosis. ζ in Brasiliæ prov. Sancti-Pauli. Species habitu à cæteris diversissima, sed caract. non separanda. Fol. 9 lin. longa, vix lineam lata, internodio multò breviora. Cor. pallidè flavescences. Pappus rufus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 505 miss.)
2. *M. VIMINEA*, herbacea? erecta, caule tereti, ramis oppositis striatis virgatis glabris subaphyllis, foliis oppositis sessilibus linearibus integris remotissimis, corymbo terminali subpaniculati ramis ramulisque pubescenti-hirtellis, capituli pedicellati bracteolâ setacæ hirtelli, invol. squamis lanceolatis acuminatis glabris. ζ in Brasiliæ prov. Rio-Grande. Habitus ferè Cynanchi aphylli. Folia pollicem longa lineam lata. Capit. 4 lin. longa. Pappus rufus. Achænium junius pubescens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 919.)
3. *M. PINNATILoba*, herbacea erecta? glaberrima, caule tereti striato parçè folioso, foliis sessilibus pinnatilobatis, lobis lineari-oblongis integerrimis minutè strigoso-ciliatis, corymbo terminali longè pedunculato, pedicellis involucrique squatis dorso pubescentibus, achæniis glabris. ζ in Brasiliæ prov. Rio-Grande. Si caulibus scandens, ad sect. ultimam prope *M. apiifolia* transferenda. (v. s. in h. Mus. Par. à Mus. Bras. sub n. 927 miss.)
4. *M. AMBROEA*, collo suffruticoso, caulibus pluribus herbaceis erectis striatis puberulis, apice nudo, foliis sessilibus lineari-oblongis acutis aut obtusis aliis

- integerrimis aliis subdentatis glabriusculis subtriplinerviis, corymbo paniculato nudo, invol. squamis lineari-oblongis acutis, bracteolâ iis æquali et conformi. 2 $\frac{1}{2}$ in Brasiliæ prov. Rio-Grande. Herba 7-8 pollic. Folia 2 poll. longa, 4 lin. lata. Pappus rufescens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 944 miss.)
5. *M. OBLONGIFOLIA*, subherbacea? erecta glaberrima, caule tereti striatulo, foliis sessilibus oblongis utrinquè vix subacutis integerrimis penninerviis subreticulatis utrinquè sublucidis rigidulis, paniculæ terminalis ramis erectis angulatis, capitulis secus ramos racemosos pedicellatis, bracteolâ lineari acuminatâ ad pedicelli basin, invol. squamis linearibus acuminatis glabris, achænio juniore rufo-pubero. — in Brasiliæ prov. Sancti-Pauli. Folia sesquipoll. longa, semipoll. lata. Pappus rutilanti-rufus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 415 miss.)
6. *M. MYRICÆFOLIA* (Bojer! in litt. 1835. sub Trizide), fruticosa erecta, ramulis foliorum projecturis exsertis acutè angulatis araneosis, foliis petiolatis oblongis à medio ad apicem dentatis obtusis basi cuneatis utrinquè glabris coriaceis, paniculâ terminali corymbosâ, invol. glabro cylindraceo obtuso, bracteolâ minimâ subaraneosâ. 3 in sylvis prov. Saccalavorum insul. Madagascar legit cl. Bojer. Frutex 6-8-pedalis Myricæ habitus. Species ex projecturis fol. distinctissima. (v. s. flor. nondum expl.)
7. *M. SÆSILIFOLIA*, herbacea erecta, caule simplici tereti hirsuto-subhispido, foliis oppositis sessilibus internodio longioribus imbricatis basi cordatis 3-5-nerviis ovato-orbiculatis obtusis subcrenatis suprâ sparsè scabris subtus reticulatis tomentoso-hirsutis, ramis apice conferto-corymbosis simplicibus basi foliolosis apice spicatis, capitulis approximatis, bracteolâ acutâ invol. dimidio breviorè, invol. squamis oblongis obtusis puberis, achænio glabro. 2 $\frac{1}{2}$? in Brasiliæ prov. Minarum General. legit cl. Vauthier (pl. exs. n. 281). Folia annua subalterna, caulina 2 poll. longa, sesquipollicem lata. Capit 2-2 $\frac{1}{2}$ lin. longa. (v. s. comm. à cl. inv.)
8. *M. LIXIOLÆNA*, caule herbaceo erecto tereti densè tomentoso-hirsuto, foliis oppositis sessilibus ovatis (infer. basi vix subcordatis) acutis hinc indè subdentatis suprâ pubescentibus subtus tomentoso-hirsutis, paniculis axillaribus paniculatis et in paniculam dispositis puberulis, invol. squamis oblongis obtusiusculis achæniisque glaberrimis. 2 $\frac{1}{2}$ in Brasiliæ prov. Minarum General. ad Serra do Frio legit cl. Vauthier (pl. exs. n. 331!). Affinis *M. sæsiliifoliæ*, sed disp. capitulorum, invol. glabritie, etc. facile differt. (v. s.)
9. *M. NUMMULARIA*, herbacea erecta, ramis teretibus glabriusculis aut pulverulento-velutinis, foliis breviter petiolatis orbiculatis basi subcordatis 5-nerviis hinc indè callosò-denticulatis utrinquè tomento minimo canescenti-velutinis subtus reticulatis, medio internodio æqualibus, paniculâ laxâ ramis elongatis ramulisque oppositis, capitulis secus ramulos spicato-congestis, bracteolâ minimâ, invol. squamis ovali-oblongis obtusis dorso velutinis, achænio glanduloso. 2 $\frac{1}{2}$ in Brasiliæ prov. Minarum Gener. ad Tejuco legit cl. Vauthier (pl. exs. n. 266). Herba rhizomate lignoso, parè ramosa, bipedalis, habitu Veronizæ schwenkizifoliæ. Folia diam. 7-8 lin. (v. s. comm. à cl. inv.)
10. *M. NEUROCAULA*, herbacea erecta tota pube minimâ pulverulentâ cinerascens, caule tereti nervis 8 prominulis notato, foliis petiolatis ovalibus basi subcuneatis apice obtusis margine subrepandis aut subdentatis 3-5-plinerviis, paniculæ amplæ ramis oppositis folio triplò circ. longioribus, capitulis fasciculatis breviter pedicellatis, bracteolâ ad basin pedicelli vix eo æquali, invol. squamis oblongo-linearibus obtusis, achænio pubero. 2 $\frac{1}{2}$? in Brasiliæ prov. Minarum Generalium prope Sabara legit cl. Vauthier (pl. exs. n. 270). Nervi caulis ex utroque folio 3 et 1 utrinquè intermedius, sed in ramulis ordo vix conspicuus. Petiol. 4-5 lin. Limbus 15 lin. longus, 10 latus. Folia ferè Halimi, sed opposita. (v. s. comm. à cl. inv.)
11. *M. TERNIFOLIA*, erecta basi suffruticosa, caule subsimplici tereti apice striato *sparsè hispido*, foliis ternato-verticillatis petiolatis ovatis acutis grossè inciso-

- dentatis suprà glabriusculis subtùs secus nervos sparè setoso-hispidis inter eos punctato-glandulosis, paniculâ terminali oppositè ramosâ, ramis apice corymbosis hispidis verticillatis, capitulis confertis brevissimè pedicellatis, bracteolâ ad basin pedicellatâ acuminatâ, invol. squamis oblongis acutis glabris. $\frac{5}{8}$ in Brasiliæ prov. Rio-Grande. Petiol. 3-4 lin. Limb. fol. 10-12 lin. longus, 6 lin. latus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 778.)
12. *M. PASTRMONOIDES*, erecta herbacea, caule simplici subpuberulo striato-angulato, foliis petiolatis hastato-lanceolatis basi cuneatis apice acuminatis grossè et inæqualiter inciso-dentatis pubescentibus, corymbo composito fastigiato, ramis elongatis nudis pubescentibus apice corymbulosis capitulis pedicellatis, bracteolâ minimâ acutâ, invol. squamis lanceolatis acutis glabriusculis. $\frac{2}{3}$ in Brasiliæ prov. Rio-Grande. Herba sesquipedalis. Folia petiolo incluso 3 poll. longa sesquipoll. lata, summa sessilia basi latiora. Pappus rufus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 791 miss.)
13. *M. THAPSOIDES*, herbacea erecta, caule simplici crasso angulato basi glabriusculo apice pedunculisque villosis, foliis confertis pubescenti-villosis acuminatis grossè et irregulariter dentatis, infer. breviter petiolatis ovatis basi cuneatis, super. subsessilibus basi dilatatis, corymbo composito conferto, ramis apice capitula breviter pedicellata fasciculata gerentibus, bracteolâ parvâ acutâ, invol. squamis lanceolatis acutiusculis dorso pubescentibus. $\frac{2}{3}$ in Brasiliæ prov. Rio-Grande. Herba pedalis aut sesquipedalis. Folia inf. 4-5 poll. longa $2\frac{1}{2}$ poll. lata. Capitula 5 lin. longa. Pappus rufus (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 904 miss.)
14. *M. LASIANDRÆ*, erecta suffruticosa, caule ramis petiolisque teretibus hirsutissimis, foliis petiolatis ovato-lanceolatis acuminatis integerrimis, suprà triplinerviis sparè setoso-scabtis, subtùs penninerviis hirsutis, ramis axillaribus terminalibus capitula glomerata subpaniculata gerentibus, bracteolâ ovatâ acutâ brevi glabriusculâ, invol. squamis ovali-oblongis obtusis subcariosis glaberrimis. $\frac{5}{8}$ in Brasiliæ prov. Sancti-Pauli. Hirsuties flavescenti-rufa. Invol. rufo-fusca. Petiol. 5 lin. longi. Limb. fol. 3 poll. longus, 15 lin. latus (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 424 miss.)
15. *M. VISMIEFOLIA*, fruticosa erecta? ramis teretibus densè et brevè velutinis petioliis ramulique ferrugineis velutino-hirsutis, foliis petiolatis ovatis basi obtusis versus apicem sensim acuminatis integerrimis subquintuplinerviis anprâ sparè scabridis subtùs ferrugineo-hirsutis, paniculis terminalibus oppositè ramosis ramulosisque, capitulis fasciculatis breviter pedicellatis, bracteolâ parvâ, invol. squamis lineari-oblongis obtusis dorso puberulis, achænio villosio. $\frac{3}{8}$ in Brasiliæ prov. Minarum Generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 267). Petiol. 8 lin. longi. Fol. limb. $3\frac{1}{4}$ poll. longus, 2 poll. latus. (v. s. comm. à cl. inv.)
16. *M. TEUCRIIFOLIA* (Spreng. syst. 3. p. 423), erecta herbacea, foliis oppositis petiolatis ovato-oblongis serratis subtùs resinoso-pilosis, paniculâ terminali contractâ. — in Brasiliâ. Cæt. ign. †.
17. *M. OFFICINALIS* (Mart. in Isis 1824. p. 587), erecta glabra, caule subimplici, foliis decussatis subtriangulâri-ovatis sinu grosso cordatis latere dentatis anticè integerrimis cernuis, paniculis corymbosis terminalibus. — in Brasiliâ ubi *Coração de Jesu* dicitur. †.
18. *M. CORDIFOLIA* (Willd. sp. 3. p. 1746). erecta fruticosa hirsuta, foliis petiolatis cordato-ovatis acutis trinerviis scabris serratis, paniculis terminalibus, invol. squamis lanceolatis. $\frac{5}{8}$ in Amer. merid. *Cacalia cordifolia* Mutis in Linn. f. suppl. 351. excl. var. *Linnaeus* filius qui solus speciem descripsit dicit in diagnosi involucri pubescentia et in descript. nunc glabra nunc glaberrima. — *Stirps Portoricensis* huc à Spreng. relata est diversa.
19. *M. NODOSA* (Spreng. syst. 3. p. 423), erecta fruticosa glabra, ramis nodosis, foliis oppositis subrhombæo-ovatis serratis triplinerviis, paniculâ terminali corymbosâ. $\frac{5}{8}$ in ins. Canb. Cæt. ign. †.

20. *M. DELTOIDES* (Poepp. ex Spreng. syst. 3. p. 423), erecta fruticosa glabra, ramis teretibus, foliis oppositis deltoideo-ovatis dentatis triplinerviis subtus pallidis, paniculâ terminali corymbosâ. $\frac{1}{2}$ in ins. Cubâ. Cæt. ign. $\frac{1}{2}$.
21. *M. AERIOSA* (Spreng. syst. 3. p. 423), erecta fruticosa, ramis teretibus hirsutis, foliis oppositis oblongis utrinquè attenuatis dentato-serratis subtus resinoso-pilosis, paniculâ terminali corymbosâ. $\frac{1}{2}$ in Bras. Cæt. ign. $\frac{1}{2}$.
22. *M. VISCOSA* (Spreng. neu. entdek. 1. p. 277), fruticosa erecta glabra, foliis apathulatis coriaceis apice crenatis confertis venoso-reticulatis, corymbo composito subglutinoso. $\frac{1}{2}$ in Brasiliâ. Species dubia ab anot. ipeo in system: suo omisa. $\frac{1}{2}$.

§. 2. Spiciformes, nempè volubiles, capitulis secus ramos elongatis sessilibus spicatis.

23. *M. HASTATA* (Willd. sp. 3 p. 1742), fruticosa scandens, caule tereti juniore subpubescente, foliis petiolatis subcordato-hastatis subdentatis acuminatis glabris, capitulis secus pedunculos axillares terminalesque aphyllous spicatis ad axillam bractearum involucri equalis solitariis, invol. squamis 4 oblongis lanceolato-ovatis submucronatis, achænio pulverulento-velutino. $\frac{1}{2}$ inter fruticeta montium Jamaicæ. Swartz! obs. 299. Kleinia, etc. Browne jam. 316. t. 34. f. 3. Flores albi suaveolentes. Sapor herbæ amarus. (v. s.)
24. *M. POLYSTACHYA*, fruticosa scandens, ramis teretibus petiolisque hirsutis, foliis petiolatis ovalibus lanceolatisve acuminatis integerrimis penninerviis supra scabris subtus præcipuè secus nervos hirsutis seu villosis, paniculâ terminali compositâ spicis pluribus distantibus oppositis elongatis et terminali longiore, capitulis secus rachin sessilibus bracteolâ lineari duplò longioribus, invol. squamis 4 lanceolato-linearibus acutis, achænio glanduloso. $\frac{1}{2}$ in convalliis et paludosis circa Bahiam legerunt cl. Salzmänn et Blanchet (n. 1033 et 1885!). Cor. pallide flavescentes. Petiol. 4-9 lin. longi. Limbus fol. 3 $\frac{1}{4}$ poll. longus, 12-18 lin. latus. Capit. 4 lin. longa. (v. s.)
25. *M. PSILOSTACHYA*, suffruticosa scandens, ramis teretibus petiolisque pube ferrugineâ subvelutinis, foliis petiolatis ellipticis acuminatis basi subacutis integerrimis supra subscabridis nervis villosis subtus puberulis et crebrè glandulosis, paniculâ terminali compositâ, spicis pluribus infernè subapproximatis, terminali longiore, capitulis secus rachin distantibus, bracteolâ lineari invol. 3° 4°ve brevior, invol. squamis dorso vix puberis apice mucronatis, mucrone patulo sæpè deciduo, achænio glandulifero. $\frac{1}{2}$ in Peruvâ? ex coll. Poeppig. Affinis *M. polystachyæ* sed pube minore ferrugineâ, spicis, pappo rufo, etc., differt. (v. s. comm. à cl. invent.)
26. *M. SCABRA*, fruticosa scandens, ramis teretibus petiolisque pube ferrugineâ scabro-villosis, foliis petiolatis ovatis basi obtusis apice acuminatis dentato-serratis penninerviis supra junioribus sparse scabris adultis subglabris subtus reticulatis secus nervulos piloso-scabris inter nervos glanduliferis, racemis spicatis elongatis nunc terminalibus subsolitariis nunc in paniculam dispositis oppositis axill. terminalibusque, bracteolis linearibus pedicelli longitudine, invol. squamis apice mucronatis subrecurvis, achænio glabro. $\frac{1}{2}$ in Cayennâ olim legit cl. Patris. Petiol. 4-6 lin. Limb. fol. 3 $\frac{1}{2}$ poll. longus, 1 $\frac{1}{4}$ poll. latus. Pappus rufus. Rami foliaque obscure ferruginei. (v. s.)
27. *M. HONSTONIA* (Willd. sp. 3. p. 1742), fruticosa glabra scandens, foliis petiolatis ovatis acuminatis integerrimis, paniculâ terminali compositâ, racemulis oppositis elongatis; capitulis pedicellatis, invol. squamis 4 equalibus. $\frac{1}{2}$ in Vera-Cruce legit olim Houston. Eupatorium Linn. h. cliff. 396. n. 6. Eup. Honstonia Linn. sp. 172. An species homonyma Swartzii hoc aut ad *M. Badieri* referenda? An. Eup. Honstonia Mill. dict. n. 16, quæ dixit folia cordata et capitula spicato-racemosa, huc pertinet? Spec. mihi dubia. $\frac{1}{2}$.
28. *M. SELLOI* (Spreng. syst. 3. p. 421), fruticosa scandens, ramis angulatis ferrugineis, foliis ovato-oblongis acuminatis integerrimis triplinerviis subtus

ferrugineis, spicis paniculato-ramosissimis, capitulis remotiusculis. 5 (in Brasiliâ. Cæst. ign.

§. 3. *Stipulatæ, nempe volubiles, petiolis utrinquē ad basin auriculā foliaceā quasi stipulatis, capitulis non spicatis sed corymbosis paniculatisve.*

29. *M. STIPULACEA* (Willd. sp. 3. p. 1745), volubilis glabra, caule tereti, ramulis substriatis, foliis cordato-hastatis, lobis apiceque acuminatis, vix hinc inde denticulatis ad basin petioli utrinquē stipulā foliaceā donatis, ramis axill. terminalibusque tripartitis corymbiferis, capitulis longè pedicellatis, invol. squamis oblongis aequalibus vix subacutis, bracteolā minimā subulatā ab invol. subdistante, achænio glabro. (in Brasiliâ ad Rio-Janeiro. Eupatorium stipulaceum Vahl symb. 3. p. 94. excl. forsan Lam. syn. p. 108 citat. Capit. 4 lin. longa. Pappus rutilanti-rufus. (v. s.)

30. *M. MANETTIA*, volubilis, caule subangulato glabro, ramis basi hexagonis villosiusculis, foliis petiolatis glabris, infer. cordato-hastatis, lobis later. brevibus acuminatis divaricatis, super. cordatis lobis subrotundis, omnibus basi repando-crenatis apice acuminatis, auriculis stipulæformibus latis obtusis subdentatis inter petiolos (ut in Rubiaceis) sitis, ramulis floridis axill. terminalibusque corymbum oligocephalum apice gerentibus, capitulis pedicellatis, bracteolis linearibus acutis foliaceis pedicellisque pubescentibus, invol. squamis apice subscariosis mucrone acuminatis. (in Brasiliâ circa Rio-Janeiro legit cl. Lund. An Eupatorium crenatum..... p. 19. t. 4. f. 1? sed nec folia seniores obtusa nec bracteolas 2 ad basin invol. video. (v. s. comm. à cl. Lund.)

31. *M. PTEROPODA*, volubilis glabriuscula, ramis striatis, fol. petiolisque alā foliaceā repando-undulatā marginatis basi in auriculam semi-reniformem auctis, limbis latè ovatis acuminatis basi et apice integris cæterum dentatis penninerviis, paniculā terminali foliis paulò longiore, ramis oppositis, ramulis capitula 3-5 subsessilia congesta gerentibus, bracteolā minimā, invol. squamis lanceolato-linearibus acutis achæniisque glabris. (in Brasiliâ circa Rio-Janeiro legit cl. Lund. Species distinctissima! Petioli 8-9 lin. longi. Limb. fol. 4 poll. longus, 2½ latus. Capit. 2-2½ lin. longa. (v. s. comm. à cl. inv.)

§. 4. *Ecordatæ, nempe volubiles exstipulatæ corymboso-paniculatæ, foliis indivisis basi non-cordatis.*

* *Foliis oblongis lanceolatisve.*

32. *M. CLEMATIDEA*, scandens?, ramis teretibus striatis tenuissimè puberulis, foliis oppositis alternisve petiolatis ovali-lanceolatis acuminatis basi subcuneatis dentibus utrinquē grossis 3-4 acutis notatis basi trinerviis glabrisculis, corymbo trichotomo laxo, capitulis pedicellatis nudis 5-floris, invol. squamis 7-9 ext. obtusis, int. brevioribus subacutis. — in Nepaliâ legit cl. Wallich. Mikania clematidea Wall cat. n. 3173. comp. n. 283. Pappus albidus. Achænium acutè 5-gonum pilis minimis sparsis subasperum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

33. *M. LIGUSTRIFOLIA*, fruticosa scandens glaberrima, ramis teretibus striatis, foliis breviter petiolatis oblongo-lanceolatis acuminatis penninerviis reticulatis suprā nitidis, pedunculis axillaribus et terminalibus racemosis in paniculam dispositis, capitulis pedicellatis, bracteolis linearibus ad basin pedicelli, invol. squamis 4 linearibus acutiusculis achæniisque glaberrimis. 2 (in Brasiliâ prov. Sancti-Pauli. Petiol. 2-3 lin. Limb. fol. 3 poll. longus, 6 lin. latus. Pappus rufus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 420 miss.)

34. *M. MYRIOCEPHALA*, fruticosa scandens glabra, ramis teretibus, foliis petiolatis lanceolatis seu ovali-lanceolatis sensim acuminatis basi subobtusis integerrimis penninerviis, paniculæ terminalis ramis pube brevi hirsutulis oppositis, ramulis capitula racemosa conferta innumera gerentibus, bracteolis ovatis

- aculis ad basin pedicelluli, invol. squamis oblongis obtusis achenisque glabris. 5 in Brasiliâ prope Bahiam ad Ilheos legit cl. Blanchet. Caulis teres nervulis subangulatus sarmentum vitis subreferens. Petiol. 4-5 lin. Limb. fol. 3-3½ poll. longus, 12-15 lin. latus. Cor ex sicco vid. pallidè flavescentes. (v. s. comm. à cl. inv.)
35. *M. ANDRACHNE*, caule..., ramis subvolubilibus striatis glabriusculis, foliis petiolatis ovalibus obtusè acuminatis integerrimis utrinquè lævibus basi quintuplinerviis, paniculâ laxâ corymbosâ, ramulis subpuberulis, bracteolis minimis aut nullis, invol. squamis oblongo-linearibus, acheniis angulatis minutè puberulis. 5 in Brasiliâ circa Bahiam ad Ilheos legit cl. Blanchet. Petiol. 6 lin. longi. Limb. fol. 3 poll. longus, 1½ latus, non malè Arbuti Andrachnis folia referentia. (v. s.)
36. *M. BUDDLEIÆFOLIA*, fruticosa scandens?, ramis teretibus striatis petiolis foliisque subtus pube tomentoso-pulverulentâ canis, foliis petiolatis lanceolatis seu ovato-lanceolatis basi subobtusis apice acuminatis integerrimis penninerviis suprâ glabris, paniculâ ramosissimâ polycephalâ, ramis oppositis, capitulis secus ramulos subsessilibus conferto-racemulosis, bracteolâ oblongâ canescente brevi, invol. squamis oblongis obtusis glabris apice ciliatis, achenio glabro. 5 (P) in Brasiliâ legit cl. Lhotsky. Petiol. 8 lin. longus. Limb. fol. 3 poll. longus, 15-18 lin. latus. Pappus sordidus. (v. s. comm. à cl. invent.)
37. *M. CLAVELLATA*, scandens?, caule tereti sparsè puberulo oppositè ramuloso, foliis breviter petiolatis elongato-lanceolatis utrinquè acuminatis distanter serratis trinerviis suprâ sparsè puberis subtus pubescenti - hirtellis, ramulis petiolisque villosis-hirtellis, paniculâ ramulorum quemquè terminantis ramis oppositis subcorymbiferis, capitulis pedicellatis, bracteolâ parvâ acutâ, invol. squamis oblongis obtusis glabriusculis, achenio pubero. (P) in Brasiliâ prov. Minarum General. ad Mariannam legit cl. Vauthier (pl. exs. n. 287). Stigmata apice insigniter clavellata! Capit. parva. Fol. 2 poll. longa, 6-7 lin. lata, petiolo 2-3 lin. (v. s. comm. à cl. inv.)
38. *M. KROCLADA*, volubilis?, ramis teretibus hirsutis confertâ rufâ petiolisque velutino-hirsutis, foliis petiolatis oblongo-seu ovali-lanceolatis acuminatis integerrimis membranaceis pube sparsa subglandulosâ suprâ secus nervos venasque dispositâ scabridis, paniculâ terminali foliis intermixtâ, ramis axillaribus, ramulis capitula 3-5 pedicellata conferta gerentibus, bracteolis ad basin pedicelli vix eo longioribus, invol. squamis oblongo-linearibus acutis dorso puberulis, acheniis glabris. (P) in Brasiliâ. Cor. ex sicco vid. pallidè flavescentes. Petiol. hirsutus 6 lin. longus. Limb. fol. 3½ poll. longus, 15 lin. latus. (v. s.)
39. *M. SMILACINA*, volubilis glaberrima, ramis striatis, foliis breviter petiolatis ovali-lanceolatis callosis-acuminatis rigidis integerrimis basi 5-7 nerviis cæterum reticulatis, paniculâ terminali distanter et oppositè ramosâ, ramis apice capitula congesta sessilia gerentibus, invol. squamis ovalibus obtusis, bracteolâ squamis simili. (P) in Brasiliâ prov. Minarum General. legit cl. Vauthier (pl. exs. n. 290). Petiol. 5 lin. longus. Limb. fol. 3 ½ poll. longus, 15 lin. latus. Capit. 8-10 arcuè congesta, 3 lin. longa. Pappus albus crassus. (v. s. comm. à cl. inv.)
- ** foliis obovatis.
40. *M. OBTUSATA*, fruticulosa scandens? glabra, ramis teretibus substriatis, foliis breviter petiolatis obovato-oblongis basi cuneatis apice obtusis subintegerrimis subtus pallidis venis reticulatis, paniculâ laxâ ramis oligocephalis oppositis, ramulis ultimis brevibus sæpè trifidis, capitulis breviter pedicellatis, bracteolâ minimâ, invol. squamis oblongis obtusis. 5 (P) in Brasiliâ prov. Sancti-Pauli. Petiol. 2 lin. Folia ferè 15 lin. longa, 5-6 lata. Pappus rufus. Invol. fusca, margine rufa submembranacea. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 472 miss.)
41. *M. OBOVATA*, volubilis glaberrima, ramis teretibus striatis, foliis petiolatis latè obovatis obtusissimis integerrimis subcarnosis, basi cuneatis trinerviis, paniculâ terminali oppositè ramosâ, ramis apice puberis paniculato-corym-

- basis, capitulis pedicellatis, bracteolâ lineari invol. dimidiò breviori, invol. squamis oblongo-linearibus subobtusis apice subciliatis, achenio glabro. (in convallibus Brasilie circa Bahiam legit ocul. Salzm. Flores albi. Petiol. 4 lin. Limb. fol. bipollic. sesquipollicem latus. (v. s. comm. à cl. inv.)
42. *M. SALZMANNIÆFOLIA*, fruticosa scandens glaberrima, ramis teretibus, foliis petiolatis obovatis ovalibusve basi subcuneatis apice obtusis integerrimis coriaceis lucidis penninerviis subtus reticulatis, paniculæ ramosissimæ laxæ ramulis subangulatis sparse puberulis, capitulis secus ramos racemosis distantibus breviter pedicellatis, bracteolâ ad basin pedicelli parvulâ, invol. squamis oblongo-linearibus obtusis, achenio glabro. ♀ in aridis Brasilie circa Bahiam legit cl. Salzm. Petiol. 3 lin. longa. Limb. fol. 20 lin. longus, 15 latus. Species affinis *M. obovatis* sed consist. fol. coriaceâ nec carnosâ et plurib. aliis notis faciliè distincta (v. s. comm. à cl. Salzm.)
43. *M. PSYCHOTRIODES*, fruticosa scandens? glabra, ramis striato-sulcatis, foliis approximatis ovalibus basi breviter cuneatis vix petiolatis triplinerviis obtusis integerrimis, paniculæ terminalis ramis pubescenti-hirtellis oppositis apice subcorymbosis, capitulis breviter pedicellatis fasciculatis, bracteolâ oblongâ invol. triplò brevior, invol. squamis oblongis obtusis glabrisculis, achenio glabro. ♀ in Brasiliâ circa Bahiam legit cl. Blanchet. Affinis *M. erithalinæ*. Folia sesquipollicem longa, ferè pollicem lata. Pappus pallidè rufescent. Cor. subflavescentes. (v. s. comm. à cl. Blanchet.)
44. *M. ERITHALINA*, fruticosa scandens? glaberrima, ramis teretibus, foliis sessilibus obovatis cuneatis subacutis integerrimis, corymbo tripartito terminali, capitulis pedicellatis confertis, bracteolâ lineari acutâ invol. brevior, invol. squamis ovali-lanceolatis mucronato-acuminatis glabris, achenio pubero. ♀ in Brasiliâ circa Bahiam in monte Serrato legit cl. Blanchet. Folia circ. 3 poll. longa, 15-16 lin. lata. Capit. 4 lin. longa. Invol. subfufcescentia. Habitat ramis in herb. ferè *Erithalis* fruticosæ. (v. s. comm. à cl. inv.)

*** *Foliis latè ovalibus ovatis aut subrotundis.*

45. *M. BANISTERIÆ*, fruticosa scandens?, ramis teretibus pube sericeâ molli canis, foliis oppositis petiolatis ovatis basi obtusis apice sensim acuminatis integerrimis penninerviis, supernè junioribus setosis, adultis nervis exceptis glabris, subtus densè sericeis, paniculæ terminalis ramis sericeo-lanatis ramulis capitula paucâ subsessilia congesta gerentibus, invol. squamis et bracteolâ ovalibus obtusis glabris. ♀ in Brasiliâ? Petiol. 4-8 lin. longi. Limb. fol. 5 poll. longus, 2 $\frac{1}{4}$ poll. latus, supernè reticulo prominulo venarum notatus. Pappus rutilans. (v. s. ex coll. cl. Poeppig.)
46. *M. ARGYREÆ*, fruticosa scandens? ramis teretibus petiolisque pube densâ hirsutâ confertâ molli velutinis, foliis petiolatis latè ovatis subrotundisve submuticis aut superioribus angustè et acutè cuspidatis basi 5-nerviis aut quinuplinerviis integerrimis suprâ sparse setulosis secus nervos villosos subtus undiquè sericeo-villosis, paniculâ terminali, ramis oppositis, infer. axillaribus folii longitudine, capitulis ad apices ramulorum subternis subsessilibus, invol. squamis bracteolisque ovatis obtusis glabris. ♀ in Brasiliæ prov. Sancti-Pauli. Affinis *M. Banisteriæ*. Petiol. 9-10 lin. Limb. fol. 3 poll. longi, 2 poll. lati. (v. s. comm. à cl. Thoremin.)
47. *M. GUACO* (H. et Bonpl. pl. equin. 2 p. 84. t. 105), herbacea volubilis, ramis teretibus sulcatis hirtis, foliis petiolatis ovatis subacuminatis basi breviter angustatis remotè dentatis reticulato-venosis suprâ scabriusculis subtus hirtis, corymbis axillaribus pedunculatis oppositis, capitulis subternis sessilibus, bracteolis linearibus inv. brevioribus, inv. squamis lineari-oblongis obtusis pubescentibus, achenio glabro. ♀ in calidissimis humidis ad ripam fluvii Magdalenæ ubi *Guaco* dicitur et antidotum contra serpentum ictus præstans habetur. H. B. et Kunth nov. gen. am. 4. p. 136. *M. Huaco* Rieux in Cav. anal. n. 18. p. 316 ex Spreng.
48. *M. TAPALLANA* (H. B. et Kunth nov. gen. am. 4. p. 137), herbacea volubi-

lis, ramis sexangularibus hirtis, foliis petiolatis elliptico-ovatis acutiusculis basi brevissimè angustatis integerrimis reticulato-venosis suprâ scabris subtus pubescenti-hirtis, corymbis axillaribus et terminalibus, capitulis ternis sessilibus, bracteolis linearibus inv. brevioribus, invol. squamis lineari-oblongis obtusis pubescentibus, achænio glabro. γ in Peruvia calidis humidis ad ripam fluv. Daude prope Guayaquil. Aff. M. Guaco. M. Tafallæ Spreng. syst. 3 p. 422.

49. M. TRINITARIA, fruticosa scandens, caule tereti glabro, ramis floridis striato-angulatis apice ferrugineo-subvelutinis, foliis petiolatis ovatis acuminatis integerrimis suprâ glabris subtus penninerviis reticulatis inter venas pube brevi ferrugineâ velutinis, paniculâ corymbosâ trichotomâ, ramulis apice tricephalis, capitulis sessilibus 2 lateralibus bracteâ lanceolatâ brevi donatis, medio ebracteato, invol. squamis 4 lineari-lanceolatis acutis glabrinsculis, achænio glabro. γ in insulâ Trinitatis Sieb. f. Trin. exs. n. 182. Petiol. 9 lin. longus. Limb. 4 poll. longus, 2 poll. latus. Capit. 4 lin. longa. Pappus rufus. Nomen aliqd ad patriam et ad capitula terna. (v. s. ex coll. Sieb.)

50. M. LEVIS, fruticosa glabra scandens, caulibus teretibus, ramulis striato-angulatis, foliis petiolatis latè ovatis obtusè acuminatis integerrimis basi subtruncatis trinerviis utrinquè subnitidis, paniculâ ramosissimâ laxâ, ramulis 1-2-3-cephalis, capitulis pedicellatis minutè bracteolatis aut medio sessili ebracteolato, invol. squamis 4 oblongis vix acutis, achænio juniore subpuberulo. γ in Brasiliâ. Petiol. pollicaris. Limb. 3-3 $\frac{1}{2}$ poll. longus, 2-2 $\frac{1}{2}$ latus acuminè obtusissimo insignis. Capit. 2 $\frac{1}{2}$ lin. longa. Pappus pallidè rufescens. (v. s.)

β . angustior, foliis paulò brevius petiolatis angustius ovatis sensim et obtusè acuminatis. γ in montosis Tejuco prope Rio-Janeiro legit cl. Lund. Petiol. 8-9 lin. Limb. fol. 3 poll. longus, 15 lin. latus, nempe dimidio angustior quàm var. α . Inflorescentia similis. (v. s. comm. à cl. Lund.)

51. M. PANICULATA, subscandens glaberrima, ramis teretibus fistulosis, foliis petiolatis ovatis abruptè et obtusiusculè acuminatis integerrimis subtus pallidis, paniculâ terminali ramosissimâ laxâ, ramis oppositis glabris, capitulis æcus ramulos spicatis distantibus, bracteolâ minimâ, invol. squamis lineari-oblongis obtusis achæniisque glabris. γ in Brasiliâ circa Rio de Janeiro arboreis altissimas paniculis suis involvit. Flores albi odorati ex Lund. Caulis diam. pollic. Petiol. 4 pollic. Limb. fol. 4 poll. longus, 1 $\frac{1}{2}$ latus. Panicula pedalis. Pappus pallidè rufescens. (v. s.)

52. M. BRACTEOSA, fruticosa scandens glaberrima, ramis teretibus apice striatulis, foliis petiolatis ovatis basi obtusis apice subacuminatis puncto calloso terminatis triplinerviis nunc integerrimis nunc basi repandis dentatisve, paniculæ ovalis brevis terminalis ramis oppositis, capitulis spicato-congestis, bracteolâ amplè concavâ ovali invol. sublongiore, invol. squamis oblongo-linearibus vix acutis, achænio glabro. γ in Brasiliâ ins. Sanctæ Catharinæ leg. cl. Gandichaud et Bacle. Petiol. pollicares. Limb. fol. 2 $\frac{1}{2}$ poll. longus, pollicem latus. Pappus rufus. Cor. ex sicco purpurascens. (v. s.)

53. M. BADIERI, fruticosa scandens glaberrima, ramis teretibus, foliis petiolatis ovalibus abruptè acuminatis integerrimis subtriplo- sen quintuplinerviis reticulatis coriaceis, paniculæ terminalis ramis oppositis apice trifidis, ramulis capitula subsessilia conferta gerentibus, bracteolis linearibus ciliatis, invol. squamis lineari-oblongis apice subobtusis villosis-hirsutis, achænio glabro. γ in Guadalupâ legit olim Badier; an fortè etiâ in Jamaicâ si Eupat. Houstonia Swartz obs. p. 300 huc referendum. Species insignis foliis apice abruptè angustatis et venis singulari modo ramoso-subreticulatis. (v. s.)

54. M. LAURIFOLIA (Willd. sp. 3. p. 1746), fruticosa volubilis glaberrima, ramis angulatis, foliis breviter petiolatis ovatis obtusiusculis basi acutis integerrimis reticulato-triplinerviis subcoriaceis nitidis, corymbis paniculatis confertis axill. oppositis et terminalibus, capitulis ternis subsessilibus, bracteolis . . . ,

- invol. squamis oblongis obtusis apice ad dorsum puberulis, achenio glabro. 5 (in Amer. merid. (Mutis), temperatis Novæ Granatæ prope Mariquita (H. et B.) nec in Mexico ut incaute Persoon indicavit. Humb. B. et Kunth nov. gen. sm. 4. p. 137. t. 350. *Cacalia laurifolia* Mutis in Linn. f. suppl. 351. An stirps Mutisiana quæ dicitur erecta eadem ac Humboldtiana?
55. *M. CISSAMPYLINA*, scandens glabra, caule tereti, foliis petiolatis ovatis subcordatis obtusis integerrimis, ramis floridis axillaribus terminalibusque apice trifidis corymbiferis, invol. squamis 4 oblongo-linearibus acuminatis, 5^a ext. lineari cæteris dimidiò breviorè, achenio glabro. (in Sancto Domingo. Capit. 5 lin. longa. Petiol. 9-10 lin. (v. s.)
56. *M. PLATYPHYLLA*, scandens? glabra, ramis teretibus, foliis petiolatis latè ovatis subrotundis apice abruptè et obtusiusculè cuspidatis integerrimis, ramis flor. axillaribus oppositis paniculatis inter capitula tenuiter puberulis, ramulis capitula plurima spicata congesta sessilia gerentibus, invol. squamis linearibus glabris subobtusis, bracteolâ acutâ brevi, acheniis glabris. (in Peruvia? ex coll. Pæppig. Rami floridi in specim. meo incompleto retrofracti, folio vix longiores. Pappus rufus. Petiol. pollicares. Limb. fol. 4 poll. diam., cuspidè 3-4 lin. longo. (v. s. comm. à cl. inv.)
57. *M. ATTENUATA*, caule herbaceo tereti volubili cum petiolis et pedunculis pubè minimâ subpulverulentis, foliis petiolatis latè ovalibus utrinquè longè acuminatis integerrimis membranaceis glabris penninerviis, paniculis axillaribus et terminalibus, ramis oppositis, ultimis apice tricephalis, invol. squamis 4 oblongis obtusis bracteolis 2 acutis minimis stipatis, acheniis glabris teretiusculis. — in Peruviz montibus Guanoccensis legit cl. Hænke. Folia 4 poll. longa, 2 poll. lata. Pappus rufus. (v. s. in h. Hænke ab ill. de Sternberg comm.)
58. *M. LINDLEYANA*, scandens glabra, caule tereti, foliis petiolatis ovatis acuminatis integerrimis trinerviis, ramorum floridorum sessilibus multò minoribus, paniculæ ramis oppositis apice corymbiferis, capitulis pedicellatis, bracteolis ovali-oblongis foliaceis invol. æqualibus, invol. squamis ovali-oblongis obtusis mucronatisve acheniisque glabris. (in Guianâ. Valdè affinis *M. Hookeriana*, sed capitulis pedicellatis et bracteolis faciliè differt. (v. s. in h. Lindl. à cl. Hooker comm.)
59. *M. HOOKERIANA*, suffruticosa scandens? glabra, ramis teretibus vix striatis, foliis petiolatis ovatis ellipticisve basi obtusis apice acuminatis integerrimis trinerviis, paniculæ terminalis ovatæ ramis oppositis, capitulis secus ramos sessilibus approximatis, bracteolis parvis ciliolatis, invol. squamis oblongo-linearibus obtusis apice subciliatis, achenio glabro. 5 (in Guianâ. Petiol. 9-12 lin. longi. Limb. fol. 2 poll. longus, pollicem latus. (v. s. in h. Lindl. à cl. Hooker comm.)
60. *M. AMARA* (Willd. sp. 3. p. 1744), fruticosa scandens, caule tereti glabriusculo, foliis petiolatis ovatis subcordatisve acuminatis subintegerrimis suprà sparsè et tenuissimè scabris subtus pubescenti-hirtellis secus nervos subtomentosis, paniculis axill. terminalibusque divaricatis oppositè ramosis, capitulis confertis ad apices ramulorum ternis subsessilibus, bracteolis brevibus acutis, invol. squamis oblongo-linearibus obtusis acheniisque glabris. 5 (in Guianâ secus rivulos et ex Vahl in insul. Caribæis? *Eupat. parviflorum* Aubl. guian. 2. p. 797. t. 315. sed panic. delineata nimis laxa. *Eup. vineæfolium* Lam. dict. 2. p. 410, sed folia utrinquè glabra dicuntur. *Eupat. amarum* Vahl symb. 3. p. 93, sed pubes fol. dicitur stellata. (v. s. in h. Lindl.)
61. *M. LATIFOLIA* (Smith in Rees cycl. 23. n. 8), volubilis glabra, foliis ovatis acuminatis subintegerrimis subtus pallidis, capitulis corymbosis, bracteis lanceolatis seu ellipticis sessilibus subasperis. — in insulâ Sanctæ Luciz. Affinis dicitur *M. amara*, sed descr. non sufficit. †
62. *M. PYRIFOLIA* (Bojer! in litt. 1835 sub *Trixide*), suffruticosa scandens glaberrima, caule tereti, foliis petiolatis ovatis aut subcordatis acuminatis

dentatis coriaceis nitidis penninerviis inter nervos tenuè reticulatis, paniculâ terminali irregulari, capitulis ad apices ramulorum confertis angustis, invol. squamis acutiusculis, bracteolâ minimâ, achæniis glabris eglandulosis. ζ \curvearrowright in prov. Betsilon ins. Madagascar legit cl. Bojer. Flores pallidè flavescentes. Capitula 2 lin. longa. (v. s.)

63. *M. SPACIOSA*, caulibus suffruticosis volubilibus teretibus, ramis petiolis pedunculis folisque subtus pube confertâ brevi velutinis, foliis longè petiolatis, infer. subcordatis, super. ovatis, omnibus acuminatis suprà minutissimè puberulis subtus reticulatis, paniculâ amplâ laxâ trichotomâ, capitulis ternis, bracteolis acutis minimis, invol. squamis lineari-oblongis obtusis dorso subpuberulis, achæniis pentagonis glabris. ζ \curvearrowright in Peruanis montibus Orinocensibus legit cl. Hænke. Petioli 2 poll. longi. Limb. fol. 6 poll. longus, 3-3 $\frac{1}{2}$ poll. latus. (v. s. in h. Hænk. ab ill. de Sternberg comm.)

§. 5. *Cordiformes, nempe volubiles exstipulatæ corymboso-paniculatæ, foliis basi cordatis integris aut repando-dentatis.*

64. *M. HÆNKIANA*, caule herbaceo volubili glabro, foliis longè petiolatis cordiformibus acutis basi subdentatis membranaceis suprà glabris subtus pubescentibus subtomentosis, ramis floridis axillaribus terminalibus paniculatis, ramulis oppositis subspiciformibus involacrisque minutissimè puberulis, achæniis angulatis glaberrimis. — in montanis Peruanis Orinocensibus legit cl. Hænke. Affinis videtur *M. Orinocensi*. Inflor. media inter spiciformes et paniculatas. (v. s. in h. Hænk. ab ill. de Sternberg comm.)

65. *M. ORINOCENSIS* (H. B. et Kunth nov. gen. am. 4. p. 134), fruticosa volubilis glabra, ramis striatis, foliis petiolatis subrotundo-ovatis cordatis acuminatis basim versus angulato-dentatis reticulato-5-nerviis, corymbis paniculatis, capitulis pedicellatis solitariis, bracteolâ invol. dimid. breviorè, invol. squamis oblongo lanceolatis acutis achæniisque glabris. ζ \curvearrowright in humidis insulâ Pararum in Mission. Orinocensibus. Petiol. 8 lin. longus. Limb. fol. sesquipollic. longus, pollicem latus. Valdè affinis *M. micranthæ*.

66. *M. MICRANTHA* (H. B. et Kunth nov. gen. am. 4. p. 134), herbacea volubilis glabra, ramis striatis, foliis longè petiolatis subrotundo-ovatis cordatis acuminatis sinnato-dentatis reticulato-5-nerviis, lobis baseos divaricatis, corymbis paniculatis, capitulis pedicellatis solitariis, bracteolâ invol. duplò breviorè, invol. squamis oblongo-lanceolatis acutis achæniisque glabris. \curvearrowright in umbris prope Caripe Cumanensium. Petioli 2 $\frac{1}{2}$ pollices longi. Limb. fol. 2 pollic. longus, sesquipollic. (in textu verisim. erron. sesquilineam) latus.

67. *M. SIRBERIANA*, scandens glabriuscula, ramis teretibus striatis, foliis petiolatis subcordatis subacuminatis in parte latiore subrepandis vix tenuissimè pubescentibus, ramis apice corymbiferis, capitulis congestis parvis, invol. squamis submembranaceis dorso subpubescentibus, 3 subacutis, 4^a longiorè magis acutâ, 5^a exte. omnibus breviorè angustiorè, achænio glabriusculo subglanduloso. \curvearrowright in insulâ Trinitatis. Eupatorium Sieb. f. trin. exs. n. 225. Specim. manum, sed capitulis parvis à plerisque facillè differt. (v. s. ex coll. Sieb.)

68. *M. LOXENSIS* (H. B. et Kunth nov. gen. am. 4. p. 135), herbacea volubilis glabra, ramis sulcatis, foliis petiolatis ovatis angustato-acuminatis breviter cordatis remotè et obsolete dentatis reticulato-trinerviis, corymbis paniculatis oppositis, capitulis pedicellatis solitariis, bracteolâ invol. dimidio breviorè, invol. squamis oblongo-linearibus acutiusculis achæniisque glabris. \curvearrowright \mathcal{L} in temperatis inter Alto de Pulla et urbem Loxa Novo Granat. Petiol. 7-8 lin. longus. Limb. fol. 2 $\frac{1}{2}$ poll. longus, 15-16 lin. latus.

69. *M. MICROPTERA*, subvolubilis minutè puberula, caule hexagono, angulis in alam angustissimam subfoliaceam productis, foliis petiolatis cordatis sinu lato, aniculis rotundatis, apice acuminatis mucrone calloso, margine subintegerrimis, paniculâ terminali oppositè ramosâ laxâ, capitulis distantibus pedicel-

latis, bracteolâ oblongâ minimâ, invol. squamis oblongo-linearibus obtusis achæniisque glabris. (in Brasiliâ prope Bahiam (Blanchet! n. 1710) et in Peruvîâ? (ex coll. Poeppig!). Petioli pollicares. Limb. fol. 2 poll. longus, sesquipollicem latus. Specim. Poeppigianum ferè glabrum sed non videtur diversum. (v. s.)

70. *M. BATATÆFOLIA*, scandens glabra, ramis teretibus, foliis petiolatis cordatis acuminatis in parte latiore repando-angulatis, limbo vix petiolo longiore, ramis apice laxè corymbiferis, capitulis pedicellatis, invol. squamis 4 acuminatis, 5^a ext. cæteris brevior angustâ adpressâ, achænio glabro. (in insulâ Cubâ circa Havanam legit indef. Ramon de la Sagra. Formâ foliorum accedit ad *M. congestam*, sed inflorescentiâ faciliè differt. (v. s. comm. à cl. inv.)

71. *M. TAMOIDES*, scandens glaberrima, caule tereti, ramulis striatis, foliis petiolatis cordatis acuminatissimis in parte latiore subrepandis, ramis nudis apice corymbum confertum gerentibus, capitulis breviter pedicellatis, invol. squamis 4 oblongo-linearibus acutis, 5^a ext. lineari brevi, achænio glabrescens. (in ins. Cubâ juxta Havanam legit cl. A. de la Ossa. Petioli 7-8 lin. longi. Species glabritis perfectâ et capitulis congestis faciliè à cæteris distincta. (v. s. comm. à cl. inv.)

72. *M. CONGESTA*, scandens, caule tereti sparsè puberulo, ramis striato-sulcatis, foliis petiolatis cordatis acuminatis in parte latiore sinuato-angulatis suprâ glabriusculis subtus secus nervos pubescentibus, sinu exciso semitondo, pedunculis axill. terminalibusque capitula corymboso-congesta gerentibus, capitulis pedicellatis, invol. squamis 4 mucrone acuminatis, 5^a ext. etiam acuminatâ cæteris longiore, achænio glabro. (in Porto-Ricco legit cl. Bertero. *M. scandens* ex Porto-Ricco Spreng.! syst. 3. p. 421 in h. Balb. ipa. teste, sed excl. syn. Capitula vix 3 lin. longa. Limbus intra sinum sæpè cuneatus. Sinus nervis limitatus. (v. s. comm. à cl. Balbis.)

73. *M. GLOMERATA* (Spreng. syst. 3. p. 421), fruticosa scandens, foliis cordatis angulatis glaberrimis coriaceis sub-5-nerviis, lobis baseos divaricatis inæqualibus, capitulis glomerato-paniculatis. 5 (in Brasiliâ. Cæt. ign. †.

74. *M. P. ALATA*, volubilis glabra, caule striato, foliis petiolatis ovatis profundè cordatis acuminatis integerrimis, cymis compositis, pedicellis ad latus interior angustè alatis, bracteis lanceolatis, invol. squamis 4-5 obtusiusculis trinerviis. 2 in nemorosis Guianæ prope Essequibo. Kleinia alata Mey. esseq. 249. Pappus roseus. Achænia sulcata. Ipse cl. auctor de genere dubitat et hic ex habitu potiùs referenda videtur.

75. *M. TESTUDINARIA*, scandens glabra, ramis angulatis vix hinc indè puberulis, foliis breviter petiolatis cordatis acuminatis rigidis integerrimis sinu angusto, nervis 3 subtus prominulis, ramis tripartitis corymbosis, capitulis pedicellatis, bracteâ ovatâ acutâ paulò ab invol. distante, invol. squamis 4 oblongis acutis glabris, achænio subpilosulo. (in Brasiliæ prov. Minarum General. prope Mariannam legit cl. Vauthier (pl. exs. n. 330) et circa Rio-Janeiro cl. Lund. Folia ferè Testudinariæ, sed minora et rigidiora. Petiol. 4 lin. Limb. circ. pollicem longus, 7 lin. latus. (v. s. comm. à cl. inv.)

76. *M. CARNOSULA*, fruticosa scandens, ramis teretibus apice villosis, foliis cordatis obtusis seu reniformi-cordatis subcarnosis suprâ sparsè puberulis subtus villosa-hirsutis subintegris sinu lato ad basin petioli stipulas foliaceas duas gerentibus, pedunculis apice corymbiferis, capitulis breviter petiolatis, invol. squamis 4 ovali-oblongis dorso pubescentibus, 2 obtusis, 2 subacutis, bracteolâ subulatâ minimâ, achænio glabro. 5 (in Brasiliæ montibus Macahe prope Rio-Janeiro mense julio flor. legit cl. Lund. Petiol. 6 lin. longi. Limbus 6-8 lin. longus, 10-12 latus. (v. s. comm. à cl. inv.)

77. *M. OPIFERA* (Mart. in Isis 1824. p. 6. p. 583), volubilis glabra, caule angulato, foliis petiolatis cordatis repando-dentatis vel subintegerrimis, adnatis obtusiusculis, capitulis corymboso-paniculatis pedicellatis, invol. squa-

- mis oblongis subacutis, bracteolâ lanceolatâ invol. paulò brevior. (in Brasiliâ ubi ex Mart. dicitur *Erba de Cobra*. (v. s.)
78. *M. DENTICULATA* (Willd. sp. 3. p. 1744), scandens glabriuscula, ramis acutè angulatis, foliis petiolatis cordatis obtusiusculis subdenticulatis scabris, corymbis terminalibus, capitulis umbellatis pedicellatis, bracteolis. . . . invol. squamis linearibus. (in Surinamo (Vahl), in sylvis Xalappensibus (Less.). Eupatorium denticulatum Vahl symb. 3. p. 93. Less. in Linnæâ 1830. p. 139. Spreng. syst. 3. p. 421. excl. Raf. syn.
79. *M. NATALENSIS*, volubilis herbacea? tota breviter hirsuto-velutina, foliis petiolatis cordatis acuminatis sinuato-dentatis 5-7-nerviis, auriculis obtusis, corymbis axillaribus oppositis pedunculatis trifidis polyccephalis folio paulò longioribus, invol. squamis glabriusculis oblongis mucronato-acutis 4? (in Africâ australi-orient. ad Port-Natal legit cl. Drege! Petioli pollicares. Limb. fol. 1 $\frac{1}{2}$ poll. longus, pollicem latus. Pappus pallide rufescens. (v. s.)
80. *M. CAPEXIS*, caule suffruticoso volubili glabro, foliis petiolatis cordatis acuminatis dentatis suprâ glabris subtus hirsutiusculis, corymbis trichotomis, invol. squamis oblongo-linearibus subacutis dorso puberulis, achæniis parvis glandulos. — ad. Cap. Bonæ Spei (Dreg!) in sylvis Coloniz distr. orient. Burch! cat. geog. n. 3674. in distr. Uitenhage (Eckl!). Krebs! pl. cap. exs. n. 199. Eupatorium scandens Thunb. fl. cap. 627. non Linn. Simillima *M. scandenti*, sed diversa præsertim hirsutie paginâ fol. inferioris. Prima et usquè ad nuperrimos, sola generis species hucusque in Africâ australi nota. (v. s.)
81. *M. OXYOTA*, glaberrima, caule herbaceo volubili, foliis petiolatis acutè hastato-triangulatis basi subtruncatis quinquenerviis apice acuminatis subintegerrimis, corymbo amplo trichotomo polyccephalo, invol. squamis oblongo-linearibus lævibus acuminatis, achæniis glabris sparsè glandulos. (in Africâ australi ad Van Staades Rivier legit cl. Drege! Petioli sesquipollic. Limbus fol. 3 poll. longus, 18-20 lin. latus. Cor. albidæ. Pappus pallide rufescens. (v. s.)
82. *M. DIOSCOREÆFOLIA*, scandens herbacea, caule tereti glabriusculo, foliis petiolatis cordatis acuminatis margine repando-crenatis, sinu subrotundo, suprâ glabris seu pube sparsâ minimâ vix puberulis, subtus minutissimè glandulos, corymbis pedunculatis trichotomis, invol. squamis submembranaceis dorso subpuberulis, 3 mucronato-acutis, quarto obtuso, achæniis glandulos. (5? in ins. Madagascar legit cl. Goudot, locis *Ambanivoules* dictis. Flores nitidè albi. Petiol. pollicem long. Limb. fol. sesquipollicem longus et latus. Pappus rufescens. (v. s. comm. ab. amic. B. Delessert.)
- β ? *Bojeri*, foliorum sinu subangustiori, floribus flavis ex cl. Bojer. — in sylvâ Bé-Fouroun super arbores scandens. An flores potius ochroleuci aut albi? Trixis? dioscoreæfolia Bojer! in litt. 1835. (v. s. comm. à cl. Bojer.)
- γ ? *crenata*, caule hinc indè pilosiusculo, foliis papyraceis, floribus ex albo sublilacinis — in sylvis cum var. β , sed eâ multò minor. Trixis? *crenata* Bojer! in litt. 1835. Fortè ut var. β species propria? (v. s. comm. à cl. Bojer.)
83. *M. THUNBERGIIOIDES* (Bojer! in litt. 1835. sub Trixide), herbacea scandens glabriuscula pube nempe minimâ sparsâ pubescens præ. in paginâ folii superâ, caule tenui tereti, foliis petiolatis cordatis acuminatis latè crenato-dentatis, pedunculis terminalibus apice corymbois, invol. squamis 4 acuminatis glabriusculis, bracteolâ lineari foliaceâ dimidiam invol. longitudinem æquante, achænio angulato glabro. (1)? in ins. Madagascar prov. Bé-Tani-Mena in sylvâ Bé-Fouroun apricis super dumeta scandens. Flores flavi et herba fortè annua ex Boj. (v. s.)
84. *M. FLORIBUNDA* (Bojer! in litt. 1835, sub Trixide), herbacea scandens, caule tereti petiolisque hirsutulis, foliis cordatis subdentatis acuminatis utrinquè sparsè et minutissimè puberulis, paniculâ terminali amplâ ramosissimâ *polyccephalâ*, invol. squamis 4 acuminatis dorso subpuberulis, bracteolâ lineari

foliaceâ invol. dimidiam longit. æquante, achænio angulato glabro glandulifero. $\frac{1}{2}$ in sylvis umbrosis prov. Bé-Zonzong ins. Madagascar legit cl. Bojer. (v. s.)

85. *M. GONOCLEADA*, fruticosa scandens, ramis acutè hexagonis præsertim ad angulos villosis, foliis petiolatis cordatis acuminatis in sinu subcuneatis hinc inde dentatis suprâ sparse puberis subtus subvillosis, pedunculis axillaribus terminalibusque apice corymbosis, capitulis subfasciculatis, invol. squamis 4 oblongis dorso subvillosis, 3 acutis, 4^a obtusâ, bracteolâ lineari-acuminatâ, achænio elongato glabro acutangulo. $\frac{1}{2}$ in Cubâ juxta Havanam in sylvis humidis legit cl. R. de la Sagra et in Mexico circa Tampico de Tamaulipas cl. Berlandier. Pappus rufescens. A *M. Guaco* differt foliis cordatis et ramis angulatis. (v. s. comm. à cl. inv.)

β . *ambigua*, ramis hexagonis, foliis cordatis in sinu non cuneatis, summis vix cordatis, bracteolâ angustiori. $\frac{1}{2}$ in Mexico ad Tampico de Tamaulipas (Berl.), in Brasiliâ circa Bahiam (Salzm. ! Blanch. !). An species distincta? (v. s.)

86. *M. CONVULVULACEA*, suffruticosa scandens, caule ramisque teretibus junioribus pubescentibus, foliis petiolatis latè cordatis acuminatis subrepando-dentatis suprâ vix puberulis subtus pubescentibus subvillosis, ramis apice corymbiferis in paniculam laxè dispositis, capitulis breviter pedicellatis, invol. squamis 4 oblongis acutis dorso pubescentibus, adjectâ 5^a exteriore lineari foliaceâ cæteris dimidiò breviorè, achænio glabro. $\frac{1}{2}$ in Sancto Domingo legit cl. Poiteau. Affinis *M. gonocladæ*, sed ramis teretibus faciliè differt. (v. s.)

β . *Portoricensis*, ramulis striatis, foliis subintegerrimis summis minùs cordatis, ramis pubescenti-hirtellis, capitulis achæniisque paulò minoribus. $\frac{1}{2}$ in Porto-Ricco legit cl. Bertero. *M. amara* Bert. ! herb. non Willd. *M. cordifolia* Spreng. ! in h. Balb. syst. 3. p. 423. excl. syn. An fortè species propria? (v. s.)

87. *M. VOLUBILIS*, scandens, caule tereti subpubescente, foliis petiolatis cordatis crenatis acutis, lobis baseos æqualibus obtusis, venis subtus petiolisque subpubescentibus, corymbo paniculato trichotomo nudo, capitulis pedicellatis 4-floris, invol. squamis circ. 6-8, inter. 4 oblongis acutis, exter. brevioribus subpancioribus, pappo rubente. $\frac{1}{2}$ in ruderalis dumetis et sylvis Ind. or. ad Penang, Singapore, etc. (Wall. !), in Javâ (Blum. !). Eupatorium volubile Vahl symb. 3. p. 93. Eupatorium cordatum Burm. ind. t. 58. f. 2. Mikania volubilis Willd. sp. 3. p. 1743. Blum. ! bijdr. 904. Wall. ! cat. n. 3174. comp. 284. Fl. albi. Achæn. angulata glabra glandulosa. Variat pubescens aut glabra foliis plùs minùs crenatis aut subintegris. An satis à *M. scandenti* diversa? (v. s.)

88. *M. PARKERIANA*, volubilis, caule tereti minutissimè pubescente, ramis striato-sulcatis glabriusculis, foliis petiolatis cordatis acuminatis membranaceis pubescentibus, sparsâ minutissimâ sub lente scabriusculis in parte latiore subangulatis, sinu semirotnondo, corymbo terminali composito suprâ ultima folia subsessili, capitulis congestis subsessilibus, bracteolâ lineari, invol. squamis ovali-oblongis membranaceis acutè mucronatis achæniisque glabris. $\frac{1}{2}$ in Guianâ circa Demerari legit cl. Parker. Affinis *M. scandenti* et pubescenti, sed capitulis in glomerulos plurimos densè congestis et invol. mucronatis distincta. Petioli bipollicares. (v. s. comm. à cl. Parker.)

89. *M. SCANDENS* (Willd. sp. 3. p. 1743. excl. syn. Plum.), caule volubili glaberrimo, foliis petiolatis cordatis acuminatis basi repando-crenatis glabris, ramis axillaribus apice corymbum subfastigiatum gerentibus, invol. squamis 4 acuminatis subeiliolatis, adjectâ 5^a exteriore breviorè, achæniis glabris subeglandulosis. $\frac{1}{2}$ ad margines rivorum Amer. bor. à Canada australiori? (Hook.) ad Carolinam super. (Michx.) et fortè in Mexico ex specim. Berlandieriano (pl. exs. n. 2272) prope Victoria lectis et Bostonianis similissimâ. Pursh fl. 2. p. 517. Nutt. gen. 2. p. 136. Ell. sketch 2. p. 292. Lindl. in trans. hort. soc. Lond. 7. p. 2. p. 250. Eupatorium scandens Linn. sp. 1171. Jacq. Æ. rar. 3.

- t. 169. coll. 1. p. 108. *M. angulosa* Rafin. fl. lud. p. 62? — Fluk. alm. t. 163. f. 3. Flores albid; Pappus albus. (v. s.)
90. *M. rubescens* (Muhl. cat. 71. Nutt. gen. am. 2. p. 136), volubilis, caule tenenter pubescente, foliis petiolatis cordatis acuminatis basi repando-crenatis angulatisque, ramis axillaribus apice corymbum subfastigiatum gerentibus, invol. squamis 4 acuminatis ciliolatis et pubescentibus, adjectâ 5^a brevior ex. angustiore, achenio glabro subglanduloso. \curvearrowright in Carolinâ infer. et Georgiâ locis humidis. Ell. sketch a p. 293. Eupatorium scandens Walt. car. 198. ex h. Fraser! Flor. pallide purpurei odori. Species valdè *M. scandenti* affinis. Stirps Martinicensis huc à Spreng. relata verisim. diversa. (v. s.)
91. *M. LAXA*, volubilis, caule tereti glabro, ramis vix striatulis præsertim in paniculâ pube minimâ rufescente velutinis, foliis petiolatis cordatis acuminatis subintegerrimis trinerviis membranaceis, paniculâ oppositè ramosâ laxâ, capitulis longè pedicellatis secus ramulos subracemosis, bracteolâ oblongo-lineari acutâ invol. dimidiò breviorè, invol. squamis obtusis. \curvearrowright in Peruvîâ legit cl. Poeppig. Petioli pollicares. Limb. fol. 3 poll. longus, 2 poll. latus. Pappus rufus. (v. s. ex coll. Poeppig.)
92. *M. MICRODONTA*, suffruticosa scandens, ramis teretibus pube brevi velutino-hirsutulis, foliis petiolo brevi retrorsum hirsutissimo, limbo basi cordato ovato-lanceolato acuminato dentibus parvis raris notato suprà sparse setoso subtus densè et molliter hirsuto, ramis axillaribus terminalibusque paniculatis oppositè ramosis, ramulis capitula 3-5 congesta subsessilibus gerentibus, invol. squamis bracteolisque ovatis subobtusis glabris. \curvearrowright in Brasiliâ prov. Minarum General. prope Tejucco legit cl. Vauthier. (pl. exs. n. 291). Petiol. 3 lin. long. Limb. fol. 3-3 $\frac{1}{2}$ poll. longus, 12-15 lin. latus. (v. s.)
93. *M. MICROCEPHALA*, caule volubili tereti pube brevi subglandulosâ hispidulo, foliis petiolatis cordatis sinu angusto acuminatis dentatis basi 5-nerviis suprà sparse scabridis subtus præcipuè secus nervos subhirsutis, paniculâ amplâ oppositè ramosâ, capitulis secus ramulos spicato-racemosis subsessilibus, invol. squamis 4 mucronulatis dorso subpuberis, bracteolâ minimâ, achenio glabro. \curvearrowright in prov. Minarum General. Brasiliæ legit cl. Vauthier (pl. exs. n. 288). Pars ramorum superior ferè aphylla. Capitula vix 2 lin. longa. Fol. inf. sesquipollicem longa, et ferè lata. (v. s. comm. à cl. inv.)
94. *M. HIRsutissima*, ramis scandentibus teretibus pedunculis petiolis folisque subtus villosis hirsutissimis, foliis petiolatis cordatis acuminatissimis subintegerrimis suprà sparse puberis, ramis paniculam amplam ramosissimam myriocephalam gerentibus, invol. squamis 4 oblongis subacutis glabris, bracteolâ paulò infrà involucrium sitâ squamis simili, achenio glabro. \curvearrowright in Brasiliâ prov. Minarum General. prope Mariannam legit cl. Vauthier (pl. exs. n. 286). Petiol. 2 poll. Limb. fol. 5 poll. longus, 2 poll. latus. (v. s.)
95. *M. TRICHOPHILA*, fruticosa scandens, ramis teretibus pilis rufis confertis petiolis pedunculisque hispidissimis, foliis petiolatis suprà sparse setosis subtus villosis-hispidis, infer. subcordatis angulatis, super. ovatis, omnibus integerrimis acuminatis, paniculâ ramosissimâ, invol. squamis 4 obtusis vix puberis, bracteolâ subacutâ, achenio glabro. \curvearrowright in Brasiliâ circa Bahiam legit cl. Blanchet (n. 2097!). Affinis *M. hirsutissimæ* et forsân cum eâ necnon cum *M. vismiazefoliâ*, eriocladâ et clavellatâ sectionunculam naturalem format ex ramis hispidis distinctam. Fol. inf. 8 poll. longa, 5-6 lata. (v. s.)
96. *M. MENISPERMICA*, volubilis tota confertè breviterque hirsuta, foliis petiolatis cordatis acuminatis in parte latiore repando-dentatis utrinquè inter hirsutiem parçè glandulosis, ramis axill. terminalibusque apice corymbosis, capitulis pedicellatis, invol. squamis 4 acuminatis dorso puberis, bracteolâ lineari-subulatâ, invol. vix breviorè, achenio glanduloso. \curvearrowright in Mexico circa Bejar mense Augusto flor. legit cl. Berlandier. Capitula 2 $\frac{1}{2}$ lin. longa. Pappus subrufescens. (v. s. comm. à cl. inv.)
97. *M. POEPPIGII* (Spreng. syst. 3. p. 422), volubilis, caule subangulato gla-

brinuculo, foliis petiolatis cordatis subhastatis acuminatis repando-crenatis glabriusculis, auriculis subacutis parce divergentibus, pedunculis axillaribus oppositis folio duplō longioribus apice corymbiferis, capitulis confertis brevissimè pedicellatis, bracteolâ minimâ, invol. squamis ovali-oblongis minutissimè mucronatis, achenio glanduloso. \curvearrowright in insulâ Cubâ. Mikania denticulata Poepp. ex Spreng. non Willd. Mikania n. 59. Poepp! pl. exs. Capitula vix a lin. longa. Affinis M. menispermeæ. (v. s.)

98. *M. mollis* (H. B. et Kunth nov. gen. am. 4. p. 135), fruticosa volubilis, ramis sexangularibus canescenti-tomentosis, foliis petiolatis subrotundo-ovatis profundè cordatis subacuminatis subintegerrimis reticulato-5-nerviis supra pubescenti-hirtis subtus molliter cano-tomentosis, corymbis paniculatis, capitulis sessilibus solitariis, bracteolâ acuminatâ invol. brevior, invol. squamis lineari-oblongis obtusis extus pubescentibus, achenio glabro. \curvearrowright in temperatis Peruviz ad ripas flum. Guancabambæ. Petioli sesquipollic. Limb. fol. 2 $\frac{1}{2}$ poll. longus, 2 poll. latus.

99. *M. suaveolens* (H. B. et Kunth nov. gen. am. 4. p. 135), fruticosa volubilis, ramis sexangularibus foliisque molliter hispido-pubescentibus petiolatis subrotundo-ovatis profundè cordatis acuminatis grossè dentatis reticulato-5-nerviis, corymbis axillaribus oppositis subpaniculatis, capitulis subpedicellatis subternis, bracteolâ acuminatâ inv. brevior, invol. squamis lineari-oblongis acutis pedicellis pubescentibus. \curvearrowright in umbris humidis prope Guarumo Novo-Granatensium. Petiol. 1 $\frac{1}{2}$ 2 poll. longus. Limb. fol. 4 poll. longus, 3 poll. latus. Affinis M. scandenti.

100. *M. lanuginosa*, scandens tota ferè hirsutie molli confertâ albâ lanuginosâ, foliis petiolatis cordatis acuminatis except. sinu et acumine dentatis suprâ villosis viridibus subtus lanuginosis canis, paniculis ramos terminantibus oppositè ramosis, capitulis secus ramulos breviter racemosis pedicellatis, bracteolâ ovatâ acutâ glabriusculâ paulò infrâ invol., squamis invol. lanceolatis acutis glabris. \curvearrowright in Brasiliæ prov. Sancti-Pauli. Petioli 9 lin. longi. Limb. fol. 2 $\frac{1}{2}$ poll. longus, 2 poll. latus. Invol. rufescentia. Pappus albus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 425 miss.)

101. *M. rubiginosa* (Smith in Rees cycl. v. 23. n. 12), volubilis?, ramis velutinis, foliis petiolatis cordatis acutis undatis tenuiter velutinis utrinquè globulos resinosis gerentibus, paniculis axillaribus cymosis bracteis spatulatis, invol. squamis obovatis obtusis mucronatisve extus velutinis. \curvearrowright in Amer. merid.? *Cacalia cordifolia* var. Linn. f. suppl. 352. Numerus florum dubius. †.

§. 6. *Angulatæ, nempe volubiles exstipulatæ corymboso-paniculatæ, foliis basi truncatis seu cordatis angulato-dentatis lobatisve.*

102. *M. chenopodiifolia* (Willd. sp. 3. p. 1745 non Wall.), tota glabra, caule herbaceo? scandente tereti striato, ramulis ultimis angulatis, foliis petiolatis triangulari-ovatis acuminatis repando-dentatis 5-nerviis, corymbo paniculato trichotomo laxo, capitulis pedicellatis 4-floris, invol. squamis circ. 8 acuminatis, inter. longioribus. \curvearrowright in Sierra Leona (Willd.), in arenosis siccis circa Albreda Senegambiæ (Perrottet!). Affinis volubili. In specim. obvis capitula ferè omnia abortiva, squamis elongatis crispis. In paucissimis vidi achenia teretiuscula, pappum albidum. (v. s. comm. à cl. Perrottet!)

103. *M. brachypoda*, fruticulosa scandens, ramis teretiusculis apice subangulatis glabris, foliis brevissimè petiolatis acutè triangularibus aut rariùs subrhombeis dentatis 7-nerviis glabris suprâ minutissimè scabridis, corymbo terminali fastigiato laxo, capitulis pedicellatis, bracteolâ lanceolatâ acutâ, invol. squamis lanceolatis mucrone acutis dorso hinc indè subpuberis, achenio glanduloso. \curvearrowright in Brasiliæ prov. Sancti-Pauli. Rami tennes rigidi. Petiol. lineam longi. Limb. fol. 8-9 lin. longus latusque. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 406 miss.)

104. *M. diversifolia*, volubilis glabra, ramis teretibus subtiliter striatis, foliis

- petiolatis membranaceis, caulibus (seu ramor. præcip.) hastato-triangularibus lobis lateralibus divaricatis terminalique acuminatis integerrimis, ramis ovatis acuminatis basi truncatis, paniculâ ramis oppositis parce villosis, capitulis ad apices ramulorum congestis brevè pedicellatis, bracteolâ lineari-subulatâ parvâ, invol. squamis oblongo-linearibus obtusis glabris apice summo puberulis. (in ætibus circa Rio de Janeiro sat rara obs. cl. Lund fl. junio. Invol. straminea. Petiol. fol. caul. 8-9 lin. Fol. 3 poll. longa et lata; petiol. fol. ram. 6 lin. longus, limbus 20-21 lin. longus, 7-8 latus. (v. s. comm. à cl. Lund.)
105. *M. BIFORMIS*, volubilis, caule tereti vix striatulo glabro, ramis sparsè et ad apicem confertè pilosiusculis, foliis sparsè puberis petiolatis trinerviis, caulibus (aut ram. præcip.) subhastato-trilobis lobis et apice acuminatis, lateralibus multò brevioribus, paniculâ terminalis laxæ ramis villosiusculis, capitulis pedicellatis, bracteolâ lineari-subulatâ parvâ, invol. squamis linearibus obtusis achæniisque glabris. (circa Bahiam Brasiliæ legit cl. Blanchet. Petioli villosuli fol. caulin. 15 lin. longi, limbus 4 poll. longus, 3 latus; fol. ram. petiol. 5 lin., limbus 2 poll. longus, pollicem latus. Anthere exsertæ; faux cor. campanulata. (v. s. comm. à cl. inv.)
106. *M. ANGULARIS* (H. et Bonpl. pl. equin. 2. p. 87. t. 106), herbacea volubilis glabra, ramis teretibus sulcato-striatis, foliis petiolatis triangulari-bastatis acutis remotè et obsolete denticulatis reticulato-trinerviis, corymbis paniculatis oppositis, capitulis subsolitariis pedicellatis, bracteolâ lineari acutâ invol. dimidio breviorè, invol. squamis lanceolatis acutis. $\frac{1}{2}$ (locis humidis in ripâ fluit. Catamago Novæ-Granatæ H. B. et Kunth nov. gen. am. 4. p. 136. Petiol. bipollis. Folia 7 poll. longa. Species affinis *M. Loxensis* dicitur.
107. *M. SCANSORIA*, volubilis glabra, ramis teretibus substriatis, foliis petiolatis trilobis basi truncatis aut subcordatis 3-5-nerviis, lobis lateralibus brevibus terminali acuminato, marginibus integerrimis, paniculæ ramosissimæ ample ramis ramulisque oppositis velutinis, capitulis ad apices ramulorum densè glomeratis subsessilibus, bracteolâ et invol. squamis æqualibus oblongo-linearibus subacutis glabriusculis. (in Brasiliæ sylvis primævis super imas arbores scandens obs. cl. Lhostky. Petiol. 8-9 lin. longi, fol. limb. 2 $\frac{1}{2}$ poll. longus, 2 poll. latus. (v. s. comm. à cl. Lhostky.)
108. *M. HEDERÆFOLIA*, fruticosa volubilis glaberrima, caule striato-sulcato, foliis petiolatis cordatis hastato-trilobis, lobis acuminatis, nervis è nervii basi utrinquè 2-3 approximatis, paniculæ terminalis ramis ramulisque oppositis, capitulis ad apices ramulorum congestis subsessilibus, bracteolâ lineari-subulatâ parvâ, invol. squamis linearibus subacutis achæniisque glabris. (in Brasiliâ circa Rio de Janeiro legit cl. Lund, et circa Bahiam cl. Blanchet (n. 2111!), Caulis crassitie digiti. Folia summa ferè triangularia; infer. petioli ferè bipollicares, limb. 3 $\frac{1}{2}$ poll. diam. (v. s. comm. à cl. Lund.)
109. *M. VITIFOLIA*, scandens? herbacea, caulibus fistulosis ramisque teretibus vix striatis sparsè petiolisque pube articulatâ hispidulis, foliis longè petiolatis cordatis 7-nerviis orbiculatis 7-angulatis lobis brevibus acuminatis dentibus callosis exsertis remotis, utrinquè secus nervos hirsutis cæterum sparsè scabridis, paniculis amplis oppositè ramosis, ramis ipsis paniculatis, capitulis pedicellatis, bracteolâ lineari acutâ, invol. squamis linearibus obtusis achæniisque glabris. — in Brasiliæ prov. Sancti-Pauli. Folia Vitis sed paulò majora, ferè potius Sparmanniæ. Paniculæ pedales sesquipedalesve. Petiol. 4 poll. Limb. 4-5 poll. diam. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 426 miss.)
- §. 7. Partitæ, nempe volubiles exstipulatæ corymboso-paniculatæ, foliis in segmenta seu partitiones profundè partitis.
110. *M. APIIFOLIA*, suffruticosa scandens glabriuscula, caule tereti glabro, ramis subangulatis apice tenuissimè rufo-velutinis, foliis petiolatis trisectis, segmentis petiolulatis ovatis trifidis dentatis acutis glabris, pedunculis axillaribus terminalibusque apice laxè corymbosis, bracteolâ lineari acutâ foliaceâ à basi

invol. subdistante, invol. squamis linearibus elongatis acutis achæniisque glabris. 5 in Brasiliæ prov. Sancti-Pauli. Species distinctissima. Pappus rufus. Capit. 5 lin. long. Cor. ex sicco purpurascentes. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 421 miss.)

- III. *M. DENTATA* (Spreng. syst. 3. p. 422), herbacea scandens, ramis angulatis, foliis compositis simplicibusque ovatis lanceolatisque angulatis argutè dentatis glabris, paniculis axillaribus diffusis, capitulis solitariis. 7 in Brasiliâ. Cæt. ign. An. affinis *M. apiifoliæ*. †

† *Species dubiæ.*

Obs. 1. Ad Mikaniæ genus referendæ videntur stirpes sequentes in vol. 8^{vo} floræ fluminensis iconibus nec descriptionibus incompletè illustratæ nempe: *Cacalia cordata* t. 53 (affinis *M. micropteræ*), *Cac. trilobata* t. 54, *Cacal. angulata* t. 55, *Cac. ternata* t. 56, *Cac. obsoleta* t. 57, *Cac. villosa* t. 58?, *Cac. nitida* t. 59, *Cac. reticulata* t. 60, *Cac. pilosa* t. 61, *Cac. triangularis* t. 62, *Cac. septemata* t. 63?

Obs. 2. Specimen incompletum ex Sierra-Leona ortum, *M. tamoidi* affine vidi in. h. Lindl. Species sola hucusque in Africâ æquinoctiali nota, verisim. nova.

†† *Species exclusæ.*

- M. ARBorea* H. B. K. = *Eupatorium dendroides*.
M. ARBorea Roxb. = *Lachanodes prenanthiflora*.
M. ARTEMISIODES Cass. = *Eupatorium leptophyllum* ß.
M. AURICULATA Willd. = *Senecio deltoideus*.
M. CHENOPODIFOLIA Wall. = *Senecio chenopodifolius*.
M. MELISSÆFOLIA Willd. = *Nothites melissæfolia*.
M. ASPARAGOIDES Licht. = *Euryops*.
M. SATUREIÆFOLIA Willd. = *Nothites satureiæfolia*.
M. SERRATIFOLIA H. B. K. = *Eupatorium serratifolium*.
M. TOMENTOSA Willd. = *Senecio penicillatus*.

XC. *ADENOSTYLES* Cass. dict. 1. suppl. p. 59. Less. syn. 157 non Blum. — *Cacalia Tourn. inst. p. 451. t. 258. Vaill. Adans. Hall. Gærtn. Gaudin. non Cass. nec Less.* — *Cacaliæ* sp. Linn. Willd.

Capitulum pauciflorum discoideum. Recept. angustum nudum. Invol. cylindraceum 1-seriale oligophyllum. Cor. tubulosæ, limbo campanulato 5-dentato. Styli rami longè exserti semicylindrici undiquè papilloso-glandulosi. Achænium teretiusculum striatum. Pappus pluriserialis setis piliformibus scabris. — Herbæ Europæ montanæ perennes. Folia alterna petiolata sæpè basi auriculata reniformia. Capitula in corymbum compositum nudum disposita pedicellata. Flores purpurascentes aut pallidi imò albidii. — Evidenter ex veterum nomenclaturâ hoc genus *Cacaliæ* nomen servare debuisset, sed Cassinianam designationem admisi, ne confusionem jam maximam augeam!

- I. *A. GLABRA*, foliis cordatis dentatis utrinquè glabris, corymbis fastigiatis, capitulis 3-6-floris. 7 in montanis et subalpinis (nec in alpinis) petrosis irriguis Europæ, non rara in Galliâ, Germaniâ, Italiâ, etc., rara in Siciliâ (Guss.). *Cacalia alpina* var. ß Linn. sp. 1170. *Cacalia glabra* Vill. dauph. 3. p. 170. *Cacalia alliarifolia* Lam. dict. 1. p. 532. *Cacalia alpina* Jacq. fl. austr. t. 234. Willd. sp. 3. p. 1735. Sturm. fl. germ. icon. Tussilago *Cacalia* Scop. carn. n. 1055. *Aden. viridis* Cass. l. c. *Aden. alpina* Bluff et Fing. comp. fl. germ. 2. p. 329. *Senecio glaber* Clairv. man. val. 240. Lob. ic. t. 592. f. 1. Petioli, etiam sub-amplexicaules, non basi appendiculati. (v. v.)

2. *A. PETASITES* (Bluff et Fing. comp. fl. germ. 2. p. 329), foliis cordatis dentatis subtus incanis superne glabrescentibus, corymbis fastigiatis, involucri glabris 3-6-floris. $\frac{1}{2}$ in pascuis et sylvaticis montium Europæ. *Cacalia alpina* var. α Linn. sp. 1170 (cum seq. mixta ex syn.). *Cacalia albifrons* Linn. f. suppl. 353. *Cacalia Petasites* Lam. dict. 1. p. 531. DC. fl. fr. n. 3104. *Cacalia alliarie* Gou. illustr. 65. *Cacalia hirsuta* Vill. dauph. p. 152. *Cac. tomentosa* Jacq. fl. austr. t. 235. *Adenostyles albida* Cass. l. c. *Adenostyles albifrons* Reich. fl. exc. 2. p. 278. *Senecio hirsutus* Clairv. man. val. 241. — Clus. hist. 2. p. 115. f. 1. Petioli sæpius basi utrinque auriculâ appendiculati. (v. v.)
- β . *nuda* (DC. fl. fr. n. 3104 β), petiolis basi exappendiculatis. — Rarior. *Cac. albifrons* β Gaud. fl. helv. 5. p. 216. Invol. 3-6-flora ut in var. α nec 15-20 ut in sequenti. (v. s.)
3. *A. LEUCOPHYLLA* (Reich. fl. exc. 2. p. 278), foliis cordatis utrinque albotomentosis dentatis corymbo subrotundo coarctato, involucri dorso lanatis 15-20-floris. $\frac{1}{2}$ in summorum Alpium petrosis, in Delphinatu, Pedemontio, Sabaudia, Valesia. *Cacalia leucophylla* Willd. sp. 3. p. 1736. DC. fl. fr. n. 3105. *Cac. tomentosa* Vill. dauph. 3. p. 171. non Thunb. nec Jacq. *Adenost. candidissima* Cass. l. c. *Senecio tomentosus* Clairv. man. val. 240. non Salisb. *Cacalia alpina* var. *tomentosa* All. ped. n. 644. Petioli basi nudi aut vix subauriculati. (v. s.)
4. *A. HYBRIDA*, foliis cordatis dentatis supra glabrescentibus subtus araneosis, petiolis basi sæpius auriculatis, corymbo laxiusculo, involucri squamis dorso puberis 12-18-floris. $\frac{1}{2}$ in Alpibus Helveticis (Gaud.), Delphinensibus (Vill.) et in Sicilia montibus Nebrodensibus (Duby!). *Cacalia tomentosa* β hybrida Vill. dauph. 3. p. 171. DC. fl. fr. n. 3105. Gaud. fl. helv. 5. p. 216. Omnino media inter *Petasitem* et *leucophyllam*, priori foliis, posteriori florum numero affinis! fortè utriusque hybrida progenies? Hanc inter species donec lux fiat admisi quia omnino anxius hæreo an *Petasitis* an *leucophyllæ* sit varietas, et quia reperta fuit in insulâ ubi neutra aliarum adhuc cognita! (v. s.)

Subtrib. II. *TUSSILAGINÆ* Less. syn. 158.

Capitula heterogama aut subdioica.

Div. I. *PETASITÆ*.

Capitula heterogama aut subdioica, corollis fœmineis etiam tubulosis. — Flores (nunquam flavi) albi aut purpurascens.

XCI. *HOMOZYNE* Cass. dict. 21. p. 412. Less. syn. 158. — *Tussilago* sect. 2^a. DC. fl. fr. — *Tussilaginis* sp. Linn.

Capitulum multiflorum subheterogamum, fl. radii 1-serialibus paucis filiformibus fœmineis, disci tubulosi 5-fidis hermaphroditis. Recept. nudum. Invol. cylindracei squamæ subuniseriales oblongo-lineares acutæ disco paulò breviores. Styli fl. fœm. et herm. conformes! Achæmium oblongo-cylindraceum sulcatum glabrum. Pappus pilosus multiserialis. — Herba Europæ montanæ perennes. Folia pleraque radicalia cum flore circ. cœtanea petiolata cordato-orbiculata dentata aut angulata, 1-2 ad basin scapi. Scapus simplex 1-cephalus imâ basi 1-2-foliosus apice parcè squamosus. Flores purpurascens et variantes albi.

1. *H. SYLVESTRIS* (Cass. l. c.), foliis subglabris reniformibus acutè inciso-7-lobis, lobis intermediis 3-dentatis. $\frac{1}{2}$ in sylvaticis sabbatinis Carinthiæ et Carniolæ. *Tussilago sylvestris* Scop. carn. n. 1057. Jacq. austr. 5. app. 4. 12. Willd. sp. 3. p. 1967. Scapus circ. pedalis rarissimè 2-3-cephalus (Bluff et Fing.)

cum petiolis et foliorum marginibus parèe subhirsutus. Fol. caulis petioli basi vaginantes. (v. s.)

2. *H. alpina* (Cass. l. c.), foliis reniformibus dentato-crenatis subtus ad venas pilosis. \mathcal{L} in pratis elatis et humidis inter terminum Fagi et lineam nivalem Alpium, Jurassi, montis Generosi (Com.). *Tussilago alpina* Linn. sp. 1312. Jacq. austr. t. 246. Lam. ill. t. 674. f. 7. Schkuhr handb. t. 242. Gaud. fl. helv. 5. p. 272. Baumg. enum. 3. p. 92. *Petasites alpinus* Scop. carn. Herba circ. semipedalis. Folia scapi a vaginantiâ, limbo subnullo. (v. s.)
3. *H. nasorum* (Cass. l. c.), foliis cordato-orbiculatis undulato-dentatis supra glabris subtus tomentosis. \mathcal{L} in alpidibus Austriæ (Jacq.), Tyrolis (Hop!) Saalsburgi, Carinthiæ, Carinoliæ super. (Graaf!), etc. *Tussilago discolor* Jacq. austr. t. 247. Sturm deutschl. fl. ic. *Petasites discolor* Baumg. enum. 3. p. 92. *Tussilago alpina* Scop. carn. — Clus. hist. 2. p. 112 fig. inf. et 113 fig. int. Herba vix semipedalis, fortè *H. alpina* var? (v. s.)

XCII. *NARDOSMIA* Cass. dict. 34. p. 186. Less. syn. 139. — *Tussilaginis* sp. auct.

Capitula multiflora subdioica — Submascula, radio uniseriâli foemineo ligulato, disco bisexuali et abortu masculo sterili, cor. tubulosis 5-dentatis. — Subfoeminea radio pluriseriâli foemineo minutissimè ligulato, disco pauci (1-5) floro, cor. tubulosis 5-dentatis. Recept. nudum planum. Invol. squamæ 1-seriales floribus æquales aut breviores. Achenia teretiuscula glabra. Pappus pilosus, fl. masculorum brevior et minus copiosior quàm foemineorum. — Herba Europææ, Asiaticæ aut Boreali-Americanae, perennes. Folia cum floribus aut paulò post flores nascentia petiolata dentata basi cordata. Scapus squamosus apice thyrsus fastigiatum gerens. Flores purpurascens aut albi.

1. *N. fragrans* (Reichenb. fl. exc. 2. p. 289), foliis coarctatis basi cordatis orbiculatis cartilagineo-dentatis supra glabrisculis subtus in nervis puberis, lobis baseos subrotundis, invol. squamis acutis discum æquantibus aut superantibus. \mathcal{L} ad Neapolim (Reyn.), in Sicilia (Guss.), Sardinia (Moris!) et Mauritania (Poir.) in vallibus subhumidis et ad rivulos. *Nard. denticulata* Cass. dict. 34. p. 187. *Tussilago suaveolens* Desf. cat. Cacialia alliariaefolia Poir. t. num. 2. p. 236. non Linn. ex Poir. dict. 8. p. 151. *Tussilago racemosa* Tausch. syll. soc. ratib. 2. p. 251? ex Ten. syll. 436. Flores hybèrni suaveolentes. Herba ex eo culta sub nom. vernaculo *Héliotrope d'hiver*.

* *Submascula*, ligulis ovali-oblongis nunc invol. longioribus, nunc rariùs invol. ferè brevioribus. — *Tussilago fragrans* Vill. act. soc. h. nat. 1. p. 72. t. 12. (v. v. et s.)

** *Subfoeminea*... nondum cognita.

2. *N. lævigata*, foliis coarctatis basi cordatis orbiculatis cartilagineo-dentatis supra glabrisculis subtus in nervis puberulis, lobis baseos subrotundis, invol. squamis obtusis disco brevioribus. \mathcal{L} in Sibiria et fortè in Bohemia.

* *Submascula*, ligulis ovali-oblongis involuero paulò longioribus. *Tussilago* Gmel. fl. sib. 2. t. 69? et ideò *Tussil. lævigata* Willd. sp. 3. p. 1969. Ledeb. fl. alt. 47. p. 54. *Tussil. Bohemica* Hoppe. tasch. 1803. p. 60. *Nardosmia straminea* Cass. dict. 34. p. 189. *Petasites lævigatus* Reich. fl. exc. 2. p. 279 huc essent revocand. etiamsi flores flavescentes (forsan ex sicco) dicti! Plantam non vidi.

** *Subfoeminea*, lig. fl. foem. minutissimis. — *Tussilago saxatilis* Turcz. l. in litt. Flores ex eo sordidè rubri. Folia ferè omninò *N. fragrantis*. An discrimen ex invol. petiti ad speciem aut ad sexum pertinet? (v. s. coll. à cl. Turczaninoff in subalpinis montium Kultokensium et in alpe Chamor.)

3. *N. frigida* (Hook. fl. bor. am. 1. p. 307. var. α), foliis cordatis inaequalitè grossè et obtusè dentatis sublobatis supra glabris subtus albo-tomentosis, lobis

baseos divergentibus. \mathcal{U} in vallibus et montanis alpinis Dahuriæ (Turcz.!), Sibiriae (Gmel.!), Norwegiæ (Chr. Smith!), Lapponiæ (Linn.), Americæ maximè borealis (Pursh). — *Nardosmia angulosa* Cass. dict. 34. p. 188.

* *Submascula*, ligulis ovali-oblongis stylo longioribus. — Tussil. frigida OEd. fl. dan. t. 61. — Gmel. sib. 2. p. 150. t. 70. (v. s.)

** *Subfeminea*, ligulis minutissimis, caule et thyrsos duplò longioribus. (v. s. coll. ad salinas Ircutianas à cl. Turczaninoff.)

4. *N. corymbosa* (Hook. fl. bor. am. x. p. 307), foliis cordatis sinuatis et acutè denticulatis oblongis acutis suprà glabris subtùs tomentosis, lobis baseos divergentibus. \mathcal{U} in insulâ Melville (Br.) ad flum. Mackensie (Hook.). Tussilago corymbosa Brown chlor. melv. p. 21. Affinis *N. frigida*.

* *Submascula*. — adhuc ignota.

** *Subfeminea*, ligulis linearibus minutissimis. — Sola hucusquè descr. Corymbus pauciflorus. Invol. squamæ acutæ floribus duplò breviores. (v. s. comm. à cl. Sabine.)

5. *N. sagittata* (Hook. fl. bor. am. i. p. 307), foliis oblongis acutis sagittatis integerrimis, lobis obtusis. \mathcal{U} ad sinum Hudsonis (h. Banks), ad rupes scopulosas, à lacu superiore ad arcidulam Franklin (Hook.). Tussilago sagittata Pursh fl. bor. am. 2. p. 531. Thyrsus dicitur ovatus fastigiatus capitulis radiatis. Videtur specim. submasculum.

6. *N. palmata* (Hook. fl. bor. am. i. p. 308), foliis subtùs tomentosis palmatipartitis, lobis oblongis grossè pauci-dentatis. \mathcal{U} in Americâ bor. ad Labrador et Terram-Novam (Ait.) à lacu Huron ad lacum Bear, in rupibus scopulosis dictis, et ad ostium flum. Columbia (Hook.). Tussilago palmata Ait. kew. 3. p. 188. t. 11. Specim. submasculum. Thyrsus fastigiatus 7-9-cephalus. Ligulæ oblongæ stylo longiores.

XCIII. PETASITES *Tourn. inst. 451. t. 258. Gærtn. fr. 2. p. 406. t. 166. Cass. dict. 39. p. 199.* — Tussilago sect. Petasites DC. fl. fr. Gaud. fl. helv. Tussilaginis sp. Linn.

Capitula multiflora subdioica. — Submascula nempe radio 1-seriali pauci- (1-5-) floro femineo, corol. tubulosâ gracili, ovario ovulato, disco bisexuali sed ovario exovulato masculo, cor. tubulosâ fauce subdilatata regulariter 5-dentata. — Subfeminea nempe radio multiseriali femineo fertili, cor. filiformi subtruncatâ, stylo exserto, disco pauci- (1-5-) floro masculo, cor. fauce dilatata regul. 5-dentata. Recept. nudum planum. Invol. squamæ subuniserialis floribus breviores bracteolis sæpè basi stipatæ. Achænia cylindræa glabra. Pappus pilosus, marium multò minùs copiosior quàm femineorum. — Herbæ Europæe perennes in humidis crescentes. Folia post flores orta, radicalia petiolata reniformia aut cordata dentata ampla. Scapus sæpiùs tomentosus squamis membranaceis glabriusculis onustus. Thyrsus polycephalus terminalis. Pedicelli simplices 1-cephali aut in subfemineis interdum ramosi. Flores purpurascens aut albid.

1. *P. vulgaris* (Desf. fl. atl. 2. p. 270), foliis profundè cordatis suborbiculatis inæqualiter dentatis suprà demum glabris subtùs pubescentibus, lobis baseos approximatis. — \mathcal{U} in Europæ humidis frequens, etiâ Algeriæ (Desf.). Tussilago Petasites Hopp! tasch. 1803. p. 35. Willd. sp. 3. p. 1971. DC. fl. fr. n. 3165. Petasites officinalis Moench.

* *Submasculus*, thyrsi ovati pedicellis simplicibus. Tussilago Petasites Linn. sp. 1215. Tuss. Petasites hermaphrodita Poir. Gaud. — Lam. ill. t. 674. f. 1. 2. Bull. herb. t. 391. fl. dan. t. 842. (v. v.)

** *Subfemineus*, thyrsi elongati pedicellis plerisque ramosis. Tussilago hy-

brida Linn. sp. 1214. Sow. engl. bot. t. 430. Schkuhr handb. t. 242. Dill. elth. t. 230. (v. v.)

2. *P. ALBUS* (Gærtn. fr. 2. p. 406. t. 166), foliis latè cordatis suborbiculatis angulosis argutè dentatis suprà demùm glabratis subtùs pubescenti-villosis albidis, nervis scabris, lobis baseos subrotundis. \mathcal{L} in montosis Europæ non rarus, etiam Algeriæ (Desf.). *T. alba* Linn. sp. 1214. DC. fl. fr. n. 3166. An hùc aut ad priorem refer. *Petasites glabrescens* Hoppe?

* *Submasculus*, thyrsi fastigiati pedicellis simplicibus. — Fl. dan. t. 524. Hoppe! in Sturm deutschl. fl. ic. (v. v.)

** *Subfemineus*, thyrsi oblongi pedicellis plerisque ramosis. Tuss. ramosa Hoppe! cent. Tuss. alba β Vill. act. soc. h. nat. par. 1. p. 73. (v. v.)

3. *P. NIVUS* (Cass. dict. 39. p. 202), foliis latè cordatis subtriangularibus dentatis suprà glabriusculis subtùs niveo-tomentosis, nervis concoloribus, lobis divaricatis subacutis. \mathcal{L} in montanis Alpium et Jurasi priore rarior; etiam in Russia merid. ex specim. masculo sub nom. *T. hybrida*? ab hon. Ac. Sc. Petrop. misso. Tussilago nivea Vill. act. soc. h. nat. Par. 1. p. 73. Willd. sp. 3. p. 1970. DC. fl. fr. n. 3167. Tussilago frigida Vill. dauph. 3. p. 175. Sut. fl. helv. 2. p. 180 non Linn. Pedicelli utriusque sexus simplices. Lobi baseos fol. sæpè bifidi.

* *Submasculus*, thyrsi oblongi pedicellis brevioribus. — Hoppe! in Sturm deutschl. fl. ic. — Moris. ox. s. 7. t. 10. f. 4. (v. v.)

** *Subfemineus*, thyrsi oblongi pedicellis elongatis. — Tussilago paradoxa Retz obs. 2. p. 24. t. 3. Hoppe! in Sturm fl. ic. Tussil. frigida Sut. fl. helv. 2. p. 180 non Linn. (v. v.)

4. *P. TOMENTOSUS*, foliis profundè cordatis ovatis orbiculatisve vix dentatis suprà glabriusculis subtùs densè albo-tomentosis, nervis concoloribus, lobis baseos approximatis sublobatis subrotundis. \mathcal{L} in Germania et Scania ad ripas flum. rarior. Tussilago spuria Willd. sp. 3. p. 1972? Tuss. tomentosa Ehr. beitr. 3. p. 65. Hoppe! tasch. 1803. p. 51. *Petasites spurius* Reichenb. fl. exc. 2. p. 279. Pedicelli utriusque sexus simplices.

* *Submasculus*, thyrsi coarctato, pedicellis brevioribus.

** *Subfemineus*, thyrsi elongato. — Tuss. spuria. Retz obs. 1. p. 29. t. 2? (v. s.)

XCIV. *ADENOCAULON* Hook. bot. misc. 1. p. 19. t. 15. Less. in linnæa 1831. p. 107. syn. 159.

Capitulum 9-10-florum heterogamum, floribus tubulosis 4-5-dentatis, radii quinque 1-serialibus fœmineis, disci 4-5 masculis. Invol. uniseriale demùm reflexum pleiophyllum. Recept. nudum. Achænia oblongo-obovata glandulis stipatis supernè instructa. Pappus nullus. — Herbæ in Americâ occidentali habitantes perennes. Caules albo-tomentosi supernè glandulis pedicellatis instructi, infernè foliosi, supernè nudi laxè paniculati. Folia alterna penninervia suprà glabra subtùs tomentosa penninervia petiolata.

1. *A. BICOLOR* (Hook. bot. misc. 1. p. 19. t. 15. fl. bor. am. 1. p. 308), foliis subtrilobis angulato-dentatis. \mathcal{L} in sylvis densis ad oram occidentalem Americæ borealis prope fretum de Fuca, Fort Vancouver et in Rocky-mountains.
2. *A. CHILENSE* (Less. in linnæa 1831. p. 107), foliis subrotundo-ellipticis cordatis integerrimis aut subintegerrimis. \mathcal{L} in Chili.

Div. II. *EUTUSSILAGINÆÆ*.

Capitula heterogama radiata, ligulis fœmineis, floribus disci hermaphroditis. — Flores semper lutei, radii etiam lutei aut rarissimè rosei.

XCV. TUSSILAGO *Tourn. inst.* 487. t. 276. *Adans. fam.* 2. p. 124. *Gærn. fr.* 2. t. 170. *Cass. dict.* 34. p. 190 (*excol. spec.*). *Less. syn.* 159. — *Tussilago* sect. *Farfara* DC. *fl. fr.* — *Tussilaginis* sp. *Linn.*

Capitulum multiflorum heterogamum, fl. radii pluriserialibus angustissime ligulatis foemineis, disci paucis masculis tubulosis limbo campanulato 5-dentato. Recept. nudum. Invol. subuniseriale, squamis oblongis obtusis. Antheræ subcaudatæ. Stylus disci inclusus abortivus, radii bifidus, ramis teretiusculis. Achænium radii oblongo-cylindraceum glabrum, disci abortivum. Pappus radii pluriserialis, disci uniserialis, setis tenuissimis. — Herba perennis Gerontogea. Rhizoma crassiusculum. Folia post anthesin nascentia petiolata cordata angulata. Scapi 1-cephali tomentosi aphylli sed squamis alternis oblongis adpressis inv. squamas referentibus onusti. Flores flavi.

1. T. FARFARA (*Linn. sp.* 1224). 2½ in argillosis petrosis humidis Europæ totius, Byzantii (d'Urv.), Tauriæ (Stev.), Persiæ (Belang.) et in montibus Sibirici (Patr.), Indiæ orient. (Royle!). *OEd. fl. dan.* t. 595. *Sow. engl. bot.* 6. t. 429. *Sturm fl. germ.* ic. T. vulgaris Lam. *fl. fr.* 2. p. 71. T. rupestris Vall. cat. n. 2990. comp. n. 100. — Legi ad littora maris et in alpebus ad altit. hexapod. 1300! Variat scapo digitali et pedali, solitario aut multiplici, foliis fere cum floribus aut post eos natis, membranaceis aut rigidulis, utrinque subglabris aut subtus araneosis denseve tomentosis, basi cordatis aut ovatis, angulatis aut subdivisis, plus minus dentatis. Has variationes video et in specim. Europæis et in Asiaticis. (v. v. et s.)

N. B. Quid *Tussilago bifolia* Levy veg. orig. p. 13?

† *Species exclusæ.*

- T. ALBA L. = *Petasites albus*.
- T. ALBICANS Sw. = *Leria albicans*.
- T. ALPINA L. = *Homogyne alpina*.
- T. ALPINA Scop. = *Homogyne discolor*.
- T. ANANDRIA L. = *Anandria bellidiastrum*.
- T. ANANDRIA Willd. = *Anandria bellidiastrum* β.
- T. BOHEMICA Hopp. = *Nardosmia saxatilis* α.
- T. CACALIA Scop. = *Adenostyles glabra*.
- T. DENTATA L. = *Chaptalia dentata*.
- T. DISCOLOR Jacq. = *Homogyne discolor*.
- T. EXCAPA Pers. = *Loxodon brevipes*.
- T. FRAGRANS Vill. = *Nardosmia fragrans*.
- T. FRIGIDA L. = *Nardosmia frigida*.
- T. HYBRIDA L. = *Petasites vulgaris*. **
- T. JAPONICA L. = *Senecio Kämpferi*.
- T. INTEGRIFOLIA Michx. = *Chaptalia tomentosa*.
- T. LAEVIGATA Willd. = *Nardosmia saxatilis*.
- T. LYRATA Willd. = *Anandria bellidiastrum*.
- T. LYRATA Pers. = *Leria nutans*.
- T. NIVEA Hopp. = *Petasites niveus*.
- T. NUTANS L. = *Leria nutans*.
- T. ONORATA Vill. = *Adenostyles glabra*.

- T. *PALMATA* Ait. = *Nardosmia palmata*.
 T. *PARADOXA* Retz. = *Petasites niveus*. **
 T. *PARADOXA* Roth. = *Petasites tomentosus*.
 T. *PETASITES* L. = *Petasites vulgaris*.
 T. *PILOSELLOIDES* Pers. = *Liberkuhnia piloselloides*.
 T. *PUMILA* Sw. = *Leria pumila*.
 T. *RACEMOSA* Tausch. = ? *Nardosmia fragrans*.
 T. *RAMOSA* Hop. = *Petasites albus*. **
 T. *RUBRA* Link. = *Senecio cruentus*.
 T. *SAGITTATA* Pursh. = *Nardosmia sagittata*.
 T. *SARMENTOSA* Pet. Th. = *Chevreulia stolonifera*.
 T. *SAXATILIS* Turcz. = *Nardosmia saxatilis*. **
 T. *SINUATA* Pers. = *Leria pumila*.
 T. *SPURIA* Willd. = *Petasites tomentosus*.
 T. *SPURIA* Schrank. = *Petasites niveus*.
 T. *SUAVEOLENS* Desf. = *Nardosmia fragrans*.
 T. *SYLVESTRIS* Jacq. = *Homogyne sylvestris*.
 T. *TOMENTOSA* Hop. = *Petasites tomentosus*.
 T. *TRIFURCATA* Forst. = *Senecio trifurcatus*.

XCVI. *CELMISIA* Cass. dict. 37. p. 259 (non 7. p. 357). Gaudich. voy. 470.

Capitulum multiflorum, fl. radii ligulatis fœmineis 1-seriatis, disci tubulosi hermaphr. 5-dentatis. Recept. latum epaleaceum nudum aut alveolare. Invol. squamæ pluriserialis inæquales acutæ disco paulò longiores. Ligulæ stam. rudimentis destitutæ stylum longum ramis obtusis dorso subpapilloso gerentes. Antheræ ecaudatæ appendiculatæ. Achænium oblongo-cylindraceum basi attenuatum. Pappus pluriserialis, paleis filiformibus inæqualibus herbatis. — Herbæ Australasicæ, caule erecto simplici nudo 1-cephalo. Folia radicalia oblonga. Capitulum disco luteo, radio roseo? Contra regulas prioritatis strictas nomen huic nec sequenti generi servavi, quia fuit à cl. Gaudichaud sub hoc nomine sanctum.

1. *C. LONGIFOLIA* (Cass. dict. 37. p. 256), foliis sessilibus linearibus argenteo-lanatis longissimis. ½ in Novâ-Hollandiâ orientali ad vallem Jamieson (Gaud.!), in ericetis humidis et excelsis paludibus montium cœruleorum (Cunn.!). Gaudich. voy. Freyc. bot. p. 470. t. 91. Ovarium sessile glabrum. Flosc. ligulati staminibus omninò orbat. Invol. disco longius. Recept. alveolatum. (v. s.)
2. *C. SPATULATA* (A. Cunn. in litt.), foliis elliptico-oblongis in petiolum longè attenuatis utrinquè glabris. ½ in Novâ-Hollandiæ collibus siccis ad Oyater Harbour prope Regis Georgii sinum legit cl. A. Cunningham. Scapus basi glaber nudus, apice pubescens squamellas lineares paucas gerens, 12-14 lin. longus. Folia petiolo incluso 2-3 poll. longa, 5-6 lin. lata. Invol. basi extus lanatum, squamis acuminatis apice subinduratis. (v. s.)

XCVII. *ALCIOPE*. — *Celmisia* Cass. dict. v. 7. p. 357 (non vol. 37). — *Ligulariæ* sp. Less. — *Arnica* sp. Thunb.

Capitulum multiflorum, fl. radii 1-serialibus ligulatis fœmineis sed stam. filamenta subabortiva gerentibus, disci tubulosi 5-dentatis hermaphroditi. Recept. latum epaleaceum. Invol. squamæ pluriserialis subimbricatæ adpressæ. Styli rami breves obtusi parcè papilloso glabri,

in radio et disco sat conformes. Achænium oblongo-cylindraceum subobcompressum. Pappus pluriserialis, setis scabridis imâ basi breviter concretis. — Herbæ suffrutescentes Capenses erectæ ramosæ. Caules teretes tomento denso lanato canescentes. Folia alterna petiolata integra aut denticulata suprà glabra subtùs cum petiolis cano-lanata. Capitula ramos terminantia. Invol. cano-lanatum. Flores flavi. — Genus hinc ad Ligulariam ex ligulis stam. rudimentaria gerentibus, illinc ad Celmisiam ex structurâ stigmatis, accedens. Dixi ex Alciopè nymphâ mythologicâ Celmisii matre, alludens ad hoc quod genus primum Celmisia dictum, alterum quasi peperit.

1. A. TABULARIS, caule basi fruticoso erecto ramoso, ramis tomentosis elongatis apice 1-cephalis, foliis breviter petiolatis ovalibus obovatisve basi subattenuatis integris aut denticulatis suprà glabris subtùs lanato-tomentosis, invol. squamis oblongo-linearibus subaraneosis, exter. paucis interiores ferè æquantibus. 5 ad Cap. B.-Spei in monte Tabulari (Th. Drege! n. 173). Arnica Tabularis Thunb. fl. cap. 668. Celmisia rotundifolia Cass. dict. 7. p. 357. Ligularia Tabularis Less. syn. 390. Folia 1-1 $\frac{1}{2}$ poll. longa, 9-10 lin. lata. (v. s.)
2. A. LANATA, caule frutescente erecto subsimplici densissimè lanato, ramis paucis apice 1-cephalis bracteolis lanatis obsitis, foliis in parte caulis inferiore petiolatis cordato-ovatis obtusis margine subrevolutis coriaceis subtùs lanatis, adultis suprà glabris, invol. 2-3-seriali imbricato dense lanato. 5 ad Cap. Bonæ-Spei juxta Roode Sand in montibus prope Winterhoek (Th.), in distr. Worcester et Tulbagh (Eckl. n. 1421. 830. 1947). Arnica lanata Thunb. fl. cap. 667. Conyza tomentosa Burm. prod. 26. Ligularia lanata Less. syn. 390. (v. s.)

ß. grandis, caule basi frutescente erecto parcè ramoso densissimè lanato, ramis apice 1-cephalis bracteolis lanatis obsitis, foliis petiolatis ovatis (basi cordatis), margine planis obtusis submembranaceis suprà glabris subtùs dense lanatis, invol. 2-3-seriali imbricato dense lanato. 5 ad Cap. Bonæ-Spei (Th.), in monte prope Tulbagh (Dreg. n. 1975). Arnica grandis Thunb. fl. cap. 668. E. Mey. in Dreg. coll. (v. s.)

XCVIII. BRACHYGLOTTIS Forst. car. gen. n. 46. Cass. dict. 48. p. 464. — Senecionis sp. Less. — Cinerariæ sp. Willd.

Capitulum pluri- (9-10-) florum heterogamum, fl. radii 1-serialibus paucis fœmineis brevissimè ligulatis aut obliquè tubulosis disco ferè brevioribus, disci tubulosis 5-dentatis herinaphr. Recept. nudum. Invol. oblongo-cylindricum squamis linearibus 1-serialibus extùs calyculato-bracteolatum. Styli radii rami exserti revoluti apice clavellati obtusi; disci stylus inclusus subindivisus fortè abortivus. Achænia oblonga basi non attenuata. Pappus 1-serialis, setis densis scabris basi in anulum concretis. — Arbores Australasicæ. Folia alterna ovalia aut subrotunda petiolata subtùs tomentosa. Corymbus paniculatus.

1. B. REPANDA (Forst. l. c.), foliis ovatis repando-sinuatis, paniculâ decompositâ divaricatâ. 5 in Novâ-Zelandiâ (Forst.), frequens in sylvis densis ad littora fluminum ad Kowa-Kowa, etc. (A. Cunn. l). Cineraria repanda Willd. sp. 3. p. 2076. non Lour. Arbuscula 12-15-pedalis. Folia petiolo 2-2 $\frac{1}{2}$ poll. instructa subrotunda suprà glabra subtùs adpressè cano-tomentosa penninervia rigidula, diam. 5-6 pollic. Panicula amplissima. (v. s. comm. à cl. Cunningham.)
2. B. ROTUNDIFOLIA (Forst. l. c.), foliis ovato-subrotundis integerrimis, paniculâ pauciflorâ. 5 in Novâ-Zelandiâ. Cass. dict. 48. p. 464 descr. Forst. ined. Cineraria rotundifolia Willd. sp. 3. p. 2076.

† *Species dubiæ hic ex patriâ adnexæ.*

3. B. ? FORSTERI, foliis subrotundis angulato-dentatis basi cuneatis subtus lanatis, petioliâ alatis basi nudis, corymbo terminali. § in Novâ-Caledoniâ (Spr. pug.), Novis-Hebridis (Spr. syst.). Cineraria Forsteri Spreng. pl. min. cogn. pug. 1. p. 58. syst. 3. p. 545.
4. B. ? CALEDONIÆ, foliis petiolatis ovato-oblongis sinuato-pinnatifidis inæqualiter denticulatis subtus pallidis obsolete lanatis; petiolis amplexicaulibus subalatis, corymbis terminalibus bracteatis. § in Novâ-Caledoniâ. Cineraria Caledoniæ Spreng. syst. 8. p. 547.

Trib. III. *ASTEROIDEÆ* Less. syn. 161. — Asteræ, Inuleæ pleræque et Helianthæ paucæ Cass.

Capitula sæpius heterogama, rariùs homogama aut dioica. Stylus fl. hermaphr. supernè cylindraceus bifidus, ramis longiusculis linearibus extùs subplanis sæpiùs subacuminatis rariùs obtusis, extùs æqualiter et minutè puberulis. Stigmatis series angustæ prominulæ usque ad originem pubis externæ extensæ. Corolla pellucida, staminigera tubulosa regulariter dentata. Pollen globosum echinulatum.

CONSPECTUS TRIBUS.

Subtrib. I. *ASTERINEÆ*. Capitula homo- aut heterogama, sæpiùs radiata. Anthæræ ecaudatæ. Folia ferè omnium alterna.

Div. I. *AMELLEÆ*. Recept. paleaceum. (XCIX—CII.)

Subdiv. 1. *EUAMELLEÆ*. Capitula heterogama radiata omnia recept. paleaceo donata. (XCIX—CI.)

Subdiv. 2. *HETEROPTHALAMEÆ*. Capitula dioica, mascula recept. nudo, foemineæ recept. paleaceo. (CII.)

Div. II. *ASTEREÆ* (1). Recept. epaleaceum. Capitula radiatâ ligulis heterochromis. (CIII—CLXIII.)

Subdiv. 1. *EUASTEREÆ*. Ligulæ 1-seriales. Pappus pilosus in disco et radio conformis, 1-serialis aut pluriserialis, setis æqualibus. (CIII—CXXIV.)

Subdiv. 2. *DIPLOPAPPEÆ*. Ligulæ 1-seriales. Pappus in disco et radio conformis, biserialis, serie exteriori brevi subpaleacea, interiori setosâ elongatâ. (CXXV—CXXXI.)

Subdiv. 3. *ERIGEREÆ*. Ligulæ pluriseriales. Pappus pilosus conformis aut inæqualis. (CXXXII—CXLI.)

(1) In hac divisione totâ capissimè secutus sum et auream Asterearum enumerationem à cl. Nees (Genera et species Asterearum 8°. Vratisl. 1832) propositam, et observationes sagacissimas manuscriptas à cl. Lindley (conf. Bot. reg. t. 1527) missas. Utrique gratias ex imo pectore perplurimas ago, ob adnotationum speciosissimas amicissimas communicationes.

Subdiv. 4. **HETEROPAPPEÆ**. Ligulæ 1-seriales. Pappus diversus in disco et radio. (CXLII-CLI.)

Subdiv. 5. **BELLIDÆ**. Ligulæ 1-seriales. Pappus paleaceus plurisquamellatus. (CLII-CLIV.)

Subdiv. 6. **BELLIDÆ**. Ligulæ uni- aut pluriseriales. Pappus nullus aut coroniformis. (CLV-CLXIII.)

Div. III. **CHRYSOCOMEÆ**. Recept. epaleaceum. Capitula lutea, nunc homogama floribus omnibus hermaphroditis, nunc radiata, ligulis cum disco homochronis (1). (CLXIV-CXCIII.)

Subdiv. 1. **GYMNOSPERMEÆ**. Pappus nullus. Ligulæ 1-seriales. (CLXIV-CLXVII.)

Subdiv. 2. **ACHYRIDÆ**. Pappus paleaceus. Ligulæ 1-seriales. (CLXVIII-CLXXI.)

Subdiv. 3. **HETEROTHECÆ**. Pappus in disco et radio diversus. Ligulæ 1-seriales. (CLXXII.)

Subdiv. 4. **PSIADIDÆ**. Ligulæ pluriseriales. Pappus pilosus in disco et radio conformis. (CLXXIII-CLXXIX.)

Subdiv. 5. **CHRYSOPSIDÆ**. Ligulæ 1-3-seriales aut nullæ. Pappus biserialis, ext. brevis subpaleaceus, inter. pilosus. (CLXXX-CLXXXI.)

Subdiv. 6. **SOLIDAGINÆ**. Ligulæ uniseriales aut nullæ. Pappus pilosus in disco et radio conformis, 1-serialis aut pluriserialis, setis æqualibus. (CLXXXII-CXCIII.)

Div. IV. **SOLENOGYNEÆ**. Capitula discoidea, floribus omnibus tubulosis, exterioribus fœmineis. (CXCIV-CXCVIII.)

Subtrib. II. **BACCHARIDÆ**. Capitula heterogama aut dioica nunquam radiata. Cor. omnes tubulosæ, fœmineis sæpius in ambitu multiseriales. Antheræ ecaudatæ. Recept. epaleaceum. Folia alterna.

Div. 1. **CONYZÆ**. Capitula heterogama monoica.

Subdiv. 1. **SPHÆRANTHÆ**. Capitula in capitulum sphæricum congesta ad bractearum axillas sessilia.

Subdiv. 2. **GRANGEINÆ**. Capitula in glomerulum non congesta. Pappus nullus aut subnullus.

Subdiv. 3. **EUCONYZÆ**. Capitula in glomerulum non congesta. Pappus pilosus.

Div. II. **BACCHARÆ**. Capitula dioica.

(1) Si quis autumat me, in divellendis Asteris et Chrysocomeis, colori nimium credidisse, errorem facile corrigat. Etenim in serie homochromæ Gymnospermæ repræsentant Bellideas, Achyridæ Bellideas, Heterothecæ Heteropappeas, Psiadidæ Erigereas, Chrysopsidæ Diplopappeas, Solidaginæ Euaustereas. Divisio nostra à colorum classicâ diversitate petita multò magis cum affinitatibus veris convenit, quam ulla alia mihi cognita.

† *Species dubiæ hic ex patria adnexæ.*

3. B. ? FORSTERI, foliis subrotundis angulato-dentatis basi cuneatis subtus lanatis, petiolis alatis basi nudis, corymbo terminali. § in Novâ-Caledoniâ (Spr. pug.), Novis-Hebridis (Spr. syst.). Cineraria Forsteri Spreng. pl. min. cogn. pug. 1. p. 58. syst. 3. p. 545.
4. B. ? CALÉDONIÆ, foliis petiolatis ovato-oblongis sinuato-pinnatifidis inæqualiter denticulatis subtus pallidis obsolete lanatis, petiollis amplexicaulibus subalatis, corymbis terminalibus bracteatis. § in Novâ-Caledoniâ. Cineraria Calédoniæ Spreng. syst. 8. p. 547.

Trib. III. *ASTEROIDEÆ* Less. syn. 161. — Asteræ, Inulæ pleræque et Helianthæ paucæ Cass.

Capitula sæpius heterogama, rariùs homogama aut dioica. Stylus fl. hermaphr. supernè cylindræus bifidus, ramis longiusculis linearibus extus subplanis sæpiùs subacuminatis rariùs obtusis, extus æqualiter et minutè puberulis. Stigmatis series angustæ prominulæ usque ad originem pubis externæ extensæ. Corolla pellucida, staminigera tubulosa regulariter dentata. Pollen globosum echinulatum.

CONSPECTUS TRIBUS.

Subtrib. I. *ASTERINEÆ*. Capitula homo- aut heterogama, sæpiùs radiata. Anthæræ ecaudatæ. Folia ferè omnium æterna.

Div. I. *AMELLEÆ*. Recept. paleaceum. (XCIX—CII.)

Subdiv. 1. *EUAMELLEÆ*. Capitula heterogama radiata omnia recept. paleaceo donata. (XCIX—CL.)

Subdiv. 2. *HETEROTHALAMEÆ*. Capitula dioica, mascula recept. nudo, foemineæ recept. paleaceo. (CII.)

Div. II. *ASTEREÆ* (I). Recept. epaleaceum. Capitula radiata ligulis heterochromis. (CIII—CLXIII.)

Subdiv. 1. *EUASTEREÆ*. Ligulæ 1-seriales. Pappus pilosus in disco et radio conformis, 1-serialis aut pluriserialis, setis æqualibus. (CIII—CXXIV.)

Subdiv. 2. *DIPLOPAPPEÆ*. Ligulæ 1-seriales. Pappus in disco et radio conformis, biserialis, serie exteriori brevi subpaleacea, interiori setosâ elongatâ. (CXXV—CXXXI.)

Subdiv. 3. *ERIGEREÆ*. Ligulæ pluriseriales. Pappus pilosus conformis aut inæqualis. (CXXXII—CXLI.)

(*) In hac divisione totâ sapiissime secutus sum et anteam Asterearum enumerationem à cl. Nees (Genera et species Asterearum 8°. Vratisl. 1832) propositam, et observationes sagacissimas manuscriptas à cl. Lindley (conf. Bot. reg. t. 1527) missas. Utrique gratias ex imo pectore per plurimas ago, ob adnotationum specimina et amicissimas communicationes.

squamulosi r-cephali. Capitula radio cœruleo, disco luteo. Cor. disci exhis pubescentes.

* Fruticosi.

1. *A. LYCHNITIS* (Linn. sp. 1276. mant. 476), caule fruticoso ramoso subdiffuso, foliis oppositis alternisve spathulato-lanceolatis obtusis integerrimis utrinque ramisque strigis minimis sericeo-canescens. ⑤ ad Cap. Bonæ-Spei ad Riet-Valley (Ecklon!), etc. Lam. ill. t. 682. f. 1. Jacq. coll. 5. t. 10. f. 1. Verbesina asteroides Linn. spec. ed. 1. p. 902. Buphthalmum Linn. hort. cliff. p. 415. — Batsyn. prod. 3. t. 15. f. 2. ex Willd. (v. v. c. et s. sp.)

2. *A. ruscifolius* (Burm. prod. 28. Lam. dict. 1. p. 129), caule fruticoso erecto ramoso, ramis tenuissimè pubescentibus, foliis alternis linearibus subacutis integerrimis involucrique tenuissimè hirtello-pubescentibus. ⑤ ad Cap. Bonæ-Spei (Burm. Eckl. in h. Dunant), ad Riebeck kastel (Drege!). *A. lychnitis* E. Mey. in Drege coll. Frutex pedalis. Folia $1\frac{1}{2}$ poll. longa, lineam lata. Capitula omnino *A. Lychnitis*. (v. s.)

3. *A. hispida*, caule basi suffruticoso ramoso erecto setis patentibus hispido, foliis alternis lato-linearibus setuloso-asperis aliis integerrimis dente (imo lobulo) utrinque r-a-notatis, invol. squamis dorso strigillosis apice acuminatis paleas referentibus. ⑤ in Africa Capensi ad Camiaberger alt. 3-400, et ad Zw. Kouse et Gariep alt. 1-200 ped. legit cl. Drege! Differt ab *A. anisato* Cass. ob paleam adpressam inter setas nullam et caulem fruticosum. (v. s.)

β. *angustissimus*, pilis patulis confertis aliis brevibus aliis longis setosis, foliis angustissimis omnibus ferè integerrimis. ⑤ ad Olifant rivier legit cl. Drege! (v. s.)

4. *A. flosculatus*, caule fruticoso ramosissimo, ramis foliisque pube brevi subvillosis, foliis linearibus integerrimis subconvolutis vix acutis, capitulis terminalibus solitariis subcorymbosis, invol. squamis dorso puberulis acuminatis, ligulis nullis. ⑤ in Africa austr. ad Klein Namaqualand, Zwarte Kouse et Gariep legit cl. Drege! Similis *A. hispida* sed discoidens et ex omni parte potius pubescens quàm hispida. (v. s.)

5. *A. scabridus*, caule fruticoseo ramoso sparsè et subadpresso setuloso, foliis linearibus acutis integerrimis setis subadpressis rigidis sparsis utrinque et ad marginem scabris, invol. squamis dorso hirtello-scabris acuminatis. ⑤ in distr. Zwelendama legit cl. Burchell (n. 6815), in deserto Karro (Eckl. Zeyh.). Caulis rigidulus tenuis. Rami non atriati. Folia 7 lin. longa vix lin. lata. Capitula trita anisum spirantia, ferè dimidiò minora quàm *A. Lychnitis*. Ligulæ pallidæ. Achænia pleraque compressa margine callosa demùm ferè calva. (v. s.)

** Herbacei annui.

6. *A. anisatus* (Cass. dict. 37. p. 489), caule herbaceo erecto simplici pube minimè adpressâ setisque longis articulatis sparsis instructo, foliis oppositis alternisve linearibus integerrimis utrinque pubescenti-scabris. ① ad Cap. Bonæ-Spei. Nees ést. p. 243.

7. *A. strigosus* (Less. in Linnæa 1831. p. 110. syn. p. 162), caule herbaceo ascendente flexuoso debili breviter et patentim piloso, foliis linearibus lanceolatis basi attenuatis integerrimis hirsutis. ① ad Cap. Bonæ-Spei in arenosis campis, Graona Kloof Swartlandiæ (Thunb.), in regione Trans-Gariepianâ (Burch. cat. geogr. n. 2276), in Karro prope Gannitz rivier (Eckl. et Zeyh.). *Aster strigosus* Thunb. fl. cap. 687. (v. s.)

8. *A. corymbosus*, radice subsimplici, caulibus pluribus herbaceis teretibus pubescentibus, centrali erecto, lateralibus ascendentibus, foliis linearibus integerrimis adpressè pubescentibus, pedunculis sub capitulo incrassatis cavis, invol. squamis adpressis pubescentibus. ① in Africa australi regione Gariepianâ legit cl. Drege! Caulis 5-6 poll. Folia 1-1 $\frac{1}{2}$ poll. longa, lineam lata. Ligulæ numerosæ cœmææ. Recept. paleæ pleraque flores disci superantes. (v. s.)

9. *A. NANUS*, caule herbaceo à basi ramoissimo cespitoso foliisque confertè et brevè puberulis, foliis elongato-lineari-lanceolatis integerrimis basi attenuatis, capitulis brevissimè pedicellatis foliis 1-2 stipatis, invol. squamis linearibus subtorulosis, ligulis amplis obtusis involucri duplò longioribus, rec. paleis paucis obtusis. (1) in Africæ Capensis regione Gariepina legit cl. Drege! Radix simplex perpendicularis. Herba 3 poll. longa, 2 lin. lata. Pappus duplex, ext. per anthesin vix perspicuus paleis ovatis, inter setis scabris deciduis. (v. s.)
10. *A. ANNUUS* (Willd. sp. 3. p. 2215), caule herbaceo subadscendente ramoso patentim et molliter pilosiusculo, foliis linearibus pubescentibus utrinquè attenuatis plerisque medio utrinquè unidentatis. (1) ad Cap. Bonæ-Spei. Nees ast! p. 224. excl. syn. Lam. et Spr. Am. alternifolius Roth cat. bot. 2. p. 111. Kaulfussia amelloides Zeyh. ! pl. exs. n. 99. Radix fibrosa caulis à basi ramosus. (v. s. in h. Moric.)
11. *A. TRIDACTYLUS*, caulibus è collo pluribus diffusis, foliis pedunculis involucrique setis patentibus rigidis confertè hispidis, foliis petiolatis tripartitis lobis linearibus oblongis integerrimis. (1) in Africâ australi legit cl. Drege! *Leysera tridactyla* E. Mey. ! in Dreg. coll. Rec. paleæ flores disci superantes. (v. s.)
12. *A. MICROGLOSSUS*, caule diffuso ramoissimo tereti parcè et patentim puberulo, foliis linearibus oblongis basi attenuatis integerrimis glabriusculis, invol. squamis linearibus complicatis acuminatis dorso hispidulis, ligulis minimis. (1) in Africâ Capensi ad Olifant rivier, Ziverfontain, et Klein Namaqualand legit cl. Drege! *Achænia* radii et exteriora disci compressa pubescentia pappo brevi paleaceo dentato coronata; disci intima abortiva. Paleæ ad latus fl. disci exter. angustissimæ, in centro disci annullæ. (v. s.)

† Species exclusæ.

- A. CAROLINIANUS* Walt. = *Eclipta brachypoda*.
A. DIFFUSUS Forst. = *Chiliotrichium amelloideum*.
A. FLORIBUNDUS Willd. h. = *Andromachia igniaria*.
A. PEDUNCULATUS Ort. = *Tridax procumbens*.
A. ROSMARINIFOLIUS Poepp. = *Chiliotrichium amelloideum* γ.
A. SPINULOSUS Pursh. = *Aplopappus spinulosus*.
A. UMBELLATUS Linn. = *Liabum Brownei*.
A. VILLOUS Parsh. = *Chrysopsis villosa*.

C. CORETHROGYNE.

Capitulum multiflorum, fl. radii ligulatis neutris 1-serialibus disci tubo brevi fauce cylindricâ apice brevissimè 5-dentatâ. Invol. squamæ 2-3-seriales lanceolato-lineares subæquales. Recept. planum paleis linearibus apice scabridis inter flores onustum. Styli rami exserti basi lineares margine glandulosi apice setas rigidas vel penicilliformes gerentes. Achænium turbinatum densè sericeum, villis supremis pappi exterioriorem simulantibus. Pappus pluriserialis setosus scabridus. — Herba Californica multicaulis tota molliter incano-tomentosa. Folia alterna acuta, infer. oblongo-lanceolata basi attenuata apice serrata, super. linearia integerrima. Capitula ad apices ramorum solitaria. Invol. parcè villosa. Ligulæ angustæ pallidæ ex sicco purpurascentes. Discus flavus. Pappus rubens. — An *Amello*! an *Osmiti* affinis? — Nomen ex *κορηθρον*, scopæ, et *γυνή*, fœmina.

1. *C. CALIFORNICA*. ♀ in Californiâ anno 1833 legit indef. Douglas. (v. s. comm. à cl. Benthani.)

CI. CHILIOTRICHUM Cass. bull. philom. 1817. p. 69. dict. 4. suppl. 576. 8. p. 576. 37. p. 463 et 489. Less. syn. 162. Nees ast. 239. — *Tropidolepis Tausch in florâ 12* (1829). p. 68. — *Amelli* sp. Forst.

Capitulum multiflorum, fl. radii ligulatis fœmineis 1-seriatis, disci 5-dentatis hermaphr. Invol. squamæ imbricatæ oblongæ acutæ. Recept. convexum paleis linearibus apice barbatis inter flores onustum. Stigmata fl. disci subulato-lineariter elongata pubera. Achænia gracilia cylindracea angulato-striata. Pappus pluriserialis, setis filiformibus scabridis inæqualibus persistentibus. — Frutices in extremâ Amer. austr. spontanei parvi ramosi. Folia alterna sessilia coriacea integerrima margine revoluta suprâ glabra subtus plus minus tomentosa. Pedunculi solitarii 1-cephali tomentosi. Ligulæ albæ subtus purpurascens. — *Aster phyllicoides* H. B. et Kunth fortè ex Nees huic generi adjungendus?

1. C. AMELLOIDES (Cass. dict. 8. p. 576), foliis oblongo-ovatis basi angustatis planiusculis. ♂ ad fretum Magellanicum (Forst. Comm.!), ad portum Egmont (Née!). Amellus diffusus Forst. comm. Gœtt. 9. p. 39. *Tropidolepis diffusa* Tausch in florâ 12, p. 68. *Aster magellanicus* Spreng. syst. 3. p. 526. Ch. amelloides γ Nees ast. 240. Folia subtus densè et fulvo-tomentosa. (v. s.)

β. *lanceolatum* (Nees l. c.), foliis lanceolatis acutis basi attenuatis. ♂ in insulis Maclovianis ubi arborum gigas interdum 4-5-pedalis, sed in monte Chatellux vix semipedalis d'Urv. ! fl. mal. n. 80. Gaudich. ! fl. mal. p. 16. *Arbuste à feuilles de romarin* Pern. mal. 2. p. 61. Eadem in Staten-Island legit cl. Webster ex h. Lindl. ! (v. s.)

γ. ? *rosmarinifolium* (Nees l. c.), foliis linearibus intensius margine revolutis basi non angustatis. ♂ in Andibus Chilensium (Pœpp.!), ad Talca-regué (Cl. Gay!), Amellus rosmarinifolius Pœpp. ! exs. Ch. rosmarinifolium Less. in linnæâ 1831. p. 109. An species propria? (v. s.)

Subdiv. II. HETEROThALAMÆ.

Capitula dioica; masc. recept. nudum, ligulæ 1-seriales; fœm. recept. paleaceum, ligulæ pluriseriales.

CH. HETEROThALAMUS Less. in linnæâ 1830. p. 145. 1831. p. 149 et 503. 179. p. 205.

Capitula multiflora radiata dioica. Involucrum pluriserialis, squamis siccis lanceolatis æqualibus disco subbrevioribus. Pappus 1-serialis setaceus caducus corollâ brevior. ♂ Recept. planum nudum. Ligulæ uniseriales fœmineæ minimæ, fructu abortivo. Cor. disci tubulosæ 5-dentatæ, antheris ecaudatis, stylo bifido abortivo, pappo clavato. ♀ Recept. conicum paleis invol. similibus onustum. Cor. ferè omnes? ligulatæ breves fœmineæ, centrales paucissimæ fortè 5-dentatæ masculæ? Achænia angulata glaberrima basi angustata, pappo apice attenuato. — Frutices Brasilienses glaberrimi erecti Athanasia aut Chrysocoma facie. Folia alterna lineari- aut oblongo-acuta integerrima aut serrata. Capitula subglobosa, ♂ corymbosa parva, ♀ in axillis ramorum steriliū breviter pedunculata duplò majora. Corollæ utriusque sexus flavæ, quo caract. et corollis plurimis ligulatis et recept. ? ♀ paleaceo à Baccharideis genus alienum videtur.

1. H. BRUNNIOIDES (Less. ! in linnæâ 1831. p. 504), foliis linearibus integerrimis. ♂ in Brasiliæ (Sellow!) prov. Rio-Grande (Gaudich.!), Marschallia (nec Melanthera) aliena Spreng. syst. 3. p. 446. (v. s. ♂ et ♀.)

2. *H. PSIADIOLDES* (Less. in linnæa 1831. p. 504), foliis oblongo-obovatis serratis. 3 in Brasiliæ (Sellow) prov. Rio-Grande (h. Mus. imp. Bras. n. 958). (v. s. in h. Mus. reg. Par.)

Div. II. ASTEREÆ.

Recept. epaleaceum nunc nudum nunc alveolatum aut fimbrilliferum. Capitula heterogama, fl. radii femineis ligulatis heterochromis (nempe albis purpureis cœruleisve seu cyanicis, disco luteo seu xanthico.)

Subdiv. 1. EUASTEREÆ.

Ligulæ 1-seriales. Pappus conformis pilosus uni- aut pluriserialis. Achænia compresso-plana.

CHII. MAIRIA. — *Mairia* et *Zyrphelis* Nees ast. p. 248 et 250. — *Pteropappus* (sect. *Asteris*) et *Zyrphelis* (sect. *Elphegæ*) Less. syn. 166 et 183. — *Asteris Arnice* et *Leysseræ* sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis femineis fertilibus, disci tubulosis 5-dentatis hermaphroditis nunc fertilibus nunc sterilibus. Recept. nudum subalveolatum. Involucrum pauciserialis subimbricatum. Antheræ basi omnino muticæ. Achænia plano-compressa glabra aut parcè pilosa erostria. Pappus 1-serialis, setis plumosis. — Herbæ rhizomate suffrutescente aut suffrutices Capenses. Folia alterna oblongo-obovata aut linearia, subrigidula, dentata aut integra. Rami caulesve apice subnudi capitula solitaria gerentes. Discus luteus. Radius violaceus aut albus. — Inter nomina 3 æquali jure admittenda *Mairiam* elegi, ne peregrinator de botanicâ Capensi bene meritis honore debito fraudetur.

Sect. 1. PTEROPAPPUS Less. syn. 166. — *Mairia* Nees ast. 248.

Flores disci fertiles. Ligulæ stam. filamenta sterilia nonnulla sæpius gerentes. — Herbacæ rhizomate perenni sublignoso.

1. *M. CRENATA* (Nees ast. 348), foliis subradicalibus obovato-oblongis basi attenuatis carnosiusculis apice grossè crenato-dentatis glabris, pedunculo hirsuto, involucri squamis dorso æqualibus aut longioribus. 2/ ad Cap. Bonæ-Spei in mont. Langkloof (Th.). Tabulari (Mundt et Maire Dreg.), circa urbem et in distr. Zwelendani (Eckl. et Zeyh.) *Arnica crenata* Thunb. fl. cap. 668. Ecklon! it. 1. n. 54. b. Lodd. bot. cap. t. 901. *Gerbera crenata* bot. reg. t. 855. Eckl. in h. Dunant. *Aster crenatus* Less. syn. 166. Pedunculus foliis brevior, æqualis aut duplò longior, nempe 1-4 pollicaris. Ligulæ violacæ. Capitulum magn. *Asteris* alpini. Variat foliis glabris aut pilosiusculis. (v. s.)
2. *M. HIRUTA*, caule adscendenti erecto subsimplici hirsutissimo, foliis in parte caulis ferè dimidiâ inferiore obovatis in petiolum attenuatis grossè dentatis supra juv. subhirsutis demùm glabris subtus densè hirsutis, superioribus paucis linearibus integerrimis hirsutis, capitulis caulem aut ramos paucos elongatos terminantibus, invol. hirsuto disco æquali. 2/ ad Cap. Bonæ-Spei in distr. Zwelendani in sylvis montosis prope Puspas-Valley leg. cl. Ecklon! et Zeyh. Affinis *M. crenatæ* sed differt caule ferè ad medium folioso, hirsutie omnium partium, et achænio ipso hirsuto. Ligulæ stam. rudimentis orbatæ. (v. s.)
3. *M. PEREZIOIDES* (Nees ast. 249), foliis subradicalibus lineari-spathulatis strigilloso-hirsutis subintegerrimis membranaceis, pedunculo-hispido-pubescente, involucri squamis disco sublongioribus. 2/ ad Cap. Bonæ-Spei in Grœnekloof et Swartland (Th.). *Aster perexioides* Less. syn. 167 *Leyssera pilosella* Thunb. herb. ex parte juxta Less. Facies *Perezia* squarrosæ.

Sect. II. *ZYRPHELIS* Cass. ann. sc. nat. vol. 17 (1829). p. 420. repet. in diss. segr. p. 36 et dict. sc. nat. 60. p. 597. Nees ast. 259. — *Elphegeæ* sp. Less. in linnæd.

Flores disci plerique steriles. Ligulæ stam. filamentis destitutæ.

§. 1. *Acaules.*

4. *M. BURCHELLII*, subacaulis, foliis radicalibus oblongo-linearibus glabris rigidè ciliatis, invol. squamis acuminatis ciliatis. $\frac{1}{2}$ ad Cap. Bonæ-Spei ad orientem urbis legit cl. Burchell et mecum comm. sub n. 8200 cat. geogr. Folia infernè nervo subtitis prominulo subtriquetra apice semicomplicata 4-5 pollic. longa. Caulis foliis paulò breviores. Invol. biseriale. Achænia subcompressa villosissima basi attenuata. Species ex habitu ad sect. primam primâ fronte referenda, sed ex disco sterili huc verè pertinens et sic utramque colligens. (v. s.)

§. 2. *Fruticosæ caulescentes.*

5. *M. TAXIFOLIA*, caule fruticuloso erecto ramoso, foliis linearibus acutis basi rigidè ciliatis, invol. squamis acuminatis, achæniis glabris $\frac{1}{2}$ ad Cap. Bonæ-Spei in collibus montis Tabularis (Th. ! Dreg. !) *Aster taxifolius* Linn. sp. 1225. Berg. cap. 286. *Leyssera ciliata* Thunb. fl. cap. 690. *Aster ciliaris* Thunb. ! in h. L'Hér. *Zyrphelis amœna* Cass. dict. 60. p. 598. *Elphegea ciliata* Less. in linnæd 1831. p. 126. *Zyrphelis taxifolia* Nees ast. p. 250. *Aster filiformis* Eckl. ! coll. exs. n. 103. b. in h. Moric. sed excl. syn. — Burch. ! cat. geogr. n. 592. Achænia matura compressa obovata glabra. Ligulæ sublilacino-violaceæ. (v. s.)
6. *M. MICROCEPHALA*, caule fruticuloso erecto ramoso, foliis linearibus basi parè ciliatis aut pube brevi hirsutulis, invol. squamis interioribus obtusis, achæniis pilosiusculis. $\frac{1}{2}$ ad Cap. Bonæ-Spei prope Kamnario (Mundt et Maire), ad orientem urbis (Burch. ! cat. geogr. n. 8159), ad Zeederberge (Dreg. !) in Zwellendam (Eckl. ! et Zeyh.). *Elphegea microcephala* Less. syn. 184. *Zyrphelis microcephala* Nees syn. 252. *Aster nanus* Ern. Mey. ! in Dreg. coll. Folia 4 lin. longa. Capitula dimidiò minora quàm prioris cui simillima. (v. s.)
7. *M. LASTOCARPA*, caule fruticuloso erecto ramoso, foliis linearibus acutis basi rigidè ciliatis, invol. squamis acuminatis, achæniis sericeo-villosis. $\frac{1}{2}$ ad Cap. Bonæ-Spei in distr. Caledon leg. cl. Ecklon ! et Zeyh. Simillima *M. taxifoliæ*, sed achæniis densè et adpressè sericeo-villosis facillè differt. (v. s.)

CIV. *FELICIA*. — *Felicia* et *Polyarrhena* Cass. bull. philom. 1818. p. 165. dict. 37. p. 464. et 490. 56. p. 172. Nees ast. 207 et 205. — *Polylepis* (1) (sect. *Asteris*) et *Polyarrhena* (sect. *Elphegeæ*) Less. syn. 173 et 184. — *Asteris* sp. Linn. et auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis 1-serialibus, disci tubulosis 5-dentatis hermaphr. sæpius omnibus fertilibus. Recept. nudum areolatum. Invol. squamæ bi- aut pluriseriales adpressæ angustæ. Achænia compressa glabra aut pubescentia. Pappus 1-serialis brevis caducus, setis flexuosis filiformibus serrulatis. — Herbæ aut frutices Capenses. Caulis ramosi. Rami 1-cephali sæpè fastigiati. Folia alterna angusta crassiuscula. Discus luteus, radius cœruleus aut albus.

(1) *Polylepis* fl. Per. est Rosacea-Sanguisorbea, et *Polylepis* Cass. est sect. *Sphaeranthi*.

Sect. I. HEBECARPÆA. — Feliciæ Cass. seu Polylepdis Less. species plurimæ.

Achaenia pilosiuscula aut subhirtella. — Folia linearia.

Herbaceæ, annuæ, radice fibrosâ.

1. *F. TENELLA* (Nees ast. 208), herbacea, foliis sparsis linearibus patulis setuloso-ciliatis, ramis floridis apice nudis 1-cephalis, invol. squamis linearibus acutis dorso subhirtellis, achæniis parvis et minutè pilosiusculis. (1) ad Cap. Bonæ-Spei, in collibus circa urbem ad Paarlberg (Dreg.), in reg. Gariepina (Dreg.), in distr. Caledon et Stellenbosch (Eckl. et Zeyh.). Aster tenellus Linn. mant. 471. Jacq. obs. 4. p. 88. Curt. bot. mag. t. 33. Less. syn. 173. Sieb. pl. exs. 253. Felicia fragilis Cass. dict. 16. p. 315. Aster dentatus et Aster tenellus Thunb. fl. cap. 686 Kaulfussia ciliata Spreng. f. suppl. syst. p. 25. Zeyh. pl. exs. n. 362. Cineraria tenella Link. enum. 2. p. 334. — Fluk alm. t. 271. f. 4. Variat caule erecto aut diffuso, foliis remotè aut crebrè serrato-ciliatis, radio pallidiùs aut intensius cœruleo. Fortè huc pertinet Cotula linifolia Burm. prod. fl. cap. 27. certè ex herb. radio donata. (v. sp. et v. c.)

♂ *longifolia*, caule basi suffruticuloso, ramis glabris teretibus, floriferis apice nudis 1-cephalis, foliis angustè linearibus sparsis vix acutis setulas breves dentiformes rigidas hinc inde ad margines gerentes, invol. squamis linearibus acutis dorso subhirtellis, achæniis maturis minutè et parvis pilosiusculis. ♂ ♀ in Africa Capensi ad Bergviver et Giffberg legit cl. Drege! ad Stellenbosch cl. Ecklon! et Zeyh. Variat foliis 1-1 ½ longis, semilineam latis, plus minus setuloso-ciliatis; pedicellis 1-4 poll. longis. An var. *F. tenellæ* frutescens? An spec. propria? (v. s.)

2. *F. COTULOIDES*, caule herbaceo tenello glabro à basi ramosissimo, ramis apice longè denudatis 1-cephalis, foliis angustè-linearibus acutis glabriusculis integerrimis, invol. squamis linearibus glabris acutis, achæniis vix sub lente puberulis, pappo multiseto. (1) in Africa Capensi ad Bergviver (Dreg.) in distr. Worcester (Eckl. et Zeyh.). Radix exilis. Herba 5-6 pollic. Folia 5-6 lin. longa 1 ½ lin. lata. Affinis *F. tenellæ*, sed præter not. ind. capitula duplò ferè minora. (v. s.)

3. *F. MICROSPERMA*, radice simplici, caulibus à collo pluribus herbaceis erectis apertibus versus apices glanduloso-puberulis viscosis (?) basi indivisis apice corymbosis, foliis linearibus integerrimis obtusiusculis, invol. squamis linearibus acutis achæniisque glabriusculis. (1) ? in Africa Capensis regione Gariepina legit cl. Drege! Radix et collum sublignescens. Caules 8-12 pollic. Folia 8-10 lin. longa, semilineam lata. Capitula *F. tenellæ*, sed laxè corymbosa. Achaenia pube brevi sparsa adpressa donata. (v. s.)

4. *F. ADPENSIS* (Nees ast. 209), herbacea, foliis obovato-oblongis hirsutis, inferioribus oppositis. (1) ad Cap. Bonæ-Spei. Aster affinis Less. syn. 174. Proxima *F. tenellæ*, sed caulis totus hirsutus nec basi glaber. Folia latiora basi angustata utrinquè hirsuta; invol. squamæ 3-4-seriales nec biseriales. Pappus disco corollæ æqualis nec è brevior. Fel. dubia Cass. dict. 16. p. 315. (v. s. in h. Cass. Mus. reg. Par.)

* * *Caule herbaceo simplicissimo rhizomate perenni orto.*

5. *F. MACRORHIZA*, caule herbaceo simplicissimo è rhizomate perenni orto, foliis alternis strigilloso-hirsutis obovato-linearibus confertissimis integerrimis, achæniis pilosis ciliatis. ♀ ad Cap. Bonæ-Spei. Aster macrorhizus Thunb. fl. cap. 687. Less. syn. 175. Nees ast. 32. Habitus *Erigerontis* graminei seu *Mairiæ peracoidis*.

6. *F. ERIGEROIDES*, caulibus herbaceis erectis teretibus tenuissimè pubescentibus, foliis elliptico-oblongis utrinquè glabris minutissimè serrato-ciliatis reticulato-nervatis, axillis foliiferis, ramis apice nudis 1-cephalis in paniculam ferè

dispositis, invol. squamis acutis glabriusculis, achæniis puberis. \mathcal{U} in Africâ austr. orient. ad Port-Natal legit cl. Drege! Habitu accedit ad *Erigeron foetidum*. Folia poll. longa, 3 lin. lata. (v. s.)

*** *Caule fruticoso.*

7. *F. FASCICULARIS*, caule fruticoso erecto ramoso glabriusculo, foliis linearifiliformibus patulis integerrimis rigidulis setis sparsis plus minus scabridis axillis folia novella fasciculata gerentibus, capitulis pedicellatis solitariis, invol. squamis linearibus acutis dorso pedicellisque minutè glandulosis. \mathcal{Z} in Africâ Capensis distr. Carro (Dreg.! Eckl.! et Zeyh.) in Uitenhage et territorio cesso (Eckl. et Zeyh.). *Aster fascicularis* E. Mey.! in Dreg. coll. *Felicia muricata* Nees ast. 210? *Aster filifolius* Zeyh.! exs. in h. Moric. Suffrutex circ. semipedalis spithameusve. Folia var. ferè glabra, caulina 3 lin., axill. 2 lin. longa achænium villosum. Pappus scaber. (v. s.)

β . *pubescens*, ramis pubescentibus elongatis, foliis rameis linearibus, ad axillas nudis. \mathcal{Z} in distr. Uitenhage (Eckl.! et Zeyh.) et prope Basche (Drege!). (v. s.)

8. *F. ANGUSTIFOLIA*, caule fruticoso erecto, ramis pubescentibus aut incanis, foliis linearibus planis ciliatis acutiusculis adultis glabris, capitulis longè pedunculatis \mathcal{Z} ad Cap. Bon.-Spei.

α . *hyssopifolia*, ramis canescentibus, foliis junioribus pube brevi molli subhirsutis. \mathcal{Z} Burm.! Dreg.! Eckl.! et Zeyh. *Aster hyssopifolius* Berg. cap. 287. Less. syn. 175. *Felicia hyssopifolia* Nees ast. 211. *Aster villosus* et *hirtus* Thunb. fl. cap. 687 et 688. ex Less. *Aster fruticosus* Burm.! prod. 27. *Conyza chrysocomoides* Desf. arb. 1. p. 232. excl. syn. *Aster chrysocomoides* Desf.! cat. h. par. 1815. p. 121. non Willd. *Aster conyzoides* Desf.! cat. h. par. 1829. p. 175 non Willd. (v. s.)

β . *glabra*, ramis vix pubescentibus, foliis etiam junioribus glabris. \mathcal{Z} ad Zeederbergen et Stormberg Dreg.! *Aster angustifolius* Jacq. h. Schoenbr. t. 370. Less. syn. 177. *Felicia angustifolia* Nees ast. 213. (v. s. et c.)

9. *F. HIRUTA*, caule fruticoso, ramis virgatis hirsuto-canescens, foliis linearibus planis obtusiusculis utrinquè strigilloso-hirsutis, capitulis solitariis breviter pedunculatis, achæniis hirsutiusculis. \mathcal{Z} ad Cap. Bonæ-Spei in regione Trans-Gariepinâ (Burch.! cat. geogr. n. 2331), in territorio cesso et Graf Reynet (Eckl.! et Zeyh.) ad Klaarprivier (Dreg.!). Folia 5-6 lin. longa erecta per totos ramos disposita, apice ramuli breviter denudato. Invol. campanulatum, squamis 2-3-ser. exter. dorso hirtis omnibus margine membranaceis. An fortè *Aster lanuginosus* Wendl. beytr. 2. p. 9. (v. s.)

10. *F. RIGIDULA*, caule fruticoso, ramis subadgregatis rigidulis puberis, foliis linearibus planis obtusis utrinquè adpressè strigilloso-hirsutis, capitulis solitariis pedunculatis, invol. squamis dorso pulverulento-mucronatis discum ferè superantibus, achæniis puberulis \mathcal{Z} ad Cap. Bonæ-Spei (h. Boiss.!) in distr. Uitenhage (Eckl.! et Zeyh.). Valdè ex characteribus accedit ad *F. hirsutam*, sed habitus satis diversus et inv. squamæ extus non hirsutæ divisionem suadent. (v. s.)

β . *subcanescens*, foliis ramulisque pube confertiore subcanescentibus. \mathcal{Z} in Africâ Capensi (Dreg.!), in distr. Uitenhage (Eckl.! et Zeyh.). (v. s.)

11. *F. CANA*, caule fruticoso adscendente ramosissimo, ramis foliisque pube confertâ brevi velutino-canis, foliis linearifiliformibus obtusis integerrimis, axillis sæpè foliiferis, pedicellis elongatis nudis 1-cephalis pubescentibus, invol. squamis linearibus acutis puberis, achæniis sparse pilosisculis. \mathcal{Z} in Africâ Capensis distr. Zwellendam et deserto Karro leg. cl. Ecklon! et Zeyh. Suffrutex 5-6 pollic. totus canus. Folia 5 lin. longa. Ligulæ albæ lineares. (v. s.)

12. *F. ZEYHERI* (Nees ast. 215), caule fruticoso, ramis adultis glabris, foliis oblongo-cuneiformibus planis glabris dense strigilloso-ciliatis, capitulis longè pedunculatis, invol. squamis glabris disco subbrevioribus acuminatis, achæniis

hirsutiusculus. ♂ ad Cap. Bon.-Spei prope Uitenhagen. *Conyza ciliaris* Spreng. in Zeyh. coll. exs. (a. 1828) n. 126. *Aster Zeyheri* Less. syn. 177. (v. s. in h. Moric.)

13. *F. DREGÆ*, fruticosa ramosa, ramulis foliis pedunculisque minutè et confertè pubescentibus, ramis apice nudis 1-cephalis, foliis oblongis utrinquè acutis integerrimis aut vix dentibus 1-2 minimis acutis notatis, invol. squamis linearibus apice subciliatis, achæniis villosiusculis. ♂ ad Cap. Bonæ-Spei in Klein Namaqualand legit cl. Dregè! Folia 7-8 lin. longa, 2 lin. lata. Cor. disci nascentes flavæ demum purpureæ. Ligulæ cœruleæ.

β. *dentata*, foliis obovatis obtusis dentatis, corollis disci flavis. ♂ cum var α legit cl. Dregè! (v. s.)

14. *F. LASIOCARPA*, caule fruticoso demisso glabro apice glanduloso pulverulento-subhirtello, foliis linearibus integerrimis acutis ad basin attenuatis et infernè semi-amplexicaulibus, infer. glabris, summis cum caule et invol. squamis extus glanduloso-pulverulentis, achæniis adpressè sericeo-villosissimis. ♂ in Africâ australi legit cl. Dregè! Caulis 4 pollic. Folia pollicem longa, lineam lata. Pappus rufescens densè serrulatus. Capit. 9-10 lin. diam. (v. s.)

Sect. II. *ANHEBECARPÆA*. — *Feliciæ Nees* seu *Polylepidis Less.* species et *Polyarrhena Cass. Nees et Less.*

Achænia glaberrima. Flores disci marginales in unâ aut duabus speciebus steriles, centrales fertiles. — Ipso suadente Neesio p. 217, *Polyarrhenam* Cass. cum *Feliciâ* habitu jubente junxi.

* *Foliis linearibus basi subattenuatis.*

15. *F. VICOIDEA*, glaberrima, caule basi radicante, foliis oppositis carnosis lineari-semiteretibus obtusis integerrimis basi subattenuatis, pedicellis terminabilibus nudis folia paulò superantibus, pappo caducissimo 5-6 seto. ♀ ad Cap. Bonæ-Spei in coloniæ parte orient. legit cl. Burchell. n. 5311. Etiam vidi in herb. Boiss. sine loci proprii design. Ligulæ albidæ. Herba 4-8 pollic. Folia serè pollicem longa, 1-1½ lin. lata. Capit. 3-4 lin. diam. (v. s.)

16. *F. CILIARIS*, caule fruticoso ramoso glabro, ramis floridis basi subfoliosis apice nudis 1-cephalis glanduloso-hirtellis, foliis alternis linearibus acutis utrinquè glabris margine plùs minùs setis rigidis ciliatis, invol. squamis linearibus acutis dorso glanduloso-hirtellis, achæniis glabris pulverulento-scabris, pappo paucisetò caducissimo scabro. ♂ in Africæ Capensis collibus distr. Zwelendama legit cl. Dregè! Habitus *Mairiæ taxifoliæ*. Ligulæ rubentes fertiles. Discus flavus sterilis. (v. s.)

β. *leiopoda*, pedicellis elongatis glabris. ♂ in Africâ austr. (Dregè!) fortè in Albany (Eckl. et Zeyh.). Achænia matura ignota et ideò stirps dubia. (v. s.)

17. *F. CAFFRORUM* (Nees ast. 214), caule fruticoso ramosissimo ramisque glabrusculis, foliis cuneiformi-linearibus planis acutiusculis glabris glanduloso-punctatis margine glanduloso-scabris, pedunculis brevibus corymbosis. ♂ ad Cap. Bon.-Spei in terrâ Caffrorum. *Aster Caffrorum* Less. syn. 179. Krebs coll. n. 178 ex Less.

β. *validior*, foliis angustioribus margine minùs glandulosis Nees l. c. *Aster linoides* hort. Gaybac. 1816.

18. *F. RTORTA*, glaberrima, caule fruticoso, ramis usquè ad capitula foliosis, foliis linearibus acutis revoluto-reflexis, nervo ad basin in angulum decurrentem productò, invol. squamis exterioribus revoluto-reflexis, intimis longioribus erectis. ♂ ad Cap. Bonæ-Spei non longè à Genadendal legit cl. Burchell et mecum sub n. 7793 cat. geogr. comm. Species distinctissima et fortè genericè separanda. Achænia vix compressa angustiora et longiora quàm in *Asteris* aliis Capensibus. (v. s.)

19. *F. IMBRICATA*, fruticosa, ramis foliis imbricatis densè vestitis apice summo vix nudis hirsutis 1-cephalis, foliis linearibus subacutis basi non attenuatis

crassiusculis nervo subprominulo ferè trigonis ad margines et rarius ad nervi apicem setis rigidis ciliatis, invol. squamis linearibus ext. dorso hirsutulis, intimis glabris subscariosis. ζ ad Cap. Bonæ-Spei in distr. Worcester leg. cl. Ecklon! et Zeyh. Folia 4 lin. longa, lineam lata. Capitulum ferè F. echinatae. Achænium glabrum. Pappi setæ densè et breviter barbellatæ. (v. s.)

* * *Foliis ovato- aut ovali-sublanceolatis basi latiusculis.*

20. F. PARALIA, caule fruticoso ramoso glabro aut vix sub capitulo pubescente, foliis confertis basi latâ sessilibus oblongis acutis glabris margine lævibus aut summis parcè ciliatis demùm reflexis, capitalis brevissimè pedunculatis, invol. squamis dorso lævibus. ζ ad Cap. Bonæ-Spei in col. orient. non longè ab Uitenhagen legit cl. Burchell n. 3733 et 3814, in distr. Albany (Eckl.! et Zeyh.). Folia ferè Euphorbiæ Paraliæ. Valdè accedit ad F. echinatum; ramis glabris, foliis margine lævibus et invol. squamis dorso glabris faciliè Burchellii n. 3733 differt, sed specim. 3814 foliis superioribus parcè ciliatis et ramulo florifero subhirsuto transitionem ferè indicat. (v. s.)

21. F. ECHINATA (Nees ast. 216), caule fruticoso, ramis hirsutis 2-3-chotomis, foliis confertis ovato-oblongis acuminatis basi latâ sessilibus glanduloso-punctatis margine ciliato-serratis demùm reflexis, capitalis brevissimè pedunculatis, invol. squamis dorso setoso-scabris. ζ ad Cap. Bon.-Spei in Plettenbergsbay arenosis (Mundt et Mair.), in Antiqualand (Burch. ! cat. geogr. n. 5516), in Outniqualand (Dreg.!), in distr. urbis (Eckl.! et Zeyh.) Pteronia echinata Thunb. fl. cap. 629. Aster echinatus Less. syn. 177. Facies F. reflexæ sed præter not. indic. capitula majora, folia suprâ subconca, inv. squam. acuminatæ. (v. s.)

22. F. REFLEXA, caule fruticoso ramisque glabris, foliis basi latâ sessilibus ovato-sublanceolatis acutis revolutis-reflexis utrinquè glabris margine serrato-ciliatis, capitalis subsessilibus, invol. squamis subobtusis dorso glabris aut parcè setoso-scabris, floribus disci marginalibus sterilibus. ζ ad Cap. Bonæ-Spei inter frutices montis Tabularis, Leonis, etc. (Burch. ! n. 8478. Dreg. ! Eckl. ! et Zeyh.). Aster reflexus Linn. sp. 1225. Berg. cap. 285. Burm. ! prod. 26. Andr. bot. rep. 2. t. 93. Curt. bot. mag. t. 884. Aster reflexus et Chrysocoma scabra Thunb. fl. cap. 689 et 627 ex Less. Polyarrhena reflexa Cass. dict. 56. p. 173. Nees ast. 206. Elphegea reflexa Less. syn. 184. Cullumia setosa Sieb. ! fl. cap. exs. n. 27 excl. syn. — Comm. hort. amst. 2. t. 28. Variat foliis plus minus ciliatis serratisve, pedunculis plus minus hirsutis, invol. squamis dorso lævibus aut parcè hispidis. Capitula 9-10 lin. diam. (v. s.)

† Species exclusæ.

F. BRACHYGLOSSA Cass. = Munychia cymbalariae.

F. DURLA Cass. = Munychia cymbalarioides ?

CV. MUNYCHIA Cass. dict. 37. p. 462 et 483. — Felicia Asteris subgenus Less. syn. 181. non Nees. — Feliciæ sp. Cass. dict. 25. p. 97. — Asteris et Cinerariæ sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis brevibus fertilibus 1-serialibus fœmineis, disci tubulosos hermaphrod. 5-dentatis fertilibus aut abortu sterilibus. Recept. nudum. Invol. squamæ subtriseriales adpressæ. Stigmata filiformia elongata. Achænia compressa subhirtella. Pappus, setis pilosis subserrulatis 1-serialibus brevibus caducis. — Herbæ Capenses diffusæ pilosiusculæ ramosæ. Folia petiolata opposita ovata parcè et grossè dentata papyracea. Pedunculi terminales et axillares aphylli elongati. Capitula solitaria, disco flavo, radio albo. Achænia matura nigricantia, pappo albisimo.

F. M. HIRUTA, fruticosa, ramis decumbentibus, foliis petiolatis ovatis parcè

dentato-sinuatis hispidulis, involucri 2-3-seriali, floribus disci fertilibus, ligulis apice grossè dentatis. 3 ad Cap. Bonæ-Spei. Cineraria hirsuta Vent. malm. n. et t. 95.

2. *M. CYMBALARIÆ* (Nees ast. 218), herbacea, foliis petiolatis ovatis dentatis, floribus capituli omnibus fertilibus, invol. glabro subbiseriali. 2 ad Caput Bonæ-Spei inter saxa montis Tabularis (Mundt et Mair. Dreg.) etc. Aster cymbalarie Thunb. prod. fl. cap. 689. Felicia brachyglossa Cass. dict. 25. p. 97. Mun. brachyglossa Cass. dict. 37. p. 483. Aster oppositifolius Linn. amœn. 5. p. 369? (v. s.)

β. *microcephala*, capitulis paulò minoribus, floribus centralibus subabortivis, caule laxè diffuso. — Burch. ! cat. geogr. n. 647 et 8466. (v. s.)

3. *M. CYMBALARIOIDES*, foliis oblongis subdentatis utrinquè attenuatis, floribus disci abortu sterilibus, involucri uniseriali. ① ad Cap. Bonæ-Spei in arenosis lapideosis jugi Leonis (Eckl. ! in h. Dunant. it. 2. n. 1050). Folia pleræque radicalia basi angustata in petiolum latum trinervium, ciliis longis distantibus ornatum. Pedunculus scapiformis. Achænia radii plano-compressa subelliptica nigricantia margine ciliato-villosa cæterum vix pilosiuscula, omninò ut in priora, sed fl. disci abortivi. An fortè Felicia dubia Cass. dict. 16. p. 315? (v. s.)

CVL. AGATHEA Cass. bull. philom. 1815. p. 175. dict. i. suppl. p. 77. 13. p. 116. et 37. p. 463 et 489. — Detris Adans. fam. 2. p. 131. — Detrioides (sect. Elphegæ) et Detris (sect. Asteris) Less. — Detridium et Agathæa Nees. — Asteris sp. Mill. — Cinerariæ sp. Linn.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis, disci tubulosis 5-dentatis hermaphr. fertilibus aut rariùs sterilibus. Recept. nudum planum vix alveolatum. Invol. campanulati squamæ 1-2 seriales adpressæ complicato-carinatæ subherbaceo-rigidulæ. Achænia matura plano-compressa nervo tumidulo cincta, pilis rigidulis scabra (in A. microphyllâ lævia), sterilia glabra. Pappus 1-serialis, setis caducis aut persistentibus serrato-scabris. — Herbæ aut suffrutices Capenses. Folia omnia aut saltem infima opposita, integra aut parçè dentata, hispidula aut rariùs glabra. Pedunculi nudi 1-cephali. Discus luteus. Radius cœruleus.

Sect. I. DETRIDIIUM — Detridium Nees syn. 255. — Elphegæ sect. Detrioides Less. syn. 182.

Flores disci steriles. Pappus disci et radii pauci- (5-6) setus.

1. *A. BERGERIANA*, caule herbaceo simplici aut ramoso erecto molliter hispido, foliis plerisque alternis oblongo-obovatis acutis integris, capitulis longè pedunculatis. ① ad Cap. Bonæ-Spei in arenosis lapideosis jugi Leontini. Cineraria Bergeriana Spreng. syst. 3. p. 551. Elphegæ Bergiana Less. syn. 182. Detridium Bergianum Nees syn. 256. Kaulfussia amelloides Eckl. ! in h. Dunant. Variat caule simplici 1-cephalo et parçè ramoso 2-7-cephalo. Habitus omninò sequentium. (v. s. in h. Boiss. et Dunant.)

Sect. II. DETRIS. — Detris Adans. l. c. — Asteris sect. Detris Less. syn. 168. — Agathæa Cass. ll. cc. Nees syn. 257.

Flores disci pleriquè fertiles. Pappus multi- (10-15) setosus.

* *Herbaceæ annuæ.*

2. *A. ROTUNDIFOLIA* (Nees ast. 258), caule herbaceo adscendente hirsutiusculo, foliis ovato-ellipticis obtusis integerrimis strigillosis uninerviis. ① ad Cap. Bonæ-Spei. Aster rotundifolius Thunb. fl. cap. 689. Less. syn. 172. Achænia flavo-fusca.
3. *A. SERRATA* (Nees ast. 259), caule herbaceo adscendente infernè glabro superne hispidulo, ramis usquè ad capitula densè foliosis, foliis verticillatis lan-

ceolatis acutis calloso-serrulatis glabris trinerviis. ① ad Cap. Bonæ-Spei. Aster serratus Thunb. fl. cap. 689. Less. syn. 173. Similis priori et magis adhuc ex et habitu Heteropappo hispido.

4. *A. DIFFUSA*, caule herbaceo tenui ramoso diffuso densè et breviter hirtello, foliis oppositis sessilibus elliptico-oblongis integerrimis hirtellis, pedicellis elongatis 1-cephalis nudis parè et brevissimè puberulis, invol. squamis linearibus acutis dorso hirtellis, achæniis maturis in disco utroquè ferè lævibus. ①? in Africâ Capensi legit cl. Drege! Pappus radii parvus caducissimus. Folia 2-3 lin. longa, lineam lata. (v. s.)
5. *A. STRICTA*, caule herbaceo erecto basi simplici sparsè hispido, foliis oppositis linearilanceolatis acutis basi attenuatis remotè serratis utrinquè hispidulis, ramis subpaniculatis confertè et brevè hirtellis, involucri squamis acutis subhirtellis, ligulis parvis. ① in Africâ Capensi ad Riebekkastel legit cl. Drege! Ex habitu stricto et radice exili fibrosâ ab *A. hispida* et *A. diffusa* facillè differt. Ligulæ minores quàm in aliis speciebus. Achænia juniora pallida puberula. (v. s.)
6. *A. TENERA*, caule herbaceo erecto oppositè ramoso hispidulo, foliis oppositis remotis sessilibus trinerviis ovatis seu ovato-oblongis distanter et calloso-serratis membranaceis subhispidulis, pedicellis elongatis nudis hispidulis, invol. squamis linearilanceolatis dorso hispidis apice membranaceis glabris. ① in Africâ Capensis regione Gariepina legit cl. Drege n. 2802! Herba 8-10 pollic. Capitula magnit. Agathææ amelloidis. Achænim oblongum pubero-villosum. (v. s.)

** *Frutescentes.*

7. *A. ZYTHERI* (Nees ast. 260), fruticosa, foliis oppositis alternisque linearilanceolatis pellucido-punctatis setoso-serratis, involucri squamis trinerviis, achæniis scabris. 3 ad Cap. Bonæ-Spei. Cineraria linifolia Zeyh. coll. exs. n. 53 in h. Moric. Kaulfussia amelloides var. Spreng. ex Nees. Folia 7-9 lin. longa, lineam sesquilineamve lata glabrescens. Ligulæ lineares elongatæ. (v. s. in h. Moric.)
8. *A. BARBATA*, caule fruticoso erecto ramoso glabro, ramulis setis longis hispidis, foliis oppositis rariùs alternis lineariloblongis basi attenuatis integerrimis margine et hinc indè subtus setis longis patulis ciliatis, ramis floridis basi foliolosis apice nudis glanduloso-hirtellis et subsetosis, invol. squamis dorso hirtellis margine subscariosis, achæniis utrinquè hirtis. 3 in Africâ Capensi ad Graaf-Reynet alt. 2-300 ped. legit cl. Drege! Fruticulus vix pedalis. Folia 6-8 lin. longa, 1-2 lata. Capitula vix *A. amelloidi* minora. (v. s.)
β? subalbescens, setis paucioribus, foliis pube adpressâ præter setas confertâ canescentibus minoribus. 3 ad Sneeuwbergen alt. 4-500 ped. legit cl. Drege! (v. s.)
9. *A. HISPIDA*, caule imâ basi fruticuloso ramosissimo cum foliis pedunculisque undiquè setis longis patulis rigidulis hispido, foliis alternis (imis paucis oppositis) linearilanceolatis obtusiusculis basi attenuatis integerrimis, invol. squamis setis et pube pulverulentâ extus scabris. 3 in Africâ Capensi ad Zeederbergen alt. 2-300 ped. legit cl. Drege! Cineraria hispida E. Mey. in Dreg. coll. Achænim atrum in disco utroque sparsè puberum. Invol. disco æquale. Ligulæ lineares invol. duplò longiores. (v. s.)
β. barbiger, pedunculis densius hispidis, invol. squamis apice floccoso-barbatis. 3 in regione Gariepina (Dreg.!). Flocci squamas terminantes etiam in var. *α* hinc indè obvii, sed in var. *β* majores et constantes videntur. (v. s.)
10. *A. HIRTA*, caule fruticoso ascendente ramoso, ramis elongatis teretibus infernè foliosis pube minutissimâ et setis patulis hispidis, supernè longè aphyllis 1-cephalis pube glandulosâ et setulis rigidis brevibus hirtellis, foliis lineariloblongis integerrimis margine subrevolutis setis rigidis ciliatis et hirtis, infer. oppositis, super. alternis, invol. squamis setulis ut in pedicello coloratis hirtel-

- lis, achæniis subglandulosis. ζ in Africâ Capensâ ad Breederivier legit cl. Drege! Pedicelli 5 poll. longi. Folia 5-7 lin. longa. 2 lin. lata. (v. s.)
11. *A. ECKLONIS* (Nees ast. 262), fruticosa, foliis alternis elliptico-oblongis subamplexicaulibus integerrimis subciliatis substrigosis suprà asperis, achæniis hirtulis, involucri squamis trinerviis. ζ ad Cap. Bonæ-Spei in calcariis inter Krede rivier et Duyrenhoks rivier. Aster Ecklonis Less. syn. 170. Setulæ paucæ secus folior. margines ut in *A. Zeyeri* cujus fortè varietas latifolia (v. s. in h. Dunant comm. à cl. Ecklon.)
12. *A. SCABRIDA*, caule fruticoso erecto ramoso glabro, ramulis pedicellisque minutè pulverulento-scabridis, foliis alternis cuneatis basi attenuatis integris ciliatis apice subtruncatis 3-5-dentatis utrinquè echinulato-scabridis, pedicellis foliola linearia integrâ paucâ gerentibus, invol. glabriusculis ζ in Africâ Capensâ legit cl. Drege! Aster scabridus E. Mey. ! in Drege. coll. (v. s.)
13. *A. BRAVIFOLIA*, caule fruticoso erecto ramoso glabro, ramis nascentibus subvelutinis, senescentibus subspinosis, floridis apice nudis 1-cephalis, foliis alternis pube confertâ brevi velutinis basi cuneatis apice obtusè 3-5-dentatis ferè lobatis, invol. squamis 3-serialibus linearibus glabriusculis, achæniis maturis parè in disco pilosiusculis. ζ in Africâ Capensâ ad Zilverfontain legit cl. Drege! Frutex 1-1 $\frac{1}{2}$ -pedalis rigidus. Folia 2 lin. longa. Cor. disci apice ru-bentes. Stigmata valdè hirta. Pappus rigidulus longus scabridus. (v. s.)
14. *A. LEOCARPA*, caulibus à collo pluribus diffusis basi suffruticosis hispidis, foliis alternis oblongo-cuneatis apice acutè 3-5-dentatis basi integris sessilibus utrinquè puberulis basi ciliatis pedicellis glanduloso-puberulis folio triplò longioribus, invol. squamis 1-2-seriatis lineari-lanceolatis acutis vix puberis, achæniis maturis glaberrimis. ζ \mathcal{L} in Africâ Capensâ ad Drakeenstensberge alt. 3-400 ped. legit cl. Drege! Ligulæ cœruleo-purpureæ. Achænia disci sterilia puberula, radii matura pappo deciduo calvescentia. Folia 7-9 lin. longa. (v. s.)
15. *A. MICROPHYLLA* (Cass. ! bull. philom. 1817. p. 183. dict. suppl. p. 63), fruticosa, foliis oppositis alternisque obovato-oblongis sessilibus suprà et subtus ad nervum strigosis, achæniis glaberrimis, invol. squamis trinerviis. ζ ad Cap. Bonæ-Spei, inter saxa montis Tabularis (Eckl. ! ad Znoëreberge (Dreg. !) in distr. Caledon et circa urbem (Eckl. ! et Zeyh.) et in distr. Albany (Borch. ! cat. geog. n. 3966) Nees ast. 261. Aster æthiopicus Burm. prod. 27. Less. syn. 169. Cineraria trachyphylla Spreng. syst. 3. p. 547. Cineraria microphylla Vahl herb. ex Less. sed habui ex eod. herb. sub nom. Cinerariæ amelloïdia. Habitus sat varius. Rami hispiduli aut glabri. Folia interdum subglabra, sed achæniis glaberrimis imò lævibus à genere distinctissima. (v. s.)
16. *A. STRIGOSA* (Nees ast. 263), fruticosa, foliis oppositis oblongo-cuneatis integerrimis erecto-adpressis apice recurvis utrinquè ramisque densè strigosis, achæniis scabris, pappi setis subplumosis, invol. squamis 1-nerviis. ζ ad Cap. Bonæ-Spei in Roggeveld, Karro, Uitenhagen, etc. Leyseria ovata Thunb. fl. cap. 692. Kauffussia strigosa Spreng. f. suppl. syst. p. 25. Zeyh. ! coll. ex n. 270. Aster strigosus Licht. herb. non Thunb. Aster Charicis Less. syn. 171. (v. s. in h. Moric.)
17. *A. AMELLOIDES*, fruticosa, foliis oppositis ovato-vel orbiculato-ellipticis in petiolum brevem attenuatis integerrimis ramisque strigosis, achæniis punctatis et scabris, invol. squamis 1-nerviis. ζ ad Cap. Bonæ-Spei in lateribus montis Tabularis. Frequens in hortis Europæis. Cineraria amelloïdes Linn. sp. 1245. Berg. cap. 290. Agathæa cœlestis Cass. bull. philom. 1815. p. 175. dict. 1. suppl. p. 77. opusc. phyt. 6. Nees ast. 264. Aster Capensis Less. syn. 168. — Mill. icon. t. 76. f. 2. (v. v. c.)
18. *A. ELONGATA* (Nees ast. 266), fruticosa, ramis alternis glanduloso-hispidis, foliis oppositis sessilibus oblongis acutis integerrimis utrinquè patentè-hirsutis achæniis hirsutiusculis, invol. squamis angustis uninnerviis ligulis unicoloribus. ζ ad Cap. Bonæ-Spei in arenosis Swartlandiæ et Saldanhabay (Th.). ad flum. Gariep (Dreg. !). Aster elongatus Thunb. fl. cap. 688. Less. syn. 170. Cineraria

ria hirta Vahl herb. ex Less. Invol. squamæ ext. hispidulæ, inter. glabræ submembranaceæ. (v. s.)

19. A. TRICOLOR (Nees ast. 267), fruticosa?, foliis oppositis oblongo-lanceolatis integerrimis utrinque patenti-hirsutis, invol. squamis angustis 1-nerviis, achæniis setuloso-scabris, ligulis bicoloribus. ♀ Pad Cap. Bonæ-Spei.

CVII. BELLIDIASTRUM Mich. nov. gen. t. 29. Cass. dict. sc. nat. 4. suppl. p. 70. 37. p. 494 non Vaill. nec Less. — Margarita Gaud. fl. helv. 5. p. 335. — Doronicum sp. Linn. — Arnica sp. Hall. — Asteris sp. Less.

Capitulum radiatum, ligulis 1-2-seriatis heterochromis fœmineis, disci tubulosis hermaphroditis. Involucrum subcampanulatum, squamis 1-2-seriatis linearibus acutis subfoliaceis. Recept. conicum nudum punctatum. Stam. in fœmineis omninò abortiva. Achænia omnia oblonga subcompressa hispidula. Pappus 1-2-serialis, pilis numerosis flexuosis scabris. — Herba Europæa perennis, trunco brevissimo radiciformi, foliis radicalibus petiolatis obovato-oblongis dentatis, scapo foliis longiore villosa 1-cephalo. Capitulum ferè Bellidis. — Cum Astere fortè non immeritò jungunt cl. Lessing et Lindley.

1. B. MICHELII (Cass. l. c.) 2½ in sylvis montosis Europæ. Doronicum Bellidiastrum Linn. sp. 1247. Jacq. fl. austr. t. 400. Arnica Bellidiastrum Vill. dauph. 3. p. 212. Sturm deutschl. fl. f. 38. t. 15. Lam. et DC. fl. fr. n. 3201. Aster Bellidiastrum Scop. carn. n. 1044. Less. syn. 166. Margarita Bellidiastrum Gaud. l. c. Bellidiastrum montanum Hoppe! pl. exs. 1832. Variat ligulis albis aut extus plüs minüs, interdum omninò rubris. (v. s.)

CVIII. ASTER Nees ast. p. 16. Lindl. adn. mss. in ast. Nees. — Asteris sp. Linn. Cass. Less. — Amellus Adans. non Linn.

Capitulum radiatum, fl. radii ligulatis fertilibus 1-serialibus, disci hermaphr. 5-dentatis. Recept. planum alveolatum, alveolorum marginibus plüs minüs dentatis. Involucri squamæ pluriseriales laxè vel imbricatæ apice plüs minüs herbacæ imò interdum foliaceæ. Achænum compressum. Pappus pilosus persistens pluriserialis, setis scabridis subinæqualibus cæterum inter se similibus. — Herbæ perennes pleræque ex Americâ boreali, rariùs ex orbe veteri aut ex Amer. austr. ortæ, interdum suffruticosæ aut scaposæ. Folia alterna simplicia integra aut dentata. Capitula solitaria aut plurima corymbosa seu paniculata. Discus flavus demum interdum purpurascens. Radius albus cæruleus purpureusve.

Ser. I. ALPIGENI Nees ast. p. 24. — Oritrophium sect. Asteris Kunth nov. gen. am. 4. p. 91.

Invol. squamæ herbacæ angustæ longitudine subæquales. Recept. nudum scrobiculatum. Pappus pauciserialis. Achænia sæpiùs hirta. Stigmata brevia triangularia. — Caules mono- aut oligocephali, sæpiùs hirsuti. Capitula majuscula.

1. A? HOLOSERRICUS (Forst. prod. p. 56. n. 296), foliis subtus albo-tomentosis, radicalibus oblongo-lanceolatis angustè serratis, caulinis linearibus integerrimis, caule 1-cephalo compresso, invol. squamis lanceolato-linearibus laxis. 2½ in Novâ-Zelandiâ ad rupes muscosas in Duskybay. Nees ast. p. 25. Rich. et Less. fl. nov. Zee. p. 248. Capitulum a poll. diam. Invol. squamæ non nihil

glutinosa et lanuginosa discum æquantes. Ligulæ pollicares lineâ angustiores purpureæ. Pappus ferrugineus.

2. *A. P. CORIACEUS* (Forst. prod. p. 56. n. 297), foliis radicalibus lato-lanceolatis sessilibus coriaceis integerrimis subplicatis subtus candido-pellitis, caule lanato 1-cephalo superne squamato, invol. squamis laxis lanceolatis longissimè acuminatis apice pilosis. $\frac{1}{2}$ in Novâ-Zeelandiâ. Spreng. syst. 3. p. 528. Nees ast. p. 26. n. 3. Valdè affinis *A. holosericeo* et cum eo subsectionem ferè formans ex Nees. An uterque ad *Euribiam* referendus?
3. *A. ALPINUS* (Linn. sp. 1226), caule 1-cephalo, foliis radicalibus spathulato-oblongis caulinisque integerrimis lanceolatis pilosis, invol. squamis lanceolatis laxis ciliatis dorso glabriusculis obtusis aut acutiusculis disco æqualibus, achæniis hirsutis. $\frac{1}{2}$ frequens in pratis apricis montium Pyrenæorum etiam Hispanorum, Jaraasi, Alpium omnium, Caucasi, Sibirie in pratis montis Karagiol prov. Karabagh (Szow. ex Fisch. et Mey.). Nees ast. p. 26. n. 4. Jacq. fl. austr. t. 88. Sturm deutschl. fl. io. Ligulæ cœruleæ lineam latæ obtusiusculæ 6-8 lin. longæ. Var. glabriusculus, pilosus aut hirsutus (*A. hirsutus* Host fl. austr. 2. p. 485). (v. v. et s.)

β. brachyglossus, humilis, foliis linearibus, radicalibus vix apice latioribus, ligulis brevissimis discum non superantibus. $\frac{1}{2}$ in Vallisæ Alpe excoelâ dictâ Fluve ad glacies æternas. Habitus ab *A. alpino* diversissimus, sed post iterata examina ad hanc speciem reduxi. (v. s. comm. à cl. Phil. Dumanant.)

γ. nivalis, foliis angustioribus magis canescentibus, capitulo minore, ligulis angustioribus, invol. squamis canescentibus brevibus obtusis apice purpurascens. $\frac{1}{2}$ in Sibiria orient. *A. nivalis* Adams in Web. et Mohr beytr. 1. p. 67. *A. pulchellus* quoramd. Conf. Nees ast. p. 27. *β.* excl. secundo Willd. syn.
4. *A. ARGENTENSIS*, caule 1-cephalo subnudo sparsè villosus, foliis radicalibus oblongis basi angustatis obtusiusculis integerrimis villosis, caulinis paucis oblongis acuminatis, invol. squamis biserialibus dorso lanuginosis acuminatis, exter. lanceolatis, int. linearibus, achæniis glabris. $\frac{1}{2}$ in Dahuriâ ad flum. Argunum legit cl. Razgildew. Habitus ferè Hieracii. Invol. purpurascens. Ligulæ lineares angustæ 9 lin. longæ, lineam latæ. Involucrum Erig. uniflori Linn. Ligulæ Asteris alpini. (v. s. comm. à cl. Turczaninow.)
5. *A. SILENEFOLIUS* (Turcz. l. in litt. 1834), glabriusculus, caule erecto simplici 1-cephalo apice subnudo, foliis infer. oblongo-linearibus basi attenuatis, super. linearibus attenuatis, omnibus integerrimis 1-nerviis, invol. squamis linearibus acutis æqualibus puberulis, ligulis linearibus disco duplò longioribus. $\frac{1}{2}$ in Sibiria orient. inter Ochotiam et Saculiam legit cl. Turczaninow. Herba semipedalis vix pilosiuscula. Variat ramis laxè racemosis ramulis 1-cephalis et ramis simplicibus necnon caulibus foliisve glabriusculis aut scabris. (v. s.)
6. *A. GAYANUS*, pube setulosâ adpressâ scabriusculus, caule erecto spice subcorymboso, ramis nudiusculis 1-cephalis, foliis oblongo-linearibus acutis sessilibus integerrimis, invol. squamis linearibus acutis disco æqualibus pauciserialibus dorso piloso-scabris, ligulis linearibus disco duplò longioribus. $\frac{1}{2}$ in excelis Andibus Chilensium legit indef. Cl. Gay. Habeo etiam ex itin. Neesi sine loci designatione. Ligulæ ex sicco videntur albæ aut pallidè purpurascens. Achænia juniora vix pilosiuscula. Variat ramis laxè racemosis ramulis 1-cephalis et ramis simplicibus necnon caulibus foliisve glabriusculis aut scabris. (v. s.)
7. *A. GRAMINIFOLIUS* (Pursh fl. am. sept. 2. p. 545), glaber, caule surperne ramoso, foliis angusto-linearibus enerviis integerrimis erectis ramulis nudiusculis 1-cephalis, invol. squamis laxis discum æquantibus. $\frac{1}{2}$ ad sinum Hudsonis (Pursh) in Canadâ (Lindl.) Nees ast. 277, sed ad *Asteres* alpidgenos nec ad *Tripolia* referendus ex Lindley adn. mss. Ligulæ pallidè purpureæ.
8. *A. REPENS* (H. B. et Kunth nov. gen. 4. p. 90. t. 334 bis), rhizomate repente, foliis radicalibus lanceolatis obtusis integerrimis reticulato-venosa coriacea

glabris, caulinis minutis, caule scapiformi erecto 1-cephalo, invol. squamis lanceolato-acuminatis laxè imbricatis. \mathcal{L} in frondosis circa urbem Loxa Novo-Granatensium. Nees ast. 30. Ligulæ albæ. Rhizoma lanuginosum. Habitus Bellidis.

9. *A. CROCIFOLIUS* (H. B. et Kunth nov. gen. am. 4. p. 89. t. 332. n. 1), foliis radicalibus angustè linearibus uninerviis integerrimis margine revolutis subtus cano-tomentosis coriaceis, caulinis linearibus multò minoribus, caule scapiformi 1-cephalo, invol. squamis laxè imbricatis lanceolato-linearibus lanato-ciliatis. \mathcal{L} in frondosis circa Loxam Novo-Granat. Nees ast. p. 31. Ligulæ albæ. Habitus Bellidis.
10. *A. PELMITUS* (H. B. et Kunth nov. gen. am. 4. p. 91), caule erecto simplici 1-cephalo basi lanato, foliis lineari-lanceolatis acutis integerrimis margine revolutis suprâ glabris subtus albido-lanatis radicalibus petiolatis, invol. lanatis squamis linearibus obtusis imbricatis. \mathcal{L} in montosis Andium Novo-Granatensium locis uliginosis. Nees ast. p. 31. Herba 6 pollic. Capitulum magn. Inulæ britannicæ. Ligulæ albæ. Achænia pubescentia. Pappus rufescens.
11. *A. PYGMÆUS* (Lindl. adn. mss.), caule villosio 1-cephalo, foliis ovato-oblongis obtusis integerrimis glabris, invol. hemisphærici squamis laxè villosissimis linearibus obtusis. \mathcal{L} ? in Americâ arcticâ. Caules biunciales. Capitula maxima. (Lindl.)
12. *A. CALIFORNICUS* (Less. in linnæâ 1831. p. 33), caule ascendente oligocephalo, foliis inferioribus spathulatis crenato-serratis margine pubescentibus, caulinis lingulatis integerrimis, invol. subæqualis squamis lineari subulatis hirsutis. \mathcal{L} in Californiâ. Nees ast. p. 33. Herba facie Bellidis. Ligulæ purpureæ.
13. *A. JAPONICUS* (Less. in Nees ast. p. 34), caule erecto 1-cephalo supernè nudo magisque piloso, foliis elliptico-acuminatis integerrimis pilosis multiplinerviis, invol. pauciserialis squamis lineari-acuminatis hirtis, exterioribus subsquarrosis. \mathcal{L} in Japoniâ. Inula dubia Thunb. fl. jap. 318. Rhizoma repens? Caulis teres striatus. Capitulum A. Sibirici. Anthera cæcudatæ. Ovaria compressa hirsuta. Ligulæ cœruleæ?
14. *A. UNALASCHKENSIS* (Less. in linnæâ 1831. p. 122), caule erecto 1-3-cephalo supernè sparse villosio folioso, foliis infimis obovatis obtusis, cæteris lanceolatis ovatove lanceolatis acuminatis subserratis pubescenti-scabris ciliatis, invol. subæqualis squamis lineari-lanceolatis herbaceis villosis. \mathcal{L} in insulâ Unalaschkâ (Less.) et in Amer. bor. montibus scopulosis (Lindl. mss.). Nees ast. p. 34. (v. s. ex comm. ab Acad. imp. Petrop.)
15. *A. FOLIACEUS* (Lindl. adn. mss.), caule simplici? racemoso, foliis oblongo-lanceolatis amplexicaulibus subserratis glabris, capitulis ramulos axillares terminantibus, foliis caulinis imò ramealibus superatis, invol. squamis patentibus foliaceis glabris. \mathcal{L} in Unalaschkâ (Fisch.). Species notabilis involucri squamis folia ramealia referentibus, pedunculo tomentoso. — in specim. meo etiam ex Unalaschkâ à Fischero misso, invol. squamæ erectæ nec patentis, ciliatæ nec omnino glabræ, folia basi obtusa, canles pilosiusculi, pilis hinc indè subseriatis. Fortè A. peregrinus Less. in linnæâ 1831. p. 123 huc excl. syn. referendus? (v. s.)
16. *A. PEREGRINUS* (Pursh fl. bor. am. 2. p. 556. non Less.), caule erecto sub-simplici glabro supernè hirtio summitate 1-3-floro, foliis sessilibus remotis oblongo-lanceolatis sensim attenuatis glabris, invol. squamis lanceolato-linearibus villosis acutissimis laxè longitudine disci, ligulis numerosis. \mathcal{L} ad Oras occid. Americæ borealis et in insulâ Unalaschkâ. Nees ast. 33. Ast. Tilesii Wikstr. in act. holm. 1822. p. 13. Pili caulis ex Less. in linnæâ 1831. p. 123? seriatiim sæpius trifariam dispositi. Pappus flosculis disci æqualis ex Nees, sed valdè dubium mihi videtur an stirps in Unalaschkâ crescens eadem ac Americanâ?
17. *A. ESPENBERGENSIS* (Nees ast. p. 36. n. 19), caule simplici vel subcorymboso hirsuto basi repente ascendente, foliis oblongo-lanceolatis acutis argutè mucronato-serratis subtus villosis, invol. squamis subæqualibus lanceolatis acumi-

nalis patulis villosis, achæniis elongatis. $\frac{1}{2}$ ad sinum Eleholtzii in prom. Es-
penberg et Kotzebue Sund legit cl. Chamisso. Aster salunginosus? Cham. in
Litt. Less. in linnæa 1831. p. 124. Capit. A. Alpini sed squamæ inv. magis
foliaceæ et patulæ. (v. s.)

β .? *dubius*, invol. squamis lanceolatis æqualibus erectis villosis. $\frac{1}{2}$ in Una-
laschkâ. A. peregrinus herb. Acad. imp. Petrop. sed non videtur A. pere-
grinus Pursh ob caulem hirsutum nec seriatiim pilosum. (v. s.)

18. A. *PYRÆNEUS* (DC. fl. fr. n. 3139), caule erecto hispido striato rarò simplici
1-cephalo sæpius apice ramoso corymboso, ramis 1-cephalis foliosis, foliis
lanceolatis semialexicaulibus acuminatis apice remotè serratis utrinquè
hispido-scabris, invol. squamis laxis lanceolato-linearibus acuminatis dorso
pubero-scabris. $\frac{1}{2}$ in Pyrenæis montibus (Tourn.), ad montem Esquierri
(Lapeyr.), nec certissimè in Sibiria. (Lindl.). Desf. cat. h. par. 102. Lapeyr.
abr. 514. Ast. Sibiricus Lam. dict. 1. p. 305 et pler. hort. bot. non Linn. Ast.
Sibiricus var. α et γ Nees ast. p. 37. ex syn. Ast. Pyrenæus, etc. Tourn. inst.
482. n. 3. Jam à tempore Tourn. in hort. bot. Paris. cultus. (v. v. c.)

19. A. *QUITENSI* (Willd. in Spreng. syst. 3. p. 533), caule sulcato villosulo dupli-
cato-racemoso oligocephalo, foliis cordato-oblongis amplexicaulibus basi api-
cæque denticulatis pubescenti-hirtis, invol. laxi æqualis squamis 3-serialibus
lineari-subulatis discum superantibus, achæniis pubescentibus. $\frac{1}{2}$ in regno.
Qaitenai Humb. et Bonpl. Nees ast. p. 38.

Ser. II. AMELLI Nees ast. 39.

Invol. squamæ ciliatæ squarrosæ, exteriores herbacæ, interiores totæ
vel apice saltem membranacæ coloratæ. Recept. alveolatum marginibus
lacero-dentatis. Achænia hirta rariùs glabra. Stigmata apice lanceolata. —
Caulis corymbosus aut racemosus rigens. Folia scabrida subintegra. Capi-
tula ampla.

20. A. *IRCUATIENSIS*, caule erecto tereti sparsè hirsuto 1-5-cephalo sub capitulis
tomentoso-hirsuto vix incrassato, foliis lanceolatis utrinquè acuminatis à me-
dio ad apicem serratis suprâ glabris subtùs puberulis, invol. squamis lineari-
lanceolatis subhirsutulis acutis nunc erectis nunc apice patente squarrosis
radio imò disco brevioribus, ligulis circ. 20, achæniis pilosis. $\frac{1}{2}$ in sabulosis
Dahuriæ ad ripam fl. Ircuti legit cl. Turczaninow. Aster conspicuus Turcz. in
Litt. 1830 non Fisch. Ligulæ purpureæ 4-5 lin. longæ, lineam latæ. Affinis
A. Richardsonii sed ex hujus descr. facilè distinctus videtur. (v. comm. à cl.
Turczaninow.)

21. A. *RICHARDSONII* (Spreng. syst. 3. p. 528. Nees ast. 30), caule erecto ramoso
1-2-cephalo sulcato villosulo-canescens sub capitulis incrassato tomentoso,
foliis latè oblongis remotè dentatis subtùs breviter pilosis, invol. squamis lineari-
lanceolatis squarroso-patulis villosis, interioribus radium subæquantibus, li-
gulis circ. 30 angustis, achæniis pilosis. — in Amer. borealis desertis à Point
Lake ad mare Polare (Richards.) et in Sibiria (Br. ex h. Pall.). Aster montanus
Brown in Frankl. it. p. 749. non Nutt. A. Ammanni Lindl. adn. mss. ipso doc.
in Litt. 1835. Folia sesquipedem longa. Radix repens. Ligulæ ex Nees albæ?

22. A. *SALSUGINOSUS* (Richards. ex Brown fl. reg. pol. in Frankl. it. p. 749.
app. ed. 2. p. 32), caulibus simplicibus 1- (seu oligo-) cephalis, foliis lineari-
obovatis acutis subintegerrimis venosis, invol. squamis laxè imbricatis lineari-
bus acutis disco ferè duplò brevioribus, achæniis hirtis. $\frac{1}{2}$ in regione sylvestri
inter 54 et 64° lat. bor. Americæ sept. Hook. in bot. mag. t. 2942. Nees ast. 29.
Ligulæ purpureæ lineares 6-8 lin. longæ sæpè emarginatæ. Pappus scaber sub-
fuscescens. Species invol. brevissimo præsertim distincta.

23. A. *BIFLORUS* (Michx. fl. bor. am. 2. p. 114), caule glabro sæpius dicephalo,
foliis lanceolatis serratis scabris, invol. squamis arctè imbricatis ovatis. $\frac{1}{2}$ in
Labradoriâ ad sinum Hudsonianum et in Pennsylvania montibus altissimis. Desf.

- biflorus Nees syn. p. 15. ast. p. 39. *Aster strictus* Pursh fl. 2. p. 556. Nutt. gen. am. 2. p. 157. Mey. labr. p. 62. Ligulæ pallide lilacinæ.
24. *A. surculosus* (Michx. fl. bor. am. 2. p. 113), caule simplici subpubescente laxo apice oligocephalo, foliis lineari-lanceolatis acuminatis setaceo-serrulatis margine scabris, infer. in petiolum attenuatis, super. vaginanti-amplexicaulibus, invol. squamis cuneiformi-spathulatis mucronatis apice patulis ramulisque pubescentibus. 2/ in sylvis Carolinæ septentr. (Michx.), in marginibus pratorum paludosorum in Tennessee et Virginiâ (Nutt.). Nees ast. 40. Radix stolonifera. Capit. *A. amelli*. Ligulæ violaceæ. Pappus rufescens.
- β. Foliis obovato-lanceolatis. — in pinetis humidis Carolinæ circa Puryburg. Ast. surculosus Ell. sketch 2. p. 352. Ast. liatroides Muhl. An à var. α distinctus?
25. *A. conspurcatus* (Lindl. adn. mss.), caule stricto apice corymboso, ramis erectis pubescentibus 1-cephalis aphyllis, foliis oblongis acutis basi angustatis grosse serratis pubescentibus ambitu scabris, invol. hemisphærici squamis squarrosis acuminatis, infer. minoribus. 2/ in Americâ arcticâ. *A. conspicuus* Lindl. adn. mss. 1833 non Fisch. nec Turcz. Ad *A. spectabilem* accedit, sed foliis latis ferè omnibus grosse serratis faciliè distinguendus. Radius cœruleus. (Lindl. mss.)
26. *A. spectabilis* (Ait. kew. 3. p. 209), caule stricto apice corymboso, ramis rigidis incurvis corymbosis, foliis oblongo-lanceolatis acutis scabris sessilibus inferioribus serratis, invol. squamis obtusis copiosissimis squarrosis ramulisque glanduloso-pubescentibus. 2/ in Amer. bor. (Ait.), in Novâ-Cæsareâ vulgaris (Nutt.). Nees ast. p. 41. Lindl. bot. reg. t. 1527. *A. elegans* Willd. sp. 3. p. 2042 ex parte et *A. spectabilis* Willd. sp. 3. p. 2048 excl. syn. ex Nees. An *A. speciosus* Horn. h. hafn. 2. p. 816? Ligulæ cœruleo-purpureæ. (v. v. c.)
- β. *bellidifolius* (Nutt. gen. am. 2. p. 157), foliis oblongo-obovatis serratis corymbo subsimplici ramulis 1-cephalis. 2/ in Novâ-Cæsareâ.
27. *A. RADULA* (Ait. kew. ed. 1. v. 3. p. 210. non Less.), caule angulato glabrisculo corymboso, ramis oligocephalis nudiusculis, foliis lanceolatis utrinque attenuatis mediò argutè serratis rugosis scaberrimis, invol. squamis plurifariam imbricatis oblongis acutiusculis apice patulis. 2/ in Novâ-Scotiâ (Ait.), montibus altioribus Novi-Eboraci et Pennsylvania (Pursh). Nees ast. p. 43 Lindl. pl. exs. Ligulæ 5-nerviæ ex Pursh albæ, ex Nees pallide lilacinæ. — Ast. nudiflorus Nutt. gen. 2. p. 157 fortè ex Nees non differt? (v. s.)
28. *A. nudiflorus* (Nutt. gen. am. 2. p. 157), caule simplici glabro, foliis sessilibus ovato-lanceolatis subacuminatis et argutè serratis, corymbo simplici oligocephalo, pedunculis pubescentibus nudis sub-1-cephalis, invol. hemisphærici disco brevioris squamis densè imbricatis lineari-oblongis ciliatis. 2/ in paludibus Novæ-Cæsareæ rarior. Capitula magna, ligulis pallide violaceis. Affinis *A. radula* et ab eo vix distinctus ex Nees, sed patria diversissima.
29. *A. LIMA* (Lindl. adn. mss.), caule scabro virgato sub-1-cephalo, foliis linearibus scaberrimis acutis ciliatis sensim in involucrium decrescentibus, invol. squamis linearibus rigidis acutis arcuè imbricatis basi albidis. 2/ in Mexico ex herb. Bentham. Radius purpureus. Habitus *Diplopappi* linearifolii sed pappus involucrium et achænium *Asteris Amelli*. An frustula descripta sint caules aut rami? (Lindl. mss.)
30. *A. TATARICUS* (Linn. f. suppl. 373), caule erecto striato hispido apice ramoso-corymboso, ramis subnudis, foliis lanceolatis seu ovali-lanceolatis basi attenuatis integerrimis apice sæpius grosse serratis acuminatis superne glabrisculis subtus hispidis summis linearibus integerrimis parvis, invol. squamis linearibus glabrisculis subæqualibus margine et apice purpureis. 2/ in Sibiria à Jeniseo fluvio ad Kamtchatkam (Gmel.), in montosis apricis Dahuriæ (Turcz.). *A. rhomboidens* Lindl. in adn. mss. 1833 et ad *A. tataricum* rel. anno 1835. Gmel. sib. 2. p. 186. t. 80. f. 1? *A. Sibiricus* β Nees ast. p. 37. An *Grindelia Sibirica* Link. enam. 2. p. 336? Ligulæ cœruleæ aut albæ? (v. a. comm. à cl Turcz.)

31. *A. SPATHULATUS* (Lindl. adn. mss.), caule simpliciter ramoso, foliis spathulatis obtusis integerrimis ramorum basi dilatatis amplexicaulibus, pedunculis subaphyllis foliis brevioribus, invol. squamis linearibus acutis erectis tomentosis. $\frac{1}{2}$ in Americâ arcticâ. Facies *Ast. pulchelli* (Lindl.)
32. *A. ASCENDENS* (Lindl. adn. mss.), caulibus ascendentibus simpliciter racemosis vel corymbosis, foliis oblongo-linearibus subaphyllatis elongatis integerrimis glabris margine ciliato-scabris, involucri glabri squamis numerosis acutis patulis. $\frac{1}{2}$ in Americâ arcticâ. (Lindl. mss.)
33. *A. SIBIRICUS* (Linn. sp. 1226), caule pubescente apice corymboso, ramis simplicibus 1-cephalis vel compositis apice incanis, foliis oblongis grosse serratis pubescentibus, superioribus sessilibus, involucri squamis linearibus acuminatis, omnibus imbricatis squarrosis. $\frac{1}{2}$ in Sibirîâ (Prescott.) *Aster inuloides* Fisch. nec Don. *A. lacerus* Lindl. adn. mss. 1833 et ad *A. Sibiricum* deductus 1835. — Gmel. sib. 2. p. 186. t. 80. f. 1? Species pulchra, invol. squamis interioribus purpureis squarrosis, radio verosimiliter albo. (Lindl. mss.)
34. *A. PRESCOTTII* (Lindl. adn. mss. 1835), caule simplici scabro ascendente 1-2-cephalo apice incano, foliis oblongo-lanceolatis argutè serratis basi angustatis scabris, superioribus sessilibus integerrimis, invol. squamis linearibus acuminatis, omnibus imbricatis squarrosis. $\frac{1}{2}$ in Dahuriâ prope Irkutsk (Prescott.). *Aster pauciflorus* Lindl. adn. mss. 1833 non Nutt. *Ast. biflorus* Presc. nec Michx. *Asteri lacero* proximus, foliis inflorescentiâ radioque purpureo diversus. (v. s. comm. à cl. Lindl.!)
35. *A. AMELLUS* (Linn. sp. 1226), caule erecto 1-cephalo aut sæpius simpliciter corymboso, foliis oblongo-lanceolatis acutis subserratis subtripplinerviis pubescenti-scabris, invol. squamis 4-5-seriatis oblongis patulis obtusis aut vix acutis apice membranaceis. $\frac{1}{2}$ in Europæ mediæ et australis Asiæque occid. siccis. Nees ast. p. 44. Adest ex Lindl. adn. mss. in herb. Linn. sub nominibus *A. Amelli* et *A. Sibirici* var. Variat 1 aut polycephalus, caule breviter scabro aut hispidio, foliis infer. obtusis aut mucronatis, etc. Præcipuæ varietates aut fortè species distinguendæ sunt:
- a. *communis*, foliis plerisque integerrimis obtusis cum mucrone, invol. squamis obtusis. $\frac{1}{2}$ *A. Amellus* Jacq. fl. austr. t. 435. DC. fl. fr. n. 3136. *A. amelloides* Roem. arch. 2. p. 298. Wender. ind. sem. h. marb. app. n. 3? Koch! et Ziz obs. *A. elegans* Nees syn. 20. (v. v. sp.)
 - β. *latifolius*, foliis inf. spathulatis obovatis obtusis, invol. squamis obtusis ciliatis. $\frac{1}{2}$ *A. Amellus* bot. reg. t. 340. *A. amelloides* Reichenb. fl. exc. 246. *A. pseudo-amellus* DC. fl. fr. suppl. n. 3136 in adn. (v. v. c.)
 - γ. *hispidus*, foliis infer. oblongis basi attenuatis cauleque patentim hispidis. — in Thuringiâ ex Wallroth. (v. s.)
 - δ. *Bessarabicus*, foliis oblongis basi attenuatis utrinquè cauleque breviter scaberrimis, invol. squamis subobtusis. — *Ast. amelloides* Bess.! enum. p. 33 et ic. 8. *Ast. Bessarabicus* Bernh. ex Reichenb. fl. exc. 246. (v. s.)
 - ε. *Ibericus*, foliis integerrimis acutis, invol. squamis interioribus vix coloratis. *Ast. Ibericus* Stev. in Bieb. fl. n. 1732.
36. *A. MODESTUS* (Lindl. adn. mss.), caule simplici folioso, corymbo paucifloro foliis supremis brevioribus, foliis lanceolatis acuminatis grosse serratis glabris, invol. squamis linearibus acuminatis. $\frac{1}{2}$ in Americâ arcticâ. *A. brumale* nomen refert sed affinitate ad *Ast. Novæ-Angliæ* proximè accedit. (Lindl. mss.)
37. *A. NOVÆ-ANGLIÆ* (Ait. kew. 3. p. 201. Nees ast. 46), caule stricto corymboso-composito, ramis arcuatis, foliis lanceolatis attenuatis auriculato-amplexantibus integerrimis hispidio-scabris, invol. squamis laxis angustis. $\frac{1}{2}$ in Amer. bor. ad rivulorum ripas et margines à Canadâ ad Carolinam. *Ast. altissimus* Mœnch. Ligulæ purpureo-violaceæ. *Achænia hirta*.
- a. *agrestis* (Nees l. c.), caule humiliori polycephalo, ramulis elongatis, invol. squamis lanceolatis venoso-trinerviis. $\frac{1}{2}$ *Ast. spurius* Willd. sp. 3. p. 2032. *Ast. amplexicaulis* Lam. dict. 1. p. 304. non Willd. nec Michx.

- A. Novæ-Angliæ Ell. sketch 2. p. 351. Lindl. ! pl. exs. A. grandiflorus hortul. (v. s.)
- β. *hortensis* (Nees 1. c.), caule altiori oligocephalo, ramulis brevibus ad apicem confertis, invol. squamis lineari-lanceolatis. ♀ Aster Novæ-Angliæ Linn. sp. 1229. Lindl. ! bot. reg. t. 183. — Asteripholis Pont. diss. 224. — Volk. fl. norib. p. 50. ic. Knorr del. 1. p. 5. 2. Herm. parad. Lugd. b. p. 98. ic. Variat in hortis caule purpureo subviscido, etc. (v. v. c.)
- γ. *roseus*, ligulis roseis. Ast. roseus Desf. ! cat. h. par. ed. 3. p. 401. non Stev. — bot. reg. t. 183. f. d. (v. v. c.)
38. A. CAROLINIANUS (Walt. car. 208), caule suffruticoso flexuoso ramosissimo pubescente, foliis oblongis acuminatis subintegerrimis in petiolum brevem auriculato-amplexicaulem contractis pubescenti-scabris, ramis apice simpliciter racemosis, invol. squamis apice reflexis, achæniis lævibus 10-striatis. ♀ in dumetis umbrosis et paludibus Carolinæ et Georgiæ. Nees ast. p. 48. Michx. fl. bor. am. 2. p. 111. Ell. sketch 2. p. 353. Ast. scandens Michx. ! herb. L'Hér. Jacq. f. in Spreng. syst. 3. p. 524. Ligulæ latæ purpureæ roseæ aut carneæ. Achænia pubescentia aut adulta glabra. Folia dicuntur integerrima, sed in spec. meo hinc indè acutè serrulata. (v. s. sine flor.)
39. A. OBLONGIFOLIUS (Nutt. ! gen. am. 2. p. 156), caule herbaceo scabro divaricato-ramoso, ramis subcorymbosis 1-cephalis, foliis lineari-oblongis subamplexicaulibus acutis integerrimis scabris, invol. squamis lineari-oblongis foliaceis squarrosis, achæniis brevè villosis. ♀ in Amer. bor. ad flum. Missouri Nees ast. p. 48. A. graveolens Nutt. ! in Journ. ac. nat. sc. philad. 1821. oct. p. 116. in linnæa 1829. p. 39. ! Ramuli fastigiati. Ligulæ pallidè purpureæ. Herba viscosa graveolens. (v. s. comm. à cl. Nutt.)
40. A. PATENS (Ait. kew. 3. p. 201), caule ramoso hirtio, ramis elongatis oligocephalis, foliis oblongo-ovatis integerrimis cordatis amplexicaulibus scabris, pedunculis 1-cephalis subnudis, invol. squamis imbricatis lanceolatis patulis. ♀ in Virginiâ et Pensylvaniâ Willd. sp. 3. p. 2034, in sylvis circa Bostoniam (Big.). A. patens α Nees ast. 49. excl. Michx. syn. Ast. amplexicaulis Michx. fl. 2. p. 114. Big. ! fl. hort. ed. 2. p. 312. Ast. amplexicaulis Michx. est omnino A. patens ex Lindl. A. undulatus Linn. sp. ed. 1. p. 1228 ex Sm. Ell. ! sketch 2. p. 361. Folia subobtusâ minutè mucronulata basi non angustata, auriculis obtusis. Pappus rufus. Achænia sericeo-villosa. (v. s. sp.)
41. A. PHLOGIFOLIUS (Muhl. in Willd. sp. 3. p. 2034), caule simplicissimo pubescente apice paniculato, foliis lanceolatis integerrimis utrinquè attenuatis infernè dilatatis cordato-amplexicaulibus subtus pubescentibus, invol. squamis laxis imbricatis lanceolatis, achæniis pilosiusculis. ♀ in Pensylvaniâ (Muhl.) à Novâ-Cæsareâ ad Virginiam (Pursh ?). Ast. patens β. Nees ast. p. 49. excl. syn. Michx. sed ab A. patente satis diversus mihi videtur ex foliis ad A. Carolinianum accedentibus, molliter pubescentibus nec scabris, achænio multò minùs villoso, capitulis majoribus. (v. s. et v. c.)
42. A. PATENTISSIMUS (Lindl. adn. mss.), caule racemoso-composito, ramis longissimis foliosis, foliis ovato-oblongis cordatis amplexicaulibus suprâ glabris subtus pilosiusculis, pedicellis brevibus, involucri turbinati lanati squamis obtusis. ♀ in Louisianâ. Radii cyanei. A. patenti proximus. (v. s. comm. à cl. Lindl.)
43. A. AURATUS (Lindl. ! adn. mss.), caule racemoso-composito, ramis rigidis apice densè racemosis, foliis oblongis acuminatis basi cordatis amplexicaulibus suprâ lævibus subtus pubescentibus, involucri parvum imbricati squamis linearibus acuminatis ramulisque glandulosis. ♀ in Americâ sept. ? Cultus in hort. anglicis sed rariùs. Radius discolor. Species A. patentissimo proxima. (v. s. comm. à cl. Lindl.)
44. A. GRANDIFLORUS (Linn. sp. 1231), caule erecto hispido racemoso, ramis elongatis 1-cephalis foliosis, foliis lanceolato-lingulatis mucronatis integerrimis hispidis reflexis, invol. squamis lineari-lanceolatis squarrosis, achæniis striguloso-hirtis. ♀ in sylvis siccis sabulosis à Virginiâ ad Carolinam (Pursh) bot.

reg. t. 273. Hoffm. phyt. t. 65. t. A. f. 1. Nees ast. p. 50. — Mill. ic. t. 282. Dill. elth. p. 41. t. 36. f. 41. Capitula ampla ligulis violaceis. (v. v. c.)

45. *A. sericeus* (Vent. h. Cels. n. et t. 33), basi suffruticulosus totus pube sericeâ adpressâ argenteus, ramis laxis apice 1-cephalis, foliis lanceolatis integerrimis, invol. squamis foliaceis imbricatis squarrosis argenteis, achæniis glabris. 2/ 3 ad ripas fluminum præruptas inter rupes in prov. Illinoensi et ad flum. Mississippi et Missouri. Nees ast. p. 51. Lindl. ! pl. exs. *Aster argenteus* Michx. fl. bor. am. 2. p. 111. Ligulæ cœruleæ. (v. v. c.)

β. *microphyllus*, foliis capitulisque triplò minoribus. 3 in Mexici distr. orientalibus prov. Texas legit cl. Berlandier (pl. exs. n. 1876). An species propria? (v. s.)

46. *A. montanus* (Nutt. gen. am. 2. p. 156), caule erecto gracili, ramis laxis 1-cephalis, foliis lineari-lanceolatis integerrimis subsericeo-villosis oboletè 3-nerviis, invol. hemisphærici squamis imbricatis lineari-lanceolatis erectis. — in montibus distr. Tennesæe et Carolinæ septentr. prope Ashville et Morganton rarior (Nutt.). Nees ast. p. 51. Valdè accedit ad *A. sericeum*. Ligulæ 2-5 violaceo-purpureæ.

Ser. III. GENUINI *Nees ast. p. 52.*

Invol. squamæ imbricatæ vel laxæ, margine magis minùsve membranaceæ, membranaceâ parte basin versus incremente et totam ferè basin in iis quæ fortius sibi incumbunt occupante. Recept. alveolato-dentatum. Achænia glabra vel puberula. Pappus pluriserialis. — Caulis altior compositè corymbosus racemosus paniculatusve.

A. Heterophylli *Nees ast. p. 52.*

Folia radicalia seu inferiora petiolata cordata vel rariùs rotundata.

47. *A. urophyllus* (Lindl. adn. mss.), caule racemoso-paniculato, ramis thytsoideis, foliis ovato-lanceolatis acuminatissimis argutè crenato-serratis suprâ scaberrimis subtùs pilosis, invol. imbricati squamis subulatis. 2/ in Louisianâ. Radii albi involucri longiores. Discus purpureus. Species *A. hirtello* proxima. (Lindl. mss.)
48. *A. hirtellus* (Lindl. adn. mss.), caule racemoso-paniculato, ramis apice conferto-racemosis paucifoliis, foliis cordato-ovatis mediò crenato-serratis suprâ scaberrimis subtùs pilosis, involucri laxè imbricato. 2/ in Louisianâ. Radius verosimiliter lilacinus, discus purpureus. (Lindl. mss.)
49. *A. cordifolius* (Linn. sp. 1229. non Lam. nec Nutt.), caule erecto racemoso-paniculato subhirsuto, foliis petiolatis cordatis acuminatis argutè serratis suprâ scabris subtùs pilosis, ramis laxis paucifoliis, invol. squamis arcuè imbricatis. 2/ in sylvis umbrosis montium à Canadâ ad Floridam. Nees ! ast. p. 52. Lindl. ! bot. reg. 1597. *Ast. cordifolius* var. foliis subtùs pubescentibus Desf. ! in hort. Par. 1815. *Solidago cordifolia* Moench. Caulis rubens nunc ferè glaber nunc subhirsutus. Folia etiam superiora cordata, petiolis subhirsutis. Ligulæ pallidè purpurascentes aut albidæ. Achænia glabra seu ad costas setulosa. (v. v. c. et s. sp.)
50. *A. paniculatus* (Ait. kew. ed. 1. v. 3. p. 205), caule glabro stricto ramisque thyrsoideis foliosis, foliis petiolatis infer. cordatis summis, ovato-lanceolatis, omnibus acuminatis serratis suprâ scabris subtùs glabris, invol. squamis imbricatis, 2/ in pratis, sepibus et sylvarum marginibus à Canadâ ad Carolinam. Nees ast. p. 53. Ell. sketch 2. p. 363. Lindl. ! pl. exs. *Ast. cordifolius* Lam. dict. 1. p. 307. *A. cordifolius* var. foliis subtùs glabris Desf. ! hort. Par. *Ast. heterophyllus* Nees syn. p. 15. non Willd. — Moris. ox. 3. s. 7. t. 22. f. 34. Ligulæ violacæ. (v. v. c.)
51. *A. heterophyllus* (Willd. sp. 3. p. 55), caule racemoso-decomposito paniculato, foliis radicalibus caulinisque infer. cordatis alate petiolatis serratis apice

integerrimis suprà in ambitu totiusve scabris subtus glabris, supremis lanceolato-linearibus minutis, invol. arcè imbricato. $\frac{1}{2}$ in fruticetis Pensylvaniæ Nees! ast. p. 55. Lindl. ! pl. exs. Ast. cordifolius Nutt. gen. am. 2. p. 156. non Linn. A. diversifolius Spreng. syst. 3. p. 533. non Michx. (v. v. c.)

β . *glabriusculus*, caule piloso-lineato, foliis glabris vel subpubescentibus. — A. pubescens Horu. h. hafn. suppl. 98. Nees! syn. p. 16. (v. v. c.)

γ . *scaber*, caule hirsuto, foliis suprà scaberrimis, capitulis majoribus. An ex Nees species propria?

52. A. *SAGITTIFOLIUS* (Wedem. in Willd. sp. 3. p. 2065), caule glabro racemoso-composito, foliis radicalibus petiolatis cordato-subasagittatis (vel et profunde cordatis) oblongis glabris inæqualiter serratis margine scabris, caulinis ovato-oblongis in petiolum alatum contractis serratis glabris, involucre arcè imbricato. $\frac{1}{2}$ in Americæ bor. (Wed.), distr. superioribus utriusque Carolinæ (Ell. sketch 2. p. 362). Ligulæ cœruleæ (Willd.), lilacinæ (Nees ast. p. 56).

53. A. *DAUMMONDII* (Lindl. adn. mss.), totus incanus, caule ramisque racemoso-thyrsoideis strictis, foliis cordato-ovatis crenato-serratis, supremis sessilibus, ramulorum lineari-oblongis acuminatis, invol. squamis subulatis. $\frac{1}{2}$ in Louisiana. Inter A. paniculatum et undulatum quasi medius, canitie facillimè distinguendus. Radius verosimiliter cyanei. (Lindl. mss.)

54. A. *DIVERSIFOLIUS* (Michx. fl. bor. am. 2. p. 113), caule puberulo racemoso-thyrsoideo, ramulis virgatis apice secundifloris, foliis radicalibus longè petiolatis cordatis crenato-serratis acutis pubescentibus margine scabris, petiolo dilatato basi amplexante, superioribus sessilibus ovato-lanceolatis ramulorum subulatis, invol. arcè imbricato. $\frac{1}{2}$ in Amer. bor. à Canadâ ad Carolinam, in Novâ-Angliâ (Pursh), in siccis Carolinæ frequens (Ell. sketch 2. p. 361), in Canadâ (Michx.), circa Bostoniam (Big. fl. bost. ed. 2. p. 312. Lindl. adn. mss. A. undulatus Linn. sp. ed. 2. p. 1228. Nees ast. p. 57. A. scaber Ell. sketch 2. p. 363. non Thunb. Ligulæ cœruleæ. (v. s. sp.)

B. Homophylli Nees ast. p. 58.

Folia radicalia basi acuta magis minùsve attenuata.

* Petiolati Nees ast. p. 58.

Folia latiuscula serrata, radicalia et caulina inferiora in petiolum attenuata, summa basi angustiore sessilia.

55. A. *PATULUS* (Lam. diet. 1. p. 308), caule glabro racemoso-paniculato, ramis patulis, foliis oblongis profunde serratis in petiolum attenuatis suprà pube minutissimâ aspersis glabrisve subtus glabris margine scabris, invol. laxè imbricato. $\frac{1}{2}$ in pratis sepius sylvarumque marginibus à Canadâ ad Virginiam usquè. Desf. ! cat. h. par. 102. A. paniculatus Willd. sp. 3. p. 2035, ex parte Nees ast. p. 16. excl. syn. A. Cornuti Wendl. in Nees ast. p. 58. nec Mull. A. Tradescanti Hofm. phyt. bl. p. 68. t. d. f. 2. ex Nees. Ligulæ cœruleæ. Nomen multò vetustius, nec recentius et verisim. decipiens, admissi. (v. v. c.)

β . Radio pallido, foliis glabrescentibus. — A. pallens Willd. enum. suppl. p. 58. Lindl. bot. reg. 1509. A. tardiflorus Linn. 1. p. 1231, hæc aut ad var. β pertinet ex cl. Lindley.

56. A. *ABBREVIATUS* (Nees! syn. p. 16), caule erecto glabro seu seriatim piloso racemoso, ramis brevibus thyrsoideis simplicibusve, foliis infer. ovali-lanceolatis serratis secus petiolum adnato-decurrentibus suprà scabris subfús lævibus, sup. oblongo-lanceolatis subintegerrimis, invol. squamis laxè imbricatis lanceolato-linearibus. $\frac{1}{2}$ in Amer. bor. sed ex confus. syn. locos proprios non citare audeo. A. Cornuti Mull. ex Nees. A. acuminatus Nees Ast. p. 60. excl. syn. Variat ramis abbreviatis aut in plantâ cultâ elongatis. Ligulæ pallidè cœruleæ. (v. v. c. à cl. Nees ad hort. Genev. miss.)

57. A. *PARNATHOIDES* (Muhl. in Willd. sp. 3. p. 2046), caule paniculato, ramis

brevibus corymbiferis, foliis ovali-lanceolatis serratis basi in petiolum auriculato-amplexicaulem attenuatis suprâ scabris subtus lævibus, invol. squamis linearibus apice squaroso-patulis. \mathcal{L} in Pennsylvania paludibus. Nees ast. p. 61. nec syn. nec hortul. Ligulæ cœrulesc. (Willd.) lilacinæ (Nees). Ramuli bifariam pilosi (Willd.). Pappus albidus. Species in hortis sub hoc nom. frequens est A. Novi-Belgii ex Nees.

58. A. PRÆCOX (Willd. enum. suppl. p. 58), caule racemoso, ramis oligocephalis basi nudis, foliis ovato-lanceolatis patenti-serratis in petiolum adnatum attenuatis glabris suprâ in ambitu scabris basin versus ciliatis, radicalibus ovatis petiolatis, involucri laxo subæquali. \mathcal{L} in Amer. septentr. Nees ast. p. 62. Invol. squamæ apice purpurascens lanceolata. Achænia pubescentia. Herba vix sesquipedalis.

59. A. CILIOLATUS (Lindl. adn. mss.), caule simplici, foliis omnibus ovatis nudis argutè serratis in petiolum abruptè angustatis ciliatis in ambitu scabris capitulis axillaribus sessilibus vel brevi pedicellatis subsolitariis, invol. squamis erectis apice membranaceis. \mathcal{L} in Americâ arcticâ. Valdè affinis A. præcoci, differt foliis abruptè nec sensim angustatis caulinisque radicalibus conformibus conspicuè per totum ambitum ciliatis, capitulis subsessilibus dimidiò minoribus, invol. squamis non apice herbaceis, deniquè staturâ spithamæ. (Lindl. mss.)

60. A. HÆMEX (Nees ast. p. 63), caule hirsuto simpliciter racemoso oligocephalo, foliis inferioribus ovato-ellipticis petiolatis, superioribus oblongis, omnibus serratis suprâ hispidè-scabris subtus hirtis, invol. squamis imbricatis oblongo-lanceolatis acutiusculis, radio brevi. \mathcal{L} in regione montanâ Chilensî. Habitus ferè A. Radulæ.

** Amelloidei Nees ast. 65.

Folia sessilia, inf. basi attenuata, sup. basi latiora. Invol. squamæ erecto-patula læx acuminatæ vel lineari-spathulatæ, int. sæpè coloratæ, ext. foliaceæ. Recept. alveolato-dentatum. Pappus mollisculus. Caulis racemosus vel racemoso-decompositus. Folia lanceolata vel lineari-lanceolata serrata.

61. A. THYRSIFLORUS (Hoffm. phyt. bl. 1. p. 83. t. d. f. 1. ex Nees ast. p. 65), caule racemoso-composito patulo, ramis elongatis apice spicato-racemosis, foliis lineari-lanceolatis attenuatis amplectentibus margine scabris angustè serratis, invol. laxi squamis lanceolatis inæqualibus recurvis. \mathcal{L} in Virginiâ? A. recurvatus Willd. sp. 3. p. 2047. ex parte. Ligulæ lilacinæ. Achænia glabra.

β . *squarrosus* (Lindl. ! adn. mss.), foliis omnibus linearibus squamoso-recurvis, ramis apice racemosis, capitulis majoribus. \mathcal{L} in hortis Angliæ cultus. (v. s. comm. à cl. Lindl.)

62. A. FIRMUS (Nees syn. 25. ast. 66), caule stricto anguloso infernè glaberrimo racemoso-composito, ramis apice oligocephalis coarctatis, foliis lanceolatis acuminatis amplectentibus in ambitu scabris medio serratis, invol. laxi squamis subæqualibus lanceolato-subulatis. \mathcal{L} patr. ign. verisim. Amer. bor. flor. aug. 7^{bris}. Humilis strictus. Folia glabrescens. Ramis apice sparsè seu lineatim pilosi. Ligulæ cœrulescenti-lilacinæ. Achænia sparsè puberula. A. puniceus Ell. sketch 2. p. 255. descr. nec diagn. hæc pertinere videtur et ideo stirps in Carolinâ et Georgiâ esset indigena. Sæpè sub nom. A. punicea et ameni in hortis vagat. (v. v. c.)

63. A. CORYMBIGENS, caule erecto lævi, ramis erectis foliosis racemosis, foliis sessilibus ferè semiamplexicaulibus lato-linearibus lingulatisve acutis subcoriaceis integerrimis lævisculis margine et costâ subtus scabris ramulorum r-cep-phalorum sensim in squamas inv. abeuntibus, invol. squamis oblongis acutis læx imbricatis subsquarrosis. \mathcal{L} in Mexici districtibus orientalibus provinciæ Texas legit cl. Berlandier (pl. exs. n. 1885). Folia caul. 3-4 poll. longa 4 lin. lata. Capitulum magn. A. punicea. Ligulæ pallidè cœrulesc. Achænia glabrescens. Pappus rufus. (v. s. comm. à cl. inv.)

64. *A. PUNICEUS* (Linn. sp. 1229), caule hispido patulo racemoso, ramis infer. racemosis, super. simplicibus, foliis oblongo-lanceolatis amplectentibus acuminatis medio adpresso-serratis suprâ scabris invol. laxi squamis lineari-subulatis æqualibus. $\frac{1}{2}$ in Amer. bor. à Canadâ ad Virginiam. Nees ast. p. 67. *A. hispidus* et *A. amœnus* Lam. dict. 1. p. 306 et 307. — Herm. h. Lugd. b. p. 651. ic. Ligulæ purpureo-violaceæ. Achænia sparsè puberula. Caules purpurei seu rufescentes 4-6-pedales. (v. v. c. et s. sp.)

β. ramis brevibus. (Nees l. c.)

γ. *demisus* (Lindl. bot. cab. t. 1636), caule sesquipedali corymboso, ligulis cœruleo-violaceis.

65. *A. VIMINEUS* (Willd. sp. 3. p. 2046 non Lam.), caule lævi racemoso-composito pyramidalis, ramis virgatis apice racemosis, foliis lanceolatis amplectentibus longè acuminatis in medio subulato-serratis suprâ et margine scabris, invol. laxi squamis lanceolato-linearibus subæqualibus. $\frac{1}{2}$ in Canadâ. Nees ast. p. 68. Spreng. syst. 3. p. 538. excl. syn. Ligulæ pallide lilacinæ. Achænia pubescentia. Sæpè vagat in hortis sub nom. *A. Tradescanti*. Flor. autumn. (v. v. c.)

β. *glaber*, caule minùs diviso, ramis brevibus, foliis in disco lævibus. — *A. glaber* Meyer ex Nees.

66. *A. ARGUTUS* (Nees ast. p. 69), caule lineis pilosis notato recurvato-racemoso-composito, ramis virgatis apice floriferis, foliis lanceolatis basi apiceque attenuatis amplectentibus medio argutè serratis suprâ in ambitu scabris, invol. imbricati squamis subulatis apice recurvis. $\frac{1}{2}$ patr. ign. verisim Amer. bor. Ligulæ cœruleæ. Achænia sparsim pubescentia.

67. *A. LAXIFOLIUS* (Nees ast. p. 69), caule scabro racemoso-composito angusto, ramis apice racemosis vel subcompositis, foliis angustè lanceolatis utrinquè attenuatis mucronato-serrulatis flaccidis suprâ scabris, invol. squamis linearibus squarrosis, capituli primarii subæqualibus. $\frac{1}{2}$ in Americâ boreali. *Aster longifolius* Nees! syn. p. 26. non Lam. Ligulæ incarnatæ angustæ. Achænia pubescentia (Nees). Omnium vulgatissimus in Amer. arct. Variat foliis serrulatis et integerrimis, caule 1-cephalo, oligocephalo, corymboso et composito-racemoso. Pappus rufescit (Lindl. mss.). (v. s.)

β. *salicifolius* (Lindl. adn. mss.), caule gracili angustissimo, foliis acuminatis, capitulis minoribus.

γ. *carneus* (Lindl. adn. mss.), caule piloso-lineato, invol. ovato, ligulis carneis. — *A. carneus* Nees ast. p. 96. ex Lindl.

68. *A. AMPLUS* (Lindl. adn. mss.), caule simplici subcorymboso, ramis 1-cephalis subaphyllis, foliis oblongo-lanceolatis margine scabris subintegris, involucre laxo imbricato. $\frac{1}{2}$ in Americâ arcticâ. *A. brumali* affinis. Foliis capitulisque multò majoribus, caule simplici subcorymboso ramisque rigidis pilosis ferè aphyllis nec virgatis glabris foliosis benè distinguitur. (Lindl. mss.)

69. *A. BRUMALIS* (Nees ast. p. 70), caule glabro racemoso, ramis sub-1-cephalis, foliis lanceolatis acuminatis amplectentibus margine scabris medio argutè adpresso-serratis, invol. laxo subæquali. $\frac{1}{2}$ in Amer. septentr. et rariùs in Germaniâ quasi spontè nascens. *Aster Novi-Belgii* Willd. sp. 3. p. 2048. Nees syn. p. 24. non Linn. *Aster lævigatus* Lam. dict. 1. p. 307. *A. serotinus* Nutt. gen. am. 2. p. 158? Ligulæ cœruleæ. Achænia pubescentia. Variat ramis 1-cephalis et racemosis ramulis 1-cephalis. (v. v. c.)

70. *A. FRÆULTUS* (Poir. suppl. 1. p. 493), caule basi glabrato racemoso-composito, ramis virgatis coarctatis summis fastigiatis, foliis lineari-lanceolatis attenuatis subamplectentibus integerrimis margine scabris, infer. obsolete subserratis, ramulorum linearibus recurvis, invol. laxè squamis lanceolatis inæqualibus apice sæpè recurvis. $\frac{1}{2}$ in Amer. bor. ad fossarum margines à Noveboraco ad Virginiam. *A. salicifolius* Ait. kew. ed. 1. v. 3. p. 203. *A. virgatus* Nees ast. p. 27. non Moench. An Ell. sketch 2. p. 353. — Rob. icon. t. 307. Ligulæ cœruleæ. Achænia pubescenti-scabra. Caulis supernè pilis lineatus. (v. s.)

- ♂. caule toto glabro foliis saturatius viridi-lucidis infer. oblongo-lanceolatis, radio disco paulò longiore. — In solo pingui.
71. *A. PESTHINENSIS*, caule piloso-lineato racemoso virgato, ramis sub-1-cephalis, foliis lanceolatis utrinque modicè angustatis acutis subintegerrimis margine scabris, ramorum ramulorumque conformibus, invol. laxi squamis linearibus subæqualibus. ♀ in Hungariâ circa Pesthinum. *A. riparius* Nees ast. p. 73. nec p. 280. Ligulæ purpurascens? *Achænia hirta*. Species dubia ex Nees.
72. *A. MULTICEPS* (Lindl. adn. mss.), undiquè glanduloso-pubescent, caule racemoso, ramis erectis 1-cephalis, foliis oblongo-linearibus acutis pubescentibus scabriusculis, invol. squamis linearibus subulatis basi albidis squarrosis. ♀ in Louisianâ. Species distinctissima nulli cognitæ arctè affinis. Amelli sectionem diceret si squamæ invol. interiores membranaceæ et coloratæ essent. (Lindl. mss.)

** Versiformes Nees ast. p. 73.

Folia sessilia, lanceolata vel oblongo-lanceolata inf. basi attenuata sup. basi latiora. Invol. squamæ erecto-patulae laxæ erectæ vel recurvæ. Capitula medio-cria, radio pallidè cœruleo vel albo. — Sectio vix à præcedente distincta.

73. *A. æSTIVUS* (Ait. kew. ed. 1. v. 3. p. 203), caule paniculato-ramoso vel racemoso, ramulis sparsis laxis 1-cephalis, foliis lanceolatis ciliatis, radicalibus adpresso-serratis, caulinis integerrimis, invol. angusti obconici squamis interioribus subulatis. ♀ in Amer. bor. uliginosis siccis sylvis cœdens à Noveboraco ad Pensylvaniam. Pursh fl. bor. am. 2. p. 549. Nees ast. p. 74. Lindl. ! pl. exs. Ligulæ albidæ. *Achænia puberula* basi attenuata. In specim. vivo à cl. Nees ad hort. Genev. misso caulibus glaber ut vult cl. auct. sed rami lineato-hirsuti ut innuit Willd. sp. 3. p. 2030. (v. v. c.)
74. *A. SUBSPICATUS* (Nees ast. p. 74), caule piloso-lineato, ramis virgatis ramulis hirsutis subspicatis, foliis oblongo-lanceolatis acutis adpresso-serratis glabris margine scabris amplexicanlibus, invol. subæqualis laxi squamis exter. spathulato-lanceolatis latioribus. ♀ in promontorio Mullgrave ad oram bor. occid. Americæ sept. An eadem ac *A. blandus* Pursh? Ligulæ purpurascens. *Achænia* densè incano-strigulosum.
75. *A. BLANDUS* (Pursh fl. bor. am. 2. p. 555), caule pyramidato-ramoso, ramis axillaribus racemiferis folio vix longioribus, foliis subamplexicanlibus oblongo-lanceolatis acuminatis serratis glabris, pedunculis tomentososis nudis, involucriis laxis subæqualibus disco brevioribus. ♀ in Canadâ ex herb. Banks. Nees ast. p. 76. Ligulæ pallidè purpureæ.
76. *A. OBLIQUUS* (Nees syn. 1818. 27. ast. p. 76), caule infernè glabro supernè racemoso-composito stricto, ramis apice subcorymbosis, foliis lineari-lanceolatis sessilibus mucronatis subintegerrimis suprâ scabris obliquis, ramulorum patentibus, invol. laxi squamis exter. majoribus spathulato-lanceolatis patulis. ♀ in Amer. bor. ad Noveboracum. *A. obliquus* Lindl. pl. exs. *A. junceus* 2 *A. lanceolatus* Nutt. gen. 2. p. 158. *A. rigidulus* Desf. cat. h. par. 1815. p. 122. An *A. lanceolatus* auct. americæ? Ligulæ lilacinæ et var. albæ. (v. v. c.)
- ♂. caule humiliore oligocephalo corymboso, pappo rufescente Nees. l. c.
77. *A. SUBASPER* (Lindl. adn. mss.), caule infernè glabro supernè pubescente racemoso-composito, capitulis ramorum secundis, foliis angustio-lanceolatis subsessilibus acuminatis uniformibus suprâ per totam superficiem scabris, superioribus gradatim minoribus, involucri hemisphærici squamis linearibus acutis disco conspicuè brevioribus, exterioribus laxis. ♀ in Louisianâ. Ab *A. obliquo* distinctissimus foliis latioribus uniformibus semper ut videtur integerrimis, capitulisque majoribus. Ast. tenuifolium etiam mentitur. (Lindl. mss.)
78. *A. HIRMALIS* (Nees ast. p. 77), caule glabro laxo racemoso-composito, foliis lineari-lanceolatis amplexicantibus acutis integerrimis margine scabris, ramulis foliaceis, invol. laxi squamis lanceolatis subæqualibus pluriseriatis ~~discreta~~

æquantibus ciliatis. \mathcal{U} patr. dub. verisim Amer. bor. In hortis ultimus florens. A salicifolius Nees syn. 26. nec alior. A. lævis Vahl herb. ex h. Puer! — Dill. elth. 1. p. 39. t. 35. f. 39. Ligulæ albæ subviolacæ. Achænia pubescenti-scabra. (v. s.)

79. A. TARDIFLORUS (Nees! ast. 78 non Linn. ex Lindl.), caule glabro corymboso-divaricato, foliis oblongo-lanceolatis oblique amplexentibus apicem versus serratis circa margines scabris ramulorum obtusis, invol. squamis imbricatis squarroso-patulis crassis, infer. sæpè elongatis subradiantibus. \mathcal{U} in Amer. bor. locis depressis inundatis à Noveboraco ad Virginiam (Pursh.). Ligulæ lilacinæ. Achænia glabra. — Ab hoc, monente cl. Lindley in litt. et confirmante specim. authent. à cl. Neesio comm. non differt Symphyotrichum unctuosum Nees! ast. p. 135. Species insignis ob pappi setas in anulum solubilem subcoactas. (v. v. et s. c.)

β . *cæspitosus* (Lindl. adn. mss.), caule humiliore, ramis densè corymbosis, foliis ramorum caulinis parum minoribus. \mathcal{U} cult. in hort. Angl. Aster adulterinus Lindl. bot. reg. t. 1571 non Nees. Symphyotrichum unctuosum Nees ast. p. 135. ex. Lindl. Achænia paucissimas setulas sub pappo gerentia. Pappus in anulum solubilem basi cohærens ex Nees.

80. A. NOVI-BELGII (Nees! ast. p. 79), caule corymboso-composito vel suprâ decomposito, ramis pilosulis rigidis, foliis lanceolatis amplexentibus acutis circa margines scabridis, inferioribus medio serratis, invol. laxi squamis intimis acuminatis. \mathcal{U} in Amer. bor. sponte et hinc inde in Europâ ex hortis emigratus.

α . *ampliflorus* (Nees l. c.), ramis simpliciter racemosis subcorymbosis, foliis oblongo-lanceolatis lanceolatisve, capitulis amplioribus, ligulis densis lætioribus, invol. squamis laxis erectis sæpè æqualibus lanceolatis, omnibus acutis. — A. Novi-Belgii Linn. sp. 1231. A. serotinus Nees syn. 21. A. floribundus Nutt. gen. 2. p. 158. — A. Novæ-Belgiæ, etc. Herm. h. Lugd. b. p. 69. ic. (v. v. c.)

β . *squarrosus* (Nees l. c.), invol. squamis exterior. squarroso-patulis sæpè foliaceo-elongatis. — Cætera ut in var. α .

γ . *serus* (Nees l. c.), caule exaltato, ligulis densis carneis discum versus albidis.

δ . *floribundus*, caule humiliori, foliis lanceolatis attenuatis, ramis confertis dichotomo-corymbosis polycephalis, radio minùs denso breviori, invol. laxè imbricatis squamis inæqualibus. \mathcal{U} A. floribundus Willd. sp. 3. p. 2048. Nees! syn. 21. A. Novi-Belgii minor Nees ast. p. 81. (v. v. c.)

81. A. LUXURIANS (Nees ast. 83), caule paniculato, ramis apice corymboso-racemosis, ramulis 1-cephalis, foliis ovali-oblongis vel lanceolatis acuminatis basi attenuatè amplexentibus, invol. squamis laxè imbricatis lineari-lanceolatis acutis. \mathcal{U} in Amer. bor. sepibus et agris neglectis à Novâ Angliâ ad Virginiam (Nees) A. Novi-Belgii Nutt. gen. 2. p. 158. A. prenanthoides Nees! syn. p. 23. (v. v. c.)

β . caule magis composito magisque piloso-lineato. A. luxuriana Spreng. syst. 3. p. 538. excl. syn. ex Nees.

82. A. ASPER (Nees ast. p. 84), caule supernè paniculato-composito glabro tuberculis aspero, ramis racemoso-corymbosis, foliis oblongo-lanceolatis acuminatis amplexentibus suprâ scabris, omnibus in medio serratis, invol. imbricati squamis apice squarroso-patulis. \mathcal{U} patr. ign. verisim. Amer. bor. Ligulæ ferè duplici serie pallidè cœrulesc. seu lilacinæ. Achænia obsoletè puberula.

83. A. ADULTERINUS (Willd. enum. 2. p. 884), caule à basi paniculato-decomposito, ramis corymboso-racemosis, foliis oblongo-lanceolatis mucronatis amplexentibus lævibus margine scabris, invol. obovati subæqualis squamis spatulato-lanceolatis squarrosis post florescentiam magis acutis. \mathcal{U} in Americâ septentr. Nees! syn. 22. ast. 85. Ligulæ albæ, demum pallidè violacæ. Achænia glabra. An var. A. Novi-Belgii ex Nees? (v. v. c.)

84. *A. squarrosulus* (Nees ast. p. 86), caule corymboso-decomposito, ramis apice corymbosis erectis, foliis lanceolato-acuminatis subamplectentibus medio argutè serratis in ambitu supernè scabris, invol. ovati squamis angustis imbricatis apice recurvis. \mathcal{U} in Americà boreali. *A. spectabilis* Willd. sp. 3. p. 2048. Nees syn. 21. *A. mutabilis* Linn. sp. 1280? ex Nees. *A. recurvatus* Spreng. in Schrad. Journ. bot. 1800. p. 195.

β . *albiflorus*, elatior, radio albo dein subviolaceo, disco mox saturatè purpureo. Nees l. c.

85. *A. lowerrolii* (Lam. ! dict. 1. 1782. p. 306), caule glabro paniculato patulo, ramis apice simpliciter corymbosis, foliis lanceolato-acuminatis subamplectentibus, infer. in medio argutè serratis, suprà in ambitu latè scabris, invol. turbinati subæqualis squamis lineari-lanceolatis patulis. \mathcal{U} in Amer. bor. paludosis à Noveboraco ad Carolinam. *A. juncens* Ait. kew. 3. (1789). p. 204. Nees! spec. miss. *A. eminens* Nees ast. p. 87. (v. v. c.)

β . *major* (Willd.), caule 3-4-pedali glabro, paniculà densiore, foliis ramorum ovato-lanceolatis, disco latè coerulesco. Nees.

γ . *eminens* (Nees ast. 87), caule 3-4-pedali laxè paniculato ad hastam lineis à foliis decurrentibus exasperato, fol. ram. lanceolatis, radio parciore pallidior. Ast. *eminens* Willd. enum. (1809) p. 886. Nees syn. 22.

δ . *densiflorus*, paniculà densiore, caule 2-3-pedali subpurpurascente. Nees l. c.

ϵ . *lævigatus*, caule elato 5-6-pedali, foliis oblongo-lanceolatis lævibus solo margine scabris rigidis saturatè viridi-lucidis. Nees ast. 88 (excl. verisim. syn. Ait. alibi ut species propria admissio). Ast. *lævigatus* Pursh fl. bor. am. 2. p. 553. nec Willd.

ζ . *brevirameus*, caule racemoso, ramis abbreviatis Nees l. c.

η . *albiflorus*, ligulis sæpius albis, rariùs ad violaceo-rubrum tendentibus, foliis latiusculis suprà magis scabris. Ast. *virginicus* Nees! syn. p. 22. Ast. *bicolor* Dietr. ined. Ast. *eminens* virginicus Lindl. bot. reg. t. 1656. ex foliis etiam superioribus grossè serratis et invol. squamis erectis nec patulis videtur altera species. (v. v. c.)

86. *A. Douglasii* (Lindl. adn. mss.), caule glabro racemoso-composito, ramis parcè foliosis oligocephalis laxè corymbosis, foliis lineari-lanceolatis versu basin angustatis glabris, omnibus supremis exceptis serratis, invol. laxi subæqualis squamis linearibus acutis. \mathcal{U} in Amer. septentr. occid. campis apricis prope flum. Columbia confluentem legit cl. Douglas.

α . ramis abbreviatis pleiocephalis, capitulis, minoribus.

β . ramis gracilibus elongatis oligocephalis subaphyllis, capitulis majoribus. (v. s. comm. à Soc. hort. Lond.)

87. *A. salicinus* (Willd. sp. 3. p. 2040), caule glabro paniculato-racemoso, ramis virgatis apice ramulisque corymbosis, foliis lanceolatis margine scabris infer. basi attenuatis in medio serratis, superioribus subamplectentibus, ramulorum pancis linearibus erectis, invol. laxi squamis subæqualibus. \mathcal{U} in Europæ mediæ secus fluminum ripas, in Alsatia, Germanià, Hungarià. Nees! ast. p. 90. DC. fl. fr. suppl. 470. *A. salicifolius* Scholl. fl. barb. suppl. 328. cum ic. ex W. et N. Ast. Hungaricus Poir. suppl. 1. p. 496. Ligulæ albæ. (v. s.)

*** *Leucanthi*. — *Leucanthi* et *ericoidei* Nees ast. p. 91 et 106.

Invol. squamæ laxè imbricatæ lineares vel lineari-lanceolatæ acutæ parùm inæquales, intimæ longiores angustiores. — Caulis laxè corymbosus paniculatusve. Folia caulina oblongo-lanceolata, linearia aut subulata, ramulorum minima. Capit. parva seu mediocria, ligulis albis seu pallidis.

88. *A. simplex* (Willd. enum. 2. p. 887), caule glaberrimo racemoso-decomposito, ramis apice subcorymbosis ramulisque contractis oligocephalis, foliis lanceolatis acuminatis lævissimis margine scabris, infinis subserratis, invol. laxi squamis lineari-subulatis imbricatis. \mathcal{U} in Amer. septentr. à Novâ-Cæsarâ

- ad Virginiam Nees! syn. 27. ast. p. 91. Spreng. syst. 3. p. 535. excl. syn. Similis *A. saligno*. Ligulæ albæ. *Achænia puberula*. (v. v. c.)
89. *A. LANCEOLATUS* (Willd. sp. 3. p. 2050), caule piloso-lineato à basi racemoso-decomposito, ramis erectis, ramulis corymbosis, foliis lanceolato-acuminatis sessilibus margine serrulato-scabris, infimis argutè serrulatis, ramulorum patulis, invol. laxi squamis imbricatis lineari-subulatis. $\frac{1}{2}$ in America boreali? Nees! syn. 27. ast. 92. Ligulæ albæ. (v. v. c.)
90. *A. RECURVATUS* (Nees ast. 93), caule piloso-lineato à basi racemoso-decomposito recurvato, ramulis apice subcorymbosis, foliis angustè lanceolatis acuminatis sessilibus margine scabris, caulinis rameisque argutè serrulatis, ramulorum integerrimis patentibus, involucri laxè imbricati squamis linearibus acutis. $\frac{1}{2}$ in paludosis Pennsylvaniae. An *Ast. recurvatus* Willd. et auct. ab hoc diversus? An ab *Ast. lanceolato* satis distinctus? †.
91. *A. LUCANTHERMUS* (Desf.! cat. h. par. ed. 3. p. 401), caule piloso-lineato racemoso-decomposito, ramulis circ. dicephalis laxis, foliis lineari-lanceolatis acuminatis sessilibus in ambitu scabris, inferioribus medio argutè subserrulatis, invol. squamis adpressè imbricatis apice subrecurvis. $\frac{1}{2}$ in America bor. depressis à Novâ-Cæsareâ ad Carolinam, nunc ex hortis migr. in Belgio et Germania hinc indè. Nees ast. 93. *Ast. dracunculoides* Willd. sp. 3. p. 2050. Nees syn. 28. non Lam. *A. bellidiflorus* Nees syn. 21. non Willd. *Ast. lanceolatus* Lejeune! fl. Spa 175. non Willd. *A. artemisiæflorus* Poir. suppl. 1. p. 500. Capitula parva, ligulis albis. (v. v. c. et s. ex Belgio.)
92. *A. LAXUS* (Willd. enum. 2. p. 857), caule glabro racemoso-composito vel decomposito, ramis laxis apice subcorymbosis, ramulis elongatis, foliis angustè lanceolatis acuminatis ambitu scabris, inferioribus subserratis, ramulorum linearibus obliquè patentibus, capituli terminalis squamis invol. laxis subæqualibus linearibus, reliquorum imbricatis apice reflexis. $\frac{1}{2}$ in Amer. bor. sabulosis demissis à Novâ-Cæsareâ ad Virginiam. Nees! syn. 27. ast. 95. Ligulæ pallidè carnes subalbæ. *Achæn. puberula*. (v. v. c.)
- β. caule humiliori supernè racemoso-composito, foliis caulinis summis rameisque basi dilatato-amplexantibus. Nees l. c.
93. *A. BELLIDIFLORUS* (Willd. enum. 886), caule stricto racemoso-decomposito seu suprâ decomposito angusto infernè glaberrimo albo-variegato purpureo-centeo, ramis apice racemosis racemosove-compositis, foliis lanceolatis adpresso-serrulatis margine scabris, ramulorum lanceolato-acuminatis subulatisve apice recurvis, capitulis confertis, invol. squamis imbricatis linearibus acutis inter. sæpè apice coloratis. $\frac{1}{2}$ in Amer. bor. Ligulæ albidæ, dein lilacinæ in γ amcenè purpureæ.
- α. *rubricaulis*, caule purpureo scabro, ramis brevibus apice simpliciter racemosis, capitulis approximatis, radio specioso, foliis pedicellorum subimbricatis linearibus. — *A. bellidiflorus* Willd. enum. 2. p. 886. Nees! pl. viv. miss. Lindl. pl. exs. (v. v. c. et s.)
- β. *angustus* (Nees l. c.), caule viridi alboque striato, ramis brevibus apice simpliciter racemosis, foliis ramulorum et pedicellorum acuminato-subulatis remotioribus. — *A. angustus* Nees syn. 26. (v. s. ex h. Gœtt.)
- γ. *rigidulus* (Nees l. c.), ramis longioribus racemoso-decompositis et suprâ decompositis. Cæt. ut in var. β. *Ast. rigidulus* Nees! syn. 29. (v. v. c.)
- δ. *salicinus*, racemo terminali subsimplici, foliis caulinis ramorumque inferioribus latioribus. Nees l. c.
94. *A. PARVIFLORUS* (Nees syn. 29. ast. 99), caule pubescente racemoso-decomposito ramisque apice corymbosis strictis, foliis lanceolatis acuminatis medio argutè serratis suprâ in ambitu scabris, ramulorum linearibus acutis patentibus, involucri squamis imbricatis linearibus acutis, ligulis linearibus involucri paulò longioribus. $\frac{1}{2}$ in Amer. bor. à Canadâ ad Novam-Angliam (Pursh), in Kentucky (Raf.). *Aster miser* Ait. Kew. ed. 1. v. 3. p. 205. non Linn. Ligulæ pallidè lilacinæ dein purpurasc. *Achænia pubescenti-scabra*. (v. s. sp. comm. à cl. Raf. et c. à cl. Lindl.)

95. *A. LAMARCKIANUS* (Nees ast. 100), caule piloso-lineato racemoso-decomposito coarctato, ramis paniculato-racemosis, foliis lanceolatis acuminatis sessilibus adpresso-serratis suprā scaberrimis, ramulorum lanceolato-acuminatis patentibus, invol. laxē subæqualis squamis linearibus. 2½ in Canadā (Lam.). *Ast. paniculatus* Lam. dict. 1. p. 306. non *Ait. A. Tradescanti* Nees! syn. 28. Affinis *A. divergenti*. Ligula pallidæ, ex Nees invol. breviores, in plantā vivā ab eo missā evidenter eo longiores. (v. v. c.)
96. *A. DIVERGENS* (Ait. kew. ed. 1. v. 3. p. 205), caule piloso-lineato racemoso-composito, ramis æqualibus patentibus, foliis lanceolatis acuminatis medio mucronato-serratis in ambitu scabris, ramulorum oblongo-lanceolatis basi attenuatis integerrimis mucronulatis patentibus, invol. squamis imbricatis lanceolatis. 2½ in agris neglectis et sylvis siccis Amer. bor. à Novo-Eboraco ad Carolinam (Pursh), et in dumetis prope Jalapam (Schied. et Dep.). Nees syn. 28. ast. p. 101. *A. divaricatus* Raf. ! in litt. ? (v. s.)
- β. *humilior*, foliis brevioribus magis confertis, paniculæ ramis infer. elongatis. Nees l. c.
97. *A. FRAGILIS* (Willd. sp. 3. p. 2051), caule piloso-lineato racemoso-decomposito, ramis subpaniculato-racemosis, foliis infer. oblongis adpresso-serratis; super. linearibus, omnibus in ambitu scabris, ramulorum oblongo-lanceolatis integerrimis mucronulatis patentibus, invol. arcetē imbricati brevis squamis lanceolato-linearibus. 2½ in Americā bor. (Muhl.), in paludosis pinetis Novæ Cæsareæ et Alabamæ (Torr.) Lindl. ! bot. reg. t. 1527. *Desf. fl. philad.* 2. p. 117. *Ast. scoparius* Nees syn. 28.
- β. *subinteger*, foliis omnibus linearibus elongatis undulatis subintegerrimis. — *A. fragilis* Nees syn. p. 29. *Desf. ! cat. h. Par. ed. 3. p. 176.* (v. s.)
98. *A. FALCATUS* (Lindl. adn. mss.), caule piloso angusto racemoso-subcomposito, ramulis brevibus 1-cephalis paucifoliis, foliis linearibus obtusis falcatis pilosisculis, invol. laxi parum imbricati squamis linearibus acutiusculis. 2½ in Americā arcticā. Facies omnino *A. hiemalis* sed affinitas vera cum *A. fragili*. Folia infer. Bupleuro falcato similia. Radius albus. (Lindl. mss.)
99. *A. TRADESCANTI* (Linn. sp. 1230), caule glabriusculo racemoso-subcomposito patulo apice simpliciter racemoso, ramis virgatis, ramulis subsecundis, foliis linearibus acuminatis margine ciliato-scabris, infer. à medio argutē remotē serratis, ramorum ramulorumque patentibus mucronatis, involucrio imbricato. 2½ in Amer. bor. sæpibus et paludibus exsiccatis à Canadā ad Virginiam. Nees ast. p. 103. *Ast. vimineus* Lam. dict. 1. p. 308. *Desf. ! cat. h. Par. 176.* *Ast. pinifolius* Nees! syn. 29. Ligulæ albæ demum subviolaceæ. Capitula parva. (v. v. c.)
100. *A. CORDIFOLIUS* (Michx. fl. bor. sm. 2. p. 112), caule glabro racemoso-composito decompositoque, ramis patulis intricatissimis, foliis glabris, inferioribus lineari-lanceolatis suberrulatis, rameis depauperatis squamiformibus patuloreflexis, ramis apice 1-cephalis, invol. imbricati squamis linearibus obtusiusculis, ligulis acutis, ovario glabro. 2½ in aridis sylvarum Carolinæ infer. (Michx. Bosc! Lindl. !), circa Noveboracum (Don) Lindl. bot. reg. t. 1487. *Desf. ! cat. h. Par. ed. 2. p. 122.* *A. foliolosus* Ell. sketch 2. p. 345. excl. ? syn. *Species Willd. sp. 3. p. 2028*, ex Nees syn. p. 30, est fortē diversa. Ligula pallide carnea. Pappus flosc. disci æqualis. (v. v. c.)
101. *A. DUMOSUS* (Linn. sp. 1227), caule glabriusculo racemoso-composito decompositoque patulo, ramis apice corymbosis, foliis linearibus integerrimis margine serrulato-scabris, super. internodiis longioribus, rameis ramulorumque parvis acutis, invol. squamis imbricatis linearibus acutis. 2½ in Amer. bor. campis siccis sterilibus à Novā Angliā ad Carolinam (Pursh). Nees ast. p. 99. *Ast. foliolosus* Nees syn. 30. *A. foliosus* Pers. ench. n. 46 ? *A. Novæ Angliæ*, etc. Herm. parad. p. 95. icon opt. Capitula parva. Ligulæ albæ. (v. v. c.)
102. *A. ERICOIDES* (Linn. sp. 1227) caule glabro racemoso-composito, ramis virgatis, ramulis subsecundis ab apice remotis, foliis linearibus angustis late-

- gerrimis glaberrimis, invol. brevis squamis imbricatis subulato-patulis infer. cuspidatis. \mathcal{U} in Amer. bor. sterilibus et ad margines viarum à Canadâ ad Carolinam (Pursh), in Kentucky (Raf.) — Nees syn. 31. ast. 107. Ast. tenuifolius Willd. sp. 2. p. 2026. excl. syn. ex Nees. Rami apice non corymbosi sed producti et t-cephali. Capitula parva, ligulis albis. (v. s. comm. à cl. Rafin.)
103. *A. GLABRELLUS* (Nees syn. 31. ast. 107), caule racemoso-subcomposito, ramis filiformibus, foliis caulinis infer. lanceolatis serratis, super. lineari-subulatis integerrimis recurvatis, invol. subhemisphærici squamis imbricatis à basi oblongo-cuspidatis subciliatis. \mathcal{U} in campis petrosis Amer. bor., in Pennsylvania, Novo Eboraco.
- α. caule folisque infer. suprâ pilosis. — *A. sparsiflorus* Willd. enum. 880 ex parte sec. Nees.
- β. glaber. — *A. dumosus* Nees syn. 31 non Linn. *A. sparsiflorus* Michx. fl. 2. p. 112. non Willd. *A. ramosissimus* et ni fallor *A. leptophyllus* et *A. subulatus* hort. Par. (v. s.)
104. *A. PILOsus* (Willd. sp. 3. p. 2025), caule villosa racemoso-composito, foliis lineari-lanceolatis integerrimis albo-villosis, capitulis secundis, invol. oblongi squamis lanceolato-subulatis imbricato-patulis. \mathcal{U} in terrâ Illinoisensium Amer. bor. Nees syn. 30. ast. 109 Ast. villosus Michx. fl. bor. am. 2. p. 113. Cl. Nees dubitat num reverâ ab *A. glabello* differat.
105. *A. HERBECULUS*, caule fruticuloso gracili pube patulâ confertâ brevi hirsuto, ramis laxis parçè ramosis ramulisque foliosis t-cephalis, foliis sessilibus linearibus integerrimis mucronato-acutis utrinquè villosa-scabris, super. ciliatis, invol. laxè imbricatis squamis linearibus puberulis acutis. \mathcal{J} in Mexicanâ prov. Texas districtibus orientalis legit cl. Berlandier (pl. exs. n. 2085). Affinis *A. scopario* sed rami hirsuti laxiores oligocephali, capitulo triplò ferè minore (v. s. comm. à cl. inv.)
106. *A. SCOPARIUS* (DC. non Nees), caule fruticuloso erecto ramosissimo glabro apice summo vix ac ne vix puberulo, ramis erectis racemiferis, foliis sessilibus linearibus scabridis, super. ciliatis, ramulis foliolosis, invol. squamis laxè imbricatis linearibus acutis ciliatis. \mathcal{J} in Mexicanâ prov. Texas districtibus orientalis legit cl. Berlandier (pl. exs. n. 1868, 1871 et 1894). *Habitus ferè Ericæ scopariæ*. Folia 4-5 lin. longa semilineam lata. Pappus rufus. Achænium puberulum. (v. s. comm. à cl. inv.)

**** Densiflori Nees ast. 110.

Invol. squamæ arcte imbricatæ. Rami densè racemiflori, capitulis confertis subsecundis. Folia integra aut serrata.

107. *A. HIRSUTICAULIS* (Lindl. adn. mss.), caule racemoso hirsutissimo, ramis foliis brevioribus vel parùm longioribus, foliis caulinis lineari-lanceolatis integerrimis acuminatis, costâ subtus hirsutissimâ, capitulis subsecundis, invol. squamis arcte imbricatis subulatis. \mathcal{U} in Amer. septentr. juxta Albany (Torrey.). Radius albus. Discus purpureus. (Lindl.)
108. *A. DIFFUSUS* (Ait. kew. ed. 1. v. 3. p. 205), caule pubescente racemoso-composito, ramis recurvo-patentibus, foliis caulinis lanceolatis, ramis oblongo-lanceolatis medio serratis suprâ scabriusculis marginè scaberrimis, capitulis subsecundis confertis, invol. squamis imbricatis acutis. \mathcal{U} in Amer. bor. agris et sæpibus à Canadâ ad Carolinam (Pursh). Nees syn. 28. ast. 110. Lindl. ! pl. exs. Spreng. syst. 3. p. 586 excl. syn. *A. divergens* β Nutt. gen. 2. p. 159. *A. secundiflorus* Deaf. ! h. par. 1815. Capitula parva, ligulis albis seu pallidè lilacinis, disco luteo mox purpur. Achænia ad costas pubescenti-scabra (Nees). (v. v. p.)
109. *A. STENOPHYLLUS* (Lindl. adn. mss.), caule glabresculo racemoso, ramis patentibus apice densissime racemosis, foliis caulinis lineari-lanceolatis acuminatis scabriusculis, capitulis secundis, invol. squamis linearibus acumina-

- tis, interioribus membranaceis coloratis. $\frac{1}{2}$ in Amer. septentr. ? in hortis anglis occurrit. Radii dilute carnei. Discus mutabilis (Lindl. mss.)
110. *A. miser* (Linn. sp. 1232. excl. syn.), caule pubescenti-scabro racemoso-composito, ramis erectis, foliis caulinis oblongo-cuneiformibus acuminatissimis medio serratis rameisque ovalibus subintegerrimis supra scabris, capitulis subsecundis confertis, invol. laxè imbricatis squamis acuminatis. $\frac{1}{2}$ in pratis udis ad Philadelphiam. Nutt. gen. am. 2. p. 158. Nees ast. p. 111. non alior. Capit. a lin. lata. Ligulæ lineares albæ. Achæn. pubescentia.
- ß. myrsifolius*, caule altiori, capitulis majoribus. $\frac{1}{2}$ in hortis cultus. *A. myrsifolius* Willd. enum. suppl. 59.
- γ. abbreviatus*, ramis abbreviatis capitulisque ideò magis confertis Nees l. c.
111. *A. pendulus* (Ait. kew. 3. p. 205), caule pubescenti-hirto laxo diffuso racemoso-composito, ramis elongatis flaccidis, foliis elliptico-lanceolatis pubescenti-scabris, caulinis medio adpresso-serratis, ramulorum ellipticis integerrimis, invol. imbricatis. $\frac{1}{2}$ in Amer. bor. agris neglectis (Pursh) ad Novoboracum (Bern.) Nees syn. 28. Ast. 114. Hoffm. phyt. bl. p. 80. t. c. f. 2. Ast. horizontalis Desf. l. cat. h. par. ed. 3. p. 402 ? Capitula parva, radio pallidè purpurascente seu albo. Achænia pubescenti-scabra. (v. v. c. in h. Par.)
112. *A. bifrons* (Lindl. adn. mss.), caule scabro laxo racemoso, ramis apice racemosis, foliis ovali-lanceolatis supra scabris ferè omnibus serratis, capitalis secundis, invol. squamis subulatis. $\frac{1}{2}$ in Amer. septentr. *A. pendulo* affinis, sed ramulis oligocephalis et capitulis amplioribus distinctus. (Nees in litt. ad Lindl.)
113. *A. Menziesii* (Lindl. adn. mss.) caule incano apice racemoso, foliis oblongo-lanceolatis sessilibus integerrimis scabriusculis incanis, inv. squamis obtusis squarrosis. $\frac{1}{2}$ in Americâ septentr. occid. legit cl. Menzies. Ab *A. concolore* vix differt nisi pilis rigidis nec mollibus, necnon invol. squamis squarrosis nec adpressis. (Lindl. mss.)
114. *A. concolor* (Linn. sp. 1228), caule pubescente apice racemoso-composito, foliis oblongo-lanceolatis sessilibus integerrimis cano-pubescentibus, capitulis ad axillas fol. ram. solitariis subsessilibus racemos foliosos constituentibus, invol. squamis imbricatis lanceolatis sericeis. $\frac{1}{2}$ in Amer. bor. pinetis quercetisque littoreis arenosis à Novâ Cæsareâ ad Floridam. Nees syn. 30. ast. 114. Lindl. ! pl. exs. Ell. sketch 2. p. 350. *Virgaria concolor* Raf. ! in litt. 1831. Radix in arenosis tuberosa. Ligulæ 12-15 lætè cœruleæ. Discus etiam cœruleus (Ell.). Radius discusque purpureus (Nees). Achænium densè sericeo-villosum. Pappus rufus. (v. s. sp. ex Carolinâ et Ohio.)
115. *A. squarrosus* (Walt. car. 209), caule scabro composito-racemoso, ramis oligocephalis foliis minutis è basi cordatâ amplexicanli acuminatis reflexis margine scaberrimis, invol. imbricatis squamis lineari-lingulatis mucronatis laxè adpressis apice subfoliaceis. $\frac{1}{2}$ in pinetis sterilibus Carolinæ infer. (Pursh) et Georgiæ (herb. Dnby!) Nees ast. 115. Ell. sketch 2. p. 350. Rami virgati tennes. Folia parva rigidula. Ligulæ 15-20 lætè cœruleæ. Achænia pilosa. (v. s. sp. coll. à cl. Bosc.)
116. *A. ramulosus* (Lindl. adn. mss. non Labill.), caule piloso erecto angusto racemoso, ramulis corymbosis 1-cephalivæ, foliis linearibus serrulato-scabris basi dilatatis sessilibus, ramulorum rectis minoribus, invol. squamis acutissimis mucronatis squarrosis. $\frac{1}{2}$? in Americâ arcticâ. Capitula duplò majora quàm *A. multiflori*.
- ß. incano-pilosus*. — in Amer. arcticâ. (Lindl. mss.)
117. *A. multiflorus* (Ait. kew. ed. 1. v. 3. p. 116), caule pubescente racemoso-composito diffuso, foliis linearibus integerrimis margine serrulato-scabris subtrinerviis, ramulorum recurvis, invol. squamis laxè imbricatis apice recurvis subspathulatis mucronatis. $\frac{1}{2}$ in Amer. bor. campis aridis ad viarum margines à Canadâ ad Carolinam (Pursh). Nees ast. p. 116. Lindl. ! pl. exs. *A. ericoides* Lam. dict. 1. p. 304. Michx. fl. bor. am. 2. p. 113. ex Lindl. *A. ericoides multiflorus* Pers. ench. 2. p. 443. *A. multiflorus* et *A. ciliatus* Nees syn.

30 et 29. — Dill. elth. 40. t. 36. f. 40. Capitula parva lignis albis. Achen. pubescenti-scabra. (v. s.)

β. ciliatus (Nees l. c.), foliis latioribus ramealibus patulis nec recurvis, omnibus ciliatis, caule stricto. — *A. ciliatus* Willd. sp. 3. p. 2027. (v. s. comm. à cl. Lindl.)

118. *A. RACEMOSUS* (Ell. sketch 2. p. 348), caule leviter pubescente diffuso, ramis gracilibus elongatis, foliis lineari-lanceolatis subtus subpubescentibus margine scabris, capitulis subsessilibus confertis juxta summitem ramorum, invol. squamis arcuè imbricatis lineari-lanceolatis. $\frac{1}{2}$ in sole pingui insulæ Paris Amer. bor. Capitula parva. Ligulæ pallidæ purpureæ. Affinis vid. *A. multifloro*.

119. *A. CANESCENS* (Pursh flor. bor. am. 2. p. 547), caule subsimplici racemoso-villoso, ramis 1-cephalis, foliis linearibus integerrimis margine serrulato-scabris subtrinervibus, ramulorum recurvis, invol. squamis acuminatis squarrosis. ② in Massachuset (Lindl.), ad ripas fl. Missouri (Pursh), in argillosis denudatis ab Arikarees ad Fort-Mandan (Nutt.). Aster biennis Nutt. gen. am. 2. p. 155. non Ledeb. Calimeris biennis Nees ast. 231. Capit. magnit. Bellidia. Ligulæ circ. 20 purpurascens-cærulesc. Hic nec ad Calimeridem relatus ex aut. cl. Lindley in adn. mss.

120. *A. MICROPHYLLUS* (Torr. in Lindl. adn. mss. non Vent.), caule glaberrimo composito racemoso, ramis 1-cephalis strictissimis, foliis minutis ovatis glaberrimis cauli adpressis, invol. turbinatis squamis linearibus acutis basi albidis arcuè imbricatis. ① in Amer. bor. ditione Alabamæ legit cl. Torrey. Radii cyanei (Lindl. mss.)

***** Concinni Nees ast. p. 118.

Invol. squamæ imbricatæ basi latiores pallidæ apice virides ciliatæ. Recept. alveolæ dentiformes. Achænia sparsim puberula. Folia latiuscula. Radius speciosus raro albus.

121. *A. PURPURATUS* (Nees ast. p. 118), caule subsimpliciter racemoso-virgato, ramis elongatis 1-cephalis, foliis angusto-lanceolatis amplexicaulibus medio argutè serratis margine scabris, super. ovato-lanceolatis, ramorum pluribus lanceolatis patulis integerrimis sequè distantibus æqualibusque, invol. turbinatis squamis imbricatis lanceolatis apice coloratis. $\frac{1}{2}$ in Amer. septentr. ? Aster miser hort. Par. Lam. dict. 1. p. 308. excl. syn. Caulis spec. mei glaber nec pilis seriatis scaber. Discus mox purpureus. Ligulæ saturatè purpureæ. Achænia glabra. (v. s. coll. in h. Par. anno 1798. sub nom. *A. miserl*.)

122. *A. AZUREUS* (Lindl. adn. mss.), caule racemoso-composito virgato, ramulis elongatis 1-cephalis, foliis lanceolatis utrinquè acuminatis scaberrimis subseriatis, superioribus integris, ramorum subulatis, invol. hemisphærici squamis imbricatis apice tantum patulis. $\frac{1}{2}$ in Louisiana. Radius azureus. Habitus *A. purpurati*. (Lindl. mss.)

123. *A. RETROFLEXUS* (Lindl. adn. mss.), caule composito-racemoso, ramis brevibus 1-cephalis, foliis lineari-lanceolatis integerrimis glaberrimis, ramulorum lineari-subulatis recurvis distantibus, invol. hemisphærici squamis lineari-lanceolatis squarrosis. $\frac{1}{2}$ in Amer. septentr. Discus albidus parum mutabilis. Radii cærulei. *A. squarrosus* hort. Angl. est species diversissima. (v. s. comm. à cl. Lindl.)

124. *A. TURBINELLUS* (Lindl. adn. mss.), caule subsimpliciter racemoso-virgato, ramis elongatis filiformibus sub-1-cephalis, foliis oblongo-lanceolatis subamplexicaulibus integerrimis, ramorum oblongis obtusis sensim in subulata decrescentibus, invol. turbinatis squamis linearibus obtusiusculis apice herbaceis. $\frac{1}{2}$ in Louisiana. Species ab *A. purpurato* diversa foliorum formâ, involuacro magis turbinato, ramisque filiformibus nec rigidis strictis. (Lindl. mss.)

125. *A. TENUIFOLIUS* (Linn. sp. 1227), caule glabro racemoso-decomposito ra-

motque virgatis, foliis angusto-lanceolatis margine scabris, infer. medio subserratis, ramorum plaribus lanceolatis integerrimis patentibus sequè distantibus aequalibusque, ramulis 1-cephalis, invol. ovato-hemisphaerisci squamis imbricatis è basi lineari-oblongà acutis apice patulis. \mathcal{L} in Americà septentr. Nees ! ast. p. 119. Lindl. ! pl. exs. Ast. polyphyllus Willd. enum. 2. p. 888. Ast. junceus Nees syn. 26. Ast. junceus et polyphyllus Pursh. Nutt. Ligulæ albo-violaceæ. Achænia pubescentia. Caules erecti. (v. v. c. in h. Par. et s. ex h. Gært.)

β . caule recurvato, ramis adscendentibus, invol. laxiore radio densiore. Ast. recurvatus Nees syn. 26. A. tenuifolius Mōse handb. 2. p. 1485 ex Nees.

γ . caule ramisque erectis, capitulis paulò minoribus, invol. magis imbricato. Nees l. c.

126. A. *concinus* (Willd. enum. 884), caule glabriusculo laxo subcorymboso, ramis virgatis dichotomo-paniculatis, foliis lanceolatis amplexantibus remotè argutè serratis, ramulorum oblongis integerrimis, invol. arcetè imbricatè squamis margine albidis. \mathcal{L} in Amer. bor. campis et sylvis à Novo-Eboraco ad Pensylvaniam (Pursh). Nees syn. 21. ast. 121. Lindl. bot. reg. t. 1619. Colla hort. rip. app. 3. p. 22. t. 12. act. sc. Tur. 33. p. 134. t. 12. Capit. mediocria, radio pulchrè cœruleo.

β . ramis strictioribus magis racemosis, foliis parvis orebrioribus. Nees l. c. (v. s. c. in h. Par. 1815.)

127. A. *onustus* (Nees ast. 122), caule glabriusculo supernè seriatim hirsutulo racemoso-decomposito, ramis apice ramosis laxis, foliis lanceolatis utrinquè attenuatis amplexantibus medio adpresso-serratis margine scabris, capitulis subsecundis breviter pedicellatis, invol. brevis squamis imbricatis lanceolatis discois subaequantibus. \mathcal{L} in Amer. bor. ? A. confertus Desf. ! cat. h. Par. ed. 3. p. 401 non Nees. Ligulæ albæ violaceo subinctæ. Achænia glabra. Stricto jure nomen Desf. antiquius servandum esset, sed nomen Neesianum admisi ne confusio generis augeatur. (v. s. c.)

β . *squarrosus* (Lindl. adn. mss.), foliis ramulorum linearibus squarrosis. Cultus in hortis Angliæ sub nomine A. Tradescanti cœrulei. (Lindl. mss.)

128. A. *Chilensis* (Nees ast. 123), caule basi glabro apice seriatim piloso racemoso-decomposito, ramis apice densè corymbosis racemosive, foliis lanceolatis amplexicaulibus acutis crenato-serratis utrinquè scabris, pedicellorum parvis oblongis squarrosis, invol. squamis arcetè imbricatis oblongis obtusis \mathcal{L} ? in regione montanà agri Chilensis (Hænk.), in Californiâ (Cham.) A. Radula Less. in linnæâ 1831. p. 125. excl. syn. ex Nees.

129. A. *strictus* (Poir. suppl. 1. p. 498), caule glabro stricto ramoso-decomposito, ramis strictis coarctatis, foliis lanceolatis attenuatis subamplexantibus medio subserratis ambitu supino-scabris, invol. arcetè imbricato. \mathcal{L} in Amer. bor. Ligulæ elongatæ lilacinæ. Similis dicitur A. præalto. Nees ast. p. 124. A. strictus Pursh. A. strictus Wender. et A. strictus hortul. sunt diversi.

β . *angustifolius* (Lindl. adn. mss.), foliis lineari-lanceolatis vix serratis. in Amer. arcticâ.

130. A. *mutabilis* (Ait. kew. ed. 1. v. 3. p. 205. an Linn. ?), caule lævi à basi paniculato decomposito patulo, ramis densè corymbosis, foliis oblongo-lanceolatis acuminatis lævibus margine scabris subtile glaucis, infer. medio serratis, super. amplexicaulibus integerrimis, invol. disco breviter squamis imbricatis lanceolatis. \mathcal{L} in Amer. bor. agris neglectis et sylvis à Pensylvaniâ ad Virginiam (Pursh). Nees syn. 22. ast. 125. Willd. sp. 3. p. 2045. excl. syn. Pluk. A. versicolor hortul. — Herm. lpgd. 6. p. 67. ic. Variat foliis omnibus aut serratis aut integerrimis. Discus luteus mox purpureus. Ligulæ cœruleæ seu purpureæ. (v. v. c.)

131. A. *confertus* (Nees ! syn. 21. ast. 126. non Desf.), caule lævi apice subcorymboso-decomposito, ramis confertissimis corymbosis, foliis oblongo-lanceolatis acuminatis basi cordato-amplexantibus medio adpresso serratis ram-

gine scabris, invol. imbricato. $\frac{1}{2}$ in Amer. bor. Ligulæ albæ. Achænia glabra. Variat foliis magis distinctè serratis. (v. v. c. ab hort. Bonn. mss.)

132. *A. VERSICOLOR* (Willd. sp. 3. p. 2045), caule lævi paniculato-composito, ramis apice simpliciter corymbosis densis, foliis oblongo-lanceolatis acuminatis lævissimis concoloribus, infer. medio serratis, super. amplexentibus integerrimis, invol. squamis imbricatis lanceolatis. $\frac{1}{2}$ in Amer. bor. campis neglectis et ad margines sylvarum à Novâ-Cesareâ ad Carolinam. Nees! syn. 22. ast. 127. Spreng. syst. 3. p. 335, excl. syn. Capitula majuscula radio primum albo demum saturatè violaceo; in iisdem indiv. adsunt flores violacei et albi. Adeat proles monstruosa radio abbreviato lacero à Nees obs. (v. v. c. ab h. Bonn. mss.)

133. *A. LÆVIS* (Linn. sp. 1230), caule lævi racemoso-composito, ramulis brevibus in apice ramorum approximatis, foliis lanceolatis amplexentibus glabris ambitu scabris viridibus, inferioribus subserratis, invol. imbricato disco brevior. $\frac{1}{2}$ in Amer. bor. sylvis et ad paludum latera à Novo-Eboraco ad Virginiam (Pursh). Nees syn. 24. ast. p. 128. Ait. kew. 3. p. 206. Lindl. bot. reg. t. 1500. Capitula ampla, ligulis cœruleis. Achænia pubescentia. Nomen subdeciens, nam folia non sunt omninò lævia. (v. v. c.)

134. *A. LÆVIGATUS* (Willd. sp. 3. p. 2046), caule lævi racemoso-composito virgato, foliis oblongo-lanceolatis amplexentibus lævibus margine scabris saturatè viridibus, infer. in petiolum alatum attenuatè parçè serratis, super. sensim minoribus integerrimis, invol. discum æquantis squamis imbricatis lanceolatis. $\frac{1}{2}$ in Amer. bor. paludosis sylvisque uliginosis à Pensylvaniâ ad Virginiam (Pursh). Nees syn. 24. ast. 129. Hook. bot. mag. t. 2995. Caulis purpureus. Ligulæ elongatæ cœruleæ. Sæpè in hortis nunc cum lævi nunc cum mutabili confusis. (v. v. c.)

135. *A. AMPLEXICAULIS* (Muhl. in Willd. sp. 3. p. 2046), caule lævi paniculato-corymboso, ramis dichotomo-corymbosis coarctatis, foliis oblongis acutis basi cordatè amplexentibus lucidis margine scabris infer. subserratis, invol. arcuè imbricatè squamis latè linearibus. $\frac{1}{2}$ in Pensylvaniâ (Muhl.), in sabulosis ad margines sylvarum à Novâ-Cesareâ ad Virginiam (Pursh). Nees syn. 22. ast. 130. Capitula mediocria, ligulis saturatè cœruleis.

136. *A. RUBRICAULIS* (Lam. dict. 1. p. 305), caule latè stricto racemoso-composito-subpaniculato, foliis oblongo-lanceolatis amplexentibus margine scabris glaucis, radicalibus ovatis in petiolum contractis obtusis, caulinis infer. basi attenuatis subserratis, super. integerrimis, invol. disco brevioris squamis imbricatis lanceolatis. $\frac{1}{2}$ in Amer. septentr. (Lam.). Nees ast. p. 131. excl. fortè var. α . Lindl. ! pl. exs. Capitula saturatè violacea. Recept. fimbriatè, fimbriis in pilum desinentibus. Achænia glabra. (v. v. c.)

137. *A. CYANÆUS* (Hoffm. phyt. bl. 1. p. 71. t. B. f. 1), purpurascens-glaucus, caule lævi racemoso-composito subpaniculato, foliis oblongo-lanceolatis amplexentibus lævibus margine scabris glaucis, super. integerrimis, radicalibus spathulatis obtusis crenato-serratis, invol. disco brevioris squamis arcuè imbricatis ex ovato-triangularibus. $\frac{1}{2}$ in Americâ boreali. Nees ast. 182. Lindl. bot. reg. t. 1495. Recept. alveolato-dentatum epilosum. Ligulæ cœruleæ.

α . invol. squamis angustioribus. — *A. glaucus* Nees syn. 23. *A. decorus* Desf. ined. ex Nees.

β . invol. squamis latioribus, ext. ovato-dilatatis. — *A. politus* Nees syn. 23. et *A. cyanæus* ibid. *A. rubricaulis cyanæus* Poir. suppl. 1. p. 493.

Species non satis notæ.

138. *A. ANSIF* (Mart. in act. ac. Monac. phil. 6. p. 195), fruticosus, foliis sessilibus angusto-linearibus acutis subtus hirsutis, capitulis solitariis axillaribus subsessilibus. $\frac{1}{3}$ in monte Kurmâ Arabiæ. Aster ericæfolius Forsk. arab. p. 150. *Ansif*, arab.

139. *A. CARICIFOLIUS* (H. B. et Kunth nov. gen. sm. 4. p. 91. t. 833), caule

- herbaceo erecto simplici supernè paniculato polycephalo, foliis linearibus amplexicaulibus integerrimis acuminatis glabris, invol. squamis linearibus puberulis subæqualibus ligulis linearibus circ. 35. $\frac{1}{2}$ in ruderatis prope urbem Mexico. Ligulæ albæ, an 1-2-seriales? An *Stenactis*? a cl. Neesio omisus.
140. *A. P. DISCORDEUS* (Ell. sketch 2. p. 358), caule erecto subvillosa, ramis panicis virgatis, foliis spathulato-ovatis, acutis serratis pilosis subtus pallidioribus, paniculæ ramis virgatis axillaribus oligocephalis, invol. squamis subulatis villosis laxè adpressis subsquarrosis, ligulis nullis! — frequens in pinguibus inter flum. Alabama et Chatahouchie. — Discus purpureus. Achænia glaberrima. An *Vernonia*, an *Chrysocoma* species?
141. *A. EXILIS* (Ell. sketch 2. p. 344), glaberrimus, caule gracili elato parè ramoso, foliis prælongis lineari-subulatis margine subscabridis, capitulis racemosis, invol. squamis lineari-lanceolatis radio dimidiò brevioribus adpressis. $\frac{1}{2}$ in Georgiae occidentalis humidis. Ligulæ circ. 20 pallidè purpureæ. Fortè affinis *A. glabello*.
142. *A. FISCHERIANUS*, caule 1-2-cephalo, foliis lanceolatis sessilibus pubescentibus supernè dentato-serratis, invol. squamis linearibus acutis coloratis. $\frac{1}{2}$ in Sibiria. *A. flexuosus* Fisch. mem. soc. mosc. 3. p. 73. non Nutt. Caulis plius minus flexuosus. Radius amènè purpureus. An inter *A. alpigenos* admittendus?
143. *A. P. GRACILIS* (Nutt. gen. 2. p. 158), caule minutim pubescente, ramis 2-3-cephalis, foliis oblongo-lanceolatis remotè inciso-serratis acutis subamplexicaulibus glabrisculis margine scabris, invol. subcylindrici squamis imbricatis lineari-oblongis partim acutis partim squarrosis, ligulis circ. 12. $\frac{1}{2}$ in savannis prov. Kentucky et Tennesse. Nees ast. 41. Affinis dicitur *A. spectabili* et conyzoidi, sed prior est *Aster* verus posterior *Sericocarpus*. Ergo affinitas dubia. Ligulæ pallidè cœruleæ.
144. *A. LUCIDUS* (Wender. obs. bot. n. 3. ex flora 1829. erg. p. 25), caule elato supernè ramoso asperimo, foliis radicalibus oblongo-lanceolatis integerrimis in petiolum decurrentibus glaberrimis splendentibus, superioribus lanceolatis subserratis capitulis paniculatis, invol. squamis linearibus discum subæquantibus. $\frac{1}{2}$ patr. ign. Radius cœrulescens, omnium speciosissimus. Videtur diversus ab *A. lævi* Dietr. lex. 2. p. 13 quem plurimi ad *A. lævem* referunt. Conf. Nees ast. p. 282.
145. *A. MARGINATUS* (H. B. et Kunth nov. gen. am. 4. p. 91), herbaceus cæspitosus diffusus procumbens ramosus, foliis petiolatis oblongis spathulatis apicem versus dentatis adpresso-pilosis, capitulis solitariis terminalibus, invol. hemisphærici squamis imbricatis lanceolato-linearibus colorato-marginatis. $\frac{1}{2}$ in altâ planitie Novo-Granatensi juxta urbem Santa-Fe de Bogota. Ligulæ 20-25 violacæ? Achænia compressa piloso-sericea. An verè generis? Affinitas dubia. Nees ast. 279.
146. *A. NITIDULUS* (DC. in Wight contrib. bot. ind. p. 9), glaber, caule erecto tereti apice corymboso-ramoso, foliis sessilibus subsemialexicaulibus lanceolato-linearibus acuminatis 1-nerviis margine serrulato-acabris, invol. squamis linearibus acuminatis margine scariosis disco paulò brevioribus. $\frac{1}{2}$ in India orient. verisim. ad montes Neelgerry legit cl. Wight. Ligulæ lineares ex sicco cœruleo-purp.; stylifera steriles. Achænia disci compressa glabra tenula. Pappus rufus sæpè crispatus 1-serialis. An ad *Galatella* refer.? (v. s. in h. Wight.)
147. *A. PRATENSIS* (Raf. fl. lud. p. 67), caule frutescente paniculato foliolato, foliis semialexicaulibus oblongis integris uninerviis. — in Louisiana. Capitula diam. pollic., ligulis violaceis. †
148. *A. FUSILLUS* (Horn. h. hafn. 2. p. 812), caulibus filiformibus corymbosis, foliis linearibus obtusis trinerviis integerrimis strigosis reflexis, involucris squamis æqualibus radio duplò brevioribus. $\frac{1}{2}$ patr. et cæt. car. ign. An *Galatella* spec.? †
149. *A. RETICULATUS* (Parsh fl. bor. am. 2. p. 548), undiquè cano-tomentosus, caule supernè ramoso, ramis apice corymboso-racemiferis, foliis lanceolato-

oblongis utrinque acutis sessilibus margine revolutis, subhis reticulato-venosis triplinerviis, pedunculis subaphyllis, invol. imbricati laxiusculi squamis acutissimis. 2 $\frac{1}{2}$ in paludibus exsiccatis Georgiæ et Carolinæ (Pursh). Nees ast. 277. Disci radii albi. Capit. mediocria. An Galatellæ sp.?

150. *A. RIPARIUS* (H. B. et Kunth nov. gen. am. 4. p. 93), herbaceus caespitosus procumbens ramosus microphyllus, ramis glabris floriferis elongatis, foliis sessilibus linearibus acutis integerrimis 1-nerviis glabris, capitulis solitariis terminalibus, invol. hemisphaerici squamis lanceolato-linearibus imbricatis, exter. subquarrosis. 2 $\frac{1}{2}$ in humidis Mexici juxta lacum Cuiseo. *A. riparius* Nees ast. p. 280 non p. 73. Ligulæ 20-25 albæ. Pappus rufescens pilosus. Achæn. ign.

†† *Species exclusæ.*

- A. ACRI* Linn. herb. = *Tripolium vulgare*.
A. ACRI Linn. = *Galatella punctata* a.
A. ACRI Willd. = *Galatella rigida*.
A. ACRI Nees. = *Galatella dracunculoides*.
A. ACULATUS Labill. = *Eurybia aculeata*.
A. ACUMINATUS Michx. = *Diplostephium acuminatum*.
A. AFFINIS Less. = *Felicia affinis*.
A. ÆTHIOPICUS Burm. = *Agathæa microphylla*.
A. ALBESCENS Wall. = *Amphirepis albens*.
A. ALBUS Willd. = *Heleastrum album*.
A. ALPINUS γ Poir. = *Galatella Lusitanica*.
A. ALTAICUS Willd. = *Calimeris Altaica*.
A. ALYSOIDES Turcz. = *Calimeris alyssoides*.
A. AMBIGUUS Bernh. = *Biotia Schreberi*.
A. AMYGDALINUS Lam. = *Diplostephium amygdalinum*.
A. AMYGDALINUS Ell. = *Diplostephium umbellatum*.
A. ANGUSTIFOLIUS Jacq. = *Felicia angustifolia* L.
A. ANNUUS Linn. = *Stenactis annua*.
A. ARGOPHYLLUS Labill. = *Eurybia argophylla*.
A. ARRAGONENSIS Asso. = *Galatella Arragonensis*.
A. ASPER Ham. = *Calimeris asperrima*.
A. ASPERULUS Wall. = *Stenactis asperula*.
A. ASPERRIMUS Wall. = *Calimeris asperrima*.
A. AURANTIUS Linn. = *Clomenoclema aurantia*.
A. BELLIDIASTRUM Scop. = *Bellidiastrum Michellii*.
A. BELLIDIODES Don. = *Stenactis bellidioides*.
A. BERBERIFOLIUS Cunn. = *Eurybia erubescens*.
A. BIENNIS Ledeb. = *Callistephus*? *biennis*.
A. BIFIDUS Nees. = *Galatella dracunculoides*.
A. BIFRONS All. = *Inula bifrons*.
A. BONARIENSIS Spreng. = *Erigeron glaucum*.
A. CANUS W. et Kit. = *Galatella cana*.
A. CASTRORUM Less. = *Felicia Castorum*.
A. CAPENSIS Less. = *Agathæa amelloides*.
A. CAUCASICUS Muss. = *Diplopappus Caucasicus*.
A. CHABERIS Less. = *Agathæa strigosa*.
A. CHAMÆDRIFOLIUS Lam. = *Printzia Bergii*.
A. CHINENSIS Linn. = *Callistephus Chinensis*.

- A. CHRYSANTHEMIFOLIUS hortul. = *Diplopappus filifolius*.
- A. CHRYSOPHYLLUS Cunn. = *Eurybia chrysophylla*.
- A. CHRYSOCOMOIDES Desf. = *Felicia angustifolia*.
- A. CILIARIS Thunb. mss. = *Mairia taxifolia*.
- A. CONCOLOR Linn. = *Sericocarpus concolor*.
- A. CONTYZOIDES Desf. = *Felicia angustifolia*.
- A. CONTYZOIDES Willd. = *Sericocarpus contyzoides*.
- A. CORDIFOLIUS Michx. = *Biotia corymbosa*.
- A. CORDIFOLIUS hortul. = *Biotia macrophylla*.
- A. CORIACEUS hortul. = *Diplostephium extenuatum*.
- A. CORNIFOLIUS Muhl. = *Diplostephium cornifolium*.
- A. CORYMBOSUS Ait. = *Biotia corymbosa*.
- A. CORYMBOSUS Moench. = *Inula germanica*.
- A. CRENATUS Less. = *Mairia crenata*.
- A. CRINITUS Linn. = *Athrixia heterophylla*.
- A. CRINITUS Thunb. = *Athrixia Capensis*.
- A. CRISPUS Forsk. = *Francoeuria crispa*.
- A. CYDONIAEFOLIUS Cunn. = *Eurybia cydoniaefolia*.
- A. CYMBALARIE Thunb. = *Munychia Cymbalarie*.
- A. DEBILIS Boj. = *Athrixia debilis*.
- A. DECURRENS Moench. = *Inula bifrons*.
- A. DECURRENS Cunn. = *Eurybia decurrens*.
- A. DENTATUS Thunb. = *Felicia tenella*.
- A. DENTATUS Andr. = *Olearia dentata*.
- A. DICHOTOMUS Ell. = *Diplostephium dichotomum*.
- A. DIVARICATUS Linn. = *Biotia macrophylla*.
- A. DIVARICATUS Spreng. = *Diplostephium amygdalinum*.
- A. DIFFUSUS ACUMINATUS Pers. = *Diplostephium acuminatum*.
- A. DRACUNCULOIDES Lam. = *Galatella dracunculoides*.
- A. DRACUNCULOIDES MINOR Ledeb. = *Linosyris glabrata*.
- A. DRACUNCULOIDES Bess. = *Galatella punctata*.
- A. DYSENTERICUS All. = *Pulicaria dysenterica*.
- A. ECHINATUS Less. = *Felicia echinata*.
- A. ECKLONIA Less. = *Agathæa Ecklonia*.
- A. ELATUS Bert. = *Biotia corymbosa*.
- A. ELLIPTICUS Cunn. = *Olearia elliptica*.
- A. ELÆAGNIFOLIUS Cunn. = *Eurybia furfuracea*.
- A. ELÆOPHYLLUS Cunn. = *Eurybia elæophylla*.
- A. ELONGATUS Thunb. = *Agathæa elongata*.
- A. ENSIFOLIUS All. = *Inula ensifolia*.
- A. ERICAFOLIUS Forsk. = *Athrixia ericaefolia*.
- A. ERUBESCENS Sieb. = *Eurybia erubescens*.
- A. EXASPERATUS Linn. = *Eurybia aculeata*.
- A. FASTIGIATUS Fisch. = *Turczaninovia fastigiata*.
- A. FASTIGIATUS Lehm. = *Galatella fastigiata*.
- A. FASTIGIATUS Ledeb. = *Galatella Hauptii*.
- A. FERRUGINEUS Wendl. = *Olearia dentata*.
- A. FILIFOLIUS Vent. = *Diplopappus filifolius*.

- A. *FILIFOLIUS* Willd. h. = *Felicia muricata*.
 A. *FLEXUOSUS* Nutt. = *Galatella flexuosa*.
 A. *FOLIOLOSUS* hortul. = *Eurybia aculeata*.
 A. *FRUTICOSUS* Linn. = *Diplopappus fruticulosus*.
 A. *FRUTICULOSUS* Burm. = *Felicia angustifolia* α.
 A. *FRUTICULOSUS* α Willd. = *Diplopappus fruticulosus*.
 A. *FRUTICULOSUS* β Willd. = *Diplopappus obtusatus*.
 A. *FRUTICULOSUS* Krebs. = *Diplopappus filifolius* β.
 A. *FUGAX* Brot. = *Galatella Lusitanica*.
 A. *FURFURACEUS* Rich. et Less. = *Eurybia furfuracea*.
 A. *GLANDULOSUS* Labill. = *Eurybia glandulosa*.
 A. *GLUTINOSUS* Bernh. = *Biotia glomerata*.
 A. *GLUTINOSUS* Roxb. = *Commidendron rugosum*.
 A. *GRACILENTUS* Banks. = *Chrysopsis falcata*.
 A. *GRANDIFLORUS* Walt. = *Tripolium paludosum*.
 A. *HAUPTII* Fisch. = *Galatella Hauptii*.
 A. *HETEROPHYLLUS* Thunb. = *Athrixia heterophylla*.
 A. *HIRTUS* Thunb. = *Felicia angustifolia* α.
 A. *HISPIDUS* Thunb. = *Heteropappus hispidus*.
 A. *HUMILIS* Ell. = *Diplostephium cornifolium*.
 A. *HUMILIS* Willd. = *Diplostephium amygdalinum*.
 A. *HYGROPHILUS* Cunn. = *Eurybia hygrophila*.
 A. *HYSSOPIFOLIA* Linn. = *Galatella hyssopifolia*.
 A. *HYSSOPIFOLIUS* Cav. = *Galatella punctata* α.
 A. *HYSSOPIFOLIUS* Berg. = *Felicia angustifolia* α.
 A. *ILICIFOLIUS* Cunn. = *Eurybia erubescens* β.
 A. *INCISUS* Fisch. = *Calimeris incisa*.
 A. *INDICUS* Linn. = *Asteromæa indica*.
 A. *INDICUS* Heyn. = *Callistephus Heyneanus*.
 A. *INFIRMUS* Michx. = *Diplostephium cornifolium*.
 A. *LANUGINOSUS* Wendl. = *Felicia*.
 A. *LATIFOLIUS* Desf. = *Biotia latifolia*.
 A. *LATIFOLIUS* Mill. = *Galatella rigida*.
 A. *LAVANDULACEUS* Willd. = *Diplostephium lavandulæfolium*.
 A. *LEDIFOLIUS* Cunn. = *Eurybia ledifolia*.
 A. *LEDIFOLIUS* Parsh. = *Eurybia ledifolia*.
 A. *LEPIDOPHYLLUS* Pers. = *Eurybia lepidophylla*.
 A. *LINARIFOLIUS* Linn. = *Diplopappus linarifolius*.
 A. *LINARIFOLIUS* Michx. = *Diplopappus rigidus*.
 A. *LINIFOLIUS* Burm. = *Senecio linifolius*.
 A. *LINIFOLIUS* Linn. = *Galatella linifolia*.
 A. *LINOINUS* hortul. = *Felicia Cafrorum* β.
 A. *LIRATUS* Sims. = *Eurybia lirata*.
 A. *LITHOSPERMIFOLIUS* Desf. = *Calimeris canescens*.
 A. *LONGICAULIS* Desf. = *Tripolium vulgare* α.
 A. *LONGIFOLIUS* Desf. = *Tripolium subulatum*.
 A. *LUSITANICUS* Brot. = *Galatella Lusitanica*.
 A. *MACROPHYLLUS* Linn. = *Biotia macrophylla*.

- A. *MACRORHIZUS* Cunn. = *Eurybiopsis macrorhiza*.
A. *MACROPHYLLUS* Raf. = *Biotia latifolia*.
A. *MACRORHIZUS* Thunb. = *Felicia macrorhiza*.
A. *MAGELLANICUS* Lam. = *Clarionea magellanica*.
A. *MAGELLANICUS* Spreng. = *Chilotrichium amelloideum*.
A. *MARYLANDICUS* Michx. = *Sericocarpus conyzoides*.
A. *MAXIMUS* Less. = *Erigeron maximum*.
A. *MESPILOFOLIUS* Less. = *Munychia? mespilifolia*.
A. *MICROPHYLLUS* Vent. = *Eurybia microphylla*.
A. *MICROPHYLLUS* Labill. = *Eurybia lepidophylla*.
A. *MONTANUS* All. = *Inula montana*.
A. *MORANENSIS* H. B. K. = *Diplostephium Moræense*.
A. *MONTICOLA* Lindl. = *Erigeron monticola*.
A. *MURICATUS* Thunb. = *Felicia muricata*.
A. *MYRSINOIDES* Labill. = *Eurybia myrsinoides*.
A. *NEMORALIS* Ait. = *Galatella nemoralis*.
A. *NEMORALIS* Willd. = *Galatella rigida*.
A. *NUDICAULIS* Lam. = *Lagenophora Commersonii*.
A. *OBOVATUS* Ell. = *Diplostephium obovatum*.
A. *OBTUSATUS* Thunb. = *Diplopappus obtusatus*.
A. *OBTUSATUS* Willd. = *Diplostephium obtusatum*.
A. *OCRACEUS* H. B. K. = *Tetramolopium ochraceum*.
A. *ODORUS* All. = *Pulicaria odora*.
A. *ODONTOPHYLLUS* Wall. = *Doronicum linifolium*.
A. *OFFICINALIS* All. = *Inula Helenium*.
A. *OPPOSITIFOLIUS* Linn. = *Munychia Cymbalaria?*
A. *PALUDOSUS* Ait. = *Heleastrum paludosum*.
A. *PALUSTRIS* Lam. = *Tripolium vulgare*.
A. *PANNONICUS* Jacq. = *Tripolium vulgare* f.
A. *PAPPUS* Willd. = *Printzia Bergii*.
A. *PAUCIFLORUS* Nutt. = *Tripolium pauciflorum*.
A. *PEDUNCULARIS* Wall. = *Amphirepis peduncularis*.
A. *PEREZIOIDES* Less. = *Mairia perezoides*.
A. *PERSOONIOIDES* Cunn. = *Eurybia persoonioides*.
A. *PHLOGOPAPPUS* Labill. = *Olearia phlogopappa*.
A. *PHLOGOTRICHUS* Spreng. = *Olearia phlogopappa*.
A. *PHYLICOIDES* H. B. K. = *Tetramolopium phyllicoides*.
A. *PIMELEOIDES* Cunn. = *Eurybia pimeleoides*.
A. *PINNATUS* Cav. = *Clomenocoma? pinnata*.
A. *PLANTAGINEUS* Pœpp. = *Senecio plantagineus*.
A. *POLIFOLIUS* Linn. = *Printzia Bergii*.
A. *PROPINQUUS* Cunn. = *Eurybia propinqua*.
A. *PUBESCENS* Moench. = *Inula squarrosa*.
A. *PULCHERRIMUS* Lodd. = *Diplopappus linarifolius*.
A. *PULCARIUS* Scop. = *Pulicaria vulgaris*.
A. *PULVERULENTUS* Hoffmans. = *Eurybia lirata*.
A. *PUMILUS* Fisch. = *Calimeris Altaica*.
A. *PUNCTATUS* W. et Kit. = *Galatella punctata* γ.

- A. PYRENAICUS Pourr. = *Jasonia steula* β.
 A. QUERCIFOLIUS Sieb. = *Olearia quercifolia*.
 A. RAMULOSUS Labill. = *Eurybia ramulosa*.
 A. REFLEXUS Michx. = *Felicia reflexa*.
 A. RIGIDUS Michx. = *Diplopappus linearifolius*.
 A. RIGIDUS Linn. = *Diplopappus rigidus*.
 A. RIGIDUS Moench. = *Inula salicina*.
 A. ROSRUS Stev. = *Calimeris rosea*.
 A. ROSMARINIFOLIUS Spreng. = *Diplopappus obtusatus*.
 A. ROTUNDIFOLIUS Thunb. = *Agathæa rotundifolia*.
 A. RUPESTRIS H. B. K. = *Tetramolopium rupestre*.
 A. RUTÆODORUS Cunn. = *Eurybia glandulosa*.
 A. SALICINUS All. = *Inula salicina*.
 A. SALINUS Schrad. = *Tripolium vulgare* δ.
 A. SATURNIÆFOLIUS Banks. = *Lorentea saturioides*.
 A. SCABER Thunb. = *Doellingeria scabra*.
 A. SCHREBERI Nees. = *Biotia Schreberi*.
 A. SCORPIOIDES Scop. = *Aronicum scorpioides*.
 A. SEDIFOLIUS Linn. = *Galatella punctata* α.
 A. SERRATUS Thunb. = *Agathæa serrata*.
 A. SERRATUS Lag. = *Grindelia inuloides*.
 A. SINUATUS Cunn. = *Olearia quercifolia*.
 A. SOLIDAGINEUS Michx. = *Sericocarpus solidaginæ*.
 A. SPATEULARIS Cav. = *Grindelia*.
 A. SPATHULATUS Cels. = *Inula salicina* β.
 A. SQUARROSUS All. = *Inula spireæfolia*.
 A. SQUARROSUS Banks. = *Diplopappus linearifolius*.
 A. STELLULATUS Labill. = *Olearia stellulata*.
 A. STRICTUS Wendl. = *Galatella dracunculoides*.
 A. STRIGOSUS Thunb. = *Amellus strigosus*.
 A. STRIGOSUS Licht. = *Agathæa strigosa*.
 A. SUBULATUS Michx. = *Tripolium subulatum*.
 A. SUBCYMOSUS Bernh. = *Biotia Schreberi*.
 A. TANACETIFOLIUS H. B. K. = *Machæranthera tanacetifolia*.
 A. TATARICUS Linn. f. = *Calimeris incisa* ?
 A. TAXIFOLIA Linn. = *Mairia taxifolia*.
 A. TENELLUS Linn. = *Felicia tenella*.
 A. TENERIMUS Less. = *Tetramolopium tenerimum*.
 A. TOMENTOSUS Schr. et Wendl. = *Olearia tomentosa*.
 A. TORTIFOLIUS Michx. = *Sericocarpus tortifolius*.
 A. TRINERVIS Deaf. = *Galatella rigida*.
 A. TRINERVIS α Nees. = *Galatella punctata*.
 A. TRINERVIS Roxb. = *Galatella asperifolia*.
 A. TRIPOLIUM Linn. = *Tripolium vulgare*.
 A. TRIPOLIUM Walt. = *Galatella flexuosa*.
 A. UMBELLATUS Ait. = *Diplostephium umbellatum*.
 A. UNDULATUS Moench. = *Pulicaria dysenterica*.
 A. UNIFLORUS Michx. = *Galatella nemoralis* β.

- A. verna** Linn. = *Stemactis verna*.
A. villosus Thunb. = *Felicia angustifolia* a.
A. viscosus Labill. = *Eurybia viscosa*.
A. Zeyheri Less. = *Felicia Zeyheri*.

CIX. TRIPOLIUM Nees ast. 152. Lindl. obs. mss. — *Aster Adams* fam. 2. p. 124. non Nees. — *Asteris* sp. Linn.

Capitulum radiatum, ligulis fœmineis uni-rarissimè biseriatis, disco hermaphr. Invol. squamæ jam secus pedunculum ortæ bracteiformes sensim aut imbricatæ aut subbiseriales, inter. longiores læves. Recept. alveolatum, alveolis dentatis. Styli fl. herm. rami apice crassiores subulati. Achænia oblonga compressa glabra aut sparsim pilosa basi setulis cincta. Pappus simplex pluriserialis, setis tenuibus remotè denticulatis. — Herbæ subcarnosæ perennes rariùs annuæ, in uliginosis salsuginosis crescentes, glabræ aut tantùm ad margines foliorum scabræ. Folia alterna angusta integra aut vix serrata. Capitula corymbosa. Ligulæ cœrulesc. aut albæ. — Genus recognoscendum, habitu plùs quàm caracteribus distinctum, hic juxta Lindleyi admonitiones dispositum.

§. 1. *Involucro subbiseriali, squamis omnibus obtusis, exterioribus brevioribus.* — *Tripolia vera*.

1. **T. vulgare** (Nees ast. 153), caulibus simplicibus ramoso-ve diffusis aut adscendentibus glabris, foliis subcarnosis lineari-lanceolatis margine subscabris basi ciliatis, radicalibus latioribus triplinerviis obtusis in petiolum attenuatis, ramis corymbosis, invol. squamis imbricatis interioribus maximis obtusis levibus. *W. Linn. Schrad.* ② Willd. In paludosis salsis Europæ totius. *Aster Tripolium* Linn. sp. 1226. fl. dan. t. 615. engl. bot. t. 87. DC. fl. fr. n. 3137. *Aster acris* Linn. herb. mon. cl. Lindley. *Aster palustris* Lam. fl. fr. 2. p. 143. Ligulæ lilacinæ, rariùs albæ. (v. v.)

β. *Pannonicum* (Nees ast. 153), foliis infer. lanceolatis basi attenuatis, caulinis angustis ciliato-serratis, caule adscendente aut erecto ramosissimo, invol. coloratis. — in paludosis subsalsis Hungariæ et Moraviæ. *Aster Pannonicus* Jacq. h. vind. 1. p. 3. t. 8. (v. v.)

γ. *diffusum*, foliis infer. lanceolatis basi attenuatis, caulinis angustis ciliato-serratis, caulibus à basi ramosissimis diffusis, invol. coloratis. — in paludosis salsis agri Chersonensis. *Aster Tripolium* Bieb. fl. taur. n. 1733. *Tripolium vulgare* *β*** Nees ast. 154? (v. s. comm. à cl. Beaupré).

δ. *salinum*, caule erecto subsimplici, foliis etiam infer. lanceolatis aut lineari-lanceolatis ciliato-serratis acutis, invol. colorato obtuso. ② in salsuginosis Sibirici et Dahuriæ. *Aster salinus* Schrad. hort. Goett. 1. t. 3. neu journ. 1810. st. 2. p. 138. *Aster Tripolium* Ledeb. fl. alt. 4. p. 99. — Gmel. fl. sib. t. 80. f. 2. Ligulæ cyaneæ paulò angustiores quàm priorum. An fortè spec. propr. ? (v. s.)

ε. *longicaule*, caule erecto subsimplici, foliis etiam infer. lineari-lanceolatis acutis margine ciliato-serratis, invol. squamis subcoloratis apice obtusiusculis. — in Hispaniæ salsis maritimis loco dicto La Rapila legit cl. L. Dufour! et rep. etiam in Græciâ (ex h. ac. Monac.). *Aster longicaulis* Dufour! in litt. 1824. Fol. sup. valdè acuminata. Pappus rufus. Vix à var. δ differt. (v. s.)

§. 2. *Invol. pluriseriali, squamis imbricatis, interioribus saltem aut omnibus acuminatis.* — *Oxytripolia*.

2. **T. angustum** (Lindl. adn. mss.), caule angusto strictissimo, ramis erectis ramosis, foliis linearibus serrulatis scabris glabris, invol. hemisphærici imbricati squamis linearibus acutis, ligulis inconspicuis. — in Amer. sept. A. T. subulato

differt caule nullo modo flexuoso, foliis serrulatis, squamis invol. capitalisque duplò majoribus. (Lindl.)

α. ramis oligocephalis. — in montibus saxosis.

β. ramis polycephalis. — ad lacum Slave dictum.

3. *T. FLEXUOSUM* (Nees ast. p. 155), glabrum, caule erecto tereti flexuoso, foliis linearibus subulatis crassis lævibus impunctatis, ramis 1-cephalis squamigeris, invol. squamis imbricatis lanceolatis acutis, infer. subpatulis, ligulis invol. circiter aequalibus, achenio glabrescente. 2ℓ in paludibus salis Amer. bor. à Novâ-Cassara ad Noveboracum (Nutt.), in Carolinâ (Nutt. Ell.). Aster Tripolium Walt. car. 210 non L. A. sparsiflorus Pursh fl. 2. p. 547 non Michx. Ast. flexuosus Nutt. gen. am. 2. p. 145 non Fisch. Foliis carnosius accedit ad *Tr. vulgare*, sed invol. regulariter imbricatis ligulisque brevissimis facile differt. Folia infer. trinervia. (v. s.)

4. *T. ? PAUCIFLORUM* (Nees ast. 167. Lindl. obs. mss. 1835), caule humili simplici 3-6-cephalo, ramis 1-cephalis viscido-pubescentibus, foliis linearibus subcarnosis glabris, caulibus subulatis amplexicaulibus, invol. hemisphaerici squamis subæqualibus acutis. 2ℓ ad margines fontium aquæ salis prope castellum Mandam ad Missouri fluvium. Aster pauciflorus Nutt. gen. am. 2. p. 154. Habitus Tripolii sed caractere diversus, potius fortè ad *Asteres* alpigeros referendus ex Lindl. adn. mss. 1834.

5. *T. ? SUBULATUM* (Nees ast. 157. Lindl. adn. mss.), caule erecto racemoso-decomposito, foliis linearibus acuminatis integerrimis, squamis ferè subulatis, panicula laxa, invol. squamis imbricatis linearibus acuminatis exter. laxiusculis, ligulis biserialibus vix involucre longioribus. ① in maritimis paludosis Americæ. Achenia compressa oblonga puberula. Pappus 1-serialis. Recept. nudum. Invol. demum reflexum. Species in orbe novo dispersa et polymorpha, nempe:

* *boreale*; à Pensylvaniâ ad Carolinam. Aster subulatus Michx. fl. bor. am. 2. p. 111. Aster longifolius Desf. cat. h. par. ed. 3. p. 400. Poir. dict. 7. p. 480. (v. s. ex Carol. comm.)

** *Mexicanum*. — circa urbem Matamoros legit Berlandier pl. exs. n. 2339. (v. s.)

*** *Cubense*. — in insulâ Cubâ (Poepp. La Sagra!), *Erigeron* expansus Spr. syst. 3. p. 518. (v. s.)

**** *Domingense*. — in Sancto-Domingo ad flum. Yague. *Erig. linifolium* Bert. herb. (v. s.)

***** *Brasilianum*. — in Brasiliæ prov. Rio-Grande. (h. Mus. Par. n. 859) et in Bandâ orientale (Bacle!). (v. s.)

***** — in ins. O-Wahu (Cham.). Aster subulatus Less. in linn. 1831. p. 120. Capitula paulò minora. (v. s.)

6. *T. CONSPECTUM* (Lindl. obs. mss. 1835), caule erecto racemoso, foliis linearibus acuminatis canaliculatis, ramis distantibus subæquilongis 3-6-cephalis rarius decompositis, bracteis ramorum et pedicellorum acutis cucullatis squamiformibus, invol. turbinato, squamis interioribus basi pallidis apice herbaceis vel coloratis acutis. — in Chili legit cl. Bridges pl. exs. n. 183. Caulis bipedalis glaber, internodiis inflorescentiæ ramis longioribus vel parum brevioribus. — Specim. Bridg. non vidi. Huc verisim. referenda *Baccharis asteroides* Bert. n. 83 et 828, in pascuis et hortis circa Rancagua et Valparaiso lecta. (v. s.)

† *Species exclusa.*

T. PALUDOSUM Nees. = *Döllingera paludosa*.

CX. *GALATELLA* Cass. dict. 37 (1817) p. 463 et 488. Nees ast. 158. Lindl. adn. mss. — *Galatea* Cass. dict. 18. p. 56. bull. philom. 1818. p. 165. Less. syn. 187. — *Asteris* sp. auct.

Capitulum radiatum, ligulis 1-serialibus neutris aut organa foeminea abortiva gerentibus, fl. disci hermaphroditis tubulosis fertilibus. Invol.

disco brevius squamis imbricatis inappendiculatis, exter. sæpè trinerviis. Recept. alveolatum, alv. marginibus dentatis. Styli fl. herm. rami apice spatulati obtusi. Achænia densè hirsuta vel strigoso-sericea. Pappus pluriserialis, setis filiformibus rigidis scabridis. — Herbæ perennes per hemisphærium boreale sparsæ. Caulis simplex apice corymbosus. Folia alterna integra oblonga aut linearia omnia basi saltem trinervia sessilia, sæpè punctata, punctis in superficie superâ præsertim obviis impressis subglandulosis. Capitula ferè Asteris sed ligulæ neutræ distantes et invol. breve. Discus flavus. Ligulæ cœrulæ purpurascens aut albæ.

* *Foliis punctatis.*

1. *G. LIMIFOLIA* (Nees ast. p. 170), caule stricto tereti apice corymboso, foliis linearibus integerrimis utrinquè punctatis subscabris 1-nerviis, ramulis 1-cephalis folia lineari-subulata parva gerentibus, invol. squamis lanceolatis acutis margine submembranaceis disco brevioribus, ligulis paucis discum non superantibus. $\frac{1}{2}$ in Amer. bor. umbrosis siccis à Virginiâ ad Pensylvaniam. *Aster limifolius* Linn. sp. 1228. *Galatella albiflora* Cass. dict. 18. p. 58. *Chrysopsis limifolia* Nutt. gen. am. 2. p. 152. Ligulæ steriles albæ. Cl. Nees qui optimè speciem in syn. ast. p. 18 deminavit, incautè in ast. folia impunctata dixit. (v. s.)
2. *G. BYSSOPIFOLIA* (Nees ast. p. 160), caule stricto apice coarctato-corymboso, ramis patulo-corymbosis, foliis lanceolato-linearibus acutis integerrimis trinerviis punctatis scabris, ramulorum lineari-subulatis, invol. squamis omnibus acutiusculis, ligulis elongatis. $\frac{1}{2}$ in campis sabulosis et sylvis à Novâ-Cæsareâ ad Carolinam. *Aster byssopifolius* Linn. mant. 114 (non Cav.) et ejus herb. ex Lindl. ! Ell. sketch 2. p. 342. Ligulæ albæ purpuro tinctæ, 5 ex Ell., 10 in specim. mihi obvis. (v. s. comm. à cl. Lindl.)
3. *G. PUNCTATA*, caule erecto tereti scabrido simplici aut apice corymboso, foliis linearibus aut lineari-lanceolatis trinerviis aut summis 1-nerviis punctatis subscabris, ramulis folia parva lineari-subulata gerentibus, invol. obconici squamis lanceolatis acutis margine membranaceis disco brevioribus, ligulis invol. duplò circ. longioribus. $\frac{1}{2}$ in Europâ ab Hispaniâ ad Caucasum. Hæc mihi videtur verus *Aster acris* Linn. (conf. fl. fr. n. 3138 et suppl. p. 469) ex syn. loco natali et acretudine, sed cum altera species sit in ejus herb. servata et plurimi Botanici nomen ad alter. spec. transtulerint, hoc omninò rejicendum potius videtur, ne confusio augeatur. Staud. cl. Lindley huc trahi var. sequentes :
 - a. *angustifolia*, foliis omnibus linearibus angustis. — in apricis Hispaniæ, Galliæ-Mediterranæ, Italiæ. *A. acris* Linn. sp. ed. 2. p. 122 (excl. syn. et patr. Hungaria) non Willd. *A. byssopifolius* Cav. ic. 3. p. 17. t. 232 excl. syn. *A. sedifolius* Linn. sp. ed. 1. p. 874. *Chrysocoma Linosyris* Asso ! fl. arrag. p. 126 excl. syn. — Lob. ic. t. 349. f. 2. Barr. ic. t. 606. Simillima *G. limifoliæ*. Ligulæ lilacinæ. (v. v. et s.)
 - b. *intermedia*, foliis lineari-lanceolatis basi trinerviis, rameis linearibus 1-nerviis. — in Italiæ et Galliæ anstr. collibus. *Aster acris* hort. Par. Reich. fl. exc. 1. p. 246. *Galatea intermedia* Cass. dict. 18. p. 58. *Galat. punctata* Nees ast. p. 161. — Garid. Aix t. 11. Pluk. alm. t. 271. f. 3. Videtur omninò eadem in solo pigniori orta. (v. s. et v.)
 - γ. *insculpta*, foliis latius lineari-lanceolatis ferè omnibus evidentius trinerviis. — in salis subuliginosis Hungariæ (W. et K. Graaf!), Podoliæ anstr. (Bess. !), Tauriæ (Bieh.), Caucasi (Mey. !), Persici prov. Karabagh (Scow. !). *Aster punctatus* Waldst. et Kit. hung. 2. t. 109. Willd. sp. 3. p. 2022. excl. syn. Cav. *Aster dracunculoides* Bess. ! enum. 33. *Galatella punctata* Cass. dict. 18. p. 57. *Galatella insculpta* Nees ast. p. 162. *Aster trinervius* var. α Nees syn. 19 (v. s. sp. et v. c.)
4. *G. CANA* (Nees ast. 163), tota lanugine molli caulescens, caule erecto à mediò

- ramoso-corymboso, foliis lanceolato-oblongis mucronatis integerrimis trinerviis punctatis, ramis subcorymbosis virgatis, invol. squamis 4-seriatis ovato-lanceolatis acutis vix margine membranaceis, ligulis disco forè duplò longioribus. $\frac{1}{2}$ in fruticetis Banatùs Hungariæ. Aster canus W. et K. r. p. 28. t. 30. Gal. canescens Cass. dict. 18. p. 57. Ligulæ lilacinæ 8-10. Achænia albo-lanata. (v. s. sp. et v. c.)
5. *G. DAKURICA*, caule glabrescentulo subangulato erecto apice corymboso, ramis foliosis, foliis lineari-lanceolatis acuminatis trinerviis margine scabris punctulatis, invol. hemisphærici squamis pluriseriatis imbricatis acuminatis subciliatis disco paulò brevioribus, ligulis 20-30 elongatis. $\frac{1}{2}$ in sabulosis Dahuriæ ad ripas Iræuti flum. detexit cl. Turczaninow. Aster Hauptii Turcz. l in litt. non alior. Folia 3 lin. lata. Achænia villosa. Ligulæ purpurascens, ab omnibus prioribus monente cl. Lindl. diff. invol. hemisphærico nec obconico. (v. s.)
6. *G. DRACUNCULOIDES*, caule stricto tereti puberulo apice breviter densèque corymboso, foliis lineari-lanceolatis mucrone calloso apiculatis integerrimis trinerviis utrinquè punctatis subtus scabridis, ramealibus angustis subfalcatis, ramulis di- aut ramoso-polycephalis, invol. squamis margine scariosis, ext. acutis int. obtusis, ligulis paucis disco duplò longioribus aut nullis. $\frac{1}{2}$ in campestribus apricis Sibirici, Taurici, Caspici. Aster dracunculoides Ledeb. fl. alt. 4. p. 97.
- a. *radiata*, ligulis paucis. — Aster dracunculoides Lam. dict. 1. p. 303. Aster acris Nees syn. 19. Galatella pauciflora Nees ast. p. 164. Ast. strictus Wend. act. soc. Marb. 2. f. 6. p. 54 ex Nees. Ligulæ cœruleæ lineares. (v. s.)
- β. *discoidea*, ligulis nullis. — Chrysocoma biflora Linn. sp. 1178. Chrysocoma dracunculoides Lam. dict. 2. p. 192. Crinitaria biflora Cass. dict. 37. p. 476. Galatella biflora Nees ast. p. 159. Aster bifidus Nees syn. 18. — Gmel. sib. 2. p. 189. t. 82. f. 1. Cl. Ledebour monet ligulas in iisdem individuis nunc paucas nunc nullas.

** *Foliis impunctatis.*

7. *G. RIGIDA* (Cass. dict. 18. p. 58. sub Galatella), caule erecto subtriato puberulo apice corymboso, foliis lanceolato-linearibus mucronatis acutis integerrimis trinerviis impunctatis subtus margineque scabris, ramis floris subtrifidis pedicellisque rigidis, invol. squamis lanceolatis subacutis margine membranaceis, ligulis disco longioribus. $\frac{1}{2}$ in vineis circa Meldas Occitanorum legit cl. Prost. Aster acris Willd. sp. 3. p. 2023. non Linn. Gall. acris Nees ast. 171. Aster trinervis hort. Par. l Nees syn. 18. Aster acris β Lam. dict. 1. p. 303. excl. ubiquè synon. hic non citatis. Ligulæ cœruleæ. Herba junior subarachnoidea. (v. v. c.)
8. *G. LEPTOPHYLLA* (Nees ast. 167), caule angulato apice simpliciter corymboso juniore lanuginoso, foliis lanceolato-linearibus integerrimis acutis trinerviis carinatis impunctatis, juvenilibus ad costas marginesque sublanuginosis demùm scabris, ramis gracilibus minutè foliosis, invol. squamis ovatis obtusis ciliatis, ligulis disco longioribus, achæniis hirtis. — in Americâ septentrionali? Ligulæ saturatè cœruleæ. Aster leptophyllus à cl. Desfontaines ipso mecum comm. est diversissimus, nec in Sprengelii opere etiam à Neesio citatus reperitur. Fortè eadem ac *G. rigida*. Species valdè obscura. †.
9. *G. HAUPTII* (Lindl. adn. mss.), foliis oblongo-linearibus acuminatis trinerviis margine scabris, caule basi subsimplici superne in ramos corymboso-fastigiatus monocephalos densè foliosos diviso, invol. turbinati squamis pluriseriatis, exter. linearibus acuminatis, intermed. latioribus acutis, intimis membranaceis rubescentibus, omnibus costâ fuscâ notatis. $\frac{1}{2}$ in Sibirici altaicæ collibus et locis apricis frequens. Aster fastigiatus Ledeb. ic. fl. ross. alt. t. 161. non Fisch. nec. Lehmann. Aster Hauptii Ledeb. fl. alt. 4. p. 100.
10. *G. FASTIGIATA* (Nees ast. 169), foliis angustè lanceolatis integerrimis subulato-acuminatis trinerviis planis impunctatis ad costam subtus marginesque

scabris, caule apice corymboso-composito, ramis ramulisque gracilibus patentibus apice minutè foliolosis, invol. squamis ovatis obtusiusculis ciliatis. $\frac{2}{2}$ in Sibiria. Aster fastigiatus Lehm. mss. ex Nees.

11. *G. nemoralis* (Nees ast. 173), caule puberulo scabro apice corymboso, foliis lanceolato-linearibus integerrimis enerviis impunctatis scabris, invol. squamis linearibus acutis pauciseriatis, ligulis elongatis. — in sphagnosis Amer. bor. à Canadâ ad Novam Cesaream. Aster nemoralis Ait. kew. v. 3. p. 198. Aster ledifolius Pursh fl. 2. p. 544.

ß. monocephalus. — Aster uniflorus Michx. fl. 2. p. 110. ex Nees et Lindl. qui specim. auth. vid. (v. s.)

12. *G. squamosa*, glabra, caule apice angulato, foliis lineari-lanceolatis acuminatis trinerviis margine et sæpè costâ mediâ serrato-scabris impunctatis, summis lineari-subulatis in squamas parvas subulatas secus pedunculum desinentibus, invol. imbricatis squamis infer. acuminatissimis, intima. longioribus margine membranaceis lanceolatis minùs acuminatis. — in Sibiria jugo Altaico et circa Barnaul legit cl. Patrin. Variat caule simplici et ramoso, ramis paucis 1-cephalis. Ligulæ amplæ cœruleæ. Styli rami acuminati longè exserti. (v. s. comm. ab amic. Delessert.)

13. *G. tenuifolia* (Lindl. ! adn. mss.), caule erecto tenui glabro simplici 1-cephalo ant apice subcorymboso, foliis linearibus glabris vix margine subscabris uninerviis aut infimis trinerviis, ramulis folia subulata parva subimbricata gerentibus, invol. squamis lanceolatis acutis subciliatis siccis, ligulis elongatis. $\frac{2}{2}$ in Sibiria (h. Lindl.!), in regione Kirghisorum (Morici!). Aster tenuifolius Linn. herb. ex Lindl. ! Aster leptophyllus Fisch. ex Lindl., sed non est Aster homonymus à Neesio citatus. Valde accedit ad priorem sed folia vix 1-1 $\frac{1}{2}$ lin. lata fere glabra. Styli rami minùs exserti obtusiusculi. (v. s.)

14. *G. juncea* (Lindl. ! adn. mss.), caule stricto junceo subfastigiato, ramulis subaphyllis 1-cephalis, foliis linearibus integerrimis enerviis impunctatis laeviusculis, invol. hemisphærici squamis linearibus acutis disco vix brevioribus. $\frac{2}{2}$? in India septentrionali legit cl. Royle sub n. 139. (v. s.)

15. *G. nudiflora* (Lindl. adn. mss.), caule corymboso, ramulis elongatis aphyllis, foliis lineari-lanceolatis acuminatis margine scabris, involucrio brevissimo, ligulis invol. triplò longioribus. $\frac{2}{2}$ in Sibiria. Amman. Ramulis longè aphyllis primo aspectu cognoscitur. Coet. mihi ign.

† *Species non satis notæ (fortè, mon. cl. Lindl., ad Asteres alpinos rejiciendæ?)*

16. *G. ? lusitanica* (Nees ast. 165), caule erecto pubescente 1-3-cephalo, foliis glabriusculis, radicalibus spathulatis apice dentatis, caulinis linearibus integerrimis punctatis suprâ bisulcis, invol. squamis laxè imbricatis lanceolatis disco subæquantibus, ligulis lanceolatis disco superantibus. $\frac{2}{2}$ in dumetis Lusitanie. Aster fugax Brot. fl. lus. 1. p. 385. Aster lusitanicus Brot. phyt. 63. t. 29. Aster alpinus γ fugax Poir. suppl. 1. p. 489. Ligulæ cœruleæ aut purpureæ sæpè steriles. Achæmia subvillosa. †

17. *G. ? arragoneensis* (Nees ast. 167), caule erecto apice corymboso oligocephalo lanato, foliis radicalibus spathulatis integerrimis, caulinis linearibus acutis subtrâ bisulcis, invol. squamis imbricatis disco subæquantibus, ligulis elongatis. $\frac{2}{2}$ in Arragonia. Aster Arragoneensis Asso fl. arr. 121. t. 8. f. 2. Lam. dict. 1. p. 302. ill. t. 681. f. 5. Ligulæ cœruleæ. †

†† *Species exclusa.*

G. glandulosa Nees. = *Eurybia glandulosa*.

CXI. TURCZANINOVIA.

Capitulum pluriflorum radiatum, ligulis 7-8 uniserialibus fœminis

fertilibus integris apice obliquis, disci 7-8 tubulosi 5-dentatis hermaproditis fertilibus paucisque sterilibus mixti. Recept. alveolatum. Invol. squamæ circ. biseriales lineares obtusiusculæ disco multò breviores. Stigm. radii elongata, disci brevia inclusa. Achænia compressa glabra colorata. Pappus 1-serialis setosus, setis sericeis subscabris. — Herba Sibirica aut Dahurica simplex erecta apice corymboso-fastigiata, basi glabra, apice pubescenti-scabrida. Folia alterna lineari-lanceolata integerrima acuta uninervia. Corymbus compositus fastigiatus polycephalus. Ligulæ albæ. Discus flavus. — Habitus Galatellæ sed ligulæ fertiles et achænia glaberrima. Caract. ferè Asteris sed involucrum disco multò brevius pauci-seriale, squamis non foliaceis, etc. — Dicavi cl. N. Turczaninof, qui Dahuriæ regionumque vicinarum botanicam egregiè illustrat et mecum hanc ipsam stirpem cum pluribus aliis rarissimis communicavit.

1. *T. FASTIGIATA*. (2) sen 2 in Sibiria (Fisch.), in Dahuriæ paludosis ad flum. Argunum (Turcz.). Aster fastigiatus Fisch. mem. soc. mosc. 3. p. 74 non Ledeb. (v. s. comm. à cl. Turczaninof.)

CXII. *CALIMERIS* Nees ast. p. 225. — *Kalimeris* Cass. dict. 24. p. 324. 37. p. 464 et 491. — *Asteris* spec. auct.

Capitulum radiatum, radio foemineo fertili, ligulis 1-serialibus latiusculis, disco hermaphr. fertili. Stigm. apice conico lanceolata angustiora. Invol. squamæ 2-4-seriales subæquales herbaceæ margine membranaceæ laxè imbricatæ. Recept. convexi vel conici alveolæ subquadratæ ad angulos obtusè dentatæ. Achænia compressa marginata hirta. Pappus sub-1-serialis setaceus inæqualis scaber. — Herbæ perennes Asiaticæ. Caulis angulatus apice corymbosus. Folia integerrima vel inciso-dentata. Discus flavus. Ligulæ albæ seu cœruleæ.

2. *C. FLEXUOSA* (Royle! in Lindl. adn. mas.), caule flexuoso glabro (aut hirsuto-scabro), foliis ovato-lanceolatis basi attenuatis apice acuminatis grossè serratis suprâ et margine setoso-scabris subtilius pallidis secus nervos vix scabris, invol. squamis lanceolatis biserialis margine hyalino-scariosis, achæniis sparè hirtellis. 2 in Indiæ orient. prov. Mussonæ legit cl. Royle pl. exa. n. 113. Ligulæ cyanæ lineares obtusæ. Pappus subrufescens corollulæ ferè æqualis. (v. s. comm. à cl. Royle.)

3. *C. INCISA*, caule erecto striato secus angulos scabro aut (in pl. cultâ) lævi apice corymboso, foliis lanceolatis utrinquè attenuatis remotè inciso-serratis margine subtusque scabris, invol. squamis apice rufo-marginatis, pappo cor. tubo duplò breviorè inæquali. 2 in Sibiria (Nees) et Dahuriæ (Turcz.) humidis. Aster incisus Fisch. mem. soc. mosc. 3. p. 76 (1812). *Grindelia incisa* Spreng. syst. 3. p. 575. *Kalimeris platycephala* Cass. dict. 24 (1822). p. 325. Nees ast. 226. Caulis 1-2-pedalis. Ligulæ cœruleæ 6 lin. longæ, lineam latæ. Variat ex Lindl. adn. mas., invol. squamis immarginatis. (v. s. et v. c.)

3. *C. ALTAICA* (Nees ast. 228. Lindl. adn. mas.), caule pubescenti-scabro sen hispido irregulariter corymboso, foliis super. lineari-lanceolatis integerrimis ciliatis, capitulis paucis subcorymbosis, invol. pubescentis squamis 3-4-seriatis lanceolatis acuminatis albo-marginatis, pappo piloso corollâ breviorè. 2 in montibus Sibiria Altaica (Ledeb.), Tanaica? (Fisch.), ad flum. Schersee et in campis Barahensibus (Pall. ex Lindl. mas.). Aster pumilus Fisch. ! cat. h. Gor. p. 62. *A. Altaicus* Willd. enum. p. 881. Ledeb. ! fl. alt. 4. p. 99. qui fortè à seq. non separat. Ligulæ cœruleæ angustissimæ. Pappus subviolaceus sericens. Folia var. longiora et obtusiora. (v. s. comm. à cl. Fischer qui etiam ad Willd. misit.)

4. *C. CANESCENS* (Nees ast. 229), pubescenti-cana, caule erecto irregulariter corymboso infernè fasciculato-folioso, foliis lanceolato-linearibus obtusis integerrimis variis suberratis, ramulis foliosis, capitulis solitariis, invol. squamis glabresculis lanceolatis acuminatis albo-marginatis, pappo piloso corollâ paulò breviorè. \mathcal{L} in Sibiria Altaicâ (Gebl. ex Lindl. mss.) ad Sagai (Fisch.), in arenosis ad Selengam (Pall. ex Lindl.). Aster Gebleri Bess. in litt. A. lithospermifolius hort. Par. 1821. Aster Altaicus plur. auct. et fortè à Cal. Altaicâ vix satis differt. Capitula paulò majora. Ligulæ latiores. (v. s.)
- γ ? *latifolia* (Lindl. adn. mss.), caule erectiusculo scabrido-hispidulo, foliis caulinis oblongis retusis, ligulis latioribus. \mathcal{L} in India orient. ad Kunawur legit cl. Royle (pl. exs. n. 103). Folia 8-10 lin. longa, 2-3 lin. lata, distantia. (v. s. comm. à cl. Royle.)
5. *C. TATARICA* (Lindl. adn. mss.), caule corymboso scabritusculo, foliis linearibus acutis pubescentibus margine ciliato-scabris, ramis patentibus sæpius simplicibus 1-cephalis, invol. squamis acuminatissimis hirsutis herbaceis (nec albo-marginatis). \mathcal{L} in Dahuriâ (Fisch. Presc. ex Lindl.), in arenosis Mongoliæ Chinesis (Turcz.). Ligulæ azureæ 7-8 lin. longæ. (v. s. comm. à cl. Turcz.)
6. *C. HISPIDA* (Lindl. adn. mss. non Nees), caule stricto apice composito-ramoso foliisque linearibus acutis integerrimis hirsutis, invol. hispidissimo squamis acuminatis, exterioribus squarrosis. \mathcal{L} ? in Sibiria (Gmel. ex Lindl.). Calimeris hispida Nees est *Heteropappi* sp. Capitula magnitudine et facie *C. Altaicæ*. Invol. squamæ interiores vix membranaceæ.
7. *C. EXILIS* (Lindl. adn. mss.), caulibus caespitosis simplicibus 1-cephalis corymbosive pubescentibus, foliis adpressis linearibus acutis pubescenti-scabris incanis sæpius in invol. squamis laxas angustas subhispidas abeuntibus. \mathcal{L} in Sibiria C. Altaicæ proxima; foliis acutis, invol. squamis acuminatis capitalisque minimis, radio breviori diversa. (Lindl.)
8. *C. INTEGRIFOLIA* (Turcz. in litt. 1834), caule erecto simplici apice corymboso, ramis 5-7 foliosis 1-cephalis, foliis oblongo-linearibus sessilibus acutis integerrimis canleque pube brevi pulverulento-velutinis, invol. squamis laxis glabresculis acuminatis, ligulis ovali-oblongis. ① ②? in Sibiria ad flumen Amur legit cl. Turczaninof et ad flum. Argunum cl. Razgildeef, etiam à Chinâ in herb. Staunton video. Ligulæ cœruleæ latiores et breviores quàm *C. Altaicæ*. (v. s.)
9. *C. ROSA*, glabrescula basi suffruticosa, caulibus erectis ramosissimis, foliis linearibus integerrimis acutis, ramis 1-cephalis, invol. squamis circ. biserialis lanceolatis acuminatis margine membranaceis ciliatis, achænio sericeo-villoso, pappo piloso. \mathcal{L} in Caucasii orientalis monte Schadagh et in saxis circa Schudur legit cl. Steven. Ast. roseus Stev. act. mosc. 3. p. 245. et 4. p. 64. Bieb. suppl. n. 1727. Nees ast. p. 29 non hort. Par. Caulis 8-9 poll. longus. Folia ferè 2 poll. longa, sesquilineam lata. Capit. Ast. alpino vix minùs. Ligulæ 5 lin. longæ, lineam latæ. (v. s. comm. à cl. inv.)
10. *C. ALYSOIDES*, fruticulosa, ramis foliosis adpressè cano-tomentosis, foliis oblongis integerrimis araneoso-tomentosis subvis canis, infer. obtusis, cæteris mucrone acutis, capitulis solitariis subcorymbosis, invol. squamis oblongis acuminatis subaraneosis erectis margine subscariosis uninerviis, achæniis puberis, pappo piloso. \mathcal{L} in Mongoliâ Chinesis locis subarenosis detexit cl. Turczaninof. Aster alyssoides Turcz. chin. n. 21. Ligulæ albæ 5-7, 3-4 lin. longæ, lineam latæ. Folia 3-4 lin. longa, vix ultra lineam lata. (v. s. comm. à cl. inv.)

† *Species exclusæ.*

C. BIENNIS Nees. = *Aster canescens*.

C. HISPIDA Nees. = *Heteropappus hispidus*.

CXIII. EURYBIOPSIS.

Capitulum pluriflorum radiatum; ligulis foemineis 1-serialibus, fl. disci tubulosis 5-dent. hermaphr. Invol. squamæ pauciseriales imbricatæ lineares acuminatæ. Recept. angustum alveolatum. Achænia elongata compressa basi subangustata hispidula. Pappus 1-serialis multisetus, setis scabridis acuminatis. — Herba Australasica glabra multicaulis, rhizomate crasso lignoso, caulibus tenuibus virgatis parçè foliosis demùm frutescentibus. Folia alterna integerrima, radicalia lineari-oblonga obtusa, caulina linearia subacuta. Capitula ad apices caulium solitaria. — Fructus compressus Asteris, cætera Eurybiæ.

1. *E. MACROBYTZA*. $\frac{1}{2}$ $\frac{3}{4}$ in Australiæ orâ boreali occid. ad radices collitum prope Prince Regent's rivier, octob. flor. legit cl. A. Cunningham. Aster? macrobyzus A. Cunn. in litt. 1834 non Thunb. (v. s. comm. à cl. invent.)

CXIV. *PODOCOMA* Cass. bull. philom. 1817. p. 137. dict. 37. p. 462 et 484. 42. p. 60. Less. syn. 187. — *Erigerontis* sp. Poir.

Capitulum radiatum, ligulis foemineis 1-serialibus (Cass.) seu 1-2-ser. (Less.), disco hermaphrodito regulari. Recept. nudum planum. Invol. irregul. imbricatum disco brevius. Achæmium oblongum compressum rostratum. Pappus pilosus pluriserialis conformis. — Herba perennes multicaules Bonarienses. Folia infima obovata dentata pubescentia, caulina sparsa oblongo-lanceolata. Capitula ramulos terminantia subcorymbosa. Ligulæ lineares albæ (Less.), forsan flavæ (Cass.). Pappus rufus. — Plures ex Less. adsunt in herbariis species indescriptæ.

1. *P. HIERACIFOLIA* (Cass. dict. 42. p. 60), capitulis subcorymbosis. $\frac{1}{2}$ in Bonariâ (Comm.) *Erigeron hieracifolium* Poir. dict. 8. p. 491.
2. *P. PRIMULÆFOLIA* (Cass. l. c. p. 61), capitulis majoribus solitariis. In Monte-Video. †.

CXV. ASTEROPSIS Less. syn. p. 188.

Capitulum radiatum, ligulis foemineis 1-serialibus, disco hermaphrodito regulari. Recept. epaleaceum. Invol. pauciserialæ. Achæmium breviter rostratum. Pappus pilosus 1-serialis conformis. — Herba Brasiliensis. Folia alterna integra. Capitula solitaria.

1. *A. MACROCEPHALA* (Less. l. c.). — in Brasiliâ (Sellow). Cæt. ign.

CXVI. ARCTOGERON. — *Erigerontis* sp. Linn. — *Asteris* sp. Gmel. Ann.

Capitulum radiatum, ligulis foemineis 1-serialibus, disco hermaphr. fertili. Recept. angustum planum subalveolatum. Invol. imbricatum squamis 3-serialibus adpressis lanceolatis acuminatis carinâ viridibus et puberulis margine albo-scariosis. Ligulæ ovali-oblongæ apice dentatæ invol. duplò longiores. Stigm. disci et radii crassa brevia. Achæmium oblongum subcompressum sericeo-villosissimum. Pappus simplex pluriserialis persistens, setis subinæqualibus scabris. — Herba Sibirica habitu Armeriæ, rhizomate caudiciformi frutescente multicipiti brevi. Folia radicalia conferta lineari-subulata margine scabro-ciliolata cæterum glabra rigida erecta pollicaria semilineam lata. Scapi vix foliis longiores hirsuti monocephali. Discus flavus. Radius albus. — Nomen ex *αρκτος*, septentrion, et *γενει*, seuex, quasi *Erigeron boreale*.

1. *A. GRAMINEUM*. $\frac{1}{2}$ in campis sabulosis Sibiriae Transbaikalisensis (Gmel.) et Dahuriae (Fisch.), ad Tchita (Patr.), frequens in desertis Selenge, Udm., Ononis et Arguni fl. adjacentibus (Amm.). *Erigeron gramineus* Linn. sp. 1212. Lam. ill. t. 681. f. 4. Poir. dict. 7. p. 490. — Gmel. fl. sib. 2. p. 174. t. 76. f. 2. — Amm. ruth. 215. *Erig. gramineus* est omnino diversus. (v. s. ex Selenga et Tchita à cl. Turcz. et Patr. coll.)

CXVII. *SERICOCARPUS* Nees ast. p. 148. — Asteris sect. 1. Ell. sketch 2. p. 340. — Asteris sp. auct.

Capitulum radiatum, ligulis 1-serialibus paucis latiusculis foemineis fertilibus, disco hermaphrodito paucifloro fertili. Stigmata apice subulata. Invol. imbricatum ovatum vel oblongum, squamis multiseriis latis basi cartilagineis enerviis apice herbaceis. Recept. alveolae saepius lacero-ciliatae. Achænia obpyramidata brevia dense sericeo-strigosa. Pappus simplex pauciserialis rigidus scaber. — Herbae perennes ex America boreali ortae. Caulis apice corymbosus. Folia impunctata integerrima basi attenuata. Capitula ad apices ramorum adgregata, conyzoides. Ligulae albæ.

1. *S. SOLIDAGINEUS* (Nees ast. 149. Lindl. ! adn. mss.), caule erecto tereti nervis prominulis subangulato glabro, foliis lineari-lanceolatis basi attenuatis apice obtusis margine scabris obscure trinerviis glabris, corymbo fastigiato, invol. oblongi squamis adpressis basi pallidis apice obtusis viridi-callosis, ligulis 5 linearibus invol. duplò longioribus. $\frac{1}{2}$ in pinguibus paludosis et marginibus sylvarum Amer. boreal. à Pennsylvania ad Carolinam. *Conyza linifolia* Linn. sp. 1205. *Aster solidaginoides* Willd. sp. 3. p. 2024. Ell. sketch 2. p. 340 non Raf. *Aster solidaginus* Michx. fl. bor. am. 2. p. 108. — Pluk. alm. t. 79. f. 2. bene. (v. s. olim ex sem. à Michx. missis sub nom. *A. corymbosi*.)
2. *S. CONYZOIDES* (Nees ast. 150), caule erecto subangulato puberulo, foliis cuneato-oblongis subtrinerviis apice serratis acutis subtus sparsè et suprà secus nervos puberulo-scabris, corymbo compositi ramis apice 3-cephalis, capitulis subsessilibus, invol. obovato-oblongi squamis ext. squarrosis int. erectis acutis, ligulis involucri aequalibus. $\frac{1}{2}$ in campis et dumetis à Pennsylvania ad Floridam (Pursh). *Conyza asteroides* Linn. sp. 1206. *Aster Marylandicus* Michx. fl. bor. am. 2. p. 108. *Aster conyzoides* Willd. sp. 3. p. 2043 non Desf.

β. ? plantaginifolius (Nees ast. 150), foliis radicalibus spathulatis, capitulis subpedicellatis, ligulis invol. aequalibus aut longioribus. $\frac{1}{2}$ in sylvis Novæ-Cæsareæ (Nutt.) et ex Noveboraco (Mitch.), Albany, Niagara et Montreal (Janin!), Kentucky (Raf.), recepti. *Aster conyzoides β* Nutt. l. c. Hanc semper ex America recepi sub nomine *A. conyzoidis*. An species propria? An ligulae longitudine variantes? (v. s.)

3. *S. RIGIDUS* (Lindl. adn. mss.), foliis obovatis subtriveniis integris cauleque scaberrimis, corymbo terminali simplici vel composito contracto, involucri turbinati squamis adpressis basi nudis, radio pappo brevioris aut nullo. $\frac{1}{2}$ in America septentr.-occident. *S. conyzoidis* proximus. (Lindl.)
4. *S. TORTIFOLIUS* (Nees ast. 151), totus pube brevi densa rigidulà canescens, caule erecto tereti apice corymboso, foliis obovato-cuneiformibus mucronatis integerrimis subaveniis, capitulis pedicellatis basi folia duo parva gerentibus, invol. obovati squamis oblongis imbricatis acutiusculis basi pallidis, ligulis involucri paulò longioribus. $\frac{1}{2}$ in Carolinà infer. et Georgià. *Conyza bifoliata* Walt. car. 204. *Aster tortifolius* Michx. fl. 2. p. 109. Ell. sketch 2. p. 341. (v. s.)

CXVIII. *MACHÆRANTHERA* Nees ast. 224.

Capitulum radiatum, ligulis foemineis (neutrisve?) linearibus 3-nerviis. Invol. pluriserialè imbricatum, squamis herbaceis margine pallidis

apice patulis. Recept. nudum? Stigmata disci apice lineari elongato sterili. Antheræ corollâ longiores basi muticæ appendice terminali cultriformi. Achænia compressa (conica?) sericeo-hirsuta, pilis pap-pum exter. simulantibus. Pappus pilosus inæqualis scaber. — Herba perennis pilis capitatis pubescens facie Anthemidis. Folia alterna pinna-tifida lobis dentatis. Capitula terminalia solitaria ramo apice nudo pe-dunculata. Ligulæ albæ ex H. et B., in sicco purpurascentes. Discus luteus.

1. *M. TANACETIFOLIA* (Nees l. c.). ? in hort. Mexic. culta. Aster tanacetifolius H. B. et Kunth nov. gen. am. 4. p. 95. Aster chrysanthemoides Willd. herb. Spreng. syst. 3. p. 538. Affinis dicitur *A. pianato* Cav.

CXIX. TETRAMOLOPIUM Nees ast. 202.

Capitulum radiatum, ligulis angustis foemineis fertilibus 1-serialibus, fl. disci hermaphr. abortivis? masculisve. Recept. nudum punctatum. Invol. imbricatum, squamis pauciserialibus angustis. Achænia 4-nervia, radii compressa cuneiformia, disci lineari-tetragona. Pappus 1-serialis setosus scabridus. — Species habitu variz, fortè non omnino con-generes.

§. 1. *Involucro pauciseriali.*

1. *T. TETRAPHYLLUM* (Nees l. c. p. 203), herbaceum caespitosum, foliis inferioribus confertis lineari-spathulatis acutis ciliato-serratis, ramis apice nudis. ? in insulâ O-Wahn. Aster tenerimus Less. in Linnaea 1829. p. 120. Ligulæ albæ? Capitula 3 lin. alta et lata.

§. 2. *Involucro multiseriali. — An species verè congeneres?*

2. *T. ? RUPESTRIS* (Nees l. c.), fruticosum erectum infernè nudum, circinnatum, foliis lineari-lanceolatis margine revolutis subtus tomentosis. ? in frigidissimis montium Ruen-Pichinchensis et Antisana. Aster rupestris H. B. et Kunth nov. gen. am. 4. p. 94. t. 334. Aster Pichinchensis Willd. herb. ex Nees. Invol. squamæ multiserialis. Ligulæ circ. 30 albæ. Achænia glanduloso-punctata.
3. *T. ? PHYLLICOIDES*, fruticosum ramosissimum, ramis cano-lanatis, foliis approxi-matis patenti-reflexis sessilibus oblongo-linearibus margine revolutis supra glabris subtus lanatis, capitulis corymbozo-congestis, invol. campanulati squamis imbricatis lanceolato-linearibus lanato-ciliatis. ? in Mexico? Aster phyllioides H. B. et Kunth nov. gen. am. 4. p. 98. Capit. magis. Solidaginis Virgæ-aureæ. Ligulæ circ. 20 albæ. Discus 8-10-florus, fl. masculis et hermaphr. mixtis. Ovarium lineare compressum hispidulum. Pappus pilosus. aefescens scaber. — Affine ex Kunth *A. Chiliotricho* sed rec. epaleaceum, aut potius Tetramolopio rupestri.
4. *T. ? OCHRACEUM*, fruticosum, ramulis angulatis tomentosis, foliis petiolatis lanceolato-oblongis remotè et obsolete denticulatis supra glabris subtus ochra-ceo-tomentosis, corymbis ramosis, invol. subhemisphaerici squamis laxè im-bricatis margine diaphanis. ? in montibus Quiézensibus? Diplostephium? ochraceum Nees ast. 201. Aster ochraceus H. B. et Kunth nov. gen. am. 4. p. 94. Ligulæ circ. 5 albæ. Discus 20-florus, fl. masc. et hermaphr. intermixtis. Achænia obovata compressa, sterilia linearia. Pappus pilosus scaber. Habitus omninò Eurybiæ lyratæ,

CXX. *HENRICIA* Cass. bull. phil. 1817. p. 11. et 1818. p. 123. dict. 20. p. 567 et 37. p. 491. Nees ast. 219.

Capitulum radiatum, ligulis foemineis fertilibus 1-serialibus, disco hermaphr. Invol. hemisphaericum disco aequale, squamis biserialis

æqualibus adpressis, ext. foliaceis ovali-acutis, inter. scariosis obtusis. Recept. convexum nudum. Achænium pilosiusculum cylindraceum. Pappus 1-serialis setosus scabridus. — Suffrutex Madagascariensis ramosus. Folia alterna breviter petiolata ovato-lanceolata serrata rigida utrinque setuloso-scabra. Corymbus aphyllus oligocephalus. Pedicelli pilosi. Invol. subhispidum. Discus flavus. Ligulæ albæ.

1. *H. AGATHOIDES* (Cass. l. c.). ♂ in Madagascar. (v. s. in h. Juss.)

CCXI. DOELLINGERIA. — Doellingeriæ et Tripolii sp. Nees. — Asteris sp. auct.

Capitulum radiatum, flor. radii 1-ser. ligulatis fœmineis, disci hermaphr. 5-dentatis. Recept. alveolatum. Involucrum imbricatum. Stigmata disci brevia apice obtusa extus subhispida. Achænia angulata subteretia. Pappus biserialis pilosus, setis scabridis, aliis filiformibus, aliis apice incrassato-clavellatis. — Herbæ rhizocarpicæ, Japonicæ aut Chineses. Folia alterna serrata aut integra. Capitula fere Asterum ex habitu. — Genus certè apud cl. Nees species heterogeneas amplectens, hic ex *D. scabra* stabilitum.

1. *D. SCABRA* (Nees ast. 183), caule herbaceo tereti glabro supernè dichotomopaniculato, foliis ovalibus utrinque attenuatis medio-groscè serratis rigidulis, suprâ sparsè subtus secus nervos scabris, invol. squamis imbricatis obtusis, ligulis circ. 3, achæniis glabris. ♀ in Japoniâ. Aster scaber Thunb. fl. jap. 316. *Diplopappus scaber* Less. in Nees. Pappus pallidè rufus plurisetosus, setis scabridis, aliis apice incrassatis, aliis acutis. Stigmata brevia apice obtusè coriacea subhispida. Achænium angulatum. (v. s. à Thunb. olim mis. in h. Delessert.)
2. *D. TRICHOGARPA*, caule herbaceo tereti, ramis pedicellisque velutino-scabris, foliis elliptico-oblongis utrinque attenuatis latè dentatis suprâ muriculato-scabris, subtus secus nervos nervulosque villosis subhirtis, capitulis secus ramulos axillares laxè racemosis longè pedicellatis, invol. squamis oblongo-linearibus glabris acutis subobtusis, ligulis circ. 12, achæniis pilosis. ♀ in Chinâ legit cl. Rottb. Pappus pallidè infuscatus, setis aliis filiformibus, aliis parè subincrassatis acutis. Habitu ad *D. scabram* accedit, sed ex pappo media inter Doellingeriam et Asteram. (v. s. in h. Lindley.)

+ Species exclusæ.

D. AMYGDALINA Nees. = *Diplostephium amygdalinum*.

D. CORNIFOLIA Nees. = *Diplostephium cornifolium*.

D. OBOVATUM Nees. = *Diplostephium obovatum*.

D. PYRAMICOIDES Nees. = *Heleastrum album*.

D. UMBELLATA Nees. = *Diplostephium umbellatum*.

CCXII. HELEASTRUM. — Asteris sp. auct.

Capitulum radiatum, ligulis fœmineis 1-serialibus, fl. disci hermaphr. Invol. imbricatum, squamis laxis acuminatis apice foliaceis subsquarrosis. Recept. alveolatum, alveolis exsertis acutè dentatis. Stylis ramis elongati acuminatissimi dorso hispidi. Achænia oblonga angulata utrinque attenuata glabra. Pappus biserialis setis longitudine æqualibus, aliis filiformibus, aliis in ensem subclavellatam dilatatis. — Herbæ boreali-Americanae perennes. Caules simplices. Folia alterna elongata acuminata integerrima. Ligulæ coruleæ aut albæ. — Differt à Doellingeriâ stigmate acuminatissimo nec obtuso, à cæteris Euastoreis

pappo bifirmi setis aliis subclavellatis, à Diplopappeis pappo exteriori interiori æquante. — Nomen ex ἰλος, ἱλος, palus, et ἀστρῶν, astrum.

1. *H. PALUDOSUM*, caule apice subcorymboso et tenuissimè subpuberulo, foliis linearibus amplexicaulibus subconcavis apice subulatis, invol. squamis subquarratis. 2*l* in paludibus et uliginosis Carolinæ infer. (Pursh), Georgiæ prope Savannah (Herb. in h. Dnby), à Novâ-Cassareâ ad Floridam (Nutt.). Aster paludosus Ait. kew. ed. 1. v. 3. p. 201. Nutt. gen. 2. p. 155. Ell. sketch 2. p. 343. Aster grandiflorus Walt. ex Ell. Tripolium paludosum Nees ast. 155. Diplopappus paludosus Lindl. adn. mss. Capitula 3-5. Ligulæ latè coerulææ numerozæ. Caulis pubescens. Folia margine scabra. (v. s.)
2. *H. ALBUM*, caule apice in ramos 1-cephalos corymboso-fastigiato diviso, foliis oblongo-lanceolatis scabris infernè attenuatis, involucri lævi. 2*l* in Amer. borealis planitie Missonriensi (Nutt.), frequens videtur in Canadâ et Amer. arcticâ (Lindl.), Chrysopsis alba Nutt. gen. am. 2. p. 152. Aster albus Willd. herb. ex Spreng. syst. 3. p. 528. Doellingeria ptarmicoides Nees ast. 183. Diplopappus ptarmicoides Lindl. adn. mss. Ligulæ lineari-oblongæ albæ (Nutt.), Pappus biformis clavatus et setaceus (Lindl.).
- β. caule 1-cephalo. — ad lacum superiorem.
- γ. corymbo composito, foliis subdentatis. — cum var. β legit Pilcher ex Lindl. mss.

CXXXIII. BIOTIA DC. non Cass. — Eurybiæ sp. Cass. dict. 37. — Eurybiæ genuinæ Nees ast. 137. — Asteris sp. auct.

Capitulum radiatum, ligulis fœmineis fertilibus 1-serialibus latiusculis, disco hermaphr. fertili. Recept. alveolæ obtusè dentatæ. Invol. ovati squamæ pluriseriales arcuè imbricatæ adpressæ non appendiculatæ, int. gradatim longiores et angustiores. Stigmata disci hispida acuminata retorta. Achænia glabra vel pubescenti-scabrida elongata tricostrata. Pappus, setis rigidulis filiformibus scabridis subinæqualibus. — Herbariæ rhizocarpicæ Boreali-Americanae Asteris facie. Caules erecti apice corymbosi. Folia serrata, radicalia ampla petiolata, caulina alato-petiolata ovata vel oblonga. Ligulæ pallidæ. Discus flavus. — Biotia Cassinii est monstruositas Madizæ. Eurybia Cass. genuina (dict. v. 16) ex speciebus Novo-Hollandicis tantum constat. Genus ex Asteribus sejunctum dicavi cl. amicis. Biot qui ex Astrorum studio ad Botanicam physiologicam feliciter migravit.

1. *B. SCHREBERI*, caule glabra apice decomposito-corymboso, foliis grossè serratis suprâ subscabris subtus glabris, radicalibus longè petiolatis ovatis aut suborbiculatis, caulinis inf. secus petiolum subdecurrentibus, super. sessilibus ovatis acutis, capitulis pedicellatis, invol. squamis arcuè imbricatis oblongis obtusis, ligulis invol. duplò longioribus. 2*l* in Amer. bor. (Bernh.). Aster Schreberi Nees! syn. (1818). p. 16. Aster subcymosus et A. ambiguus Bernh. ex Nees et Cav. cat. h. taur. 1821. p. 12. A. macrophyllus var. Van Roy. hort. lugd. ex h. Delaroche. Eurybia Schreberi Nees! ast. 137. Sæpè cum B. macrophyllâ confusa sed caulibus glabris facillè distincta, ab etiam et à B. latifoliâ foliis radicalibus non cordatis differt. (v. v. c. à cl. Nees mss.)
2. *B. LATIFOLIA*, caule glaberrimo apice corymboso, foliis glabris, radicalibus caulinisque infer. petiolatis cordatis, super. alato-petiolatis sessilibusve, omnibus serratis acuminatis, capitulis pedicellatis, invol. squamis arcuè imbricatis oblongo-lanceolatis obtusis, ligulis elongatis. 2*l* in Amer. bor. (Desf.) ad montes Alleghanijs (Raf.) et circa Bostoniam (Lindl. herb.). Aster latifolius Desf. cat. h. par. p. 176. Aster macrophyllus Raf. in litt. Eurybia macrophylla γ Nees ast. 140? fortè suad. Lindley mera B. Schreberi var. foliis cordatis distincta? (v. s.)

3. *B. GLOMERATA*, caule glabro apice thyrsoideo, foliis infer. sinu lato cordatis alato-petiolatis, super. ovato-ellipticis subsessilibus, omnibus serratis supra in ambitu subtus ad costas scabris, capitulis subsessilibus, invol. squamis arcuè imbricatis ovatis obtusis, ligulis vix involucri æquantibus. $\frac{1}{2}$ in Pennsylvania (Bernh.). *Aster glomeratus* Bernh. ex Nees. *Eurybia glomerata* Nees ast. 139. — *Aster* in hort. Par. anno 1818 sub nom. erroneo *A. corymbosi* cultus huc pertinere videtur, sed ligulæ parùm invol. longiores nec breviores. (v. s.)
4. *B. MACROPHYLLA*, caule plùs minùs hirsuto apice patulo-corymboso, foliis radicalibus petiolatis cordatis, caulinis alato-petiolatis sessilibusve ovatis, omnibus subcaberis serratis acuminatis, capitulis pedicellatis, invol. squamis arcuè imbricatis oblongo-lanceolatis obtusis, ligulis elongatis. $\frac{1}{2}$ in Amer. bor. umbrosis rupestribus à Canadà ad Virginiam. *Ast. macrophyllus* Linn. sp. 1232. Michx. fl. bor. am. 114. *Eurybia macrophylla* Cass. dict. 37. p. 487. Nees ast. 140. Desf. hort. Par. ed. 3. p. 176. *A. cordifolius* hort. plur. Folia floralia acuta. (v. v. et s.)
- β . *divaricata*, foliis floralibus rotundatis. $\frac{1}{2}$ in Amer. bor. (Linn.). *Ast. divaricatus* Linn. sp. 1226. *Ast. macrophyllus carneus* hort. angl. ex herb. Linn. obs. Lindl. in adn. mss.
5. *B. COMMIXTA*, caule infernè glabro supernè pubescenti-scabro patulo-corymboso, foliis ovatis serratis, radic. petiolatis, caulinis super. oblongo-lanceolatis sessilibus viscido-scabris, capitulis pedicellatis, invol. squamis imbricatis subquarrosis acutis, ligulis elongatis. $\frac{1}{2}$ patr. ign. verisim. Amer. bor. *Eurybia commixta* Nees ast. 142. An. *E. Jussiei* Cass. dict. 37. p. 487? Ligulæ lilacinæ. (v. s. c. olim in hort. angl. sub nom. *A. ornati*.)
6. *B. CORYMBOSA*, caule glabro gracili apice laxè corymboso, foliis petiolatis serratis acuminatis subtus secus nervos subsetosis, infer. petiolatis cordatis, mediis ovatis, sup. alato-petiolatis aut subsessilibus, capitulis pedicellatis, invol. squamis imbricatis obtusis, exter. ovatis, ligulis 6-8 invol. duplò longioribus. $\frac{1}{2}$ in sylvis montanis à Canadà ad Virginiam Lindl. bot. reg. t. 1532. *Aster corymbosus* Ait. kew. ed. 1. v. 3. p. 207. Nees syn. 17. *Eurybia corymbosa* Cass. dict. 37. p. 487. Nees ast. 143. *Aster cordifolius* Michx. fl. bor. am. 2. p. 114. ex. ejus herb. obs. cl. Lindley. (v. s.)
- β . *alata* (Bartl. fl. phil. 2. p. 114), petiolis latius et frequentius alatis. Varietas ex cl. auct. permanens.

CXXIV. *EURYBIA* Cass. bull. philom. 1818. p. 166. dict. 16. p. 486. non fl. mex. — *Eurybiæ* sp. Cass. dict. 37. p. 485. — *Enrybiæ* sp. et *Spongotrichum* Nees. — *Asteris* sp. Labill.

Capitulum pluriflorum radiatum, ligulis 1-serialibus fœmineis fertilibus, fl. disci tubulosi 5-dent. hermaphr. Recept. parvum planum aut subconvexum alveolatum. Invol. oblongum imbricatum, squamis chartaceo-membranaceis. Styli radii ramis elongatis erectis obtusis vix hispidulis. Achænium nunquàm compressum obovatum aut oblongo-cylindraceum striato-angulatum aut angulato-alatum glabrum vel apice puberulum. Pappus uniserialis, setis scabris præsertim ad apicem, subæqualibus, interdum apice incrassatis. — Frutices omnes à Novâ-Hollandiâ orti. Folia alterna (in 2-3 speciebus opposita) coriacea petiolata aut sessilia, integerrima dentata aut sinuata, sæpè subtus tomentosa. Inflorescentia varia. Discus flavus. Ligulæ albæ violacæ aut carneæ. — *Eurybia* in veterum mythologiâ erat stellarum mater.

SECT. I. BRACHYGLOSSA.

Ligulæ stylos suos non excedentes.

1. *E. BRACHYGLOSSA*, fruticosa, foliis alternis subsessilibus obovatis margine re-

volutis subintegerrimis suprà glabris subtus tomentosis, capitula ad axillas foliorum solitariis subsessilibus, ligulis circ. 3 stylo sue fere brevioribus. \S in Novæ Hollandiæ orâ orientali. Simillima icon. t. 202 Labill. sed diversa capitulis solitariis nec ternis, sessilibus nec pedunculatis, ligulis brevissimis, invol. squamis nec acuminatis. Achænium glabrum. Pappus α -serialis. (v. s. comm. à Mus. reg. Par.)

2. *E. AXILLARIS*, fruticosa, ramis tomentosis, foliis alternis oblongo-cuneatis obtusis integerrimis, junioribus suprà araneosis demum glabris subtus tomentosis, capitulis in axillis fol. solitariis subsessilibus paucifloris, ligulis 3-4 stylo non superantibus. \S in Novæ Hollandiæ littore orientali. Folia 9-10 lin. longa, 2 lin. lata. Capitula parva. Folia minima obtusa cava extus involucri bracteiformia. Invol. squamæ obtusæ. Pappus pallidè rufescens subaniserialis. (v. s. comm. à Mus. reg. Par. ex itin. Baudin.)

3. *E. LINEARIFOLIA*, fruticosa, ramulis tomentoso-canis, foliis alternis linearibus utrinquè acutiusculis integerrimis suprà demum glabris subtus tomentosis, capitulis in axillis foliorum subsessilibus, solitariis et secus ramos foliatis racemum longum constituentibus, ligulis stylo fere brevioribus. \S in Novâ Hollandiâ australi. Rami conferti. Folia 10 lin. longa, lineam lata, margine revoluta. Invol. squamæ oblongo-lineares apice summo ad dorsum tomentosæ. Pappus pallidè rufescens. Capit. triplò majus quàm in axillari, 9-florum. Ligulæ 4 parvæ. Achænium villosissimum. (v. s. comm. à Mus. reg. Par. ex itin. Baudin.)

4. *E. CAPITELLATA*, fruticosa, ramulis tomentoso-canis, foliis linearibus basi attenuatis vix acutiusculis integerrimis suprà junioribus subpubescentibus adultis glabris subtus tomentosis, ramulis apice capitula parva pauciflora distinctè paniculata paucâ axillaria et terminalia gerentibus, ligulis paucis stylo non superantibus. \S in Novâ Hollandiâ orientali. Valdè accedit ad *E. axillarem* sed capitulis distinctissimè pedicellatis egregiè differt. Invol. squamæ extus subcanescentes lineares obtusæ demum patentes. Recept. planum vix scrobiculatum. (v. s. comm. à Mus. reg. Par. ex itin. Baudin.)

5. *E. DAMPIERI* (A. Cunningham! in litt. 1834. sub Conyzâ), fruticosa, ramulis tomentoso-canis, foliis alternis linearibus obtusis integerrimis margine subrevolutis suprà glabris subtus tomentosis, capitulis axillaribus breviter pedicellatis solitariis geminisve, invol. squamis obtusis, ligulis stylo subæqualibus. \S in insulis Archipelagi Dampieri frequens (King voy. 1818), in ins. Enderby unâ ex insulis Rosemary anglis dictis (A. Cunn.!), Conyzæ Novæ Hollandiæ angustis rosmarini foliis Woodward in Damp. voy. ed. 1729. vol. 3. p. 111. t. 4. f. 3. ex Cunn. Species foliis pollicem longis vix lineam latis ab omnibus affinis distincta! (v. s. comm. à cl. Cunn.)

6. *E. OLIGANTHA*, fruticosa, ramis canescentibus, foliis oblongo-cuneatis obtusis integerrimis suprà glabris subtus canescentibus, capitulis 3-4-floris ad axillas foliorum solitariis subsessilibus. \S in Novâ Hollandiâ ad sinum Canum mariorum legit cl. Gaudichaud. Specimina valdè manca ex ejus herb. habui cæcè tantum ex habitus analogiâ refero. (v. s. sine fl. et fr.)

Sect. II. ARGOPHYLLÆ. — Eurybiæ heteroclytæ Nees ast. 144.

Ligulæ stylos suos longè superantes. Achænium subcylindricum.

7. *E. CHRYSOPHYLLA*, fruticosa, foliis oppositis petiolatis oblongis basi attenuatis apice acutiusculis integerrimis suprà glabris nitidis sublaevibus subtus flavido-æresco-tomentosis, pedunculis axillaribus folio duplò longioribus α -cephalis in corymbum laxum dispositis. \S in montanis Novæ Cambriæ australis ad septentr. urbis Bathurst, mense septemb. 1822. omninò defloratam legit cl. All. Cunningham. Aster chrysophyllus Cunn. in litt. 1834. Frutex robustus 6-pedalis, affinis *E. viscosæ* sed foliis suprà nec reticulatis nec viscosis et pedicellis α -cephalis differt. (v. s. comm. à cl. invent.)

8. *E. VISCOSA* (Cass. dict. 37. p. 487. Nees ast. 146), fruticosa, foliis oppositis

petiolatis ellipticis oblongiusve utrinque attenuatis apice obtusiusculis integerrimis supra glabris reticulatis subtus albo-tomentosis, corymbis terminalibus trichotomis, acheniis villosissimis. \S in Novâ Hollandiâ ad oram merid. et ins. Van-Diemen (Cunn.!) ad Hobart-Town (Cunn.!). Aster viscosus (Labill.!) Nov. Holl. sp. 2. p. 53. t. 203. Balbisia Caledoniæ Spreng. syst. 3. p. 569. Balbisia suffruticosa Dietr. Folia recentia supra viscosa (Labill.) 3 poll. longa, 5-12 lin. lata, petiol. 2-3 lin., apice obtusa aut vix acuta. Capitula parva pauciflora. Invol. squamæ lineares acutæ. Ligulæ violaceæ apice 3-dentatæ. Achenia pilis sursum tendentibus villosissima teretiuscula. (v. s.)

9. *E. PERSOONIOIDES*, fruticosa, foliâ alternis et suboppositis obovatis basi cuneatis brevissimè petiolatis integerrimis coriaceis supra venosis glaberrimis lucidis, subtus pube sericeâ adpressâ confertâ canis, capitulis pedicellatis paucis subcorymbosis, invol. squamis ciliatis, achenio hirtello. \S in petrosis insula Van-Diemen (Cunn.!) ad montem Wellington 4100 ped. alt. supra mare (Cunn.!). Aster persoonioides Cunn. in litt. 1834. Affinis *E. myrsinoidi*. Suffrutex ramosus ut videtur humilis. Rami angulati canescentes. Folia sesquipollicem longa 9 lin. lata, obtusa aut submucronata. Pappus albidus copiosus basi intortus. Flosc. in invol. pauci. (v. s.)

10. *E. FURFURACEA*, fruticosa, foliis alternis petiolatis ovatis ovalibuve coriaceis integerrimis supra glabris viridibus subtus tomento adpressissimo argenteis, corymbo composito. \S in Novâ Zeelandiâ ad ripas fluminum et spec. ad flumen Tamesim legerunt cl. D'Urville et Lesson et A. Cunningham qui mecum human. commun. specimina. Aster furfuraceus Rich. et Less. fl. Nov. Zeel. p. 216. Aster elæagnifolius Cunn. in litt. 1834. Species elegans 6-10-pedalis *E. argophyllæ* affinis sed foliis integris faciliè distincta, affinis etiam *E. chrysophyllæ* et *E. viscosæ* sed foliâ alternâ. Invol. serè turbinatum, squamis 4-seriatis paucis. Pappi setæ apice longius barbatae. (v. s.)

11. *E. ARGOPHYLLA* (Cass. dict. 37. p. 487), fruticosa, foliis alternis petiolatis oblongo-lanceolatis acuminatis inæqualiter et argutè serratis coriaceis supra glabris subtus ramisque argenteo-sericeis, capitulis paniculato-corymbosis. \S in sylvis ambrosiis montanis Nov. Holland. oræ merid. et orientalis et ad Cap. Van-Diemen. Aster argophyllus Labill. Nov. Holl. spec. 2. p. 52. t. 201. Sims bot. mag. t. 1563. Arbor 20 ped. alta, truncus 8 poll. diam. (Cunn.!). Folia friatâ odorem moschi redolent! Capitula parva. Invol. squamæ imbricatæ adpressæ extus velutinæ. Ligulæ 3-5 albae, stylo evid. exserto bifido. Cor. tubulosæ ad medium 5-fidæ, lobis linearibus. Ovaria glabra. Pappus rufescens acabes, radii hi, disci uni-serialis. Recept. angustum alveolatum. (v. v. c. et n. sp.)

12. *E. CYDONIÆFOLIA*, fruticosa, foliis alternis petiolatis ellipticis obtusis basi attenuatis integerrimis supra glaberrimis subtus petiolis ramis pedunculis involucriisque molliter velatino-tomentosis, pedunculis squamulosos parè ramosos, invol. turbinatâ squamis imbricatis multiserialibus, inter. gradatim longioribus linearibus. \S ad oram orientalem insulæ borealis Novæ Zeelandiæ octob. 1806. legit cl. A. Cunningham. Aster cydoniæfolius Cunn. in litt. 1834. Flores in æpæcin. juniores mihi non satis noti. (v. s. comm. à cl. inveni.)

13. *E. LIRATA*, fruticosa, ramis junioribus tomentosis adultis hirato-striatis, foliis alternis petiolatis lanceolatis basi obtusis apice longè acuminatis repando-subdentatis supra glabris subtus tomentosis, corymbo paniculato. \S in Novâ Hollandiâ. Aster liratus (nec lyratus) Sims bot. mag. t. 1509. Diplostephium liratum Nees ast. 189. Aster pulverulentus Hoffmannsegg ver. p. 1824. p. 165. Folia 3 poll. longa, 5-6 lin. lata, petiolo 3 lin. Ligulæ albae circ. 15 lineares. Achenia obovata pentagona glabra atrigulis minimis secus angulos apice imbr. Pappus albidus meo sensu 1-serialis. (v. s. c.)

14. *E. RAUWASCEUS*, fruticosa, foliis alternis vix petiolatis oblongis seu obovatis rigidè et acutè sinuato-serratis coriaceis supra glabris reticulatis subtus ramisque tomentosis, capitulis 15-20-floris paniculatis. \S in Novâ Hollandiâ (Sieb.!), in collibus petrosis montium Cœruleorum (Cunn.!) et in ins. Van-

Diemen prope Hobart-Town (D'Urville). *Aster crubescens* Sieb. ! pl. exs. n. 389. *Aster herberifolius* Cunn. ! in litt. Folia 12-15 lin. longa, 4-5 lata, petiolo lineam longo. Tomentum rufescens. Pedunculi axillares et terminales paniculati foliis parvis oblongis subintegris ad ramific. instructi. Invol. et pappus *E. argophyllæ*. Ligulæ paniceæ lineares fœmiuæ ex sicco albæ. Tubuli 5-dentati nec 5-fidi. Achænidium teretiusculum 8-10-striatum, adultum glabrum. (v. s.)

β. *ilicifolia*, foliis oblongo-lanceolatis utrinquē acutè attenuatis. β in sylvis ad septentr. urbis Bathurst in Novâ Cambriâ australi legit cl. Cunningham nov. 1822. *Aster ilicifolius* Cunn. in litt. 1834. Folia 2-3 poll. longa, 6-8 lin. lata. Specim. Sieberianum omnino medium inter stirpes duas Cunninghami. (v. s. comm. à cl. inv.)

15. *E. MYRSINOIDES* (Nees ast. 146. excl. Sieb. syn.), fruticosa, foliis alternis subsessilibus obovatis margine revolutis et subcrenatis suprâ glabris subtus tomentosis, pedunculis 3-cephalis folio paulò longioribus 5-6-floris, ligulis 3 stylos suos duplò ferè excedentibus. β in Novâ Hollandiâ ad Cap. Van-Diemen. *Aster myrsinoides* Labill. Nov. Holl. sp. 2. p. 53. t. 202. Folia 4-5 lin. longa, 2. lin. lata. Invol. imbricatum. Flores 6-8. Ligulæ angustæ acutæ. Achænia glabra.

β. *serrata*, foliis argutè serratis subtus sericeo-tomentosis, ligulis 2 elongatis cor. disci duas multò superantibus. β in insulâ Van-Diemen legit cl. Gunn. An spec. propria ? (v. s. comm. à cl. Lindley.)

16. *E. PIMELEOIDES*, fruticosa, foliis alternis obovatis obtusis basi cuneatis margine integris subrevolutis suprâ glabris subtus ramulisque tomentosis, capitulis terminalibus solitariis, invol. subhemisphærici multiflori squamis imbricatis, ext. brevibus subtomentosis, inter. longioribus scariosis glabris, ligulis circ. 10 disco duplò longioribus, achæniis sericeo-villosissimis. β in sterilibus dumosis propè Peel's range in Nov. Holl. inter. ad occid. urbis Port-Jackson legit cl. Cunningham jun. 1817. *Aster pimeleoides* Cunn. ! in litt. 1834. Habitus *Pimeleæ*. Folia referunt icon. Labill. t. 202. sed capitula solitaria multò majora, achænia villosissima, etc. Villi achæniî superiores pappum exteriorem simulant. (v. s. comm. à cl. Cunningham.)

17. *E. SUBREPANDA*, fruticosa, ramis tomentoso-canis, foliis alternis oblongo-linearibus obtusis basi attenuatis margine obtusissimè subrepandis suprâ junioribus pulverulento-scabris demùm glabris subtus tomentosis, capitulis pedicellatis ad apices ramorum racemoso-corymbosis. β in Novâ Holl. insulâ King prope fretum Bass (itin. Baudin !) et in insulâ Van-Diemen prope Hobart-Town (D'Urville ! Gunn !). Folia 8-9 lin. longa, lineam lata. Pedicelli tomentosi 3 lin. longi. Invol. squamæ oblongæ acutæ dorso tomentosæ flosculis breviores. Pappus rufescens. Achænidium pubescens. Ligulæ 8-10. (v. s.)

18. *E. GUNNIANA*, fruticosa, ramis tomentoso-canescens, foliis alternis oblongo-linearibus obtusè sinuato-dentatis subcoriaceis suprâ glabris venoso-reticulatis subtus densè tomentoso-canis, capitulis longè pedicellatis ad apices ramulorum racemoso-corymbosis β in insulâ Van-Diemen legit cl. Gunn. Valdè accedit ad *E. subrepandam* sed folia longiora, etiam juniora suprâ glabra, regulariter dentata, 15 lin. longa, 3-4 lin. lata, capitula dimidiò minora longius pedicellata, ligulæ 12-13 (albæ) paulò breviores, fl. disci pauciores (10-12), invol. squamæ ferè glabræ. (v. s. comm. à cl. Lindley.)

19. *E. ROSMARINIFOLIA*, fruticosa, ramulis canescenti-tomentosis, foliis oppositis linearibus mucronatis margine revolutis integerrimis suprâ submuricatis subtus tomentosis, pedunculis corymbosis ultimis trichotomis, pedicellis elongatis, invol. squamis lanceolato-linearibus subcarinatis apice callosis. β in Novâ Hollandiâ prope Bathurst in collibus calcareis mense novemb. legit cl. Cunningham. *Aster rosmarinifolius* Cunn. ! mss. 1822. An fortè *Solidago rosmarinifolia* Cass. dict. 56. p. 167 ? Folia 1 $\frac{1}{2}$ -2 poll. longa. Capitula parva. Ligulæ albæ. Achænidium glabrum. (v. s. comm. à cl. invent.)

20. *E. LEDIFOLIA*, fruticosa, ramis glabris subangulatis junioribus pube adpressâ subsericeis, foliis alternis margine revolutis integerrimis obtusis subcylindricis suprâ glabris subtribus in sulco sericeo-subhirsutis, pedicellis paucis 1-cephalis, invol. squamis oblongis patulis dorso subhirsutis. $\frac{1}{2}$ in insulâ Van-Diemen (Gunn!) in rupestribus montis Wellington januario flor. (Cunn.). *Aster ledifolius* Cunn. in litt. 1834. Capitula majora paucissima et folia dimidiò breviora quàm *E. rosmarinifolia*. Pappus sordide albus pluriserialis, setis exterioribus nonnullis brevissimis. Achænia hispidula. (v. s.)
21. *E. DECURRENS*, fruticosa glabra glutinosa, ramis angulatis, foliis remotis basi angustissimè decurrentibus linearibus coriaceis nunc integerrimis nunc versus apicem dentibus 1-2 repando-dentatis, capitulis ramulos terminantibus subracemosis, invol. squamis linearibus glabris, flosculis paucis, achæniis turbinatis sericeo-villosis. $\frac{1}{2}$ in sterilibus partium Novæ Hollandiæ interiorum ad austr.-occid. flum. Lachlan legit cl. A. Cunningham. *Aster? decurrens* Cunn. in litt. 1834. Fruticulus ferè *Ephedræ* facie. Stigmata longè exserta apice lanceolata. (v. s. comm. à cl. invent.)

Sect. III. SPONGOTRICHUM *Nees ast.* 176.

Ligulæ stylos suos superantes. Achænium tetragono-conoideum scabrum. Capitulum campanulatum. Invol. squamæ glanduloso-ciliatæ. Suffrutices foliis linearibus. — An genus proprium?

22. *E. GLANDULOSA*, fruticulosa glabra, foliis alternis sessilibus linearibus margine glandulas protuberantes gerentibus, capitulis pedicellatis corymbosis, invol. squamis margine angustè membranaceis tenuissimè ciliato-serratis. $\frac{1}{2}$ in paludosis Novæ-Hollandiæ prope lacum Georgii ad Cap Van-Diemen et fretum d'Entrecasteaux. *Aster glandulosus* Labill. Nov. Holl. sp. 2. p. 50. t. 197. *Galatella glandulosa* Nees *ast.* 174. *Aster rutæodorus* Cunn. in litt. Folia pollicem et ultrâ longa vix semi-lineam lata patentia. Ligulæ 10-20 albæ oblongæ integerrimæ. Achænia cuneiformi-tetragona pubescenti-scabra. Pappus 1-serialis pallidè rufescens, setis apice paulò longius barbellatis. (v. s.)
23. *E. HYGROPHILA*, suffruticosa, caule infernè glabro secus ramos apice pubescente, foliis alternis linearibus utrinquè attenuatis medio paucidentatis aut subrepandis suprâ glabris nervo medio impresso, subtus pube stellatâ brevi velutinis margine subrevolutis, capitulis longè pedicellatis longè racemoso-paniculatis, invol. squamis acutis, pappo biseriali vix inæquali. $\frac{1}{2}$ abundè in fossâ paludosa Novæ Hollandiæ, ad litus boreale insulæ Stradbroke (lat. 27 $\frac{1}{2}$) mense septembr. floridam legit cl. Cunningham. *Aster hygrophilus* Cunn. in litt. 1828. Ligulæ albæ. Achænium glabrum. Caulis rubescens. (v. s. comm. à cl. invent.)
24. *E. TENUIFOLIA*, fruticulosa araneosa aut glabrata, foliis alternis sessilibus eglandulosis integerrimis, ramulis foliosis apice 1-cephalis subcorymbosim dispositis, invol. squamis linearibus acuminatis puberulis. $\frac{1}{2}$ in Novâ Hollandiâ orientali (h. Mus. Par.) et interiori ad Cabbageangriviè (Cunn.). Accedit ad *E. glandulosam* sed folia glandulis orbata. Ligulæ 9-10-lineares elongatæ. Achænia scabra. Pappus 2-serialis subrufescens, serie ext. brevi multisetâ. (v. s.)
- β . *Bathurstiana*, invol. squamis interioribus subobtusis acutiusculis nec acuminatis. $\frac{1}{2}$ in sterilibus circa Bathurst legit cl. A. Cunningham. Frutex florihundus fortè ex cl. invent. distinguendus. (v. s. comm. à cl. Cunningham.)
25. *E. ELÆOPHILA*, suffruticosa glabra, caule erecto, ramis striato-angulatis, foliis alternis linearibus integerrimis margine subrevolutis, capitulis ad apices ramorum ramulorumque solitariis, invol. squamis lanceolatis acutis dorso subpuberulis, achænio pubescente subangulato. $\frac{1}{2}$ in paludosis ad ripas lacus dicti Lagoon in sinu Regis Georgii ad oram merid.-occid. Novæ Hollandiæ, januario flor. legit cl. Cunningham. *Aster elæophilus* Cunn. in litt. 1834. *Affinis E. tenuifoliæ*. Capitula dimidiò minora. (v. s. comm. à cl. inv.)

26. *E. RAMOSISSIMA*, fruticosa ramosissima, ramulis villoso-subtomentosis, foliis lineari-lanceolatis integerrimis acutis minimis supra glabris subtus tomentosis, capitulis terminalibus solitariis sessilibus, invol. squamis imbricatis apice ovato-lanceolatis dorso sublanatis margine nigro ciliato-verratis, achenio villosissimo, pappo biseriali. ζ in Novâ Hollandiâ prope Port-Jackson legit cl. Gaudichaud, et in Nov. Camb. interiore ad septentr. flum. Damarep in climate ferè tropico collegit cl. Cunningham. Species distinctissima! Folia 2-3 lin. longa. Capitula 5-6 lin. longa et lata. Ligulæ 10-12. Recept. alveolatum. Pappus copiosus albus dentato-scaber. Achenium subangulato-turbinatum vilis albis erectis copiosis ornatum. (v. s.)
27. *E. LEPIDOPHYLLA*, fruticosa, foliis alternis conferto-imbricatis exiguis oblongis crassis margine revolutis integerrimis coriaceis subtus tomentosis, ramulorum abortivorum fasciculatis ramos obtegentibus, capitulis ad apices ramulorum brevissimorum solitariis. ζ in Novâ Hollandiâ ad littus australe et Cap. Van Diemen. Aster microphyllus Labill. ! Nov. Holl. sp. 2. p. 51. t. 199 non Vent. Aster lepidophyllus Pers. ench. 2. p. 442. Invol. squamæ vix acutæ. Achen. villosum. Pappus albidus, setis exter. dimidiò brevioribus paucioribus scabridis ut interiores. (v. s.)
28. *E. MICROPHYLLA*, fruticosa, ramis setuloso-hirtis, foliis obovatis integerrimis obtusis basi attenuatis supra muricatis subtus tomentosulis vix subincanis, capitulis ad apices ramulorum axillarium brevissimorum solitariis secus ramos foliosos racemosis, pappo exteriori brevissimo. ζ in Novâ Hollandiâ orientali. Aster microphyllus Vent. malm. 83 (non Labill.). Sieb. ! pl. exs. Nov. Holl. n. 338. Diplostephium microphyllum Nees ast. 191. Folia ex ramulis axill. abortivis fasciculata lineam sesquilineamve longa. Capit. 2 lin. longa. Invol. squamæ subobtusæ apice dorso puberulæ. Ligulæ albæ angustæ. Achenium puberulum. Pappus duplex, ext. brevis. (v. s. ex Sieb. Gaudich. et itin. Baudin.)
29. *E. PROPINQUA*, fruticosa, ramis glabris teretibus, ramulis pubescenti-subcaescentibus, foliis alternis minimis linearibus integerrimis margine subrevolutis supra glabris subtus subpubescentibus, capitulis ramulos terminantibus glabris, squamis linearibus vix acutis, achenio villosio. ζ in dumosis plantilei Liverpool Novæ Hollandiæ legit cl. Cunningham julio flor. Eandem fortè habeo ex orâ merid. Novæ Holl. in Baudin. expedit. lectam. Aster propinquus Cunn. ! in litt. 1834. Media inter *E. microphyllam* et *E. ramulosam*. (v. s.)
30. *E. RAMULOSA*, fruticosa, ramis pubescenti-scabris, foliis alternis lineari-lanceolatis acutis integerrimis supra muricato-scabridis subtus tomentosis albidis, capitulis ad apices ramulorum axill. brevissimorum solitariis secus ramos foliatis racemosis, pappo subuniseriali. ζ in Novæ Hollandiæ littore merid., ad Port-Jackson, circa Bathurst et Cap. Van-Diemen. Aster ramulosus Labill. Nov. Holl. sp. 2. p. 50. t. 196. Diplostephium ramulosum Nees ast. 193. Aster australis Hortulanorum. Folia 4 lin. longa, semilin. lata. Capit. 2 lin. longa. Invol. squamæ lanceolatæ acutæ membranaceæ dorso vix pubescentes. Ligulæ 6-7 albæ ferè lineares. Cor. disci 7-3 albæ ! Pappi setæ ext. 1-3. (v. s. sp. et v. c.)
31. *E. ACULEATA*, fruticosa, ramis setuloso-hirtis, foliis alternis lineari-lanceolatis integerrimis supra muricatis subtus tomentosis, capitulis ad apices ramulorum axill. brevissimorum solitariis secus ramos foliatis densè racemosis, pappo subuniseriali. ζ in Novæ Holl. Cap. Van-Diemen. Aster aculeatus Labill. ! Nov. Holl. sp. 2. p. 52. t. 200. Aster exasperatus Litk. enum. 2. p. 328. Aster foliolosus Hortul. Haxtonia foliosa hort. Bonn. Diplostephium aculeatum Nees ast. 192. Invol. squamæ oblongo-lineares obtusiusculæ apice ciliatæ. Ligulæ paucæ lineari-lanceolatæ. Pappus albus. (v. s.)
32. *E. SPILÆIA*, fruticosa, ramis tomentoso-canis, foliis alternis lineari-lanceolatis basi attenuatis integerrimis supra glabris lævibus nervo medio impresso, subtus tomentosis canis, capitulis ad apices ramulorum axill. brevissimorum solitariis secus ramos foliatis densè racemosis, pappo subuniseriali. ζ in Novæ

Holl. terrâ Van-Diemen. Aster aculeatus herb. Mus. Par. Accedit reverâ ad E. aculeatata sed ramis tomentosâ nec hirtellis, foliis suprà levibus nec muricatis paulò majoribus etc. differt. Achæmium glabrescens. Pappus ferè 1-ser., setis nempe ext. paucissimis subnullis. (v. s. comm. à Mus. reg. Par.)

† *Species exclusæ.*

- E. COMMIXTA Nees. = *Biotia commixta*.
 E. CORYMBOSA Cass. = *Biotia corymbosa*.
 E. FULVIDA Cass. = *Olearia stellulata*.
 E. JUSSÆI Cass. = *Biotia commixta*.
 E. GLOMERATA Nees. = *Biotia glomerata*.
 E. QUERCIFOLIA Cass. = *Olearia phlogopappa*.
 E. SCHREBERI Nees. = *Biotia Schreberi*.

Subdiv. 2. DIPLOPAPPÆ.

Radix 1-serialis. Pappus disci et radli conformis duplex, exterior squamosus, interior pilosus.

CXXV. OLEARIA Moench suppl. 254. Nees ast. 184. — *Diplostephia eurybioidea* Nees ast. 193. — *Diplopappi* sp. Less. — *Haxtonia hortul.* — *Asteria*, sp. auct.

Capitulum multiflorum heterogamum, fl. radii 1-seriatis ligulatis fœmineis, disci tubulosi hermaphroditis 5-dentatis. Recept. alveolatum planiusculum pedunculo apice cavo obconico insidens. Invol. squamæ imbricatæ adpressæ post fructif. patentes. Stam. radii abortiva, disci filamentis brevissimis donata. Stigmata disci cochleariformia. Achæmium cylindraceo-tetragonum villosum. Pappus duplex, exterior brevis paleaceo-setosus, interior setosus longus scaber. — Frutices Novo-Hollandici facie Eurybiæ. Folia alterna petiolata ovata repanda seu obtusè dentata aut integra suprà sparsè villosa subtùs ramique densè tomentosa. Pedunculi 1-cephali rigiduli ad apices ramorum 1-3. Disquis flavus. Radius albus seu carneus.

* *Foliis integerrimis subtùs lepidotis.*

1. O. ELLIPTICA, fruticosa, foliis alternis petiolatis ellipticis vix acutis basi attenuatis integerrimis suprà junioribus araneosis demùm glabris subtùs adpressè lepidotis subalbidis, corymbo terminali, capitulis longè pedicellatis. 3 in sylvis siccis umbrosis jugi montium Illawara Novæ Hollandiæ detexit cl. A Cunningham. Aster ellipticus A. Cunn. ! in litt. 1834. Habitus ferè Eurybiæ viscosæ sed pappus duplex Oleariæ (v. s. comm. à cl. invent.)

** *Foliis dentatis sinuatisve subtùs tomentosis.*

2. O. DENTATA (Moench l. c. Nees l. c.) fruticosa, foliis alternis petiolatis ovatis repando-dentatis crassis, pedunculis paucis 1-cephalis folio brevioribus. 3 in Novâ Hollandiâ. Aster dentatus Andr. bot. rep. 1. t. 61. Aster tomentosus Schrad. et Wendl. sert. hannov. 8. t. 24. Sieb. ! pl. exs. Nov. Holl. n. 341. Aster ferrugineus Wendl. in florâ 1819. p. 676. Hoffmans. verz. 1826. p. 61. (v. v. c. et s. sp.)
3. O. ROTUNDIFOLIA, fruticosa, foliis alternis breviter petiolatis ovato-rotundis, inferioribus basi cordatis suprà sparsè hirsutis subtùs ramisque densè strigiloso-hirsutis hinc indè subrepandis, pedunculis paucis ad apices ramorum 1-cephalis folio triplo longioribus. 3 in Novâ Hollandiâ ad Port-Jackson legit

- cl. Gaudichaud. *Diplopappus rotundifolius* Less. in *linnæa* 1831. p. 216. Proximè ex habitu ad priorem accedit. (v. s. comm. à cl. Gaudichaud.)
4. *O. QUERCIFOLIA*, fruticosa, foliis alternis subsessilibus obovato-oblongis obtusis suprà glabris rugosis subtùs pube stellatâ tomentosis sinuato-dentatis, dentibus latis obtusis, pedunculis axillaribus folio longioribus 1-3-cephalis. § in Novâ Hollandiâ (Sieb.), in paludosis rori cataractæ Campbell in montibus Cœruleis expositis (Cunn.). *Aster quercifolius* Sieb. pl. exs. Nov. Holl. n. 340. *Aster sinuatus* A. Cunn. in *lit.* 1834. Folia 15-18 lin. longa, 6-8 lin. ad apicem lata. Pedunculi bipollic. Achænia acutè pentagona adalta glabra. Pappi setæ subbiser. pallidè rufescentes scabræ, setæ exter. breves paucissimæ. (v. s.)
5. *O. PHLOGOPAPPA*, fruticosa, foliis alternis breviter petiolatis oblongo-lanceolatis suprà glabris subtùs tomentosis sinuato-dentatis, dentibus obtusis, corymbis paniculatis. § in Novâ Hollandiâ ad Cap. Van-Diemen. *Aster phlogopappus* Labill. Nov. Holl. sp. 2. p. 49. t. 195. *Aster phlogotrichus* Spreng. *syn.* 3. p. 515. *Eurybia quercifolia* Cass. dict. 16. p. 47. *Diplostephium phlogotrichum* Nees ast. 186. Ligulæ circ. 10 violaceæ 3 dentatæ. Achænia 5-angulata pubescenti-scabra. Pappus rufescens biserialis scaber. (v. s.)
6. *O. STELLULATA*, fruticosa, foliis alternis in petiolam brevem attenuatis oblongo-lanceolatis acuminatis suprà subscabridis subtùs pube stellatâ densè tomentosis grossè dentatis, dentibus mucronatis, corymbis paniculatis. § in Novâ Hollandiâ ad Cap. Van-Diemen. *Aster stellulatus* Labill. Nov. Holl. sp. 2. p. 50. t. 196. *Eurybia fulvida* Cass. dict. 16. p. 48. *Diplostephium stellulatum* Nees ast. 187. Proximè ad *O. phlogopappam* accedit. Folia 3-4 poll. longa, 4-6 lin. lata. Invol. squamæ imbricatæ lineares acutæ. Ligulæ ex Labill. circ. 15, et tubuli circ. 20. Achænia matura pauciora obovato-teretiuscula obtusè tetragona villosa. Pappus, setis biserialibus scabris subrufescentibus, exter. brevibus subpaleaceis. Recept. parcè convexum alveolatum. (v. s.)

CXXXVI. *DIPLOSTEPHIUM* Cass. dict. 37. p. 486. — *Dœllingeria* sp. Nees. — *Diplopappi* sp. Less. — *Asteris* sp. auct.

Capitulum multiflorum radiatum, ligulis fœmineis fertilibus 1-seriatis, disco hermaphrodito. Invol. subcylindraceum disco brevius, squamis pauciseriatis imbricatis adpressis coriaceis margine membranaceis. Recept. planum alveolatum, alveolis dentatis. Stigmata subulata hispida. Achænia elongata angulata hispidula. Pappus duplex, ext. setulosus brevis, inter. elongatus filiformis aut apice clavellatus scaber. — Herbæ perennes aut frutices asteriformes. Species omnes Americanæ. Folia alterna. Discus flavus. Ligulæ albæ aut cœruleæ. Sectiones duæ fortè in posterum genericè distinguendæ.

SECT. I. EUDIPLOSTEPHIUM.

Pappi interioris setæ plures apice clavellatæ. — Herbæ rhizocarpicæ Boreali-Americanæ facie Asteris.

1. *D. UMBELLATUM*, caule angulato scabro apice corymboso composito, foliis oblongo-lanceolatis utrinquè attenuatis margine scabridis, invol. imbricato, achæniis pubero-hirtis. 2½ in paludosis et ad fluminum ripas à Canadâ ad Carolinam. *Aster umbellatus* Ait. *kew.* ed. 1. v. 3. p. 199. Hoffm. *phytogr.* bl. p. 74. t. b. f. 2. ex Pers. *Chrysopsis amygdalina* Nutt. *gen.* 2. p. 153. excl. *syn.* *Aster amygdalinus* Ell. *sketch* 2. p. 367. *Dœllingeria umbellata* Nees ast. 178. Pappus rufus. Ligulæ albæ. (v. s.)
2. *D. AMYGDALINUM* (Cass. dict. 37. p. 486), caule subangulato glabro apice corymboso-paniculato, foliis lanceolatis oblongisve margine scabris, involucre laxo, ligulis 8-10, achæniis hirtellis. 2½ in umbrosis petrosis à Canadâ ad Ca-

rolinam. *Aster amygdalinus* Lam. diet. 1. p. 305. Lindl. bot. reg. t. 1517. *Chrysopsis humilis* Nutt. gen. 2. p. 153. excl. syn. *Aster humilis* Willd. sp. 3. p. 2038. *Aster divaricatus* Spreng. syst. 3. p. 529. excl. syn. *Dœllingeria amygdalina* Nees ast. 179. Ligulæ albæ. Pappus rufus, setis plerisque apice incrassatis. (v. v. et s.)

2. *humilis*, caule angulato glabro simplici dense folioso, foliis sessilibus utrinquè attenuatis, corymbo simplici oligocephalo. $\frac{1}{2}$ in Terrâ Novâ legit cl. de Lapilaye. (v. s.)

3. *D. cornifolium*, caule tereti glabro parçè paniculato, foliis elliptico-oblongis utrinquè acuminatis breviter petiolatis ciliatis integerrimis glabris, corymbi ramis nudis 2-3-cephalis, invol. laxè imbricato, achæniis glabris. $\frac{1}{2}$ in Pennsylvania (Muhl.), à Canadâ ad Carolinam (Michx.). *Aster cornifolius* Muhl. in Willd. sp. 3. p. 2039. *Aster infirmus* Michx. fl. Bor. Am. 2. p. 109 (certè ex Lindl.). *Aster humilis* Ell. sketch 2. p. 366. — Pluk. alm. t. 79. f. 1. Ligulæ 7-8 albidæ. Pappus rufus, ext. brevis. E fol. basi decurrentiæ tenuissimæ interdum puberulæ. Achænia obovata subcompressa glabra. (v. s. comm. à cl. Torrey.)

4. *D. acuminatum*, caule simplici angulato-flexuoso scabro, foliis lato-lanceolatis basi angustatis et integerrimis apice grossè serratis acuminatis margine scabridis cæterum glabriusculis subsessilibus, paniculâ corymbosâ laxâ, pedunculis nudis folio longioribus, invol. squamis linearibus laxis disco brevioribus, achænio sparse glanduloso-pubero. $\frac{1}{2}$ in sylvis Amer. bor., in Canadâ (Michx.), agro Bostoniensi (Big.), Noveboracensi (Tor.), ad saltus Niagaræ. (Janin!) in montibus Alleghany (Raf.). *Aster acuminatus* Michx. fl. Bor. Am. 2. p. 109. Hook. bot. mag. t. 2707. *A. diffusus acuminatus* Pers. Ligulæ albæ. Pappus pallidè rufus, setis exter. brevibus paucis. Affinis *D. cornifolio*. (v. s.)

5. *D. dichotomum*, caule pubescente, foliis arcè sessilibus ovalibus obtusis pubescentibus, corymbi dichotomi ramulis nudis elongatis, invol. squamis lineari-lanceolatis pubescentibus achænio valdè hirsuto vix longioribus. — in humidis pinguibus insulæ Paris Amer. bor. *Aster dichotomus* Ell. sketch 2. p. 366. Ligulæ 10-16 albæ purpureo tinctæ. Pappus duplex.

6. *D. obovatum*, totus brevè et molliter pubescens, caule tereti subsimpliciter corymboso, foliis obovatis obtusis integerrimis subsessilibus margine scabris, invol. imbricato pubescente, ligulis circ. 10, achænio angulato hispido. $\frac{1}{2}$ in Georgia fruticetis ad margines paludum propè Savannah. *Chrysopsis obovata* Nutt. gen. 2. p. 152. *Diplostephium boreale* Spreng. syst. 3. p. 544. excl. syn. *Dœllingeria obovata* Nees ast. 182. *Aster obovatus* Ell. sketch 2. p. 368. Ligulæ et disci cor. ochroleucæ. Pappus ferrugineus duplex.

Sect. II. AMPHISTEPHIUM. — *Diplostephium Kunth nov. gen. am. 4.*

p. 97.

Pappi interioris setæ filiformes nec clavellatæ. — Frutices Mexicani foliis linearibus. — Sectio media inter *Diplostephium* et *Diplopappum*, fortè genericè separanda, sed mihi non satis nota.

7. *D. MORANENSE* (Nees ast. 198), fruticosum, ramis foliosis glabriusculis, foliis linguato-linearibus calloso-mucronatis planis subcarnosis, infer. apicem versus mucronato-dentatis, ramis florigeris subcorymbosis minutè foliosis, invol. hemisphærici imbricati disco brevioris squamis inf. subulatis, cæt. lauceolato-ensipidatis. $\frac{1}{2}$ in montibus Mexicanorum propè fodinam Moran rarissimum. *Aster Moranensis* H. B. et Kunth nov. gen. am. 4. p. 93. Ligulæ paucæ albæ siccitate revolutæ. Achænia compresso-quinquangularia hirta-sericea. Pappus duplex exterior setulosus.

8. *D. LAVANDULÆFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 97. t. 335), fruticosum, foliis lineari-lanceolatis margine revolutis integerrimis subtus caule involucrique albo-tomentosis, capitulis racemoso-confertis. $\frac{1}{2}$ in arenosis

exustis ad radices montium ignivomorum Ilinisæ et Cotopaxi prope Mulalo. Nees ast. 189. Aster lavendulacens Willd. herb. Frutex orgyalis. Capit. 9-10 lin. diam. Ligulæ 30-35 albæ. Discus pallidè lentus. Stigm. lineari-subulata. Pappus duplex, ext. paleaceus brevis, int. setosus ferrugineus. Achæn. verisimil. non compressa. An hujus generis?

† *Species exclusæ.*

- D. ACULEATUM Nees = Eurybia aculeata.
- D. EXTENUATUM Nees = Diplopappus extenuatus.
- D. FILIFOLIUM Nees = Diplopappus filifolius.
- D. FRUTICULOSUM Nees = Diplopappus fruticosus.
- D. GRANDIFLORUM Spr. = Andromachia grandiflora.
- D. HISPIDUM Nees = Diplopappus hispidus.
- D. IGNIARIUM Spreng. = Andromachia igniaria.
- D. LIRATUM Nees = Eurybia lirata.
- D. MICROPHYLLUM Nees = Eurybia microphylla.
- D. MELASTOMOIDES Spreng. = Andromachia melastomoides.
- D. NUBIGENUM Spreng. = Andromachia nubigena.
- D. OBTUSATUM Nees = Diplopappus obtusatus.
- D. OCHRACEUM Nees = Tetramolopium ochraceum.
- D. PHLOGOTRICHUM Nees. = Olearia phlogopappa.
- D. RAMULOSUM Nees = Eurybia ramulosa.
- D. SESSILIFLORUM Spr. = Andromachia sessiliflora.
- D. SOLIDAGINEUM Spr. = Andromachia solidaginea.
- D. STELLULATUM Nees = Olearia stellulata.
- D. TERETIFOLIUM Nees = Diplopappus filifolius β .
- D. VOLUBILE Spr. = Andromachia volubilis.

CXXXVII. CALLISTEPHUS Cass. dict. 37. p. 464 et 491. Nees ast. 221. — Calistemma Cass. bull. philom. 1817. p. 32. dict. 6. suppl. 45. atl. et op. phyt. t. 7. — Asteris sp. Linn. — Diplopappi sp. Less. — Chrysopsidis sp. Nutt.?

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis in statu natur. 1-serialibus, disci hermaphroditis tubulosis 5-dentatis. Recept. latum subconvexum breviter alveolatum. Invol. 3-4-seriale patens, squamis ciliatis obtusis, bracteis foliaceis involucri brevioribus cinctum. Stigmata disci apice ovata. Achænia obovato-cuneiformia compressa hirtella. Pappus duplex uterque 1-serialis, exter. brevissimus paleaceo-setaceus setis in coronam subconcretis, inter. setis longis filiformibus scabris deciduis. — Herba Chinensis annua erecta ramosa, ramis apice 1-cephalis. Folia alterna sessilia. Capitula ampla aut floribus omnibus ligulatis aut omnibus tubuloso-ampliatibus plena, disco flavo, ligulis tubulisve ampliatis omnes colores flavo excepto exhibentibus.

§. 1. *Invol. squamis exterioribus patulis.*

1. C. CHINENSIS (Nees l. c.), foliis infer. spathulatis petiolatis, mediis rhombo-lanceolatis, summis oblongis, omnibus grosse dentatis plus minus ciliatis, ramis 1-cephalis elongatis, invol. squamis foliaceis subquarrosis longè ciliatis. (1) in Chinâ et Japoniâ, nunc in hortis totius orbis cultus, gallicè *Reine Marguerite*. Aster Chinensis Linn. sp. 1232. Calistemma hortense Cass. bull. phi-

- lom. 1817. p. 32. dict. et opusc. atl. t. 7. *Callistephus hortensis* Cass. dict. 37. p. 491. *Diplopappus* Chineseus Less. syn. 165. — Dill. elih. t. 34. (v. v. c.)
2. *C. P. BERNIS* (Lindl. in litt. 1835), foliis integerrimis oblongo-linearibus ciliatis, ramis elongatis monocephalis, invol. squamis lanceolato-linearibus acuminatis ciliatis æqualibus disco longioribus squaroso-patentibus. (2) in Sibiria. *Aster* Gmel. fl. sib. 2. p. 189. t. 81 excl. descr. Species mihi planè ignota. †.

§. 2. *Invol. squamis exterioribus erectis.* — *Poloa* DC. in *Guill.* ann. 1833. v. 2. p. 514 et in *Wight contr.* p. 10.

3. *C. WIGHTIANUS*, foliis sessilibus oblongo-linearibus integris aut subserratis breviter mucronatis, ramis foliosis apice compressis minutè puberulis, invol. squamis exter. foliaceis lineari-oblongis eciliatis vix interiores superantibus. — in Ind. orient. ad Dindigul legit cl. Wight. *Erigeron* Wightianum Wall. cat. n. 3284. comp. n. 394. *Erigeron* dubium Spreng. in litt. ad Wall. *Buphthalmum* Wight! herb. *Poloa* Wightiana DC. l. c. Variat glabriusculus aut subpubescens. Ligulæ fertiles aut rarius substeriles, pappo ext. nunc cupuliformi nunc subnullo. (v. s. comm. ab hon. Cœt. merc. angl. Ind. or.)
4. *C. HEYNEANUS*, foliis sessilibus oblongis acutè serratis pube minutissimâ utrinquè ramisque hirtellis, invol. squamis exter. foliaceis lineari-oblongis eciliatis inter. paulò superantibus. — in Indiâ orient. legit cl. Heyne. *Aster* Indicus Heyne in Wall. cat. n. 2971. comp. n. 81 non Linn. Species priori proxima sed hirsutæ brevi et confertâ, foliis serratis, ramis non compressis et habitu diversa. (v. s. comm. ab hon. Cœt. merc. angl. Ind. or.)

† *Species exclusa.*

C. INDICUM Don. = *Asteromæa Indica*.

CCXXVIII. *DIPLOPAPPUS*. — *Diplopappi* sp. *Cass. dict.* 13. p. 308. *Less. syn.* 163. — *Rhinactina* Less. syn. 162, non Willd. — *Asteris* sp. *auct.* — *Diplostephii* sp. *Nees*.

Capitulum multiflorum radiatum, ligulis 1-serialibus fœmineis, fl. disci hermaphr. 5-dentatis. Recept. planum subfoveolatum. Invol. squamæ imbricatæ. Achænia oblonga compressa. Pappus disci et radii duplex scabridus, ext. brevis setulosus, inter. longus setosus. — Herbæ aut suffrutices patriâ et habitu varii. Genus (forsan idè in posterum dividendum) differt à *Diplostephio* achæniis compresso-planis nec teretiusculo-angulatis; à *Callistepho* achænio evidentius compresso et pappo exter. setoso nec coroniformi; à *Chrysopside* pappo exter. potiùs setoso quàm paleaceo et ligulis à disco discoloribus albis cœruleis seu purpureis nec coloribus flavis. — An fortè *Asterosperma* Less. syn. 389 mihi ignotum hùc etiam referendum?

§. 1. *Pedunculares.* — *Diplostephia peduncularia* *Nees ast.* 195; *caulibus fruticosis, invol. imbricati squamis adpressis lined dorsali elevatâ et glandulis 1-2 secus lineam donatis margine subscariosis foliis 1-nerviis glabris integerrimis.* (1)

1. *D. FRUTICULOSUS* (Less. in linnæâ 1830. p. 171. syn. 164), foliis lineari-cuneiformibus integerrimis planis glanduloso-punctatis glabris, ramulis lateralibus apice nudis 1-cephalis, pedunculis folia triplò superantibus, involucrio discum æquante, ligulis lanceolatis involucrio duplò longioribus. 3 ad Cap. Bonæ-Spei

(1) Si aliquis vellet aut quinque hujus divisionis species in unicam conjungere aut ex iis genus proprium conficere, non repugnarem.

(Eckl. et Zeyh.). *Diplostephium fruticosum* Nees ast. 194. *Diplostephium longipes* Cass. dict. 56. p. 171. *Aster fruticosus* Linn. sp. 1225. Thunb. fl. cap. 687. Burm. prod. 27. *Aster fruticosus* Willd. sp. 3. p. 2018. Jacq. fragm. p. 9. t. 5. f. 4. Sims bot. mag. t. 2286. — Comm. h. Amst. 2. t. 27. Pedunculi pollicares ramos præsertim laterales terminantes fastigiatum dispositi. Ligulæ filicinæ seu albidæ. (v. s.)

β. ligulis lateraliter ad apicem dente acuto utrinquē incisus elongatis linearibus. — Hook. bot. mag. t. 2718.

2. *D. EXTENUATUS* (Nees ast. 195 sub *Diplostephio*), foliis lineari-cuneiformibus integerrimis planis glabris grossè glanduloso-punctatis, ramulis lateralibus apice nudis 1-cephalis, pedunculis folia triplò superantibus, invol. disco breviorè, ligulis lanceolatis involucri duplò longioribus. ♂ ad Cap. Bonæ-Spei ad Glænkloof (herb. Deless.) et in monte Tabulari (Eckl. in herb. Donant). *Aster fruticosus* var. auct. *Aster coriaceus* hortul. fortè *Diplostephii fruticosi* mera varietas loco pinguiori nata. Huc potiùs quàm *D. obtusatus* pertinet icon Pluk. mant. t. 330. f. 19. (v. s.)

3. *D. OBTUSATUS* (Less. in linnæa 1831. p. 118), foliis lineari-cuneatis integerrimis planis glabris pellucido-punctatis, ramulis lateralibus apice 1-cephalis, pedunculis subnudis vix folia superantibus, involucri disco breviori, ligulis lineari-oblongis involucri vix longioribus. ♂ ad Cap. Bonæ-Spei ad Blear-Bery ex herb. Deless. *Diplostephium obtusatum* Nees ast. 196. *Aster obtusatus* Willd. sp. 3. p. 2018. excl. syn. Thunb. *Aster rosmarinifolius* Spreng. syst. 3. p. 526. — *Aster fruticosus* β Willd. sp. 3. p. 2018 et reverà fortè *D. extenuatus* varietas brevi-pedunculata. Variat foliis obtusis aut acutiusculis. (v. s.)

4. *D. FILIFOLIUS*, foliis lineari-filiformibus glanduloso-punctatis glabris, ramis lateralibus apice 1-cephalis, pedunculis subnudis folia duplò superantibus, involucri disco subæquante, ligulis involucri duplò longioribus. ♂ ad Cap. Bonæ-Spei (Eckl. et Zeyh.) Dreg. ! *Diplostephium filifolium* Nees ast. 197. *Aster filifolius* Vent. malm. (1804). t. 82. — *Aster* Eckl. coll. exs. Ligulæ albæ. Folia 6-8 lin. longæ. Pedunculi pollic. et sesquipollic. Glandulæ foliorum ut scrobuli tantum apparent. (v. s.)

β. *teretifolius*, foliis pedunculis ligulisve brevioribus. ♂ ad Cap. Bonæ-Spei. *Diplost. teretifolium* Nees ast. 198. *Diplopappus teretifolius* Less. in linnæa 1831. p. 118. *Aster fruticosus* Krebs! coll. exs. n. 192. *Aster fruticosus* var. Thunb. fl. cap. 687. (v. s.)

5. *D. ELONGATUS*, foliis lineari-filiformibus eglandulosis glabris, axillis nudis, ramis apice longè denudatis monocephalis folia triplò superantibus, invol. squamis dorso ad apicem glandulosis, ligulis involucri duplò longioribus. ♂ ad Cap. Bonæ-Spei ad Zneuroberge alt. 200-300 ped. legit cl. Dreg. ! Affinis *D. filifolio* sed folia non fasciculata pollicem nec semipollicem longæ. Pedunculi tripollicares. Invol. disco dimidio brevius, etc. (v. s.)

§. 2. *Calimeridei, caulibus herbaceis, foliis latis sæpiùs trinerviis et dentatis, invol. squamis pauciserialibus vix imbricatis eglandulosis.*

6. *D. ASPER* (Less. syn. 163), caule herbaceo adscendenti-erecto simplici scabrido apice longè aphyllò 1-cephalo, foliis oblongo- aut ovali-lanceolatis triplinerviis integerrimis aut dente grosso 1-2 hinc indè notatis utrinquē asperis ciliatis. invol. squamis disco subæqualibus planis acuminatis margine scariosis. ♀ ad Cap. Bonæ-Spei in graminosis prope Galgebosch (Th.), prope Hangklipp (Mundt et Mair.), in Cafraria (Krebs! coll. n. 173), in col. orient. non longè à flum. Ganritz (Burch! n. 4740), in distr. Albany et in territorio cesso (Eckl. et Zeyh.), in Zneuroberge et Willberge (Dreg. !). *Calendula hispida* Thunb. cap. 704 et *Leyssera pilosella* ejusd. herb. sp. dextra. et læv. ex Less. *Diplostephium hispidum* Nees ast. 163. *Capitulum* ferè *Asteris alpini*. (v. s.)

7. *D. ROYLEI*, caule herbaceo erecto subsimplici pedunculisque hispidis, foliis

oblongis acuminatis sessilibus aliis integerrimis aliis apice 3-5-dentatis supra glabris subtus et margine scabris obscure trinerviis, invol. squamis glabris disco brevioribus margine scariosis. — in Indiâ prov. bor. occid. legit cl. Royle. Valdè accedit ad *D. asperum*. Ligulæ extus revolutæ ex sicco albidæ. Achænia villosa. Pappus crassè barbellatus. Cl. Lindley in adn. mss. habet *Diplopappum* quemdam Roylei fortè à nostro diversum sed mihi ignotum. (v. s. comm. à cl. invent.)

8. *D. molliusculus* (Lindl. adn. mss.), caulibus simplicibus adscendentibus moncephalis pubescenti-scabris, foliis oblongis integerrimis scabriusculis subtus strigoso-pilosis, capitulis longè pedunculatis, invol. hemisphærici squamis acuminatissimis, exter. basi pubescentibus, inter. membranaceis. $\frac{1}{2}$ in Indiâ orient. ad Sirmore. *Aster molliusculus* Wall.! cat. et herb. n. 2972. comp. n. 82. (v. s. comm. ab hon. Coet. merc. angl. Ind. or.)
 9. *D. ASPERULUS*, caule herbaceo erecto dichotomè ramoso tereti glabro, foliis ovatis acuminatis grossè serratis penninerviis ciliatis supra sparè scabris, radicalibus petiolatis, caul. basi cuneatis, corymbo laxo, pedicellis nudis, invol. squamis oblongo-linearibus obtusis glabris, ext. foliaceis laxis, int. disco brevioribus subscariosis. $\frac{1}{2}$ in montibus Indiæ orientalis ad Kamaon. *Aster asperulus* Wall.! cat. et herb. n. 2968. comp. n. 78. Nees ast. 281. Achænia sparè villosa. Ligulæ elongatæ. (v. s. comm. ab hon. Coet. merc. angl. Ind. or.)
 10. *D. ASPERRIMUS*, caule herbaceo stricto ramoso scabro, foliis elliptico-ovatis lanceolatisve acuminatis sessilibus trinerviis utrinquè scaberrimis, corymbo patulo foliato, invol. squamis oblongis obtusis adpressis ciliatis, achæniis sericeo-villosis, pappi setis exter. paucis brevibus. $\frac{1}{2}$ ad Narrain-Hotty et alibi in Nepaliâ legit cl. Wallich. *Aster asperimus* Wall.! cat. et herb. n. 2970. comp. n. 80. *Aster trinervius* Roxb. cat. h. calc. 61. Don prod. fl. nep. 177. *Aster asper* Ham. herb. Galatella asperima Nees ast. n. 173. Cineraria canescens Wall.! herb. Ligulæ fertiles à Galatellâ excludunt. Recept. in centrò alveolato-simbrilliferum. Achænia plano-compressa nec ut in *Diplostephio* angulata. Pappus rufus primâ fronte 1-serialis sed, mon. cl. Lindl., verè sed obscure biserialis, setulis inter achæniâ villosis super. subconfusis. Habitus ferè *Dœllingeriæ* sed setæ nec clavellatæ nec 1-seriales. (v. s.)
 11. *D. CAUCASICUS*, caule herbaceo erecto striato pubescenti-scabro, foliis ovato-sen oblongo-lanceolatis serratis acuminatis basi trinerviis utrinquè (præs. margine et secus nervos) scabridis, capitulis 1-7 caulem ramosos terminantibus, invol. subæqualis squamis lineari-acuminatis glabriusculis ciliatis, achæniis subtrigonis hirtis, pappi setis exter. paucis brevibus. $\frac{1}{2}$ in alpinis Caucasi (M. Puschk.) ad Gurjel (Szow.! ex Fisch. et M.). *Aster caucasicus* Willd. sp. 3. p. 2020. Bieb. fl. et suppl. n. 1729. Nees ast. p. 35. Pappus duplex ritè observ. cl. Meyer. Ligulæ purpurascens. Herba pedalis. (v. s. comm. à cl. Fisch. et Mey.)
- §. 3. Amelloidei, Nees ast. p. 199, sub *Diplostephio*; caule herbaceo-suffrutescente, foliis linearibus rigidulis ciliato-scabris, involucris eglandulosis imbricatis, capitulis subcorymbosis.
12. *D. LINARIÆFOLIUS* (Lindl.! adn. mss.), caule stricto apice racemoso, foliis linearibus rigidis glabris margine scaberrimis serrulato-ciliatis, invol. turbinati squamis linearibus acutis patulo-imbricatis margine coloratis. $\frac{1}{2}$ in Amer. bor. spec. in agro Noveboracensi. *Aster linariifolius* Linn. sp. 1227. *Aster rigidus* Michx. fl. bor. am. 2. p. 110. *Diplostephium linariifolium* var. Nees ast. 199. *Chrysopsis linariifolia* Nutt. gen. 2. p. 152? Ligulæ cœruleæ. Achænia compressa villosa. Caulis imâ basi suffrutescens. (v. s.)
 13. *D. RIGIDUS* (Lindl. adn. mss.), caule stricto racemoso, ramis elongatis 1-cephalis, foliis linearibus rigidis margine scaberrimis serrulato-ciliatis, invol. hemisphærici squamis linearibus acutis, exterioribus laxis à foliis pedicelli parum diversis. $\frac{1}{2}$ in Amer. bor., spec. ex Carolinâ et Georgiâ recepti. ~~Nees~~

rigidus Linn. sp. 1228. Aster linarifolius Michx. fl. 2. p. 110. Aster pulcherrimus Lodd. bot. cab. 1. t. 6. Diplostephium linarifolium var. Nees ast. 199. — Icon Pluk. t. 14. f. 7. huc fortè pertinet. — Cl. Lindleyum qui specim. Linnæana vidit secutus sum in dist. hujus et linarifolii. (v. s.)

14. *D. INCANUS* (Lindl. bot. reg. t. 1693), totus incanus, caule suffruticoso, ramis longis corymbosis 1-cephalis apice nudis, foliis linearibus mollibus distantibus obtusis mucronulatis, invol. ovati glandulosi squamis linearibus multiseriatis, exter. squarrosis. $\frac{1}{2}$ in Californiâ et ad oram occid. Americæ bor. Hook. bot. mag. t. 3382. Caulis $2\frac{1}{2}$ -pedalis à basi ramosus pube brevi mollis, foliis valde denudatis. Ligulæ amœnè violaceæ. Proximus *D. rigido* et *linarifol.* Ovarium teres villosum, undè potius ad *Diplostephium* referendus? (v. s.)

15. *D. LUTESCENS* (Lindl. adn. mss.), caule simplici corymboso, ramulis sub-1-cephalis subaphyllis, foliis oblongo-linearibus basi longè angustatis subintegris scabris, invol. pubescentis squamis oblongo-lanceolatis obtusis. $\frac{1}{2}$ in Americâ arcticâ. Radius lutescens. Proximus *Heleastro albo* ex Lindl. An fortè *Chrysopsidis* sp.?

16. *D. LEUCOPHYLLUS* (Lindl. adn. mss.), fruticosus? undiquè lanatus, ramis brevibus 1-cephalis, foliis crassis ovalibus acutis crenatis in petiolum angustatis, invol. squamosi squamis linearibus membranaceis, superioribus calvis. $\frac{1}{2}$? in Californiâ. Pappus exter. brevis, inter. valde inæqualis subulatus. Achænia tomentosa fusiformia. (Lindl.)

+ Species exclusæ.

- D. CHAMISSONIS* Less. = *Aplopappus Chamissonis*.
D. CHINENSIS Less. = *Callistephus Chinensis*.
D. CHRYSANTHEMIFOLIUS Less. = *Aplopappus chrysanthemifolius*.
D. CORONOPIFOLIUS Less. = *Aplopappus coronopifolius*.
D. DELPHINIFOLIUS Cass. = *Polyactidium delphinifolium*.
D. DUBIUS Cass. = *Stenactis annua*.
D. ERICOIDES Less. = *Aplopappus ericoides*.
D. GRAMINIFOLIUS Less. = *Neja gracilis*.
D. PRUNELLOIDES Less. = *Aplopappus prunelloides*.
D. ROTUNDIFOLIUS Less. = *Olearia rotundifolia*.
D. SERICEUS Less. = *Aplopappus sericeus*.

CXXXIX. RHINACTINA Less. syn. 162 non Willd.

Omnia *Diplopappi* sed corollæ disci inæqualiter 5-dentatæ et ideo ferè subbilabiata, pappi serie externâ paucisetâ. — *Herbæ Sibiricæ* perennes, rhizomate lignoso. Folia glaucescentia brevissimè subhirtella integerrima, radicalia obovata in petiolum attenuata obtusa aut submucronata, caulina pauca. Caulis simplex 1-cephalus aut parçè ramosus oligocephalus. Invol. squamæ siccæ glabræ. Achænia villosa. Ligulæ cœruleæ. — An genus à *Diplopappo* satis diversus? Habitus ferè *Asterum* alpigenorum sed pappus duplex.

1. *R. LIMONIFOLIA* (Less. in linnæâ 1831. p. 119. syn. p. 162), foliis radicalibus obovatis obtusis trinerviis longè petiolatis, caulinis ovali-oblongis sessilibus, caulibus capitula 2-4 longè pedicellata gerentibus, invol. squamis obtusiusculis. $\frac{1}{2}$ in rupestribus graminosis Sibiricæ Altaicæ ad montem Kurtzschum. *Aster obovatus* Mey. in Ledeb. fl. alt. 4. p. 95. Ledeb. ic. alt. 5. p. 492. (v. s. comm. à cl. Meyer.)

2. *R. UNIFLORA* (Bunge ex herb. Acad. imp. Petrop.), foliis radicalibus oblongo-obovatis acutis uninerviis breviter petiolatis, caulinis paucis oblongis basi attenuatis, caulibus simplicibus r-cephalis, invol. squamis acuminatis. ♀ in montibus Altaicis. Caules floridi vix bipollicares folia radicalia paululum superantes.

CXXX. NOTICASTRUM.

Capitulum radiatum, ligulis r-serialibus fœmineis fertilibus linearibus, fl. disci tubulosis 5-dentatis bisexualibus sterilibus. Invol. 3-seriale, squamis linearibus acuminatis, extimis brevibus, intermed. coloratis et intimis subscariosis disco ferè longioribus. Recept. planum alveolatum. Antheræ ecaudatæ. Styli rami in radio et in disco elongati subulati dorso hispidi. Achænia oblongo-linearia pube sericeâ villosa sessilia subrostrata, radii fertilia compressa nervoso-striata, disci angusta sterilia. Pappus pluriserialis persistens, setis rigidulis scabris extimis multò minoribus. — Herba perennis Chilensis puberulo-glandulosa. Rhizoma frutescens. Caules subsimplices teretes adscendentes 6-7 poll. Folia radicalia in petiolum attenuata, caulina sessilia subsemiamplexicaulia lanceolata mucrone calloso apiculata integerrima. Capitula pauca ad apices ramulorum solitaria subcorymbosa. Ligulæ rubentes disco duplò longiores integræ, juniores convolutæ. Discus flavus. — Genus ex disco sterili et stigmatibus subulatis distinctissimum. Affinitas subdubia. Nomen ex *νοτίκος*, australis, et *αστρον*, quasi Aster australis.

1. *N. ADSCENDENS*. ♀ in Cordilleris Chilensibus legit indef. Claud. Gay et mecum commun. (v. s.)

CXXXI. DISTASIS.

Capitulum multiflorum heterogamum, fl. radii r-serialibus ligulatis fœmineis, disci hermaphrod. tubulosis 5-dentatis. Recept. angustum breviter alveolatum. Invol. squamæ 2-3-seriales subæquales lineares acutæ. Antheræ ecaudatæ? Achænia subcompressa hirtella. Pappus duplex uterquè 4-5-paleaceus, ext. paleis ovali-oblongis brevibus, inter. setis piliformibus scabris. — Herba Mexicana ramosissima puberula verisim. annua. Folia caulina alterna sessilia linearia obtusa integerrima basi attenuata. Capitula ramos terminantia solitaria laxè paniculato-corymbosa. Ligulæ albidæ. Discus flavus. — Genus si verè antheræ ecaudatæ affine Neja sed diversum ligulis uni- nec pluriserialibus, pappo pauciseti et habitu: si anth. forsan caudatæ affine Pentanemæ et Francoeurix. Ex hac dubitatione Distasis (gracè dubitatio) dicta!

1. *D. MODESTA*. (1)? in Mexico ad Laredo legit cl. Berlandier (pl. exs. n. 1416). Capitula vix 3 lin. diam. Folia 3-4 lin. longa. Ligulæ revolutæ. (v. s. comm. & cl. inv.)

Subdiv. 3. ERIGERÆ.

Ligulæ pluriserialis. Pappus pilosus, nunc r-serialis, nunc bi- seu pluriserialis, serie externâ equali aut breviori.

CXXXII. MELANODENDRON. — *Solidaginis* ap. Roxb.

Capitulum multiflorum heterogamum, fl. radii pluri- (3-5-) serialibus fœmineis angustè ligulatis, disci tubulosis hermaphrod. 5-dentatis. Recept. planum ebracteolatum tenuiter alveolare. Invol. pluriserialis imbric., squamis lanceolatis acuminatis disco subbrevioribus. Stigm

exserta Asterearum. Anthæræ basi muticæ. Achænium compresso-tetragonum glabrum erostre. Pappus 1-serialis, setis scabridis. — Arbor insulæ Sanctæ-Helenæ maxima glabra. Folia alterna oblonga basi attenuata integerrima obtusa coriacea obscure viridia et ideò arbor anglicè dicta *Black-Cabbage-tree* undè græcè *Melanodendron* dixi. Capitula numerosa pedicellata in corymbum amplum disposita. Ligulæ albæ. — Genus *Erigeronti* valdè affine sed habitu diversissimum et achæniis classicè distinctum.

1. *M. INTEGRIFOLIUM*. 5 in insulæ Sanctæ-Helenæ arenosis. *Solidago integrifolia* Roxb. in *Beats* app. 323 non Desf. Arbor insulæ maxima, trunco 5-6 ped. ambitu, comâ ramosâ amplâ, ramis lævibus, foliis nigricantibus, floribus albis, ligulis 20-30. Achænia subtetragona glabra. (v. s. comm. à cl. Burchell.)

CXXXIII. *LEPTOCOMA* Less. in *linnæâ* 1831. p. 130. syn. p. 188. Nees ast. p. 220. — *Laveniæ* sp. Wall.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus foemineis minimis ligulatis integris, disci paucis tubulosis 5?-dentatis. Recept. planum brevissimè fimbriatiferum. Invol. campanulatum subbiseriali, squamis ellipticis acutis. Stylus ♀ bifidus ligulæ æqualis, ♂ brevis inclusus. Achænia plano compressa glaberrima costâ marginata disco parvo calloso, marginalia rostrata, centralia erostrata. Pappus 2-4-setis tenuissimis flexuosis deciduis constans. — Herba erecta rigidula vix puberula Nepalensis. Folia alterna brevissimè petiolata elliptico-lanceolata utrinquè acuminata integerrima aut hinc indè calloso-serrulata triplinervia suprâ sparsè puberula subtus pallidiora glanduloso-punctata in nervis pubentia. Pedunculi axillares distantes 1-cephali juniores breves reflexi, adulti longiores patentes racemum foliaceum rigidum terminalem constituentes, sed capitula infer. seriis florida! Flores albi parvi. — Genus affine *Lagenophoræ* et *Brachycomi* sed pappo diversum.

1. *L. RACEMOSA* (Less. et Nees ll. cc.). 2½ in Nepaliâ detexit cl. Wallich. *Lavenia* rigida Wall.! cat. et herb. n. 325. Rhizoma lignosum subtherosum. (v. s. comm. ab hon. Cœtu merc. angl. Ind. or. et à cl. Royle.)

CXXXIV. *VITTADINIA* Rich. et Lesson fl. nov. Zeel. 250. — *Brachycome* Gaudich. ? non Cass.

Capitulum multiflorum heterogamum, fl. radii 1? 2-3-serialibus ligulatis foemineis, disci tubulosis cylindricis 5-dentatis hermaphroditis aut styli abortu masculis. Recept. planiusculum nudum. Invol. subbiseriali, squamis subæqualibus linearibus margine scariosis. Anthæræ basi ecaudatæ. Achænium elongatum basi attenuato substipitatum terejusculum striatum. Pappus 1-serialis, setis confertissimis scabridis. — Herbæ Australasicæ diffusæ aut erectæ. Folia alterna dentato-lobata rariùs integra. Rami apice 1-cephali. Ligulæ purpurascèntes aut albæ. Pappus rufus.

§. 1. *Novo-Zeelandicæ, ligulis fortè 1-serialibus.*

1. *V. AUSTRALIS* (Rich. et Less. fl. nov. Zeel. p. 250), caule à basi ramoso diffuso, foliis obovato-spathulatis grossè dentatis basi attenuatis ciliatis, summis subintegerrimis, ramis floridis foliosis. 2½ in Novâ-Zeelandiâ loco dicto *Passo-des-Français* (Lesson!) et in valle Wangarôâ (A. Cunn.). *Brachycome* spathulata

Gaudich. voy. Freyc. bot. p. 467? Ligulæ 1-2-seriales. Herba variat forè glabra aut puberula. (v. s.)

§. 2. Novo-Hollandicæ, ligulis 2-3-serialibus.

2. *V. TRILoba*, caule erecto ramoso puberulo-hirtello, foliis glabriusculis basi cuneatis apice trifidis, lobis oblongis integerrimis submucronulatis, ramis floridis apice nudis. $\frac{1}{2}$ in Novæ-Hollandiæ orâ orientali prope Port-Jackson. Brachycome triloba Gaudich. voy. Freyc. bot. p. 467. (v. s.)
3. *V. CUNEATA*, caule erecto ramosissimo vix pubescente, foliis angustè cuneatis obtusis integerrimis pubescentibus, pedunculis apice nudis. $\frac{1}{2}$ in herbosis ad flumen Lachlan in parte interiore Novæ-Cambiæ australis legit cl. A. Cunningham aprili flor. Ligulæ parvæ 2-3-seriales. Pappus pallescens. Folia 3-6 lin. longa ad axillas sæpè fasciculata. (v. s. comm. à cl. A. Cunningham.)
4. *V. SCABRA*, caule ascendente minutè puberulo, foliis linearibus aut infer. elongato-lanceolatis integerrimis aut infer. denticulatis confertis utrinquè pubero-scabris, ramis floridis subnudis. $\frac{1}{2}$ in collibus scopulosis ad Cleveland-bay in Novâ-Hollandiâ tropicâ junio flor. legit cl. A. Cunningham. Ligulæ 2-3-seriales forè albæ. Folia pollicem et ultra longa, 1-2 lin. lata. (v. s. comm. à cl. Cunningham.)
5. *V.?* *DENTATA*, caule. . . , foliis linearibus integerrimis vel remotè denticulatis apice tricuspidatis sublanatis. — cum priore. Brachycome dentata Gaudich. l. c.

CXXXV. FULLARTONIA.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus fœmineis angustè ligulatis, disci tubulosi 5-dentati hermaphrod. Recept. nudum. Invol. polyphyllum squamis linearibus acuminatis 1-2-seriatis extus bracteis foliaceis proximè cinctum. Antheræ ecaudatæ. Stylus radii apice bifidus glaber, disci indivisus apice capitatus truncatus subbarbatus. Achænia erostria teretiuscula, radii glabra calva, disci pilosula pappo 1-seriali scabro rufo coronata. — Herba ex Indiâ boreali-occid. orta erecta supernè pilis sparsis apice glandulosis hispida. Folia alterna semiamplexicaulia ovalia, super. oblonga grossè dentata. Capitula ad apicem caulis et ramorum axillarium solitaria. Flores ex sicco pallidè flavi seu albid. — An ex ligulis pluriserialibus et achænia radii calva ad Astereas referenda? An ex stigmatè apice truncato et subbarbato potiùs ad Senecioneas? — Nomen ex cl. Fullarton peregrinatore Anglo qui Indiam totam et speciatim prov. Kamaonensem peragravit et sæpè à Geographis citatur.

1. *F. KAMAONENSIS*. — in Kamaon prov. Indicâ ad radices Himalayæ. (v. s. sine nom. et nom. comm. ab ill. Aulæ merc. Ind. or.)

CXXXVI. POLYACTIDIUM. — *Polyactis Less. syn.* (1832). p. 188 non *Link* 1809. — *Stenaectis Cass. dict.* 50. p. 483 excl. *sp. plur.*

Capitulum multiflorum radiatum, ligulis fœmineis angustissimis numerosissimis circ. 2-serialibus, disco hermaphr. Recept. nudum platum minutissimè punctulatum. Invol. squamæ lineares 2-3-seriales subæquales. Achænium compressum, disco epigyno minuto. Pappus duplex, ext. brevissimus coroniformis dentatus setulis subadpressis, interior paucisetosus elongatus scaber. — Herba Mexicana strictè erecta ispidula. Folia alterna, pleraque pinnatipartita, lobis linearibus dentatis, superiorum integris, summa linearia. Capitula corymboso-paniculata. Radius albus seu pallidè cœruleus. Discus flavus. — Differt ab

Erigeronte ut *Diplopappus* ab *Astere*. Valdè affine *Stenactidi* sed distinctum ob florum omnium pappos conformes.

1. *P. DELPHINIFOLIUM*. ② in Mexico (Alam.! Mairët!) circa urbem (Berl. pl. exs. n. 659), in Amer. merid.? (Humb. et Bonpl.). *Erigeron delphinifolium* Willd. hort. berol. n. et t. 90 enum. p. 873. *Diplopappus delphinifolius* Cass. dict. 13. p. 96. *Stenactis delphinifolia* Cass. dict. 50 p. 483. *Polyactis delphinifolia* Less. l. c. (v. s. sp. et c.)

CXXXVII. HETEROCHÆTA.

Capitulum multiflorum radiatum, ligulis pluriserialibus linearibus fœmineis, disco hermaphr. 5-dentato. Recept. nudum punctulatum. Invol. squamæ 2-3-seriales acuminatæ subæquales. Achænia pubera compressa? Pappus disci et radii duplex, ext. brevis paleaceus, int. setiformis scaber longus. — Herbæ Indicæ (et fortè Americane) pilosæ villosæ aut canescentes. Capitula solitaria terminalia. Ligulæ albæ aut purpurascentes. — Genus medium inter *Stenactidem* et *Erigeropsidem*, sed differt pappo disci et radii conformi. Dist. ab *Erigeronte* pappo duplici, à *Diplopappo* et *Diplostephio* ligulis pluriserialibus. Nomen ab *ἡτερος*, diversus, et *χαιρα*, seta.

§. 1. *Species gerontogæ.*

1. *H. DIPLOSTEPHIODES*, caule erecto folioso hispido tereti striato, foliis radic.... caulinis lineari-lanceolatis sessilibus acuminatis integerrimis utrinquè sparsè hispidis, invol. squamis subbiseriatis dorso hispidis lanceolatis acuminatis disco longioribus, achæniis plano-compressis obovatis hispidis, pappi serie externâ paleis lanceolatis constante. 2. in Indiæ bor. occid. montibus legit cl. Royle. Capitulum ligulis exceptis diam. pollicare. Ligulæ angustæ pollicem longæ purpureæ numerosissimæ. (v. s. comm. à cl. Royle sub n. 147.)
2. *H. HISPIDA*, caule erecto ramoso folioso setis patentibus rigidis patulis hispido, foliis lato-linearibus acutis integerrimis secus margines et nervum iisdem setis sparsè hispidis, invol. confertè et brevius hispido, achæniis plano-compressis glabriusculis, pappi paleis ext. squamiformibus lanceolatis. — Specim. habeo ex itin. Neæano, an ex Amer. calid.? an ex insulis Philippinis? Species distinctissima. Capitula ad apices ramorum denudatorum solitaria. (v. s.)
3. *H. ERIGEROIDES*, cæspitosum totum hirtello-canescens, caulibus erectis scapiformibus 1-cephalis basi tantum foliosis, foliis radicalibus petiolatis obovato-oblongis obtusis aut calloso-submucronatis integerrimis, caulinis paucis sessilibus oblongis, invol. hirsutissimi squamis biserialibus linearibus acuminatis disco longioribus, achænio villosa, pappi serie externâ multipaleacea brevi. 2. in montibus Indiæ boreali-occidentalis legit cl. Royle. Habitus ferè *Erigeri* Alpini aut nivalis. Ligulæ numerosæ angustissimæ. Collum suffruticosum multiceps. Caulis 3-4 pollic. (v. s. comm. à cl. Royle sub n. 114. 1.)
4. *H. ASTEROIDES*, caule erecto 1-cephalo folioso hirsutissimo, fol. radic...., caulinis ovalibus sessilibus apice callosis integerrimis utrinquè hirsutis, infer. basi attenuatis, invol. squamis 2-3-seriatis lanceolatis linearibusque puberis disco longioribus subcoloratis, ligulis subuniseriatis linearibus obtusiusculis, achænio villosa, pappo ext. brevissimo subcoroniformi. 2. in montibus Indiæ boreali-occidentalis. Habitus omninò *Asteris* Alpini. (v. s. comm. à cl. Royle sub n. 114. 2.)

§. 2. *Americane, congeneres dubie.*

5. *H. ? GNAPHALIOIDES*, caule erecto infernè simplici apice paniculato polycephalo albo-lanato, foliis sessilibus lanceolato-linearibus apice dentato-serratis subtus tomentoso-lanatis albis, radicalibus suprâ glabris, caulinis lanato-pubescentibus. 2. in Mexici convalle Guanajuatensi et ad montem Cabilate. *Eriger-*

geron gnaphalioides H. B. et Kunth nov. gen. am. 4. p. 88. t. 331. *Stenactis gnaphalioides* Cass. dict. 50. p. 484. Ligulæ pappo breviores albidæ circ. 100, quo caract. hæc et sequens differunt à *Stenactide*.

6. H. ? *PUBESCENS*, caule erecto ramoso polycephalo foliisque pubescentibus, radicalibus petiolatis oblongis lobato-dentatis, caulinis sessilibus integerrimis, capitulis longè pedunculatis. 2/ in temperatis juxta Moran Mexicanorum. *Erigeron pubescens* H. B. et Kunth nov. gen. am. 4. p. 88. *Stenactis pubescens* Cass. dict. 50. p. 484. Affine *Polyactidio* ex Kunth, *Stenactidi* ex Cass. Ligulæ circ. 80 albæ subtitis pallide violacæ.

CXXXVIII. THEROGERON.

Omnia *Erigerontis* sed pappi dispares, ligularum pluriserialium 1-serialis pilosus, disci duplex, ext. paleis brevibus lanceolatis acutis circ. 5, inter. setis 5-7 elongatis. — Suffrutices erecti ramosi glabri in Novâ-Hollandiâ interiore spontanei. Capitula ramos terminantia solitaria. Ligulæ albæ tenues. — Nomen ex *θερος*, æstas, et *γέρων*, senex, quasi *Erigeron æstivus*.

1. T. *DENTICULATUM*, foliis linearibus apice sublatioribus dentibus 3-5 notatis. 3 in Novâ-Hollandiâ interiore in Molle's planitiibus prope Lachlan flumen julio 1817. flor. legit cl. A. Cunningham. Caulis tennis teres. Ramuli apice nudi foliola indivisa acuta paucissima gerentes. Invol. squamæ lineares acutæ. Fl. ligulati innumeri, tubulosi pauci latiores sæpius steriles. Achænia radii compressa glabra. (v. s. comm. à cl. A. Cunningham.)
2. T. *INTEGRUM*, foliis lanceolato-linearibus utrinquè acuminatis integerrimis. 3 in planitiibus apricis Novæ Hollandiæ interioris rarissima ad Lachlan flumen julio 1817 flor. legit cl. A. Cunningham. Caulis teres. Folia ferè Lini sed angustiora. Rami apice nudi vix foliolosi. Invol. squamæ lineares acuminatæ. Ligulæ numerosissimæ. Flores tubulosos verisim. paucissimos in capitulo non inveni, sed ex analogiâ cum priore summâ huc refero. (v. s. comm. à cl. A. Cunningham.)

CXXXIX. ERIGERON Linn. gen. n. 951. (excl. spec.) Less. syn. 190.

Capitulum multiflorum radiatum, ligulis fœmineis pluriserialibus linearibus disco æqualibus aut longioribus, fl. disci tubulosos regularibus, nunc omnibus hermaphr., nunc exterioribus fœmineis centralibus hermaphroditis, nunc fortè abortu submasculis. Recept. nudum foveolato-punctatum (nec nisi in E. maximo) alveolato-fimbrilliferum. Achæmium compressum eostre. Pappus pilosus scaber 1-serialis (aut 2-serialis?, serie interiore brevissima?). — Herbæ rarius suffrutices. Folia alterna integra dentata aut lobata. Capitula subhemisphærica, disco flavo, ligulis albis cœruleis purpurascensibusve, rarissimè ochroleucis. — Genus apud auctores heterogeneas species non satis notas plurimas colligens.

Sect. I. *EUEBIGERON*. — *Erigeron* Cass. dict. 15, p. 181. 37. p. 462 et 482.

Ligulæ disco longiores aut eo æquales. Pappi setæ in anulum basi non concretæ.

§. 1. *Ligulis disco longioribus.*

* *Fruticosa seu perennia.*

1. E. *FRUTICOSUM*, glabrum, caule fruticoso ramoso, foliis oblongo-lanceolatis utrinquè acuminatis in petiolum attenuatis medio dentibus magnis acutis utrin-

quæ 2-3 incis, capitulis pedicellatis corymbosis, invol. squamis linearibus acutis imbricatis pedicellisque pubescentibus, ligulis disco duplò ferè longioribus. $\frac{1}{2}$ in insulâ Juan Fernandez legit cl. Larrain qui ad cl. Bertero misit. Affine *E. paucifloro* sed facillè foliis inciso-dentatis differt. Ligulæ albæ. (v. s. comm. à cl. Bertero.)

2. *E. PAUCIFLORUM* (Hook. et Arn. bot. Beech. f. 2. p. 87), caule fruticoso ramoso glabro, ramulis pubescentibus, foliis confertis glabris lineari-lanceolatis basi attenuatis atque ciliatis integerrimis aut hinc inde serratis, junioribus subpubescentibus, paniculæ terminalis corymbosæ ramis 2-4-cephalis, ligulis lineari-oblongis disco paulò longioribus, squamis invol. oblongis acutis discum subæquantibus. $\frac{1}{2}$ in insulis Sandwich speciatim in ins. Ouahu. Habitus Railiardiæ linearis. Folia pollicem longa sesquilineam lata. Ligulæ lineares albidæ obtusiusculæ pappo subrescente longiores. Stigmata oblonga linearia papillosa. (v. s. coll. à cl. Macrae ab hon. Soc. hort. Londin. comm.)

3. *E. LEPIDOTUM* (Less. in linnæâ 1831. p. 502), fruticosum, ramis striatis junioribus ad apicem foliatis, foliis linearibus acutis integris serrativè glabris adpressè ciliatis lepidotis, corymbis compositis pedicellis filiformibus hirsutopubescentibus, invol. campanulati squamis linearibus acuminatis. $\frac{1}{2}$ in Ouahu ins. Sandwicensium. Achænia hirsuta punctata. Ligulæ reflexæ albæ?

4. *E. SULCATUM*, caule erecto striato-sulcato pubero-scabrido apice ramoso, foliis sessilibus auriculis 2 obtusis semiamplexicaulibus ovalibus acuminatis grossè et distanter mucronato-dentatis ntrinque setulis sparsis aspero-scabris summis lineari-lanceolatis integerrimis, capitulis ad apices ramorum solitariis subcorymbosis, invol. squamis glabris lanceolatis inter. scariosis acuminatis, ligulis disco 3^o-4^{ve} longioribus. $\frac{1}{2}$? in Brasiliæ pascuis uliginosis legit cl. Lund. Ligulæ biseriales lineares albæ. Cor. disci flavæ. Anth. ecaudatæ. Pappus 1-serialis. Achænia compressa glabra. Folia 6-7 pollic. longa, 1 $\frac{1}{2}$ poll. lata. Habitus Inulæ aut Doronici. Car. Asteris aut Erigeri. Species distinctissima, socialis, putridè fœtens, 8-12 pedes alta. Radix ex Lund subglobosa fibrosa. (v. s. comm. à cl. inventore et ex prov. Sancti-Pauli in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 401 miss.)

β . caule glabriusculo, foliis involucriisque vix pilosis $\frac{1}{2}$ in Brasiliæ prov. Sancti-Pauli. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 413.)

5. *E. GLAUCUM* (Ker. bot. reg. t. 10), caulibus ascendenti-erectis pilosulis, foliis glaucis viscosis ciliato-villosis, infer. obovatis in petiolum alatum attenuatis paucidentatis, super. oblongis sessilibus, capitulo terminali subsolitario, ligulis circ. 100 linearibus disco triplò ferè longioribus. $\frac{1}{2}$ in Americâ ad Buenos-Ayres? (Ker.). Aster Bonariensis Spreng. syst. 3. p. 528. *Stenactis glauca* Nees ast. 275. Habitus *Asterum alpinorum*. Rhizoma caudiciforme breve persistens. Herba ingrâtè odora. Ligulæ lilacinæ 2-3-seriales. Capitulum 15-18 lin. diam. Pappus simplex mihi videtur. (v. v. c.)

6. *E. SPECIOSUM*, caule erecto ramoso pilis eglandulosis sparsis distanter hispido, foliis integerrimis ciliatis glabris, radicalibus spatulatis in petiolum attenuatis, caulibus sessilibus acuminatis, capitulis ramos apice nudos terminantibus, invol. hispido-ciliato, ligulis numerosissimis (circ. 120) disco triplò ferè longioribus. $\frac{1}{2}$ in Californiâ. *Stenactis speciosa* Lindl. bot. reg. t. 1577. Valdè affinis *E. glauco* et *E. labello*. Capit. diam. bipollic. Ligulæ atro-violaceæ.

7. *E. GLABELLUM* (Nutt. gen. am. 2. p. 148), caule erecto vix sparsè piloso, foliis integerrimis glabris eciliatis, radicalibus obovato-oblongis spatulatis, super. sessilibus lanceolatis, capitulis 3-5 pedicellatis corymbosis, invol. pubescente, ligulis numerosissimis (circ. 100) linearibus disco duplò longioribus. $\frac{1}{2}$ in planitiebus ad fl. Missouri prope Fort Mandan (Nutt.), in sylvis Amer. arcticæ (Richards. app. p. 30). Hook. bot. mag. t. 2923, sed caulis pilis capitatis hispidulus pingitur! quod in meis non occurrit. Ligulæ pallidè cæruleo-purpureæ. Affine *E. glauco* et *specioso*. Capit. diam. pollicari. (v. v. c.)

8. *E. MAXIMUM* (Otto hort. berol.), caule herbaceo erecto striato setulis callosis scabro, foliis amplexicaulibus dentato-serratis suprè exasperatis, radicalibus

elongato-lanceolatis, sup. cordato-lanceolatis acuminatis, capitulis ad apices pedunculorum solitariis, invol. squamis biserialibus acuminatis disco subaequalibus, ligulis biserialibus, recept. alveolato-fimbriifero. \mathcal{L} in Mexico. *Leptoselma maximum* Don in Sweet br. fl. gard. ser. 2. t. 38. *Aster maximus* Less. syn. 182. Rec. non est paleaceum sed fimbriiferum et ideò genus ab *Erigerone* non satis differt. Ligulæ ex albo purpurascens circ. semipoll. longæ.

9. *E. LOWGERÆ*, caule simplici imò basi folioso longissimè aphylo 1-cephalo basi hispidò apice pubescente, foliis subradicalibus sessilibus obovato-cuneatis grossè dentatis utrinquè sparsè hispidis, dentibus mucronatis, invol. squamis linearibus dorso villosis, ligulis numerosissimis disco duplò longioribus. \mathcal{L} in Mexico legit cl. Karvinski. Caulis 12-13 poll. longus per 10-11 nudus. Ligulæ albæ lineares. (v. s. in h. acad. reg. Monac.)

10. *E. PHILADELPHICUM* (Linn. sp. 1211 non Willd.), pubescens, caule adscendenti-erecto laxè corymboso, foliis infer. cuneato-obovatis sinuato-dentatis, caulinis oblongo-lanceolatis semiamplexicaulibus, capitulis longè pedicellatis, ligulis numerosissimis ultrà 100 tenuissimis disco duplò longioribus. \mathcal{L} in Amer. bor. agris et pascuis à Canadà ad Carolinam. Pursh fl. bor. am. 2. p. 533. Ell. sketch 2. p. 396, descr. optima. Er. amplexicaule Poir. suppl. 5. p. 464. Willdenowius pessimè phrasim Linnæanam correxit et pluribus dubiis ansam dedit. Herba circ. pedalis. Invol. post anthesim reflexum. Ligulæ pallidè purpureæ. (v. v. c. et s. sp.)

β . *quercinum*, foliis infer. ultra medium pinnatifidis sublyratis, lobis hinc indè dentatis. \mathcal{L} in Carolinà (Ell.) et Floridà (Pursh). *Erig. quercifolium* Pursh fl. bor. am. 2. p. 533 (excl. syn. Lam.). Ell. sketch 2. p. 397. Var. *E. Philadelphici* monente opt. Elliot. (v. s.)

11. *E. KARVINSKIANUM*, caule virgato tereti suffruticuloso puberulo, foliis caulinis inf. obovatis longè cuneatis medio triplinerviis utrinquè glabris basi ciliatis apice tri-rariùs quinquedentatis, dentibus grossis obtusis mucronatis, super. oblongis subintegerrimis, capitulis pedunculatis laxè corymbosis, invol. squamis glabriusculis lineari-subulatis, ligulis disco duplò longioribus biserialibus. \mathcal{L} in Mexico legit cl. Karvinski. (v. s. in h. acad. reg. Monac.)

12. *E. QUERCIFOLIUM* (Lam. ill. t. 681. f. 4), glabriusculum, caulibus erectis à basi ramosis, foliis inferioribus lyrato-dentatis basi in petiolum attenuatis, superioribus sessilibus oblongo-linearibus integerrimis, capitulis pedicellatis laxè corymbosis, invol. squamis linearibus acutis glabriusculis, ligulis tenuissimis circ. 100 disco duplò longioribus. \mathcal{L} in Floridà (Lam.), in territorio Arkansano (Nutt.) Poir. dict. 7. p. 491. Proximè accedit ad *E. Philadelphicum* sed glabritie ferè totà jam primà fronte differt. (v. s. comm. à cl. Nuttall.)

13. *E. MUCRONATUM*, caule erecto tereti glabriusculo aut subpubero ramoso, foliis lanceolatis basi attenuatis glabriusculis ciliatis integerrimis aut utrinquè lobulo ad medium auctis apice lobulise mucrone calloso longo superatis, capitulis paucis pedunculatis, invol. squamis linearibus subulatis puberulis, ligulis biserialibus disco duplò longioribus. \mathcal{L} in Mexico legit. cl. Karvinski. Ligulæ albæ. (v. s. in h. acad. reg. Monac.)

14. *E. BELLIDIFOLIUM* (Muhl. in Willd. sp. 3. p. 1958), hirsutum, caule florido erecto paucifolio è basi stolonifero, foliis radicalibus obovatis subserratis, caulinis distantibus sessilibus oblongo-lanceolatis, capitulis 3-5 pedicellatis corymbosis, ligulis linearibus (circ. 50) disco duplò longioribus. \mathcal{L} in siccis umbrosis Amer. bor. à Canadà ad Carolinam. Pursh fl. bor. am. 2. p. 532. Ell. sketch 2. p. 393. Sims bot. mag. t. 2402. Er. pulchellum Michx. fl. bor. am. 2. p. 124. Variat incanum aut minus hirsutum. Ligulæ numerosissimæ pallidè cœrulæ. Pedicelli sub capitulis obconici. (v. s. sp.)

β . *scariosum*, hirsuto-incanum, invol. squamis ferè omninò albo-scariosis vix basi ad carinam pilosiusculis. \mathcal{L} in Kentucky legit cl. Rafinesque. Ex bellidifolium Raf. ! in litt. (v. s. sp.)

15. *E. PURPUREUM* (Ait. kew. ed. 1. v. 3. p. 186), caule erecto sparsè piloso,

- foliis caulinis oblongis dentatis amplexicaulibus, summis integerrimis, pedunculis corymbosis sub capitulo paulò incrassatis, invol. squamis carinâ pilosis margine membranaceis, ligulis tenuissimis numerosissimis (ultra 100) disco duplò longioribus. $\frac{1}{2}$ in pascuis et fertilibus collibus ad ripas fluminum Amer. bor., in Canadâ ad sinum Hudsonis (Ait.), à Canadâ ad Virginiam (Pursh 2. p. 583), in Kentucky (Raf.!), Noveboraco (Lud. Bonap.!) ad ripas fl. Mississipi (Berl.!), in Californiâ (Douglas!). Erig. amplexicaule Torr. in litt. Ligulæ purpureæ (Pursh) in sicco pallidæ imò albæ. (v. s. sp.)
16. *E. ASPERUM* (Nutt. gen. am. 2. p. 147), hirsuto-scabrum, caule gracili circ. 2-cephalo, foliis lanceolatis acutis integris, radicalibus subspathulatis, pedunculis 2-3-subterminalibus, capitulis hemisphæricis. — in Amer. bor. planitiibus ad fl. Missouri. Ligulæ albæ. Pappus duplex, int. simpliciter pilosus circ. 20-setus achænio lævi multò longior leviter rufescens. An ex pappo duplici *Stenactis* aut *Heterochaeta*?
17. *E. INTEGRIFOLIUM* (Bigel. fl. houst. ed. 2. p. 302), caule simplici folioso glabro, foliis lanceolatis integris 3-nerviis pubescentibus, capitulis corymbosis, inv. hemisphærici squamis acutis. $\frac{1}{2}$ circa Bostoniam in sylvis. Ligulæ albæ. Inter *E. bellidifolium* et *E. strigosum* collocat cl. auctor.
18. *E. PUMILUM* (Nutt. gen. am. 2. p. 147), hirsutum, caulibus è collo plurimis adgregatis 1-cephalis, foliis oblongo-linearibus integris sessilibus, capitulis hemisphæricis ante anthesin nutantibus, involucrio hirsutissimo. — in Amer. bor. planitiibus ad flum. Missouri. *Erigeron hirsutum* Pursh fl. suppl. 742, non Lour. Ligulæ albæ invol. duplò longiores. Capit. magn. *Bellidis* perennis. Pappus duplex, inter. brevis 12-setus.
19. *E. HUMILE* (Grah. in Jam. new. phil. journ. Ed. 1828. decemb. p. 175), caule brevissimo 1-cephalo, foliis integerrimis ciliatis sursum pubescentibus patulis, radicalibus spathulatis basi longè attenuatis, caulinis lineari-lanceolatis acuminatis, invol. campanulati calyculati squamis 5 laxis subimbricatis pilosis, ligulis invol. æqualibus disco longioribus, tubulis disci 4-dentatis. $\frac{1}{2}$ in Americæ arcticæ littoribus (Richards.). Ligulæ albæ.
20. *E. BERTERIANUM*, caule gracili erecto foliisque setis minutissimis adpressis puberulo apice laxè paniculato, ramis subfoliosis apice 1-3-cephalis, foliis distantibus linearibus integerrimis, invol. squamis glabriusculis linearibus acuminatis, inter. margine membranaceis disco æqualibus, ligulis disco duplò ferè longioribus. — in fruticetis sabulosis collium prope Valparaiso Chiliensium jan. fl. legit cl. Bertero. Habitu accedit hinc ad *Asterem Gayanum* illinc ad *Er. spinulosum*, à priori ligulis 2-3-seriatis, à poster. ligulis disco duplò longioribus faciliè distinctum. (v. s. comm. à cl. Bertero.)
21. *E. WIGHTII*, caule erecto hispido breviter ramoso, foliis oblongis, infer. basi attenuatis subserratis obtusiusculis, sup. integerrimis acutis, omnibus utrinquè puberulo-hirtis, capitulis pedicellatis subracemosis, invol. squamis dorso scabris lineari-subulatis disco æqualibus, ligulis tenuissimis disco longioribus, achænio glabriusculo. $\frac{1}{2}$? in peninsulâ Indiæ orient. legit cl. Wight (herb. $\frac{1}{2}$). Ligulæ pallidè purpurascens multiseriales. Rami valdè hispidi. Herba cinerascens (v. s. in h. Wight.)
22. *E. ANGUSTISSIMUM* (Lindl. ! adn. mss.), caule basi suffruticoso ascendente ramosissimo, ramis apice aphyllis 1-cephalis, foliis linearibus integerrimis glabris, invol. squamis linearibus acuminatis disco brevioribus, inter. membranaceo-marginatis, ligulis angustè linearibus biserialibus, achænio villosulo. $\frac{3}{4}$ $\frac{1}{2}$ in Indiæ or. montibus Peerpunjal legit cl. Royle. Palmaris. Ligulæ cyanæ. (v. s. comm. à cl. Royle sub n. 115.)
23. *E. MONTICOLA* (Wall. ! cat. et herb. n. 2977. comp. n. 87), caule imâ basi frutescente erecto simplici cum foliis sparsè villosis, foliis radicalibus lineari-lanceolatis basi attenuatis, caulinis sessilibus linearibus, omnibus subintegerrimis acutiusculis, capitulis 2-5 pedicellatis corymbosis, invol. squamis linearibus acutis disco brevioribus, ligulis linearibus multiserialibus, fl. tubulosis

paucis in centro. $\frac{1}{2}$ in montibus Nepaliæ ad Kamaon et Deyra D'hoon. Er. humilis Spreng. in litt. ad Wall. Ast. monticola Lindl. adn. mss. Herba 3-6 poll. longa. Ligulæ albæ. Achænia oblonga compressa puberula. Pappus rufus. (v. s. comm. ab ill. Cur. merc. angl. Ind. or. et à cl. Royle.)

24. E. POEPPIGII, totum hispido-scabrum, caule erecto subangulato racemoso, foliis linearilanceolatis elongatis acutis integerrimis, infer. basi in petiolum attenuatis, super. sessilibus, ramis erectis paucifoliis apice nudis 1-cephalis, invol. squamis linearibus acutis dorso hispido-scabris disco brevioribus, ligulis disco duplò ferè longioribus, achæniis elongatis glaberrimis. $\frac{1}{2}$ in Chili? legit cl. Poeppig (pl. exs. n. 21). Herba ferè pedalis. Capitula circ. 5 distantia. Ligulæ ex sicco albidæ. Valdè accedit præsertim ex achæniis formâ ad E. Andicolam. (v. s. ex coll. Poeppig.)

25. E. ANDICOLA, cæspitosum, caulibus plurimis erectis molliter hispidis 1-cephalis apice subaphyllis, foliis linearibus acutis longè ciliatis et hispidis integerrimis, radicalibus erectis basi attenuatis, caulinis sessilibus paucis, invol. squamis lanceolato-linearibus acuminatis hispidis disco æqualibus, ligulis linearibus disco duplò longioribus, achæniis elongatis glaberrimis. $\frac{1}{2}$ in excel-sis Andibus Chilensium legit cl. Gay. Invol. purpureascens setis albis hispidum. Ligulæ ex sicco albæ. Folia radic. 2 poll. longa, 2 lin. lata. Caulis 3-4 pollic. (v. s. comm. à cl. Gay.)

β . *pubescens*, setis omnibus brevissimis, capitulo minore minùs hispido. — in Andibus Chil. etiam legit cl. Gay. An spec. propria? An E. Poeppigii var. 1-cephala? (v. s.)

26. E. GLABRIFOLIUM, glaberrimum subcaule, rhizomate suffruticulis, foliis radicalibus linearilongis basi subattenuatis, caulinis paucis semialexicaulibus, omnibus integerrimis acutis, caule adscendente 1-cephalo, invol. squamis disco æqualibus oblongis obtusis adpressis, ligulis disco duplò longioribus. $\frac{1}{2}$ in summis Andibus Chilensium legit cl. Gay. Etiam adest sub n. 638. in plant. Poeppigianis. Habitus Leontodontium alpinorum. Accedit ad E. Andicolam sed distinctissimum. Caulis 2-3 pollic. Folia radic. pollicem longa, 3 lin. lata. Ligulæ albidæ ex sicco. Pappus albus. Achænia puberula-hirtella. (v. s.)

27. E. SCAPOSUM, caule basi decumbente surculoso-ramoso folioso, apice longè denudato 1-cephalo puberulo, foliis pubescenti-hirtellis, infer. subcongestis oblongo-lanceolatis dentatis, summis distantibus paucis semialexicaulibus lanceolatis integerrimis, invol. squamis linearibus acuminatis dorso hirtellis disco æqualibus, ligulis disco duplò longioribus. $\frac{1}{2}$ in Mexico circa Tolucaam aprili flor. legit cl. Andrieux (pl. exs. n. 277). Folia infer. sesquipollic. longa, 3-4 lin. lata. Caulis floridus spithamæus. Ligulæ albidæ. Discus in sicco fuscus. (v. s.)

β . *latifolium*, foliis lato-oblongis minùs hirtellis, capitulo majore, invol. squamis glabresculis sen minùs hirtellis. $\frac{1}{2}$ circa Mexico legit cl. Berlandier pl. exs. n. 375. (v. s.)

28. E. PULCHELLUM (DC. non Michx.), caulibus basi subcæspitosis simplicibus 1-cephalis sparse pilosis apice confertè et longè hirsutis, foliis glabresculis ciliatis, infer. obovatis spatulatisve, caulinis sessilibus oblongis, invol. hirsutissimo subcolorato, ligulis tenuissimis disco duplò longioribus. $\frac{1}{2}$ in Alpibus Iberico-Caucasicis (Stev.!), Sibiriz et Dahuriæ speciatim ad montem Schebet (Turczaninoff!). Aster pulchellus Willd. sp. 3. p. 209. Aster Alwartensis Lodd. cat. 1807. ex Sims bot. mag. t. 2321. Erig. alpinus β ericalyx Ledeb. fl. alt. 2. p. 91. Ligulæ purpureæ. Fl. fœm. tubulosi nulli, undè ab E. alpino faciliè differt. (v. s.)

β . *Unalaschkense*, minor, foliis radicalibus petiolatis obovatis obtusis, caulinis linearibus acuminatissimis hispidis, caule ad apicem involucrique setis longis rufidis patulo-tortuosis hirsutis, ligulis albidis. $\frac{1}{2}$ in ins. Unalaschka et ad sinum Sancti-Laurentii legit cl. de Chamisso. Erig. Alpina Less. in linnæa 1831. p. 126. Fl. fœm. tubulosi ut in var. α nulli. (v. s.)

29. *E. COMPOSITUM* (Pursh fl. bor. am. 2. p. 535), subcaule, foliis radicalibus longè petiolatis triplicato-tripartitis, lobis linearibus divaricatis petiolisque puberis, caule folia 1-2 linearia distantia gerente vix puberulo 1-cephalo. 2/ in Amer. bor. ad ripas fl. Kooakoosky (Lewis ex Pursh), in reg. arcticis ad Point-Lake (Rich. ex Hook. in trans. Linn. soc. 14. p. 374. t. 13). Cineraria Lewisii Richards. in Frankl. journ. app. p. 148 ex Hook. Capitulum Bellidis. Ligulæ ex albo demùm rubellæ invol. duplò longiores.
30. *E. PAPPOCHROMA* (Labill. nov. holl. 2. p. 47. t. 193), glabrum acule, foliis radicalibus obovatis subserratis basi attenuatis ciliatis, caulinis integris linearibus distantibus, caulibus scapiformibus simplicissimis 1-cephalis. 2/ in insulâ Van-Diemen. Erigeron phlogotrichus Spreng. syst. 3, p. 520. Herba 3 pollic. Ligulæ dilutè violacæ disco numerosiores circ. 2-seriales. Achaemia glabra compressa. Pappus rufus. An certè hujus generis? (v. s. comm. à cl. Labill.)

*** *Herbaceæ annuæ.*

31. *E. RIVULARÆ* (Swartz prod. 113. fl. ind. occ. 1346), caule erecto parè ramoso pilosiusculo folioso oligocephalo, foliis radicalibus canlinisve oblongo-cuneatis membranaceis ciliatis puberulis, aliis integerrimis, aliis dentatis, aliis pinnatifidis, dentibus lobisve mucronatis, capitulis pedicellatis subcorymbosis, invol. squamis linearibus disci longitudine, ligulis linearibus disco paulò longioribus. ① in petrosis secus rivulos Jamaicæ (Sw.). Erig. runcinatum Bert. herb. Folia ferè Capellæ Bursæ-Pastoris. Species *E. Jamaicensi* affinis. Ligulæ albæ parvæ parvæ. (v. s. ex h. Bert. sine loci design.)
32. *E. JAMAICENSIS* (Linn. amœn. 5. p. 406), caule erectiusculo pubescenti-subvillosa paucifolio et oligocephalo, foliis radic. rosulatis puberulis membranaceis cuneiformi-lanceolatis apice paucidentatis, caulinis distantibus squamiformibus, invol. squamis linearibus glabrisculis disci longitudine, ligulis linearibus disco paulò longioribus. ① in agris Jamaicæ (Sloan. jam. 1. p. 260. t. 152. f. 3. Swartz obs. 305. t. 8. f. 2) et Sancti-Domingi (Bert. l.). Herba semipedalis. Folia radical. 2 poll. longa 6-7 lin. lata. Capit. parva, ligulis albis. (v. s. ex Sancto-Domingo.)
33. *E. CUNEIFOLIUM*, rhizomate obliquo, caulibus erectis oligocephalis rigidulis nudis glabris vix basi puberulis, foliis radicalibus obovato-cuneatis obtusis subintegerrimis glabrisculis, caulinis distantibus squamiformibus, invol. squamis linearibus disci longitudine, exter. patulis, ligulis linearibus disco paulò longioribus. ① in ins. Porto-Ricco legit cl. Wydler (pl. exs. n. 303). Affine *E. Jamaicensi*. Folia pollicem longa, 5 lin. lata. Caulis 7-8 pollic. Capitula longè pedicellata parva. Ligulæ in sicco purpurascentes. (v. s.)
34. *E. BELLIOIDES*, caulibus debilibus adscendentibus gracillimis 1-2-cephalis glabris, foliis radic. rosulatis orbiculatis ovatisve subdentatis glabrisculis membranaceis in petiolum alatum longè cuneato-attenuatis, capitulis longè pedicellatis minimis, invol. squamis linearibus disci longitudine. ① in Porto-Ricco decemb. flor. legit cl. Bertero. Erig. rivulare Spreng. in h. Bert. Herba pusilla Bellii seu Bellidis annuæ facie. *E. Jamaicensi* affinis sed distinctissima. Folia 9 lin. longa 3 lin. apice lata. Caules 2-3 pollic. (v. s. ex h. Bert.)
35. *E. TENELLUM*, pubescens, caule à basi ramoso gracili, ramis apice nudis 1-cephalis, foliis oblongis integerrimis, infer. obtusis basi attenuatis, super. sessilibus acutis, invol. squamis linearibus puberulis disco subæqualibus, ligulis disco duplò longioribus. ① in Mexico circa Tamaulipas in campis Matamoros legit cl. Berlandier martio flor. Herba 2-4 pollic. Bellidis annuæ ferè facie sed pubescens hinc inde villosa. Ligulæ albido-cæruleæ (v. s. comm. à cl. inv.)
36. *E. LEPTORHIZON*, caule herbaceo patentim hispidis apice ramoso, foliis membranaceis infernè ciliatis utrinquè puberulis, infer. basi in petiolum attenuatis obovatis seu subspathulatis apice parè dentatis, super. sessilibus oblongo-lanceolatis basi cuneatis apice grossè serratis, capitulis pedicellatis paucis subcorymbosis, invol. squamis linearibus puberulis disco æqualibus,

ligulis disco duplò longioribus. ① in Amer. austr. (Demb.!) circa Limam (Abadia!) Herba semipedalis. Folia 2 et ultrà pollices longa, 6-6 lin. lata. Ligulæ albæ. Pappus paucisetus. An fortè Stenactidis species, sed in capit. junior. seriem externam non ritè vidi. (v. s.)

37. *E. AFFINE*, caule erecto basi patentim hirsuto supernè pubero simplici apice in ramos 1-2 nudos elongatos 1-cephalos diviso, foliis infer. elliptico-oblongis basi longè cuneatis integris apice dentibus grossis mucronatis utrinquè 4 incisus utrinquè pubero-hirtis, summis semiamplexicanibus serratis acuminate, invol. pubescenti-hirto, ligulis circ. 100 linearibus disco duplò longioribus. ① circa urbem Mexico junio flor. legit cl. Berlandier (pl. exs. n. 522). Er. Philadelphico affine sed caules 1-2-cephali, capitula duplò majora circ. Bellidis perennis. (v. s. comm. à cl. inv.)

§. 2. *Ligulis discum non superantibus.*

38. *E. STRICTUM*, caule strictè erecto simplici sparè setifero tereti, foliis confertis erectis linearibus glabris secus nervum subtus hinc indè setosis margine longè et rigidè ciliatis aut demùm subserrulatis, paniculæ spiciformis confertæ terminalis ramulis brevibus foliolosis, invol. squamis glabris linearibus, ligulis brevissimis. — in Mexico circa Bejar Augusto flor. legit cl. Berlandier (pl. exs. n. 1760). Habitus ferè Linariæ. Caract. Erig. Canadensis sed panicula contracta nec expansa, folia stricta magis ciliata, etc. (v. s. comm. à cl. Berlandier.)

39. *E. SPINULOSUM* (Hook. et Arn. bot. Beech. 1. p. 32), caule herbaceo erecto apice laxè corymboso-paniculato hispido, setis rigidis internè sparsis patulis in pedicellis confertis adpressis, foliis lineari-oblongis acutis integris margine cartilagineo setis rigidulis præsertim scabro, invol. squamis circ. discum æquantibus, ligulis minimis disco æqualibus. — in Chili (Bridg.), in pascuis sterilibus collium circa Valparaiso Chil. (Bert.!). *E. fasciculatus* Colla pl. chil. f. 2. p. 26. n. 67. Affine videtur *E. Canadensis* et Bonariensis sed setis rigidis ferè spinellosis facillè differt. (v. s. comm. à cl. Bert.)

40. *E. CANADENSE* (Linn. sp. 1211), caule erecto hispido paniculatim ramosissimo, foliis lineari-lanceolatis ciliatis, invol. cylindricis, ligulis confertis involucro vix longioribus, achænio oblongo breviter pilosulo. ① frequens in campis et agris Americæ borealis; etiam crescit in Mexico (Alam.!) ad Xalapam (Less.) in Brasiliâ (Ch.), Cubâ (La Sagra!), Guadalupâ (Bert.!), ina. Trinitatis (Sieb.), Maderâ (Low.!), Indiâ orient. (Lamb.!), Persiâ boreali (Stev.!), Africâ austr.-orient. (Dreg.!) et Europâ totâ! Fl. dan. t. 292. Sencio ciliatus Walt. fl. car. 208? Erig. paniculatum Lam. fl. fr. Herba peregrina polymorpha. Variat erecta aut diffusa, caule plus minùs hispido interdum glabro, paniculâ racemiformi aut laxâ, etc. An species plures hic confusæ? (v. v. et s.)

41. *E. PUSILLUM* (Nutt.! gen. am. 2. p. 148), gracile glabrum, foliis lineari-lanceolatis, omnibus integris margine scabri, paniculæ subsimplicis ramulis divaricatis, ligulis paucis, achænio oblongo vix puberulo. ① frequens in Novâ Cesariâ et Pensylvaniâ (Nutt.!) in Carolinâ? (Ell. sketch 2. p. 398). *E. Canadense* Æ pusillum Bart. fl. phil. 2 p. 108. An Erig. Canadensis var. macilentâ? Herba 4-5 poll. longa. Flores disci 4-dentati ex Nutt. (v. s.)

42. *E. DIVARICATUM* (Michx. fl. bor. am. 2. p. 123), decumbens ramosissimum diffusum, ramulis fastigiatis, foliis subulatis, ligulis brevissimis, corollis disci 4-fidis. ① in pratis Illinoensium (Michx.), in Kentucky et Louisianâ (Nutt. gen. 2. p. 140). Affine *E. Canadensi* et *E. pusillo* dicitur. An *E. ramosum* Raf. fl. lud. p. 66. satis differt?

43. *E. BONARIENSE* (Linn. sp. 1211), caule hirsuto erecto, foliis pubescenti-hirsutis elongato-lanceolatis utrinquè attenuatis, infer. grossè inciso-dentatis, super. integerrimis, paniculâ elongatâ racemiformi, ligulis tenuissimis discum non superantibus. ① in Amer. austr. ad Bonariam (Dill. Bacle!), in Brasiliâ ad Rio-Janeiro in cultis et ad vias (Lund!). Erig. undulatum Moench meth.

598. — Dill. elth. 2. f. 334. Herba subpurpurascens seu obscure virens. Folia sæpè ad basin margine reflexa aut plana. Pappus simplex. Aff. E. Canadensi. (v. v. c. et s. sp.)
44. E. MULTIFLORUM (Hook. et Arn. bot. Beech. f. 2. p. 87), glabrum, caule herbaceo, foliis lineari-lanceolatis basi in petiolum longum attenuatis integerrimis, paniculâ ramosissimâ compactâ, ramis nudis, ramulis squamatis, squamis involucri oblongo-lanceolatis radii subæquantibus. — in insulis Sandwich. An fortè E. Canadense in ins. Owahu à cl. Less. citatum in Linnaë 1831. p. 126 ?
45. E. HIRTELLO, caule suffruticoso ramoso tereti setis sparsis patulis strigillosis hirtello, foliis lineari-oblongis integerrimis ad margines et subtus in nervo adpressè hirtello-ciliatis, capitulis paucis pedicellatis, invol. squamis linearibus glabriusculis, acheniis pubescentibus ligulis vix disco subæqualibus. § in Chili circa Valparaíso legit cl. Gaudichaud.
46. E. CHINENSIS (Jacq. h. Schœnb. 3. p. 30. t. 303), caule erecto hispido apice paniculato, foliis villosis lyratis in petiolum attenuatis, lobis inf. parvis paucis, terminali maximo ovali dentato acuto, summis lanceolatis, invol. hemisphærico demum reflexo. — in Chinâ. Ligulæ albæ breviores disco vix longiores.
47. E. ? TRIDACTYLUM, caule fruticoso brevi, ramis ferè à collo ortis erectis teretibus apice nudis 1-cephalis foliis involucriisque pubescentibus, foliis in parte ramorum infer. confertis in petiolum attenuatis apice trifidis, lobis et parte indivisâ linearibus, summis remotis linearibus acutis, invol. squamis linearibus 1-2-seriatis disco brevioribus, ligulis minimis, acheniis sericeo-villosis, pappo scabrido corollis longiore. § in Brasiliæ provinciâ Rio-Grande (h. imp. Bras. n. 1054 !). Fortè genus proprium ? Pappus rufus multisetus. Cor. ex sicco purpurascens; ligulæ paucae. (v. s. in h. Mus. reg. Par.)

Sect. II. TRIMORPHÆA. — Trimorphæa Cass. dict. 37. p. 462 et 482.
55. p. 348.

Flor. fœm. innumeri, exteriores 1-3-seriales angustissimè ligulati, interiores pluri-seriales corollâ tubulosâ tenuissimâ truncatâ donati. Flor. centrales tubulosi hermaphrod. 5-fidi. — Erig. Alpini var. aliæ plurimis aliæ paucissimis floribus fœmineis tubulosis donatæ demonstrant strictam hujus sectionis cum priore affinitatem.

48. E. KAMTSCHATICUM, caule erecto glabriusculo paniculato, ramis erectis apice racemosis, foliis infimis dentatis, inferioribus oblongis obtusis basi longè in petiolum alatum et ciliatum attenuatis, superioribus sessilibus lanceolatis, invol. squamis acuminatis puberulis pappum æquantibus, ligulis disco æqualibus. 2. in Kamtschatkâ. E. Villarsii Willd. sp. 3. p. 1958. excl. syn. et patr. Herba ferè bipedalis. Caulis purpurascens. Folia vix triplivenia. Capitula triplò minora quàm E. Villarsii veri. (v. s. comm. ab ill. Acad. sc. Petrop.)
49. E. ACER (Linn. sp. 1211), piloso-hirsutiusculum, caule erecto folioso superne racemoso, pedunculis 1-cephalis, foliis oblongis integerrimis inferioribus basi angustatis, super. sessilibus, pappo achenio duplò longiore. ② seu 2. in siccis apricis montosis Europæ totius, Caucasi et imp. Rutheni usque ad Dahuriam. DC. fl. fr. n. 3131. Er. vulgare Linn. fl. lapp. 308. Trimorphæa vulgaris Cass. dict. 55. p. 324. Utraque varietas variat plus minus hirsuta, ligulis purpurascens vel cœruleis albidisve, pappo rufo aut albedo.
- a. brachyglossum, radio discum non superante. Er. acre Sow. engl. bot. t. 1158. Schkuhr handb. t. 241. Bess. ! enum. p. 33. Ledeb. fl. alt. 3. p. 89. E. murale Lapeyr. suppl. 133. (ex specim. necum ut Lapeyr. à cl. Dufour comm.). Er. Philadelphicum Willd. sp. 3. p. 1958 ? (v. v. et s. olim in hortis sub nom. E. Philadelphici cult.)
- β. asteroides, radio discum superante. Er. asteroides Andr. ! ex Bess. Link. enam. h. herol. 2. p. 324 ? Er. Podolicus Bess. ! enum. p. 33. Ledeb. fl.

- alt. 4. p. 190. var. α . Er. Dræbachense Retz prod. n. 1010 ex fl. dan. t. 874. — Col. cephr. 2. p. 25. t. 26. f. 2. (v. v.)
50. *E. ELONGATUM* (Ledeb. ic. alt. t. 31. fl. alt. 4. p. 91), glabrum, caule subsimplici aut paniculato, foliis integerrimis, infer. spatulatis obtusis, super. lanceolatis oblongiæ acutis, pedunculis elongatis axillaribus 1-cephalis, invol. glabro, ligulis disco paulo longioribus. (2) seu \mathcal{L} . in siccis lapideis Sibiriae Altaicæ (Ledeb.) et Baicalensis (Turcz.). Affine *E. acris* et ab eo vix diversum mon. cl. Fischer in litt. sed glabritie distinctum. *E. elongatum* Moench suppl. 247. est verisim. diversum sed non satis notum. (v. s. sp. et c.)
51. *E. ARMERIFOLIUM* (Turcz. in litt. 1830), caespitosum basi multicaule humile, caulibus ramisve pubescentibus 1-cephalis, foliis omnibus linearibus integerrimis glabris basi setoso-ciliatis, invol. glabriusculi squamis linearibus apice coloratis acuminatis, ligulis disco paulo longioribus. \mathcal{L} . in uliginosis subaeualis prope Ircutiam in Dahuriâ legit cl. Turczaninof et in deserto Kuraico cl. de Bunge. *Erigeron podolicus* β pusillus Ledeb. fl. alt. 4. p. 90. (ex sp. à cl. Meyer misso). *Erig. graminens* Stev. et Falk. non Linn. ex Ledeb. Variat caulibus foliosis brevibus et apice elongatis subnudis. Ligulae pallidae tenuissimæ. Cor. fl. foem. tenuissimè truncatæ vix à stylo distinguendæ. (v. s. comm. à cl. inv.)
52. *E. ALPINUM* (Lam. fl. fr. 2. p. 140. et DC. ed. 3. n. 3132), caulibus basi subcaespitosis pilosis mono- seu oligocephalis, foliis lanceolatis acutiusculis villosis ciliatisve integerrimis, invol. squamis apice subulatis plus minus hirtis, ligulis disco duplò longioribus, pappo paulo achænio longiore. \mathcal{L} . in pascuis montanis imò alpinis Europæ et Sibiriae, imò fortè Americæ arcticæ. *Stenactis alpina* Cass. dict. 50. p. 485. Species polymorpha nec tamen meo sensu divellenda.
- α . caule 1-cephalo, invol. glabriusculo, foliis villosis, cor. tubulosis foemineis numerosis. — frequens in pascuis elatis apricis. *Erig. Alpinum* Sow. engl. bot. t. 464. Hæc et sequens var. sub *Er. alpino* Linn. sp. 1211. Sturm fl. germ., etc. militant. (v. v. et s.)
- β . caule sæpius oligocephalo, invol. glabriusculo, foliis glabris ciliatis, cor. tubulosis foemineis paucis. — in pascuis pinguioribus. *Erig. glabratum* Hoppe! pl. exs. et *E. rupestre* ejusd. vix differre videtur. (v. v. et s.)
- γ . caule 1-cephalo, invol. hirsutissimo, cor. tubulosis foemineis paucis. — in pascuis editis siccioribus. *Erig. uniflorum* Linn. sp. 1212. fl. lapp. t. 9. f. 3. (v. v. et s.)
- δ . caule oligocephalo, invol. hirsutissimo, cor. tubulosis foemineis paucis. — in pratis et agris montanis. *Erig. hirsutum* et *Erig. grandiflorum* Hoppe! pl. exs. (v. v. et s.)
53. *E. VILLARII* (Bell. app. ad fl. ped. p. 38. t. 7), caule erecto pubescente subviscoso apice racemoso, foliis lanceolatis triplinerviis integerrimis scabridis sessilibus, ramulis 1-cephalis folio longioribus, invol. scabrido, ligulis disco duplò longioribus, pappo achænio vix longiore. \mathcal{L} . in collibus montanis Delphinatûs (Will.), Galloprov. super. (Honorat!), Pedemontii (Bell.), Sabaudia (J. DC.), Vallesia (Sehl.). DC. fl. fr. n. 3137. *Erig. atticum* Vill. dauph. 3. p. 237. Herba subviscosa pedalis polycephala rigida. Species Willd. hortensis videtur diversa! Villars junior nomen suum Villar scribebat undè Bellardi dixit *E. Villarii*. (v. v. et s.)
54. *E. SPATHULATUM* (Vest St. Cr. p. 303), caule erecto simplici hispido folioso apice corymboso, foliis infer. in petiolum subulatum attenuatis ovalibus (subspatulatisve) grossè dentatis, sup. gradatim subsessilibus serratis acutis, omnibus ciliatis villosis subhirsutis, invol. squamis linearibus acuminatis disci longitud., ligulis minutissimis. — in ins. Sancti Tomasi locis siccis aprili flor. legit cl. Bertero et fortè etiam in Porto-Ricco? St. Crucis (Vest). *Erig. incisum* var. caribæa Spreng. syst. 3. p. 516 (excl. syn. ad spec. 3. refer.). Accedit ad *E. Chinense* Jacq. sed folia infernè non lyrata. Ligulae albæ vix disco longiores. Cor. foem. tubulosæ tennes, hermaphrod. latiores. Invol. demum reflexum. Ambigit inter *Nidorellam* et *Trimorphæam*. Nomen specif. minus aptum tamen retinui. (v. s. ex h. Bert.)

55. *E. SEMIBARBATUM*, caule erecto subsimplici hispido, foliis oblongis callosomucronatis, infer. basi attenuatis glabriusculis, cæteris sessilibus semiamplexicaulibus ciliatis et pubescenti-hirtis dentatis integerrimis, capitulis 1-5 ad apicem caulis ramorumve solitariis subcorymbosis, invol. pedicellisq. hispidissimis, corollis fœmineis ligulatis tubulosive longe barbatis, hermaphr. glabriusculis. — in Indiâ orient. legit cl. Royle. Herba vix pedalis. Folia 2 poll. longa, 4-5 lin. lata. Capit. 9-10 lin. diam. Ligulæ cœrneæ disco duplò longiores. Cor. tubulosæ fœmineæ brevissimæ, herm. 5-dentatæ glabræ. Achænia præm. radii villosissima. Pappus basi setis squamulisve cinctus; an ex achænio productus, an duplex? (v. s. comm. à cl. Royle sub n. 138.)
56. *E. ROYLEI*, caule erecto hispidulo, foliis ovali-oblongis subserratis callosomucronatis ciliatis scabridis triplinerviis, infer. basi attenuatis, super. sessilibus, capitulis multifloris pedicellatis racemosis, invol. pedicelloque hispidissimo, ligulis tubulosisq. fœmineis parçè barbatis, corollis hermaphr. hirsutis, achæniis puberulis. — in Indiâ orientali legit cl. Royle. Folia 2 poll. longa, 6-7 lin. lata. Ramuli erecti vix folio suo longiores. Ligulæ ex sicco albidæ. Pappus rufus cor. tubulosis æqualis. (v. s. comm. à cl. Royle sub n. 140.)
57. *E. LESCHENAUTII*, totum piloso-hispidum, caulibus herbaceis erectis teretibus striatis, foliis sessilibus oblongo-linearibus lingulatisve obtusiusculis mucronatis integerrimis, pedunculis axillaribus folio paulò longioribus 1-cephalis, summis brevioribus oligocephalis subcorymbosis, invol. squamis linearibus disco brevioribus, achæniis villosis. — in Indiâ australi montibus Neelgherry dictis legit cl. Leschenault. Villi fructus superiores calyculorum circa pappum interdum simulant. (v. s. comm. à cl. Leschenault.)
58. *E. MULTICAULE* (Wall.! cat. et herb. n. 3286. excl. litt. B.), glabriusculum, caulibus herbaceis à collo multiplicibus erectis teretibus, foliis lanceolatis basi attenuatis sessilibus acutis integerrimis, paniculâ terminali laxâ, ramis floridis elongatis capitula solitaria paucave gerentibus, invol. squamis linearibus acuminatis, ligulis linearibus 1-seriat. invol. paulò longioribus, tubulis fœm. pluriseriatis, masc. circ. 20. (1) ? in Indiâ orient. prov. Kamaon legit cl. Wallich. Herba 2-pedalis. Caulis subpurpurascens. Folia 12-15 lin. longa, 3 lin. lata. Ligulæ albæ. Pappus rufescens. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
- ℓ. Madurensis* (DC. in Wight contrib. bot. ind. p. 9), caule foliisque pube brevi sparsâ scabridis, foliis super. integerrimis, capitulis omnibus solitariis. — in provinciâ Madurensi legit cl. Wight. *Erigeron multicaule* C. Wall. cat. n. 3286. (v. s. in h. Wight.)
- γ. Sirmorensis*, caule foliisque pube brevi sparsâ scabridis, foliis super. serratis. — in agro Sirmorensi legit cl. Gerard. *Er. multicaule* B. Wall.! l. c. Fortè species propria? (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
59. *E. HISPIDUM* (DC. in Wight contrib. p. 9), caule basi suffruticuloso erecto ramoso hispido, foliis oblongis hinc indè grossè dentatis subpuberulis, inferioribus obtusis basi attenuatis, superioribus acutis basi dilatatis cordato-amplexicaulibus, capitulis ad apices ramorum 1-2, invol. squamis linearibus acuminatis, ligulis innumeris angustissimè linearibus elongatis 1-2-seriatis, fl. fœm. pluriseriatis vix ligulatis, centralibus hermaphrod. 5-dentatis. *♂ 2* in Benghalo et fortè agro Maderaspetano in arenosis (Wight!). *Aster Benghalensis* Heyn.! herb. Erig. hirsutum Wall.! cat. et herb. n. 2976. comp. n. 86. non Lour. nec Persh et Conyza graveolens Wall.! cat. n. 3106. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
60. *E. SUBLYRATUM* (Roxb. ined. ex Wight.! herb. DC. in Wight contr. bot. ind. p. 9), caule ascendenti-erecto molliter hispido subsimplici, foliis utriusque villosis obovatis grossè inciso-dentatis dentibus obtusis mucronatis, infer. basi attenuatis incisis sublyratis, summis cordato-amplexicaulibus, corymbo terminali 7-9-cephalo, capitulis pedicellatis, invol. squamis linearibus acuminatis glabriusculis disco subæqualibus. (2) in arenosis et apricis ad Negapatam prov. Tanjore legit cl. Wight, ad Rampur in Rohilkund et in Nepaliâ Wall.! cat. et herb. n. 3007. comp. 117. *Conyza hirsuta* Wall. non Linn. *Conyza lacera*

Burm. herb. Ligulæ albæ Achænia glabra parva elliptica compressa. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

Sect. ? IV. PTERIGERON.

Ligulæ pluriseriales disco longiores fœminæ. Fl. centrales pauci tubulosi 5-dentati hermaphrod. aut stylo subdiviso abortientes et ferè masculi? Cor. tubulosæ, fauce non dilatata. Anther. basi ecaudatæ coadunatæ. Stylus fl. herm. inclusus apice vix bifidus. Pappus scaber rigidulus. Recept. alveolatum. Habitus ferè Conyzæarum decurrentium. Sectio dubia.

61. E. ? DECURRENS, sparse hirsutum, caule suffrutescente?, foliis breviter adnato-decurrentibus ovatis acutis integris aut ad basin dentatis, capitulis ad apices ramulorum solitariis subpaniculatis dispositis, invol. squamis imbricatis pluriserialibus acuminatis, ext. ovatis dorso scabris, int. scariosis glabris, achæniis sericeo-villosis. — in Novâ Hollandiâ orient. aut in Timor. Ligulæ lineares pappo scabro longiores apice tridentatæ pluriseriales. Fl. herm. seu stylo indiviso masculi centrales pauci tubulosi fauce vix dilatata. Antheræ concretæ. Recept. alveolatum. Pappus densè barbellatus. An prioris congener? An genus proprium? (v. s. comm. à Mus. reg. Par. ex itin. Bandin.)

Sect. v ? ELEUTHEROGERON.

Ligulæ pluriseriales disco longiores fœminæ. Fl. centr. pauci hermaphr., tubo tereti, fauce dilatata campanulata 5-dentata. Anth. liberæ. Styli rami subforcipati acuti hispiduli. Recept. punctatum. Achænia villosa, pappo barbellulato. — Folia breviter decurrentia.

62. E. ANOMALUM, caule erecto glabrinsculo, ramis puberulo-scabris, foliis caulinis semiamplexicaulibus breviter decurrentibus obovato-oblongis basi attenuatis grossè serratis utrinquè subscabridis, summis oblongis non decurrentibus, capitulis breviter pedicellatis ad apices ramorum subcorymbosis, invol. 1-2-serialis squamis oblongo-linearibus subæqualibus, ligulis disco paulò longioribus. — in Indiâ orientali legit cl. Royle. Flor. ligulati fœminei 20-25, ligulâ lineari integrâ vix stylo longiore. Fl. hermaphr. 5, fortè stigm. abortivo submasculi. Antheræ liberæ certè ecaudatæ. Faux cor. abruptè dilatata. Recept. non sensim alveolatum. Achænia fertilia compresso-plana obovato-oblonga-in disco pubera. (v. s. comm. à cl. Royle sub n. 144.)

+ Species non satis notæ.

63. E. ? CANESCENS (Willd. sp. 3. p. 1953), caule erecto ramoso cano-pubescente, foliis lanceolatis integerrimis pubescentibus, pedunculis axillaribus 1-cephalis. — Patr. ign. Cor. radiata flava et ideo à genere verisim. removenda?
64. E. CAUCASICUM (Stev. act. soc. h. nat. mosc. 4. p. 63), hispidum, caule oligocephalo, foliis integerrimis obtusè mucronatis, radicalibus oblongis petiolatis, caulinis cordato-ovatis sessilibus, capitulis longè pedunculatis, achænio hispido longitudine pappi. — in Caucasi alpe Kaischaur. Invol. valdè hirtum apice coloratum. Capit. E. acri majus.
65. E. ? COCHINCHINENSE (Lour. fl. coch. 2. p. 611, in obs. Pers. euch. 2 p. 431), caule erecto ramoso polycéphalo, foliis lanceolatis sinuatis subserratis, caulinis semiamplexicaulibus, paniculâ terminali, capitulis oblongis. — in Cochinchinâ. Erig. Philadelphicum Lour. l. c. non Linn. Flores radii tenuissimi (an ligulati?) fœminei erecti cum disco lutei.
66. E. DIVERSIFOLIUM (Cav. anal. scienc. nat. 1801. v. 4. p. 88), caule herbaceo glabro, foliis inferioribus lato-lanceolatis serratis petiolatis, rameis linearibus sessilibus. — in insulâ Teneriffâ.
67. E. DUBIUM (Spreng. syst. 3. p. 519), caule erecto simplici villosa, foliis sessilibus linearibus acutis pilosisculis, infimis subserratis, pedunculis paucis

- erectis 1-cephalis, radio discum superante. — ad flum. Magnum (Rio-Grande) Brasilæ. Cæt. ign.
68. *E. FASTIGIATUM* (Horn. h. hafn. add. p. 971), foliis spathulatis basi subpinatifidis apice remotè serratis, ramis paniculatis, pedunculis 1-cephalis. ① patr. et cæt. car. ign. †.
69. *E. FILIFORME* (Spreng. syst. 3. p. 520), caule herbaceo subramoso, foliis filiformibus subflexuosis longè pilosis, pedunculo solitario 1-cephalo. — ad flum. Magnum (Rio-Grande) Brasilæ (Sell.). Cæt. ign. An fortè *Stenactis pilosa*?
70. *E. ? FRAGRANS* (Spreng. syst. 3. p. 518), caule ramoso foliisque lanceolatis serratis incano-villosis, ramis 1-cephalis subviscosis. — patr. et cæt. car. ign. *Senecio fragrans* Lehm. ex Spreng. †.
71. *E. HIRSUTUM* (Lour. fl. coch. ed. Willd. 2. p. 611. non Pursh), caule erecto hispidò, foliis lineari-lanceolatis subserratis pilosissimis, capitulis paniculatis, invol. subsquarroso. — circa Cantonem Sinarum. Herba 2-pedalis. Discus luteus. Ligulæ cœruleæ lineares multæ.
72. *E. LÆVIGATUM* (Rich. in act. soc. h. nat. Par. 1790. p. 105), caule angulato debili piloso, ramis paniculatis, foliis subspathulato-linearibus integris lævigatis subpilosis ciliatis. ② (ex Horn. h. hafn. 2. p. 807). Hab. in Cayennâ.
73. *E. LEHMANNI* (Spreng. syst. 3. p. 517), caule humili arachnoideo-lanato, foliis petiolatis spathulatis mucronatis integerrimis subtus subtomentosis, capitulis corymbosis aggregatis densè lanatis. ① in Ægypto ex hort. angl. Er. pusillum Lehm. sem. h. hamb. 1823. p. 8. non Nutt. Er. *Lehmanni* Lehm. in act. nov. nat. cur. 14. p. 2. p. 809. Recept. convexum. Flor. et fr. indescr.
74. *E. LINEARIFOLIUM* (Cav. anal. scienc. nat. 1801. v. 4. p. 89), herbaceum villosum, foliis linearibus, pedunculis foliosis mono- aut polycephalis. — Circa Mogador. †.
75. *E. MONTEVIDENSIS* (Spreng. syst. 3. p. 519), fruticosum, ramis 1-cephalis, foliis confertis lineari-filiformibus flexuosis strigillois, pedunculis villosis, radio discum superante. — in Montevideo (Sell.). Cæt. ign.
76. *E. MYROSOTIS* (Pers. ench. 2. p. 431. n. 33), caule simplici sub-1-cephalo foliisque linearibus hirsutis. — ad fretum Magellanicum. Er. *myosotoides* Juss. herb. Caulis ferè pedalis. Flores purpurascens. Cæt. ign. †.
77. *E. ? PILOSUM* (Pers. ench. 2. p. 431. n. 10. non Walt.), caule simpliciusculo piloso, foliis brevi-ovatis petiolatis grossè serratis pubescentibus, capitulis subpaniculatis. — patr. et cæt. car. ign. †.
78. *E. ? PINIFOLIUM* (Poir. dict. 8. p. 490), glabrum, caule adscendente parèe ramoso, foliis confertis filiformibus acutis submucronulatis integerrimis, pedunculis nudis 1-cephalis, invol. hemisphærico vix pilosiusculo, ligulis invol. longioribus. — in Bonariâ (Comm.). Flores lutei. Certè à genere videtur alienum.
79. *E. ? RESINOSUM* (Spreng. syst. 3. p. 519), villosum granis resinosis asperum, foliis sessilibus linearibus acutis paucidentatis, pedunculis paucis erectis, radio discum superante. — ad flum. Magnum (Rio-Grande) Brasilæ. Ad *Asteris* genus relatum cum dubio à cl. Less. syn. 182.
80. *E. RETROFLEXUM* (Poir. suppl. 5. p. 464), caule glabro paniculato-ramoso, foliis angustissimè linearibus glabris vetustate retroflexis, pedunculis erectis subpubescentibus alternis sub-1-cephalis, invol. brevi imbricato glaberrimo. 2/ in Amer. bor. Flor. et fruct. indescr.
81. *E. ? ROTTLERI* (Spreng. syst. 3. p. 518), suffruticosum, foliis oblongo-lanceolatis subdentatis hispidiusculis, caulinis amplexicaulibus, ramis elongatis filiformibus 1-cephalis, invol. glabro. — in Indiâ orient. An *Conyza* seu *Blumæ* species?
82. *E. ? SCANDENS* (Thunb. fl. jap. 313), caule scandente, foliis ovatis serratis

villosis, capitulis axillaribus. — in Japoniâ. Certissimè alius generis est sed frustula suppetentia in herb. Thunb. nimis manca sunt. (Less. syn. 190.)

83. E. SUMATRENSE (Retz. obs. 5. p. 28), tomentosum, foliis sessilibus lanceolatis dentatis, paniculâ racemosâ. — in Sumatrâ. Flosc. radii dicuntur capillares undè verisim. species ad Conyzam seu ad Blumeam referenda videtur, sed vix nota.

84. E. WAHLII (Gaudich. ann. sc. nat. 1825 mai p. 104. voy. Freyc. 468), caulis herbaceis subsimplicibus erectis folisque glabris obovato-lanceolatis acutiusculis remotè serratis infernè angustatis basi vaginatis. — in insul. Maclovianis. Capit. ignota.

†† *Species exclusa.*

E. ÆGYPTIACUM Linn. = Conyza ægyptiaca.

E. ALATUM Don = Blumea alata.

E. AMBIGUUM Nutt. = Stenactis ambigua.

E. ANNUUM Linn. = Stenactis annua.

E. ASTEROIDES Roxb. = Blumea pubiflora.

E. CAMPHORATUM Linn. = Blumea camphorata.

E. CANADENSE Ten. = Conyza ambigua.

E. CAPENSE Houtt. = Webbia pinifolia.

E. CAROLINIANUM Linn. = Phalacrocoma obtusifolium.

E. CILIATUM Ledeb. = Conyza Altaica.

E. CONTORTUM Desf. = Conyza ambigua.

E. CRISPUM Pourr. = Conyza ambigua.

E. DELPHINIFOLIUM H. et B. = Polyactidium delphinifolium.

E. DIFFUSUM Pers. = Aplopappus diffusus.

E. DUBIUM Spr. = Callistephus Wightianus.

E. FOLIOSUM Don = Conyza flavida.

E. GLANDULOSUM Poir. = Chrysopsis graminifolia.

E. GLUTINOSUM Linn. = Jasonia glutinosa.

E. GLUTINOSUM Desf. = Psiadia glutinosa.

E. GNAPHALIOIDES H. B. K. = Heterochaeta gnaphalioides.

E. GOUANI Linn. = Conyza Gouani.

E. GOUANI Jacq. = Conyza ægyptiaca.

E. GRAMINEUM Linn. = Arctogeron gramineum.

E. GRAVEOLENS Linn. = Inula graveolens.

E. HETEROPHYLLUM Willd. = Stenactis annua.

E. HIERACIFOLIUM Don = Blumea hieracifolia.

E. HIRTUM Thunb. = Conyza incisa β.

E. HYSSOPIFOLIUM Michx. = Phalacrocoma obtusifolium.

E. INCANUM Wahl = Conyza incana.

E. INCISUM Thunb. = Conyza incisa.

E. INULOIDES Poir. = Pulicaria inuloides.

E. JAPONICUM Thunb. = Conyza Japonica.

E. JOSSLEYI Pers. = Conyza aspera.

E. LEUCANTHEMUM Don = Blumea leucanthema.

E. LINIFOLIUM Willd. = Conyza ambigua.

E. LINOSYRIS Clairv. = Linosyris vulgaris.

E. LONGIFOLIUM Desf. = Jasonia longifolia.

- E. MOLLE* Don = *Blumea trichophora*.
E. NERVOSUM Willd. = *Chrysopsis graminifolia*.
E. NUDICAULE Michx. = *Stenactis verna*.
E. OBLIQUUM Linn. = *Blumea amplexans*.
E. PINNATIFIDUM Don = *Conyza absinthifolia*.
E. PINNATIFIDUM Thunb. = *Conyza pinnatifida*.
E. PINNATUM Linn. f. = *Conyza pinnatilobata*.
E. PUBESCENS H. B. K. = *Heterochaeta pubescens*.
E. RUTILUM Poir. = *Conyza callocephala sericea*.
E. SCABRUM Thunb. = *Pulicaria erigeroides*.
E. SCABRUM Pers. = *Conyza aspera*.
E. SCARIOSUM Moench. = *Conyza Gouani*.
E. SERRATUM Forsk. = *Conyza ægyptiaca*.
E. SICULUM Linn. = *Jasonia sicula*.
E. SQUARROSUM Clairv. = *Inula conyza*.
E. STIPULATUM Schum. = *Blumea Guineensis*.
E. TRICUNEATUM Linn. = *Baccharis cuneata*.
E. TRISULCUM Don = *Blumea trisulca*.
E. TUBEROSUM Linn. = *Jasonia tuberosa*.
E. VISCOSUM Linn. = *Inula viscosa*.
E. VISCOSUM Desf. = *Psiadia glutinosa*.
E. WIGHTIANUM Wall. = *Callistephus Wightianus*.

CXL. RHYNCHOSPERMUM Reinw. in *Blume bijdr.* p. 902.

Capitulum multiflorum radiatum, ligulis 2-3?-serialibus fœmineis vix disco longioribus, fl. disci tubulosis 5-dent. herm. Invol. hemisphæricum, squamis numerosis imbricatis adpressis. Recept. subconvexum scrobiculatum. Antheræ ecaudatæ. Styli rami oblongi breves. Achænia compressa glabra basi attenuata apice in rostrum brevem producta. Pappus 1-serialis setosus, setis apice tortilibus seu inflexis. — Herba perennis Javanica glabriuscula. Caulis teres, ramis ultimis verticillatis. Folia alterna lanceolata utrinquè acuminata subpetiolulata remotè serrulata minutissimè scabra. Capitula ad axillas foliorum solitaria brevè pedunculata, summa nuda subterminalia. Color flor. in sicco pallidus. — Fructu rostrato accedit ad Myriactidem, sed pappo facile differt.

1. *R. VERTICILLATUM* (Reinw. et Blum. l. l. c.). 2l in montanis provinciæ Javanicæ Tjanjor flor. aug. Planta rarissima et distinctissima. Folia 3 poll. longa, 5 lin. lata. (v. s. comm. ab amic. Blume.)

CXLI. MICROGYNE Less. syn. 190. non Cass. (1)

Capitulum multiflorum.... flor. in ambitu pluriserialibus fœmineis, in disco hermaphroditis regularibus. Achænium erostre. Pappus coniformis pluriserialis. — Herba Brasiliensis annua ramosa canescens. Folia alterna trifurcata, laciniis linearibus. Capitula solitaria ramos

(1) MICROGYNE Cass. = *Eriocephalis* sect. *Cryptogyne*.

supernè aphyllis terminantia. Pappus rufescens. — An fl. radii ligulati ut in *Erigeris* aut filiformes ut in *Solenogyneis*?

1. M. *TRIFURCATA* (Less. l. c.). ① in *Brasiliâ* (Sellow.). Cæt. ign.

Subdiv. 4. HETEROPAPPEÆ.

Radius 1-serialis. Pappus diversus in disco et radio.

CXLII. SIMBLOCLINE.

Capitulum radiatum, ligulis 1-serialibus fœmineis linearibus elongatis, fl. disci tubulosis 5-dent. hermaphr. aut verisim. abortu styli sterilibus. Invol. multiseriale, squamis exter. brevibus lanceolatis acutis dorso tomentosis imbricatis, inter. multò longioribus glabris discum superantibus squarrosis glabris. Recept. alveolatum, alveolis elevatis dentatis ferè fimbriiferis. Styli radii elongati apice bifidi glabri, disci inclusi brevè bifidi. Anthæræ ecaudatæ. Achænia radii oblonga subcompressa parçè pilosa et subglandulosa, disci tenuia. Pappus radii 1-serialis setosus scabridus, disci biserialis, serie utrâque setiformi exter. brevi, inter. setis radii simili. — Frutex Peruvianus. Folia oblongo-lanceolata integerrima supernè glabra atro-viridia subtùs ramulis rufescenti-tomentosa. Capitula pedicellata laxè corymbosa. Flor. ex sicco vid. purpurascens? Ligulæ numerosæ circ. 30-40. — Genus differt à *Diplostephii* sect. *Amphistephio* pappo radii simplici nec cum centrali conformi, à *Tetramolopio* pappo disci duplici nec simplici et recept. profundè alveolato. — Nomen ex *σμβλος*, alveus apum, et *κλίση*, receptaculum.

1. S. *HÆNKEI*. ♂ in Peruviâ legit cl. Hænke ex herb. reg. Acad. Monac. Folia 2 poll. (petiolo 3 lin. incl.) ferè longa, 6 lin. lata. (v. s. in h. suprâ cit.)

CXLIII. *HETEROPAPPUS* Less. *syn.* 189. — *Asteris* sp. Thunb. — *Calimeris* sp. Nees.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus fœmineis ligulatis, disci tubulosis 5-dentatis hermaphroditis. Recept. Invol. biseriale squamis patenti-divergentibus. Achænium erostre. Pappus radii coroniformis, disci pilosus 1-serialis. — Herba Japonica erecta ramosa hispida. Folia alterna, infer. oblongo-obovata crenata, super. lanceolata. Capitula terminalia solitaria parva, radio albo.

1. H. *HISPIDUS* (Less. l. c.). — in Japoniâ. *Aster hispidus* Thunb. fl. jap. 315. Banks ic. Kämpf. t. 29. non Lam. — *Keitsja* Kämpf. am. ex. 5. p. 876. *Calimeris* hispida Nees ast. p. 227.

CXLIV. *PHALACROLOMA* Cass. *dict.* 39. p. 404 (excl. sp. 2). Less. *syn.* 188. — *Erigerontis* sp. Linn.

Capitulum pluriflorum radiatum, ligulis fœmineis 1-serialibus disco hermaphrodito fertili regulari multò longioribus. Recept. nudum planiusculum. Invol. cylindræo-campanulatum 2-3-seriale imbricatum. Achænia compressa. Pappus radii coroniformis brevis 1-serialis, disci biserialis, exterior radiali conformis, interior setis multis longis scabris. — Herba boreali-americana erecta supernè ramosa. Folia alterna remota sessilia oblonga aut linearia integerrima hispidula. Capitula laxè paniculata. Discus flavus. Ligulæ ex Dill. et Pursh flavæ, ex Cass. et Less. forsan albæ.

1. *P. obtusifolium* (Cass. l. c.). \mathcal{H} in demissis Amer. bor. ad sinum Hudsonis et juxta lacum Mistassins (Michx.), à Canadâ ad Carolinam (Pursh), sed ab Elliot in Carolinâ non indicatur. Virga-Aurea Caroliniana Dill. elth. 2. p. 412. f. 394. Erigeron Carolinianum Linn. sp. 1210. Erigeron hyssopifolium Michx. fl. bor. am. 2. p. 123. An stirps Cassiniana certè eadem ac Dilleniana seu Linnæana? Dillenius flores ligulatos pappo eodem ac tubulosos donatos exhibet!

† *Species exclusa.*

P. acutifolium Cass. = *Stenactis annua*.

CXLV. MINURIA.

Capitulum multiflorum, fl. radii 1-serialibus fœmineis ligulatis, disco tubulosis 5-dentatis styli abortu masculis. Invol. pauciserialè imbricatum, squamis linearibus acuminatis margine scariosis. Recept. planum nudum. Achænia disci subturbinata sericeo-villosissima, pappo multisetiformi scabro, villis achæniî summis pappum duplicem simulantibus; disci gracilia glabra inania pappo 1-seriali, paleis aliis brevibus inter se concretis aliis paucis setiformibus longis scabris. — Suffrutices Novæ-Hollandiæ interioris indigenæ erecti glabri ramosissimi. Folia alterna acerosa seu linearia acuta integerrima. Rami apice summo plûs minûs nudi 1-cephali. Ligulæ albæ. — Genus pappo heterogeneo distinctissimum, habitu Erigeropsidis, caractere ferè Stenactidis. Nomen ex $\mu\upsilon\rho\omicron\varsigma$, exilis.

1. *M. leptophylla*, ramis usque ad capitulum folia parca gerentibus, ligulis disco triplo longioribus. \mathcal{H} ad apices collium sterilem Novæ-Hollandiæ interioris ad flumen Lachlan aprili flor. legit cl. Cunningham. Ligulæ circ. 20 lineares. (v. s. comm. à cl. A. Cunningham.)
2. *M. tenuissima*, ramis apice longiusculè nudis, ligulis disco duplo longioribus. \mathcal{H} in collibus scopulis sterilibus Novæ-Hollandiæ interioris ad flum. Lachlan aprili flor. legit cl. Cunningham et mecum comm. Ligulæ pauciores et paulo longiores ex sicco videntur. Capitula dimidiò ferè minora. (v. s.)

CXLVI. STENACTIS *Nees ast. p. 273. Less. syn. 189.* — Diplopappi et postea Stenactidis et Phalacrolomatis sp. *Cass. dict. 37. p. 462 et 485 et 39. p. 405.* — Asteris sp. *Linn.* — Erigerontis sp. *Ait.* — Pulicariæ sp. *Gærtn.*

Capitulum multiflorum heterogamum, fl. radii uniserialibus fœmineis angustè ligulatis, disci hermaphr. 5-dentatis. Recept. nudum punctulatum planum seu convexum. Invol. squamæ 2-3-seriales imbricatæ angustæ. Achæniû oblongum compressum. Pappus radii 1-serialis setaceus deciduus, disci duplex, exter. brevissimus, interior pappo radii similis. — Herbæ erectæ ramosæ. Folia alterna dentata. Capitula ad apices ramorum solitaria disco luteo, radio albo aut pallidè violaceo. — Nomen ex $\sigma\tau\epsilon\nu\omicron\varsigma$, angustus, et $\alpha\pi\tau\iota\varsigma$, radius.

* *Americana.*

1. *S. annua* (*Nees ast. p. 273*), caule erecto apice polycephalo corymboso, foliis infer. ovatis, super. lanceolatis dentato-serratis pilosiusculis, invol. setoso. ① in Americâ bor. secus vias et in ruderalis frequens, undè migravit in Europam. Habeo lectam circa Bassano à fratre meo, inter Pataviam et Vicenzam à cl. Duby, indic. in Belgio à Dekin et Passy fl. brux. p. 55. Diplopappus dubius *Cass. bull. philom. 1817. p. 137. 1818. p. 77.* Stenactis dubia *Cass. dict. 37. p. 485.* Phalacroloma acutifolium *Cass. dict. 39. p. 405.* Pulicaria annua *Gærtn. fr. 2. p. 462.* Cineraria corymbosa *Moench.* Erigeron annuum *Pers. ench. 2.*

- p. 431. *Erigeron heterophyllum* Willd. sp. 3. p. 456. Nutt.! gen. 147. *Aster annuus* Linn. sp. 1229. fl. dan. t. 486. — Corn. can. 193. t. 194. Herba valde quoad habitum et formas diversas foliorum varians. Ligulæ albæ. (v. v. et s.)
2. *S. AMBIGUA*, vix pubescenti-scabrida, caule erecto apice ramoso subcorymboso, foliis elongatis oblongo-linearibus, infer. basi in petiolum longè attenuatis subserratis, super. linearibus integerrimis, invol. squamis puberulo-scabris. — in Georgiâ (Nutt.! gen.) et in territorio Arkansano (Nutt.! in litt.). *Erigeron ambiguus* Nutt. gen. am. 2. p. 147. Ell. sketch 2. p. 395? Ligulæ numerosæ 1-seriales. Pappus radii brevis, disci duplex, ext. paleaceus brevis. (v. s. comm. à cl. Nutt.)
3. *S. PILOSA*, caule erecto apice corymboso-ramoso tereti pilis patulis subhispido, foliis linearibus acutis integerrimis pilis longis et pube breviori hirtellis, invol. squamis acuminatis margine membranaceis dorso puberis, achæniis villosis. ① in Brasiliæ prov. Rio-Grande. Ligulæ albidæ tenues. Pappus exter. disci parçè conspicuus. Herba 6-10 pollic. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1046 misc.)
4. *S. VERNA* (Nees ast. 275), caule subaphyllo glabrinusculo composito-subcorymboso, foliis radicalibus rosulatis oblongis acutis glabris subdentatis, caulinis distantibus linearibus integerrimis, invol. vix pubescente. ② in depressis humidis Carolinæ et Georgiæ. *Aster vernus* Linn. sp. 1230. *Erigeron nudicaule* Michx. fl. bor. am. 2. p. 134. Ell. sketch 2. p. 332. *Aster vernus* et *Erigeron nudicaule* Pers. ench. *Doronicum lævifolium* Walt. fl. car. p. 205? Ligulæ lineares 1-seriales albæ ex Ell., purpureæ in spec. meo Georgico, non invol. æquales ut dicunt cl. Nees, Michx., etc., sed duplò longiores ut dixit cl. Elliot. An ideo longitudine varia? An duæ species confusæ? Pappus duplex, in radio depauperatus. (v. s.)
5. *S. STRIGOSA*, caule erecto sparse et patentium piloso, foliis lanceolatis utrinquè attenuatis dentatis strigoso-hispidis, capitulis laxè corymboso-paniculatis, invol. glaberrimo, ligulis numerosis filiformibus. ②? in Pennsylvania (Muhl.), Noveboraco (Torrey!). *Erigeron strigosus* Muhl. in Willd. sp. 3. p. 1956. *Erig. australe* Horn. ex Spreng. An *Erig. strigosus* Ell. sketch 2. p. 394 et *Doronicum ramosum* Walt. fl. car. 205 in Carolinâ cresc. ad hanc speciem referenda? (v. s. comm. à cl. Torr.)

†* Asiaticæ.

6. *S. BELLIDIODES* (Nees ast. p. 274), caule adscendente angulato-striato oligocephalo pilosiusculo, foliis lanceolatis acutis apice subserrulatis pubescenti-scabridis, infer. hinc indè grossè serratis, summis sublinearibus integerrimis, invol. setoso. ② in Nepalîæ superioris montanis apricis et in Kamaon. *Aster bellidioides* Don prod. fl. nep. p. 177. Wall.! pl. asiat. rar. 2. p. 1. t. 101. cat. n. 2978. comp. 88. Ligulæ lilacinæ. (v. s. comm. ab hon. Cæt. merc. angl. Ind. or.)
7. *S. MULTIRADIATA* (Lindl. in litt. 1835), caulibus à basi ramosis erectis infer. glabris sup. villosis sulcatis, foliis sessilibus ovalibus basi subcuneatis et integerrimis apice sparse et grossè serratis mucrone callosio terminatis utrinquè puberulis, ramis 1-cephalis sub capitulis brevi spatio nudis, invol. squamis linearibus acuminatis hirtellis disco æqualibus, ligulis linearibus numerosissimis, achæniis puberulis. ② in Indiæ montibus boreali-occid. leger. cl. Wallich et Royle. *Aster multiradiatus* Wall.! cat. et herb. n. 2969. comp. n. 79. *Erigeron radiosus* Spreng. in litt. ad Wall. *Aster inuloides* Don prod. fl. nep. 178? Nees ast. 36? sed ex Lindl. adn. descr. Neesii fortè è pluribus speciebus petita? (v. s.)

† *Species exclusæ.*

- S. ALPINA* Cass. = *Erigeron alpinum*.
S. DELPHINIFOLIA Cass. = *Polyactidium delphinifolium*.
S. GLAUCA Nees = *Erigeron glaucum*.

S. GNAPHALIOIDES Cass. = *Heterochæta gnaphalioides*.

S. PUBESCENS Cass. = *Heterochæta pubescens*.

S. SPECIOSA Lindl. = *Erigeron speciosum*.

CXLVII. GYMNSTEPHIUM Less. syn. 185. Nees ast. 253.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis uniserialibus, disci tubulosis 5-dentatis masculis sterilibus. Recept. nudum areolatum. Invol. squamæ imbricatæ subtriseriales. Achænia radii plano-compressa erostria subpilosa, disci linearia incana. Pappus disci obsoletus paucisetosus, setis plumoso-serratis, radii nullus. — Plantæ herbacæ Capenses ramosæ. Folia alterna linearia integerrima. Capitula pedunculata solitaria parva, radio cœruleo, disco luteo. Involucra sæpius glandulis oblongis ut in *Tagete* notata. — Affine dicitur *Chariei*, sed differt disco masculino. Affinis fortè *Feliciæ* et *Mairiæ*, sed differt achæniis radii epapposis, disci paucisetis. — Genus fortè nimis artificiale?

§. 1. *Glabra, invol. glandulis oblongis notatis.*

1. G. GRACILE (Less. l. c.), herbaceum? glabrum, ramis floridis supernè aphyllis, invol. squamis ellipticis glandulis oblongis notatis. 2½? ad Cap. Bonæ-Spei in Tradouw. Facies *Mairiæ* microcephalæ.
2. G. FRUTICOSUM, glabrum fruticosum, ramis floridis apice villosis ferè ad apicem folia seu squamas elongatas gerentibus, invol. squamis lineari-oblongis exter. acutis, inter. obtusis et glandulas oblongas gerentibus. 5 in Africa Capensi ad Zwelendam alt. 200-300 ped. legit cl. Drege. Fruticulus circ. pedalis. Folia 9-12 lin. longa, 1-1½ lin. lata margine subrevoluta, uninervia. (v. s.)

§. 2. *Hirsuta, involucris eglandulosis.*

3. G. HIRSUTUM (Less. l. c.), herbaceum? hirtum, invol. squamis linearibus acuminatis. 2½? ad Cap. Bonæ-Spei in Steenbaks rivier ad Swarteberge. Capitula magn. Bellidis perennis.

CXLVIII. CHARIEIS Cass. bull. philom. 1817. p. 68 et 1821. p. 12. Dict. 8. p. 191. 24. p. 369. 37. p. 463 et 489. Less. syn. 389. — *Kaulfussia* Nees hort. ber. p. 53. t. 11. ast. p. 268 non Blume.

Capitulum multiflorum, heterogamum, fl. radii ligulatis fœmineis, disci hermaphr. tubulosis 5-dentatis. Recept. alveolatum, alveolorum marginibus subdentatis. Invol. 2-seriale campanulatum, squamis exter. planis, interioribus complicato-carinatis. Stylus disci ramis apice quasi trapezoideis depressis et ibi dorso subpuberulis. Achænia obovato-cuneiformia compressa nervo calloso cincta cæterum subnervia hirtella, radii sæpè inania semper epapposa aut 1-setosa. Pappus disci setis capillaribus plumosis 1-serialibus. — Herba monocarpica Capensis. Folia infima opposita, cætera alterna, oblonga lanceolatave basi trinervia cum ramis subhirtellis. Pedunculi terminales aphylli. Capitula nascentia subflava demùm cœrulea, disco et radio sæpè discoloribus aut concoloribus. Potiùs ex habitu et stigmate ad *Astereas* quàm ad *Senecioneas* meo sensu pertinere videtur.

1. C. HETEROPHYLLA (Cass. ll. cc.). ① ② ad Cap. Bonæ-Spei (Nees), ad Zilverfontain (Dreg!). Char. cœrulea Nees ast. p. 269. cum var. 3 nempè:

- α. caule erecto oligocephalo, disco luteo ligulis angustioribus. — *C. heterophylla* Cass.
 β. caule erecto oligocephalo disco radioque cœruleis, ligulis latioribus. — *C. cœrulea* Cass. dict. 24. p. 370.
 γ. caule ramosiore sæpè diffuso, disco radioque cœruleis, ligulis latioribus. *C. Neesii* Cass. dict. 24. p. 372. *Kaulfussia amelloides* Nees hor. ber. 53. t. 11. (v. v. c. et s.)

CXLIX. CHÆTOPAPPA. — *Chætophora* Nutt. in litt. 1825 non Schrank nec Brid., nec Hornsch. (1).

Capitulum multiflorum radiatum, fl. radii 1-serialibus fœmineis ligulatis, disci hermaphrod. tubulosis 5-dentatis. Recept. angustum nudum. Invol. squamæ laxè imbricatæ apice scariosæ acuminatæ. Achænia teretiuscula. Pappus nunc duplex nunc subduplex, ext. latè et hyalinè paleaceo, in fl. disci brevi in fl. radii longiore, inter. setis circ. 5 rigidis scabris constante. — Herba pusilla annua Boreali-Americana. Tota pubescenti scabra, sæpius multicaulis ramosa, rariùs in siccis simplex. Folia caulina alterna linearia aut lineari-spathulata integerrima, radicalia obovato-spathulata in petiolum attenuata. Rami apice nudi 1-cephali. Invol. glabrum nitens. Ligulæ albæ demùm subpurpurascens. Discus flavus. — *Boltoniæ* paulò affinis videtur.

1. *C. ASTEROIDES* (Nutt. in litt. sub *Chætophora*). (1) in territorio Arkansano. Herba 2-5 poll. longa. (v. s. comm. à cl. invent.)

CL. BOLTONIA *L'Hér. sert. angl. p. 27. Cass. dict. 37. p. 464 et 491. Nees ast. 234.* — *Madea Soland. ined. ex L'Hér. mss.* — *Matricariæ* sp. Linn. — *Chrysanthemi* sp. *Walt.*

Capitulum radiatum, ligulis 1-serialibus linearibus fœmineis fertilibus, disco hermaphr. fertili. Receptaculum hemisphæricum alveolatum. Involucri squamæ discum æquantès biserialès adpressæ margine membranaceæ. Achænia plano-compressa callosò-marginata glabriuscula seu hispidula. Pappus brevissimus setulosus nunc subæqualis, nunc in floribus disci setis duabus elongatis subulatis crassioribus constans. — Herbæ rhizocarpicæ Boreali-Americanae erectæ ramosæ Asteris facie. Folia alterna subverticalia lanceolata glaucescentia lævia margine scabra integerrima aut inferiora dentata. Capitula laxè corymbosa. Discus luteus. Radius albus aut pallidè purpurascens.

1. *B. GLASTIFOLIA* (*L'Hér. l. c.*), foliis lanceolatis glaucescentibus, inferioribus serratis, capitulis brevius pedicellatis, achæniis obovatis, pappi disci setis plurimis inæqualibus, 2 oppositis crassis elongatis rigidulis. 2/ in Amer. bor. à Pennsylvania ad Carolinam ad ripas rivulorum et paludum. Willd. sp. 3. p. 2162. Nees ast. 235. bot. mag. t. 2381. Ell. sketch 2. p. 399 (excl. signo?). *Madea glauca* Soland. in hort. Lee ex *L'Hér.*! *Matricaria glastifolia* Hill. h. kew. 19. t. 3? (v. v. c.)
 2. *B. DIFFUSA* (Ell. sketch 2. p. 399), glaberrima, foliis lineari-lanceolatis integerrimis margine scabris, paniculæ diffusæ pedunculis elongatis, achæniis obovatis, pappi disci setis plurimis inæqualibus 2 oppositis crassis elongatis rigidulis. 2/ in Amer. bor. pinguibus humidis inter Chatahouchie et Alabama. Anonymos asteroides Michx. herb. *Boltonia asteroides* bot. mag. t. 2554 excl. syn. Capitula minora quàm *B. glastifoliæ*, cui accedit ex pappo disci, sed sæpè

(1) *Chætophora* Schrank est genus à plerisque inter Algas admissum. *Chætophora* Brid. est *Leskeæ* sp. *Chætophora* Hornsch est *Hookeriæ* sp.

ob folia integerrima et capitula longè pedicellata, in hortis cum *B. asteroide* confusa. (v. s.)

3. *B. ASTEROIDES* (L'Hér. l. c.), foliis integerrimis lineari-lanceolatis, capitulis longè pedicellatis laxè subcorymbosis, achæntiis ovalibus glabris, pappo brevissimo deciduo in radio et disco subæquali. \mathcal{U} in paludosis et ad lacuum ripas Carolinæ et terræ Illinoensium? Willd. sp. 3. p. 2162. Nees ast. 236. *Matricaria asteroides* Linn. mant. 116. *Chrysanthemum Carolinianum* Walt. fl. car. 204. (v. v. c.)

CLL. SOMMERFELTIA Less. gen. 180 non Schum. (1)

Capitulum radiatum, ligulis fœmineis 1-serialibus, disco sterili regulari. Recept. epaleaceum. Invol. imbricatum. Achænium erostre. Pappus pilosus, radii pluri-, disci 1-serialis. — Fruticulus Brasiliensis. Folia conferta pinnatisecta rigida glabra, segmentis filiformi-subulatis spinescentibus. Capitula solitaria ramos supernè aphyllis terminantia.

1. *S. SPINULOSA* (Less. l. c.). \mathfrak{f} in Brasiliâ. *Conyza spinulosa* Spreng. syst. 3. p. 510. Folia cinerea. Pedunculi aggregati (Spr.).

Subdiv. 5. BELLIÆ.

Radius 1-serialis. Pappus squamellatus.

CLII. CALOTIS Brown in bot. reg. n. 504.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus fœmineis, disci tubulosis 5-dentatis masculis. Recept. epaleaceum, convexum scrobiculatum. Invol. floribus disci subæquale, squamis 1-2-seriatis distinctis. Anthæræ basi muticæ. Stigmata 2 gracilia divergentia. Achænium verticaliter compressum cuneatum coronatum paleis 2 auriculæformibus aristisque 1-3-glochidiatis. — Herbæ Novo-Hollandicæ ramosæ pilosiusculæ. Folia alterna sessilia inciso-dentata. Capitula ramos apice aphyllis terminantia solitaria. Discus flavus, Radius cœruleus.

1. *C. CUNEIFOLIA* (Brown in bot. reg. t. 504), glabriuscula, foliis cuneatis apice inciso-dentatis, invol. biseriali circ. 15-16-phylo, aristis per totam ferè longitudinem retrorsum serratis. — in Novâ-Hollandiâ interiore ad ripas fluminis Lachlan Lodd. bot. cab. t. 825.
2. *C. DENTEX* (Brown in bot. reg. t. 504 in obs.), glabriuscula, foliis oblongo-lanceolatis utrinquè incisuris 2-3 profundis acutis notatis, invol. subuniseriali 8-9-phylo, aristis apice summo glochidiatis. — in Novâ-Hollandiâ circa Port-Jackson. (v. s. ex hort. kew.)
3. *C. DILATATA* (Cunn. l. in litt. 1834), tota pubescens, foliis latè subrhomboideis grossè dentatis basi cuneatis in petiolum basi obtusè auriculatum attenuatis, involuero subbiseriali, aristis lævibus. \mathcal{U} in collibus lapideis Novæ-Hollandiæ interioris ad Peel's rage, Exley, etc. junio 1817 florent. legit cl. A. Cunningham. Folia pollicem lata. (v. s. comm. à cl. invent.)

CLIII. ASTEROMÆA Blum. bijdr. p. 901. Nees ast. 232. Less. syn. 191. — *Asteris* sp. Linn.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci hermaphr. tubulosis 5-dentatis. Recept. convexum alveolatum. Invol. biseriale, squamis subæqualibus margine abra-

(1) *SOMMERFELTIA* Schum. = *Drepanocarpus*. Conf. Prod. 2. p. 420.

naceis apice herbaceis. Stigm. disci apice lanceolata. Achænium 4-costatum compressiusculum basi attenuatum subpuberulo-glandulosum. Pappus paleaceus 1-serialis, paleis brevibus basi subcoalitis apice setaceo-multifidis. — Herba Asiatica erecta glabriuscula. Folia alterna ovali-oblonga utrinquè acuminata sessilia integra aut hinc inde serrata ramea obovata. Capitula ad apices ramorum solitaria. Ligulæ cœruleæ aut albæ, discus luteus. — Habitus Asteris.

1. *A. INDICA* (Blum. l. c.). — in Indiâ or.? (Linn.), in imp. Chinensi? (Nees), culta in hortis Indicis. *Aster indicus* Linn. sp. 1230. Burm. fl. ind. 181. Willd. sp. 3. p. 2041 excl. descr. et car. ex Nees ast. 233. *Aster Javanensis* Garc. herb. ex Burm. *Callistemma indicum* G. Don in Lond. hort. brit. t. 348. (v. s.)

CLIV. *BELLIUM* Linn. mant. p. 157. DC. fl. fr. 4. p. 923. Cass. dict. 4. suppl. p. 41. et 37. p. 454. Less. syn. 191. — *Bellidis* sp. Tourn. — *Pectidis* sp. Schreb.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci hermaphr. tubulosis 4- (rariùs-5) dentatis. Rec. ovoideo-conicum nudum. Invol. 1-2-seriale, squamis oblongis. Anth. inclusæ. Stigm. subforficata. Achænium compressum obovale subpuberulum. Pappus squamis 4-5 membranaceis truncatis et setis totidem cum squamis alternantibus constans. — Herbæ in Mediterraneis regionibus indigenæ annuæ aut perennes. Folia subradicalia obovata integra aut rariùs dentata. Pedunculi scaposi 1-cephali. Capit. discus luteus, radius albus. — Habitus *Bellidis*.

1. *B. BELLIDIODES* (Linn. mant. 285. excl. syn. Triumf.), caule stolonifero, foliis radicalibus spathulatis integris basi longè attenuatis, scapis folia ferè triplo superantibus. ① in rupestribus humidis maritimis Corsicæ!, Sardinia (Moris), ins. Balearicarum (Camb. bal. 104), Therasiæ (D'Urv.), Chalcì et Rhodi (Sibth.) Viv. fragm. f. ital. 1. p. 8. t. 10. f. 1. Lam. ill. t. 684. DC. fl. fr. 4. p. 923. non Desf. *Bellis droseræfolia* Gon. ill. 69. — Bocc. mus. t. 107. (v. v. et s.)
2. *B. NIVALE* (Requien! in ann. sc. nat. 5. p. 383), caule non stolonifero, foliis radicalibus spathulatis integris breviter attenuatis, scapo apice villosa, foliis paulò longioribus aut iis subæqualibus. ①? in editoribus montibus Corsicæ legerunt cl. Requien et Soleirol, et in Sardinia cl. Badaro. *Herbula pollicaris acaulis*. An ex cl. Duby (bot. gall. 1. p. 266) à priori satis differt? (v. s.)
3. *B. CRASSIFOLIUM* (Moris! pl. sard. elench. 1. p. 26), caule non stolonifero basi multicipiti subadscendente, foliis subradicalibus crassis obovatis integerrimis basi longè attenuatis subpubescentibus, scapis folia triplo quadruplo superantibus pubescentibus. ② in rupestribus maritimis Sardinia ad S. Eliam et Sancti-Petri ad Cala-Vinagra legit cl. Moris. Species distinctissima habitu *Bellidis* perennis. (v. s.)
4. *B. ROTUNDIFOLIUM*, caule non stolonifero, foliis radicalibus obovato-subrotundis crenato-dentatis in petiolum longum desinentibus pubescentibus, scapo foliis triplo quadruplo longiore pubescente. ② in Atlante. *Doronicum rotundifolium* Desf. fl. atl. 2. p. 279. t. 235. f. 1. *Arnica rotundifolia* Willd. sp. 3. p. 2110. *Bellium giganteum* Cass. dict. 37. p. 456. Habitus ferè *Bellidiæstri* sed pappus paleis 5 setisque totidem alternantibus ut in *Bellio*.
5. *B. MINUTUM* (Linn. mant. 286), caule folioso capillari adscendente lævi, foliis obovatis integerrimis petiolatis. ① ad Orientis fontes. *Pectis minuta* Linn. sp. 2. p. 1250. Schreb. nov. act. ups. 1. p. 81. t. 5. f. 2. *Bellis cretica fontana* omnium minima Tourn. cor. 37.

† *Species exclusæ.*B. ANNUUM Less. = *Bellis annua*?B. DENTATUM Viv. = *Bellis dentata*.

Subdiv. 6. BELLIDÆ.

Radius 1- aut rarissime 2-serialis. Pappus nullus aut coroniformis.

CLV. BELLIS. Linn. gen. 962. Gaertn. fr. 2. t. 168. Cass. dict. 37. p. 493. Less. syn. 192. — *Bellis* et *Kyberia* Neck. elem. n. 80 et 81. — *Bellidis* sp. Tourn. Willd.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci tubulosi hermaphrod. 4-5-dentatis. Recept. conicum. Invol. campanulatum 1-2-seriale, squamis obtusis foliaceis æqualibus. Achænia compressa obovata hispidula omnino epapposa. Herbæ Europæ acaules caulescentesve. Folia radicalia aut alterna oblonga aut obovata sæpius dentata in petiolum attenuata. Pedunculi axillares aut scapiformes 1-cephali. Discus luteus. Ligulæ albæ extis subroseæ aut subviolaceæ ante explicationem capituli in globum conniventes.

* *Species caulescentes verisim. annuæ.* — *Kyberia* Neck. elem. n. 81.

1. B. DENTATA, caule sæpius basi ramoso diffuso hispidulo, foliis obovatis grossè dentatis basi cuneatis, invol. squamis vix acutis sæpè pilosis, corollis omnibus ad basin tubi pilorum articulatorum fasciculo barbatis. (1)? in humidis Italiæ ad Pisas (Savil), Veronam (Reich.), Neapolim (Guss.), Siciliæ (Guss.), Melitæ (Beaupré!), Mili (Beaup.), Leucadæ (Paroll), Constantinopolis (Castagne!), Barbariæ (Desf.). *Bellium dentatum* Viv. l. fragm. p. 8. t. 10. f. 2. (excl. syn. plerisq. et descr. fruct.). *Bellium bellidioides* Desf. l. atl. 2. p. 279. non Linn. — *Bellis minima annua* Triumf. obs. p. 80. ic. Simillima sequenti sed corollâ imâ basi barbigerâ undè incautè ad *Bellium* hinc indè relata. (v. s.)
2. B. ANNUA (Linn. sp. 1249), caule sæpè à basi ramoso diffuso hispidulo, foliis obovatis integris subdentatis basi in petiolum attenuatis, invol. squamis vix acutis sæpè pilosis, corollis basi nudis. (1)? in pratis humidis Occitanie, Galloprov., ins. Balearicarum (Camb.), Melitæ (D'Urv.), Corsicæ (Thom.), Teneriffæ (Brouss.), Barbariæ (Desf.), Orientis, Italiæ, Hispaniæ (Willd.). DC. fl. fr. n. 3220. — Bocc. mus. 2. t. 35. B. ramosa et B. repens Lam. fl. fr. 2. p. 122. *Bellium annuum* Less. syn. 192? Habitus varius: acaulis, erecta aut sæpius diffusa. (v. v.)

β *minuta*, caule basi multiplici, scapis erectis folio quadruplò longioribus. — circa Byzantium locis humidis legit cl. L. Castagne et in Græciâ (h. acad. Monac!). Habitus omnino *Bellii bellidioidis* sed achænia calva. (v. s.)

3. B. INTEGRIFOLIA (Michx. fl. bor. am. 2. p. 131. non Lam.), caulescens divaricato-ramosa, foliis integerrimis longè ciliatis inferioribus obovatis, super. lanceolatis, invol. squamis glabriusculis lævibus acuminatissimis et quasi in aristam desinentibus. (1) ad ripas rivulorum et in collibus umbrosis Tennessee (Michx.), in Kentucky occidentali (Raf.) in agro Arkansano (Nutt.). Radix exilis. *Eclipta integrifolia* Spreng. syst. 3. p. 602. ex syn. (v. s.)

** *Species subacaules perennes.* — *Bellis* Neck. elem. n. 80.

4. B. PERENNIS (Linn. sp. 1248), foliis subradicalibus spathulato-obovatis crenato-dentatis pubescentibus venosis invol. squamis pilosis vix acutis. 2/ in Europæ pascuis vulgatissima usquè ad Tauriam (Bieb.), etiam in Maderâ sed dicitur introducta (Low.) Lam. ill. t. 677. fl. Dan. t. 502. engl. bot. t. 424.

Schkuhr handb. t. 251. Variat glabriuscula aut hispidula, acaulis aut subcaulescens, etc. Reperitur, in hortis præcipuè, scapis fasciatis, capitulis proliferis (J. Bauh. hist. 3. p. 1. p. 113. f. 2), flosculis omnibus ligulatis vulgò (bot. mag. t. 228), et rariùs flosc. omnibus tubulosis ampliatis (Blackw. t. 530), ligulis albis aut purpurascensibus, etc. (v. v.)

β. integrifolia (DC. fl. fr. ed. 3. v. 4. p. 185. var. s.), caulescens elongata, foliis integerrimis, capitulis minoribus. — *Bellis integrifolia* Lam. ! dict. 5. p. 6. non Michx. diù quasi spontanea in colle horti Parisiensis. (v. s.)

5. *B. sylvestris* (Cyr. pl. rar. 2. p. 22. t. 4), foliis subradicalibus obovatis crenatis trinerviis, invol. squamis pilosis vix acutis. $\frac{1}{2}$ in campis apricis Sicilia (Guss.), agri Neapolitani (Cyr. Ten.), Pisani (Savi!), Nicæensis (C. Maun.), Galloprov. (Rob.), Avenionensis (Req.), in Istriâ (Reich.), in Apenninis Appuanis (Bertol.), Sardinia (Moris!), agro Byzantino (Cast.), Smyrnæ (Parol.), et in Mauritaniâ Mogadoriensi (Brouss.). Pedunculi longiores et capitulum majus quàm *B. perennis*. Invol. squamæ magis foliaceæ et disco paulò longiores. (v. s.)

β. hybrida (Ten. ! syll. p. 436. fl. med. 2. p. 64), caule basi subramoso, foliis elliptico-oblongis magnitud. *B. perennis*, invol. squamis disco subbrevioribus. $\frac{1}{2}$ cum. var. *α*. Omnino media inter *B. perennem* et *B. sylvestrem* et fortè vera hybrida ? (v. s.)

† *Species exclusæ.*

- B. ciliaris* Labill. = *Brachycome ciliaris*.
B. droseræfolia Gou. = *Bellium bellidioides*.
B. graminea Labill. = *Paquerina graminea*.
B. javanica Blum. = *Lagenophora javanica*.
B. magellanica Lam. = *Lagenophora Commersonii*.
B. pilosa Blum. = *Lagenophora pilosa*.
B. spinosa G. alp. = *Plagus ageratifolius*.
B. stipitata Labill. = *Lagenophora Billardieri*.

CLVI. *BRACHYCOME* Cass. dict. 37. p. 464. et 491. — *Brachycomes* sp. Less. syn. 192. — *Brachyscome* Cass. bull. philom. 1816. p. 199. dict. v. 5. suppl. p. 63. — *Bellidis* sp. Labill.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis 1-seriatis, disci tubulosi 5-dentatis hermaphroditis. Recept. conicum subalveolatum epaleaceum. Invol. campanulatum pauciseriale, squamis margine submembranaceis. Achæmium lateraliter plano-compressum erostre pappo subsetiformi brevissimo coronatum. — *Herbæ Novo-Hollandicæ*. Folia alterna glabra parcè dentata pinnatilobata aut trifida. Discus luteus. Radius albus.

1. *B. stricta*, caule strictè erecto folisque pube brevi hirtellis, foliis oblongis basi attenuato-cuneatis grosse dentatis, invol. squamis apice membranaceis obtusissimis, fructu alà integrâ marginato in disco subtuberculato. $\frac{1}{2}$ in Novæ-Hollandiæ terrâ Van-Leuwin. Bellis aculeata Labill. ! Nov. Holl. specim. 3. p. 55. t. 206. Brach. aculeata Less. syn. 192. excl. syn. Sieb. Herba rigidula erecta tota pubero-hirta. Rami apice aphylli. Achænia vix tuberculata nec verè aculeata. (v. s. comm. à cl. Labill.)
2. *B. heterodoxa*, glabriuscula, caule adscendente aut erecto, foliis radicalibus petiolatis pinnatifidis, caulinis sessilibus oblongis apice grosse et acutè trilobulatis basi subdenticulatis medio integerrimis, invol. squamis apice membranaceis acutis, fructu alà dentatâ marginato in disco tuberculato. $\frac{1}{2}$ in planis humidis Novæ-Hollandiæ interioris circa Bathurst et ad Lachlan river legit cl. A.

Cunningham. Br. Billardieri Cass. dict. 5 suppl. p. 63. excl. syn. Folia forma et incisuris valde varians. Pappus setis longioribus et confertioribus quam cæter. specierum. (v. s. comm. à cl. invent.)

3. B. *SCAPIFORMIS*, caule erecto 1-cephalo apice nudo glabro sub capitulo confertè et brevè velutino, foliis radicalibus spathulatis obtusis crenatis in petiolum attenuatis, caulinis sessilibus oblongis linearibus, infer. apice serratis, cæteris integerrimis, invol. squamis margine membranaceis acutis, achænio obovato integro et tuberculato. 2½.

α. *puberula*, foliis pube brevi scabridis. — in Smith's Plains ad Lachlan river in Novæ-Hollandiæ inter. junio flor. legit cl. Cunningham. (v. s. comm. à cl. inv.)

β. *glabra*, foliis glabris. — in insulâ Van-Diemen legit cl. Gunn. (v. s. comm. à cl. Lindl.)

4. B. *SIEBERI*, glabra adscendens, foliis oblongis basi attenuatis acutè paucidentatis, invol. squamis subobtusis vix membranaceis, achænio obovato margine callosò integro. — in Novæ-Hollandiæ, Sieb. coll. exs. n. 485. B. aculeata Less. syn. 192. excl. syn. Labill. — Priori affinis sed differt glabritie totâ, caule adscendente debili gracili, capitalis dimidiò minoribus, invol. squamis angustis vix obtusis. Achænia obovato-compressa puberula margine callosa integra. (v. s.)

β. *Gunnii*, herba tota brevissimè subpuberula. 2½ Cl. Gunn legit in insulâ Van-Diemen. (v. s. comm. à cl. Lindl.)

5. B. *CILIARIS* (Less. syn. 192), glabra, foliis pinnatipartitis, lobis linearibus distantibus integerrimis, squamis invol. obtusis apice membranaceis, achæniis margine ciliatis. — in Novæ-Hollandiæ terrâ Van-Leuwin (Labill.) ad oram orientalem (Mus. Par.). Bellis ciliaris Labill. Nov. Holl. specim. 2. p. 56. t. 207. In specim. juniore achænia ciliata non vidi et ideò species mihi dubia. (v. s. comm. à Mus. reg. Par.)

6. B. *MULTIFIDA*, erecta glaberrima, foliis pinnatipartitis, lobis lineari-subulatis integerrimis acutis, invol. squamis obtusis, achænio oblongo immarginato utrinquè tuberculato. 2½ in umbrosis rupestribus Novæ-Hollandiæ int. ad Peel's Range junio flor. legit cl. A. Cunningham. Valdè accedit ad iconem 207 Labill. sed achænia certissimè non sunt ciliata, lobi foliorum tenuiores acutiores. (v. s. comm. à cl. invent.)

7. B. *ANGUSTIFOLIA* (Cunn. ! in litt. 1834), glabra adscendens aut erecta, foliis oblongo-linearibus integerrimis 1-nerviis, nervo valido, invol. squamis acutis, achænio integerrimo tuberculato, pappo brevissimo. 2½ in siccis glareosis apricis Novæ-Hollandiæ ad Goulburn Plains, maio flor. legit cl. Cunningham, et in ins. Van-Diemen cl. Gunn. Folia rigidula nervo prominulo a poll. longa, 2 lin. lata. Pappus minutissimus. (v. s. comm. à cl. Cunn. et Lindl.)

8. B. *LINEARIFOLIA*, glabra, foliis linearibus integerrimis indivisis. — in Novæ-Hollandiæ. Omnino refert figuram sinistram tab. 204. sp. fl. Nov. Holl. Labill. sed achænia certè pappo prioribus simili coronata nec nuda. (v. s. comm. à cl. Lambert. et Thibaud.)

†. *Species exclusæ.*

B. *DENTATA* Gaudich. = Vittadinia dentata.

B. *SPATHULATA* Gaudich. = Vittadinia spathulata.

B. *TRILOBA* Gaudich. = Vittadinia triloba.

B. *XANTHOCOMOIDES* Less. = Keerlia linearifolia.

CLVII. *PAQUERINA* Cass. dict. 37. p. 464 et 492. Less. syn. 192 — Bellidis sp. Labill.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœminicis 1-seriatis, disci-hermaphroditis tubulosis 5-dentatis. Recept. conicum

breviter paleaceo-fimbrilliferum. Invol. campanulati squamæ subuni-seriales. Achænium erostre epapposum plano-compressum margine crassiusculum è scrobiculis recept. oriundum. — Herba Novo-Hollandica glabra gracilis, tota foliata, simplex aut ramosa, ramis 1-cephalis. Folia alterna lineari-oblonga integerrima basi attenuata.

1. *P. GRAMINEA* (Cass. l. c.). — in Novâ-Hollandiâ ad prom. Van-Diemen (Labill.) et ad fretum d'Entrecasteaux. Bellis graminea Labill. Nov. Holl. sp. 2. p. 54. t. 204. excl. fortè fig. sinistra. (v. s. comm. à Mus. reg. Par.)

CLVIII. *LAGENOPHORA* Cass. bull. philom. mart. 1818. Less. syn. 193. — *Lagenifera* Cass. bull. philom. 1816. decemb. — *Microcalia* A. Rich. — *Bellidis* sp. DC. — *Asteris* sp. Lam. — *Calendulæ* sp. Forst.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis 1 seriatim, disci tubulosis 5-dentatis hermaphroditis masculive sterilibus; antheræ subliberæ. Recept. planum nudum. Invol. disco subæquale, squamis circ. biseriatis acutis basi coriaceis adpressis, apice submembranaceis. Achænia radii plano-compressa oblonga epapposa rostrata, disci abortiva. — Herbæ graciles, ad collum folia pauca propè basin gerentes, scapo squamulato aphylo 1-cephalo. Folia basi attenuata aut petiolata. Flores disci lutei, radii purpurei.

1. *L. FORSTERI*, foliis glabris obovato-orbiculatis acutè dentatis petiolatis, petiolis ciliatis. $\frac{1}{2}$ in Novâ-Zeelandiâ (Forst.) ad flumen Hawa-Hawa (D'Urv. 1), loco gallicè dicto Havre de l'Astrolabe (Lesson!). *Calendula pumila* Forst. prod. 305. Willd. sp. 3. p. 2344. *Microcalia australis* A. Rich. fl. Nov. Zeel. p. 231. yoy. astr. 30. Petiolus limbo paulò longior, pilis articulatis rigidulis basi præsertim instructus. Limbi dentes infimi profundiores. Capitulum majus quàm in sequente. (v. s. comm. à cl. D'Urville, Lesson et Rich.)

2. *L. COMMERSONII* (Cass. dict. 25. p. 110), foliis glabris obovatis obtusis integerrimis aut apice pauci-dentatis in petiolum brevem attenuatis. \odot ? ad fretum Magellanicum (Comm.!), in Staten-Island (Webster. in h. Lindl.), in summis Andibus Chilensibus (C. Gay!), et in insulâ Tristan d'Acunhâ (Pet. Th.). *Aster nudicanlis* Comm. herb., Lam. dict. 1. p. 308. ill. t. 681. f. 4. *Calendula pumila* var. β . Forst. prod. 305. *Calendula Magellanica* Willd. sp. 3. p. 2344. *Calendula pusilla* Pet. Th. flor. Trist. d'Ac. p. 40. t. 9. *Bellis Magellanica* DC. in Lam. dict. enc. 5. p. 7. (v. s.)

3. *L. BILLARDIERII* (Cass. dict. 25 p. 111), foliis utrinquè hirtellis oblongis basi in petiolum brevem attenuatis sinuato-dentatis. $\frac{1}{2}$? in Novâ-Hollandiâ ad prom. Van-Diemen et alibi. Fortè species plures hic commixtæ nempè:

α . *pusilla*, foliis utrinquè hirsutis vix basi attenuatis, scapo sesquipollicari, invol. squamis exterioribus lineari-subulatis subpatulis. — Specim. habeo à cl. Labillardière sub nom. *Bellidis stipitata*. (v. s.)

β . *media*, foliis utrinquè sparse hirtellis basi sensim attenuatis, scapo 4-5-pollic., invol. squamis linearibus subacutis subadpressis. *Bellis stipitata* Labill. Nov. Holl. spec. 55. t. 205. — Habui ex coll. Sieb. n. 505. ex herb. Lamb. et cult. in hort. Kew. (v. s.)

γ . *glabrata*, foliis utrinquè glabriusculis basi sensim attenuatis et verè petiolatis, scapo 4-6 pollic., invol. squamis erecto-adpressis. (v. s. comm. à cl. Lambert.)

4. *L. HIRSUTA* (Poepp. ex Less. in. linnæâ 1831. p. 131), foliis utrinquè hirsutis petiolatis obovatis grossè dentatis, dentibus macronulatis, scapo vix foliis duplò longiore. $\frac{1}{2}$ in Chili Poepp. coll. exs. n. 715. Folia cum petiolo 9-10 lin. ferè bipollicaria, 9-10 lin. lata membranacea. Invol. squamæ erectæ acuminatæ disco paulò longiores. Achæni rostrum glanduloso-viscosum. (v. s.)

CLIX. IXAUCHENUS *Cass. dict.* 56. p. 176. — *Ixiauchenus Less. syn.* 193.

Capitulum multiflorum heterogamum, fl. radii biseriatis foemineis ligulatis longissimis, disci tubulosi 5-dentatis hermaphroditis aut abortu masculis. Recept. nudum latum planum. Invol. subhemisphæricum disco æquale, squamis pluriseriatis laxè adpressis foliaceis. Achænia ovali-oblonga compressa glabra in collum brevem glutinosum attenuata epapposa. — Herba Novæ-Hollandiæ?, glabra basi tantum foliata. Caulis scapiformis aphyllus 1-cephalus. Folia alterna subradicalia valdè inæqualia semiamplexicaulia ferè lyrata. Discus flavus. Radius albus. An genus à *Lagenophorâ* aut à *Myriactide* satis diversum?

1. *I. SUBLYRATUS* (Cass. l. c.). — in Novâ Hollandiâ propè Port-Jakson? *Ixiauchenus lyratus Less. l. c.*

CLX. MYRIACTIS *Less. in linnæâ* 1831. p. 127. *syn.* 193. — *Laveniæ sp. Wall. list.* — *Balsamitæ sp. Wall. herb.* 1821.

Capitulum multiflorum heterogamum, fl. radii bi- aut multiserialibus foemineis angustè ligulatis integerrimis, fl. disci hermaphr. tubulosi 5-lobis. Recept. nudum. Invol. 1-2-seriale, squamis linearibus acutis. Achænia compresso-plana calva glabra erostria apice sæpè glandulifera. — Herba Indiæ aut Persiæ erectæ subdichotomo-ramosæ. Folia alterna ovata aut lanceolata dentato-serrata, suprà pube minimâ scabrida. Pedunculi elongati 1-cephali subpaniculatim dispositi. Capitula globosa albida aut flavida.

§. 1. *Ligulis circiter biseriatis, achænio apice vix subattenuato. — Species Indiæ australioris aut insularis, mediæ inter Lagenophoram et Myriactidem veram.*

1. *M. WIGHTII* (DC. in Wight contrib. bot. ind. p. 10), parcè pilosa, foliis inferioribus ovatis basi longè cuneato-attenuatis grossè inciso-dentatis, superioribus oblongis integris sessilibus dentium et folii ipsius apicibus calloso-mucronatis. ½ in Indiâ orientali ad montes Neelgherry legit cl. Wight. *Carpesium nov. sp. Wight! herb.* Folia radicalia in petiolum attenuata ovata, canliu. infer. basi cuneata parcè dentata, summa subsessilia integra utrinquè acuminata. Capitula terminalia solitaria 4 lin. diam. Invol. hirsutiusculum post. anthesin reflexum. Ligulæ albide subbiseriales per exsic. revolutæ. (v. s. in h. Wight.)

2. *M. JAVANICA*, caule erecto aut adscendente glabriusculo apice ramoso, foliis ovalibus in petiolum attenuatis distanter et calloso-serratis sparsè pilosis ciliatis. ½ in insulâ Javæ sylvis altioribus montis Gede legit cl. Blume. *Bellis Javanica Blum. l. bijdr.* p. 917. *Lagenophora Javanica Blum. l. in litt.* 1831. Herba sesquipedalis. Rami subaphylli. Capitula ferè prioris. Ligulæ plurimæ. Recept. convexum. (v. s. comm. à cl. Blume.)

3. *M. PILIOSA* (Blum. l. in litt. 1831), tota pilosa, caule suffruticuloso adscendente, foliis ovato-lanceolatis acuminatis profundè dentatis, racemo terminali paucifloro. — in summo montis Gede ins. Javæ. *Bellis pilosa Blum. l. bijdr.* 917. *Lagenophora pilosa Blum. l. in litt.* 1831. Capitula prioris sed minora. Recept. minus elevatum. (v. s. sine flor. comm. à cl. Blume.)

§. 2. *Ligulis pluriserialibus, achænio apice omninò erostrato. — Species Nepalenses.*

4. *M. NEPALENSIS* (Less. ll. cc. in Lin. t. 2. f. F. 1), caule glabro, involucreo biseriali capitulo multò breviori, receptaculo convexo ad centrum subconico.

— in Nepaliâ. *Lavenia sphærantha* Wall.! cat. et herb. n. 327. Capitula globosa Pisi magnitudine. (v. s.)

5. *M. WALLICHII* (Less. ll. cc.), caule hirsutiusculo, involucrio subniseriali capitulo subæquali, receptaculo plano. — in montibus excelsis Nepaliæ borealis. Balsamita dentata Wall.! cat. n. 326. Capitula dimidiò quàm prioris minora. (v. s.)

ß. *hispidioides*, caule et foliis utrinquè pilis sparsis hispidissimo. — in montibus Indiæ bor. occid. legit cl. Royle. (v. s.)

6. *M. GMELINI*, erecta pilis articulatis setulisque parçè hirsuta, foliis petiolatis membranaceis grossè dentatis, invol. squamis 2-3-seriatis lineari-oblongis ciliato-fimbriatis, achæniis basi glandulosis in collum globulo glanduloso terminatum productis. 2½ in Persiâ prov. Gilan legit Gmelin jun. et in sylvis montium Talusch rev. Hohenhaker. Botriyadenia Gmelini Fisch. et Mey. in litt. 1835. Folia ferè *Callistephi Chinensis*. Habitus *Dicrocephalæ latifoliæ*. Descr. ex F. et M. litt. (v. s. capitulum à cl. auct. miss.)

CLXI. *GARULEUM* Cass. dict. 18. p. 162. 30. p. 324, Less. syn. 194. excl. syn. Tourn. — *Osteospermum* sp. Ait. *L'Hér. Thunb.*

Capitula radiata, fl. radii longè ligulatis fœmineis, disci tubulosis 5-dentatis masculis, ovulis abortivis. Recept. convexum nudum. Invol. squamæ biseriales. Stylus disci ramis apice liberis divergentibus, extùs piliferis, intùs ad margines glandulosis. Ovaria disci oblonga. Achænia radii obovato-oblonga 3-4-costata erostrata epapposa sicco-coriacea, pericarpio tenui extùs rugoso et subaspero, disci plano-compressa lævia. — Herbæ suffrutescentes Capenses erectæ glabræ aut subviscido-pubescentes. Folia alterna pinnatisecta, lobis plus minùs dentatis. Capitula terminalia pedunculata, radio cœruleo, disco luteo, corollis basi puberulis.

1. *G. PINNATIFIDUM*, foliis pinnatifidis, lobis ovatis dentatis. ½ ad Cap. Bonæ-Spei (Th.) ad Sneurobergen alt. 4-500 ped. (Dreg.!). *Osteospermum pinnatifidum* Thunb. prod. 167. fl. cap. 717. *L'Hér.*! st. nov. 11. t. 6. Ost. cœruleum Alt. kew. 3. p. 275. Jacq. ic. rar. 1. t. 179. coll. 1. p. 78. *Garuleum viscosum* Cass. dict. 18. p. 325. *Chrysanthemoides cœrulea* Moench. meth. 584. — Zeyh.! pl. exs. cap. n. 102. in h. Moric. Herbæ odor gravis. (v. v. c.)
2. *G. BIPINNATUM* (Less. l. c.), foliis bipinnatisectis, lobis lineari-setaceis acutè lobulatis aut integris. ½ ad Cap. Bonæ-Spei, in Hantum (Th.), in Carro (Dreg.! Eckl.!), ad Zondag rivier et Graaf rivier (Dreg.!). Achænia radii trigona irregulariter transversè sulcata, disci inania plano-compressa lævia. *Osteospermum bipinnatum* Thunb. fl. cap. 717. (v. s.)

CLXII. *KEERLIA*. — *Brachycomes* sp. Less.

Capitulum multiflorum radiatum, ligulis circ. 15-25 fœmineis 1-2-serialibus linearibus, fl. disci hermaphr. aut abortu masculis 5-dentatis brevibus. Invol. latè campanulatum imbricatum pluriseriale, squamis lanceolatis acuminatis margine submembranaceis. Recept. planum nudum. Styli radii breves obtusi, disci erecti dorso subpuberuli. Achænia glabra obpyramidata seu cylindræa sæpiùs longitudinaliter sulcata. Pappus paleis brevissimis pluribus acutis basi subconcretis et ideò ferè coronam minutissimam conficientibus. — Herbæ Mexicanæ glabræ aut pube brevi velutino-villosulæ erectæ parçè ramosæ. Folia alterna oblonga aut cuneata sessilia seu semiamplexicaulia integra aut apice dentata. Rami apice nudi 1-cephali. Flores albi aut flavi. — Genus à *Brachycomi* (confirmante habitationum diversitate) recept. plano et achæniis teretiusculis nec lateraliter plano-compressis distinctum, dicavi cl.

§. 1. *Foliis alternis.*

1. *G. GLUTINOSUM* (Less. l. c.), suffruticosum, ramis teretibus, foliis lineari-lanceolatis integerrimis trinerviis glutinosis, paniculâ fasciculatâ, ligulis minimis, disco sterili. $\bar{\zeta}$ in Brasiliâ. *Selloa glutinosa* Spreng. pug. p. 36. n. 35. nov. prov. 36. syst. 3. p. 496. Ker. bot. reg. t. 462. *Molina viscosa* hort. berol. ex Spreng.
2. *G. CORYMBOSUM*, fruticosum, ramulis subangulatis dichotomo-corymbosis, foliis oblongis utrinquè attenuatis subviscosis trinerviis, venis laterilibus tenuissimis, capitulis ad apicem ramuli cujusque subternis 8-floris, ligulis 5, disco circ. 3-floro. $\bar{\zeta}$ in Mexico circa Matamoros legit cl. Berlandier (pl. exs. n. 2266). Folia 12-15 lin. longa, 2 lin. lata. (v. s.)
3. *G. MULTIFLORUM*, fruticosum, glutinosum, ramulis angulatis, foliis linearibus acutis trinerviis venis laterilibus tenuissimis, capitulis ad apices ramulorum 3-5 confertissimè corymbosis 13-14-floris, ligulis 5, fl. disci 7-8. $\bar{\zeta}$ in Mexico (Alam.! Maire!) circa urbem (Berl.! pl. exs. n. 50). Flor. septemb. *Molina glutinosa* Alam.! in litt. Priori affine. Folia 2 poll. longa, $1\frac{1}{2}$ -2 lin. lata. (v. s.)
4. *G. SCOPARIUM*, fruticosum subviscosum ramosissimum, ramis erectis confertis apice paniculato-corymbosis, foliis oblongo-linearibus acutis trinerviis, venis laterilibus tenuibus, capitulis ad apices ramulorum plurimis confertis circ. 10-floris, ligulis et tubulis 5. $\bar{\zeta}$ in Mexicanæ confederationis statu Sancti-Ludovici de Potosi legit cl. Berlandier (pl. exs. n. 1338). Fol. 9-10 lin. longa, $1\frac{1}{2}$ -2 lata. (v. s.)

§. 2. *Foliis oppositis.*

5. *G. NUDATUM*, caule herbaceo lævi angulari, foliis oppositis crassis integerrimis linearibus, capitulis in corymbum compositum trichotomum dispositis, bracteis filiformibus, tubulis 5-8, ligulis subnullis. \mathcal{U} in Floridâ orientali. *Selloa nudata* Nutt. in Sillim amer. journ. 5. 1822. p. 300. Affinis dicitur *Chrysocome* et potius ut videtur *Bigelowiæ*.
6. *G. OPPOSITIFOLIUM*, caule herbaceo lævi tereti, foliis oppositis linearibus submembranaceis integerrimis aut remotè denticulatis, capitulis in corymbum compositum trichotomum dispositis, bracteis lineari-filiformibus, floribus 10 omnibus tubulosis. — in Mexico inter San-Ferrado et Santander legit cl. Berlandier (pl. exs. n. 2263!). Herba erecta 1-1 $\frac{1}{2}$ -pedalis. Folia 3-4 poll. longa, 3 lin. lata. Corymbus polycephalus flavus. Habitus omninò sect. 1, sed folia opposita et capitula discoides. (v. s.)

Subdiv. 2. *ACHYRYDÆ.*

Pappus paleaceus multiradiatus (ut in *Belliis*).

CLXVIII. *BRACHYRIS* Nutt. *gen. am.* 2. p. 163. Less. *syn.* 191. — *Brachyachyris* Spreng. *syst.* 3. p. 574.

Capitulum pluriflorum radiatum, fl. radii 5-10 ligulatis 1-serialibus fœmineis, disci 3-10 tubulosis 5-dentatis hermaphroditis. Recept. nudum. Invol. ovati aut cylindranei squamæ arcuè imbricatæ. Achænia obconica erostria. Pappus paleis 5-8 oblongis persistentibus. — Herbæ Occident.-Americanae rhizocarpicae aut suffruticosae glabræ erectæ. Folia alterna linearia seu lanceolata integra, sæpè punctata. Capitula solitaria aut terna ad apices ramulorum corymbosa. Invol. squamæ pallidè apice maculâ viridi notatæ. Cor. luteæ. — Habitus hinc *Euthamiae*, illinc *Bigelowiæ* sed pappo paleaceo nec piloso distincta, ex *ἄραχυνς*, brevis, et *ἀχυρίς*, palea.

Sect. 1. *EUBRACHYRIS.*

Achænia radii pappo fl. disci simili donata. Fl. disci numero ligularum æquales aut pauciores.

* *Suffruticosa.*

1. *B. MICROCEPHALA*, caule suffruticuloso corymboso ramoso tereti vix striatulo, foliis angustissimè linearibus patulis margine subrevolutis, corymbo fastigiato, fl. radii 5 invol. paulò longioribus, fl. disci 1-3. $\frac{1}{2}$ in Mexico prope locum dictum *Sallito* legit cl. Berlandier januar. florent. Invol. parva oblonga glutinosa. Ligulæ in sicco revolutæ. Folia pollicem longa vix semilin. lata. (v. s.)
2. *B. PANICULATA*, caule suffruticoso paniculato ramoso teretiusculo, foliis linearibus acutis punctatis planis crassiusculis, capitulis oblongis in paniculam dispositis, fl. radii 5 invol. paulò longioribus, disci 5-6. $\frac{1}{2}$ in Chili circa Coquimbo legit cl. Gaudichaud et circa Acapulco cl. Hænke (ex h. Sternb.). Etiam habui sine loci designatione olim à cl. Née lectam. Folia 7-9 lin. longa, ferè lineam lata. (v. s.)

** *Herbacea.*

3. *B. EUTHAMIA* (Nutt. gen. am. 2. p. 163), caulibus numerosis marginato-angulatis et scabris, foliis adproximatis angustè linearibus integris punctatis scabris, corymbo dichotomo fastigiato, fl. radii 5 invol. circ. longitudine, fl. disci 5. $\frac{1}{2}$ in collibus aridis ad fl. Missouri, ab Arikarees ad montes. *Solidago Sarothræ* Pursh fl. bor. am. 2. p. 540. *Brachyachyris Euthamiæ* Spreng. l. c. Herba balsamica subfœtida strenuè diuretica.
4. *B. CALIFORNICA*, caulibus teretibus apice subpaniculatis, foliis linearibus acutis basi attenuatis et subciliatis planis crassiusculis, paniculâ racemosâ oligocephalâ, capitulis ovatis, ligulis circ. 8-10 invol. paulò longioribus, fl. disci 8-10. $\frac{1}{2}$ in California legit cl. Douglas. Folia pollicem longa, vix lineam lata. (v. s. comm. ab hon. soc. hortic. Londin.)
5. *B. NEERANA*, rhizomate lignoso, caulibus pluribus subangulatis apice mono-aut oligocephalis, foliis linearibus acutis crassiusculis viscosis, infer. oblongo-linearibus, capitulis obovatis ligulis 8-10 disco duplò ferè longioribus, fl. disci 10-12. $\frac{1}{2}$ habeo ex herb. Neei sine loci natalis design. verisim. ex Mexico aut ex Chili. Invol. squamæ minùs quàm in prioribus adpressæ. Herba vix digiti longit. superans. Folia pollicem longa, 1-2 lin. lata. (v. s.)
6. *B. OVATIFOLIA*, herbacea, caule erecto tereti, foliis ovatis acutis petiolatis argutè serratis, capitulis subsessilibus glomeratis in racemum elongatum subaphyllum simplicem unilateralem dispositis. $\frac{1}{2}$ in agro Kentuckensi ad ripas fluminum legit cl. Rafinesque. *Solidago sphacelata* Raf. ! a. n. 1820 ex specim. miss. 1834. Species distinctissima ! Capit. oblongum squamæ inv. subsphacelatæ maculâ viridi non notatæ. Ligulæ 5. Fl. disci circ. 4-5. Pappus paleis 5-6. (v. s.)

Sect. II. AMPHIACHYRIS DC. not. VII.

Achenia radii ferè calva aut coronâ brevissimâ donata. Flor. disci ligulis numerosiores.

7. *B. DRACUNCULOIDES* (DC. not. VII. jard. gen. p. 1. t. 1), caulibus erectis ramosis teretibus, foliis lanceolato-linearibus punctatis uninerviis aut basi subternerviis margine scabridis, corymbo laxo, ramis oligocephalis, invol. obovato foliis parvis bracteato, ligulis 10-11 ovali-oblongis invol. longioribus, fl. disci circ. 20. $\frac{1}{2}$ in territorio Arkansano semina legit cl. de Pourtalès. Herba prioribus major habitu Galatellæ aut *Euthamiæ* graminifoliæ. Folia 3 lin. longa. Recept. alveolatum. (v. v. in hort. Genève. comm. à cl. de Charpentier.)
- β . *angustissima*, foliis linearibus angustissimis uninerviis. $\frac{1}{2}$ in terr. Arkansano etiam semina legit cl. de Pourtalès. Folia lineam lata punctata. Capit. omninò prioris. (v. v. in h. Gen.)

CLXIX. HEMIACHYRIS.

Capitulum pluriflorum, fl. radii circ. 5 ligulatis fœmineis, disci 7-8 tubulosi 5-dentati hermaphr. Invol. ovatum, squamis oblongis adpressè imbricatis. Recept. nudum. Styli fl. disci appendiculati hirtelli.

Achænia obovato-teretiuscula villosiuscula, radii calva, disci squamellis ovatis subacutis brevibus coronata. — Herba suffruticosa Texana erecta ramosissima glabra. Rami juniores sulcato-angulati, adulti teretēs. Folia alterna linearia integerrima uninervia, summa ferē subulata. Capitula parva ramulos terminantia solitaria. Flores lutei. — Affinis *Brachyridi* sed pappi paleæ brevissimæ squamiformes nec lineares. An *Euheleniis* potius referenda? — Nomen alludit ad *achænia* alia calva, alia squamellas gerentia.

1. *H. TEXANA*. \tilde{f} in Mexici prov. Texas legit cl. Berlandier prope Bejar. Flor. aug. (Berl. pl. exs. n. 1763 et 1765). Caulis 1 $\frac{1}{2}$ -2-pedalis parcē foliosus. Folia infer. pollicem longa, 1 $\frac{1}{2}$ lin. lata, summa multò minora. (v. s.)

CLXX. *LEPIDOPHYLLUM* Cass. bull. philom. 1816. p. 199. dict. 26. p. 36. Less. syn. 191. — *Conyzæ* sp. Lam.

Capitulum pluriflorum radiatum, ligulis 2-3 fœmineis brevibus 1-serialibus, disco 4-6-floro hermaphr. regulari. Invol. oblongo-cylindraceum, squamis paucis obtusis imbricatis. Recept. parvum nudum. *Achænia* oblonga striata. Pappus 1-serialis, setis paleaceis confertis inæqualibus margine subfimbriatis. — Frutex glaber ramosus habitu ferē *Pteroniæ*, caract. *Brachyridis*, ex Amer. austr. extremâ ortus. Folia opposita quadrifariam imbricata ovato-subtrigona coriacea parva integerrima. Capitula ramulos terminantia sessilia solitaria. Cor. flavæ.

1. *L. CUPRESSIFORME* (Cass. dict. 26. p. 37). \tilde{f} ad fretum Magellanicum inter oram Patagonicam et sinum Boncaut leg. Commerson, et ad portum Desideratum Née! *Conyza cupressiformis* Lam. dict. 2. p. 91. ill. t. 697. f. 3. *Baccharis cupressiformis* Pers. ench. 2. p. 425. n. 50. *Athanasia*? *cupressiformis* Comm. ined. (v. s.)

CLXXI. *GRINDELIA* Willd. mag. naturforsch. Berl. 1807. p. 261. Dunal in mem. mus. h. n. Paris. 5. p. 48. — *Demetria* Lag. gen. nov. in el. k. madr. 1814. p. 30. — *Donia Brown* in h. kew. ed. 2 (1813) v. 5. p. 82 non Don. — *Grindelia* et *Aurelia* Cass. dict. 37. p. 468. — *Grindelia* et *Donia* Desf. Link. etc. *Grindelia* et *Doniæ* sp. Less. — *Asteris* sp. Cav.

Capitulum multiflorum heterogamum, fl. radii 1-seriatis ligulatis (in unicâ spec. nullis), disci tubulosi 5-dent. hermaphr. Recept. foveolatum. Invol. basi hemisphæricum, squamis pluriserialibus imbricatis. Antheræ basi nudæ nec caudatæ. *Achænia* obovata aut subangulata glabra. Pappus pauciaristatus deciduus, aristis crassis rigidis. — Suffrutices aut Herbæ Americane ad apices et præsertim ad involucria glutiniferæ. Folia alterna serrata aut integra, radicalia sæpius spathulata, caulina sessilia aut semiamplexicaulia. Capitula ad apices ramorum solitaria. Flores flavi. — Genus naturalissimum etiamsi ab auctoribus etiam gravissimis variè vexatum.

1. *G. GLUTINOSA* (Dunal in mem. mus. h. n. Par. 5. p. 49), fruticosa glabra apice glutinosa, foliis sessilibus breviter decurrentibus obovatis obtusis apice serratis punctato-pellucidis, capitulis terminalibus solitariis præsertim ante explicationem valdè glutinosi, pappo 5-8-seto. \tilde{f} in Mexico. *Aster glutinosus* Cav. ic. 2. p. 53. t. 168. *Doronicum glutinosum* Willd. sp. 3. p. 2115. *Inula glutinosa* Pers. ench. 2. p. 452. *Donia glutinosa* Brown h. kew. ed. 2. v. 5. p. 82. *Demetria glutinosa* Lag. gen. in elench. h. madr. 1814. p. 30. *Aurelia decurrens* Cass. dict. 37. p. 468. Pappi setæ rigide triquetre secus angulos acabridæ. (v. v. c.)

2. *G. SQUARROSA* (Dunal l. c. p. 50), herbacea glabra, caule parçè ramoso. ramis apice 1-cephalis, foliis semiamplexicaulibus oblongis serrulatis, invol. squamis glutinosis in acumen filiforme recurvato-squarrosorum productis, pappo 2-4-setoso. ♀ in pratis apricis secus rivos fluminis Missouri (Pursh). *Donia squarrosa* Pursh fl. bor. am. 2. p. 559. bot. mag. t. 1706. *Anrelia amplexicaulis* Cass. dict. 37. p. 468. (v. v. c.)
3. *G. SUBDECURRENS*, suffruticulosa glabra parçè ramosa, ramis 1-cephalis, foliis ovato-oblongis per totam longitudinem calloso-serratis, infimis breviter decurrentibus, super. cordato-sessilibus, capitulis ebracteatis glutinosis, invol. squamis lineari-subulatis squarrosis, pappo 2-4-seto. ♂ in Mexico circa Villalpando legit cl. Mendez. Suffrutex circ. pedalis. Rami albidì glabri. Pappus caducissimus. Habitus hinc *G. squarrosa*, illinc *G. pulchellæ*. (v. s.)
4. *G. INULOIDES* (Willd. mag. nat. Berl. 1807. p. 261. enum. 894), caule basi suffruticoso apice pubescente aut sparsè hirsuto, foliis radicalibus spatulatis, caulinis subovatis serratis basi latioribus subamplexicaulibus, ramis apice 1-cephalis subhirtellis, invol. squamis glabris in acumen lineari-subulatum patulum productis, pappo 1-3-seto. ♂ in Mexico (Lag.), planitie altà (H. et B.), inter Vittoria et Tula (Berl. n. 2186), in ditone Texas (Berl. n. 1542 et 2186) et Oaxaca (Andr. n. 271). Dunal! l. c. p. 50. t. 1. Cass. dict. 37. p. 468. bot. reg. t. 248. *Ast. spatularis* Brouss. est serratus Lag. ex Steud. *Grind. spatulata* Link enum. h. ber. 2. p. 336. *Aster spatulatus* hort. madr. 1805. *Inula serrata* Pers. ench. 2. p. 451. *Demetria spatulata* Lag. el. h. madr. 1814. p. 30. (v. v. c. et s. sp.)
5. *G. MICROCEPHALA*, caule basi suffruticuloso glabro, ramis apice parçè pubescentibus, foliis sessilibus subsemiamplexicaulibus ovato-oblongis serratis subpubescentibus, capitulis ramos terminantibus solitariis, invol. squamis glabris in acumen lineari-subulatum productis, pappo disci 4-5-seto, achæniis radii ovatis vix angulatis lævibus. ♂ in sylvis prov. Mexicanæ Texas legit cl. Berlandier (pl. exs. n. 2057) Affinis *G. inuloidi* sed folia et capitula dimidiò minor. Flores disci steriles in specim. meo videntur. Achænia radii matura grossa calva. (v. s.)
6. *G. ARGUTA* (Schrad. in litt.), herbacea subcæspitosa pubescenti-hirtella, foliis infer. spatulatis in petiolum attenuatis, super. oblongis basi angustatis semiamplexicaulibus, omnibus distanter et acutè dentatis, capitulis terminalibus solitariis pedicellatis, invol. squamis linearibus acutis, pappo trisetato. ♀ in Mexico circa Valladolid de Mechoacan et Pazcuaro (H. et B.). *Grindelia angustifolia* H. B. et Kunth nov. gen. am. 4. (1820) p. 309. bot. reg. t. 781. non Dunal. An *Grind. spatulata* Bern. in Link enum. h. ber. 2. p. 336. sed nostra est nec annua nec glabra. (v. s. c.)
7. *G. ANGUSTIFOLIA* (DC. in Dunal l. c. 1819. p. 51. t. 3. non Kunth), caule subherbaceo glabro, foliis angustis integerrimis, radicalibus spatulatis, caulinis distantibus ovatis acutis sessilibus, capitulis pedunculatis, invol. squamis lanceolatis adpressis. ♀ ♂ in Mexico. *Asteris* sp. Moc. fl. mex. ined. *Grindelia Duvalii* Spreng. syst. 3. p. 575. (ubi Duval loco Dunalii ter citatur.)
8. *G. PULCHRELLA* (Dunal l. c. p. 51. t. 2), suffruticosa glabra erecta apice glutinosa, ramis teretibus, foliis caulinis sessilibus oblongis acutis hinc inde acutè dentatis, capitulis ad apices ramorum solitariis foliis 2-3 proximè bracteatis, invol. glutinosi squamis lanceolatis acutis adpressis, pappo 3-aristato. ♀ in Amer. legit cl. Née! circa Mendoza Chilensium ex herb. hort. monsp., sine loci design. in h. Thib. nuno meo. An verè Chilensis? (v. s. ex itin. Née.)
9. *G. INTEGRIFOLIA*, caule herbaceo erecto ramoso sparsè hirsuto, foliis caulinis semiamplexicaulibus ovato aut oblongo-lanceolatis acutis integerrimis membranaceis puberulis margine scabridis, capitulis ad apices ramorum solitariis inter folia suprema bracteantia sessilibus, invol. squamis extus puberis intus glutinosis in acumen filiforme subulatum productis, pappo 1-3-aristato. ♀ in Americæ borealis parte occid. (v. s. ex hort. Soc. hortic. Lond. à filio lect.)

10. *G. RUBRICAULIS*, caule herbaceo basi glabro apice sparsè piloso, foliis sessilibus semiamplexicaulibus oblongis rigidis grossè inciso-serratis glabris, invol. squamis intimis confertè glutinosis, exter. laxiusculis, pappo oligochaeto caducissimo, ligulis circ. 20 basi angustatis. $\frac{1}{2}$ in Californiâ legit cl. Douglas. Caulis ruber. Folia circ. 3 poll. longa 8-9 lin. lata. (v. s. comm. ab. hon. Soc. hort. Lond.)
11. *G. BUPETHALMOIDES*, fruticosa glabra, foliis oblongo-cuneatis obtusis serratis secus ramos floridos confertis, capitulis terminalibus solitariis sessilibus, invol. squamis acuminatis exterioribus amplis foliaceis, recept. demum convexo vix areolato, pappi setis circ. 10 corollæ disci longitudine. $\frac{1}{2}$ in Brasiliæ prov. Rio-Grande. Species patriâ et formâ distinctissima. Ligulæ amplæ flavæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1027 miss.)
12. *G.?* *CORONOPHILIA* (Lehm. ind. sem. h. Hamb. 1828. p. 16. et in linnæâ 1830. p. 376 descr.), herbacea, foliis sessilibus crassiusculis linearibus pinnatifidodentatis rugosis glabris pellucido-venosis, capitulis solitariis, invol. glutinoso squamis ext. linearibus apice patentibus, pappo coronæformi membranaceo setaceo aristato. $\frac{1}{2}$ in terris Mexicanis. Capitula magn. *Matricariæ Chamomillæ*. Achænium obovatum compresso-quadrangulum. An ex pappi structurâ omnino congener?
13. *G.?* *ANOMALA*, fruticosa glabra, foliis lineari-lanceolatis utrinquè acuminatis acutè et remotè serratis subviscosis, capitulis ramos terminantibus solitariis discoideis, invol. squamis glutinosis in appendicem filiformem disco æqualem productis, achænio obcompresso glabro, pappo 2-3-p-seto crasso caducissimo. $\frac{1}{2}$ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1031!). Folia 1 $\frac{1}{2}$ poll. longa, 2-3 lin. lata. (v. s. in h. Mus. reg. Par.)

† *Species dubiæ.*

- G. FRUTICOSA* (Desf. cat. h. par. ed. 2. p. 269), sub *Donia* indescrita et in edit. 3^a omitta.
- G. LAMBERTII* (G. Don in Lond. hort. brit. p. 350). $\frac{1}{2}$ in Mexico. Descr. si edita non novi.

†† *Species exclusæ.*

- G. INCISA* Spreng. = *Calimeris incisa*.
- G. SIBIRICA* Link = ? *Aster conspicuus* Fisch.
- DONIA CILIATA* Nutt. = *Aplopappus ciliatus*.
- DONIA CANARIENSIS* Less. = *Jasonia lævigata*.

Subdiv. 3. *HETEROTHECÆ.*

Pappus diversus in disco et in radio. Ligulæ uniseriales.

- CLXXII. *HETEROTHECA* Cass. bull. philom. 1817. septemb. dict. 21. (1823) p. 130. 51. (1827) p. 460. — Calycium Ell. sketch 2. (1824) p. 339. non Pers. — *Diplocoma* Don in Sweet br. fl. gard. (1828) ser. 1. t. 246. Less. syn. 162. — *Doronici* sp. Cerv. Link.

Capitulum radiatum, fl. radii ligulatis fœmineis 1-serialibus, disci tubulosis 5-dentatis hermaphroditis. Rec. planum alveolatum. Invol. squamæ pauciseriales imbricatæ lineares adpressæ apice subpatulæ. Stigmata radii teretiuscula glabra, disci apice lanceolata hispida. Achænia radii oblonga lævia apice calva, disci cuneiformia villosa papposa aut pauciseta, pappo duplici, exter. brevissimo setoso, inter. longo setoso subscabro. — Herbæ Bor.-Americanae perennes (biennesve?) erectæ paniculato-ramosæ villosæ aut pilosæ. Folia caulina sessilia integra rariùs dentata. Flores flavi. Pappus fl. disci rufus, ext. brevis albidus.

Sect. I. CHÆTACTIS.

Achænia radii pappo simplici pauciseto donata. — Sectio (forsan genus) inter *Chrysopsidem* et *Heterothecam* media.

1. *H. CHRYSOPSIDIS*, tota hispida, caule erecto paniculato-ramoso, foliis caulinis obovato-oblongis lanceolatisve subdentatis, inf. basi in petiolum brevem attenuatis, super. sessilibus, ramis 1-3-cephalis, capitulis pedicellatis, invol. latè campanulati squamis linearibus acutis puberis, achæniis radii glabriusculis pappum paucisetum 1-serialem gerentibus. $\frac{1}{2}$ in Mexico circa Saktillo januar. flor. legit cl. Berlandier. Flores flavi, ligulæ punctulatæ. Pappus disci duplex intense rufus, ext. brevis albidus. Species omninò habita et caract. media inter *Heterothecam* et *Chrysopsidem*. (v. s. comm. à cl. Berlandier.)

Sect. II. GYMNACTIS.

Achænia radii omninò calva.

2. *H. INULOIDES* (Cass. dict. 51. p. 460), villosa-hispida, foliis infer. petiolatis ovatis dentatis, super. sessilibus lanceolatis imò sublinearibus integerrimis, capitulis cymoso-corymbis. $\frac{1}{2}$ (Sw.), ② (Link), in Mexico. *Doronicum Mexicanum* Cerv. in Link et Otto neue abbild. t. 22. *Diplocoma villosa* Don in Sweet! br. fl. gard. ser. 1. t. 246. Less. in Linnæa 1831. p. 144. t. 2. f. 17. (v. v. in h. Genev. et s. sp.)
3. *H. SCABRA*, caule hispidulo paniculato-ramoso, foliis infer. ovalibus dentatis petiolatis, caulinis cordato-ovatis sessilibus, omnibus scabris punctatisque, capitulis divaricato-paniculatis, invol. dorso pilosiusculis, ligulis oblongis. $\frac{1}{2}$? in arenosis maritimis et pascuis siccis Carolinæ et Georgiæ. *Inula scabra* Nutt. gen. 2. p. 151. *Inula punctata* Muhl. cat. p. 76. *Chrysopsis scabra* Ell. sketch 2. p. 339. (v. s. comm. à cl. Elliott.)
4. *H. LAMARCKII* (Cass. dict. 21. p. 130), hispidulo-scabra, foliis lineari-lanceolatis acutis subserratis ciliatis, paniculâ divaricatâ, pedunculis involucrisque viscido-pubescentibus, infer. axillaribus. $\frac{1}{2}$ in Carolinâ (Fraser!), Georgiâ circa Savannah (Nutt.), Marylandiâ (Pluk.)? *Inula subaxillaris* Lam. dict. 3. p. 259. *Inula divaricata* Nutt. gen. 2. p. 152. *Chrysopsis divaricata* Ell. sketch 2. p. 338. — Pluk. alm. t. 340. f. 1. sub n. 5. Capitula triplò ferè minora quàm priorum. (v. s.)
5. *H. LEPTOGLOSSA*, caule erecto sparsè hispido paniculato-ramoso, foliis infer. ovalibus basi attenuato-subpetiolatis, super. sessilibus lanceolatis, omnibus hinc indè serratis utrinquè scabro-villosis subcinereis, capitulis pedicellatis paniculato-corymbosis, invol. squamis linearibus elongatis dorso pilosiusculis, ligulis linearibus. $\frac{1}{2}$ in Mexici prov. Leoninâ ad occid. urbis Guanajuato legit cl. Mendez. Valdè affinis *H. scabræ* sed certè invol. squamis elongatis et ligulis linearibus distincta. (v. s. comm. à cl. inv.)

Subdiv. 4. PSIADIEÆ.

Pappus pilosus conformis sæpius 1-serialis, aut setis saltem inter se similibus. Ligulæ pluriseriales.

CLXXII. ERATO.

Capitulum multiflorum heterogamum, flor. radii pluriserialibus angustè ligulatis fœmineis, disci tubulosis hermaphroditis fauce dilatata limbo 5-dentato. Invol. imbricatum, squamis lanceolatis, exter. rigidè ciliatis, int. paulò longioribus subeciliatis. Recept. planum nudum. Cor. radii secus tubum brevissimè puberulæ, disci ad apices dentium rigidè pilosæ. Anth. ecaudatæ. Stylus apice puberulus, breviter bi- (interdum 3-) fidus. Achænium tetragonum glabrum. Pappus

1-serialis setiformis subscaber. — Herba Oronocensis erecta Polymniam ferè referens. Caulis glaber tetragonus. Folia opposita, petiolis basi latè auriculatis, auriculis in vaginam connatis, limbo latè ovato acuminato grossè inciso-dentato basi 5-7-nervio, suprà juniora adpressè villosa adulta glabra, subtùs secus nervos adpressè pubera. Capitula in corymbum terminalem compositum densum disposita. Ligulæ ex sicco flavæ, cor. tubi colore intensiores. — Dixi ex nomine musæ Eratois alludens ad habitum Polymniæ affinem.

1. E. POLYMNIOIDES. *W* in Peruviae montanis Oronocensibus legit cl. Hænke. Folia petiolo 2-3 pollic. longo, limbo 5-6 poll. longo, 2-4 lato. (v. s. in h. Hænk. à cl. de Sternberg miss.).

CLXXIV. WOODVILLEA.

Capitulum multi- (80-100) florum radiatum, ligulis circ. 3-serialibus linearibus fœmineis sterilibus disco et invol. duplò longioribus, fl. disci tubulosis 5-dent. hermaphr. Invol. campanulatum, squamis 2-3-serialibus linearibus-oblongis æqualibus. Recept. epaleaceum. Ligulæ basi longè tubulosæ hispidæ. Anthæræ ecaudatæ. Stylus radii longè bitidus glabriusculus, disci ramis brevibus crassiusculus, dorso puberis. Achænia villosa, radii linearia sterilia, disci oblonga compressa. Pappus setis pluriserialibus tenuissimis subscabris. — Herba Californica Calendulæ facie glanduloso-villosa. Caulis adscendens teres. Folia alterna oblonga basi attenuata sessilia obtusa integerrima subuliter trivenia. Capitula solitaria terminalia magn. Calendulæ. Ligulæ verisim. flavæ. — Locus in tribu subdubius, artificiali normâ admissus. — Dixi in memoriam cl. Gulielmi Woodville auct. operis meritissimi cui titulus *Medical Botany*. (3 vol. 4°. 1790-1793.)

1. W. CALENDULACEA. — in Californiâ legit cl Douglas. Herba spithamea. Folia 2 poll. longa, 6-7 lin. lata. Ligulæ 6 lin. longæ, lineam latæ. (v. s. comm. ab hon. Soc. hort. London.)

CLXXV PSIADIA *Jacq. hort. Schœnbr. 2. p. 13. t. 152. Less. syn. 190.* — Psiadia et Elphegea *Cass. dict. 43. p. 503 et 14. p. 361.* — Conyzæ et Baccharidis sp. *Lam.* — Thouarsia *Vent. herb.* — Alix et Glutinaria *Comm. herb.*

Capitulum multiflorum; heterogamum, primâ fronte discoideum reverà radiatum, fl. radii pluriserialibus fœmineis ligulâ brevissimâ integrâ, disci tubulosis 5-dentatis hermaphroditis et ovarii abortu masculis paucis interdum subnullis. Recept. epaleatum subalveolatum. Invol. imbricatum pauciserialia squamis margine subscariosis. Achænia radii oblonga glabra erostria. Pappus conformis 1-serialis setosus scabriusculus. — Frutices Mascarenenses junioribus partibus glutinosi seu viscosi aut Madagascarienses non viscosi. Folia alterna petiolata oblonga grossè dentata aut integra. Corymbi compositi polycephali subaphylli.

Sect. I. GLUTINARIA *Comm. mss.*

Frutices Mauritiani glutinosi, foliis eglandulosis sæpius triplinerviis, ligulis numerosis pluriserialibus brevissimis.

* *Foliis serratis aut dentatis.*

1. P. GLUTINOSA (*Jacq. h. schœnbr. 2. t. 152*), ramis teretiusculis, foliis petiolatis lanceolatis utrinquè acuminatis serratis penninerviis, summis integerri-

- mis, capitulis 25-30-floris. ♂ in insulâ Mauritiî (Lam. Ait.), sed specim. Maurit. nunquàm vidi. *Solidago viscosa* Schrad. sert. hannov. 12. t. 6. *Conyza glutinosa* Lam. dict. 2. p. 89. *Erigeron viscosum* et nunc *Erigeron glutinosum* hort. Par. ex Desf. cat. p. 174. Folia eglandulosa juniora valdè glutinosa, 3-4 poll. longa, pollicem lata petiolata. (v. v. c.)
2. *P. BALSAMICA* (DC. in Wight contrib. bot. ind. p. 10), ramulis angulatis, foliis vix petiolatis basi cuneatis trinerviis integerrimis apice penninerviis serraturis grossis et paucis instructis, capitulis 25-30-floris. ♂ in insulis Mauritianis. Variat foliis lanceolatis glabris (*Baccharis arguta* Pers. ench. 2. p. 423. *Conyza Balsamica* Wall. cat. n. 3068. comp. n. 178. *Eupatorium flexuosum* Roxb. fl. ind. 3. p. 415), lanceolatis villosulo-glutinosi, ovali-lanceolatis subacutis (*Elphegea crenata* Cass. dict. 14. p. 362?) et obovatis subretusis (*Elphegea latifolia* Cass. dict. 14. p. 362). Folia et ramuli viscid. Similis Psiad. glutinosa nec certissimè ab eâ separanda. Vulgò in hortis *Banane africain*. (v. s. sp.)
3. *P. RETUSA* (Lam. dict. 2. p. 90. sub Conyzâ), suffruticosa pilis sparsis subhirsutula, foliis longè cuneatis retusis infernè integerrimis triplinerviis apice dentatis, pedunculis axillaribus terminalibusque folio paulò brevioribus corymbam 5-7-cephalum gerentibus, invol. globosi squamis subglabris margine latè membranaceis obtusis. ♂ in insulâ Borboniæ maritimis (Comm. Bontou!). Folia sapore salino donata et aceto ad usum culinarè præparata. Alix salisifolia Comm. mss. et ic. Vulg. *la saliette*, *la bien salée*. Folia 3 poll. longa, 9 lin. lata. (v. s.)
4. *P. DENTATA*, ramis teretiusculis, foliis breviter petiolatis coriaceis linearilanceolatis basi cuneatis integerrimis et subtrinerviis apice serraturis utrinquè 2-5 notatis, capitulis multifloris. ♂ in montanis insulâ Borboniæ (Bontou!). *Elphegea dentata* Cass. dict. 14. p. 363. Folia 1-3 pollic. longa, 3-5 lin. lata viscosa. (v. s.)
- ** *Folius integerrimis.*
5. *P. LINEARIFOLIA*, foliis petiolatis linearilanceolatis utrinquè acuminatis triplinerviis integerrimis, capitulis 10-12-floris. ♂ in insulis Mauritianis, nempè in ins. Borboniæ (Comm.), Mauritiî (Bontou in h. Lindl.). *Conyza linearifolia* Pers. ench. 2. p. 92. *Baccharis linearifolia* Pers. ench. 2. p. 425. *Elphegea lanceolata* Cass. dict. 14. p. 562. Species distinctissima evidenter duabus prioribus affinis et caract. Psiadiæ exhibens. Ligulæ paulò majores quàm in prioribus. Flores flavi. Folia et ramuli glutinosi. Folia cum petiolo 3 pollicaria, 4-5 lin. lata. Sequentis var. videtur ex Bojer obs. in litt. (v. s. in h. Lindley.)
6. *P. INTEGERRIMA* (DC. in Wight contrib. p. 10), foliis distinctè petiolatis ovalibus obtusis mucronatis integerrimis subtripplinerviis, capitulis 8-12-floris. ♂ in montanis insulâ Mauritiî et culta? in Indiâ orientali (Wight) et Moluccis (D'Urv.) Ligulæ albidæ paulò majores quàm prioris. Fl. tubulosi paucissimi imò nulli, undè stirps subdioica evadit. Petioli circ. semipollicares.
- α. *minor*, foliis triplinerviis, nervo medio vix ac ne vix nervulos pinnatos gerente. *Baccharis viscosa* Lam. dict. 1. p. 345. journ. phys. 1. p. 62. t. 3. *Elphegea minor* Cass. dict. 14. p. 362. (v. s.)
- β. *trinervia*, foliis triplinerviis nervo medio penninervio. Psiadia trinervia Willd. mag. naturf. ges. Berl. 1807. p. 139 huc pertinere videtur etiamsi folia dicantur opposita cum in nostrâ evidenter alterna.
- γ. *quinquenervia*, foliis elliptico-lanceolatis integerrimis 5-nerviis, corymbis densis, capitulis multifloris. ♂ in ins. Mauritianis. *Elphegea quinquenervia* Cass. dict. 14. p. 363 huc relata ex auct. cl. Bojer in litt. sed mihi ignota.
7. *P. PENNINERVIA*, foliis distinctè petiolatis ellipticis integerrimis penninerviis, capitulis 10-12-floris. ♂ in montibus ins. Mauritiî legit cl. Bojer. Psiadia glutinosa Bojer! in litt. Differt à Psiadiâ glutinosâ Jacq. et à divisi. prim. foliis in-

tegerrimis nec serratis, à *P. integerrimâ* et *linearifoliâ* foliis penninerviis nec tri aut triplinerviis (v. s. comm. à cl. Bojer.)

Sect. II. GLANDULARIA DC. non Gmel.

Species Madagascariensis, herbacea? non viscosa, foliis sessilibus penniveniis subtus minutè glanduliferis, pedunculis axillaribus, floribus purpurascens, ligulis paucis oblongis 1-seriatis.

8. *P. MADAGASCARIENSIS*, foliis sessilibus oblongo-aut ovali-lanceolatis acuminatis subintegerrimis penniveniis glaberrimis subtus sub lente glandulosis, pedunculis axillaribus apice 8-10-cephalis folio circ. longitudine æqualibus in paniculam interruptam digestis. 2 $\frac{1}{2}$ in insulâ Madagascar. *Conyza Madagascariensis* Lam. dict. 2. p. 85. *Baccharis Madagascariensis* Pers. ench. 2. p. 423. Folia imâ basi angustè semiamplexicaulia ex sicco non glutinosa. Invol. squamæ glaberrimæ margine membranaceæ. Ligulæ oblongæ. Pappus rufescens. (v. s.)

CLXXVI. MICROGLOSSA. — Conyzæ sp. auct.

Capitulum multiflorum minutissimè radiatum, flor. radii pluriseriatis ligulatis foemineis, ligulis linearibus pappo brevioribus, fl. disci paucis tubulosis 3-5-fidis abortu masculis. Recept. nudum. Invol. imbricatum, squamis lanceolato-linearibus subscariosis erectis. Antheræ basi ecaudatæ. Achænia oblonga compressa? glabra. Pappus pilosus 1-serialis. — Suffrutes Indici aut Africani, erecti aut subvolubiles glabriusculi apice ramosi quasi proliferi. Folia petiolata ovata acuminata integra aut denticulis exsertis acutis exsertè subserrata subtus pubescentia et glandulosa adulta subglabrata. Corymbi aphylli fastigiati divaricati terminales. Pappus rufescens. — Differt à *Conyzâ* fl. radii non tubulosis sed brevissimè ligulatis, ab *Erigeronte* floribus disci masculis paucis et habitu toto.

* Volubiles.

1. *M. VOLUBILIS*, caule volubili, foliis petiolatis ovatis acuminatis, ligulis stylo suo ferè brevioribus, foliis ramulisque glabriusculis. 3 $\frac{1}{2}$ in Indiæ orient. montanis humidis, in insulis Javâ (Blum!) et Penang (Wall!). *Conyza prolifera* Blum! bijdr. 896. *Conyza volubilis* Wall! cat. et herb. n. 3057. comp. n. 167. *Sonchus volubilis* Rumph. amb. 5. t. 104. f. 1 (v. s.)

ℓ. *ambigua*, ligulis stylo suo longioribus, ramulis foliisque subtus brevissimè pubescenti-hirtellis. — in Sierra-Leonâ? Simillima specimenibus prioris à Wallichio in ins. Penang lectis, sed folia subtus densius velutina et confertius glandulosa. An certè diversa? (v. s. ex h. L'Hér.)

γ. *Madagascariensis*, ligulis stylo suo brevioribus, ramulis foliisque subtus brevissimè pubescenti-hirtellis. — in insul. Madagascar ad margines sylvarum legit cl. Bojer, (v. s.)

N. B. An fortè *Conyza pyrifolia* Lam. dict. 2. p. 89 etiam ex ins. Javâ orta esset hujus generis altera species? *Conyza prolifera* Lam. videtur diversissima ex Decaisn. nouv. ann. mus. p. 408.

2. *M. SESSILIFOLIA*, volubilis glabra, ramis teretibus subherbaceis, foliis sessilibus basi obtusis ellipticis acuminatis integerrimis, ligulis stylo suo circ. æqualibus. 3 $\frac{1}{2}$ in insulæ Madagascar (Forb! in Lindl! herb.) sylvis super arbores scandens (Bojer.). Folia 2 $\frac{1}{2}$ poll. longa, pollicem lata. Rami flor. axillares et terminales subpaniculati. Invol. squamæ acutæ. Flores lutei. Pappus rufescens. (v. s. comm. à cl. Bojer.)

** Non volubiles.

3. *M. ALTISSIMA*, arborea glabra viscosa, ramis confertis flexilibus subangulatis, foliis sessilibus lanceolatis basi attenuatis, inf. serratis, super. integerrimis,

- corymbis compositis irregulariter subfastigiatis, invol. squamis obtusis, ligulis stylo suo æqualibus. 5 in insulæ Madagascar prov. Emirnensi legit cl. Bojer. Baccharis altissima Bojer! in litt. 1835. Arbor 25-30-pedalis speciosa. Folia super. 2 poll. longa, 4 lin. lata, cum ramulis glutine viscoso illinita. Flores lutei. (v. s. comm. à cl. Bojer.)
4. *M. HISPIDA*, fruticosa, ramis erectis teretibus subherbaceis hispidis, foliis ovato-oblongis grosse dentatis hispidis in petiolum alatum basi obtusè auritum attenuatis, corymbo composito laxo, invol. squamis acuminatis, recept. convexo alveolato, ligulis stylo suo æqualibus. 5 in insulæ Madagascar prov. Tananarivou circa urbem ad fossarum margines legit cl. Bojer. Frutex 4-5-pedalis. (v. s.)
5. *M. PETIOLARIS*, fruticosa, ramis teretibus subtiliter striatulis et puberulis, foliis petiolatis ovatis acuminatis subintegerrimis suprâ secus nervos subpuberulis subius pallidis pubescentibus, corymbo composito fastigiato, invol. squamis glabris linearibus subacutis, ligulis stylo subæqualibus. 5 in Sierrâ-Leonâ legit cl. Don. Petioli 6 lin. longi. Folia 2 poll. longa, 10 lin. lata. (v. s. in h. Lindl.)

CLXXVII. *NIDORELLA* Cass. dict. 37. p. 459 et 469. v. 56. p. 166. — *Erigerontis*, *Conyzæ*, etc., sp. auct.

Capitulum multiflorum heterogamum, fl. homochromis, radii foemineis minutissimè ligulatis uni- bi- aut pluriseriatis, disci circ. 20 tubulosis 5-fidis bisexualibus, centralibus interdum sterilibus. Invol. campanulatum imbricatum. Recept. alveolatum. Anth. ecaudatæ. Achænia oblonga teretiuscula aut subcompressa, sæpius minutè puberula. Pappus 1-serialis, setis subscabridis basi in annulum subconcretis. — Herbarum suffrutices Australi-Africana, et una ex Arabiâ. Caules erecti. Folia alterna. Capitula corymbosa aut rariùs solitaria. Corollæ omnes flavæ, ligulis interdum disco pallidioribus.

§. 1. Genuinæ, *corymbis polycephalis compositis*.

* *Foliis incisis dentatisve.*

1. *N. MESFILIFOLIA*, suffruticosa ramosa glabriuscula, foliis petiolatis ovatis sen cordatis profundè et acutè inciso-dentatis, capitulis corymbosis. 5 in Africâ australi inter dumeta et sylvulas, in distr. Uitenhage, Stellenbosch et Albany (Eckl. et Zeyh.), ad Zneëreberge et Omsanculo (Drege!). Aster mespilifolius Less. syn. 180. Eupatorium Eckl. ! Rami teretes. Folia aut glabra aut pube minutissimâ scabrida. Corymbi compositi 15-20-cephali. Invol. squamæ oblongo-lineares imbricatæ. Recept. foveolatum. Cor. omnes pallidè flavæ, radii ligulatæ stylo longiores, disci fertiles. Achænia pube minutissimâ scabrida. (v. s.)
2. *N. PINNATILOBATA*, basi suffruticosa subsimplex tota puberulo-hirtella, foliis in petiolum attenuatis pinnatilobatis, lobis utrinquè 2-4 lanceolatis acutis, corymbo composito polycephalo. 5 in Africâ austr.-orient. ad Zw. Key et Basche legit cl. Drege ! Folia pollicem longa, inf. pinnatifida, sup. pinnatipartita lobis angustioribus. Invol. squamæ vix acutæ dorso subscabræ margine angustè scariosæ. Ligulæ minimæ obovatæ stylo longiores. Achænia teretiuscula pubescentia. (v. s.)
3. *N. TRILOBA*, rhizomate suffruticoso, caulibus ramisque erectis foliisque pubescentibus, foliis 3-5-lobatis basi in petiolum angustatis, lobis oblongis terminali (præs. in fol. infer.) majore obtusiore, corymbo subpaniculato laxo, invol. squamis puberis oblongis acutis ext. dorso puberis, int. subscariosis ciliatis, omnibus demum reflexis 5 2 in desertis circa montem Sinai legit cl. Bonpl. (n. 101 !); mens. junio flor. Arabis dicitur Nefflé. An *Erigeron incisus* Sieb. et Spreng. citati? *Conyza triloba* Decaisn. ann. scienc. nat. 2. p. 261. Herbar.

sempedalis pallida, invol. substraminea. Cor. flavidæ. Ligulæ brevissimæ vix perspicuæ transeunt gradatim in tubos, undè species media inter Trimorphæam et Nidorellam. Pappi setæ albæ basi in anulum ab achænio separabilem concretæ. (v. s. comm. à cl. inv.)

4. *N. CHRYSOCOMA*, herbacea, ramis teretibus virgatis villosis, foliis rameis basi obtusè aurito-subcordatis lineari-oblongis grossè serratis aut subincisis pubescentibus, corymbo composito polycephalo, invol. squamis dorso puberulis apice acuminatis. ① in ins. Madagascar ad agros cultos legit cl. Bojer. *Chrysocoma aurea* Bojer! in litt. 1835. Herba strictè erecta speciosa 2-3 pedalis. Flores aurei, ligulæ minutissimæ. Affinis *N. pinnatilobatæ* sed herbacea, folia vix incisa potius dentata, invol. squamæ multò magis acuminatæ. (v. s. comm. à cl. inv.)
 5. *N. RESEDIFOLIA*, herbacea erecta, caule foliorumque petiolis parcè strigillosis, foliis subpuberulis in petiolum longè attenuatis pinnatipartitis, lobis 1-2-jugis oblongis, terminali obovato-oblongo paulò majore, omnibus mucronatis, corymbo composito, invol. squamis elliptico-oblongis obtusiusculis subciliatis. — in Africâ australis regione Gariepina, Klein Namaqualand legit cl. Drege! Ligulæ minutæ pallidius quàm tubuli flavidæ ovales subacutæ. Achænia glabriuscula. (v. s.)
 6. *N. HIRTA*, herbacea subsimplex erectiuscula, caule strigilloso-hirto, foliis hirtello-scabris, infer. oblongis in petiolum longè attenuatis integris aut hinc indè lobulatis, cæteris basi dilatato-amplexicaulibus pinnatipartitis, lobis 1-2-jugis oblongo-linearibus, corymbo composito conferto, invol. squamis linearibus acutis dorso subhirtellis. — in Africâ australi ad Stormberg legit cl. 5-600 ped. legit cl. Drege! Affinis *N. resedifoliæ*. Cor. etiam radii saturatis flavæ. Ligulæ minimæ. Folia in axillis sæpè fasciculata. (v. s.)
 7. *N. AURICULATA*, herbacea erecta subsimplex, caule tereti confertè hirtello, foliis basi rotundè auriculatis pube minimâ pubescenti-scabris, infer. petiolatis latè ovatis grossè dentatis, super. basi suprâ aurículas attenuatis oblongis integris, corymbo composito fastigiato, capitulis congestis, invol. squamis obtusiusculis subciliatis dorso subhirtellis. — in Africâ austr.-orient. ad Zw. Omtata et Qmsamwubo legit cl. Drege! Folia inf. petiolo incl. 4 poll. longa, ferè 2 poll. lata. Corymbus ferè *N. fœtidæ*. Ligulæ minimæ flavæ. (v. s.)
 8. *N. OBOVATA*, herbacea erecta tota brevè et confertè hirtella, foliis obovatis basi cuneato-attenuatis subsessilibus grossè et obtusè dentatis reticulato-nervis confertis, corymbo composito polycephalo, capitulis confertis, invol. squamis oblongis ciliolatis obtusis. — in Africâ austr. distr. Kafferland legit cl. Drege! Folia sesquipollicem longa, 9-10 lin. lata, summa oblonga integra. Ligulæ minimæ obtusæ flavæ. (v. s.)
 9. *N. SENECIONEÆ*, herbacea erecta tota minutè puberulo-hirtella, foliis oblongis infernè attenuatis imâ basi vix auriculatis, infer. grossè paucidentatis, super. integris, corymbo composito fastigiato, invol. squamis oblongis subacutis dorso puberis et subciliatis. ② in Africæ Capensis distr. Uitenhage et Caledon legit cl. Ecklon! et Zeyher. Ligulæ minutæ flavæ. Folia sesquipollie. longa, 4 lin. lata. (v. s.)
- β. *Albanensis*, foliis obovatis magis dentatis in axillis confertis. — in distr. Albany. Eckl.! (v. s.)
10. *N. LINIFOLIA*, herbacea erecta simplex glabra, foliis lineari-lanceolatis acuminatis, infer. basi longè attenuatis grossè serratis, super. sessilibus integerrimis, corymbo composito, invol. squamis acutis dorso glabris margine subciliatis. ①? in Africâ austr.-orient. prope Port-Natal legit cl. Drege! Folia infer. 3 pollic. longa, 3 lin. lata. Ligulæ minimæ obtusæ. (v. s.)

** *Foliis integerrimis.*

11. *N. SOLIDAGINEA*, collo subliguoso, caulibus herbaceis erectis subsimplicibus infernè sparse setosis supernè minutè et confertè puberulis, foliis linearibus

integris acutis minutè puberulis basi subciliatis, infer. basi attenuatis, super. basi subdilatis, corymbo composito, pedicellis involucrique minutè puberulis, ligulis angustis acutis invol. paulò longioribus. \mathcal{U} in Africâ australi ad Wittbergen et Stormberg alt. 4-500 ped. legit cl. Drege. Capitula in genere majora nempe 3-4 lin. diam. Folia 2 poll. ferè longa, $\frac{1}{2}$ lin. lata. (v. s.)

12. *N. HOTTENTOTICA*, caule basi sublignoso erecto villosulo-sublanuginoso, foliis integerrimis subtus laxè lanatis suprâ subglabris, infimis obovatis in petiolum attenuatis mediis oblongo-cuneatis obtusis basi angustis, summis oblongis basi latiusculis apice mucronatis, corymbo composito. \mathfrak{z} ad Cap. Bonæ-Spei, longè ab urbe in regione Transgariepina inter Klaarwater et Liktakun prope Kosi-Fountain legit cl. Burchell et sub n. 2528. cat. geogr. mecum comm. Capitula omninò *E. foetidi* sed habitu et lanugine valdè differt. (v. s.)

13. *N. POLYCEPHALA*, herbacea erecta paniculato-ramosissima tota pube brevi cinerascenti-pubescenti, foliis caulinis auriculato-semiamplexicaulis elongatis lineari-lanceolatis triplinerviis, rameis linearibus uninerviis, omnibus integerrimis acuminatis, corymbo cujusque rami compositio congestifloro, invol. squamis acutis subhirtellis. — in Africâ australi legit cl. Drege! Herba bipedalis. Folia caul. 3 poll. longa 3 lin. lata, ramea pollicaria lineam lata. Ligulæ parvæ disco pallidiores parvæ obtusæ subtridentatæ. (v. s.)

14. *N. MUCRONATA*, caule herbaceo tereti striato villosulo, foliis integerrimis lineari-oblongis mucronatis basi obtusè subauritis sessilibus glabriusculis 1-nerviis, corymbo composito congestifloro, invol. squamis obtusiusculis disco dimidiò brevioribus. (\mathfrak{z}) in insulâ Madagascar savannis ad radices montis Autoungun legit cl. Bojer. *Chrysocoma lappulina* Bojer! in litt. 1835. Affinis *N. rapunculoidi*. (v. s. comm. à cl. Bojer.)

15. *N. RAPUNCULOIDES*, caule herbaceo erecto simplici infernè setis hispido supernè glabriusculo aut minutissimè velutino, foliis elongatis lineari-cuneatis obtusis basi longè attenuatis integerrimis minutè mucronulatis basi ciliatis cæt. sparsè puberulis, corymbo composito, invol. squamis ovatis obtusiusculis, ligulis obtusis integris minimis. \mathcal{U} ? in Africâ australi legit cl. Drege. *Baccharis rapunculoides* E. Mey.! in Dreg. coll. Radix obliqua subsimplex. Caulis bipedalis. Capitula parva. Ligulæ disco pallidiores. Folia 2 $\frac{1}{2}$ poll. longa 4 lin. lata. (v. s.)

16. *N. HYSSOPIFOLIA*, caule suffruticuloso ramoso tereti plùs minùs hirtello, foliis lineari-cuneatis integerrimis obtusis basi attenuatis minutissimis pubescentibus, corymbis compositis, invol. squamis oblongis acutis dorso subhirtellis, ligulis minimis obtusis subcucullatis. \mathfrak{z} in Africâ Capensi (Drege!) ad Albany, Uitenhage, territ. cessum et circa urbem (Eckl.! et Zeyh.). *Baccharis hyssopifolia* E. Mey.! in Dreg. coll. Valdè accedit ad *N. foetidam* sed folia remotiora, capitula minora, etc. (v. s.)

ß. glabrata, caule foliisque glabris. — in Uitenhage. Eckl.! (v. s.)

17. *N. FOETIDA*, caule herbaceo erecto simplici pilis hispido, foliis confertis linearibus basi cuneato-attenuatis obtusis integerrimis submucronatis utrinquè subscabridis, axillis ferè omnibus foliiferis, corymbis pluribus compositis pedunculatis confertis, invol. squamis linearibus acutis dorso subhirtellis, ligulis minimis obtusis. \mathcal{U} in Africâ (Pluk. Herm.), ad Cap. Bonæ-Spei (Th. L'Hér.! herb. Burm.! herb. Eckl.!). *Erigeron foetidum* Linn. sp. 1213. Less. syn. 190. *Inula foetida* Linn. sp. 1241 (ex Bocc. syn.), Willd. sp. 3. p. 2105. Thunb. fl. cap. 667. Burm. prod. 26. *Solidago retusa* Burm.! prod. 27. *Nidorella foliosa* Cass. dict. 37. p. 469. — Mill. dict. t. 233. Pluk. t. 223. f. 3. Herm. lugd. b. t. 662. *Stirpis Melitensis* Bocconi hùc falsò à pluribus relata est *Jasonia glutinosa*. (v. s.)

18. *N. PUNCTULATA*, glaberrima, caule fruticoso, ramis viscosis, foliis in petiolum attenuatis elongato-lanceolatis acuminatis integerrimis utrinquè paucis glandulosè minutissimis notatis, corymbo composito, pedunculis compressis, involucrio pappo et etiam disco breviori, squamis obtusiusculis. \mathfrak{z} ad Cap.

Bonæ-Spei in regione Transgariëpinâ inter montes Asbestinos et Witte-water legit cl. Burchell (cat. geog. n. 2061). Species distinctissima. Folia 3-4 poll. longa, 3-5 lin. lata, crassiuscula. Flores radii feminei brevissimè ligulati. (v. s.)

19. *N. AMPLEXICAULIS*, caule herbaceo glabro simplici apice corymboso, foliis oblongo-ligulatis basi obtusè auriculato-cordatis breviter decurrentibus apice subacutis hinc indè subdentulatis glabris, corymbi compositi ramis pubescentibus. 2½ ad Cap. Bonæ-Spei ad orientem urbis legit cl. Burchell et mecum sub n. 7135. cat. geogr. comm. Herba bipedalis. Folia radicalia basi subangustata, caulina infer. longius decurrentia, 4 poll. longa, 6-8 lin. lata, summa vix decurrentia acuminata. Capitula parva. (v. s.)
20. *N. LONGIFOLIA*, herbacea erecta glaberrima, caule ad collum subadscendente apice angulato, foliis integerrimis radicalibus longissimis oblongis obtusis basi longè attenuatis, caulinis sessilibus semiamplexicaulibus acutis, corymbo composito polycephalo, invol. squamis acutis, ligulis minimis. 2½ in Africâ australi ad Zneenrebergen legit cl. Drege! et fortè in Kafferland cl. Ecklon et Zeyher. Herba ferè bipedalis, similis *N. amplexicauli*, sed folia radicalia pedem et ultrâ longa, 10-13 lin. lata. (v. s.)

§. 2. *Spuriæ, capitulis solitariis terminalibus, foliis sessilibus linearibus integerrimis hirtis.*

21. *N. EXILIS*, caule herbaceo basi suffruticoso erecto exili apice subnudo, ramis paucis elongatis 1-cephalis, foliis linearibus subacutis integerrimis, omnibus ramisque pube minutâ subglanduliformi puberulis, inferioribus majoribus setas longas patulas sparsas insuper gerentibus, capitulis longè pediculatis solitariis, invol. squamis subimbricatis planis linearibus acutis disco æqualibus, lignis 8-10 in sicco revolutis, achæniis villosis. ♂ et fortè (I) in Africâ Capensi ad Clam-Williams leg. cl. Ecklon et Zeyh. Achænia disci in specim. meo pleraque abortiva. Stylorum disci rami conici undiquè hirtelli. Pappus 1-serialis-plumosus. Habitus Senecionis puberali sed genericè diversissima. Flor. disci et radii flavi isochromi. (v. s.)
22. *N. ? GARIËPINA*, caule fruticoso erecto divaricato-ramoso tereti hirtello, foliis linearibus integerrimis utrinquè hirtis, ramis floridis apice breviter denuatis, invol. squamis linearibus acuminatis dorso hirtis apice ad marginem scariosis ciliatis, achæniis vix puberis, ligulis minimis. ♂ in Africâ Capensis regione Gariëpinâ locis asperis legit cl. Drege. Recept. omninò nudum. Inv. biseriale demùm reflexum. Folia 6 lin. longa, semilineam lata. (v. s.)

CLXXXVIII. HOMOCHROMA.

Capitulum multiflorum, fl. radii ligulatis 1-seriatis homochromis, disci tubulosi 5-fidis bisexualibus sed sterilibus. Invol. 2-3-seriale, squamis linearibus acuminatis adpressis. Recept. subconvexum foveolatum. Anth. basi ecaudatæ. Stylus Asteris. Achænia radii plano-compressa obovata villosa, disci filiformia glabra. Pappus 1-serialis setosus, setis radii circ. 15, disci circ. 10, omnibus præsertim ad apicem plumosis. — Suffrutex Capensis ramosus hirtellus, ramis foliosis apice nudis 1-cephalis glanduloso-subhirtellis. Folia alterna sessilia oblongo-linearia integerrima hirtella et ciliata. Capitula flava ligulis majoribus quàm in *Nidorellâ*.

1. *H. ECKLONIS*. ♂ in Africâ Capensi ad Clam-Williams leg. cl. Ecklon et Zeyher. Folia 6 lin. longa, lineam aut sesquilineam lata. (v. s.)

CLXXIX. NEJA *D. Don in Sweet brit. fl. gard. ser. 2. n. et t. 78.* — *Diplopappi* sp. Less.

Capitulum multiflorum heterogamum, fl. radii 1-3-serialibus angustè ligulatis fœmineis, disci 5-dentatis hermaphrod. Recept. planum favulosum. Invol. latè hemisphæricum imbricatum 3-seriale, squamis linearibus subulatis. Cor. extus papilloso-pilosæ, stigm. radii obtusa lævia, disci hirtella acuta. Achænium lineari-oblongum compressum sericeo-villosum. Pappus duplex, ext. brevis angustè paleaceus, interior longior pilosus scaber. — Suffrutices Americani, pilis longis sparsis patulis hispidi. Rami foliosi aut apice nudi 1-cephali. Folia sparsa linearia subacerosa callosomucronata integerrima. Invol. hirtellum. Cor. omnes flavæ. Pappus *Diplopappi*. Ligulæ *Erigeri*. Color florum *Chrysopsidis*. Habitus proprius.

Sect. I. PODONEJA.

Recept. vix favulosum. Cor. papilloso-pilosæ. Capitula longè pedunculata. — Folia impunctata.

1. *N. GRACILIS* (Don in Sweet! l. c.), ramosa, foliis subulato-linearibus parçè et rigidulè pilosis impunctatis, ramis apice longè aphyllis, involucri glabriusculo disco ferè breviorè, ligulis 2-3-seriatis. ζ in Mexico. Lodd. bot. cab. t. 1814. *Diplopappus graminifolius* Less. syn. p. 165. Folia angustissima ferè subulata. Capitulum 8-9 lin. diam. (v. s. ex hort. Barclay et Sweet.)

Sect. II. PHYLLONEJA.

Recept. alveolat. Corollæ glabræ. Capitula sessilia. Folia punctata.

2. *N. MACROCEPHALA*, caule subsimplici per totam longitudinem densè folioso, foliis lato-linearibus punctatis molliter villosis, involucri villosissimo disco longiore, ligulis 2-3-seriatis elongatis. ζ in Brasiliæ provinciâ Rio-Grande. Caulis circ. pedalis foliis obtectus. Folia 12-15 lin. longa, $1\frac{1}{2}$ -2 lata. Invol. squamæ multiseriales elongato-acuminatæ, apice purpurascente rigidulo. Capit. diam. ferè sesquipollicari. Recept. breviter alveolatum. Pappus ext. brevissimus. Habitus ferè inularum sed antheræ omninò ecaudatæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1001, 1006 et 1047 miss.)

Sect. III. MONOGYRIA.

Ligulæ 1-seriales. Capitula ad apices ramorum apice nudorum solitaria. Cor. glabræ. — Sectio *Chrysopsidi* affinis sed achæniâ oblonga nec obovata.

3. *N. NIDORELLOIDES*, caule basi fruticoso multicauli cæspitoso, ramis adscendentibus molliter subhispidis ferè ad apicem usque foliosis 1-cephalis, foliis linearibus aut lineari-sublanceolatis ciliato-subhispidis et glanduloso-puberis, invol. squamis linearibus subpuberulis pappo ext. parco brevi setiformi. ζ \mathcal{U} in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 10481). Pars rami pedunculiformis vix foliis longior. Folia pollicem longa aut paulò longiora, 1-2 lin. lata. Pappi series ext. vix à pilis achæniî distinguenda. (v. s. in h. Mus. reg. Par.)
4. *N. SUBVILLOSA*, caule fruticoso ramoso, ramis villis subadpressis subhirsutovillosis apice longè nudatis vix hirtellis 1-cephalis, foliis subvillosis linearibus eciliatis, invol. squamis linearibus glabriusculis, pappo ext. brevi subpaleaceo. ζ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1015!). Sequenti affinis nec ex specim. manco satis certa. Villositas caulis et foliorum et ciliorum verorum absentia satis notabiles characteres. (v. s. in h. Mus. reg. Par.)
5. *N. LINEARIFOLIA*, caule fruticoso ramoso glabro, ramis hispidulis apice subhirtellis aphyllis 1-cephalis, foliis linearibus elongatis hinc indè ciliatis, invol.

- squamis linearibus subhirtellis, pappo ext. minuto vix paleaceo. ♂ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1034 et 1045). Primâ fronte videtur var. *N. tenuifolia* sed folia planiuscula linearia ferè pollicem longa, et pappus exterior vix oculo nudo facillè perspicuus triplò prioris ferè brevior. (v. s. in h. Mus. reg. Par.)
6. *N. tenuifolia*, caulibus fruticosis ramosis glabris, ramis longè foliosis apice nudis vix subpubescentibus 1-cephalis, foliis lineari-subulatis tortilibus angustissimis longè ciliatis, invol. glabriusculo, ligulis linearibus, pappo utroque circ. 10-seto. ♂ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1009). Valdè *N. ciliari* affinis, sed folia 8-9 lin. longa subtortilia nec stricta. Pedunculi glabriusculi. Caulis magis fruticosus. Pappus pauciorisetus. (v. s. in h. Mus. reg. Par.)
7. *N. ciliaris*, basi suffruticosa multicaulis, caulibus adscendenti-erectis infernè glabratiss apice subaphyllis hirsutis 1-cephalis, foliis lineari-subulatis sessilibus acutis erectis præsertim ad basin longè ciliatis inv. squamis linearibus dorso hirtellis disco paulò brevioribus, ligulis angustè linearibus. ♂ ♀ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1004!). Caulis subcæspitosi semipedalea. Folia 5 lin. longa. Pappus duplex uterque circ. 15-setosus, ext. paleaceus, inter. setiformis. Capit. ferè Nejæ gracilis sed ligulæ 1-ser. (v. s. in h. Mus. reg. Par.)

Subdiv. 5. CHRYSOPSIDÆ.

Pappus duplex, exter. brevis subpaleaceus, inter. setosus. Ligulæ 1-seriales aut rarissimè 2-3-seriales.

CLXXX. CHRYSOPSIS *Nutt. gen. am. 2. p. 150 (sect. Inulæ). Ell. sketch 2. p. 333. excl. sp. — Diplogon Raf. non Brown. — Diplopappi sp. Less. — Inulæ sp. Pursh.*

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci herm. 5-dent. Recept. planum subalveolatum. Invol. imbricatum, squamis linearibus acuminatis. Anth. ecaudatæ. Achænia obovata compressa. Pappus disci et radii duplex, ext. brevis subpaleaceus, int. setiformis scaber elongatus. — Herbæ Americanæ rhizocarpicæ Inuloideæ sæpius ramosæ et villosæ. Folia integra aut dentata alterna. Capitula ramos terminantia. Cor. omnes luteæ. — Differt ab *Inulâ* antheris ecaudatis, ab *Heterothecâ* achæniis omnibus eodem pappo donatis, ab *Oleariâ* achæniis compressis, à *Diplopappo* ligulis flavis.

§. 1. *Herbaceæ.*

1. *C. PINIFOLIA* (Ell. sketch 2. p. 335), glaberrima, caule rigido, foliis linearibus confertis rigidis integerrimis acutis, pedunculis corymbosis 1-cephalis, invol. squamis margine submembranaceis apice pubescenti-lanatis, achænio oblongo villosa. ♀ in collibus arenosis Carolinæ inter flumina Flint et Chatahoochee legit cl. Elliot. Pappus int. rufus, ext. brevis albidus. (v. s. comm. à cl. Inv.)
2. *C. GRAMINIFOLIA* (Ell. sketch 2. p. 334), tota adpressè sericeo-villosa nitens, caule supernè etiâ folioso, foliis lanceolato-linearibus acuminatis erectis integerrimis nervosis, corymbo composito, invol. squamis acuminatis dorso subpubescentibus et distinctè glandulosis, achænio oblongo utrinquè attenuato villosa. ♀ in arenosis siccis Carolinæ et Georgiæ, Floridæ (Nutt.) *Inulâ graminifolia* Michx. fl. bor. am. 2. p. 122. *Nutt. gen. 2. p. 151. Erigeron nervosus* Willd. sp. 3. p. 1953. *Erigeron glandulosus* Poir. dict. 8. p. 487 non Walt. Pappus rufus, ext. brevis setiformis. Caulis apice polycephalus, laxè corymbosus. Specimen habeo è Georgiâ simplex 1-cephalus. An dist. var.? (v. s.)
3. *C. ARGENTEA* (Ell. sketch 2. p. 334), tota adpressè sericeo-villosa nitens, caule supernè subnudo, foliis lanceolato-linearibus erectis acuminatis integerrimis vix subtiliter nervulosis, corymbo subpaniculato, invol. squamis acumi-

natis dorso pubescentibus non aut vix glandulosi, achenio oblongo villosi. \mathcal{L} in siccis Georgiæ, Carolinæ (Ell.), agri Kentuckensis orient. (Raf.!), à Virginîa ad Floridam (Nutt.) Inula argentea Pers. ench. 2. p. 452. Nutt. gen. 2. p. 151. Diplogon stenophyllum Raf. in litt. Valdè accedit ad *C. graminifoliam*. Nomina facile decipiunt. Hæc potius graminea, et altera sericea dicendæ. Nostra interdum ferè glabra lucens et tunc verè graminum folia refert. (v. s.)

4. *C. FALCATA* (Ell. sketch 2. p. 336 in adn. n. 4), lanato-villosa, foliis sessilibus linearibus acutissimis subfalcato-patentibus nervosis utrinquè pilosis, pedunculis paucis axillaribus corymbosis involucriisque villosis. \mathcal{L} in arenosis pinetis Novæ-Cæsareæ. Inula falcata Pursh fl. bor. am. 2. p. 532. Aster gracilentus Banks herb. ex Pursh. Inula Mariana var. β . Nutt. est verisim. diversa. Herba 7-9 poll. longa, meo sensu *C. graminifoliæ* potius quàm Marianæ affinis. (v. s. comm. à cl. Torrey.)

5. *C. MARIANA* (Ell. sketch 2. p. 335), caule villosi, foliis sessilibus oblongo-ellipticis inf. subovatis obtusis distanter dentatis mucronulatis laxè subsericeo-villosis, corymbo simplici, pedunculis fastigiatis, invol. squamis linearibus acutis viscido-pubescentibus, achenio obovato villosi. \mathcal{L} in arenosis siccis Novæ-Cæsareæ (Raf. Torr.), Carolinæ (Ell.), Georgiæ, etc. Inula Mariana Linn. sp. 3. p. 1240 excl. syn. Pluk. Pursh fl. 2. p. 531. Inula Mariana var. α Nutt. gen. sm. 2. p. 151. Diplogon Marianus Raf. in litt. Inula glandulosa Lam. dict. 3. p. 259 ex syn. — Mill. dict. ic. t. 57. Nomen fortè falsum nam provenit ex syn. Pluk. nunc excluso et incertum manet an stirps in Marylandiâ cresc. (v. s.)

6. *C. TRICHOPHYLLA* (Ell. sketch 2. p. 336), caule basi laxè lanuginoso superne glaberrimo, foliis oblongis obtusis laxè lanuginosis margine subscabris, superioribus minoribus glabriusculis, pedunculis corymbosis fastigiatis 1-cephalis glaberrimis, invol. hemisphærici squamis oblongo-linearibus glabris, apice subfoliaceo parè recurvo, ligulis 14-16, achenio oblongo villosi. \mathcal{L} in Carolinâ infer. (Fraser!) et Georgiâ (Nutt.). Inula trichophylla Nutt. gen. 2. p. 150. Valdè accedit ad *C. gossypinam* sed facilè glabritie partium super. differt. (v. s.)

7. *C. GOSSYPINA* (Ell. sketch 2. p. 237), tota laxè lanuginoso-candicans, foliis sessilibus oblongo-spathulatis obtusis integerrimis, pedunculis corymbosis fastigiatis 1-cephalis puberulis, invol. squamis linearibus acutis puberis, achenio oblongo villosi-hirtello. \mathcal{L} à Virginîa ad Floridam et Georgiam, frequens in pinetis. Inula gossypina Michx. fl. 2. p. 122. Nutt. gen. 2. p. 150. Inula glandulosa Lam. dict. 3. p. 259 ex herb. sed excl. syn. Habitus Inularum. Ligulæ 20-25. Pappus int. rufus, ext. brevis albus. (v. s.)

8. *C. DENTATA* (Ell. sketch 2. p. 337), tota lanuginosa, foliis cuneato-obovatis obtusis sinuato-dentatis, superioribus oblongo-ovalibus integris, corymbo simplici, ligulis 30-35 nervosis, achenio hispido oblongo. \mathcal{L} in Georgiâ circa Louisville. Affinis *C. gossypinæ*.

9. *C. VILLOSA* (Nutt. gen. 2. p. 151), subdecumbens hirsuto-villosa, caule ramoso corymboso, foliis sessilibus integris lineari-oblongis subspathulatis acutis, infernè ciliatis margine scabris, capitulis fastigiatis, achenio obovato-villosi. \mathcal{L} in planitie Missouriianâ à fonte ad confluentem (Nutt.). Amellus villosus Pursh fl. 2. p. 564. Sideranthus integrifolius Fras. ex Steud. Species ex Nutt. distinctissima ob caulis pubescentiam duplicem villosam nempe et pilosam. Ligulæ flavæ circ. 25 lineares oblongæ.

§. 2. Fruticosæ.

10. *C. HIRTELLA*, caule basi fruticuloso ascendente ramoso, foliis ad basin ramosum confertis linearibus integerrimis hirtellis, ramis basi hispidulis foliosis apice glabriusculis aphyllis 1-cephalis, invol. demùm omninò reflexi squamis linearibus dorso hispidulis. \mathcal{L} in Amer. bor. occident. ad flumen Columbia legit cl. Douglas. Herba 5-6 pollic. Folia $1\frac{1}{2}$ poll. longa, $\frac{1}{2}$ -1 lin. lata. Ligulæ circ. 25 lineares flavæ. Achenia oblonga compressa parè pubera. Pappus duplex.

uterque panicetosus. Recept. sine alveol. rudim. nudum, setulam ex areolis præ. marginalibus proferens. (v. s. comm. ab hon. soc. hort. Lond.)

11. *C. CANESCENS*, caule basi fruticoso erecto ramosissimo, ramis junioribus tomentoso-canis adultis foliisque adpressè puberis cinereo-subcanescentibus, foliis lineari-subulatis integerrimis infernè confertis supernè nudis, pedunculis 1-cephalis subaphyllis laxè corymbosis, invol. squamis linearibus acutis disco brevioribus dorso puberis. $\bar{\zeta}$ in Amer. boreal. occid. ad flumen Columbia legit cl. Douglas. Folia sesquipoll. longa. Ligulæ circ. 25 lineares ex sicco pallidè flavæ disco duplò triplòve longiores. Achænia tenuia sparsè pilosa. Recept. scrobiculatum. (v. s. comm. ab hon. soc. hort. Lond.)

† *Species exclusæ.*

- C. ALBA* Nutt. = *Doellingeria ptarmicoides*.
C. AMYGDALINA Nutt. = *Diplostephium umbellatum*.
C. DIVARICATA Nutt. = *Heterotheca Lamarckii*.
C. HUMILIS Nutt. = *Diplostephium amygdalinum*.
C. LINARIFOLIA Nutt. = *Diplopappus linarifolius*.
C. LINIFOLIA Nutt. = *Galatella linifolia*.
C. OBOVATA Nutt. = *Diplostephium obovatum*.
C. SCABRA Ell. = *Heterotheca scabra*.

CLXXXI. FRESENIA.

Capitulum multi- (15-16) florum homogamum. Invol. squamæ circ. 3-seriales imbricatæ lineares margine submembranaceæ dorso glandulas 1-3-seriatim dispositas gerentes. Recept. areolatum angustum. Cor. tubulosæ glabræ 5-dentatæ fauce non ampliata. Anth. ecaudatæ. Stylus *Chrysocomæ*. Achænia compressa erostria sericeo-villosa. Pappus duplex (more *Vernoniæ*), exter. brevis paleiformis, inter. setaceus scabridus longus. — Suffrutices Australi-Africani glabri, ramis oppositis apice nudis 1-cephalis. Folia opposita linearia acuta ferè subulata integerrima sæpè in axillis fasciculata. Cor. pallidè ex sicco flavæ. — Genus omninò *Diplopappis* peduncularibus simile, diversum ligulis nullis et foliis oppositis, cæterum à *Pteroniâ* et *Chrysocomâ* pappo biseriali inæquali faciliè distinctum, à *Vernoniâ* stigmate *Asteroides*, à *Pegolettiâ* antheris ecaudatis diversum, dicavi cl. Fresenio botanico Francofurtensi qui *Ruppelliâ* plantas Arabicas et Africanas egregiè descripsit.

1. *F. LEPTOPHYLLA*, foliis oppositis lineari-subulatis glabris. $\bar{\zeta}$ in Africâ Capens. ad Zeederbergen legit cl. Drege. Folia 3-4 lin. longa, $\frac{1}{4}$ lin. lata. (v. s.)
2. *F. ? SCAPOSA*, foliis alternis ad basin ramorum confertis linearibus adpressè villosis. $\bar{\zeta}$ in Africâ australi legit cl. Drege! Caulis fruticosus demissus brevis. Rami veteres reliquiis foliorum vestiti. Folia 6-8 lin. longa aut omninò linearia aut versis apicem subliora. Rami floridi 3 pollic. 1-cephali molliter hirsutuli. Invol. biseriale squamis linearibus dorso villosis. Cor. flavæ glabræ. Achænia oblongum sericeo-villosum. Pappus duplex scaber, exter. brevis. An *Diplopappi* sp. discoidea? An *Freseniæ* sp. *alternifolia*? An genus proprium? (v. s.)

Subdiv. 6. *SOLIDAGINEÆ*. — *Chrysocomearum* et *Asterearum* gen. *Less.*

Ligulæ nunc nullæ nunc 1-seriales. Pappus disci et radii conformis pilosus 1-serialis. Achænia (ferè omnium) angulata seu teretiuscula nec compresso-plana. — *Generum discoideorum* et *radiatorum* junctionem jubent 1°. Specier. ligulis

minimis paucisve imò solitariis cum discoideis analogia (*Euthamia*, *Bigelowia*); 2°. variationes in toto ordine frequentes inter capitula discoidea et radiata sæpe in iisdem generibus imò iisdem speciebus obviæ, præsertim in capitulis homochromis.

CLXXXII. *BIGELOWIA* DC. non Spreng. nec Smith nec Raf. (1). — *Chrysocoma* Americanæ auct.

Capitulum 3-5-florum homogamum aut ligulâ unicâ fœmineâ heterogamum. Recept. angustum squamam setaceam basi latiusculam achænia æquantem inter flores centralem gerens. Invol. oblongum, squamis paucis imbricatis erectis. Achænia oblonga subangulosa pubescenti-villosa. Pappus pilosus 1-serialis rigidus scaber. — Herbæ Boreali-Americanæ perennes. Folia alterna integra, caulina oblonga aut linearia. Capitula in corymbum disposita. Cor. luteæ. — Genus differt à *Linosyride* pappo 1-seriali, à *Chrysocoma* invol. squamis non foliaceis, ab utrâque capitulis paucis nec multifloris, achæniis teretiusculis nec compressis, pappo rigido, recept. subsquamigero, etc. A *Chrysocoma* separatum dicavi cl. J. Bigelow qui floræ Americanæ auream coronam florâ Bostoniensî et medicâ addidit.

§. 1. *Genuinæ, glaberrimæ, capitulis 3-4-floris.*

1. *B. NUDATA*, herbacea glaberrima, foliis radicalibus spathulato-lanceolatis obtusis tenuiter 3-nerviis, caulinis distantibus linearibus, corymbo composito fastigiato, capitulis 3-4-floris. 2℥ ad paludum margines à Virginiâ ad Floridam. *Chrysocoma nudata* Michx. fl. bor. am. 11. p. 101. Nutt. gen. 2. p. 137. Ell. sketch 2. p. 309. (v. s.)
2. *B. VIRGATA*, herbacea glaberrima, foliis angustis linearibus enerviis, caule virgato ramoso, ramis corymbiferis fastigiatis, capitulis oblongis 3-4-floris, involucri squamis glutinosis adpressis. 2℥ ad paludum margines in Novâ-Cæsareâ (Nutt.) et in Floridâ prope Savannah (h. Dub.). *Chrysocoma virgata* Nutt. gen. am. 2. p. 137. (v. s.)

§. 2. *Spuriæ, pubescenti-tomentosæ, capitulis 5-floris. — An certè congeneres?*

3. *B. MISSOURIENSIS*, herbacea tomentoso-pubescentis, foliis angustis linearibus, corymbo laxiusculo, invol. 5-floris, squamis glabris, exter. patulis. 2℥ ad ripas fluminis Missouri. *Chrysocoma nauseosa* Pursh fl. bor. am. 2. p. 517. Nutt. gen. am. 2. p. 137.
4. *B. DRACUNCULOIDES*, suffruticosa, ramis albidis pulverulento-tomentosis, foliis linearibus 3-nerviis impunctatis, corymbo fastigiato, capitulis 5-floris, invol. angulato lævi. 3 in locis denudatis ad ripas flum. Missouri. *Chrysocoma dracunculoides* Pursh fl. 2. p. 517 non Lam. *Chrysocoma graveolens* Nutt. gen. am. 2. p. 136. Odor rutaceus.

§. 3. *Solidagineæ, glabra, capitulis 5-floris, flor. 4 tubulosis, 1 ligulato. — An Solidagini an Bigelowiæ associanda?*

5. *B. ? UNILIGULATA*, glabra, foliis lanceolatis utrinquè acuminatis serratis, paniculâ compositâ multiflorâ, fasciculis erectis, invol. angusto-oblongis 5-floris, ligulâ unicâ. 2℥ in Novâ-Cæsareâ (Greene), in arenosis à Virginiâ ad Carolinam

(1) *BIGELOWIA* Smith cycl. = *Borya* Willd. — *BIGELOWIA* Spr. neu. entd. est ex ipso auct. *Samydæ* spec. et Spreng. syst. est eadem ac *Borreria*. — *BIGELOWIA* Raf. est omnibus ignota.

(Pursh). *Solidago pauciflorescens* Michx. fl. bor. 2. p. 116. Pursh fl. 2. p. 537 huc pertinere videtur etiamsi nostra non habeat folia obtusa enervia nec caulem fruticosum. Vereor ne cl. Michx. confuderit Big. nudatam et B. pauciflorescentem. Pursh qui ex Michx. dixit caulem fruticosum addit signum ♀, unde caract. non videtur satis notus. (v. s. comm. à cl. Greene.)

CLXXXIII. *SOLIDAGO* Linn. gen. n. 955. Gært. fr. 2. p. 447. t. 170. Nutt. gen. am. 2. p. 159. — *Virga-aurea* Tourn. inst. p. 483. t. 275. Vaill. act. ac. Par. 1720. — *Doria* Adans. fam. 2. p. 124. — *Solidago* et *Euthamia* Cass. Less.

Capitulum pluri- aut multiflorum radiatum, ligulis 5-15 oblongis foemineis 1-serialibus sæpè inter se distantibus, fl. disci hermaphr. 5-dentatis. Recept. epaleaceum nunc omninò nudum nunc alveolatum seu fimbrilliferum. Invol. adpressè imbricatum polyphyllum sæpè oblongum. Achænium teretiusculum multicostatum. Pappus 1-serialis pilosus scabridus. — Herbæ pleræque Americanæ, rhizocarpicæ, rarò gerontogæ, interdum basi suffrutescentes elatæ rusticæ. Folia alterna integra aut serrata. Racemi nunc terminales nunc axillares, erecti patientes imò reflexi, pedicellis secundis et cymas subscorpioideas conficientibus aut verè racemosis. Ligulæ semper luteæ exceptâ S. bicolore. Discus semper luteus.

Sect. I. *VIRGAUREA*. — *Virga-aurea* Tourn. Vaill. — *Solidago* Cass. dict. 37. p. 459 et 472. Less. syn. 163.

Involucrium imbricatum, squamis liberis. Flor. ligulati pauciores quàm fl. disci. Recept. nudum aut alveolatum, non fimbrilliferum. — Racemi paniculati aut simplices, non corymbosi, capitulis ad apices pedicellorum solitariis.

SER. I. UNILATERALES Loud. hort. brit.

Racemi secundi subscorpioidei.

* *Triplinerviæ, foliis triplinerviis.*

1. *S. CANADENSIS* (Linn. sp. 1233), caule villosa erecto, foliis lanceolatis serratis triplinerviis scabris, racemis paniculatis secundis recurvis, ligulis abbreviatis. ♀ in sepibus et agris Amer. bor. à Canadâ ad Pensylvaniam (Pursh) et in montibus Carolinæ (Ell.). Variat foliis nudiusculis et scabris, serratis aut subintegerrimis, racemis elongatis et patentibus. — Pluk. alm. t. 263. f. 1. Huc *S. humilis* Mill. dict. n. 16. Planta polymorpha, ligulis brevissimis inter omnes ferè congeneres distincta. Confr. diss. Tauschii in florâ 1828. p. 497. in quâ ex plantis hortensibus plures varietates aut species huic et seq. affines distinguuntur. (v. s. ex Kentucky à cl. Rafin. comm. et v. c.)
2. *S. PROCERA* (Ait. kew. ed. 1. v. 3. p. 211), caule hirsuto erecto, foliis lanceolatis serratis triplinerviis scabris subtus villosis, racemis spiciformibus erectis, innuptis nutantibus, ligulis abbreviatis. ♀ in sepibus et paludosis Amer. bor. à Canadâ ad Virginiam (Pursh) et fortè in Georgiâ occid. (Ell. sketch 2. p. 369?). Herba 4-6-pedalis. Olim in hort. Par. culta sub nom. *S. hirsutæ* (L'Hér. herb.). *S. procera* Desf. cat. Affinis *S. Canadensi* sed capitula multò minora. (v. v. c.)
3. *S. LONGIFOLIA* (Schrader, hort. Gœtt. ex litt. 1832), caule erecto tereti villosa paniculato-ramoso, foliis lineari-lanceolatis acuminatis triplinerviis suprà scabris subtus pubescentibus, inf. serratis, sup. integerrimis, racemis elongatis patulis secundis in paniculam dispositis, invol. glabri squamis subacutis, ligulis linearibus vix disco longioribus. ♀ in Amer. bor. ? (v. s.)
4. *S. NUTANS* (Desf. cat. h. Par. ed. 3. p. 402), caule pubescenti subhirsuto erecto, foliis lanceolatis acutis argutè serrulatis triplinerviis suprà scabris sub-

tus pubescentibus, paniculâ ante anthesin nutante, ligulis abbreviatis. $\frac{1}{2}$ in Amer. bor. ex testim. hortul. Panicula ampla. Capitula magnitudine media inter *S. proceram* et *canadensem*. (v. v. c. in h. Par. jam ann. 1815. sub hoc nom.)

5. *S. REFLEXA* (Ait. kew. ed. 1. v. 3. p. 211), caule erecto subvillosulo, foliis lanceolatis acuminatis versus apicem argutè serratis scabris margine reflexis, ramis subsecundis brevibus in paniculam angustam confertis, ligulis lineari-oblongis disco ferè duplò longioribus. $\frac{1}{2}$ in pinetis et agris vetustis Amer. bor. à Novâ-Cæsareâ ad Carolinam (Pursh). (v. v. c.)
6. *S. FRAGRANS* (Willd. enum. suppl. 59), caule erecto glabro, foliis glabris oblongis utrinquè attenuatis subtriplinerviis, infer. subserratis, superior. integerrimis, racemis subsecundis, pedicellis pubescentibus, ligulis oblongis disco paulò longioribus. $\frac{1}{2}$ in Amer. bor. (Link. enum. 2. p. 531). Bracteæ lineari-subulatæ arcuatò-recurvæ. Invol. squamæ subvirescentes. (v. s. comm. à cl. Gerhard.)
7. *S. LATRIFLORA* (Linn. sp. 1234), caule erecto pilosinaculo, foliis lanceolatis subtriplinerviis glabris margine scabris inferioribus serratis, racemiâ paniculatis brevibus subrecurvis secundis, ligulis lineari-oblongis disco vix duplò longioribus. $\frac{1}{2}$ in campis et sylvis siccis Amer. bor. à Canadâ ad Carolinam. Pursh fl. 530. Ell. sketch 370. Herba dimidiò minor quàm *S. canadensis*. Capitula latiora. (v. v. c.)
8. *S. ANGULATA* (Muhl. ex Schrad! in litt. 1813. Mart.! enum. h. Erlang. 1814), caule erecto angulato apice summo cum pedicellis scabrido, foliis inf. ovatis acuminatis semiamplexicaulibus penninerviis, super. sessilibus lanceolatis triplinerviis, omnibus serratis supra scabris subtus lævibus, racemis brevibus patentibus secundis, ligulis paucis linearibus disco ferè brevioribus. $\frac{1}{2}$ in Amer. septentr. Invol. squamæ glabræ submembranaceæ lineares. Achænia glabriuscula. Ligule 3-4, fl. disci 9-10. (v. s. comm. à cl. Schrader et Mart.)
9. *S. GLABRA* (Desf.! cat. h. Par. ed. 3. p. 402), glaberrima erecta, foliis infer. lanceolatis, sup. lineari-lanceolatis, omnibus acuminatis argutè serrulatis confertè et brevè ciliatis triplinerviis, racemis secundis revolutis, ligulis 10-12 lineari-oblongis disco paulò superantibus. $\frac{1}{2}$ in Amer. bor. In sylvis et sepiis à Novâ-Angliâ ad Pensylvaniam (Pursh). Rami paniculæ subpubescentes. Folia sensim basi attenuata. *S. serotina* Ait. kew. ed. 1. v. 3. p. 211? *S. altissima* glabra L'Hér.! herb. (v. v. c. in h. Par. anno 1819, sed altera species sub hoc nom. anno 1829 colebatur.)
10. *S. GIGANTEA* (Ait. kew. ed. 1. v. 3. p. 211), caule erecto glabro, foliis lanceolatis utrinquè attenuatis serratis margine scabris triplinerviis, racemis paniculatis secundis patentibus, pedunculis hirtis, ligulis lineari-oblongis disco paulò longioribus. $\frac{1}{2}$ in Amer. bor. In apricis ad latera collium fertilium à Novâ-Angliâ ad Virginiam (Pursh). Valdè accedit ad *S. serotinam*. Nervi foliorum laterales validiores. (v. s. ex Noveboraco comm. à cl. Torrey.)
11. *S. CILIARIS* (Muhl. in Willd. sp. 3. p. 2056), caule erecto glabro, foliis glabris, infer. ovalibus acuminatis basi longè attenuatis ciliatis serratis penninerviis, super. lanceolatis subintegris triplinerviis, pedunculis glabris, bracteis ciliatis, ligulis lineari-oblongis disco paulò longioribus. $\frac{1}{2}$ in Pensylvaniâ (Muhl.), Noveboraco (Torr.!). *S. glabra* hort. Par. 1829. non Desf. (v. s. et v. c.)
12. *S. CLELIÆ*, caule erecto glabro tereti, foliis oblongo-lanceolatis sessilibus utrinquè attenuatis argutè serratis medio triplinerviis utrinquè et ad marginem glabris et lævibus, racemis axillaribus folio sæpè brevioribus patentibus secundis, capitulis longè pedicellatis, pedicellis glabriusculis, ligulis disco duplò ferè longioribus. $\frac{1}{2}$ patr. ign. *S. dubia* Scop. ex hort. Ital. non Schult. Affinis videtur *S. patulæ* et *S. argutæ* sed folia triplinervia. Folia radicalia oblonga subobovata in petiolum attenuata. (v. v. c. anno 1808 in hort. cl. Cleliæ Girmaldi prope Genuam.)
13. *S. SCABRIDA*, caule erecto basi glabro tereti apice puberulo angulato, foliis

oblongo-lanceolatis mucrone apiculatis integerrimis aut hinc inde adpressis triplinerviis utrinque subscabridis secus nervum et ad margines rigidius scabro-ciliatis, racemis secundis, bracteolis subulatis, invol. squamis glabris acutis, ligulis lineari-oblongis disco paulò longioribus. \mathcal{L} in Mexico legit cl. Alaman. Affinis S. Canadensi et S. fragranti. (v. s.)

14. S. VELUTINA, tota pube brevi conferta pulverulento-velutina, caule erecto tereti apice paniculato, foliis oblongo-lanceolatis mucrone subapiculatis integerrimis triplinerviis, racemis erectis vix subsecundis parè bracteolatis, invol. squamis linearibus acutis, ligulis disco paulò longioribus. \mathcal{L} in Mexico circa urbem (Berl. Alam.) et var. paniculà contractà ad Real del monte (Hænk!). Sol. Mexicana Berl. pl. exs. n. 930. à bot. Mexic. plurimis cum S. limonifolià et S. paniculatà sæpè confusa sed facillè distincta. (v. s.)

15. S. DECEMFLORA, caule erecto tereti folisque pube brevissimà subscabris, foliis oblongis mucronulatis integerrimis subtriplinerviis, radicalibus basi attenuatis, racemis subsecundis in paniculam corymbosam dispositis, invol. glabri squamis linearibus, capitulis 10-floris, ligulis 5 minutissimis. \mathcal{L} in Mexicanæ prov. Texas districtibus orientalibus legit cl. Berlandier (pl. exs. n. 1924). Achænia sericeo-villosa. Herba pedalis. (v. s.)

16. S. PUNCTICULATA, caule erecto tereti glabro, ramis apice subscabridis, foliis oblongo-linearibus margine subscabris integerrimis subtriplinerviis glabris utrinque sub lente opacè puncticulatis, racemis subsecundis in paniculam oblongam dispositis, invol. glabri squamis linearibus, capitulis 10-floris, ligulis 4-5-minutis. \mathcal{L} in Mexico ad Rio de la Guadalupe legit cl. Berlandier (pl. exs. n. 1900). Affinis S. decemfloræ sed præter notas indic. achænia glabriuscula. (v. s.)

17. S. POLYGLOSSA, caule erecto tereti hirsuto, foliis elongato-lanceolatis mucrone acuminatis, super. apice parè et acutè serratis triplinerviis utrinque pubescenti-hirtellis, racemis secundis patulis subarcuatis in paniculam dispositis, invol. glabri squamis linearibus acutis, floribus circ. 20, ligulis 12-14 disco paulò longioribus. \mathcal{L} in Brasiliæ prov. Rio-Grande. Habitus et folia ferè S. giganteæ aut S. nutantis. Capitula ferè S. microglossæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 860 missa.)

18. S. MICROGLOSSA, caule basi suffruticuloso erecto tereti subhirsuto, foliis sessilibus elongato-lanceolatis acuminatis puberulis trinerviis, infer. adpressè et distanter serrulatis, super. integerrimis, racemis secundis in paniculam oblongam dispositis, pedicellis bracteolatis, invol. squamis linearibus glabris, floribus circ. 20-25, ligulis 10-15 disco æqualibus. \mathcal{L} in Brasiliæ parte dictà Banda orientale legit cl. Bacle. Achænia gracilia glabriuscula. (v. s. comm. à cl. Bacle.)

β . *Megapotamica*, paniculæ ramis longioribus magis patulis. \mathcal{L} in Brasiliæ prov. Rio-Grande. Priori similis sed major et panicula magis diffusa. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 846, 854 et 857.)

19. S. MARGINELLA, caule suffruticuloso? erecto tereti vix puberulo, foliis linearibus integerrimis acuminatis ciliolato-scabris, infer. basi attenuatis subtriplinerviis, racemis secundis patulis, pedunculis puberis, invol. squamis linearibus utrinque lineà scariosà albà marginatis, floribus circ. 25, ligulis 15 disco æqualibus. \mathcal{L} in Brasiliæ parte dictà Banda orientale legit cl. Bacle. Pappus rufus. Achænia glabra. Stirps vix pedalis. Affinis S. microglossæ. Species notabilis pilis rigidis brevibus nervulis secus marginem folii circ. biserialibus. (v. s. comm. à cl. Bacle.)

β . *sublanceolata*, caule elatiore glaberrimo, foliis lineari-lanceolatis utrinque glabris, paniculæ ramis puberis, ligulis minimis. \mathcal{L} in Brasiliæ prov. Rio-Grande. Herba sesqui- aut bipedalis. Folia 2-3 lin. lata. Pappus albus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 870 miss.)

** Venosæ, foliis penninerviis aut venosis.

20. S. ASPERA (Ait. kew. ed. 1. v. 3. p. 212.), caule erecto tereti piloso, foliis

ovato-ellipticis scaberrimis rugosis serratis, infer. penninerviis, sup. enerviis, racemis paniculatis secundis, ligulis lineari-oblongis disco paulò longioribus. 2 $\frac{1}{2}$ in Amer. bor. campis siccis et sylvis à Noveboraco ad Carolinam (Pursh fl. 530). Verisim. secus montes nec in planitie (Ell. sketch 2. p. 371). — Dill. elth. 2. t. 305. f. 392. opt. (v. v. c.)

21. *S. ALTISSIMA* (Linn. sp. 1233. excl. Dill. syn. omnibus), caule erecto hirtio, foliis lanceolatis penninerviis profundè serratis scaberrimis rugosis, paniculis secundis, ligulis 7-10 lineari-oblongis disco vix longioribus. 2 $\frac{1}{2}$ in Amer. bor. sepibus et agris frequens. — Mart. cent. t. 14. *S. rigidula* Bosc! herb. et hort. Paris. ann. 1828. Huc ex cl. Ait. pertinent *S. altissima*, pilosa, recurvata et virginiana Mill. dict. Species valde varians. *S. Canadensi* affinis. (v. v. c.)
22. *S. RUGOSA* (Mill. dict. n. 25. Willd. sp. 3. p. 2058), caule erecto hirtio, foliis infer. oyatis dentato-serratis, super. lanceolatis adpressè serratis omnibus scabris rugosis acutis, racemis paniculatis secundis patentissimis, ligulis lineari-oblongis disco vix duplò longioribus. 2 $\frac{1}{2}$ in Amer. bor. siccis à Canadà ad Virginiam (Pursh. fl. 537), in pinguib. Carolinæ (Ell. sketch 2. p. 372). An *S. altissimæ* var. ? ex Ell. — *S. altissima* E. Ait. kew. 3. p. 213. — Dill. elth. t. 308. f. 396. (v. s.)
23. *S. VILLOSA* (Pursh fl. 2. p. 538), caule villis patulis hispido tereti, foliis super. sessilibus ovato-lanceolatis mucronatis integerrimis sæpè subtus secus nervum hispidis, infer. serrulatis, omnibus 1-nerviis margine scabridis, racemis secundis folio paulò longioribus in paniculam dispositis, invol. glabri squamis acutis, ligulis vix disco longioribus. 2 $\frac{1}{2}$ in Amer. bor. sylvis et agris frequens (Pursh), in Carolinà (Ell. sketch. 2. p. 372). Habeo ex Carolinà (Bosc!.) et ex Georgià (h. Duby!), sed specim. Georgic. habet capitula minora. *S. altissima* Bosc. herb. *S. Bosciana* Moret! rel. Bosc. ined. *Achænia* vix puberula. (v. s.)
24. *S. SCABRA* (Muhl. in Willd. sp. 3. p. 2059), caule erecto hirtio sulcato, foliis oblongis utrinquè attenuatis acuminatis suprà glabris subtus rugosis scabris medio adpressè serratis, racemis secundis, ligulis lineari-oblongis disco paulò longioribus. 2 $\frac{1}{2}$ in agris et limitibus Amer. bor. à Pensylvià ad Virginiam (Pursh fl. 537), in insulà longà (Rafin!). Folia subtus secus nervos nervulosque penninervios scabra. (v. s.)
25. *S. NEMORALIS* (Ait. kew. ed. 1. v. 3. p. 213), caule erecto pube brevissimà velutino, foliis radicalibus obovato-cuneatis serratis basi longè attenuatis, caulinis lanceolatis integerrimis scabris, racemis paniculatis secundis, ligulis... 2 $\frac{1}{2}$ in arenosis et agris siccis Amer. bor. à Canadà ad Carolinam (Pursh fl. 537. Ell. sketch 2. p. 373.), in Kentucky (Raf.!). Herba tota subcinerascens. (v. s.)
26. *S. PUBERULA* (Nutt. gen. am. 2. p. 162), caule simplici tereti subpubescente, foliis lanceolatis integris pubescentibus utrinquè attenuatis, radicalibus subserratis, racemis spicatis axillaribus erectis condensatis, invol. squamis lineari-lanceolatis acutis, ligulis circ. 10 elongatis. 2 $\frac{1}{2}$ in Novà-Cæsareà (Green.!) circa Amboy, etc. (Nutt.). Habitus *S. nemoralis* quoad folia sed in flor. diversa. (v. s. comm à cl. Greene.)
27. *S. PATULA* (Muhl. in Willd. sp. 3. p. 2059), caule erecto glabro striato, foliis ellipticis serratis glabris, radicalibus oblongo-spathulatis, racemis paniculatis secundis patentibus, pedunculis pubescentibus. 2 $\frac{1}{2}$ in Pensylvià. Folia radic. exactè *Leucanthemi* vulgaris.
28. *S. ULMIFOLIA* (Muhl. in Willd. sp. 3. p. 2060), caule erecto glabro, foliis ellipticis profundè serratis glabris, radicalibus subtus villosis, radicalibus obovatis, racemis paniculatis secundis, ligulis abbreviatis. 2 $\frac{1}{2}$ in paludosis et umbrosis sylvis à Noveboraco ad Virginiam (Pursh fl. 2. p. 538), in Georgià occid. (Ell. sketch 2. p. 375), ex quo et ex Muhl. caulis villosus striatus, folia suprà subscabra, pedunculi villosi.
29. *S. ARGUTA* (Ait. kew. ed. 1. v. 3. p. 313), caule erecto glabro, foliis glabris

- ad marginem subscabrido-ciliolatis argutè et inæqualiter serratis, radicalibus ovato-oblongis, basi attenuatis, caulinis ellipticis acutis, racemis paniculatis secundis, ligulis lineari-oblongis elongatis. $\frac{1}{2}$ in sylvis et pratis Amer. bor. à Canadâ ad Virginiam (Pursh), in Carolinâ (Ell.), ad Noveboracum (Torr.!)-S. argentea Horn. suppl. p. 99 (saltem stirps sub hoc nom. in hortis vagans), non differt, ni fortè capitulis paulò majoribus. (v. s. sp. et v. c.)
30. *S. CINERASCENS* (Schwein in Ell. sketch 2. p. 375), caule erecto gracili pubescente, foliis elongatis lineari-lanceolatis basi attenuatis serratis utrinquè subscabris pubescentibus, racemis secundis recurvis, pedunculis ligulisque elongatis. $\frac{1}{2}$ in Carolinâ boreali prope Salem et in Georgiâ occid. (Ell.)
31. *S. JUNCEA* (Ait. kew. ed. 1. v. 3. p. 213), caule erecto glabro, foliis lanceolatis glabris margine scabris, infer. serratis, racemis paniculatis secundis. $\frac{1}{2}$ in agris et sylvis arenosis à Novâ-Cæsareâ ad Carolinam (Pursh fl. bor. am. 2. p. 538), in Carolinâ superiori (Ell. sketch 2. p. 375 *), in Georgiâ circa Savannah (h. Duby!). Invol. squamæ plerumque apice virides et obtusæ. Ligulæ paucæ (1-4) parvæ disco paulò longiores. Achænia sublentè pubescentia subcompressa. Habitus ferè *Erigerontis* subulati sed ligulæ paucæ flavæ. (v. s.)
32. *S. ELLIPTICA* (Ait. kew. ed. 1. v. 3. p. 2060), caule erecto glabro, foliis ellipticis lævibus serratis penninerviis, nervis utrinquè 1-2 validioribus, racemis paniculatis secundis, ligulis oblongis disco duplò longioribus. $\frac{1}{2}$ in umbrosis rupestribus Amer. bor. à Noveboraco ad Virginiam (Pursh fl. 538). *S. latissimifolia* Mill. ex Ait. *S. plantaginea* Desf. ! cat. h. Par. ed. 3. p. 402. An *S. elliptica* Ell. sketch 2. p. 376, in pinguibus insulæ Paris lecta omninò eadem? Folia sup. subtriplinervia, Racemi suberecti. (v. v. c.)
33. *S. VERRUCOSA* (Schrader. hort. Gœtt. p. 12. t. 6. et in nen. jour. 1810. p. 140), caule verrucoso glabro erecto apice reflexo, foliis ovato-lanceolatis acutis serratis glabris, super. angustioribus, racemis axillaribus folio æqualibus compositis apice reflexis, ligulis ellipticis. $\frac{1}{2}$ in Amer. bor. Affinis *S. argutæ* et *S. ellipticæ* dicitur.
34. *S. RECURVATA* (Willd. enum. p. 889), caule erecto pubescentè, foliis lanceolatis serratis acuminatis margine scabris suprâ glabris subtus secus nervos subscabridis, racemis elongatis secundis recurvatis paniculatis, ligulis oblongo-linearibus disco paulò longioribus. $\frac{1}{2}$ in sylvis umbrosis Amer. bor., in Pennsylvania et Virginiâ (Pursh). (v. s. c. in h. Gœtt. à cl. Schrad. comm.)
35. *S. TORTIFOLIA* (Ell. ! sketch 2. p. 377), caule erecto pubescente, foliis lineari-lanceolatis subserratis patulis tortuosis suprâ nervoque scabris subtus subglabris, paniculâ pyramidatâ, racemis recurvis, capitulis 8-10-floris, ligulis lineari-oblongis disco ferè duplò longioribus. $\frac{1}{2}$ in pascuis siccis Carolinæ et Georgiæ. *S. odora* Michx. fl. 2. p. 118. non Ait. Affinis *S. odoræ* sed folia minora tortilia non pellucido-punctata sed venis ramosis pellucidis notata. (v. s.)
36. *S. PYRAMIDATA* (Pursh fl. bor. am. 2. p. 537), caule erecto tereti hirtò, foliis oblongis acutis subsemialexicauli-sessilibus glabris margine scabris rariter obsolete dentatis, paniculâ nudâ secundâ pyramidatâ, ramis reflexis pedunculis glabris, ligulis minutis. $\frac{1}{2}$ in pinetis Georgiæ (Pursh). Vix à *S. retrorsâ* differt Nutt. gen. 2. p. 159. Ell. sketch 2. p. 378. Achænia lævia.
37. *S. ODORA* (Ait. kew. 3. p. 214), caule erecto pubescente, foliis lineari-lanceolatis integerrimis glabris pellucido-punctatis margine scabris, racemis paniculatis secundis, ligulis oblongo-linearibus disco paulò longioribus. $\frac{1}{2}$ in siccis collibus Amer. bor. à Canadâ ad Carolinam (Pursh fl. 539), in montibus Alleghany (Raf.!). Big. med. bot. 1. p. 188. t. 20. fl. bost. 304. non Willd. nec Michx. ? Sol. lanceolata Bosc ! herb. *S. retrorsa* Michx. fl. 2. p. 117. Ell. ! sketch 2. p. 377. Vulgò dicitur *Goldenrod* et ab Anglo-Americanis ut *Theæ* succedanea admittitur. Folia fragrantia, distillata oleum volatile aromaticum præbent. — Pluk. alm. t. 116. f. 6. (v. s.)
38. *S. GONOLADA*, caule erecto tereti glabro, ramis angulatis glabriusculis pa-

- niculatis, foliis elongato-linearibus sessilibus integerrimis acuminatis subpenninerviis glabris margine scabris pellucido-venosis, racemis secundis, invol. squamis glabris acutis, ligulis vix disco longioribus. $\frac{1}{2}$ in urbis Mexici valle legit cl. Berlandier (pl. exs. n. 728). Folia infer. 3-4 poll. longa, 5-6 lin. lata sup. sensim angustiora et breviora. Invol. stramineum. (v. s.)
39. *S. SEMPERVIRENS* (Linn. sp. 1232), caule erecto glabro, foliis glabris lævibus subcarnosis margine scabris integerrimis, radicalibus ovalibus basi in petiolum attenuatis, caulinis lineari-lanceolatis, racemis paniculatis secundis, pedunculis puberulis, ligulis lineari-oblongis disco duplò longioribus. $\frac{1}{2}$ in humidis et ad ripas rivorum à Canadà ad Pensylvaniam (Pursh), in Noveboraco (Linn. Torr.), in Carolinà (Ell.). *S. Noveboracensis* et *carnosa* Mill. *S. tardiflora* Moench. — Pluk. alm. t. 235. f. 5. Moris. ox. s. 7. t. 23. f. 15. Caulis ruber orgyalis. (v. v. et s.)
40. *S. CORYMBOSA* (Ell. sketch 2. p. 378), caule erecto glabro, ramulis hispidis, foliis inferioribus oblongo-lanceolatis, super. ovatis, omnibus carnosius rigidius glabris margine asperimis ciliatisque, racemis secundis corymbosis, inferioribus recurvis, ligulis elongatis. $\frac{1}{2}$ in Georgià medià, circa Louisville (Ell.). Affinis *S. lævigatæ* et *S. limonifoliæ*. An *S. corymbosa* Poir. suppl. 5. p. 461 eadem aut diversa?

SER. II. ERECTÆ Loud. loc. cit.

Racemi erecti nec secundi.

* *Serratæ, foliis serratis.*

41. *S. BICOLOR* (Linn. mant. 114), caule erecto pilosiusculo, foliis utrinquè pilosis, infer. ellipticis acutis dentatis in petiolum attenuatis, super. elliptico-oblongis mucronatis integerrimis, racemis erectis brevibus, invol. squamis obtusis, ligulis ovalibus disco paulò longioribus, acheniis glabris. $\frac{1}{2}$ in sylvis et collibus siccis Amer. bor. à Canadà ad Carolinam (Pursh). — Pluk. alm. t. 114. f. 3. Achenia in iisdem capitulis compressa et teretiuscula. Racemi nunc foliis paulò longiores nunc breves in axillis conferti. Ligulæ albæ, quæ notà ab omnibus Solid. differt et ad Asterem vergit. (v. v. c.)
42. *S. AXILLARIS* (Pursh fl. bor. am. 2. p. 542), caule glabro tereti stricto, foliis lanceolatis acuminatis remotè serratis glabris margine scabris, racemis in foliorum axillis sessilibus subglobosis erectis, ligulis ovali-oblongis disco duplò longioribus. $\frac{1}{2}$ in sylvis umbrosis à Canadà ad Virginiam (Pursh), in Kentucky (Raf.). *S. lateriflora* Raf. in litt. Dispositione capitulorum accedit ad *S. bicolor*, sed ligulæ flavæ et folia diversissima. Ad hanc alludere videtur Ell. sketch 2. p. 387. in notà sub *S. flexicauli*. (v. s.)
43. *S. PETIOLARIS* (Ait. kew. ed. 1. v. 3. p. 216), caule erecto villosa, foliis ellipticis scabrusculis petiolatis, racemis erectis, ligulis elongatis. $\frac{1}{2}$ in siccis arenosis à Novà-Cæsareà ad Carolinam (Pursh fl. 2. p. 539), in montanis Carolinæ. (Ell. sketch 2. p. 383.)
44. *S. FLEXICAULIS* (Linn. sp. 1234), caule erecto flexuoso glabro angulato, foliis ovatis acuminatis serratis glabris, racemis axillaribus erectis folio brevioribus, summis in paniculam spiciformem confertis, ligulis oblongis disco paulò longioribus. $\frac{1}{2}$ in sylvis et agris Amer. bor. à Canadà (Pursh) ad Georgiam (Ell.). *S. flexicaulis* var. α Willd. sp. 3. p. 2064. Pursh fl. 542. — Pluk. alm. t. 235. f. 3. Variat foliis angustè ovatis et latis imò interdum latioribus quàm *S. latifoliæ*. Caule regulariter flexuoso facillè distincta. Folia secus caulem in nervos 2 decurrentia. *S. scrophulariæfolia* Mill. hujus var. ex Stend. (v. v. c.)
- β ? *ciliata*, foliis latè ovatis profundè argutè et inæqualiter serratis villosociliatis infernè longius acuminatis, racemis folio multò longioribus. $\frac{1}{2}$ in Amer. bor. Noveboraco (Torr.) et Canadà (Bosc.) *S. flexicaulis* β ? *latifolia* Torr. in litt. An fortè spec. propria? (v. s.)
45. *S. LATIFOLIA* (Linn. sp. 1234), caule erecto glabro angulato non flexuoso, foliis ellipticis utrinquè attenuatis serratis glabris, racemis axillaribus folio

- brevioribus, ultimis in paniculam angustam concretis, ligulis oblongis disco paulò longioribus. $\frac{1}{2}$ in Amer. boreali ad Canada (Pluk. alm. t. 235. f. 4). Videtur à *S. flexuosa* (jam mon. Linn. Desf., etc.) sat distincta, caule stricto, foliorum dentibus minoribus magis adpressis. Icones Pluk. optimè hujus et flexicaulis habitum expriment. (v. v. c.)
46. *S. HUMILIS* (Desf. l. cat. h. Par. ed. 3. p. 402, non Pursh), caule erecto subangulato villis patulis hispido, foliis sessilibus confertis ovato-lanceolatis acuminatis nervoso-rugosis serratis suprâ glabris subtus secus nervos villosis, racemis erectis in paniculam dispositis, invol. squamis glabris subacutis, ligulis disco vix longioribus. $\frac{1}{2}$ in Amer. bor. ad Terram-Novam legit cl. Lappylæ. In specim. spontaneis rami breviores conferti, in hortensibus longiores nunc erecti nunc expansi imò reflexi. (v. v. c. et s. sp.)
47. *S. FLABELLIFORMIS* (Wendl. ex hort. Gœtt.), caule erecto glabro striato, foliis caulinis oblongo-lanceolatis utrinquè acuminatis ferè petiolatis subtriplicinerviis glabris margine scabridis hinc indè serratis aut integerrimis, racemis axillaribus erectis paniculatis folio brevioribus, pedicellis glabriusculis bracteolatis, invol. squamis linearibus margine subhyalinis, ligulis ovali-oblongis disco longioribus. $\frac{1}{2}$ patr. ign. Involucra glabra substraminea. (v. s. comm. à cl. Wallroth.)
48. *S. AMBIGUA* (Ait. kew. ed. 1. v. 3. p. 217), caule erecto subflexuoso glabro angulato ramoso, foliis oblongo-lanceolatis densè serratis subtus pilosiusculis, racemis erectis in paniculam subaphyllam dispositis, ligulis oblongis disco duplò longioribus. $\frac{1}{2}$ patr. ign. verisim. Amer. bor. Willd. sp. 3. p. 2064. enum. h. Ber. 891. (v. s. ex hort. Berol. comm. à cl. Wallroth.)
49. *S. MULTIFLORA* (Desf. l. cat. h. Par. ed. 1804. p. 103. ed. 3. p. 402), caule erecto glabro tereti ramosissimo, ramis sparse pubescentibus in paniculam dispositis, foliis sessilibus lanceolatis acuminatis serratis glabris aut summis secus nervos subpuberis, racemis erectis, invol. squamis glabris acutis, ligulis disco paulò longioribus. $\frac{1}{2}$ in Amer. bor. Pers. ench. 2. p. 449. n. 34. Poir. suppl. 5. p. 461. utriq. ex hort. Paris. (v. v. c.)
50. *S. GRACILIS* (Poir. dict. 8. p. 476? DC. pl. rar. j. de Genev. not. 6. p. 6. n. 2), caule erecto glabro subsimplici, foliis lanceolatis acuminatis serratis glabris ciliolatis penninerviis, infer. ovali-lanceolatis basi attenuatis, summis subintegerrimis, pedunculis puberulis, racemis erectis oligocephalis in paniculam angustam dispositis, invol. squamis vix acutis, floribus 9, ligulis 4 disco duplò ferè longioribus. $\frac{1}{2}$ Patr. ign. verisim. Amer. bor. Herba sesquipedalis gracilis, caule rubente. Cl. Poir. dixit folia integerrima, sed in hortis sub nomine Poir. ubiquè occurrit. (v. v. c.)
51. *S. SCHRADERI*, caule erecto tereti sparse pubero paniculato-ramoso, foliis lineari-lanceolatis acuminatis apice serratis glabris margine subscabris, summis integerrimis, racemis erectis brevissimis bracteolatis, invol. squamis acutis, ligulis linearibus disco aequalibus. $\frac{1}{2}$ Patr. ign. verisim. Amer. bor. *S. gracilis* Schrad. l. hort. Gœtt. ex litt. 1832. Affinis *S. gracili* sed folia longiora, angustiora paulò rigidiora. Rami longi in paniculam dispositi et racemulos breves axillares longè gerentes. Ligulæ multò breviores, etc. (v. v. c.)
52. *S. SALICINA* (Elliot sketch 2. p. 389), caule elato gracili supernè pubescente scabriusculo, ramis virgatis elongatis erectis, foliis lanceolatis suprâ scaberri-
mis subtus glabris, inferioribus serratis, racemis subsecundis, ramulis brevibus rariè recurvis. $\frac{1}{2}$ in Georgia occid. Ligulæ 5 tenues. Achænia ferè glabra.
53. *S. CÆSIA* (Linn. sp. 1234), caule erecto lævi, foliis lanceolatis acuminatis serratis glabris, racemis axillaribus erectis, ligulis disco paulò longioribus. $\frac{1}{2}$ in sylvis et agris demissis Amer. bor. à Novâ-Angliâ ad Carolinam et Georgiam. Pursh fl. 540. Ell. sketch 2. p. 385. Ait. kew. ed. 1. v. 3. p. 217. — Dill. elth. 2. t. 307. f. 395. Caulis purpurascens. Achænia ferè glabra. Invol. squamæ obtusiusculæ. (v. s. comm. à cl. Torrey.)
54. *S. LIVIDA* (Willd. enum. 891), caule glabro paniculato, foliis lanceolatis

- serratis glabris margine scabris, ramis apice racemosis, ligulis elongatis. \mathcal{L} in arenosis agris et sylvis à Noveboraco ad Virginiam. (Pursh fl. 2. p. 541.)
55. *S. HISPIDA* (Muhl. in Willd. sp. 3. p. 2063), caule erecto hispido scabro, foliis lanceolatis scabris integerrimis, radicalibus serratis, racemis erectis, ligulis mediocribus. \mathcal{L} in campis graminosis et sylvis Amer. bor. à Novâ-Cæsareâ ad Virginiam (Pursh fl. 541). Similis *S. cæsiæ* sed folia caulina integerrima.
56. *S. HIRTA* (Willd. enum. p. 891), caule paniculato hirtio, foliis lanceolatis utrinquè scabris, caulinis serratis, rameis integerrimis, racemis erectis, ligulis elongatis. \mathcal{L} in Amer. bor. Cæt. ign.
57. *S. GRANDIFLORA* (Desf. ! cat. h. Par. ed. 3. p. 403), caule erecto villosa-scabro apice paniculato, foliis utrinquè villosa-scabris omnibus serratis acutis, infer. ellipticis basi attenuatis, super. oblongo-lanceolatis, racemis erectis compactis, invol. squamis obtusis, ligulis 8-10 elongatis. \mathcal{L} patr. ign. verisim. Amer. bor. Valde accedit ad *S. rigidam* sed præter. car. ind. differt foliis radic. vix petiolatis, limbo basi attenuato nec obtuso, achæniis puberis, fol. superioribus rariter serrulatis sed non integerrimis, etc. (v. v. c.)
58. *S. RIGIDA* (Linn. sp. 1235), caule erecto piloso-scabro apice paniculato, foliis rigidis scabris, radic. petiolatis ovalibus crenato-dentatis, caulinis sessilibus ovato-oblongis integerrimis, racemis erectis compactis, invol. squamis obtusis, ligulis 8-10 elongatis. \mathcal{L} in montanis Amer. bor. à Novâ-Cæsareâ ad Carolinam. Pursh fl. 2. p. 543. — Herm. parad. t. 243. Capitula ampla multiflora. Achænia glabra. (v. s. sp. et v. c.)
- β ? *microcephala*, capitulis multò minoribus, ligulis 11-13, disco abortivo. — Hanc varietatem verisim. monstrosam in horto Parisino lectam à cl. Adr. de Jussieu sub nom. *S. asperulæ* recepi, sed non respond. phrasi Desf. (cat. p. 302). *S. asperula* mihi manet ignota. (v. s.)
59. *S. GLOMERATA* (Michx. fl. bor. am. 2. p. 117), caule humili simplicissimo, foliis glabris oblongo-lanceolatis serratis, racemo simplici glomerulis axillaribus superioribus capitato-congestis constante, invol. turgidis multifloris. \mathcal{L} in montibus Carolinæ (Michx.), prope Salem (Schwein.). Folia infer. ovalia acuminata (Nutt.). Affinis *Asteri* (Nutt.). Distincta serraturis profundis approximatis et racemis capitatis. (Ell. sketch 2. p. 387.)
60. *S. MACROPHYLLA* (Pursh fl. bor. am. 2. p. 542), caule. . . , foliis infer. ovatis acuminatis attenuatis inæqualiter argutè serratis, caulinis lanceolatis utrinquè attenuatis subsessilibus serratis, racemis axillaribus pedunculatis foliosis longitudine foliorum, invol. oblongis turgidis multifloris, ligulis subelongatis. \mathcal{L} in Canadâ. Media inter *Solid.* et *Asterem* (Pursh). Vix à *S. glomeratâ* distincta (Nutt.). †.
61. *S. ELATA* (Ell. sketch 2. p. 389), caule piloso tereti supernè tomentoso, foliis ovali-lanceolatis acutis subintegerrimis venosis subtus tomentoso-pubescentibus, racemis erectis paniculatis, ligulis 7-10 elongatis. \mathcal{L} in pinetis Georgiæ et Carolinæ borealis. An *S. elata* Pursh fl. 2. p. 543 omnino eadem ?
62. *S. SQUARROSA* (Nutt. gen. am. 2. p. 161), caule crasso robusto supernè pubescente, foliis lævibus, inferioribus latissimis spatulato-ovalibus serratis acutis margine scabris, super. sessilibus lanceolato-ellipticis integris racemis glomeratis rigidis pubescentibus, invol. multiflora squarrosa !, ligulis elongatis 10-12. \mathcal{L} in rupestribus Novæ-Cæsareæ (Nutt.), Pennsylvaniae et Georgiæ (Muhl.). Sed fortè ex Ell. sk. 2. p. 389, stirps Georgica differt à specie Nuttalianâ verâ Affinis dicitur *S. strictæ*. Achænia glabra.
63. *S. CARINATA* (Schrad. ! in litt. 1832), caule erecto apice angulato subpuberulo, foliis radic. longè petiolatis oblongis basi attenuatis apice obtusis et mucronatis sinuato-serratis, super. sessilibus linearibus subserratis, omnibus glabris margine scabris nervo medio subtus prominulo, racemis erectis in paniculam dispositis, ligulis 7-9 disco longioribus. \mathcal{L} patr. ign. verisim. Amer. bor. Folia radicalia petiolo incluso 7 pöll. longa, 8-10 lin. lata. Pedicelli puberuli bracteolati. (v. s. c. ex hort. Gœtt.)

64. *S. VIRGATA* (Michx. fl. bor. am. 2. p. 117), caule glabro simplicissimo, foliis subcuneato-lanceolatis obtusis integerrimis glaberrimis adpressis, superioribus sensim minoribus, ramis paniculæ elongatis apice racemosis, pedunculis glabris. $\frac{1}{2}$ in umbrosis humidis sylvis Carolinæ inferioris et Georgiæ. Pursh fl. 2. p. 539. Ell. sketch 2. p. 384. Racemi secundi ex Michx., erecti ex Ell. Andræ species hic confusæ?
65. *S. PULVERULENTA* (Nutt. gen. am. 2. p. 161), caule elato simplicissimo cum foliis et pedunculis pulverulento-pubescente, foliis sessilibus impunctatis, infer. ellipticis serratis, sup. minoribus obovatis integris margine scabris, racemo erecto composito spiciformi, invol. squamis subacutis, ligulis circ. 10 elongatis. $\frac{1}{2}$ in Carolinâ et Georgiâ. Sæpè cum *S. virgatâ* confusa. Fol. inf. ferè *Spireæ salicifoliæ*.
66. *S. SIMPLEX* (H. B. et Kunth nov. gen. am. 4. p. 163), caule erecto simplici subangulato striato supernè hirtullo, foliis lineari-lanceolatis supernè obsolete crenatis infernè valdè angustatis glabris margine denticulato-scabris, paniculâ terminali, ramis circ. 3-cephalis, lignis 8-9 lanceolato-oblongis. $\frac{1}{2}$ in Mexico ad pagum Sancta-Rosa. Habitus *S. Virgæ-aureæ*.
67. *S. VIRGA-AUREA* (Linn. sp. 1235), caule erecto tereti, foliis caulinis lanceolatis utrinquè attenuatis serratis, infer. ellipticis petiolatis, racemo erecto simplici aut composito, invol. squamis linearibus acutis, ligulis circ. 8 elongatis, achæniis puberis. $\frac{1}{2}$ frequens in Europâ et Asiâ boreali, mirè varians, sed veris. in plures species non dividenda et imò varietatibus vix inter se satis diversis conflata. — Hall. helv. n. 69.
- a. *vulgaris*, caule flexuoso apice ramoso et pubescente, foliis ellipticis et lanceolatis serratis subpubescentibus, pedunculis thyrsi lateralibus polycephalis, invol. glabriusculo. — Engl. bot. t. 301. Flor. Dan. t. 663. Doria *Virga-aurea* Scop. carn. 2. p. 176. Sol. *vulgaris* Lam. fl. fr. 2. p. 145. (v. v.)
 - β. *ericetorum*, caule apice ramoso et pubescente, foliis infimis lanceolatis, caulinis lanceolato-linearibus subintegris, capitulis parvis racemosis — in arenosis et ericetis. Sol. *Virga-aurea* β *ericetorum* Duby! bot. gall. 1. p. 266. Sol. *minuta* Thor. chlor. land. (v. v. et s.)
 - γ. *alpestris*, caule glabro simplici aut ramoso, foliis glabris, inf. elliptico-lanceolatis, racemo erecto coarctato, invol. glabriusculo. $\frac{1}{2}$ in montanis. Sol. *alpestris* Waldst. et Kit. in Willd. sp. 3. p. 2065. (v. s.)
 - δ. *Cambrica*, caule simplici apice pubescente, foliis elongato-lanceolatis basi cuneatis dorso ad nervos subhirsutis, racemo coarctato. — in montanis. Sol. *Cambrica* Ait. kew. 3. p. 218. Sol. *minuta* Mill. — Dill. elth. 2. f. 393. (v. v. et s.)
 - ε. *minuta*, caule simplici glabro, foliis elongato-lanceolatis, racemi pedunculis 1-cephalis. — in montanis et alpestribus. Herba digitalis aut palmaris. Sol. *minuta* Linn. sp. 1235. Lodd. bot. cab. t. 189. (v. v. et s.)
 - ζ. *minor* (Pers. ench. 2. p. 450), caulibus cæspitosis humilibus flexuosis, foliis superioribus integerrimis, pedunculis 1-cephalis, ligulis 2-3 minimis. In umbrosis Lusitanix. Brot. fl. lus. 1. p. 382.
 - η. *littoralis*, tota breviter velutino-pubescentis, caule simplici, foliis obovato-lanceolatis subintegerrimis, racemo oblongo conferto, capitulis majusculis. $\frac{1}{2}$ in litore maris Etrusci. Sol. *littoralis* Savi! cent. 182. bot. etr. 4. p. 87. Herba 3. poll. ad 2 ped. Invol. squamæ extus velutinæ. Folia in petiolum attenuata. Ligulæ 8 longæ. Achænium hirsutum. (v. s.)
 - θ. *reticulata*, caule pubescente ramoso, foliis oblongo-lanceolatis integris rugosis pilosiusculis ciliatis subtus reticulato-venosis. — in Pyrenæis orientalibus.
 - κ. *arenaria*, caule pubescente, foliis lanceolatis acuminatis undulatis, racemis axillaribus erectis paniculatis terminalibus. — in Hungariâ. *Solid. arenaria* Horn. hafn. suppl. 1. p. 99. Poir. suppl. 5. p. 461.

- λ. *nudiflora*, glabra, foliis ellipticis acutis basin versus attenuatis, summis integerrimis, pedunculis nudis apice 1-2-cephalis. — in Corsicæ monte Talano. Sol. nudiflora Viv. fl. Cors. sp. nov. 15. fl. ital. fragm. p. 68.
- μ. *Arctica*, caulibus subadscendentibus, foliis oblongis obtusis basi angustatis dense et longè ciliatis, capitulis confertis, ligulis 6-8 minutissimis. ♀ in regionibus boreali-arcticis, in ins. Unalaschkâ (Fisch.), ad sinum Sancti-Laurentii (Cham !). (v. s.)
68. *S. MULTIRADIATA* (Ait. kew. ed. 1. v. 3. p. 218), caule erecto simplici villosulo, foliis sessilibus lanceolatis glabris ciliatis, infer. apice serratis, racemo terminali erecto, ligulis elongatis numerosis. ♀ in Terrâ Labrador et ad Hudsonianum sinum.
69. *S. LEOCARPA*, caule erecto simplici tereti glabro apice pubescenti-villoso, foliis glabris, infer. in petiolum longum attenuatis ovalibus, mediis basi attenuatis oblongo-lanceolatis acuminatis argutè subserratis, summis sessilibus subvis pubescentibus, racemo simplici erecto, invol. squamis glabris acuminatis, capitulo crasso multifloro, ligulis 8-10 disco paulò longioribus, achæniis glaberrimis. ♀ in Amer. boreal. montibus albis (White mountains), legit cl. Boot. An *S. humilis* Pursh. fl. 2. p. 543 ? non Desf. A *S. latifolia* in Pluk. icon. exhib. abundè differt. Affinis *S. minutæ* florist. Europ. sed achæniis glaberrimis differt. (v. s. comm. à cl. Boot.)
70. *S. LEPIDA*, caule erecto stricto tereti puberulo, foliis ovali-lanceolatis basi longè cuneatis integris apice grossè serratis acuminatis penninerviis suprâ sparsè et præcipuè ad margines subvis secus nervos scabridis, paniculâ erectâ terminali, invol. glabri squamis linearibus acuminatis, ligulis disco paulò longioribus. ♀ in Amer. septentr. orâ bor.-occid. ad Mullgrave seu Nootka legit cl. Hænke. Affinis *S. latifoliæ* et ambiguae (v. s. in h. Hænke à cl. de Sternberg miss.)
- β. *subserrata*, foliis parè et minutè serratis ♀ eod. loco à cl. Hænke lecta. (v. s. cum var. α.)
71. *S. CONFERTIFLORA*, herbacea glabra viscosa, caule simplici usquè ad thyrsium folioso, foliis ovali- seu oblongo-lanceolatis basi longè (præs. inferioribus) petiolatis basi integris apice serratis, capitulis in thyrsium spiciformem confertissimis, invol. squamis linearibus erectis, flor. 8-14, ligulis paucis parvis. ♀ in Americâ bor. littore occid. ad Mullgrave et Nootka legit cl. Hænke. Species distinctissima. Ligulas in specim. junior. non ritè vidi. (v. s. in h. Hænke à cl. de Sternberg miss.)
72. *S. SPATHULATA*, glabra, caule erecto tereti subsimplici, foliis obtusis obovato-spathulatis, infer. distinctè triplinerviis apice dentatis basi longè in petiolum attenuatis, super. integerrimis breviter basi attenuatis, ramulis brevibus capitula sessilia congesta pauca gerentibus, invol. squamis lineari-oblongis obtusis, flor. circ. 25 acutis. ♀ in Mexicanis terris legit cl. Hænke. Ex habitu ad *Aplopappum* seu *Grindeliâ* accedit. (v. s. in h. Hænke à cl. de Sternberg miss.)
73. *S. ROTUNDFOLIA* (DC. non Roxb.), caule erecto tereti puberulo scabro, foliis ovalibus ovatis obovatisve obtusis mucronatis utrinquè et margine scaberrimis, infer. crenatis superioribus integerrimis, racemis erectis compactis brevibus in paniculam angustam dispositis, bracteis subrotundis, invol. squamis glabris subacutis, ligulis minimis. ♀ in Mexici prov. Texas distr. orientalibus legit cl. Berlandier (pl. exs. n. 1884). Species ab omnibus formâ bractearum distinctissima. (v. s.)
- ** *Integrifoliæ, foliis integerrimis.*
74. *S. LITHOSPERMIFOLIA* (Willd. enum. 891), caule erecto ramoso pubescente subhispido, foliis lanceolatis utrinquè scabris attenuatis sub-3-nerviis integerrimis, racemis erectis, invol. squamis linearibus acutis, ligulis elongatis. ♀ in sylvis arenosis Amer. bor. à Novâ-Cesareâ ad Carolinam ex Pursh fl. 2. p. 541. Sed cl. Elliot (2. p. 386) in Carolinâ non reperit. Folia apice mucrone calloso apiculata, venis lateralibus minimis. (v. s. ex hort. Hal. comm. à cl. Walloth.)

75. *S. FUSCATA* (Desf. cat. h. Par. ed. 3. p. 402), glaberrima, caule erecto fusco lævi, foliis lanceolatis utriusque integerrimis, racemis erectis, pedicellis brevibus bracteolatis, invol. squamis oblongis glabris vix acutis, ligulis linearibus disco paulò longioribus. \mathcal{L} in Amer. bor. ? Ligulæ 5-6, fl. disci 6-7. Habitus ferè *S. ambiguæ*, sed caulis rubescens, folia integerrima, etc. (v. v. c. in hort. Par.)
76. *S. VIMINÆA* (Ait. kew. ed. 1. v. 3. p. 215), caule erecto subpubescente, foliis lineari-lanceolatis membranaceis basi attenuatis glabris margine scabris, infimis subserratis, racemis erectis, ligulis elongatis. \mathcal{L} ad ripas fluminum Amer. bor. à Canadâ ad Virginiam (Pursh fl. 541). *S. integrifolia* Mill. dict. ex Ait. Affinis *S. limonifolia* sed caulis subpubescens, folia non carnosâ.
77. *S. LIMONIFOLIA* (Pers. ench. 2. n. 36), glaberrima, caule erecto-obliquo, foliis subcarnosis integerrimis lævibus, infer. ellipticis basi longè attenuatis, cæteris lanceolatis sessilibus semialexicaulibus, racemis paniculatis erectis, pedunculis squamosis glabris, invol. squamis linearibus acutis, ligulis elongatis. \mathcal{L} in salsis et ad ripas rivulorum Amer. bor. (Willd.), in Carolinâ (Ell.), nec in Mexico. (Nutt. gen. am. 2. p. 150). *S. Mexicana* Linn. sp. 1234. *S. obliqua* Moench. — Dod. icon. 4. t. 219. Pluk. alm. t. 235. f. 2. malè. Nomen *Limonia* veteris admissum jam præeuntibus Nutt. 2. p. 159 et Ell. 2. p. 380 recepi, cum planta in Mexico non crescat. Affinis *S. sempervirenti* sed racemi non secundi, *S. lævigatæ* sed pediculi glabri. (v. v. c.)
78. *S. LÆVIGATA* (Ait. kew. ed. 1. v. 3. p. 215), caule erecto lævi, foliis sessilibus lanceolatis carnosius integerrimis undique lævibus, racemis paniculatis erectis, pedunculis squamosis villosis, invol. squamis linearibus acuminatis, ligulis angustis elongatis. \mathcal{L} in Amer. bor. paludosis salsis à Canadâ ad Virginiam. Pursh fl. 2. p. 541. Affinis *S. limonifolia* sed si specim. meum cum *Aitoniana* specie omnino congruit, facillè differt foliis triplinerviis et notis indicatis. (v. s. ex hort. Hafn.)
79. *S. INTEGRIFOLIA* (Desf. cat. h. Par. ed. 1804. p. 103. ed. 3. p. 402. non Roxb.), caule erecto simplici puberulo, foliis lanceolatis sessilibus acuminatis tenuè et obsolete triplinerviis glabris, infer. parçè denticulatis, cæteris integerrimis, racemis axillaribus erectis subpaniculatis, pedicellis puberis, invol. squamis acutis, ligulis elongatis. \mathcal{L} in Amer. bor. (Desf.). Pers. ench. 2. p. 449 n. 35. ex hort. Par. An *S. speciosa* Nutt. gen. am. 2. p. 160? et ex eo *S. sempervirens* Michx. ? Affinis *S. lævigatæ* et *limonifolia*. (v. v. c.)
80. *S. PUBESCENS* (Ell. sketch 2. p. 381), caule erecto ramoso pubescente, foliis longè lanceolatis basi attenuatis pubescentibus, inferioribus serratis, racemis erectis paniculatis, ligulis 7-10 mediocribus. \mathcal{L} in humidis Georgiae circa Louisville. *Achænia* minutè pubescentia. Habitus *S. speciosæ* Nutt. (*integrifolia* Desf.?)
81. *S. PANICULATA*, glaberrima, caule erecto apice multiramoso paniculato, foliis lineari-lanceolatis acuminatis sessilibus integerrimis margine scabris triplinerviis, venis lateralibus nervo medio longè parallelis, racemis erectis, pedicellis puberulis, invol. squamis linearibus subacutis, ligulis disco paulò longioribus. \mathcal{L} in Mexico circa urbem. *S. Mexicana* H. B. et Kunth nov. gen. am. 4. p. 104? Allam. in litt. An *Virga-aurea Mexicana* Bauh. pin. 517? Certè à *S. limonifolia* diversa. (v. s.)
82. *S. STRICTA* (Ait. kew. ed. 1. v. 3. p. 216), caule erecto glabro, foliis elongato-lanceolatis acutis glabris margine scabris triplinerviis, radic. basi longè attenuatis serratis, super. integerrimis, racemis paniculatis erectis, pedunculis glabris, invol. squamis acutis, ligulis elongatis. \mathcal{L} in America borealis sylvis arenosis à Novâ-Cæsareâ ad Carolinam (Pursh fl. 2. p. 540). Herba 2-3-pedalis stricta parçè foliosa. (v. v. c.)
83. *S. RECTA* (Pursh fl. 2. p. 542), caule erecto tereti basi glabro apice cum *pedunculis pubescente*, foliis lanceolatis subæqualibus glabris venosis integris *marginè fimbriato-scabris*, racemis erectis brevibus confertis, invol. squamis

linearibus obtusiusculis, ligulis 5-7 oblongis disco vix longioribus. $\frac{1}{2}$ in humidis Carolinæ. Nutt. gen. 2. p. 161. Ell. sketch 2. p. 385. *S. viminea* Bosc! herb. Species botan. amer. ipsis dubia. Specim. meum descr. Ell. sat convenit, à descr. Nutt. differt foliis minoribus sed folia omnia Sphæriâ quadam paraiticâ deformata. (v. s. ex h. Bosc.)

84. *S. ANGUSTIFOLIA* (Ell. sketch 2. p. 388. non Mill.), caule erecto glabro, foliis subulato-linearibus integerrimis glabris, racemis erectis paniculatis, invol. squamis acutis lineari-lanceolatis, ligulis oblongis 7-10 disco paulò longioribus. $\frac{1}{2}$ in insulâ Paris circa Beaufort (Ell.) et in Georgiâ (h. Duby). Affinis *S. vimineæ*. Rami graciles elongati. Folia sup. erecta. (v. s.)

85. *S. LINEARIFOLIA*, caule (ramisve) tereti puberulo, foliis linearibus acutis sessilibus integerrimis tenuè trinerviis glabris margine ciliolato-scabridis, racemis brevibus erectis in paniculam racemiformem confertis, invol. squamis linearibus glabris subscariosis, floribus circ. 20, ligulis 7-8 vix disco longioribus. — in campis Chilensibus dictis *Pampas* olim legit cl. Née. Ex specim. non satis apparet an sit fruticulosa seu herbacea. Invol. pallidè rufescens. Achænia glabra. (v. s.)

¶ *Pœppigii*, caule glabro, foliis margine magis scabrido-ciliatis, paniculæ ramis magis distantibus, capitulis minoribus. — in Amer. austr. legit cl. Pœppig et ex ejus coll. recepi sub. n. CCXXVII sine loci designatione. An spec. propr. ? (v. s.)

86. *S. DOMINGENSIS* (Spreng. ! syst. 3. p. 639), caule erecto subangulato glabro apice paniculato, foliis lato-linearibus rigidiusculis integerrimis obsolete trinerviis lævibus, racemis strictè erectis in paniculam dispositis foliosis, ligulis paucis minutissimis. $\frac{1}{2}$ in Sancto-Domingo legit cl. Bertero. Folia apice callosa sesquipollicem longa, 2 lin. lata, juniora fortè viscosa. Invol. glaberrima pauciflora. (v. s.)

Sect. II. EUTHAMIA. — *Euthamia* Nutt. gen. am. 2. p. 162. Cass. dict. 37. p. 471.

Capitula oblonga. Invol. imbricata squamis subagglutinatis. Flor. ligulati 10-20 nempe floribus disci numerosiores. Recept. fimbriatiferum. Achænia oblongo-ovoidea villosa. — Herbæ, capitulis ad apices ramulorum sessilibus adgregatis in corymbum dispositis. Folia linearia integerrima. — An genus proprium ut innuit Nuttall et asserit Cassini ? Habitus medius hinc inter Solidaginem et Nidorellam, illinc Brachyridem et Bigelowiam.

87. *S. GRAMINIFOLIA* (Ell. sketch 2. p. 391), caule erecto glabro angulato ramossissimo, foliis lanceolato-linearibus 3-5-nerviis secus nervos et margines scabriusculis, axillis nudis, corymbis terminalibus fastigiatis, fl. disci 8-10, ligulis 15-20 longitudine disci. $\frac{1}{2}$ in agris et pascuis ad latera sylvarum et flum. Americæ bor. à Canadâ ad Pensylvaniam (Pursh), in Carolinâ (Ell.), Noveboraco (Torr.!). *Chrysocoma graminifolia* Linn. sp. 1178. *Solidago lanceolata* Willd. sp. 3. p. 2060. *S. lanceolata* α major Michx. fl. 2. p. 116. *Euthamia graminifolia* Nutt. gen. am. 2. p. 162. *Chrysocoma Virginiana* olim in h. Par. (v. v. c. et s.)

88. *S. TENUIFOLIA* (Pursh fl. bor. am. 2. p. 540), caule erecto subscabro angulato corymboso-ramoso, foliis angustissimè linearibus patulis 1-3-nerviis scabris, axillis foliosis, corymbis terminalibus fastigiatis, fl. disci 5-6, ligulis 10 disco paulò longioribus. $\frac{1}{2}$ in pinetis Amer. bor. à Novâ-Cæsareâ ad Floridam (Pursh), in pascuis siccis Carolinæ (Ell. sketch 2. p. 392) et Georgiæ (h. Duby!). *Euthamia tenuifolia* Nutt. gen. am. 2. p. 162. *Sol. lanceolata* β minor Michx. fl. bor. am. 2. p. 540. Caulis humilis. Folia atomis resinosis tecta. (v. s.)

+ *Species non satis notæ.*

89. *S. CANTONIENSIS* (Lour. coch. ed. W. 2. p. 612), caule herbaceo simplicissimo, foliis lanceolatis sessilibus integerrimis glabris, pedunculis polycephalis

subterminalibus, invol. squamis adpressis, ligulis circ. 6. $\frac{1}{2}$ ad Cantonem Sinarum. Affinis dicitur *S. minutæ* Linn.

90. *S. DECURRENS* (Lour. coch. 2. p. 612. ed. w.), caule suffruticoso erecto simplici, foliis radic. lato-lanceolatis subcrenatis in petiolos longos decurrentibus, thyrsio oblongo bracteato, ligulis circ. 7, invol. squamis lanceolato-linearibus. $\frac{1}{2}$? prope Cantonem Sinarum. Caulis 1-ped. Radix repens.
91. *S. ASPERATA* (Pursh fl. bor. am. 2. p. 538), caule paniculato-corymboso, racemis suberectis, capitulis adscendentibus, foliis lanceolatis serratis scabris. $\frac{1}{2}$ in Canadâ. Cæt. ign.
92. *S. NOVEBORACENSIS* (Linn. sp. 1235), caule nudiusculo ramoso fastigiato, foliis radicalibus ovato-oblongis petiolatis scabris, capitulis solitariis amplis. $\frac{1}{2}$ in Amer. bor. ad Noveboracum.
93. *S. DUBIA* (Schult. obs. bot. 178), caule subflexuoso simplici glabro angulato supernè cum pedunculis subpubescente, foliis caulinis oblongo-lanceolatis utrinquè attenuatis grossè serratis utrinquè glabris margine pilis brevibus ciliatis, racemis axillaribus erectis, ligulis elongatis, invol. squamis margine auratis crispatis subciliatis. $\frac{1}{2}$ patr. ign. Affinis dicitur *S. æmbigux* et *alpestri*.

† *Species exclusæ.*

- S. ACAULIS* Spreng. = *Liabum acaule*.
S. ALATA Wall. mss. = *Senecio alatus*.
S. ANGULOSA Wall. mss. = *Senecio angulosus*.
S. ANGUSTIFOLIA Mill. = *Senecio sarracenicus*.
S. ARBORESCENS Forst. = *Steiractis arborescens*.
S. CHINENSIS Spreng. = *Senecio vagans*.
S. CORDIFOLIA Mill. = *Aster cordifolius*.
S. CUSPIDATA Wall. = *Amphirhapis cuspidata*.
S. CYMBALARIA Moench. = *Munychia cymbalaria*.
S. DENSIFLORA Wall. mss. = *Senecio densiflorus*.
S. DORONICUM Linn. = *Senecio Doronicum*.
S. GLAUCA Gmel. = *Senecillis glauca*.
S. GRAVEOLENS Lam. = *Inula graveolens*.
S. HETEROTRICHA Wall. = *Amphirhapis heterotricha*.
S. HIERACIOIDES Spreng. = *Liabum hieracioides*.
S. INTEGRIFOLIA Roxb. = *Melanodendron integrifolium*.
S. LEUCADENDRON Forst. = *Lachanodes Leucadendron*.
S. NEPALENSIS Don. = *Amphirhapis pubescens*.
S. PAUCIFLOSCULOSA Michx. = *Bigelovia uniligulata*.
S. PUBESCENS Wall. = *Amphirhapis pubescens*.
S. ROTUNDIFOLIA Roxb. = *Commidendron rotundifolium*.
S. RUBRICAULIS Wall. = *Amphirhapis rubricaulis*.
S. SPURIA Roxb. = *Commidendron spurium*.
S. SAROTHRÆ Pursh. = *Brachyris Euthamiæ*.
S. SPHACELATA Raf. = *Brachyris ovatifolia*.
S. URTICÆFOLIA Mill. = *Calca urticæfolia*.
S. VISCOSA Lam. = *Inula viscosa*.
S. VISCOSA Schrad. = *Psiadia glutinosa*.

CLXXXIV. AMPHIRHAPIS. — *Solidagin* et *Asteris* sp. Wall.

Capitulum multiflorum radiatum, ligulis 10-20 angustis homochromis, fl. disci tubulosis 5-dent. hermaphr. Invol. imbricatum, squamis pauciserialibus exappendiculatis. Recept. alveolatum. Anth. et styl. Asterearum. Achænia lineari-oblonga modicè compressa villosa aut pubescentia, villis sæpè ad basin pappi calyculum pilosum ferè simulantibus. — Herbæ perennes Indicæ inter *Solidagin*em et *Asterem* mediæ, à *Solidagine* achænio compresso, ab *Astere* ligulis homochromis et achænio lineari-elongato, à *Commidendro* achæniis glabris et habitu toto diversæ.

* *Capitulis corymbosis paniculatisve.*

1. *A. HETEROTRICHA*, caule fruticoso, ramis densè setoso-villosis, foliis brevissimè petiolatis ellipticis utrinquè acuminatis rariter callosò-serrulatis suprà sparse hirtis subtùs adpressè setoso-villosis, corymbo composito polyccephalo, invol. squamis oblongo-linearibus subobtusis villosis, ligulis circ. 10, achæniis villosis. 5 in Indiâ orient. ad Punduah prov. Silhet legit cl. Wallich. *Solidago heterotricha* Wall.! cat. n. 2275. comp. n. 335. Folia 6 poll. longa, 1-1 $\frac{1}{2}$ lata. Habitus ex capitulis densè corymbosis à *Solidagine* alienus. Achæni villi super. pappum exteriorem ferè simulant. Pappus 1-serialis rigidus scaber. (v. s. comm. ab hon. Cæt. merc. Ind. or.)
2. *A. RUBRICAULIS*, caule herbaceo glabriusculo, ramis floridis pubescenti-velutinis, foliis oblongo-lanceolatis basi attenuatis parcè serratis acuminatis glabriusculis, thyrsò elongato laxo, invol. squamis linearibus acuminatissimis dorso puberulis, ligulis 12-15, achæniis villosis. 2½ in Nepaliâ ad Sheopore legit cl. Wallich. *Solidago rubricaulis* Wall.! cat. n. 2273. comp. n. 333. Caulis rubescens. Rami floridi laterales puberuli bracteis foliaceis instructi. Capitula erecta. Pappus albus longitudine cor. tub. Achæni villi supremi pappum exter. simulant. (v. s. comm. ab hon. Aulâ merc. angl. Ind. or. et à cl. Royle.)
3. *A. CUSPIDATA*, caule herbaceo? vix apice summo puberulo, foliis petiolatis ovalibus serratis glabris basi attenuatis apice cuspidato-acuminatis, ramis floridis corymbum gerentibus in paniculam amplam dispositis, invol. squamis linearibus acutiusculis exterioribus brevioribus dorso puberulis, ligulis 12-15, achæniis villosis. 2½ in Indiâ orient. ad Srinaghur legit cl. Wallich. *Solidago cuspidata* n. 2274. comp. n. 334. Folia 5 pollic. longa, 2 ferè poll. lata, petiolo ferè semipollic. Achæni villi super. calyculum circa pappum formant et pappum exter. simulant. (v. s. comm. ab hon. Cæt. merc. angl. Ind. or.)
4. *A. ALBESCENS*, caule herbaceo? erecto ramoso tereti apice minutè pubescente, foliis lanceolatis utrinquè acuminatis suprà glabris subtùs tenuissimè pubescenti-velutinis denticulis minimis remotis subdentatis, corymbis terminalibus polyccephalis subpaniculatis, invol. squamis lineari-lanceolatis acuminatis glaberrimis margine scarioso albis disco brevioribus, ligulis 10-12 linearibus, achæniis villosis. 2½ in Nepaliâ ad Kamaon et Gossain-Than legit cl. Wallich. *Aster albescens* Wall.! cat. n. 2974. comp. n. 84. *Solidago salicifolia* Wall.! in herb. olim ex ipso. Conyza Wall. cat. n. 3066. comp. 176. *Eupatorium cinerascens* Wall.! herb. *Eupatorium coloratum* Spreng. in litt. ad Wall. Folia ramea 2 poll. longa, 6 lin. lata. Pappus rufus. Flores verisim. pallidi. (v. s. comm. ab hon. Cæt. merc. angl. Ind. or.)
5. *A. PUBESCENS*, caule herbaceo simplici erecto pubescente, foliis oblongo-lanceolatis mucrone acuminatis basi attenuatis argutè hinc inde serrulatis utrinquè pubescentibus, thyrsò elongato stricto, invol. squamis linearibus acuminatis glabriusculis, ligulis 7-8, achæniis jun. puberulis adultis glabris. 2½ in Nepaliâ ad Kamaon, Gossain-Than et ad Srinaghur (Wall.! Royle!). *Solidago Virga-Aurea* Ham. ex Don. Sol. pubescens Wall.! cat. n. 2276. comp. n. 336. *Solidago*

Nepalensis Don prod. nep. p. 177. Herba stricta variat nunc 4 poll. oligocephala, nunc bipedalis polycephala.

** *Capitulis solitariis longè pedunculatis.*

6. *A. PEDUNCULARIS*, caulibus herbaceis pluribus erectis subsimplicibus basi angulatis glabris supernè teretibus hispidis, foliis longè petiolatis ovatis subcordatis acuminatis integerrimis suprà petioloque setosis-hispidis margine ciliatis subtus pallidis glabriusculis, pedunculo elongato hispidulo nudo 1-cephalo, invol. squamis paucis foliaceis oblongis dorso hispidis, ligulis circ. 20 linearibus, achæniis villosis. $\frac{1}{2}$ in Indiâ orientali ad Kamaon. Aster peduncularis Wall. l. cat. n. 2967. comp. n. 77. Cineraria ovata Spreng. in litt. ad Wall. Flores ex sicco vid. flavi. Species distinctissima à prioribus habitu sat diversa. Rhizoma vid. repens. (v. s. comm. ab hon. Cæta merc. angl. Ind. or.)

CLXXXV. COMMIDENDRUM DC. in Guill. arch. bot. 1833. v. 2. p. 334. — Conyzæ et Solidaginis spec. Roxb.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis foemineis, disci tubulosi hermaphr. 5-dentatis. Recept. nudum angustum. Invol. imbricatum, squamis linearibus pauciseriatis acuminatis. Antheræ ecaudatæ appendiculatæ. Stylus Asterearum. Achænium compresso-planum erostre glabrum. Pappus 1-serialis pilosus scaber. — Arbusculæ Helenianæ gummiferæ (et idèd *Gum-wood* angl. dictæ, undè cl. Burchell nomen Commidendri finxit). Rami cicatricibus post foliorum lapsum crassis callosis notati, Pedunculi corymbosi polycephali ex axillis supremis aut apice ramorum orti. Flores albi.

§. 1. *Foliis glabris, pedunculis apice ramosis corymbiferis.* — Solidaginis sp. Roxb.

1. *C. ROTUNDIFOLIUM*, foliis latè obovatis basi cuneatis subdentatis longè petiolatis utrinquè glaberrimis, junioribus vernicosis, corymbis compositis polycephalis, pedunculis apice ramosis, pedicellis circ. capituli longitudine. 5 in colibus insulæ Sanctæ-Helenæ. Solidago rotundifolia Roxb. in Beats. app. p. 324. *Bastard cabbage, Gum-Wood* et *Bastard-Tree* vern. angl. Petioli limbo sublongiores. Ligulæ 3-10. Capitula parva. (v. s. comm. à cl. Burchell et Wallich.)
2. *C. SPURIMUM*, foliis cuneato-lanceolatis basi attenuatis integris subsessilibus apice serratis utrinquè glabris, corymbis polycephalis compositis, pedunculis apice ramosis, pedicellis idèd capitulo vix longioribus. 5 in insulâ Sanctæ-Helenæ. Conyza arborescens Forst. comm. Gætt. 9. p. 68? Willd. sp. 3. p. 2053? excl. Ait. syn. Folia suprema 2 poll. longa, 8 lin. lata. Ramuli vix subpuberuli. Ligulæ 5-6. Fl. tubulosi 6-7. (v. s. comm. à cl. Burchell.)

§. 2. *Pedunculis axillaribus à basi ramosis in pedicellos valdè elongatos 1-cephalos aut simplicibus.* — Conyzæ sp. Roxb.

3. *C. GUMMIFERUM*, foliis cuneato-lanceolatis basi integris attenuatis subpetiolatis apice serratis subrugosis junioribus suprà sparsè puberulis subtus subtomentosis, pedunculis axillaribus ferè à basi in pedicellos valdè elongatos monocephalos divisis. 5 in insulæ Sanctæ-Helenæ montanis. Conyza gummifera Roxb. in Beats. app. p. 304. *Gum-Wood-Tree* vern. angl. Arbor ramis 2-3-chotomis. Folia ad apices ramorum conferta, juniora gummifera Roxb. Pedicelli foliis ferè breviores. (v. s. comm. à cl. Burchell.)
4. *C. ROBUSTUM*, foliis cuneato-lanceolatis basi attenuatis integris subpetiolatis, apice crenato-dentatis rugosis suprà araneosis subtus tomentosis, pedunculis ex imâ basi in pedicellos elongatos 1-cephalos divisis, involucri arachnoideo. 5 in insulâ Sanctæ-Helenæ prope Longwood. Conyza robusta Roxb. in Beats.

app. p. 304. Arbor 20-pedalis. Capitula paulò majora et pauciora prioris ad quam valde accedit. (v. s. comm. à cl. Burchell et Wallich.)

5. *C. augosum*, foliis cuneatis ferè petiolatis acutis apice grossè serratis suprâ pubescentibus subtùs hirsuto-villosis reticulato-rugosis, pedunculis axillaribus 1-cephalis hirsutis simplicibus folli circ. longitudine, invol. squamis glabris linearibus. ♀ in insulâ Sanctæ-Helenæ. *Conyza rugosa* Ait. kew. 3. p. 184 non Vahl ed. 2. v. 5. p. 30 excl. syn. Willd. *Aster glutinosus* Roxb. in Beats. app. p. 303 ex Burch. in litt. (v. s. in h. Deless. olim à fratre suo in hort. Kew. lect. et nom. Aitoniano insig.)

CLXXXVI. STEIRACTIS.

Omnia *Solidaginis* sed ligulæ steriles. — Frutex arboreus Novo-Zeelandicus. Folia petiolata ovali-orbiculata glabra repanda subtùs albido-purpurea. Capitula paniculata. Invol. subcylindraceum, squamis linearibus imbricatis margine membranaceis. Ovarium teres pilosiusculum. Pappus flavicans fimbriato-pilosus. (Car. ex Rich.)

1. *S. ARBORESCENS*. ♀ in Novâ-Zeelandiâ legit Forster. *Solidago arborescens* Forst. prod. 208. Rich. fl. Nov.-Zeel. 252.

CLXXXVII. ROCHONIA.

Capitulum multiflorum radiatum, ligulis 1-serialibus homochromis foemineis sterilibus, fl. disci tubulosis hermaphr. 5-dentatis. Invol. imbricatum, squamis subscariosis linearibus 1-nerviis. Recept. alveolato-fimbrilliferum. Styli rami elongati semiteretes glabriusculi. Achænia teretiuscula subvillosa. Pappus setosus 1-serialis subscaber. — Frutices Madagascarienses ramulis foliisque subtùs canis. Folia alterna sessilia coriacea integerrima 1-nervia. Capitula pauca aut solitaria ramulos terminantia. Flores flavi. — Genus affine *Aplopappo* dicavi cl. Alex. Rochon physico Gallo qui insul. Madagascar adiit et descripsit. (Conf. Roch. voy. Par., in-8. 1791. et vol. 3. 1802.)

1. *R. CINERARIOIDES*, foliis ellipticis suprâ glabris læviusculis subtùs cano-tomentosis subreticulatis, capitulis 7-8 pedicellatis in racemum dispositis. ♀ in savannis interioribus insulæ Madagascar legit cl. Bojer. *Conyza cinerarioides* Bojer in litt. 1835. Frutex circ. orgyalis. Invol. squamæ extimæ tomentosæ, int. scariosæ flavide glabriusculæ. Folia 7-9 lin. longa, 4 lin. lata. Pedicelli tomentosi. (v. s. comm. à cl. Bojer.)

ℓ. *subreticulata*, foliis ovatis supernè reticulo vix elevato notatis, capitulis racemoso-corymbosis. ♀ cum priore legit cl. Bojer. An fortè species distincta? (v. s.)

2. *R. CUNEATA*, foliis cuneatis suprâ glabris nervulis elevatis reticulatis subtùs adpressissimè canis, capitulis terminalibus solitariis. ♀ in savannis interioribus insulæ Madagascar legit cl. Bojer. *Conyza cuneata* Bojer! in litt. 1835. Frutex octopedalis. Folia 10 lin. longa, 3 lin. lata. (v. s. comm. à cl. Bojer.)

CLXXXVIII. *APLOPAPPUS* Cass. dict. 56. p. 169. — *Diplopappus* Less. in *linnæâ* 1831. p. 110. — *Diplopappi* et *Doniæ* sp. auct.

Capitulum multiflorum radiatum, ligulis 1-serialibus foemineis (in unic. sp. nullis), fl. disci hermaphroditis 5-dentatis. Recept. planiusculum foveolatum seu alveolato-subfimbrilliferum. Invol. squamæ imbricatæ sublineares acutæ. Achænia oblonga teretiuscula aut turbinate densè sericea rarius glabra. Pappus setosus uni-aut pluriserialis subinæqualis sed setis inter se conformibus. — Herbæ aut suffruticeæ

Americana. Folia alterna integra aut dentata. Capitula terminalia. Cor. omnes flavæ, ligulis in *A. gymnocephalo* solo discoloribus albis seu purpurascensibus. — Genus distinctum à *Grindelia* pappo nec caduco nec pauciseti, ab *Inula* antheris ecaudatis, à *Diplopappo* et *Diplostephio* pappo simplici nec duplici, à *Solidagine* ligulis numerosioribus et habitu toto, verisimiliter in posterum dividendum!

Sect. I. LEIACHENIUM.

Achænia glaberrima elongata. Capitula radiata. — Habitus specierum ad *Grindelia* accedens.

1. *A. ciliatus*, caule herbaceo erecto glabro ramoso, foliis oblongis obtusis semiamplexicaulibus ciliato-serratis venoso-reticulatis glabris, invol. squamis lanceolatis, exter. squarrosis setulâ acuminatis, inter. submembranaceis erectis, achænio glaberrimo. (2) in Amer. bor. territorio Arkansano. — Planta mihi obvia proveniens ex semipibus à cl. de Pourtales loco dicto collectis et à cl. de Charpentier cultis, optimè resp. *Donia ciliata* Nutt. in Journ. acad. nat. sc. Phil. v. 2 (1821), p. 118. Hook. exot. fl. 1. t. 45, sed in meâ achænia radii certè papposa nec ut in icone Hook. calva, antheræ disci pleræque abortivæ nec fertiles. An duæ in eodem loco species valde similes? an eadem culturâ varians? (v. s. c.)
2. *A. foliosus*, caule fruticoso tereti erecto ramoso usque ad apices toto dense folioso, foliis latè sessilibus confertissimis obovato-oblongis grossè serratis, junioribus puberulis et subglutinosi, adultis reflexo-imbricatis, capitulis inter folia sessilibus solitariis, invol. squamis oblongo-linearibus obtusis mucronulatis glutinosis margine membranaceis, achænio parè piloso ferè glabro. 3 in rupestribus asperis maritimis loco dicto *Vina la Mar* prope Valparaiso Chiliensium legit ill. Bertero jannar. flor. Vulgò *Palo negro* dicitur. Species distinctissima! (v. s. comm. à cl. inv. sub n. 1035).
3. *A.?* *Hoorebekia*, caule suffruticoso tereti, ramis elongatis apice nudis 1-cephalis, foliis sparsis infer. lanceolatis grossè serratis utrinquè acuminatis semiamplexicaulibus, summis integris decurrentibus, invol. ventricosi squamis in acumen elongatum productis, ligulis amplis apice 3-dentatis invol. duplò longioribus, pappo circ. 9-10 seto. 3 in Chili. *Hoorebekia* Chiliensis Cornelissen in Mussche hort. Gand. 1817, p. 120 et in præf. ic. et desc. Capitulum floridum ex icone duos et ultra pollic. An hujus loci? †.
4. *A. ericoides*, fruticosus glabriusculus, ramis teretibus, foliis acerosi teretibus integerrimis obtusis in axillis fasciculatis, capit. pedicellis paniculato-corymbosis, invol. turbinati squamis siccis 1-nerviis scarioso-ciliatis laxè imbricatis ligulis 3-4 disco paulo longioribus, achænio glabro. 3 in Californiâ legerunt cl. de Chamisso! et Douglas! Adest inter Mexicanas herb. Haenk. *Diplopappus ericoides* Less.! in linnæa 1831, p. 117. Habitus *Aspalathi*. Ligulæ flavæ. Pappus albidus rufescens, setis pluriserialibus ferè æqualibus. Achænia oblonga basi attenuata. Capitula 10-12-flora. (v. s.)
5. *A. prunelloides*, herbaceus, caule procumbente villosulo-hirsuto foliato subsimplici, foliis crassiusculis glabris oblongo-obovatis in petiolum attenuatis serratis subpinatifidisve, capitulo solitario, invol. campanulati squamis imbricatis glabris adpressis siccis oblongo-ovatis, achænio glabro. 2 in Chili. *Diplopappus prunelloides* Pæpp. ex Less. in linnæa 1831, p. 111. Pappus flavescens. Cor. glabræ. Habitus *Brunellæ grandifloræ*.

Sect. II. BLEPHARODON.

Achænum sericeo-villosissimum breve turbinatum.

6. *A. gymnocephalus*, caulibus erectis pube duplici aliâ brevi velutinâ aliâ hispida scabridis, foliis sessilibus ligulatis velutino-scabris inciso-dentatis, dentibus in setam pellucidam elongatis, ramis floridis 1-cephalis apice subnudis.

foliis nempe summis gradatim minoribus à capitulo remotiusculis. \mathcal{U} \mathcal{J} in Mexico circa urbem leger. cl. Alaman et Berlandier (pl. exs. n. 588 et 805). Herba imà basi snffruticosa, semipedalis aut spithamæa. Ligulæ ex sicco pallidæ aut subtus rubentes, an ochroleucæ, an albidæ? Recept. foveolæ amplæ lacere acutæ. Pappus pallidè rufus. (v. s.)

7. *A. PHYLOCEPHALUS*, caulibus fruticosis a basi ramosis, ramis ad apicem usquè foliosis puberulo-scabris, foliis ligulatis subscabridis sessilibus dentatis, dentibus in setam pellucidam elongatis, capitulis inter folia suprema sessilibus bracteatis. \mathcal{J} in Mexico inter San-Fernando et Matamoros legit cl. Berlandier. sub n. 2278! Ligulæ flavæ! (v. s.)
8. *A. ? SPINULOSUS*, canescens scaber ramosissimus, foliis bipinnatifidis inciso-dentatis, lacinii linearibus rigido-mucronatis, capitulis lateralibus et terminalibus congestis, ligulis bidentatis. \mathcal{U} in campis apricis ad flum. Missouri Amer. bor. Amellus? spinulosus Pursh. fl. 2. p. 564. Sideranthus spinulosus Fras. ex Steud. Starkea? pinnata Nutt. gen. 2. p. 169. Certè non est Amellus et Chrysopsidi et Asteri potius ex Nuttall affinis. Ligulæ flavæ. An Asteres pinnatifolii nempe *A. aurantius* et *A. pinnatus* huic affines?
9. *A. LINEARIFOLIUS*, fruticulosus ramosissimus glaber, foliis linearibus acutis integerrimis glanduloso-punctatis, ramis apice nudiusculis 1-cephalis, invol. squamis lineari-lanceolatis acuminatis submembranaceis glaberrimis. \mathcal{J} in Californiâ legit cl. Douglas. Folia 1 $\frac{1}{2}$ poll. longa, 1 $\frac{1}{2}$ lin. lata. Ligulæ flavæ integræ invol. duplò longiores. Achænia sericeo-villosissima. Pappus pluriserialis niveus. (v. s. comm. ab hon. soc. hort. Lond.)

Sect. III. EUAPLOPAPPUS.

Achænia sericeo-villosa elongata. — Species omnes Chilenses.

10. *A. CHAMISSONIS*, fruticosus ramosissimus, ramis striatis junioribus glutinosis, foliis sessilibus confertis coriaceis glutinosis obovato-lanceolatis acuminatis callosio-denticulatis reticulato-venosis pilosiusculis ciliatis, capitulis solitariis, invol. turbinati squamis arcuè imbricatis siccis obtusis serrulato-fimbriatis mucronatis, achæniis sericeo-villosis. \mathcal{J} in Chili prope Taleaguano et Cuchacuchu. Diplopappus Chamissonis Less. in linnæa 1831. p. 111. Conyza punctata Willd. sp. 3. p. 1937 ex Less. Cor. radii medio pilosiusculæ, cæt. glabræ.
11. *A. CORONIFOLIUS*, fruticulosus basi ascendens glaber, foliis oblongo-obovatis cuneatis in petiolum attenuatis grossè inciso-serratis, nervis non prominulis, pedunculis subnudis 1-cephalis subsolitariis, invol. squamis angustè linearibus acuminatis subsquarrosis serrulato-subciliatis, achænio subsericeo-villoso. \mathcal{J} in Chili legit cl. de Chamisso. Diplopappus coronifolius Less. in linnæa 1831. p. 112. Rhizoma radicans horizontale. Folia coriacea. Cor. luteæ. Pappus rufus, setis subæqualibus. (v. s. comm. à cl. de Cham.)
12. *A. PULCHELLUS*, fruticosus ramosissimus glutinosus glaber aut miutissimè pubens, ramis teretibus basi foliosis apice in pedunculum 1-cephalum longè denudatis, foliis lanceolatis acuminatis basi attenuatis exsertè et acutè serratis subaveniis coriaceis, invol. campanulati squamis laxè imbricatis linearibus apice subulatis acuminatis, achænio sericeo-hirsuto. \mathcal{J} in Chili ad Rancagua locis siccis sterilibus (Bert. n. 314), prope Valparaíso (Gaudich.) et ad flumen Tinguirica (cl. Gay!). Grindelia pulchella Bertero! merc. chil. 1829. p. 646! non Dunal. An Aplopappus glutinosus Cass.? An var. *A. coronifolii* maior et latior? Pedunculi 3-4 pollic. Folia 12-15 lin. longa, 3 lin. lata. Pappus rufus. (v. s.)
13. *A. GLUTINOSUS*, fruticulosus, ramis teretibus foliosis puberulis apice in pedunculum longum nudum 1-cephalum productis, foliis coriaceis obovatis dentatis subaveniis punctatis subpuberulis glutinosis, capitulis solitariis, invol. hemisphericis squamis imbricatis linearibus glabris, achæniis sericeo-villosis. \mathcal{J} in Chili. Diplopappus glutinosus Less. in linnæa 1831. p. 113. Nees ast. 284. An Aplopappus glutinosus Cass. dict. 56. p. 169 huc aut suad. Less. ad *A. coronifolium* referendus? Pedunculus 3 poll. (v. s.)

14. *A. SCROBICULATUS*, fruticulosus, ramis procumbentibus foliosis, foliis spathulatis nitidis dentatis, pedunculis elongatis folia parva paucā angustā gerentibus, invol. squamis lanceolatis scrobiculatis cauleque glutinoso-nitidis, achænio strigoso-sericeo. ζ in Cordilleris Chilensibus. Affinis glutinoso Poepp. et Less. *Diplopappus scrobiculatus* Nees. ast. p. 284.
15. *A. BEATERRI*, caule suffruticuloso adscendente striato glabro apice aphylo, foliis subdistantibus obovatis basi cuneatis apice obtusis margine scabridociliatis et grossè serratis tenuè reticulatis, capitulo terminali erecto, invol. squamis linearibus acuminatis margine membranaceis ciliolatis, achænio sericeo-villosissimo. \mathcal{L} ζ in petrosis et rupestribus maritimis prope Valparaíso Chilensium legit cl. Bertero. *Grindelia glutinosa* Bert. ! merc. chil. 1829. maio. p. 646. non Dunal. Capitulum ferè *A. macrocephali*. Ligulæ glabræ nec ut in *A. Chamissonis* medio pilosæ. Caulis apex subviscosus glaber. (v. s. comm. à beat. et amic. Bert.).
- β ? *lanceolatus*, caule sub capitulo puberulo, foliis lanceolatis basi cuneatis apice acutiusculis intensius reticulatis, invol. squamis subpuberis et angustioribus. \mathcal{L} ζ in rupestribus maritimis prope Valparaíso legit Bertero et cum priore sub n. 1038 cum sign.? commemor. Folia ferè *A. pulchelli* et capitulum *A. Berterii* aut *macrocephali*. (v. s.)
16. *A. MACROCEPHALUS*, herbaceus, caule simplici erecto glutinoso tereti basi densè folioso apice aphylo 1-cephalo, foliis subcoriaceis spathulatis obtusis rigidè serratis margine hispidiusculis reticulato-venosis, invol. latè campanulatis squamis linearibus mucronato-acuminatis viscosis hispidiusculis, achænio sericeo-villosissimo. \mathcal{L} in Chili (Poepp.), prope Rancagua in monte La Leona (Bertero !). *Diplopappus macrocephalus* Poepp. ex Less. in linnæa 1831. p. 114. Caulis 10-pollic. Pappus flavescens. Corollæ glabræ luteæ. (v. s. comm. à cl. Bertero sub n. 315.)
17. *A. GRINDELIOIDES*, herbaceus, caulibus basi ramosis teretibus apice subaphyllis hirsutis, foliis sessilibus subsemialexicanalibus obovatis acutis acutè serratis glabriusculis, capitulis solitariis, invol. campanulatis squamis foliaceis linearibus acuminatis subpuberulis, ligulis elongatis, achæniis hirsuto-canescensibus. \mathcal{L} in Chili legit cl. Poeppig ! *Diplopappus grindelioides* Less. in linnæa 1831. p. 115. Capitulum ferè *Grindeliæ*. Pappus sordidus. (v. s. comm. à cl. Poeppig sub n. 916.)
18. *A. CHRYSANTHEMIFOLIUS*, herbaceus?, ramis striatis florigeris apice subnudis, foliis sessilibus membranaceis glabris subciliatis, infer. obovatis, super. lanceolato-linearibus mucronato-serratis et summis integerrimis, capitulis solitariis, invol. campanulatis squamis linearibus acuminatis glaberrimis, achæniis hirsuto-villosis. \mathcal{L} ? in rupibus maritimis prope Concepcion Chilensium. *Andromachia alternifolia* Kunze in Poepp. coll. exs. 1. n. 243. *Diplopappus chrysanthemifolius* Less. in linnæa 1831. p. 116. Pappus rufescens. Cor. glabræ, tubo ligularum hispidulo.
19. *A. HISPIDULUS*, fruticulosus, ramis teretibus, junioribus foliisque pube brevī confertè hispidulis, adultis glabris, foliis linearibus confertis linearibus obtusis integerrimis, ramis apice aphyllis 1-cephalis, invol. squamis linearibus acuminatis hispidulis, achæniis parè sericeo-villosis. ζ in Chili prope Coquimbo legit cl. Gaudichaud. Folia pollicem ferè longa, semilineam lata, minora in axillis glomerata. Pedunculus pollicaris. Ligulæ angustæ flavæ. Accedit ex foliis ad sect. iv, ex ligulis ad sect. v. (v. s. comm. à cl. Gaudichaud sub n. 87.)

Sect.? IV. LEUCOPSIS.

Achænia villosa aut hirsuta elongata. Pappus corollis disci longior. — Folia integra villosa-canescens. — An genus proprium?

20. *A.?* *ACUMINATUS*, herbaceus, totus sericeo-lanatus, caule erecto folioso, foliis oblongo-lanceolatis acuminatis integerrimis, infer. obovato-oblongis obtusis, capitulo solitario terminali, invol. campanulatis squamis laxè imbricatis dorso

- lanatis linearibus acuminatis, ligulis 25-30 invol. longioribus, achæniis albo-villosissimis. $\frac{1}{2}$ in Brasiliæ prov. Rio-Grande et fortè Sancti-Pauli (h. Mus. imp. Bras. n. 1041 et fortè 439?). Priori simillimus sed inv. squamis foliisque superioribus valdè acuminatis facilè differt. In specim. prov. Sancti-Pauli pars inferior plantæ glabrescit. (v. s. in h. Mus. reg. Par.)
21. A. ? *SERICUS*, herbaceus totus sericeo-lanatus, caule adscendente toto folioso, foliis oblongis acutis basi attenuatis integerrimis, capitulo solitario terminali, invol. campanulati squamis laxè imbricatis siccis linearibus obtusiusculis, ext. subvillosis, cæst. glabris siccis, ligulis vix invol. longioribus, achæniis elongatis hirsutis. $\frac{1}{2}$ in pascuis sabulosis ad Quintero Chilensium legit cl. Bertero (pl. exs. n. 1411) et ad Talcaguano (Cham.). *Diplopappus sericeus* Less. in linn. 1831. p. 110. Cor. flavæ. Folia pollicem longa, 2-3 lin. lata. Pappus pluriserialis rufescens setis ext. brevioribus. An *Noticastro* affinis? (v. s.)
22. A. ? *CANESCENS*, totus adpressè sericeo-villosus canescens, caulibus fruticulosus à collo plurimis apice subramosis, foliis oblongo-linearibus basi attenuatis integerrimis mucronato-acutis infernè setis rigidis paucis ciliatis, ramulis brevibus 1-cephalis, inv. campanulati squamis linearibus acutis subcanescentibus, ligulis paucis invol. paulò longioribus, achæniis villosis pappo dimidio brevioribus. $\frac{1}{2}$ in Mexicanæ prov. Texas districtibus orientalibus nov. et dec. flor. legit cl. Berlandier (pl. exs. n. 1877). Affinis A. sericeo sed capitula triplò 4-plòve minora, pappus pallidè rufescens, achænia etiam ratione capituli multò breviora etc. (v. s. comm. à cl. Berlandier.)
23. A. ? *DIFFUSUS*, suffruticosus sparsè pubero-villosus, ramis adscendentibus foliosis, foliis spathulato seu oblongo-linearibus mucronato-acuminatis subintegerrimisque scabris, capitulis solitariis, invol. squamis lanceolatis laxis, achænio adpressè villosis. $\frac{1}{2}$ in Monte Video (Pers.), et Rio-Grande (h. Mus. imp. Bras. n. 1009, 1008, 1010, 1022, 1033!). *Erigeron diffusus* Pers. ench. 2. p. 431. *Onoseris montevidensis* Spreng. syst. 3. p. 502. Affinis A. sericeo sed pube totius plantæ sparsior, caulis fruticulosus ramosior, folia scabra margine scaberrima, Less. in linn. 1831. p. 110. (v. s. in h. Mus. reg. Par.)

Sect. VI. ? PYROCHÆTA.

Capitulum radiatum, ligulis 1-serialibus angustissimè linearibus neutris seu styliferis sterilibus, fl. disci tubulosos brevissimè 5-dentatis herm. fertilibus. Invol. imbricatum, squamis linearibus pluriserialibus intimis disco circ. æqualibus. Achænia radii abortiva, disci turbinata sericeo-villosissima. Pappus radii setis paucissimis, disci multisetis scaber rigidulus. — Herba Mexicana annua erecta. An genus proprium?

24. A. ? *HÆNKEI*, radice exili, caule herbaceo simplici aut parcè ramoso glabro, foliis oblongo-cuneatis apice serratis paucis dentatis junioribus subtus arenaosis demùm glabris. ① inter plantas Regiomontanas herb. Hænkeani ab ill. de Sternberg ad studium missi conservatur. Herba pedalis. Folia remotiuscula pollicem longa, 3 lin. lata. (v. s. in h. suprà dicto.)

Sect. V. ? INULOPSIS.

Achænia villosa breviteretiuscula. Ligulæ lineares numerosissimæ. — Herbæ Mexicanæ et Brasilienses, habitu Inulæ. Fortè genus proprium? imò species duæ in genera duò fortè divellendæ?

25. A. ? *STOLONIFERUS*, caulibus à collo plurimis debilibus parcè foliosis basi hispidulis, aliis sterilibus decumbentibus, aliis 1-cephalis apice villosito-tomentosis et glanduloso-pilosis, foliis radicalibus obovatis hirsutissimis subdentatis, secus caules floridos lanceolatis acutis, secus caules steriles obovato-oblongis subobtusis, invol. squamis linearibus acuminatissimis disco longioribus hispidis. $\frac{1}{2}$ in Mexici summo monte San-Felipe supra locum ubi ex structurâ glaciæ apotheca legit cl. Andrieux. Inula stolonifera Gay! in Andr. coll. exs. n. 322. Ligulæ numerosissimæ 1-seriales lineares. Cor. tubulosæ oo. Anthere basi nudæ. Invol. subpurpurascens. Flores intimi abortu masculi. Achænia juniora villosa, adulta ign. (v. s.)

- β. *puberus*, foliis radicalibus lanceolatis utrinque attenuatis pubescenti-subhirtis. ♀ in Mexico legit cl. Alaman. Var. α simillima. Folia multò minùs pilosa et radicalia lanceolata. Receptaculum alveolato-fimbrilliferum. (v. s.)
26. A. P. *SCAPOSUS*, acaulis, foliis radicalibus obovato-oblongis basi angustato-cuneatis obtusis apice calloso-denticulatis pilosiusculis, scapis glabris folia multò superantibus 1-cephalis, invol. squamis linearibus acutis vix subpuberulis margine submembranaceis. ♀ in Brasiliæ prov. Minarum Generalium ad Serro-Frio (Vanthier! pl. exs. n. 315) et in prov. Sancti-Pauli (h. Mns. imp. Bras. n. 436!). Folia radicalia 2-2½ poll. longa, vix pollicem lata. Scapi 7-10 pollic. Cor. omnes flavæ glabræ. Pappus rufus, nunc 1-serialis nunc varius setulas parcas basi gerens. Achænia villosa. Medius inter Aplopappum et Chrysopsidem. Habitu ab utroque alienus! (v. s.)

Sect. VII. APLDISCUS.

Capitula discoidea nec radiata. Achænia plùs minùs villosa. Cor. fauce dilatata. — Sectio media inter Aplopappum et Pyrrocoma.

27. A. *DISCOIDEUS*, caule fruticuloso ramoso, ramis subangulatis apice viscosulis subhirtellis, foliis oblongis basi attenuatis dentibus grossis utrinque 3-4 acutis (interdum aristatis) dentatis utrinque punctatis glabris subviscosis, corymbis ramos terminantibus 12-15-cephalis subcompositis, capitulis eradiatis, invol. squamis glabris imbricatis acutis disco brevioribus. ♀ in Mexici montanis circa urbem leger. cl. Née, Alaman, Berlandier, Mairet, ad Tlapujahua cl. Keerl. Semper recepi sub nom. Baccharidis seu Molinæ sed flor. omnes bisexuales, nec dioici; discoidei ut in sect. Pyrochætâ. (v. s.)
28. A. *RAMULOSUS*, caule fruticuloso ramoso, ramulis brevibus striato-angulatis, foliis oblongo-linearibus subcuneatis vix apice denticulatis mucrone acutis, utrinque punctatis subviscosis, capitulis ad apices ramulorum foliosorum solitariis subracemosis eradiatis, invol. squamis imbricatis acutis glabris disco brevioribus. ♀ in Mexico ad Tlapujahua legit et Keerl. Cor. ex sicco flavæ, antheræ fortè rubescentes. Styli rami longo-appendiculati hispidi. (v. s. in h. Mart.)

CLXXXIX. PYRROCOMA Hook. fl. bor. am. 1. p. 306. t. 107.

Capitula multiflora homogama, floribus hermaphr. tubulosis 5-dentatis, dentibus brevibus erectis, fauce non ampliata. Invol. squamæ plures laxè imbricatæ plùs minùs squarrosæ. Recept. epaleaceum. Antheræ subinclusæ. Styli rami lineares basi teretes nudi apice extùs pilosi. Achænium oblongum teretiusculum glabrum. Pappus 1-serialis, setis scabris fulvis. — Herbæ (aut suffrutices) Americanæ pedales erectæ simplices. Folia alterna lanceolata seu oblonga coriacea subpubescentia parcè dentato- seu spinoso-serrata sæpè punctata. Capitula lutea.

Sect. I. EUPYRROCOMA.

Invol. squamæ squarrosæ apice subrecurvæ. Folia punctata. — Species boreali-Americana.

§. 1. Bracteosæ, involucri foliis spinoso-dentatis ad basin cincta, squamis vix squarrosis.

1. P. *CARTHAMOIDES* (Hook. l. c.), capitulo solitario terminali amplè bracteato, foliis oblongo-lanceolatis pauci-serratis, infer. in petiolum attenuatis. — ad oram boreali-occid. Americæ legit cl. Douglas. Folia 4-6 poll. longa, 7-9 lin. lata. (v. s. comm. ab hon. Soc. hort. Lond.)

§. 2. Squarrosæ, capitula plurima racemoso-spicata subbracteata; invol. squamæ squarrosæ.

2. P. *GRINDELIOIDES*, capitulis ad axillas supremas sessilibus racemoso-spicatis,

invol. squamis squarrosis obtusiusculis, foliis obovatis sessilibus acutè serratis. § 4 in California legit cl. Douglas et inter Regiomontanas herb. Hænke etiam servatur. Caulis hirtio-pubescent. Folia punctata fortè in vivo pellucido-punctata, pollicem longa, 6-7 lin. lata. Recept. alveolatum. Pappus in specim. Douglasii pallescens, in Hænke intensè rufus. (v. s.)

Sect. II? CHROMOCHÆTA.

Invol. squamis adpressis imbricata non squarrosa. Folia impunctata. — Suffrutices Chilenses. Habitus Aplopappi sed in specim. duodenis ligulas nullas vidi. An genus proprium?

3. *P. PARVIFOLIA*, caule fruticoso erecto tereti paniculato-ramoso glabrinsculo, foliis cuneatis sessilibus obtusis apice grossè et acutè dentatis rigidulis glabrinsculis, ramis novellis axillaribus densè foliosis, capitulis subsessilibus, invol. squamis linearibus acutis, achænio glabro. § in Cordilleris Chilensibus legit cl. Hænke. Ligulæ in specim. nullæ sed verisim. deciduæ. Pappus rufus. Folia 4-6 lin. longa, 2-3 lata. (v. s. in herb. Hænke à cl. Sternb. miss.)
4. *P. ANGUSTIFOLIA*, caule fruticoso erecto tereti glabro paniculato-ramoso, foliis sessilibus oblongis acutis rigidis glabris utrinquè dentes 2-3 exsertos grossos gerentibus, capitulis subsessilibus, invol. squamis oblongo-linearibus obtusis, achænio glabro. § in Cordilleris Chilensibus legit cl. Hænke. Valde affinis *P. parvifoliæ* et fortè varietas? Differt glabritie absolutà, foliis distantibus longioribus angustis, invol. squamis obtusis. (v. s. in h. Hænke ab ill. de Sternberg miss.)

CXC. *LESSINGIA* Cham. in *linnæa* 1829. p. 203. t. 2. f. 2. Less. 388. Cass. opusc. 3. p. 96.

Capitulum multiflorum homogamum, fl. radii majoribus profundè 5-lobis obconicis radiatiformibus, fl. disci cylindrico-tubulosis breviter 5-dentatis. Invol. imbricatum. Recept. nudum. Genitalia inclusa. Stylus ramis oblongis extus hispidis. Achæmium compressum erostre sericeo-villosum. Pappus 1-serialis, setis rufis scabridis rigidulis. — Herba Californica procumbens ramosa demum glabrata. Folia alterna crassiuscula, ima pinnatifida, super. apice dentata. Capitula solitaria terminalia et ramorum elongatione alaria. Flores flavi. — Facies omninò et stylus ipse potiùs meo sensu Asterearum quàm Senecionearum seu Tagetinearum.

1. *L. GERMANORUM* (Cham. l. c.). ① in arenosis ad portum Sancti-Francisci novæ Californiæ legit cl. de Chamisso. An nomen specificum servandum et species potiùs Californica dicenda? (v. s. comm. à cl. inv.)

CXCI. *LINOSYRIS* Lob. hist. 223. — *Crinitaria* Less. syn. 195. — *Crinitaria* et *Linosyris* Cass. dict. 37. p. 475 et 476. — *Crinita* Manch. meth. p. 578. non *Houtt.* — *Chrysocomæ* sp. Linn.

Capitulum multiflorum homogamum discoideum. Recept. planum alveolatum, alv. marginibus carnosissimis dentatis. Invol. pauciseriale, squamis foliaceis. Cor. limbus profundè 5-fidus. Anth. exsertæ. Stigmata exserta Asterearum. Achæmium oblongum erostre compressum sericeum. Pappus pilosus biserialis, setis scabris. — Herba erecta perennes Europæ et Asiaticæ. Folia alterna conferta linearia aut oblonga subintegerrima. Capitula ad apices ramorum solitaria sæpè in corymbos disposita. Flores lutei. — Genera duo Cassiniana vix sectiones formant. Nomen *Linosyris* elegi quia vetustius et quòd alterum vix intelligibile.

* *Herbacæ.*

1. *L. vulgaris* (Cass. ex Less. syn. 195), foliis linearibus glabris impunctatis, capitulis corymbosis, invol. squamis appendiculatis apice patulis squarrosis. $\frac{1}{2}$ in Europæ mediæ et australis et imò Tauriæ et Oelandiæ apricis rupestribus et argilosis. *Chrysocoma Linosyris* Linn. sp. 1178. DC. fl. fr. 4. p. 141. All. ped. t. 11. f. 2. Schkuhr handb. t. 238. *Linosyris foliosa* Cass. ex Desf. cat. h. Par. p. 196. *Crinitaria Linosyris* Less. syn. 195. *Linosyris nuperorum* Lob. hist. 223. *Aster Linosyris* Bernh. ex Steud. nom. phan. 193. *Erigeron Linosyris* Clairv. man. herb. val. 243. (v. v. et s.)
 β . *minor* (Wallroth! in litt.), caule 3-5 pollicari. $\frac{1}{2}$ in siccis Thuringiæ, Galliæ. C. *Linosyris tripollicaris* Vill. dauph. 3. p. 188. DC. fl. fr. n. 3130. Culta dimensiones majores mox acquirit (Savi. pug. p. 11). (v. s.)
 γ . *patula*, foliis patentibus vel reflexis. $\frac{1}{2}$ in collibus calcareis Alsatiae super. (Nestl.) et Jurassi Neocomensis (Chaill.). — An in paludosis Etruriæ circa Stagno et Liburnum si *Chrys. palustris* Savi pug. p. 11. et *Chrys. Liburnica* Spreng. syst. 3. p. 475 huc referendæ? (v. s.)
 δ ? *verticalis*, foliis lanceolatis verticalibus. $\frac{1}{2}$ ad rupium fissuras in Hispaniæ regnis Murcico, Granatensi et Valentino. *Chrysocoma verticalis* Lag. nov. gen. p. 15. n. 330.
 2. *L. punctata* (Cass. dict. 37. p. 476. sub *Crinitaria*), foliis linearibus aut lineari-lanceolatis trinerviis punctatis subscabris, capitulis corymbosis, invol. laxo. $\frac{1}{2}$ in campis apricis siccis aut subsalsis Sibiriae ad Irtim (Patr.), Ural (Fisch.), in Phrygiâ (Parol.), in campis ad Wolgam (Bieb. Stev.). *Aster acris* Gmel. itin. 1. p. 172. t. 38. f. 2. *Chrysocoma biflora* Linn. sp. 1178 (nomen omn. decipiens). *Chrysocoma dracunculoides* Lam. dict. 2. p. 192. Bieb. taur. 2. p. 288. *Crinita punctata* Mœnch. meth. 578. *Crinitaria biflora* Less. syn. 195. — Gmel. Sib. 2. t. 82. f. 1. Variat foliis angustissimis 1-nerviis et latioribus 3-nerviis. Cum *Astere* confudit Linnæus quando dixit variare cum et sine radio. (v. v. c. et sp.)
 3. *L. villosa*, foliis lanceolatis adpressè villosis, capitulis paucis subcorymbosis, involucri squamis adpressis margine scariosis. $\frac{1}{2}$ in apricis Hungariæ, Caucaei, Sibiriae. *Chrysocoma villosa* Linn. sp. 1178. Waldst. et Kit. pl. hung. 1. t. 58. *Conyza oleæfolia* Lam. dict. 2. p. 86. — Gmel. Sib. 2. t. 82. f. 2. (v. s.)
 4. *L. divaricata*, herbacea incana, foliis infer. lanceolatis trinerviis, super. lineari-subulatis diminutis, ramis 1-cephalis squamatis virgatis, invol. laxi incani squamis acutis. $\frac{1}{2}$ in campis siccis circa Astracan ex Gmel. *Chrysocoma divaricata* Fisch. cat. gor. 1816. p. 37. Bieb. suppl. 563. — Gmel. Sib. 2. p. 193. t. 82. f. 3. Minus pubescenti-cana quam *L. villosa*.
- ** *Fruticosæ, recept. sæpius punctato nec alveolato.* (Pleræque minùs notæ.)
5. *L. CANDICANS*, caule suffrutescente diffuso, foliis linearibus sericeo-pubescentibus margine replicatis, capitulis paniculis confertis 5-9-floris, invol. squamis adpressis apice coloratis. $\frac{1}{2}$ in Ægypto propè Alexandriam. *Chrysocoma candicans* Delil. fl. eg. 127. t. 46. f. 2. Invol. cylindricum nec punctatum nec ut in prioribus foveolatum.
 6. *L. HASTATA*, fruticosa, foliis oblongis sessilibus ad medium lobo horizontali hastatis incanis, capitulis solitariis, invol. imbricati squamis acutis. $\frac{1}{2}$ in Arabia felici. *Chrysocoma spatulata* Forsk. descr. æg.-ar. 147. *Stæhelina hastata* Vahl symb. 1. p. 70. *Pappus biserialis* ex Less. syn. 195.
 7. *L. MONTANA*, fruticosa, foliis oblongis villosis integerrimis, capitulis solitariis, invol. oblongis squamis ext. obtusis, int. acutis apice patulis. $\frac{1}{2}$ in monte Horeb Arabiæ et in Palestinâ si pl. n. 440 herb. Bové huc ritè referenda. *Chrysocoma montana* Vahl symb. 1. p. 70. *Pappus rufus biserialis* ex Less. l. c. (v. s.?)
 8. *L. UNIFLORA*, fruticosa, ramis glabris erectis, foliis linearibus complicatis

acutis, pedunculis 1-cephalis, invol. imbricato. ζ in *Egypto*. *Chrysocoma uniflora* Spreng. nov. prov. 1818. p. 11. Recept. nudum.

9. *L. GLABRATA* (Lindl. obs. mss. 1835, sub *Crinitaria*), foliis linearibus obtusissimis punctatis glabris, capitulis corymbosis, invol. obconici squamis ext. pubescentibus, inter. dilatatis obovatis glabris. \mathcal{Z} in *Sibiria Altaica deserto Sogoro-Kirghisico*. *Aster dracunculoides discoidens minor* Ledeb. fl. alt. 4. p. 98. Species mihi ignota et ex cl. Lindley admissa.

XCXII. *CHRYSOCOMA* Cass. dict. 37. p. 477. Less. syn. 195. — *Chrysocoma* sp. Linn. Thunb. — *Chrysocoma* et *Crinita* sp. Mœnch.

Capitulum homogamum discoideum multiflorum. Recept. nudum subaeolatum. Invol. campanulati floribus brevioris squamæ oblongo-lanceolatæ imbricatæ. Achænia lateraliter plano-compressa erostria vix subhispidula. Pappus setaceo-pilosus 1-serialis. — Frutices Capenses. Folia linearia sparsa sæpius integra. Rami apice plus minus denudati. Capitula subglobosa. Flores lutei. — Species recognoscendæ!

§. 1. *Fruticosæ, foliis linearibus, capitulis subsolitariis.*

1. *C. COMA-AUREA* (Linn. sp. 1178), foliis sparsis linearibus patulis rectis aut subrecurvis ramisque glabris aut junioribus subpubescentibus. ζ ad Cap. Bonæ-Spei frequens. *C. patula* Berg. cap. 234? *Crinita linearifolia* Mœnch suppl. 237. Pluk. alm. t. 327. f. 2. Volk. fl. norib. t. 148. *C. aurea* Thunb. fl. cap. 626. Variat ramis plus minus divergentibus, omnino glabris aut apice summo pubescentibus; foliis erectiusculis patulisve, glabris aut jun. pubescentibus; ramis apice longè aut breviter denudatis; capitulis erectis aut cernuis 30-40-floris, sed hæ variationes adeo inter se commixtæ apparent ut vix in statu scientiæ præsentî seq. var. sint distinctæ. (v. v. c. et sp.)
 - β. *patula*, ramis divaricatis. ζ Chr. *patula* Linn. mant. 28. Thunb. fl. cap. 626. (v. s.)
 - γ. *cernua*, capitulis nubilibus cernuis ζ Chr. *cernua* Linn. sp. 1177. Thunb. fl. cap. 626. — Comm. hort. Amst. 2. p. 89. t. 45. (v. s.)
2. *C. CILIATA* (Linn. sp. 1177), foliis sparsis linearibus patulis rectis serrulato-ciliatis, ramis pubescentibus vix apice nudis ζ ad Cap. Bonæ-Spei. Chr. *ciliaris* Willd. sp. 3. p. 1789. non Link. — Icon Comm. hort. 2. t. 48 habitum exhibet, sed cilia foliorum nec in descr. nec in icone indicantur, et folia (in icone integra) in descr. apice trifida dicuntur. (v. s.)
3. *C. TENUIFOLIA* (Berg. cap. 235), foliis sparsis lineari-subulatis patulis brevibus ramulisque glabris aut subpuberulis, capitulis breviter pedicellatis circ. 30-floris. ζ ad Cap. Bonæ-Spei in montibus ad Los-Tafelberg (Drege!), in Worcester, Swellendam, Uitenhage (Eckl. et Zeyh.). Chr. *microphylla* Thunb. fl. cap. 626? Olim recepi sine nom. à cl. Thunb. Folia 1-3 lin. longa, juniora erectiuscula dein patula, subacerosa. Rami folia minima ferè usque ad capitulum gerentes numerosi corymbosi. (v. s. sp.)
4. *C. MICROCEPHALA*, glabra, foliis linearibus subpatulis acutis, ramis floriferis fastigiatis ad capitulum usque foliatis, capitulis circ. 20-floris. \mathcal{Z} ad Cap. Bonæ-Spei (Boiss. herb.), in distr. Uitenhage, Worcester (Eckl. et Zeyh.), ad Nieuwewelde, Breederivier, Zwarkoprivier (Drege!). Similis *C. cernua* sed capitula fastigiata erecta dimidiò minora; differt à *C. ciliata* et *C. tenuifolia* foliis glaberrimis et à *C. pedunculata* ramis apice non denudatis. (v. s.)
5. *C. PEDUNCULARIS*, glabra, foliis linearibus erectis sparsis oppositisve, ramis floriferis vix basi foliosis elongatis nudis, capitulis circ. 20-floris. ζ ad Cap. Bonæ-Spei in Saldanhabay (Ecklon in h. Dunant), in Namaqualand (Eckl. et Zeyh.), ad Stormberg et Camisberge (Drege!). Species distinctissima. Folia 3-4 lin. longa. Pedunculi uni- vel bipollicares. Capitula dimidiò minora quàm priorum. (v. s.)

6. *C. LONGIFOLIA*, glabra, foliis angustè linearibus elongatis erectis integerrimis, capitulis pedicellatis corymbosis circ. 20-floris. \S in Africâ australi ad Olifant-rivier Camisberge legit cl. Drege! Folia 9-15 lin. longa semilineam lata. Pedicelli circ. foliorum longitudine. (v. s.)
ℓ. patula, foliis patulis, corymbis minùs confertis. \S ad Kaus et Clan-William legit cl. Drege! (v. s.)
7. *C. OBLONGIFOLIA*, fruticosa nana, foliis elliptico-oblongis rigidè ciliatis utrinquè ramisque glabris, ramis florif. apice aphyllis. invol. squamis linearibus acuminatis margine subscariosis. \S in Africâ Capensi legit cl. Drege! Chr. ciliata E. Mey. \dagger in Drege. coll. Melius quàm vera *C. ciliata* nomen meretur sed nec descr. nec icon respondent. (v. s.)
8. *C. TRIDENTATA*, glabra, foliis crassiusculis obtusè trilobis seu grossè tridentatis, paucissimis linearibus integris, capitulis pedicellatis 30-40-floris. \S in Africâ Capensis deserto Carro et ad Zwarteberge legit cl. Drege! Species distinctissima. Capitula *C. Comæ-aureæ* ferè majora. Achænia nigra minutissimè puberula. Folia variant 3-5 lin. longa. (v. s.)
9. *C. PINNATIFIDA*, fruticosa glabra, foliis sessilibus oblongo-linearibus crassiusculis pinnatifidis, lobis utrinquè 2-3 integerrimis obtusis, marginibus subrevolutis, ramulis apice subnullis 1-cephalis. \S in Africâ australi ad Zwarteberge legit cl. Drege! Caulis circ. bipedalis simplex apice breviter subpaniculatus. Invol. subpurpurentia. Flores pallidè flavi. Achænia plano-compressa sparse et minutissimè puberula (v. s. comm. à cl. inv.)

\S . 2. *Herbaceæ, foliis non linearibus, capitulis corymboso-paniculatis.*

10. *C. UNDULATA* (Thunb. fl. cap. 627), herbacea, caule pubescente, foliis cordatis lanceolatis undulatis, capitulis paniculato-fastigiatis. — Ad Cap. Bonæ-Spei. Chrysocomæ congener ex herb. Thunb. teste Less. syn. 195.
11. *C. DECURRENS*, herbacea subscabrida, foliis elongatis pinnatifidis longè secus caulem decurrentibus, corymbo paniculato-fastigiato, capitulis ad apices ramulorum congestis. — in Africâ australi ad Willbergen alt. 4-500 ped. legit cl. Drege! Herba bipedalis à genere habitu diversissima. Folia ferè Conyzæ pinnatilobatæ. Flores flavi. Capitula circ. 20-flora. Achænia compressa puberula. Pappus 1-serialis multisetus. Genitalia generis. (v. s.)
12. *C. PTEROCAULA*, herbacea erecta subsimplex hirtello-pubescent, foliis elongatis petiolatis in alas integras secus caulem longè productis, infer. lyratis, lobis lateralibus 3-4-jugis ovatis obtusis, terminali lanceolato basi lobato seu sinuato, super. lineari-lanceolatis acuminatis plus minùs pinnatilobatis summis integerrimis, corymbo paniculato laxo. — in Africâ australi legit cl. Drege! Prioribus valdè affinis et cum eis sect. naturalem (à Chrysocomis geminis habitu distinctissimam) efficiens. (v. s.)

\dagger *Non satis notæ.*

13. *C. NIVEA* (Willd. sp. 3. p. 1790), fruticosa, foliis lineari-lanceolatis tomentosis planis, corymbis terminalibus sessilibus. \S ad Cap. Bonæ-Spei. An fortè *Senecionis* sp.?
14. *C. TOMENTOSA* (Linn. syst. veg. 615. Willd. sp. 3. p. 1790), suffruticosa, foliis linearibus utrinquè tomentosis, ramis tomentosis 1-cephalis. \S patr. et cæt. car. ign. \dagger .
15. *C. ? PURPUREA* (Forst. prodr. n. 286. Willd. sp. 3. p. 1791), herbacea, foliis elliptico-lanceolatis subserratis pubescentibus, paniculâ terminali corymbosâ. — in insulâ Tannâ. \dagger .

$\dagger\dagger$ *Species exclusæ.*

C. ACAULIS Walt. = *Vernonia oligophylla*.

- C. ADPRESSA Fl. Flum. = *Vernonia monticola*.
- C. AMARA Schum. = *Decaneurum Senegalense*.
- C. ANGUSTIFOLIA Walt. = *Vernonia angustifolia*.
- C. AUREA Bojer. = *Nidorella Chrysocoma*.
- C. BIFLORA Linn. = *Linosyris punctata*.
- C. BIFLORA Lam. = *Galatella dracunculoides* β.
- C. CAMPHORATA Rob. et Cast. = *Jasonia glutinosa*.
- C. CANDICANS Delil. = *Linosyris candicans*.
- C. CAPILLACEA Michx. = *Eupatorium foeniculaceum*.
- C. CILIARIS Link. = *Agathæa Ecklonis*.
- C. CINEREA Labill. = *Ozothamnus cinereus*.
- C. CORONOPHOLIA Michx. = *Eupatorium coronopifolium*.
- C. DENTICULATA Jacq. = *Plagiis virgatus*.
- C. DICHOTOMA Jacq. = *Jasonia dichotoma*.
- C. DIVARICATA Fisch. = *Linosyris divaricata*.
- C. DRACUNCULOIDES Pursh. = *Bigelowia dracunculoides*.
- C. DRACUNCULOIDES Lam. = *Linosyris punctata*.
- C. FERRUGINEA Spreng. = *Ozothamnus ferrugineus*.
- C. FOETIDA Lam. = *Erigeron foetidum*.
- C. FRUTICOSA Stend. = *Vernonia fruticosa*.
- C. GIGANTEA Walt. = *Vernonia angustifolia*.
- C. GRAMINIFOLIA Linn. = *Solidago graminifolia*.
- C. GRAVEOLENS Nutt. = *Bigelowia dracunculoides*.
- C. HETEROPHYLLA Raf. = ? *Baccharidea ex auct.*
- C. LAPPULIFOLIA Bojer. = *Nidorella mucronata*.
- C. LIBURNICA Spreng. = *Linosyris vulgaris* γ patula.
- C. LINOSYRIS Linn. = *Linosyris vulgaris*.
- C. LINOSYRIS Asso. = *Galatella rigida*.
- C. OLIGOPHYLLA Fl. Flum. = *Vernonia cephalotes*.
- C. MONTANA Vahl. = *Linosyris montana*.
- C. MUCRONATA Forsk. = *Iphyona spinosa*.
- C. NAUSEOSA Pursh. = *Bigelowia Missouriensis*.
- C. NOVEBORACENSIS Walt. = *Vernonia noveboracensis*.
- C. NUDATA Michx. = *Bigelowia nudata*.
- C. ODORATISSIMA Rensch. = *Liatris odoratissima*.
- C. OPPOSITIFOLIA Linn. = *Pteronia divaricata*.
- C. PANICULATA Gmel. = *Liatris paniculata*.
- C. PAUCIFLORA Fl. Flum. = *Ageratum melissæfolium*.
- C. PEDUNCULATA Fl. Flum. = *Vernonia pedunculata*.
- C. PILOSA Gmel. = *Liatris hirsutissima*.
- C. PINIFOLIA Spr. = *Ozothamnus pinifolius*.
- C. PRÆALTA Desf. = *Vernonia præalta*.
- C. PUMILA Fl. Flum. = *Vernonia grandiflora*.
- C. PURPUREA Raf. non Forst. = *Liatris seu Vernoniæ sp.*
- C. RETICULATA Labill. = *Ozothamnus reticulatus*.
- C. ROSMARINIFOLIA Spr. = *Ozothamnus*.
- C. SAXATILIS DC. = *Jasonia glutinosa*.

- C. SCABRA Thunb. = *Felicia reflexa*.
 C. SCOPARIA Linn. = *Baccharis scoparia*.
 C. SERICEA Linn. f. = *Schizogyne sericea*.
 C. SPATHULATA Forak. = *Linosyris hastata*.
 C. SPHACELATA Mœnch. = *Senecio reclinatus*.
 C. SPICATA Forsk. = *Trichogyne cauliflora*.
 C. SPINOSA Delil. = *Iphyona spinosa*.
 C. SQUAMATA Labill. = *Leptorhynchos squamatus*.
 C. TOMENTOSA Ell. = *Vernonia Elliottii*.
 C. TOMENTOSA Walt. = *Vernonia noveboracensis*.
 C. UNIFLORA Spr. = *Linosyris uniflora*.
 C. VILLOSA Linn. = *Linosyris villosa*.
 C. VERTICALIS Lag. = *Linosyris vulgaris* ♂ *verticalis*.
 C. VERTICILLATA Fl. Flum. = *Vernonia capitellata*.
 C. VIRGATA Nutt. = *Bigelovia virgata*.
 C. VIOLACEA Schum. = *Vernonia cinerea*.
 C. VIRGINIANA Hort. Par. = *Solidago graminifolia*.

N. B. Chrysocomæ omnes Floræ Fluminensis è genere sunt excludendæ, aliæ ad Vernoniæ, aliæ ad Baccharidem, paucæ ad Eupatorium aut ad Porophyllum referendæ et ad calcem generum indicatæ.

CXCIII. PTERONIA *Lin. gen. n. 937. Thunb. fl. cap. 629.* — *Pterophora* et *Scepinia Neck.* — *Pterophorus*, *Scepinia* et *Pachyderis Cass.* — *Pterophorus*, *Pteronia*, *Henanthus* et *Pachyderis Less.*

Capitulum sæpius multiflorum discoideum homogamum, rariùs trimò uniflorum. Recept. alveolatum, alv. marginibus nunc brevibus nunc in fimbrillas plus minus productis. Invol. squamæ pluriseriales imbricatæ adpressæ. Achenia compressa aut turbinata, glabra aut villosa, apice rariùs in collum breve attenuata. Pappus setaceo-paleaceus multiserialis, setis crassiusculis densè et brevè barbellatis sæpè (præsertim in sect. II) in cupulam basi concretis. — Fruticul: Capenses rigidi ramosi. Folia alterna aut sæpius opposita, integerrima aut in unicâ specie subdentata, glabra aut hirsuta, sæpè ciliata. Capitula terminalia solitaria aut corymbosa. Flores lutei rariùs purpurei. — Genera Lessingiana ut sectiones retinui ob similitudinem habitûs specierum generalis et transitiones formarum fructuum plurimas.

Sect. I. SCEPINIA. — *Scepinia Neck. elem. n. 140. Cass. dict. 37. p. 475. 48. p. 44.* — *Pteronia* et *Henanthus Less. syn. 195.*

Capitulum multi- aut pauciflorum. Recept. alveolatum brevè fimbrilliferum. Achaenium teres aut subcompressum, villosum, in collum minimè attenuatum. — Folia alterna aut opposita.

§. I. *Foliis alternis sparsis aut subverticillatis.*

1. P. ASPALATHA, glabriuscula, foliis sparsis sessilibus ellipticis oblongisve enerviis subcarnosis acutis inferioribus dente 1-2 utrinquè hinc indè notatis, capitulis brevè pedunculatis terminalibus multifloris, invol. squamis lineari-lanceolatis acuminatis cum ramulis aut foliis supremis brevissimè puberulis. ♂ ad *Cap. Bonæ-Spei* inter flum. Karroc et Dwaal legit el. Burchell (n. 1427!) ad

Nieuweweld (Drege!). Rami striati vetustis fol. cicatric. ut in *Aspalathis tuberculatis*. Fol. summa parva integra, inf. apice interdum subrecurva. (v. s.)

2. *P. GLAUCESCENS*, glabra, foliis sparsis linearibus glaucis integerrimis, capitulis ramos terminantibus subsessilibus 10-12-floris, invol. squamis linearibus obtusis. ζ in Africa australi ad Nieuwewelde alt. 2-300 ped. legit cl. Drege! Rami elongati albidii inermes. Pappus pallide rufescens. Achænia villosa. Styli rami crassi valde hispidi. (v. s.)

β . *rigidor*, ramis abbreviatis, adultis sæpè spinescentibus. ζ ad Zw. Storm et Willberge alt. 4-500 ped. (v. s.)

3. *P. BACCHAROIDES* (Less. syn. 196), glabra, foliis sparsis trigonis sessilibus subtus sulco longitudinali notatis imbricato-patentibus, capitulis terminalibus solitariis. ζ ad Cap. Bonæ-Spei. Osteospermum teretifolium Thunb. fl. cap. 713. Osteosp. trigonum Spreng. syst. 3. p. 628. Similis *Pterophoro camphorato* (Less.). Rami glabri striati cinerei filiformes. (Th.)

4. *P. ACEROSA*, glabra, foliis sparsis subulatis acutis rigidulis, capitulis terminalibus solitariis sessilibus, invol. squamis oblongo-linearibus integris, interioribus paulò longioribus acutis. ζ ad Cap. Bonæ-Spei. Folia conferta 8-9 lin. longa. Achæn. villosam. Pappus subrufescens. Numerus florum variat in speciminibus cæt. simillimis:

a. triflora, capitulis oblongis trifloris. ζ non longè à flum. Gauritz (Burchell n. 4721 et 5009), ad sinum Algoa (Forb.!), in distr. Uitenhage, Albany, Swellendam (Eckl. et Zeyh.), ad Gamtos et Koem-rivier (Drege!). (v. s.)

β . *quinqueflora*, capitulia oblongis circ. 5-floris. ζ in distr. Albany Eckl. et Zeyh. (v. s.)

γ .? *multiflora*, capitulis 8-12-floris subsolitariis majoribus. ζ in colonie distr. orient. Burch. n. 4577, in distr. Albany et Uitenhage Eckl. et Zeyh. An sp. propr.? (v. s.)

5. *P. UNGINATA*, glabra, foliis ternato-verticillatis filiformibus supernè sulco exaratis apice deflexo recurvis acutis, capitulis corymbosis, invol. cylindracei-obconici squamis glabris, ext. sensim brevioribus, floribus in invol. 4-5. ζ ad Cap. Bonæ-Spei, ad Lange Valey legit cl. Drege! Similis ex inflorescentiâ *P. acerosæ*, ex dispos. foliorum *P. verticillatæ*. Achænia teretia villosa. Cor. luteæ. (v. s.)

6. *P. VERTICILLATA*, glabra, foliis ternato-verticillatis linearibus patulis apice deflexo-curvis acutis, capitulis numerosis corymbosis, invol. cylindracei-obconici squamis lævibus, exter. sensim brevioribus, floribus in invol. 5. ζ ad Cap. Bonæ-Spei. Recept. angustum fimbriiferum. Pappus pallide rufescens. Achæn. teres villosum. Folia 6 lin. longa subirreg. verticillata. Cor. ex sicco purpurascens. Spec. affinis prioribus. (v. s. comm. à cl. Lambert.)

§. 2. *Foliis oppositis distinctis.*

* *Foliis ramulivæ plùs minùsve pubentibus.*

7. *P. DIVARICATA* (Less. syn. 196), ramulis foliisve pube minimâ subpulverulentâ velutinis, foliis oppositis ovatis obtusis breviter petiolatis, capitulis in cymam corymbosam dispositis pedicellatis cylindraceis, involucri glabri disco dimidiò brevioris squamis ovatis obtusis margine scariosis integris. ζ ad Cap. Bonæ-Spei (Burm.!), ad Groenekloof Eckl. et Zeyh. Drege! *Empatorium divaricatum* Berg. cap. 229. Thunb. fl. cap. 628. *Chrysocoma oppositifolia* Lin. sp. 1177. N. L. Burm. prod. fl. cap. 24. Breyn. ic. t. 17. f. 2. ex Lin. Fol. 6-7 lin. longa, 4 lin. lata. Capit. 7 lin. long. 6-florum. Invol. 4 lin. longum. Pappus rufus. (v. s.)

8. *P. LUPULINA*, ramulis subvillosis, foliis oppositis sessilibus ovalibus suboblongis utrinquè villosis subcaberis, capitulis terminalibus solitariis sessilibus ovatis glabriusculis, invol. discum subæquantis squamis concavis obtusissimis apice scariosis infimis ferè obovatis. ζ ad Cap. Bonæ-Spei ad limitæ colonie prope

Sack-rivier legit cl. Burchell et mecum comm. sub n. 1495. cat. geogr. Fol. 6-7 lin. longa, $2\frac{1}{2}$ -3 lata. Ramuli crebri oppositi ultimi dichotomi. Capitula 16-flora, 8 lin. longa, 4 lin. lata. Stigmata longè exserta. (v. s.)

β. *rotundifolia*, foliis ovato-subrotundis glabrisculis. — ad Nieuwewelde alt. 2-300 ped. legit cl. Drege! (v. s.)

9. *P. OPPOSITIFOLIA* (Lin. syst. veg. 614), ramulis folisque pube subpulverulentâ minimâ velutinâ, foliis oppositis ovato-lanceolatis erectis internodio subæqualibus nervo medio notatis, capitulis terminalibus solitariis sessilibus ovatis glabris apice subviscosis, inv. squamis ovatis obtusis adpressis margine suberoso-ciliolatis. ♀ ad Cap. Bonæ-Spei. Thunb. fl. cap. 632. Gærtn. fr. 2. t. 167. f. 2. *Scepinia dichotoma* Cass. dict. 48. p. 45. Folia 3-4 lin. longa, sesquilinea. Capit. multiflora 9-10 lin. longa, 6-7 lin. lata. (v. s. comm. à cl. Burchell cat. geogr. n. 6837.)
10. *P. CINEREA* (Lin. f. suppl. 356), ramulis subpubescentibus mox glabris, foliis elliptico-oblongis lanceolatisve oppositis cinereo-tomentosis patentibus in ramulis approximatis, capitulis terminalibus sessilibus ovatis 15-20-floris, invol. squamis ovatis obtusis adpressis subciliatis dorso tennè tomentosis sabstriatis. ♀ ad Cap. Bonæ-Spei. Thunb. fl. cap. 632. Folia 5-6 lin. longa, juniora cana adulta cinerea. Capit. 8-9 lin. longa, 6-7 lin. lata. Pappus rufus. (v. s. ex herb. Burm. sub nom. *Stæhelinæ*.)
11. *P. XANTHOLEPIS*, ramulis ultimis folisque pube subtomentosâ cano-villosis, foliis oppositis oblongo-linearibus secus ramulos confertissimis, capitulis terminalibus solitariis sessilibus oblongis 6-9-floris, invol. glaberrimi flavi squamis oblongo-linearibus obtusiusculis integris intimis margine angustè membranaceis. ♀ ad Cap. Bonæ-Spei legit cl. Krebsn. 1901, ad Uitenhage, Swellendam (Eckl. et Zeyh.), et Kaus (Drege!). Folia 3-4 lin. longa, lineam lata. Internodia ramulorum vix semilin. longa. Capit. 6-7-florum. Squamæ seriatim imbricatæ exter. nempe suprâ interiores oppositè nec alternè sitæ. Proximè accedit ad *P. incanam*. Huc fortè *Eupatorium cinereum* Lin. f. suppl. 354. refer. Pappus albus. Squamæ flavæ. (v. s.)
12. *P. INCANA*, ramulis incanis demùm glabris, foliis oppositis linearibus suboblongisve obtusis cano-tomentosis secus ramulos confertissimis, capitulis terminalibus solitariis sessilibus oblongis 10-12-floris, invol. glaberrimi squamis oblongis obtusissimis margine scarioso angustissimo cinctis. ♀ ad Cap. Bonæ-Spei. *Pteronia rigida* Berg. cap. 231 ex Less. syn. 158. *Chrysocoma incana* N. L. Burm. ! prod. fl. cap. 24. *Eupatorium cinereum* Thunb. fl. cap. 627? et ideò *Pteronia rigida* Berg. cap. 231? Similis *P. xantholepidi* sed inv. squamæ pallidè subfuscæ nec flavæ. Capit. 10-12 nec 6-8-florum. Pappus intensè rufus nec albus. Folia multò magis incana nec cinerea. (v. s.)
13. *P. CANESCENS*, ramis glabris vix nascentibus araneosis, foliis oppositis linearibus oblongisve subtrigonis utrinquè canescenti-tomentosis integerrimis ad apices ramulorum confertis, capitulis terminalibus solitariis ovatis, invol. squamis subadpressis ovato-oblongis subacutis dorso araneoso-tomentosis margine angustè scarioso-marginatis. ♀ in Africâ Capensi ad Onderbokkweid et Zeederberge legit cl. Drege! *Achænia sericeo-villosissima teretiuscula* *erostria*. (v. s.)
14. *P. LUCILIOIDES*, demissa tota hirtello-velutina canescens, foliis oppositis linearibus obtusis basi cuneato-attenuatis integerrimis enerviis, capitulis terminalibus solitariis sessilibus 7-9-floris, invol. cylindraceis subconicis glaberrimis. ♀ in Africâ Capensis regione Gariepina et Klein-Namaqualand legit cl. Drege! Folia 5-6 lin. longa, $1\frac{1}{2}$ lin. lata. Capit. semipoll. longa. *Achænia villosa* hispida. (v. s.)
15. *P. ERYTHROCEATA*, glabra, junior velutina, foliis oppositis ovatis obtusis subglaucis superioribus internodio longioribus, capitulis terminalibus sessilibus oblongis 4-floris, invol. squamis oblongis glabris acuminatis margine subscariosis dorso lineatis, pappo rubro. ♀ ad Cap. Bonæ-Spei prope Sack-rivier legit cl. Burchell (n. 1403 et 1546 cat. geogr.), et ad Zwart-Ruggens cl. Drege! Ramuli

et folia novella subvelutina. Rami oppositi adulti rigidi subspinescentes. Folia 4 lin. longa, 2 lin. lata. Capit. 5 lin. longum pappo rubro insignitum. (v. s.)

** *Glaberrimæ.*

16. *P. MICROPHYLLA*, glabra ramosissima, foliis oppositis trigonis ad apices ramulorum confertis integerrimis glaucescentibus minimis, capitulis terminalibus solitariis plurifloris oblongis basi obconicis, invol. squamis adpressis ovatis, exter. subacutis, inter. subciliatis. ζ in Africâ Capensi ad Stellenbosch leg. cl. Eklon.! et Zeyher. Affinis *P. ciliatæ*. Folia vix ultra lineam longa. Invol. flavum apice non contractum. Achænium teres sericeo-villosum. Pappus longus basi rufescens. (v. s.)
17. *P. GLOMERATA* (Lin. f. suppl. 356), glabra, foliis oppositis sessilibus ovato-trigonis in ramulis præsertim ad apices et axillis glomeratis, capitulis terminalibus sessilibus solitariis obovato-cylindraceis viscosis, inv. squamis adpressis, exter. brevibus ovatis dorso viridibus, cæteris apice flavescens ciliatis. ζ ad Cap. Bonæ-Spei in Carro (Th. Drege!), ad Zwartberge et Zwart-Ruggens (Drege!), ad Camisberge et Onder-Bokkewelde (Drege!). Thunb. fl. cap. 63a. Burch. cat. geogr. n. 1359 et 1435. *Scepinia lepidophylla* Cass. dict. 48. p. 45. Folia 1-2 lin. longa subglauca, summa sub capitulis concava obtusa. Capitulum 8-9 lin. longum 4 lin. diam. (v. s.)
18. *P. FALLENS* (Lin. f. suppl. 357), tota glabra ramosissima, foliis oppositis trigonis elongatis subfiliformibus obtusis, capitulis solitariis brevè pedunculatis terminalibus turbinatis, invol. squamis ciliatis, inter. margine scariosis sublacero-ciliatis. ζ ad Cap. Bonæ-Spei in Carro (Eckl.! et Zeyh.), et Cannaland. Thunb. fl. cap. 63o. non Poir. Folia unguicularia ex Th., 10 lin. longa in specim. meo Thunbergiano. Capit. 5-6 lin. longum. Achæn. villis albis sericeis densè onustum. Pappus rufus. (v. s.)
19. *P. TRICEPHALA*, glabra subviscosa ramosissima, foliis oppositis linearibus complicato-trigonis integerrimis apice uncinulato-recurvis, capitulis ad apicem ramorum ternis in axillis supremis 4-5-floris, invol. oblongi squamis acutis linearibus exter. subtrigonis. ζ in Africâ Capensi ad Nieuwewelde alt. 2-300 ped. legit cl. Drege! Suffrutex vix semipedalis. Folia et invol. 4 lin. longa vix ac ne vix subconnata. Achænia teretiusscula erosaria villosissima. An forte *P. fastigiata* Thunb. fl. cap. 629? sed in nostrâ folia semilineam lata nec longa. (v. s.)
20. *P. BECKROIDES*, ramis glabris gracilibus, foliis oppositis sessilibus linearilanceolatis complicatis integerrimis pulverulento-velutinis, capitulis terminalibus solitariis sessilibus oblongo-turbinatis basi attenuatis glabris, invol. squamis lanceolatis acuminatis, inf. brevibus apice callosis, aliis apice membranaceis hyalino-marginatis. ζ ad Cap. Bonæ-Spei in distr. Swellendam collibus Carro similibus legit cl. Drege! Folia 1 $\frac{1}{2}$ lin. longa. Capit. flavicantia 8-10 lin. longa. Achænia sericeo-villosissima teretiusscula. (v. s.)
21. *P. GYMNOCLINE*, glabra, ramis virgatis striatis, foliis inf. oppositis, super. alternis sessilibus oblongo-linearibus obtusiusculis 1-nerviis cinereis minutissimè punctulatis, capitulis terminalibus solitariis sessilibus oblongo-cylindraceis circ. 15-floris, invol. squamis flavidis adpressis, exter. obtusis, intimis longioribus acuminatis. ζ in Africâ Capensis reg. Gariepina, Carro, Klein-Namaqualand, legit cl. Drege! Folia 5 lin. longa, lineam lata. Capit. flavum 9-10 lin. longum. Pedicelli apex ex squamarum cicatricibus areolatus. Recept. planum omninò nudum. Achæn. villosissimum. (v. s.)
22. *P. CILIATA*, glabra ramosissima, foliis oppositis ad apices ramorum confertis ellipticis rigidis nervo medio prominulo subcarinatis integerrimis glaucis, capitulis terminalibus solitariis oblongis utrinquè attenuatis, squamis infimis subfoliaceis carinatis, cæteris adpressis obtusissimis densè et tenuiter ciliatis. ζ in Africâ Capensi ad Namaqualand leg. cl. Ecklon.! et Zeyh. Folia 1 $\frac{1}{2}$ -2 lin. longa. (v. s.)

- β. subtrigona*, foliis oblongis evidentiùs subtrigonis. *♂* ad Olifant-rivier Dregel! *P. ciliata* Thunb. fl. cap. 632? Folia 2-3 lin. longa. (v. s.)
23. *P. succulenta* (Thunb.! prod. 143. fl. cap. 630), tota glabra ramosissima, foliis oppositis trigonis carnosis obtusis patentibus, capitulis terminalibus campanulato-ovatis, invol. squamis ovatis obtusis integerrimis. *♂* ad Cap. Bonæ-Spei in Karro et Hantum (Th.), ad Clan-William (Eckl.!), ad Olifant-rivier (Dreg.!). Achænia villis albis sericeis erectis densè vestita turbinata sub pappo attenuata. Pappus rufus. Folia in specim. meis 2-6 lin. longa, ex Thunb. unguicularia usquè pollicaria. Fl. flavi. Fimbrillæ alveolares achæniæ ferè longitudine. (v. s.)
24. *P. sœuviiifolia*, glabra ramosissima, foliis oppositis carnosis lineari-oblongis acutis planis basi uninerviis patentibus integerrimis, capitulis terminalibus solitariis multifloris, invol. campanulati squamis ovatis obtusissimis integerrimis. *♂* ad Cap. Bonæ-Spei, ad Kaus et Olifant-rivier legit cl. Dreg. Affinis *P. succulentæ* et fortè eum eâ à Thunb. confusa, sed folia plana nec trigona, acuta nec obtusa, pollicem longa, 2-3 lin. lata. Flores flavi. Achænia sericeo-villosissima. Recept. vix alveolatum. (v. s.)
- §. 3. *Foliis oppositis vaginato-connatis, capitulis multi-pauci- imò unifloris cylindraceis, invol. squamis adpressis luteis exter. ovatis.*
25. *P. connata*, glabra ad apices viscosa, foliis oppositis basi connatis lineari-filiformibus supernè sulcatis, capitulis ad apices ramorum solitariis ternisve ovato-cylindraceis 15-25 floris, invol. squamis adpressis obtusis, ext. ovatis, inter. oblongis. *♂* in Africâ Capensi in Karro (Dreg.! Eckl.! et Zeyh.), in Swellendam (Eckl.! et Zeyh.). An *P. flexicaulis* Thunb. fl. cap. 629 nec Lin. f.? Folia sesquipollicem longa, vaginâ rarissimè ruptâ. Capitula pollicem longa nunc solitaria sessilia, nunc terna lateralibus pedicellatis. Flores flavi. (v. s.)
26. *P. flexicaulis* (Lin. f. suppl. 355), glabra ad apicem viscosa, foliis junioribus basi connatis demùm ruptis margine membranaceis lineari-filiformibus, capitulis 3-floris densè corymbosis in quoque ramulo subternis pedicellatis, invol. cylindrici glabri squamis oblongis obtusiusculis. *♂* ad Cap. Bonæ-Spei in Carro (Thunb. fl. cap. 629? Eckl.! et Zeyh.), et inter fluv. Bushman et Gauritz (Burch.! cat. geogr. n. 4209), ad Zw. Onder-Bokkeweld et Zeederbergen (Dreg.!), in Swellendam et Uitenhage (Eckl.! et Zeyh.). *P. fastigiata* Zeyh.! pl. cap. exs. n. 20, non Thunb. qui de suâ dixit folia semilineam longa, Caulis flexuosus cinereus glaber. Folia pollicem longa. Capit. 5 lin. longa. (v. s.)
27. *P. fasciculata* (Lin. f. suppl. 357), tota glabra, foliis oppositis sessilibus confertis lanceolatis acutis viscosis suprâ concavis subtùs carinatis, capitulis plurimis adgregato-fastigiatis oblongis, invol. glabri squamis lanceolatis. *♂* ad Cap. Bonæ-Spei in Karro prope Hex-rivier (Th.), Gauritz-rivier (Eckl.!), in collibus distr. Swellendam (Eckl.!), ad Zeederbergen (Dreg.!). Thunb. fl. cap. 630. *Hemanthus fasciculatus* Less. syn. 195. Folia pollicem longa, 2 lin. lata, juniora vaginato-connata, vaginis demùm obliquè ruptis et ligulas triangulares stipulæformes secus ramos persistentes et oppositas relinquentibus. Invol. lutea. (v. s.)

Sect. II. PACHYDERIS. — *Pachyderis* Cass. dict. 56. p. 170. Less. syn. 196.

Capitulum multiflorum. Recept. alveolatum. Achænia compressa (exter. interdum subtrigona) apice brevissimè rostrata, nempe sub pappo in collum crassum glabrum sæpè villis occultatum attenuata. — Folia opposita rariùs alterna.

§. 1. *Achæniis plus minùs villosis hirsutisve.*

* *Corollis purpureis.*

28. *P. hirsuta* (Linn. suppl. 356), ramis hispidis, foliis oppositis lanceolatis

acutis suprà glabris subtùs hirsutis margine ciliato-scabris, capitulis terminalibus sessilibus solitariis ovatis multifloris, inv. glabri viscosi squamis ovatis integris. ζ ad Cap. Bonæ-Spei, in Karro (Thunb. fl. cap. 631.), in coloniæ distr. orient. (Burch. l. cat. geogr. n. 6380 et 6208), ad Zwartland, Paarlberg (Dreg.!), ad Clan-William, Swellendam, Worcester (Eckl. l. et Zeyh.). An *P. uniflora* Poir. dict. 5. p. 733? Folia 5 lin. longa, $1\frac{1}{2}$ lin. lata sub capitulis conferta falso-verticillata. Capit. pollicem longum, 9 lin. diam. Cor. purpureæ. Achænia villosa in collum glabrum crassum brevem sed aliis ferè longiorem attenuata. Pappus rufus. (v. s.)

29. *P. CEPHALOTES* (Linn. f. suppl. 338), ramis hirtis subfastigiatis, foliis oppositis lanceolato-trigonis rigidè ciliatis subscabris remotiusculis, capitulis terminalibus sessilibus ovatis solitariis, invol. squamis imbricatis adpressis apice subpuberulis viscosisque, exter. brevibus ovatis, mediis oblongis apice ad margines membranaceis ciliatis, ζ ad Cap. Bonæ-Spei prope urbem Burch. l. cat. n. 61. Eckl. l. et Zeyh. Thunb. fl. cap. 631. *Jacea incana*, etc. Burm. l. herb. et idèò *Centaurea incana* Burm. prod. 28 videtur. Capitulum 9 lin. longum, 5 lin. diam. Folia 4 lin. longa erecta patula aut apice recurva. Achænia compressa villosa-hispida breviter et crassè rostrata. Cor. purpureæ. Priori valdè affinis. Ramuli interdum parcè hirtelli et folia etiam parcè ciliata, cæterum glabra (Eckl.) undè ad var. β transit. (v. s.)

β . *glabra*, ramis glabris, foliis eciliatis in disco utroque glabris complicatis aut planis. ζ in Karro legit cl. Ecklon. (v. s.)

** *Corollis flavis.*

30. *P. UNDULATA*, ramis glabris ad apicem tenuissimè foliisque glanduloso-pulverulentis, foliis oppositis in petiolum attenuatis ovatis undulato-crispatis, capitulis ex axillis super. pedicellatis in corymbum 3-7-cephalum dispositis oblongis 5-floris, invol. squamis erectis oblongis glabris integris obtusiusculis. ζ in Africâ Capensi ad Zilverfontain in Klein-Namaqualand legit cl. Drege! Folia cum petiolo sesquilinea. 7-8 lin. longa 4 lin. lata. Invol. flavescens pappo maturo dimidiò brevis. Flores flavi. Achænum compressum subrostellatum parcè villosum. (v. s.)

31. *P. OVALIFOLIA*, ramis glabris vix apice summo subcanescentibus, foliis oppositis elliptico-oblongis ovalibusque integerrimis pube adpressâ subtomentosâ canis, adultis glabris, capitulis ovato-oblongis terminalibus solitariis, invol. squamis adpressis, exter. ovatis, inter. magis oblongis margine scariosis, omnibus mollè et tenuiter ciliatis. ζ in Africâ Capensi ad Olifantrivier (Dreg.!), in Worcester (Eckl.). Achænum villosissimum brevissimè rostratum. Habitus ferè *P. canescens*. Folia majora 8 lin. longa. 4 lin. lata, plurima minora. (v. s.)

32. *P. SCARIOSA* (Linn. f. suppl. 356), ramis glabris demum apice rigidis spinulentibus, foliis suboppositis sessilibus ovali-oblongis glabris, capitulis terminalibus solitariis ovatis, invol. glabri squamis ovatis margine latè scarioso-membranaceis mucrone fusco-subpungenti-acuminatis. ζ ad Cap. Bonæ-Spei ad Hantum (Thunb. fl. cap. 633), ad Kouse (Dreg.). Achænia compressa villosa breviter rostrata. (v. s.)

33. *P. MEMBRANACEA* (Linn. f. suppl. 357), ramis glabriusculis, foliis oppositis sessilibus oblongis acutis pulverulento-tomentosis, capitulis terminalibus solitariis oblongis, invol. glabri squamis lanceolatis margine latè membranaceis dorso fusco-virescentibus. ζ ad Cap. Bonæ-Spei in Karro (Thunb. fl. cap. 633) Ecklon l. et Zeyh. in Bosmanrivier et Zneureeberge (Dreg.!). Frutex pedalis, ramis demum induratis subspinescentibus, adultis obtusè tetragonis. Folia 5-8 lin. longa 1-nervia interdum basi trinervia. Invol. ferè *P. scariosæ* sed mucro ultra partem scariosam non productus. Achænum villosissimum teretè angulatum brevissimè rostratum. (v. s.)

34. *P. ACUMINATA*, tota glabra, foliis alternis lanceolatis utrinquè attenuatis mu-

cronatis subenerviis, capitulis terminalibus subsessilibus oblongis 10-12-floris, invol. squamis lanceolatis acuminatis margine scariosis. ♀ ad Cap. Bonæ-Spei extrâ coloniam inter flum. Zack et Gariep legit cl. Burchell et mecum comm. sub n. 1587 cat. geogr. Folia 10-12 lin. longa, 3 lin. lata. Achænia compressa villosissima sub pappo attenuata. Pappus pallidè rufescens. (v. s.)

35. *P. CENTAUROIDES*, caule tereti minutissimè glanduloso-pulverulento, foliis oppositis et alternis integerrimis, caulinis elliptico-oblongis acutis glabriusculis distantibus, ramulorum linearibus acutis subconvolutis utrinquè velutino-canis, capitulis terminalibus solitariis ovatis, invol. squamis adpressis integris obtusis glabris, infimis brevioribus subacutis dorso subvelutinis. ♀ in Africâ Capensi ad Dutoiskloof legit cl. Drege! Folia caulina 7-8 lin. longa, 2-3 lata, ramealia 3 lin. longa lineam lata. Capit. flava pollicem longa semipollic. lata. Achænia villosa-hirsuta compressa breviter rostrata. (v. s.)
36. *P. GLABRATA* (Linn. f. suppl. 358), tota glabra, foliis oppositis oblongo-linearibus crassiusculis in nervo medio subtitis prominulis glaucis ad apicem et in axillis confertis, capitulis terminalibus solitariis sessilibus oblongis 4-floris, invol. glabri squamis ovali-oblongis margine lato-membranaceis acuminato-submucronatis. ♀ ad Cap. Bonæ-Spei in Carro infra Bockland (Thunb. fl. cap. 630), intra coloniam prope Karroc-rivier (Burch.! cat. geogr. n. 1410). Folia 3-4 lin. longa. Capitula 6-7 lin. longa, 2-3 lata. Cor. flavæ. Achænia compressa villosa sub pappo attenuata. Pappus pallidè rufescens. (v. s. comm. à cl. Burchell.)
37. *P. ELONGATA* (Thunb. fl. cap. 631), ramis glabris, foliis oppositis sessilibus approximato-imbricatis trigonis carinâ et margine ciliato-scabris, capitulis terminalibus obconicis, invol. squamis obtusis sublaceris mucrone calloso parvo. ♀ ad Cap. Bonæ-Spei (Boiss.! herb.), in Uitenhage (Eckl.! et Zeyh.) ad Zw. Kuga et Zondag-rivier (Dreg.!). Affinis ex habitu *P. cephaloti* sed ex caract. multo affinium *P. ciliatæ*. Folia glauca 2-3 lin. longa rigide hinc inde ad margines et costam ciliata. Capitula grossa obovata. Squamæ adpressæ fimbriato-lacerae. Flores flavi. Achænia compressa parcissimè villosa, collo brevi. Pappus rufus. (v. s.)
38. *P. CALLOSA*, glabra ramosissima, foliis linearibus crassiusculis subobtusis integerrimis eciliatis subuninerviis, capitulis terminalibus solitariis sessilibus circ. 15-floris, invol. ovali-oblongi squamis adpressis margine tenuissimè et molli-ter ciliatis apice callosis obtusis. ♀ in Africâ Capensi legit cl. Drege (sed in collect. numeros desid.). Suffrutex semipedalis. Folia 5-7 lin. longa, vix ultrâ semilineam lata. Capit. 9 lin. longum. Invol. pallidè virescens ad apices squam. intensius viride. Cor. flavæ. Achænia villosissima compressa breviss. rostrata. Recept. ferè concavum areolatum brevissimè alveolatum. (v. s.)
39. *P. MUCRONATA*, glabra ramosissima, foliis oppositis lineari subtetragonis basi ciliatis rigidulis obtusis erectis, capitulis terminalibus subsolitariis sessilibus, circ. 15-floris, invol. ovali-cylindrici subviscosi squamis adpressis obtusis margine submembranaceis integris mucrone acuto sæpius patente terminatis. ♀ in Africâ austr. ad Zwart-Ruggens alt. 2-300 ped. legit cl. Drege! Frutex circ. pedalis. Ramuli albid. Folia 3-5 lin. longa. Axillæ foliiferæ. Invol. flava 7 lin. longa sæpius solitaria interdum terna. Achænia villosissima compressa brevissimè rostrata. (v. s.)
40. *P. TURBINATA*, glabra ramosissima, ramis fastigiato-corymbosis 1-cephalis, foliis oppositis linearibus obtusis subtitis subconvexis integerrimis summis gradatim in squamellas minimas bracteiformes conversis, capitulis terminalibus solitariis 15-floris, invol. obovato-turbinati squamis adpressis acutiusculis ♀ in Africâ Capensi ad Gariep., Klein-Namaqualand, legit cl. Drege! Suffrutex semipedalis. Folia 3 lin. longa. Capitula semipollic. Invol. et squamæ corollæ flavæ per exsiccationem viridescentes. Achænia compressa villosa brevissimè rostrata. (v. s.)
41. *P. LEUCOLOMA*, glabra, foliis oppositis sessilibus orbiculatis ovalibusve integerrimis uninerviis, capitulis terminalibus solitariis sessilibus multifloris, invol. cylindracei squamis adpressis obtusis margine scarioso albo integerrimo cinctis.

♂ in Africâ Capensi ad Camisberge legit cl. Drege! Folia 2-3 lin. longa, 2 lin. lata internodiis super. longiora. Capit. 10 lin. longum 4-5 lin. diam. Achænia compressa villosissima rostro brevi glabro. (v. s.)

42. *P. CYLINDRACEA*, glabra, foliis oppositis in ramulis ad axillas confertissimis patulis lineari-trigonis uncinulatis subobtusis, capitulis terminalibus solitariis sessilibus cylindraceis, invol. squamis adpressis latè ovatis margine membranaceis ciliato-laceris, plerisque apice mucrone calloso notatis. ♂ ad Cap. Bonæ-Spei in regione Cis-Gariepinâ legit cl. Burchell et mecum sub n. 1603. cat. geogr. comm. Folia subglauca 3 lin. longa. Capitulum 10 lin. longum, 3 lin. diam. 9-florum. Ovaria compressa villis rarissimis vestita apice sub pappo contracta. (v. s.)

43. *P. TENUIFOLIA*, glabra, caulibus è collo paucis simplicibus, foliis oppositis lineari-subulatis integerrimis acutis, capitulis terminalibus solitariis breviter pedicellatis 8-10-floris, invol. ovali-oblongi squamis adpressis obtusissimis integerrimis. ♂ in Africâ Capensi distr. Caledon leg. cl. Ecklon! et Zeyh. Suffrutex semipedalis. Folia 12-14 lin. longa semilineam lata erecta suprâ subcanaliculata subtus punctis seu tuberculis minimis notata. Achænia compressa brevissimè rostrata villosissima. (v. s.)

44. *P. QUINQUEFLORA*, ramis virgatis glabris junioribus subcinereo-pubescentibus, foliis oppositis oblongo-linearibus subconcavis 1-nerviis integerrimis pulverulento-velutinis subcanescentibus, capitulis ad apices ramulorum solitariis sessilibus terminalibus ob ramulos oppositos ternato-corymbosis 5-floris, invol. ovalis squamis ovalibus acutis dorso 1-nerviis pulverulentis margine hyalinis glabris. ♂ in Africâ Capensi ad Graaf-Reynet leg. cl. Ecklon! et Zeyh. Suffrutex 9-10 pollic. Internodia foliis triplò quadruplòve longiora. Folia 4 lin. longa. Capit. ferè semipollic. Achænia compressa villosa brev. rostrata. Recept. alveolatum. (v. s.)

45. *P. LEPTOSPERMOIDES*, glabra, ramis gracilibus, foliis oppositis linearibus ferè subulatis acutis enerviis integerrimis foveolato-punctatis, capitulis terminalibus solitariis sessilibus oblongis utrinquè attenuatis 7-floris, invol. squamis glabris acuminatis integerrimis. ♂ in Africâ Capensi ad Zilverfontain legit cl. Drege! ex ramis numerosis oppositis habitus paniculatus. Folia 6 lin. longa juniora glaucescentia vix lineam lata. Capitula 6 lin. longa deflorata ferè Dianthi. Achænia compressa breviter rostrata parcè villosa et glandulosa. (v. s.)

46. *P. LEPTOLEPIS*, ramis glabris, nascentibus subcanescentibus, foliis oppositis velutino-subtrigonis integerrimis ad apices ramulorum confertis, capitulis solitariis terminalibus oblongis 7-floris, invol. squamis exactè linearibus subobtusis glabris. ♂ in Africâ Capensi ad Zwart-Ruggens Carro alt. 2-300 ped. legit cl. Drege! *P. oppositifolia* E. Mey.! non *L. Pachyderis obtusifolia* Cass. dict. 56. p. 170? Folia 2-2 $\frac{1}{2}$ lin. longa. Achænia villosa breviter rostrata. (v. s.)

47. *P. LATUSQUAMA*, ramis glabris, junioribus subvelutinis, foliis oppositis lineari-oblongis subtrigonis cano-velutinis integerrimis, capitulis terminalibus solitariis cylindricis 4-floris, invol. squamis latè ovatis acutis adpressis dorso subvelutinis. ♂ in Africâ Capensi ad Camisberge et Onder-Bokkeweld legit cl. Drege! Proximè accedit ad *P. leptolepidem* sed præter caract. indic. differt foliis remotiusculis, magis patulis, brevioribus; invol. squamis majoribus paucioribus. (v. s.)

48. *P. EMPETRIFOLIA*, glabra humilis ramosissima, foliis oppositis ad apices ramulorum approximatis patulis lineari-subtrigonis integerrimis subobtusis junioribus viscosis, capitulis terminalibus solitariis sessilibus turbinato-oblongis multifloris, invol. squamis oblongis ciliato-glandulosis erectis obtusiusculis. ♂ in Africâ australi legit cl. Drege! Suffrutex 3-4 pollic. Internodia brevissima Folia sesquilinea. Capit. 7 lin. longa. Invol. flavescentia. Cor. purpurascens. Achænia compressa villosa sub pappo contracta. (v. s.)

§. 2. *Achæniis omnibus aut saltem centralibus glabris.*

49. *P. HETEROCARPA*, glabra decumbens?, foliis oppositis breviter connatis linearibus obtusis integerrimis crassiusculis, capitulis terminalibus solitariis sessilibus ovato-campanulatis, invol. foliis proximis superato glabro, squamis ovali-oblongis obtusis margine hyalinis, achæniis exter. villosis-hirsutis, centralibus glaberrimis. ζ in Africâ Capensi ad Olifant-rivier legit cl. Drege! Folia 12-15 lin. longa, $\frac{1}{2}$ lata. Capit. 8-9 lin. longam latumque. Pappus rigidus. (v. s.)
50. *P. STORRELIPOIDES*, glabra, foliis oppositis sessilibus rigidulis glaucis linearilanceolatis complicatis ad margines et sæpe subtus in carinâ rigidè ciliatis, capitulis terminalibus solitariis sessilibus ovato-subcylindraceis, invol. squamis glabris adpressis margine scariosis subfimbriatis, achæniis glaberrimis eglandulosis. ζ in Africâ Capensi ad Graaf-Reynet (Dreg.! Eckl.! et Zeyh.), in Carro et Namaqualand (Eckl.! et Zeyh.). Affinis *P. onobromoidi*. Folia 4-5 lin. longa, $\frac{1}{2}$ lata. Capit. pollicem longa. (v. s.)
51. *P. VISCOSA* (Thunb. fl. cap. 632), ramis glabris, foliis oppositis sessilibus ovatis calloso-subacutis margine scabridis patulis, capitulis terminalibus solitariis oblongis, invol. squamis viscosis ovatis obtusis integris achæniis glanduliferis. ζ ad Cap. Bonæ-Spei in Hantum (Th. Zeyh.! in h. Moric. n. 98), ad Zwart-Ruggens in Carro (Dreg.!). Capitula crassa pollicem longa. Achænia basi villis brevibus erectis (fortè fimbriis) cincta compressa oblonga utrinquè attenuata glanduloso-scabrida. Pappus rufescens. (v. s.)
52. *P. ONOBROMOIDES*, glabra, foliis alternis sessilibus elongatis lanceolato-linearibus obtusis rigidulis subconvolutis margine rigidè ciliatis, capitulis terminalibus solitariis sessilibus ovatis multifloris, invol. squamis integerrimis adpressis obtusis ext. latè ovatis, int. oblongis, achæniis glanduliferis. ζ in Africâ Capensi ad Olifant-rivier (Dreg.!) et in Namaqualand (Eckl.!). Rami albi. Folia pallida 12-18 lin. longa, 2-3 lata. Capit. 15 lin. longum, 10-12 lin. latum. Cor. flavæ. Achænia compressa glabra apice breviter rostrata. (v. s.)

Sect. III. *PTEROPHORUS*. — *Pterophorus* Vaill. act. ac. Par. Cass. dict. 37. p. 474. Less. syn. 195. — *Pterophora* Neck. elem. 1. p. 139.

Capitulum multiflorum. Recept. fimbriiferum. Achænia erostre plano-compressum glaberrimum. Pappus subuniserialis, setis basi in anulum subconcretis. — Folia sparsa linearia integra capitula terminalia solitaria, ramis ad apicem ferè foliosis. — Fortè genus inter *Pteroniam* et *Chrysocomam* medium.

53. *P. CAMPHORATA* (Lin. sp. 1176. mant. 464), ramis hirsutis, foliis sparsis fasciculatisque lineari-filiformibus ciliatis, capitulis terminalibus solitariis subsessilibus, invol. squamis serrulato-ciliatis disco brevioribus. ζ ad Cap. Bonæ-Spei frequens (Burm.! Eckl.! et Zeyh. Dreg.!). *Pteronia* camphorata N. L. Burm.! prod. fl. cap. p. 24. Thunb. fl. cap. 629. Lam. ill. t. 667. f. 1. *Pterophorus* camphoratus Cass. dict. 37. p. 474. — Pluk. mant. t. 345. fol. 56. pl. 2. Flores interdum 6-fidi. (v. s.)
54. *P. STRICTA* (Ait kew. 3. p. 162), ramis apice villosis-scabris, foliis sparsis fasciculatisque lineari-filiformibus basi ciliatis, invol. squamis integerrimis apice in folium subacerosum productis disco æqualibus longioribusve. ζ ad Cap. Bonæ-Spei ad Zwartberge (Dreg.!). *Pterophorus* strictus Less. syn. 195. *P. laurina* Houtt.! in h. Delessert. (v. s.)
55. *P. ASPERA* (Thunb. fl. cap. 631), ramis villosis-scabris, foliis fasciculatis linearibus glabris, invol. squamis glabris lanceolatis acutis apice reflexis. ζ ad Cap. Bonæ-Spei, in Stellenbosch, Swellendam, Langekloof, etc. (Eckl.! et Zeyh.), in Paarlberg et Gifberg (Dreg.!). *Pterophorus* asper Less. syn. 195. (v. s.)

† *Species non satis notæ (achæniis nempè indscriptis). Plures fortè inter priores jam recensitæ.*

* *Alternifoliæ.*

56. *P. CRASSIFOLIA* (Poir. dict. 5. p. 731), ramis.... foliis sparatis linearibus carnosè obtusis basi spinuloso-ciliatis, capitulis terminalibus solitariis sessilibus cylindricis, invol. glabri squamis ovali-lanceolatis obtusis apice subfimbriatis. ♀ ad Cap. Bonæ-Spei. Invol. squamæ flavæ.

57. *P. CHRYSOCOMIFOLIA* (Poir. dict. 5. p. 732), ramis pubescentibus, foliis fasciculatis filiformibus acutis glabris, capitulis confertis, invol. glabri squamis linearibus obtusis. ♀ ad Cap. Bonæ-Spei. Cor. in sicco cœrulescens.

** *Oppositifoliæ.*

58. *P. FASTIGIATA* (Thunb. prod. 143. fl. cap. 629), tota glabra, foliis oppositis connatis trigonis obtusis imbricatis apice recurvis, capitulis terminalibus sessilibus ternis, invol. squamis ovatis. ♀ ad Cap. Bonæ-Spei. Caulis ramosissimus, ramulis adgregatis flexuosis subfastigiatis. Folia semilineam longa.

59. *P. PANICULATA* (Thunb. fl. cap. 629), tota glabra, foliis oppositis filiformibus amplexicaulibus, capitulis in paniculam terminalem dispositis, invol. oblongi angulati squamis ovatis obtusis. ♀ ad Cap. Bonæ-Spei. Achæn. ign.

60. *P. VILLOSA* (Linn. f. suppl. 356), ramis scabridis foliis oppositis linearibus obtusis piloso-hispidis, capitulis terminalibus solitariis ovatis, invol. squamis integris summis margine membranaceis. ♀ ad Cap. Bonæ-Spei Thunb. fl. cap. 631.

61. *P. GLAUCA* (Thunb. fl. cap. 631), ramis glabris, foliis oppositis frequentissimis ovatis glauco-tomentosis subcarinatis obtusis, capitulis terminalibus solitariis cylindricis, invol. glabri viscosi squamis ovatis integris, infimis carinatis. ♀ ad Cap. Bonæ-Spei. Folia semilineam longa.

62. *P. RETORTA* (Linn. f. suppl. 356), ramis pilosis, foliis oppositis sessilibus ovatis acutis piloso-scabris reflexis, capitulis terminalibus solitariis ovatis, invol. squamis ovatis obtusis integris. ♀ ad Cap. Bonæ-Spei. Thunb. fl. cap. 633.

63. *P. INFLEXA* (Linn. f. suppl. 356), ramis glabris, foliis oppositis subsessilibus ovatis tenuissimè pubescentibus (pilosive), capitulis terminalibus solitariis ovatis inflexis, invol. glabri squamis ovato-oblongis integris apice membranaceis. ♀ ad Cap. Bonæ-Spei in Hantum. Thunb. fl. cap. 633.

†† *Species exclusæ.*

P. CHAMÆPEUCE Spreng. = *Chamæpeuce mutica*.

P. CHAMÆPEUCE Ten. = *Chamæpeuce gnaphalodes*.

P. MINUTA Linn. f. = *Nestleria minuta*.

P. FAUCIFLORA Sims. = *Helichrysum*.

P. POROPHYLLUM Cav. = *Dysodia porophylla*.

P. TOMENTOSA Lour. fl. coch. ed. W. v. 2. p. 597, certè è genere excludenda sed locus verus ignotus.

DIV. IV. SOLENOGYNÆ.

Capitulum discoidem heterogamum, fl. radii tubulosis aut vix ac ne vix subligulatis tenuissimis sub-3-dentatis fœmineis pauciseriatis, disci hermaphroditis 5-dentatis. — Subdivisio ambigua inter Astereas et Baccharideas.

CXCIV. DUHALDEA.

Capitulum multiflorum discoideum heterogamum, fl. radii paucis tubulosis fœmineis, disci circ. 20 hermaphr. 5-dentatis, fauce longâ tubulosâ. Invol. 3-seriale imbricatum ovatum, squamis linearibus, ext. dorso subtomentosis, int. glabris disco circ. æqualibus. Recept. nudum planum. Anth. ecaudatæ. Styli rami exserti lineares glabriusculi. Achænia villosa linearia aut teretiuscula. Pappus 1-serialis, setis subrigidulis acutis scabris. — Frutex aut suffrutex Chinensis ramosus, ramulis subcorymbosis confertè villosis teretibus, adultis glabratiss. Folia alterna, superior. subsessilia elliptica obtusè mucronulata penninervia integerrima aut parcè calloso-denticulata suprâ pilis sparsis scabra subtus villis subsericeis adpressis cana. Capitula ad apices ramulorum axillarium in paniculam ferè dispositorum confertè breviter pedicellata. Flores verisim. flavi. — Genus affine Rochoniæ sed capit. discoidea. Ex flor. radii paucis ambigit inter Chrysocomeas et Solidagineas. — Nomen in memoriam Joh. Bapt. Duhalde qui medio sæculo XVII Chinam adiit et descripsit.

1. *D. CHINENSIS*. ♂ in Chinâ. Folia infer. ignota, super. 20 lin. longa, 10 lin. lata. (v. s. in h. cl. Lindley.)

CXCV. MICROTRICHIA.

Capitulum multiflorum heterogamum, fl. tubulosis, radii 1-2 serialibus fœmineis minimis, disci hermaphr. 5-fidis. Invol. 1-2-seriale disco brevius. Recept. globosum nudum. Stigma Asterearum. Achænia sessilia turbinata erostria confertissima glabriuscula pappo piloso 1-seriali brevissimo superata, marginalia interdum calva. — Suffrutex diffusus Senegalensis ramosissimus cinereo-hirsutus. Folia alterna sessilia oblonga basi subcuneata integerrima apice latiora grossè dentata subacuta. Rami apice breviter aphylli 1-cephali. Flores flavi. — Genus Gymnostephio et Nolletiae vid. affine.

1. *M. PERROTTETII*. ♂ in Senegaliâ medio Januario florentem legit cl. Perrottet. (v. s.)

CXCVI. NOLLETIA *Cass. dict. 37. p. 479. Less. syn. 187. — Conyzæ sp. Desf.*

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, marginalibus fœmineis filiformibus, cæteris hermaphr. 5-dentatis. Recept. planum nudum. Invol. 2-3-seriale, squamis imbricatis linearibus acuminatis floribus vix brevioribus demum reflexis. Achænium compressoplanum erostre. Pappus 1-serialis setis corollæ subæqualibus scabris. — Frutex Mauritanicus ramosissimus. Folia alterna linearia strigillis adpressis utrinquè subvillosa integerrima. Rami apice breviter denudati 1-cephali. Capitula lutea. — Genus valdè affine Feliciæ et tantum diversum ob flores marginales tubulosos nec radiatos.

1. *N. CHRYSOCOMOIDES* (Cass. l. c.). ♂ in Mauritaniâ ad colles arenosos prope Kervan (Desf.) et circ. Mogador (Brous!). *Conyza chrysocomoides* Desf. fl. Atl. 2. p. 269. t. 232 non arb. fr. Simillima Feliciæ angustifoliæ quæcum confusa fuit etiam à clarissimis Botanicis, sed differt non tantum floribus marginalibus filiformibus nec ligulatis, sed adhuc ramis glaberrimis nec pube adpressâ subincanis, apice ferè usquè ad capitula foliosis nec longè denudatis, foliis adpressè strigillosis nec glabriusculis, 3-5 lin. longis nec sesquipoll., etc. (v. s. sp.)

CXC VII. *SARCANTHEMUM* Cass. bull. philom. 1818. p. 74. dict. 47. p. 349. Less. syn. 191. — *Conyza* sp. Lam.

Capitulum multiflorum heterogamum, flor. omnibus tubulosis, tubo infernè crasso ampliato, radii pluriseriatis fœmineis extus in ligulam minimam superatis, disci 5-dentati abortu ovarii masculis. Recept. planum paleaceo-fimbriiferum. Invol. hemisphæricum imbricatum floribus paulò brevius. Achænia glabra radii compresso-obovata pappo brevi coroniformi squamato superata, disci abortiva ad callum basilarem reducta, pappo setoso-paleaceo irregulari 1-seriali coronata. — Frutex Mauritianus glaber. Folia alterna angustè oblongo-lanceolata basi attenuata apice serrata obscure triplinervia. Corymbus compositus terminalis. Capitula ad apices pedunculorum conferta subsessilia. Flores flavi. — Genus affine *Psiadia* et *Nolletia*.

1. *S. CORONOPUS* (Cass. ll. cc.). ♂ in insulâ Rodrigue (Comm.), Mauritii (Spreng.)? Bourboniæ (Bory!). *Conyza Coronopus* Lam. dict. 2. p. 87, sed *Conyza coronopifolia* Kunth omnino diversa. Folia 2-3 poll. longa, 2 $\frac{1}{2}$ -3 lin. lata. (v. s. comm. à cl. Bory.)

CXC VIII. LEPTOTHAMNUS.

Capitulum multiflorum heterogamum, fl. radii filiformibus truncatis fœmineis 1-serialibus, disci tubulosi 5-dentati hermaphroditis. Recept. ebracteolatum planum vix subalveolatum. Invol. squamæ floribus paulò breviores imbricatæ acuminatæ circ. 3-seriales. Stylus radii apice bifidus, disci inclusus indivisus. Achænium radii gracile glabrum, disci obovato-oblongum pubescenti-villosum. Pappus pilosus deciduus in radio 1-serialis, in disco æqualiter subbiserialis seu saltem copiosior. — Suffrutex gracilis ramosus erectus Capensis *Psilothamnus* paululum referens. Folia sparsa erecto-adpressa lineari-subulata nervo subtus prominente donata, utrinquè rigidè ciliolata. Rami apice breviter apophylli hispido-scabri. Invol. glabra. Cor. è flavo purpurascens. — Accedit ad *Chrysocoma* sed fl. radii fœminei, ad *Erigeron* tem sed fl. radii 1-seriales et habitus diversus.

1. *L. CILIARIS*. ♂ ad Cap. Bonæ-Spei in regione Trans-Gariepinâ ad Klaarwater et Kosiountain (Burch. l. n. 1839 et 2512), ad Klipplatrivier alt. 3-400 ped. (Dreg.!). Frutex pedalis. Rami erecti glabri aut sub capitulis scabri. Folia 2-3 lin. longa. (v. s.)

CXC IX. SOLENOGYNE Cass. dict. 56. p. 175. Less. syn. 193.

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, radii pluriseriatis fœmineis 3-dentatis, disci circ. 10 masculis 4-5-dentatis ovario nullo. Recept. planum nudum. Invol. subhemisphæricum floribus æquale, squamis subtriseriatis inæqualibus obtusis subimbricatis adpressis. Achænia oblongo-obovata compressa erostris epapposa apice summo tuberculo terminata. — Herba Novo-Hollandica parva, Bellii facie, basi foliata cæterum nuda. Folia alterna inæqualia spatulata apice dentata. Caules scaposi filiformes 1-cephali. Capitula parva solitaria. Flores verisim. lutei. — Floribus omnibus tubulosis genus inter *Bellideas* singulare.

1. *S. BELLIOIDES* (Cass. dict. 56. p. 175). ♀? in Novâ-Hollandiâ circa Port-Jackson.

Subtrib. II. *BACCHARIDÆ* Less. syn. 200 non 204.

Capitula heterogama aut dioica nunquam radiata, corollæ omnes tubulosæ, fœminæ filiformes sæpius multiseriales in ambitu. Anthæræ ecaudatæ. Recept. epaleaceum. — Folia alterna.

Div. I. *CONYZÆ* Less. syn. 200.

Capitula heterogama monoica.

Subdiv. I. *SPHÆRANTHÆ*.

Capitula in glomerulum sphaericum congesta ad bractearum axillas sessilia. — Species gerontogæ.

CC. *BLEPHARISPERMUM* Wight herb. DC. in Wight contrib. bot. ind. p. 11. et diss. cum icon.

Capitula innumera in glomerulum globosum coacervata, biflora, fl. tubulosis, altero fœmineo apice breviter 3-dentato, altero masculo 5-dentato. Recept. capituli angustissimum paleâ linearî inter flores interjectâ. Invol. squamæ ext. paucæ (2?) breves ovatæ, int. (seu paleæ rec.) duæ flosculis æquales, altera sub fl. fœmineo plana, altera sub fl. masculum conduplicata. Stylus brevis non exsertus apice vix bifidus, infra emargin. subtimidulus puberulus. Anthæræ apice appendiculatæ basi subcaudatæ? Achænium fl. fœm. obovale-oblongum obcompressissimum subtetragonum, angulis 2 nerviformibus cum lateribus glaberrimis, 2 marginalibus longè et regulariter ciliatis; fl. masc. gracile elongatum glabrum. Pappus achæniî fœm. paleis scariosis 3-5 acuminatissimis, masc. nullus. — Suffrutices Indici glaberrimi. Rami teretes graciles. Folia alterna ovata aut ovalia integerrima penninervia. Pedunculi 1-3 ad apicem rami 1-cephali. Glomerulus globosus. — Affinitas ignota. — Car. præcipuè ex B. petiolari.

1. B. PETIOLARE (DC. l. c.), foliis petiolatis ovato-lanceolatis acuminatis, glomerulis ternis longè pedunculatis. § in Ind. orient. legit cl. Wight sed locus in herbario non indicatus. Glomeruli Piso paulò majores. Bracteolæ involucri univ. loco parvæ membranaceæ paucæ capitulis breviores. Conf. descript. Kleinii editam in Wight contrib. bot. ind. p. 12. (v. s. in h. Wight.)
2. B. SUBSESSILE (DC. l. c.), foliis ellipticis utrinquè attenuatis subsessilibus, glomerulo terminali solitario subsessili, bracteis foliaceis capitula superantibus. § in agro Maderaspano. Glomerulus ferè Platani capitulo æqualis. Capitula fortè pluriflora. (v. s. in h. Wight et inter pl. Wallichianas in h. soc. lin. Lond. sine nom. sub n. 7524.)

CCI. *ATHROISMA* DC. in ann. bot. Guill. 1833. v. 2. p. 516.

Capitula plurima in glomerulum ovatum conferta. Axis glomeruli cylindrica. Bractææ sub capitulis ovatæ acutæ concavæ. Capitula pluriflora heterogama. Recept. capituli paleas plurimas concavas membranaceas inter flores gerens. Inv. oligophyllum à paleis recept. vix distinctum. Fl. ext. 4-5 fœminei tubulosi tenues 4-5 dentati. Flor. int. pauci masculi tubulosi apice latiores 5-dentati. Anth.... Stylus fl. fœm. apice bifidus glabriusculus. Achænia nigra obcompressa ovalia hinc plana illinc convexa ad angulum marginalem apice ciliata cæterum glabra. Pappus parvus villum referens subsetosus. — Herba Indica basi suffruticosa erecta ramosa parcè subpuberula. Folia alterna petiolata lyrato-

pinnatipartita, lobis oblongis obtusis dentatis. Glomerulus terminalis subsessilis stramineus. — *Athroisma* græcè acervus ad capitulorum dispositionem alludit. — Genus videtur medium inter *Sphæranthum* et *Elephaspermum*.

1. *A. LACINIATUM*. ζ in prov. Indiæ or. Martabanâ ad Troglâ legit cl. Wallich. *Sphæranthus laciniatus* Wall.! cat. n. 3184. comp. 294. cat. birm. 1872. (v. s. comm. ab ill. Cur. merc. angl. Ind. or.)

CCII. *SPHÆRANTHUS* *Vaill. act. Par. 1719. t. 20. f. 12. Linn. gen. n. 998. Lam. ill. t. 718. Gærtn. fr. 2. t. 164. Cass. dict. 50. p. 208. Less. syn. 201 non Scop.*

Capitula in glomerulum subglobosum densè aggregata, pluriflora, heterogama, fl. omnibus tubulosis, fœmineis pluriserialibus basi subindurato-dilatatis sæpè tridentatis in ambitu, styli abortu masculis, 5-dentatis in centro. Recept. partiale nudum generale bracteolatum. Invol. partiale polyphyllum uni aut 2-3-seriale, generale varium. Anth. ecaudatæ. Stylus in capit. ambitu bifidus, in centro indivisus. Achæ-nium erostre epapposum pilosiusculum aut glabrum. — *Herbæ Indicæ* aut *Africanæ*. Folia longè secus caulem decurrentia. Pedunculi alati aut apteri 1-cephali. Glomerulus densus Pisi aut *Avellanæ* magnitudinis. Cor. violacæ.

Sect. 1. *EUSPHÆRANTHUS*.

Glomeruli axis plena nec lacunosa; bractæ omnes lineares molles capitulis subæquales, exteriores vix ac ne vix involacrum generale constituentes. — *Species Indicæ*.

1. *S. MICROCEPHALUS* (Willd. sp. 3. p. 2395), foliis lanceolatis aut subovatis serratis vix asperiusculis in alas integerrimas secus caulem decurrentibus, capitulis globosis, pedunculis sæpius apteris capitulo triplo longioribus. \mathcal{Z} frequens in uliginosis et oryzetis Indiæ orient. nempe in ins. Luzoniâ (Cham.!), Javâ (Blum.!), Timor (Riedl!), ad Prome, Penang, Tavoy, Chapedong, etc. (Wall.! Wight!), in Chinâ (Staunt!). *Sph. africanus* Wall.! cat. et herb. n. 289 excl. litt. B. *Sph. lævigatus* et *Sph. globosus* Wall.! herb. *Sph. indicus* Blum. bijdr. p. 891. var. \mathcal{E} . An *Sph. indicus* Linn. saltem fl. Zeyl. n. 312? An *Sph. africanus* ex Indiâ Linn. sp. 1314? Capitula semper in meis specim. globosa Pisi circ. magnit. Pedunculus apterus aut hinc indè rarissimè subalatus. (v. s. specim. ∞ .)
2. *S. HIRTUS* (Willd. sp. 3. p. 2395), foliis obovatis serratis utrinquè hirtellis in alas serratas decinentibus, glomerulis ovato-globosis, pedunculis glomerulo triplo longioribus alis serrato-incisis ferè semper instructis. — Frequens in humidis et oryzetis peninsulæ Indicæ (Wight!), ad Patna, Kuworkot, Travancore, etc. (Wall!), circa Pondichery (Belang!), in ins. Java circa Buitenzorg (Blum!). *Sph. hirtus* Blum.! bijdr. p. 891., Wall.! cat. n. 288. D. E. F. *Sph. africanus* Wall.! cat. n. 289. B. C. *Sph. indicus* Wight! herb. Icones Pluk. alm. t. 108. f. 7 et 312. f. 6. Burm. Zeyl. t. 94. f. 3. huc fortè pertinent. Glomerulus Piso duplo ferè major. Pedunculus rarius subapterus. (v. s.)
3. *S. MOLLIS* (Roxb. herb.), foliis obovato-lanceolatis utrinquè caulibusque hirsuto-lanuginosis grossè serratis, serraturis aristatis, subserratis, glomerulis ovato-globosis subhirsutis, pedunculis glomerulo duplo longioribus aliis hirsutissimis apteris aliis interruptè alatis hirsutis. \mathcal{Z} in Indiâ orient. prov. Silhet, Kamaon, Hindostan (Roxb. et Wall!). — Icon Rheed. mal. 10. t. 43. huc fortè refer.? — Caulis ad collum induratus ferè fruticosus. Caules è collo plurimi sesquipedales ramosissimi. An fortè *Sph. hirti* var. hirsutissima? (v. s. comm. ab hon. Anlâ merc. angl. Ind. or. et à cl. Royle.)

Sect. II. CUSPIDELLA.

Glomeruli axis subplena aut vix subcava; bracteæ apice acuminato-pungentes, exteriores invol. universæ distinctè formantes, cæteræ vix ac ne vix capitula superantes. — Species Africanæ.

4. *S. SENEGALENSIS*, foliis lanceolatis acuminatis utrinquè villosis aristato- et ferè spinuloso-serratis secus caulem in alas grossè et subspinuloso-serratas desinentibus, glomerulis globosis, pedunculis interruptè et subspinuloso-alatis glomerulo multò longioribus. (I) in Senegaliâ ad ripas inund. fluminis præsertim prov. Walo legerunt cl. Adanson, Roussillon! Perrottet! Sph. africanus seu paniculatus Cass. dict. 50. p. 211 et 212. Variat à caule digitali simplici 1-cephalo ad caulem sesquipedalem ramosissimum polycephalum. Capitulum violaceum Pisi majoris magnit. (v. s. comm. à cl. Perrottet.)
5. *S. ANGUSTIFOLIUS*, foliis linearibus acutis glabriusculis distanter serratis in alas longas subserratas desinentibus, pedunculis glomerulo duplò ferè longioribus alis integris ad latus inferius instructis, glomerulo depresso bracteis latè ovatis acuminatis cincto. — in arenosis humidis Senegaliæ ad Casamance legit cl. Perrottet. Planta abnormis et fortè novi generis pignus. Capitula parva pauciflora in glomerulum densè aggregata. (v. s. comm. à cl. Perrottet.)
6. *S. PEDUNCULARIS*, foliis linearibus acutis glabriusculis serratis in alas longas angustas subserratis productis, pedunculis folia æquantibus aut superantibus, glomerulo globoso bracteis latè ovatis breviter mucronatis cincto. — in Africa australiori ad Zw. Omsanculo et Omcomas legit cl. Drege! Proximè accedit ad *S. angustifolium* sed pedunculi 2 pollices nec semipollicem longi. Folia 2-2½ poll. longa, 2-3 lin. lata. (v. s.)

Sect. III. POLYCEPHALOS. *Polycephalos Forsk. ægypt. descr. p. 154.* — *Oligolepis Cass. dict. 50. p. 212.* — *Sphæranthus Gærtn. fr. 2. p. 413. t. 164.*

Glomeruli axis plena aut sæpiùs cavo-lacunosa; bracteæ omnes ovato-lanceolatae cuspidatæ rigidæ capitula superantes, exteriores vix ac ne vix invol. gener. constituentes. — Species Indicæ, Africanæ et Australasicæ.

7. *S. GLABER*, totus glaber, foliis lanceolatis vix subserratis in alas subintegerrimas decurrentibus, pedunculis glomerulo paulò longioribus aliis apteris aliis ad basin hinc indè angustissimè alatis. 2½ ad ripas flum. Alligatorum in sinu Van-Diemen ad oram Australiæ borealem legit cl. Cunningham. Axis obovata plena. Pedunculus gracilis. Basis corollarum callosa albida. Fl. disci pauci. Achenium glanduloso-viscidum. (v. s. comm. à cl. Cunningham.)
8. *S. SUAVEOLENS*, foliis lanceolatis serratis vix asperiusculis in alas parcè serratas decurrentibus, pedunculis glomerulo dimidio brevioribus subtrigonis angulis alatis subdentatis. 2½ in Ægypto ad ripas Nili! et fortè in Indiâ orientali sed nulla specimina Indica in herbariis reperi. — *Polycephalos suaveolens Forsk. æg. p. LXXXII et 154.* Sph. indicus Lam. ill. t. 718. f. 2. Gærtn. fr. t. 164. Delil. fl. æg. ill. p. 27. Cass. dict. 50. p. 210 et omn. ferè hort. europ. Vulgò arabis dicitur *Habagbag*. Glomerulus ovato-globosus, axi lacunosâ, 6-7 lin. longus totidemque diam. Cor. violaceo-purpureæ. (v. s. sp. et v. c.)
9. *S. AMARANTHOIDES* (Burm. fl. ind. 186), foliis lanceolatis aut lineari-lanceolatis integerrimis serratisve glabris subtus punctatis breviter decurrentibus, capitulis ovato-globosis subsessilibus, pedunculis brevissimis apteris, bracteis acuminatissimis. 2½ in Indiæ orientalis humidis, in prov. Tangore ad oppidum Negapatnam (Wight!), ad ripas Irrawaddy inter Kink-Talong et Vandoboo (Wall!) et fortè in Zeylonâ. — DC. in Wight contrib. bot. ind. p. 11. S. Zeylanicus Heyne in h. Wall! n. 290. — Pluk. amalth. t. 357. f. 6. Affinis *S. suaveolenti* sed notis indicatis distinctissimus. (v. s. sp. comm. ab hon. Aulâ merc. angl. Ind. or.)
10. *S. ? COTULOIDES*, reptans hirsutus, caulibus teretibus gracilibus, foliis ovatis

dentatis mucrone cuspidatis secus caulem in alam parvam interruptam productis, capitulis parvis terminalibus brevissimè pedicellatis. — in insulâ Madagascar ad viarum margines prope urbem Tananarivou legit cl. Bojer. Cotula reptans Bojer! in litt. Herba parva habitu ad Grangeas seu Dichrocephalas accedens, sed ex foliis decurrentibus et squamarum dispositione in anthodia equidem nimis juniore observatâ, speciem cum dubio ad Sphæranthum reduxi. (v. s. comm. à cl. Bojer.)

† *Species non satis notæ.*

11. *S. COCHINCHINENSIS* (Lour. fl. coch. 2. p. 623), foliis oblongis integerrimis lanuginosis secus caulem glabrum indè pseudotriquetrum decurrentibus, glomerulis cordato-ovatis breviter pedunculatis. — in segetibus et hortis Cochinchinæ et Chinæ. Flores albicantes parçè purpurâ conspersi.
12. *S. CHINENSIS* (Linn. mant. 119), foliis sessilibus pinnatifidis, pedunculis alas decurrentes crispas gerentibus. — in Indiâ orientali. An verè Sphæranthi congener? An affinis *S. laciniato* Wall.? An eadem? Cur Chinensis, cum ipse cel. auctor ex Indiâ ortam dicat? †.

†† *Species exclusæ.*

S. AFRICANUS Burm. = *Dichrocephala latifolia*.

S. AFRICANUS Linn. videtur ex priore et ex Sphærantho microcephalo conflatus.

S. LACINIATUS Wall. = *Athroisma laciniatum*.

Subdiv. 2. GRANGEINÆ.

Capitula heterogama monoïca in glomerulum non congesta. Pappus nullus aut subnullus.

CCIII. *DICHROCEPHALA* DC. in ann. bot. Guill. 1833. v. 2. p. 517 et in *Wight contrib. p. 11*. — *Centipeda* Less. syn. 200 non Lour. — *Cotula*, Grangæ, Ethuliæ et Artemisiæ sp. auct. — *Dichrocephalæ* sp. *L'Hér. mss.*

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, marginalibus multiseriatis vix 3-4-dentatis fœmineis, centralibus hermaphroditis abortu masculis paucis 4-dentatis. Recept. nudum conicum. Invol. expansum subuniseriale, squamis ovatis subæqualibus. Cor. fœm. cylindricæ, centrales fauce campanulatâ. Stylus inclusus. Achænium compressum erostre, fœmineis calvum, masculis setulâ 1-2 coronatum. — Herbæ annuæ ex Asiâ aut Africâ ortæ hirsutiusculæ aut subglabræ ramosæ adscendentes aut erectæ. Folia alterna dentata cuneata aut petiolata, petiolo nudo aut appendiculato. Capitula globosa parva, in racemos paniculasve pauca disposita, pedicellis propriis nudis breviora. — *Centipeda* Loureirii quam cl. Lessing non ipse vidit, differt à nostro genere, ex descr. auct., 1°. Achæniis oblongis subteretibus nec compressis. 2°. Pappo nullo nec in fl. masc. 1-2-setoso. 3°. Capitulis axillaribus solitariis sessilibus nec racemosis. 4°. Floribus purpurascens nec albo-flavis. Potius ad *Myriogynem* pertinere videtur.

1. *D. CAPENSIS*, caule demisso aut adscendente pilis sparsis hirtello, foliis ovatis grossè dentatis sparsè puberulis petiolatis, petiolis aliis nudis aliis lobulis 1-2 utrinquè auctis, capitulis longè pedicellatis racemosis. ① ad Cap. Bonæ-Spei (Th.), in Coloniz distr. oriental. (Burch. cat. geog. n. 5243). *Ethulia auriculata* Thunb. fl. cap. 623. *Centipeda Capensis* Less. syn. 201. Herba basi repens adscendens vix pedalis. Capitula parva subglobosa pauca racemosa. (v. s. comm. à cl. Burch.)
2. *D. GRACILIS*, glaberrima aut apice summo subpubescens, caule adscendente debili parçè ramoso, foliis exauriculatis pinnatipartitis lobis utrinquè 2-3-

subdistantibus oblongis apice pauci-dentatis, capitulis longè pedicellatis paucis racemosis. ① in Indiæ prov. boreali-occident. legit cl. Royle. Caulis imò basi subradiceus, pedalis. Capitula parva. Species ferè media inter *D. Capensem* et *latifoliam*. (v. s. comm. à cl. Royle.)

3. *D. LATIFOLIA*, caule erecto sparsè piloso, foliis obovatis in petiolum attenuatis grossè dentatis sæpè basi inciso-subpinnatifidis, ramis floridis ramosis subnudis, pedicellis rigidis divaricatis capitulo globoso longioribus. ① in Javâ, Indiâ or., Chinâ, ins. Madagascar. *Cotula bicolor* Roth. cat. bot. 2. p. 116. *Sphæranthus Africanus* Burm. ind. 185. t. 60. f. 2. non Linn. *Grangea latifolia* Lam. ill. t. 699. f. 1. *Cotula sinapifolia* Roxb. herb. Wall. cat. et herb. n. 34. excl. litt. *G. Ethulia glomerata* Spreng. in litt. ad Wall. *Hippia bicolor* Smith in Rees cycl. 18. n. 2. *Grangea dissecta* Bojer! in litt. Poir. suppl. 2. p. 826. *Cotula latifolia*. Pers. ench. 2. p. 464. *Dichrocephala erecta* L'Hér.! mss. Desf. cat. h. par. 1804. p. 95. *Spilanthes atriplicifolius* Linn. ed. Houtt. in h. Deless. (v. v. et s.)

β. *Javanica*, foliis superioribus profundius incisus ferè pinnatisectis, lobis lanceolatis. In Javâ montanis humidis. *Cotula latifolia* var. *Javanica* Blum.! bijdr. n. 913. (v. s.)

4. *D. SONCHIFOLIA*, caule erecto glabriusculo, foliis petiolatis lyratis, lobis lateralibus utrinquè solitariis oblongis obovatisve, terminali maximo ovato dentato, ramis floridis racemoso-ramosis, pedicellis pedicello ramoso longioribus. ① in Iberiâ (Stev.), ad flum. Alazonium (Bieb.), in Persiæ prov. Guriel prope Ozargheti et Dwabzo (Szovitz!). *Cotula sonchifolia* Bieb. fl. taur. 2. p. 328. *Ethulia paniculata* Schk. handb. t. 235. Valdè affinis *D. latifolia* sed glabrior, minus rigida; folia longius petiolata verè lyrata, limbo basi non cuneato. (v. s.)

5. *D. CHRYSANTHEMIFOLIA*, erecta ramosa tota pube brevi confertâ hirtello-scabra, fol. inf..., superioribus oblongis cordato-semiamplexicaulibus grossè serratis summis integris, pedunculis capitulo multò longioribus. — in Javâ montanis detexit cl. Blume et in Indiâ orient. legit cl. Wight. *Cotula chrysanthemifolia* Blum.! bijdr. p. 918. Capitulum piso paulò minor in parte ext. flavescens cæterum purpurascens. *Achænia glabra* ferè calva. (v. s.)

6. *D. GRANGEA*, caule erecto tereti piloso, foliis caulinis glabriusculis basi breviter auritis semiamplexicaulibus angustè oblongis pinnatifidis, lobis paucis vix subdentatis ovato-oblongis, summis linearibus subintegris, pedunculis 1-cephalis subfoliosis, capitulo obovato. — in Indiâ prov. boreali-occident. legit cl. Royle. Non malè habitum refert *Grangea* cernuoidem. Invol. parvum. *Achænia compressa glabra* fortè omnia calva. (v. s. comm. à cl. Royle.)

7. *D. LYRATA*, caule ramoso tereti piloso, foliis villosis in petiolum attenuatis lyratis, lobis lateralibus paucis parvis terminali obovato aut ovali dentatis, capitulis ad apices ramorum solitariis inter folia suprema sessilibus. — in insulâ Madagascar ad viarum margines prope urbem Tananarivon detexit cl. Bojer. *Cotula lyrata* Bojer! in litt. non Wall. *Achænia compressa glabra* apice setulis 2-3 subpuberis. Habitus *Grangeæ* sed ex *achæniis* potiùs ad *Dichrocephala* referenda. (v. s. comm. à cl. Bojer.)

CCIV. *GRANGEA* *Adans. fam.* 2. p. 121. — *Grangea* et *Pyrarda* *Cass. dict.* 19. p. 304 et 41. p. 120. — *Grangea* et *Pyrarda* *Less. syn.* 200 et 201. — *Grangea* sp. *Juss. Lam.*

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, radii pluri-aut biserialibus fœmineis tenuissimis 3-dentatis, disci hermaphroditis 5-dentatis. Invol. biseriale, squamis oblongis obtusis. Recept. hemisphæricum. Anth. basi ecaudatæ. Stylus radii bifidus, disci indivisus. *Achænium* obovatum lateraliter subcompressum basi attenuatum apice supra semen productum uti sub flore subcupuliforme. Pappus minimus piloso-dentatus cupulæ margine insidens. — *Herbæ* *Africæ vel Asiæ æquinocialis* plus minus villosæ. Folia alterna semiamplexi-

caulia sinuato-pinnatifida, lobis obtusis. Capitula solitaria subglobosa, pedunculis oppositifoliis aut terminalibus.

Sect. I. *CRASSICOLLA*. — *Grangea* *Adans. l. c. Cass. dict. 19. p. 304.*

Flores radii pluriseriales. Rostrum achæniî prominens albidum cellulosum semine ferè crassius.

1. *G. MADERASPATANA* (Poir. dict. suppl. 3. p. 825), caulibus procumbentibus diffusivè puberis apice summo villosis. ① in Indiâ orientali (Wall. cat. et herb. n. 345), præsertim in agro Maderaspatano frequens (Wight!), in Javâ prope Bataviâ (Blum. bijdr. p. 919). *Artemisia Maderaspatana* Linn. sp. 1190. Lam. dict. 1. p. 264. *Cotula Maderaspatana* Willd. sp. 3. p. 2170. *Grangea Adansonii* Cass. dict. 19. p. 304. — Pluk. alm. t. 1. f. 2. mala. *Grangea* Lam. ill. t. 699. f. 3 huc ex textu referenda fortè ad sequentem potiùs pertinet. (v. s.)

Sect. II. *LEPTODERIS*.

Flores radii pluriseriales. Achæniû compressum ovale utrinquè attenuatum, rostro gracili in cupulam expanso.

2. *G. ÆGYPTIACA*, caule procumbente foliisque hirsutis subincanis. ① in Ægypto ad canalem Alexandriæ (Delil.) et circa Kahiram in marginibus agrorum (Forak.). *Tanacetum Ægyptiacum* Jacq. hort. Vind. 3. t. 88. *Grangea Maderaspatana* Desf. cat. hort. par. ed. 3. p. 159 excl. syn. Delil. ill. fl. æg. p. 25 excl. syn. *Tanacetum humile* Forak. æg. descr. p. 148? — Lam. ill. t. 699. f. 3. Herba Gr. *Maderaspatanæ* simillima et sæpè cum eâ confusa, sed differt habitu multò magis hirsuto et canescente et achæniî collo angusto nec insigniter crasso. (v. s.)

Sect. III. *PYRARDA* *Cass. dict. 41. p. 120. non Adans.* — *Pyrarda Less. syn. 202.* — Nomen à *Pyrard* peregr. Gallo.

Flores radii biseriales. Achæniû apice truncatum, rostro calloso vix conspicuo aut nullo.

3. *G. CERUANOIDES* (Cass. dict. 19. p. 307), caule erecto ramosissimo. — in Senegaliâ secus ripas fluminis prov. Walo legerunt olim Geoffroy et recentius Perrottet. *Pyrarda cernanoides* Cass. dict. 41. p. 121. Folia basi obtusè subauriculata semiamplexicaulia nunc dentata nunc pinnatifidita, plus minù villosa, lobis acutis. Capitula pedicellata corymbosa in ramo quoque circiter quaternaria. (v. s. comm. à cl. inv.)
4. *G. PROCUMBENS*, caulibus longè procumbentibus foliisque villosis-hispidis subincanis. — in Senegaliâ flor. mense nov. legit cl. Perrottet. Habitus *Grangæ Ægyptiacæ* sed fructificatio *Pyrardæ*: achæniû nempe truncatum nec in collum sensim attenuatum. An fortè Gr. *Galamensis* Cass. dict. 19. p. 307 sine descr.? (v. s. comm. à cl. Perrottet.)

+ *Species exclusæ.*

G. CUNEIFOLIA Poir. = *Myriogyne minuta*.

G. DECUMBENS Desf. = *Myriogyne minuta*.

G. MINUTA Poir. = *Myriogyne minuta*.

G. LATIFOLIA Lam. — *Dichrocephala latifolia*.

G. LANCEOLATA Poir. = *Eclipta procumbens*.

CCV. *CYATHOCLINE* *Cass. ann. sc. nat. 1829. p. 34. dict. se. nat. 60. p. 581 et 595. Less. syn. p. 202.*

Capitulum multiflorum heterogamum, fl. marginalibus multiseriali-

bus fœmineis tenuissimis limbo parvo 3-dentato, centralibus plurimis masculis tubo obconico 5-dentato. Invol. pluriseriale disco ferè brevius. Recept. ebracteolatum suprâ invol. basin elevatum cupulæforme. Achænia epapposa glabra oblonga utrinquè attenuata, rostro brevi pelliculo. — Herbæ Indicæ puberæ odoratæ viscosæ erectæ ramosæ. Folia pinnatipartita, lobis obovatis linearibus dentatis, dentibus lobulisve secus rachin inæqualiter intermixtis, caulina basi in auriculas dentatas magnitudine varias dilatata. Capitula parva subglobosa albida, in racemos corymbosos nudos terminales disposita.

1. *C. LYRATA* (Cass. l. c.), foliis inferioribus lyratis, lobis folii superioribus majoribus obovatis. ①? in Indiâ orient. provinciis diversis nempe in Pegu (Belang.!), Nepauliâ et circa Trogla, Oude, Prome, Taong-Dong (Wall.), Mooty-Journa (Ham.). — DC. in Wight contrib. bot. ind. p. 12. Tanacetum viscosum A, B, C, D, Wall. cat. et herb. n. 323a. comp. 342. Tanacetum gratum Wall. herb. Tan. purpureum Ham. herb. Artemisia hirsuta Rottl. in Spreng. syst. 3. p. 490 ex specim. Spreng. à cl. Besser comm. Herba sesquipedalis. (v. s.)
2. *C. STRICTA*, foliis pinnatipartitis, lobis sublinearibus grossè et regulariter serratis. ①? in Indiâ orient. verisim. Maderaspatanâ. Tanacetum viscosum E et F Wall. cat. n. 323a. comp. 342. Artemisia stricta Heyn. herb. (v. s.)

CCVI. *LESTADIA* Kunth in Less. syn. 203.

Capitulum multiflorum heterogamum, fl. fœmineis pluriserialibus in ambitu, reliquis masculis. Receptaculum planum epaleaceum. Involuc. imbricatum. Corolla disci et ambitûs 5-dentata sed fœminea tenuior. Achænium oblongum rostratum calvum. — Fruticulus ramosissimus circ. 4 pollic. Americæ meridionalis. Folia hirsuta linearia sparsa confertissima. Capitula solitaria in ramis supernè aphyllis, æquè magna ac in Bellide perenni.

1. *L. PINIFOLIA* (Kunth et Less. l. c.). 5 in Amer. merid. Habitus Asteris hyssopifolii.

CCVII. *GYMNARHENA* Desf. mem. mus. 4. p. 1. t. 1. Cass. dict. 21. p. 111, 23. p. 566. Less. syn. 419. — *Gymnarhea* Steud.

Capitulum multiflorum discoideum, fl. radii multiserialibus fœmineis tubulosis 3-4-dentatis post anthesin basi incrassatis et accretis, fl. disci 10-12 bisexualibus ovarii abortu masculis fauce dilatatis 4-dentatis. Invol. nullum aut squamis paucis 1-serialibus constans, bracteis foliaceis 1-2 suppletum. Recept. convexum in parte exter. paleas flores fœmineos involventes post anthesin accretas gerens, in disco et radio fimbrias floribus interpositas ferens. Anth. basi nudæ. Stylus radii bifidus, ramis longis gracilibus semiteretibus, disci simplex subclavellatus acutus grossè papillosus. Achænia radii villosa oblonga pappo setoso pluriseriali scabro superata, disci teretia glabra inania setis 5-6 uniseriatis basi inter se concretis superata. — Herba Orientalis demissa annua ramosa. Folia alterna pauca oblonga circa capitulum bracteiformia, cæterum nulla. Capitula flava. — Genus in ordine anomalum videtur affine hinc Evaci suad. cl. Desfontaines sed antheræ ecaudatæ, illinc Grangæ suad. cl. Cassini sed pappus saltem elongatus.

1. *G. MICRANTHA* (Desf. l. c.). ① in Persiâ inter Mossul et Bagdad leg. cl. Olivier et Bruguère.

Subdiv. 3. EUCONYZEÆ.

Capitula in glomerulum non congesta. Pappus pilosus 1- aut 2-serialis.

CCVIII. THESPIIS DC. in *Guill. ann. bot.* 1833. v. 2. p. 517. — *Cotulae* sp. *Wall.*

Capitulum multiflorum heterogamum, flor. marginalibus multiseri-
libus cor. tenuissimâ subconicâ stylo vix exserto, centralibus paucis
masculis cor. limbo subcampanulato tubo brevissimo anth. inclusis
ecaudatis. Recept. nudum planum punctulatum. Involucrum pluriseriale,
squamis oblongis acutis disco subbrevioribus. Achænia compressa gla-
bra. Pappus 7-8-setus achænio brevior, setis crassiusculis barbellatis.
— Herbæ Indicæ annuæ erectæ ramosæ glabræ. Caules teretes. Folia
alterna oblonga basi attenuata apice dentata. Pedunculi axillares et
terminales capitula pauca parva subcorymbosa congesta gerentes. Flo-
res ex sicco vid. flavidi. Pappus junior albus demum subrufescens.
Genus affine Conyzæ et Grangææ. — Nomen alludit ad Thespidem
quemdam heroem mythologicum qui filias, ut receptaculum nostrarum
stirpium corollas femineas, 50 protulit.

1. T. DIVARICATA, caule à basi ramosissimo, ramis divaricatis, pedunculis folio brevioribus. ① in Indiæ or. provinciis Silhet et Nepal legit cl. Wallich. *Cotula divaricata* Wall. cat. et herb. n. 3238. comp. 348. Herba 4-5 poll. longa. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
2. T. ERRECTA, caule basi simplici strictè erecto apice corymboso-ramoso, ramis obliquè erectis, pedunculis folio longioribus. ① in Indiæ orient. prov. Silhet. *Cotula sinapifolia* var. G. Wall. cat. et herb. n. 3227. comp. 347, excl. aliis litt. ad *Dichrocephalum latifolium* referendis. Caulis 12-15 poll. longus. (v. s. comm. cum prior.)

CCIX. KARELINIA Less. ined. ex *Fisch et Mey. in litt.* 1833. — *Serratulae* sp. *Pall. Willd.*

Capitulum multiflorum heterogamum, fl. ambitus multiserialibus
fœmineis filiformibus truncatis, centralibus 8-10 hermaphroditis 5-
fidis. Recept. nudum, planum. Inv. cylindrici squamæ ovato-lanceolatæ
rigidæ adpressæ. Antheræ basi ecaudatæ. Stylus fl. fœm. gracilis glaber,
lobis divergentibus elongatis, fl. herm. inclusus longè hispidulus lobis
brevibus erectis. Achænia erostria cylindracea. Pappus fl. fœm. 1-serialis,
setis tenuissimis elongatis capillaribus, fl. hermaphr. setis subplurise-
riatis apice barbellatis. — Herba Orientalis erecta ramosa glabra. Folia
sessilia oblongo-lanceolata integra margine subscabra. Capitula corym-
bosa cylindrica non malè extus Centaureæ cerinthefoliæ seu *Acroptili*
capit. referentia. Flores purpurascens. — Genus affine Conyzæ sed
habitu toto involucro teretiusculo facilè distinctum.

1. K. CASPIA (Less. ined.). 2/ in depressis salsis prope mare Caspium olim legit Pallas, et nuperius ad ostia fluvii Ural cl. Karelina. *Serratula Caspia* Pall. voy. app. n. 397. vol. 2. t. 13. p. 470. *Serratula Caspia* Willd. sp. 3. p. 1643. Herba bipedalis, corymboso-ramosa. (v. s. comm. à cl. Fisch et Mey.)

CCX. BERTHELOTIA.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus fœmi-
neis tenuissimè tubulosis 2-3-denticulatis, disci paucis (5-12) majori-
bus 5-dentatis hermaphr. sæpè abortivo-sterilibus. Recept. planum

punctatum nudum. Invol. ovalis squamæ pluriseriales imbricatæ adpressæ, ext. latæ breves obtusæ, intimæ lineares disco subæquales acutæ scariosæ. Antheræ ecaudatæ. Stylus radii gracilis glaber exsertus apice bifidus, disci papillosus subinclusus bifidus, ramis papilloso-hispidis. Achænia sessilia erostria teretiuscula glabra. Pappus 1-serialis, radii setis tenuissimis vix scabridis basi subdistinctis, disci setis rigidis basi subpaleaceis concretis apice barbellato-scabris. — Fruticulus Indiæ et Africæ æquinoctialis indigena, erectus ramosus, in totâ superficie pube brevissimâ subadpressâ molliter canescens. Folia alterna sessilia lanceolata mucronata rigidula integerrima aut hinc indè ad apicem acutè serrata nervosa. Capitula ad apices ramorum densè corymbosa oblonga. Flores purpurascences. Pappus albus. Genus affine Conyzæ sed differt habitu et pappo fl. centralium polyadelpho, paleis nempe linearibus basi concretis. — Fruticulum elegantem dicavi cl. Berthelot floræ faunæque Canariensis strenuo illustratori.

1. B. LANCEOLATA. ♂ Frutic. pedalis. An species forsân dux admittendæ?

a. *Senegalensis*, capitulis 5-6 lin. longis, pappi disci setis evidentèr basi concretis quasi polyadelphis. In arenosis circa Lamsar et Gandiole mensæ junio legit cl. Perrottet. Conyza proteiformis Perr.! in litt. Folia infer. ex Perr. in litt. magis serrata. (v. s. comm. à cl. inv.)

β. *indica*, capitulis paulò minoribus et confertioribus pappi setis brevius et obscurius polyadelphis. — in Indiâ orientali ad ripam Gangis prope Cownpure et in ripâ fl. Jummæ leg. cl. Wallich et Hamilton. Conyza rubra Ham.! herb. Conyza lanceolata Wall.! cat. et herb. n. 2991. comp. n. 101 Saussurea mucronata Spreng in litt. ad Wall. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

CCXI. LÆNNECIA Cass. dict. 25. p. 91. 37. p. 462 et 482. Less. syn. 203. — Conyzæ sp. Kunth.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus fœmineis tenuissimis apice 2-3 dentatis, disci hermaphroditis 5-dentatis. Recept. ebracteolatum punctulatum. Inv. campanulatum imbricatum, squamis linearibus apice subulato-acuminatis glabris coloratis. Antheræ ecaudatæ. Achæmium erostre oblongo-obovatum compressum villosum. Pappus conformis setosus biserialis, serie ext. brevî multisetosâ, int. longâ pilosâ. — Herbæ Americanæ lanatæ gnaphaloideæ. Folia alterna oblongo-linearia dentata aut incisa. Capitula ad apices ramulorum axillarum sæpius solitaria.

1. L. GNAPHALIOIDES (Cass. l. c.), caulibus diffusis à basi ramosis, foliis oblongo-linearibus inciso-dentatis, capitulis hemisphæricis. ♀ in locis altis montis Cocollar Cumanensium. Conyza gnaphalioides H. B. et Kunth nov. gen. am. 4. p. 73. t. 327.

2. L. FILAGINOIDES, caule simplici stricto, foliis oblongo-linearibus parçè dentatis, capitulis ovato-hemisphæricis. ①? circa urbem Mexici legit cl. Berlandier angusto flor. (pl. exs. n. 820). Priori caracteribus evidentèr affinis sed præter notas suprâ indicatas diversa. Capitula dimidiò minorâ, pedunculis magis lanatis, etc. (v. s.)

3. L. PARVIFOLIA, caulibus diffuso-ramosissimis, foliis oblongo-lanceolatis acutè inciso-dentatis, capitulis ex axillis ortis pedicellatis subsessilibusve in racemum subdigestis. ① in Mexici valle Toluccanâ octob. flor. legit cl. Berlandier (pl. exs. n. 11241). Flor. herm. 2-3 in centro capituli, fœminei circ. 12-15. Caulibus diffusis ad L. gnaphalioidem accedit sed folia 2-3 lin. longa, capitula dimidiò et ultrâ minorâ. (v. s.)

CCXII. CONYZA *Less. syn.* 203. — *Eschenbachia Mœnch meth.* 573. — *Dimorphanthæ et Fimbrillaria Cass.* — *Conyzæ sp. Linn.* — *Conyzæ, Baccharidis et Erigerontis sp. auct.*

Capitulum multiflorum heterogamum, fl. radii multiseriis foemineis corollâ tenuissimâ filiformi truncatâ aut breviter 2-3-dentatâ, disci paucis masculis tubulosis 5-dentatis. Recept. ebracteolatum punctulatum seu fimbrilliferum planum aut convexum. Inv. squamæ pluriseries. Antheræ ecaudatæ. Achænium compresso-planum sæpius glabrum basi attenuatum. Pappus 1-serialis, setis filiformibus vix scabris. — Herbæ ferè omnes Gerontogææ. Caules erecti ramosi teretes. Folia varia sæpè variè incisa semiamplexicaulia. Capitula pedunculata corymbosa aut subpaniculata. Flores flavi. Pappus rufus aut rufescens. — Differt ab *Erigeronte* et *Nidorellâ* ligulis planè nullis, à *Blumèâ* achæniis compresso-planis et antheris ecaudatis.

Sect. I. ANDRYALOIDES.

Recept. alveolare, alveol. marginibus integris. Achænium pappi evid. scabri longitudini æquale aut ferè longius (nec ut in cæter. multò brevius). — Herba Indica subcaulis lanato-tomentosa, scapis 1-cephalis. — An fortè genus?

1. *C. ANDRYALOIDES*, tota tomento lanuginoso cana subcaulis, foliis subradicalibus petiolatis obovatis obtusis basi subcuneatis grossè et paròè dentatis, scapis 1-cephalis folia triplò circ. superantibus foliola linearia acuta integra paucis gerentibus, inv. squamis biserialibus linearibus acutis dorso puberis vix achænio longioribus. ♀ in Ind. prov. boreali-occident. legit. cl. Royle. Herba 2-3 poll. longa subcæspitosa verisim. alpina aprica. Achænia compressa elongata villosissima. Pappus 1-serialis scaber achænio ferè brevior, corollâ longior. Fl. foeminei tennes pluriseries, centrales steriles androgyni, anth. ecaudatis, stylo simplice inclus. (v. s. comm. à cl. Royle.)

Sect. II. DIMORPHANTHES. — *Dimorphanthæ Cass. dict.* 13. p. 254; 25. p. 93; 37. p. 461 et 481.

Recept. planum nudum aut brevissimè alveolato-subfimbriatum. Achænia pappo setiformi multò breviora.

* Species Americanæ. (*Pleræque disco hermaphrodito nec masculo donatæ et forsân sectionem propriam generis constituent.*)

2. *C. TRIPLINERVIA* (Less.! in linnæâ 1831. p. 137), herbacea glaberrima, caule erecto sulcato-angulato, foliis petiolatis ovato-lanceolatis basi cuneatis et integerrimis apice acuminatis serratis triplinerviis subitè punctatis, paniculâ corymbosâ polycéphalâ, capitulis pedicellatis, invol. squamis lineari-lanceolatis acutis, achænio glabro subcompresso pappo brevior. ♀ in Brasiliâ merid. (Sellow!), ad fretum Sanctæ-Catharinæ et Rio Janeiro (Less.), prov. Rio-Grande (h. Imp. Bras. n. 785). Folia 3 poll. longa, pollicem lata. Pappus ♂ apice subclavellatus, ♀ filiformis, uterque in specim. meo albus. Fl. ♂ 3-7. (v. s. comm. à Mus. reg. Berol.)

3. *C. ARGUTA* (Less.! in linnæâ 1831. p. 138), herbacea glaberrima, caule erecto subsulcato, foliis vix petiolatis lineari-lanceolatis acuminatis per totum ambitum argutè et regularitè serratis uninerviis impunctatis, paniculis corymbosis polycéphalis, capitulis pedicellatis, invol. squamis lineari-lanceolatis acuminatis. ♀ in Brasiliâ meridionali (Sellow!), prov. Rio-Grande (h. Imp. Bras. n. 783). Folia 3 poll. longa, 3 lin. lata. Capitula ferè *C. triplinervie* sed flores masculi 1-3 undè primâ fronte ad *Baccharides* vergit. (v. s. comm. à Mus. reg. Berol.)

4. *C. THESIIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 75), suffruticosa basi ra-

mosa, ramis erectis simplicibus glabris angulatis, foliis sessilibus angustè linearibus acutis integerrimis 1-nerviis utrinquè adpressè pilosis, corymbis terminalibus simplicibus, capitulis pedicellatis, invol. squamis 2-3-seriatis subæqualibus linearibus apice subulatis, ext. hispidis. $\frac{5}{8}$ in Mexico?

5. *C. ERIGEROIDES*, herbacea erecta ramosissima, caule striato villosa, foliis sessilibus lineari-sublanceolatis utrinquè attenuatis integerrimis basi subtriplicinerviis pubescentibus, paniculâ diffusâ parçè foliosâ amplâ, capitulis pedicellatis, invol. squamis 1-2-seriatis linearibus acuminatis puberis post anthesin omninò deflexis. ①? in Brasiliâ circa Bahiam. Simillima hinc *Erig. Canadensi*, illinc *Conyze fastigiatae*. Recept. foveolatum omninò nudum vix ultra lineam diam. Pili caulis sublati, ramorum adpressi. Folia 2-2 $\frac{1}{2}$ poll. longa, 3 lin. lata, infima hinc indè dente notata. (v. s.)
6. *C. SINUATA* (Ell. sketch 2. p. 323), herbacea pilosa scabrinuscula; foliis infer. lanceolatis profundè sinuatis, lobis ovalibus acutis, super. linearibus integerrimis, capitulis paniculatis, invol. imbricatis squamis lineari-subulatis. ① frequens circa Charleston Carolinæ. Flores albi centrales hermaphroditi. Achænia oblonga angulata pilosa.
7. *C. CHILENSIS* (Spreng. nov. prov. 1818. p. 14), caule herbaceo erecto subsimplici tereti striato-nervoso hirsuto, foliis radicalibus patulis oblongis basi attenuatis obtusis sinuatis, caulinis paucis semiamplexicaulibus subhirsutis lanceolatis acutis subintegerrimis, capitulis paucis corymbosis latissimis. $\frac{2}{3}$ in Chili et in Brasiliâ (Chamisso!), prope Bahiam (Salzm.! Blanch!), et ad portum Mulgrave Amer. bor. occid. (herb. Hænke!). — Link enum. 2. p. 322. Less. in linnæâ 1831. p. 136. *Conyza procera* Desf. cat. h. Par. 1829. p. 164. *Dimorphanthès procera* Cass. bull. philom. 1821. p. 175. dict. 25. p. 93. Fl. innumeri pallidè flavi centrales ferè omnes hermaphr. et plerique fertiles. Pappus in specim. spontaneis rufus, in cultis albidus. Recept. punctulatum, 4 lin. diam. (v. s. sp. et v. c.)

♀. *monocephala*, caule foliisque glabrinusculis, capitulo solitario. — in Chili cum priore legit cl. de Chamisso. An fortè sp. propria? (v. s.)

8. *C. LINEARIS*, herbacea ramosa, caule pilis patulis hispidò ad apices ramorum steriliùm confertè hirsuto, foliis linearibus integerrimis acutis sæpè spiràliter tortis utrinquè pube brevi confertâ cinereis basi plus minùs villosis, paniculâ racemosâ, capitulis longè pedicellatis, invol. squamis biserialibus linearibus dorso puberis apice glabrinusculis. $\frac{2}{3}$? in pratis et arvis sabulosis ad Quintero Chilensium legit cl. Bertero januar. flor.; etiam video sine loci design. in h. Dombey. Capit 3 lin. longa, matra 6-7 lin. lata. Recept. subalveolare nudum 3 lin. ferè diam. Affinis *C. diversifoliæ* ex foliis super. et *C. Chilensi* capit. majusculis. (v. s.)
9. *C. ALBIDA* (Willd. herb. ex Spreng. syst. 3. p. 514), herbacea erecta, caule villosa striato supernè ramoso-paniculato, foliis sessilibus lineari-lanceolatis acuminatis apice remotè et grossè serratis canescenti-pubescentibus, paniculâ foliosâ laxâ, capitulis pedicellatis, invol. squamis 1-2 seriatis linearibus acuminatis dorso pubescentibus. ① ex Salzm. in Brasiliâ (Cham.), ad Bahiam (Salzm.!), Rio-Janeiro (Beyr. Blanch!) et Sanctam-Catharinam (Mac-Rae in h. Lindl.). — Less. in linnæâ 1831. p. 136. Habitus *Erig. Canadensis* seu Bonariensis. Affinis *C. diversifoliæ*. Recept. lineam diam., alveolatum, ferè fimbriatiferum. (v. s.)
10. *C. DIVERSIFOLIA* (Weinm. in florâ 1820. p. 611), herbacea ad collum suffruticosa erecta tota villosa-cinerea, caule simplici pilis confertissimis hirsuto, foliis pube breviori velutino-villosis subtùs hirsutis elongato-linearibus acutis, infer. hinc indè grossè serratis, cæteris integerrimis, paniculâ oligocephalâ pubescente, capitulis pedicellatis, invol. squamis linearibus pube adpressâ parvâ vix cinereis. $\frac{2}{3}$ in pascuis Chilensibus ad S. Fernando legit cl. Bertero. *C. ambigua* Hook. et Arn. in bot. Beech. p. 57? excl. syn. Plantam descr. ex specim. spontaneis 1-2-pedalibus et foliis 3 poll. longis, 2 lin. latis; sed ut vid. ex

descr. Weinm. in plantâ cultâ caulis 5-6-pedalis, folia 7-9 poll. longa, radic. pinnatifida. (v. s.)

11. *C. ? CERNOPODIODES*, rhizomate suffruticoso, caule herbaceo ramosissimo basi pilosiusculo, ramis virgatis glabrinusculis, foliis lineari-lanceolatis utrinque acuminatis sparse villosis membranaceis, infer. apice trilobis, mediis remotis et grosse serratis, summis integerrimis, paniculis elongatis apice corymbosis, capitulis longè pedicellatis, invol. squamis linearibus. $\frac{1}{2}$ in Brasiliâ prov. Rio de Janeiro in Serra dos Orgaos legit cl. Vauthier (pl. exs. n. 324). Recept. angustum nudum, Flor. ambitus numerosi teretes crassiusculi stylo non exserto, masculi pauci centrales vix dissimiles. Achenium compressum, Papp. 1-ser. (v. s.)
12. *C. CORONOPIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 70), herbacea erecta canescenti-hispida, foliis sessilibus pinnatipartitis, lobis linearibus acutis basi subinciso-lobatis, capitulis pedicellatis in corymbis terminales ramosos dispositis, invol. squamis subtriseriatis æqualibus linearibus apice subulatis. $\frac{1}{2}$ in altâ planitie Mexicanâ ad littus lacus Tezucensis. Herba bipedalis. Cap. 2 $\frac{1}{2}$ lin. longa. Flor. disci 30-35.
13. *C. HISPIDA* (H. B. et Kunth nov. gen. am. 4. p. 71), herbacea erecta sparse hispida, foliis sessilibus lineari-lanceolatis mucronatis integerrimis, capitulis longè pedicellatis laxè corymbosis hemisphæricis, invol. squamis vix imbricatis linearibus acuminatis dorso hispidulis. $\frac{1}{2}$ in altâ planitie Mexicanâ ad littus lacus Tezucensis (H. B. et K.), circa urbem (Berl.! pl. exs. n. 537). Folia sesquipoll. longa, 3-4 lin. lata. Capitula 3-4 lin. diam. Discus circ. 10-florus. (v. s.)
14. *C. OBTUSA* (H. B. et Kunth nov. gen. am. 4. p. 71), herbacea subcæspitosa canescenti-hispida, caulibus sterilibus subdiffusis, floridis erectis, foliis basi petioliformibus integris apice pinnatipartitis lobis oblongis integerrimis obtusis mucronatis, capitulis racemoso-corymbosis vix pedicellatis, invol. squamis vix imbricatis linearibus acutis dorso hispidulis. $\frac{1}{2}$ inter urbem Mexici et Toluca-m (Berl.! pl. exs. n. 537). An *Conyza gnaphalioides*? Andrieux pl. exs. mex. n. 274 non Kunth ab hac satis differt? Pappus enim 1-serialis. Folia præsertim ad basin conferta. Capitula ferè *C. sophiæfoliæ*. (v. s.)
15. *C. CARDAMINIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 71), caulibus herbaceis cæspitosis procumbentibus pilosis, foliis petiolatis profundè pinnatifidis pubescenti-pilosis, laciniiis spathulato-lanceolatis acutis integerrimis, capitulis pedicellatis corymbosis, invol. semiglobosi squamis lanceolato-linearibus apice subulatis. $\frac{1}{2}$ in ruderatis frigidis ad pagum Mulalo Quitensium in radicibus montis Cotopaxi. Flores albi centrales 25-50 hermaphroditi. Affinis *C. coronopifoliæ*.
16. *C. SUBDECURRENS*, herbacea erecta stricta puberula, caule simplici tereti, foliis adnato-sessilibus breviter decurrentibus ligulato-linearibus hinc indè irregulariter et acutè dentatis confertis demùm sæpè inflexis, capitulis brevissimè pedicellatis in paniculam angustam foliis internixtam digestis, invol. squamis biseriatis linearibus acuminatis dorso pubero-hirtellis. $\frac{1}{2}$ (I) ? in Mexico (Hænke!) ad Cordilleras Guchilapanas (Berl.! pl. exs. n. 1044). Caulis circ. pedalis. Folia 10-12 lin. longa, 2-3 lin. lata, sæpè crispato-undulata. Capitula 2-3 lin. lata, sæpè crispato-undulata. Capitula 2-3 lin. diam. Flores disci circ. 10. (v. s.)
17. *C. KUNTHIANA*, herbacea pusilla cæspitosa repens, foliis confertis sessilibus lineari-spathulatis obtusis integerrimis albo-lanatis, capitulis terminalibus solitariis sessilibus, invol. squamis subscariosis obtusis, inter. oblongo-linearibus. $\frac{1}{2}$ in regione alpinâ regni Quitensis? *C. pusilla* H. B. et Kunth nov. gen. am. 4. p. 69 non Houtt. Flores centrales hermaphr. Folia 3-4 lin. longa.
18. *C. MYOSOTIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 69), caule herbaceo supernè corymboso polycéphalo hispidio-hirto, foliis sessilibus spathulato-oblongis obtusis hispidulis, capitulis corymbosis pedicellatis, invol. semiglobosi squamis linearibus subulatis. $\frac{1}{2}$ in temperatis juxta La Sequia Novo-Granaten-

sium ad ripam Smitæ. Flores albi centrales hermaphr. Caulis bipedalis superne angulatus.

19. *C. LYRATA* (H. B. et Kunth nov. gen. am. 4. p. 70), herbacea ramosissima viscosa glanduloso-hirta, caule sulcato, foliis sessilibus ovato-oblongis basi lyratis dentatis, pedunculis terminalibus et axillaribus 3-5-cephalis, invol. subcampanulati squamis linearibus acuminatis apice hispidis disco longioribus. $\frac{1}{2}$ in humidis prope Guayaquil Quitensium. Flores albi in invol. 220-250, centrales hermaphroditi. Herba 2-3 pedalis.

20. *C. SOPHILEFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 72. t. 326), herbacea stricta virgata hispidula, foliis pinnatipartitis, lobis linearibus apice 2-3-fidis obtusiusculis, capitulis glomerato-apicatis, fasciculis in paniculam dispositis, invol. squamis vix imbricatis linearibus acuminatis hispidulis. $\frac{1}{2}$ ex H. B. et K., forte ① ex spec. sicco. — in valle Mexicanâ (Berl. pl. exs. n. 694) inter Chalco et lacum Tezcucensem (H. B. K.). Radix gracilis. Caulis 1-2-pedalis. Capit. 2 lin. diam. Discus circ. 5-florus. (v. s.)

21. *C. PULCHELLA* (H. B. et Kunth nov. gen. am. 4. p. 72), herbacea paniculato-ramosa hirtella, foliis sessilibus pinnatipartitis, lobis linearibus integerrimis obtusis, capitulis pedicellatis corymboso-subpaniculatis, invol. squamis biseriatis lanceolato-linearibus apice subulatis. $\frac{1}{2}$ in regione temperatâ circa Moran Mexicanorum. Herba semipedalis. Folia 4 lin. longa. Flores in inv. circ. 100, quorum 6 centrales hermaphroditi.

22. *C. APURENSIS* (H. B. et Kunth nov. gen. am. 4. p. 73), herbacea erecta superne ramosa hispidula, foliis sessilibus spatulato-oblongis acutis argutè serratis, corymbo fastigiato polycephalo, invol. semiglobosi squamis linearibus apice subulatis. $\frac{1}{2}$ in humidis prov. Varinensis ad ripas flum. Apures. Herba pedalis. Affinis *C. coronopifolia*. Flores albi centrales hermaphroditi.

23. *C. FLORIBUNDA* (H. B. et Kunth nov. gen. am. 4. p. 73), herbacea erecta ramosissima glabriuscula, foliis petiolatis lanceolatis acutè remotè et argutè dentatis utrinquè sparsè et adpressè puberis, corymbo foliato, invol. subsemiglobosi squamis linearibus acutis glabris. $\frac{1}{2}$ frequens circa urbem Quito. Flores albi 30-35 quorum 7-8 centrales hermaphr. Herba 3-pedalis gratè odora.

$\frac{1}{2}$ caule hispidulo superne paniculato, foliis subintegerrimis. $\frac{1}{2}$ circa Guancabamba Peruvianorum.

24. *C. GRAHAMI*, caule imâ basi suffruticuloso herbaceo subsimplici patentim et sparsè piloso apice confertè hispido, foliis semialexicaulibus oblongo-ligulatis basi integerrimis apice parè et grossè serratis glabriusculis secus nervum et ad margines villosulo-ciliatis, capitulis 3-5 subsessilibus congestis, invol. squamis linearibus acuminatis dorso hispidulis. $\frac{1}{2}$ in Bonariâ ad Mendozam. Affinis *C. Gouani*. Flores innumeri, centrales circ. 20 masculi? Pappus albidus invol. paulò longior. (v. s. cult. in horto Edimb. à cl. Grahami comm.)

** Species Gerontogæ, nempe in Europâ, Africâ continentali aut Indiâ or. crescentes.

25. *C. ALTAICA*, caule herbaceo ramosissimo paniculato parè piloso-setoso, foliis sessilibus oblongo-linearibus setaceo-ciliatis, capitulis solitariis ramos terminantibus, bracteis foliaceis inv. squamas referentibus sed capitulo longioribus evid. ciliatis inv. cingentibus. invol. squamis linearibus disco æqualibus apice subrecurvis. ② in Sibirîæ altaicæ plautiebus prope Lokteusk. *Erigeron ciliatus* Ledeb. ic. alt. ill. 1. t. 100. fl. alt. 4. p. 92. Cor. radii tenues, albæ, ex cl. Ledebour brevissimè ligulatæ, ex specim. observato breves truncatæ, undè ad Conyzam species meo sensu referenda. (v. s. comm. à cl. C. A. Meyer.)

26. *C. GOUANI* (Willd. sp. 3. p. 1928), rhizomate lignoso, caule glabriusculo herbaceo erecto, foliis lanceolatis apice serratis margine ciliato-scabris, infer. obovatis, summis integerrimis, capitulis ad apices caulis et ramorum congestis, invol. squamis lanceolatis margine membranaceis. $\frac{1}{2}$ in insulis Canarien-

sibus Palmā et Teneriffā. *Erigeron Gouani* Linn. mant. 469. Jacq. hort. Vind. t. 79. Gouan illust. 66? *Dimorphanthes Gouani* Cass. dict. 13. p. 255. *Baccharis hieracifolia* Lam. *Baccharis arunca* Schrank. *Erigeron scariosum* Mönch. (v. s. et v.)

β. *diffusa*, caule demissiore ramosiore subdiffuso. 2. *Conyza diffusa* Poir. suppl. 2. p. 338. (v. s. comm. à cl. Broussonet ex insulis Canariis.)

27. *C. ROYLEI*, caulibus herbaceis adscendentibus simplicibus teretibus velutino-sabhisuris, foliis radicalibus petiolatis ovalibus, caulinis basi attenuatis obovatis, omnibus grossè dentato-incisis villosis-subhiapidis, capitulis racemosis pedicellatis subsecundis, invol. squamis oblongo-lanceolatis acutis, acheniis compressis in disco glabris margine subciliatis. 2. in Indiæ or. prov. boreali-occid. legit cl. Royle. Herba 6-8 poll. Cor. masculæ flavae. (v. s. comm. à cl. Royle sub n. 135.)

28. *C. PUBESCENS*, caulibus herbaceis simplicibus adscendentibus cum foliis et involucribus pube minimā brevissimā velutino-pubescentibus, foliis infer. in petiolum longum attenuatis obovato-oblongis, super. brevè basi attenuatis ovali-oblongis acutis, omnibus parè et minutè dentatis, capitulis racemosis breviter pedicellatis, invol. squamis oblongis mucrone subacutis, acheniis compressis utrinquè hispidulis. 2. in Indiæ or. provinc. boreali-occident. legit cl. Royle. Affinis sed diversa videtur à *C. Roylei* quācum in herb. conjuncta sub n. 135 et nī fallor sub n. 133. (v. s. comm. à cl. inv.)

29. *C. BOVÉ* (DC. in ann. sc. nat. 1834. bot. p. 251), glabra, caule erecto apice corymboso-ramoso tereti, foliis sessilibus semiamplexicaulibus oblongo-linearibus distanter dentatis obtusiusculis, corymbo laxo, invol. squamis linearibus acuminatis pedicellisquæ puberulis. — in humidis montis Sinai majore florentem legit cl. Bové. Arabis dicitur *Gazani*. *Conyza Bové!* pl. exs. n. 118. Herba fortè basi suffruticosa, circiter pedalis. Folia sesquipoll. longa, 2 lin. lata. Capitula 3-4 lin. lata. Fl. femineae innumeri tenuissimi stylo exserto bifido, cor. truncata, centrales 5-dentati pauci stylo simplici verisim. abortivo. (v. s. comm. à cl. invent. cum pl. aliis Arabicis et ideò nomine suo insign.)

30. *C. FASTIGIATA* (Willd. enum. suppl. 57.), caule herbaceo erecto tereti ramoso subcorymboso hinc inde piloso, foliis sessilibus subamplexicaulibus lanceolato-linearibus subserrulatis integerrimisve basi ciliatis, corymbo terminali oligocephalo, capitulis pedicellatis, invol. squamis subuniseriatis linearibus acutis disci longitudine. ① ex Senegal. habeo spec., etiam in Mauritania (ex Spreng. ! syst. 3. p. 514). *C. paniculata* Desf. ! cat. h. Par. 1815. p. 114. non Willd. An *Dimorphanthes angustifolia* Cass. dict. 25. p. 95? Valde affinis hinc *C. ambigua* illinc *C. Gouani*. (v. s.)

31. *C. AMBIGUA* (DC. fl. fr. suppl. p. 468), caule herbaceo erecto ramoso paniculato tereti hirsuto, foliis linearibus sessilibus acutiusculis integerrimis aut parè dentatis scabro-puberulis, paniculā polycephalā, capitulis pedicellatis invol. squamis subuniseriatis linearibus acuminatis disci longitudine hispidulis. ① in pratis et cultis Galliae occitanicæ non infrequens, in Italiā circa Romam (Sebast.!), Neapolim (Guss.) et Pisas (Savi!), in Hispaniā Gaditanā (Pic!), Græciā (h. ac. Mon!). *Erigeron crispum* Pourr. chl. narb. n. 470 ex Dnb. ! bot. gall. 1. p. 266. *Erigeron linifolium* Willd. sp. 3. p. 1955. *Erigeron contortum* Desf. h. Par. ex Pers. ench. 2. p. 40. *Erigeron Canadense* Ten. prod. ex syll. p. 428. Habitus *Eriger.* Canadensis. Sola species Europæa nunc in *Conyze* genere remanens! Recept. planum nudum. Flor. fem. innum., masc. 5-6. Achenia teretiuscula glabra. (v. s.)

32. *C. CAFFRA*, caule herbaceo erecto glabro aut pilosiusculo, foliis semiamplexicaulibus oblongo-lingulatis obtusiusculis remotè serrulatis utrinquè glabris, pedunculis velutinis in corymbum laxum dispositis, invol. squamis lineari-subulatis velutino-canescenscentibus disci longitudine, cor. masc. circ. 25 apice hirsutis, fem. innumeris. — in Africæ austr. terra Caffrorum, circa Key, legit cl. Drege! Herba sesquipedalis. Folia 30-33 lin. longa, 7 lin. lata. Recept. planum

punctulatum, punctis fl. central. majoribus distantibus, fl. fœm. confertissimis minimis. (v. s.)

33. *C. ÆGYPTIACA* (Ait. kew. 3. p. 183), caule herbaceo erecto tereti hispido, foliis oblongo-spathulatis utrinque pilosis per totam longitudinem grossè et obtusè dentatis, capitulis subpaniculatis globosis, invol. squamis linearibus acuminatis. ① in *Egypti desertis* Delil. ill. fl. æg. p. 25. *Erigeron ægyptiacum* Linn. mant. 112 (excl. Bocc. syn. et ideò patr. Sicula). Jacq. hort. Vind. 3. t. 19. *Conyza ægyptiaca* et *Baccharis Dioscoridis* Bové! pl. exs. n. 302 et 304. *Erigeron serratum* Forsk. æg. 148. *Eschenbachia globosa* Moench meth. 573. *Dimorphanthès ægyptiaca* Cass. dict. 13. p. 256. An *C. spathulata* Horn. cat. h. hafn. 2. p. 805 et Link enum. h. berol. 2. p. 322. huc, jam suad. Hord., ut varietas addenda? For. flavi. Recept. punctulatum. *Achænia compressa*. (v. s. sp. et v. c.)

34. *C. JAPONICA* (Less. syn. 204), caule herbaceo erecto simplici piloso, foliis sessilibus obovatis basi attenuatis apice obtusis serratis villosis, paniculâ terminali. ① in Japoniæ insulâ Nipon. Vulgò dicta *Foso*. Pappus ferrugineus. *Erigeron Japonicum* Thunb. fl. jap. 312.

35. *C. DENTATA* (Willd. sp. 3. p. 1928?), caulibus herbaceis erectis teretibus apice subramosis undiquè hispidissimis foliosis, foliis utrinque hirsutis oblongis, infimis basi attenuatis parcè inciso-dentatis, super. sessilibus semiamplexicaulibus à basi ad medium inciso-dentatis ferè semipinnatifidis, corymbo terminali composito polycephalo, invol. squamis linearibus acuminatis disco sublongioribus. 2/ in demissis humidis ad ripas fl. Senegal in regione Walo legit cl. Perrottet. Herba 12-15 poll. longa. Affinis *Ægyptiacæ*. Pappus rufescens. Rec. alveolato-subfimbriiferum. (v. s. comm. à cl. Perrottet.)

♀. *subnuda*, caule multò minùs præsertim supernè folioso, foliis angustioribus minùs dentatis, corymbo minore confertiore, pappo albedo. 2/ in demissis humidis et arenosis Senegambiæ ad Khoro legit cl. Perrottet. *Erigeron exstipulatum* Schum. guian. p. 387? An species propria? Flor. fœm. ad apicem obliquè truncati. (v. s. comm. cum priore.)

36. *C. AMOENA* (Link enum. h. Berol. 2. p. 322), caule herbaceo hirtò, foliis sessilibus ovalibus obtusis basi attenuatis denticulatis hirtis, paniculâ terminali arcatâ. 2/ ad fluvium Congo. Herba 2-3-pedalis pilis aliis glandulosis. Invol. squamæ hirsutissimæ æquales lineares. Cor. purpurascens. An *Blumæ* sp.? An prioris var.?

37. *C. ASTEROIDES* (Wall. cat. et herb. n. 3052. comp. 162. non Linn.), caule ascendenti erecto piloso ramoso tereti, foliis oblongis pilosis, infimis basi attenuatis obtusis grossè paucidentatis, superioribus semiamplexicaulibus acutis subintegerrimis, ramis laxis subpaniculatis, capitulis solitariis aut paucis aggregatis, invol. squamis linearibus cuspidato-acuminatis flores æquantibus, recept. in ambitu annuliformi in centro convexo. 2/ in Indiâ orient. ad Monghir, Dhupari, Assufgur, Hurdur, etc. Pappus pallidè rufus. *Achænia compressa*. Flor. fœm. innumeri filiformes, masc. 15-20. Affinis *C. veronicæ-foliæ*. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

38. *C. SEMIPINNATIFIDA* (Wall. cat. n. 3058. comp. 168), hirsuta, caule herbaceo erecto tereti apice ramoso-corymboso, foliis obovato-oblongis obtusis basi attenuatis grossè et obtusè dentatis aut infernè incis, summis oblongis hinc indè acutè dentatis, corymbo laxo, capitulis pedicellatis, invol. squamis linearibus acutis disco brevioribus ciliatis demùm reflexis, recept. valdè prominulo convexo-depresso. — in Indiæ or. prov. Martaban ad ripas flum. Saluen legit cl. Wallich. Fol. 2 poll. longa, 8-9 lin. lata. Flor. flavi, fœm. innumeri teretes obliquè truncati, stigm. non exsertis, masc. circ. 12, anth. inclusis. *Achænia compressa*. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

39. *C. VERONICÆFOLIA* (Wall. cat. et herb. n. 3005. comp. n. 115), caulibus ascendenti-erectis teretibus parcè ramosis pilosis, foliis pilosis infer. oblongis *subacutis* basi in petiolum attenuatis grossè serratis, superioribus semiamplexi-

caulibus lineari-lanceolatis acuminatis apice serratis, corymbis terminalibus congestis 12-20-cephalis, invol. squamis linearibus acuminatis dorso scabris demum reflexis disco æqualibus. — in Nepaliâ legit indef. Wallich. *Conyza* congesta Spreng. in litt. ad Wallich. Collum lignosum. Radix gracilis. Caulis 6-12 poll. Cor. fœmin. innumeræ truncatæ albide stylo bifido, masc. purpureæ circ. 10. Achænia compressa ut in *Erigeronte*. Pappus rufus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

40. *C. PETIOLARIS* (Wall. ! cat. n. 3282. comp. 292), sparsè pilosa, caule herbaceo erecto tereti vix apice ramoso, foliis infer. petiolo longo villosissimo donatis ovalibus obtusis grossè dentatis basi cuneatis, summis sessilibus oblongis serrulatis acutis, capitulis ad apices caulis et ramorum congestis subcorymbosis, invol. squamis linearibus acuminatis disco æquantibus. (1) in Nepaliâ? Recept. nudum medio pilosiusculum. Flor. fœm. innumeri filiformes stylo exserto bifido, masc. pauci. Achænia compressa obovalia. Pappus rufus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
41. *C. MULTICAULIS*, caulibus herbaceis è collo plurimis strictè erectis simplicibus teretibus basi pilosis apice puberulis scabridis, infimis basi breviter attenuatis, summis sessilibus subamplexicaulibus, corymbo racemoso conferto, invol. squamis linearibus acuminatis flores subæquantibus, recept. convexo fere hemisphærico. — in Nepaliâ legit cl. Wallich. *Conyza stricta* Wall. ! cat. et herb. n. 3044. comp. 154. non Willd. Pappus rufescens. Achænia glabra compressa. Flor. fœm. innumeri tenuissimi truncati, masc. in centro pauci. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
42. *C. VISCIDULA* (Wall. cat. et herb. n. 3006. comp. n. 116), caule herbaceo erecto tereti infernè glabro in ramulis breviter puberulo viscidulo, foliis ramis lineari-lanceolatis utrinquè acuminatis serratis puberulo-viscidulis, paniculâ laxâ foliis intermixtâ, capitulis pedicellatis, invol. 1-seriali 10-phylo squamis linearibus acuminatis disco æqualibus dorso puberulis. — in Indiâ or. ad Noakote legit cl. Wallich. Herba videtur elata sed pars inf. mihi ignota. Caulis pennæ anserinæ crassitie et ultrâ. Flor. fœm. innum. supra modum tenuissimi, masc. purpurascens circ. 5. Pappus rufescens. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
43. *C. STRIATA* (Wall. ! cat. et herb. n. 3065. comp. n. 175), glabriuscula aut supernè minutissimè pubens, caule herbaceo erecto striato-nervoso ramoso, foliis lanceolatis aut oblongo-linearibus utrinquè acuminatis integerrimis aut vix subserratis, ramis foliosis in paniculam dispositis corymbo conferto polycephalo terminatis, invol. squamis dorso puberulis lanceolato-linearibus acuminatis disco brevioribus. — in Indiâ orient. ad Prome et in prov. Silhet legit cl. Wallich. Folia 1-2 poll. longa, 3-4 lin. lata. Flores fœm. innumeri tenues truncati, masc. 4-5. Pappus in spec. Promensi pallide rufescens, in Silhetensi rufus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
44. *C. ANGUSTIFOLIA* (Ham. ! in Wall. ! cat. et herb. n. 3071. comp. n. 181. non Lam.), caule herbaceo erecto simplici tereti substriato patentim piloso, foliis lineari-lanceolatis elongatis acuminatis basi attenuatis ad medium grossè paucidentatis secus nervum medium subtus villosis cæterum pubescentibus, summis linearibus integerrimis, paniculâ elongatâ laxâ, ramis foliosis, invol. squamis linearibus acuminatis florum longitudine. — in Indiâ or. sed locus ign. Herba elata subcinerea. Fol 5 poll. longa, 4-5 lin. lata. Flor. flavi, ext. fœm. filiformes, inter. circ. 10-12. Achænia compressa. Pappus albidus subflavidus. (v. s. in h. hon. Cur. Ind. or. nunc soc. lin. Lond.)
45. *C. ABSINTHIFOLIA* (DC. in Wight contrib. p. 16), pilosiusculo-pubescent erecta ramosissima, foliis obovato-oblongis mucronatis basi cuneato-atenuatis, inf. grossè pauci-dentatis, sup. integerrimis, paniculâ corymbosâ ramosissimâ polycephalâ, inv. squamis linearibus acuminatis floribus brevioribus, fl. fœm. numerosissimis vix ligulatis. — (1) in Indiâ or. montanis in Nepaliâ, Kamaon, Sirmore, Dendigul, Neelgerry (ubi dic. ex Lesch. *Outoua-Gueda*), peninsulâ Indiæ or. (Wight ! herb. 217), in prov. Indiæ bor.-occid. (Royle !

pl. exs. n. 132 et verisim. 129, 130 et 131). *Conyza micrantha* Spreng. in litt. ad Wall. *Baccharis trifurcata* Trevir. in nov. act. acad. nat. cur. 13. p. 201. *Conyza cuneata* Ham! herb. *Conyza montana* et *C. stricta* Heyn! herb. *Conyza pinna-tifida* var. Roxb. cat. calc. 61. *Conyza trifida* Ham. ex Don. *Erigeron pinna-tifidum* var. Don. prod. 172 ex Wight! herb. Ambigit inter *Erigeron* et *Conyza*. Habitus *Erig.* aeris sed ligulæ nullæ. *Achænium* parvum glabrum lenticulare compressum. Capitula parva. Pappus rufescens. Herba variat *foliis* angustis et latioribus, integris dentatis aut subpinnatifidis, obtusis aut mucro-natis. An fortè plures hic latent commixtæ? (v. s.)

46. *C. WALLICHII*, caule herbaceo erecto tereti superne ramoso-corymboso gla-bro, ramulis ultimis pulverulento-puberis, foliis lanceolatis acuminatis basi attenuatis serratis glabris, paniculæ corymbosæ polycephalæ ramis sæpè bifidis, capitulis ad apices ramulorum sessilibus confertis, invol. squamis linearibus acuminatis disco paulò longioribus. — in Indiâ orient. *Conyza disticha* Wall.! cat. et herb. n. 3024. comp. n. 134 non Spreng. Flor. fœm. innumeri, mihi vix ex statu provento specim. noti, masc. 5-7. *Achænia* parva ovalia compressa glabra. Pappus rufus.

α. serrata, foliis ovali-lanceolatis grossè serratis. — *Conyza disticha* A. Wall.! l. c. *Caecalia disticha* Heyn.! herb. (v. s. comm. ab hon. Cur. Ind. or. et à cl. Royle n. 128.)

β. serrulata, foliis elongato-linearilanceolatis tenuissimè serrulatis. — circa Oude legit cl. Wallich cat. l. c. litt. B. (v. s. comm. cum *α.*)

47. *C. TRAVANCORENSIS* (Wall.! cat. n. 3067. comp. 177), caule herbaceo erecto tereti subtilissimè striato et puberulo, foliis subpetiolatis lanceolatis utrinquè acuminatis subserrulatis glabris cum nervo medio subtus puberulo, ramis corymbosis oligocephalis, capitulis pedicellatis, invol. 1-2-serialis squamis lan-ceolatis acuminatis flores subsequantibus. — in Indiâ or. ad Travancore. Folia 3-4 poll. longa, 9-10 lin. lata. Recept. convexum subhemispharicum. Flor. fœm. innum. filiformes, masc. 4-5. Pappus rufescens. *Achænia* ex immaturo statu non satis nota. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)

β. brevifolia, ramulis teretibus vix puberulis, foliis subsessilibus ovali-lan-ceolatis acutis, capitulis magis corymboso-confertis, invol. squamis magis acuminatis. — in Indiâ or. prov. boreali-occid. legit cl. Royle. Folia sup. pollicem longa, 5 lin. lata. *Achænia* minima compressa glabra. An species propria? (v. s. comm. à cl. Royle sub n. 120.)

48. *C. RUFA* (Wall.! cat. et herb. n. 3079. comp. n. 189), caule herbaceo erecto tereti simplici à basi multiplici piloso, foliis villosis oblongis obtusè et inæqua-liter dentatis obtusis, infer. basi in petiolum attenuatis, supremis basi cuneatis sessilibus, paniculâ corymbosâ, ramis apice corymbulosis, capitulis pedicellatis, inv. squamis ovatis latè albo-marginatis biserialibus cuspidatis floribus æqua-libus. — in Ind. orient. ad Assufghur legit cl. Wallich. Herba pedalis. Folia sesquipoll. longa, 4-5 lin. lata. Pappus rufus. Flor. fœm. innumeri, masc. 4-5. *Achænia* ex statu imperf. ignota. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

*** *Species Madagascarienses aut Mauritiânæ.*

49. *C. BOJERI*, suffruticosa erecta ramosa glabra, ramis teretibus, foliis oblongo-lanceolatis sessilibus basi attenuatis integerrimis aut hinc inde subdentatis, pedunculis axillaribus terminalibusve nudis minutissimè pubescentibus capi-tula 7-8 congesta gerentibus, invol. squamis ovatis obtusiusculis disco dimidio brevioribus. ♂ in sylvis Béfouroun dictis prov. Bé-tani-mena ins. Madagascar legit cl. Bojer. Folia 2 poll. longa, 7 lin. lata. Flores flavi. *Achænia* glabra. (v. s. comm. à cl. Bojer.)

50. *C. ATTENUATA* (DC. non Wall.), fruticosa glabra, ramis teretibus apice sub-herbaceis flexilibus, foliis sessilibus elongato-lanceolatis basi obtusis apice acuminatissimis serratis, corymbo composito polycephalo laxiusculo, invol. *squamis linearibus acuminatis nitidulis*. ♂ in sylvis umbrosis Béfouroun dictis

prov. Bé-tani-mena ins. Madagascar legit cl. Bojer. Baccharis attenuata Bojer! in litt. 1835. Folia 2½ poll. longa, 5 lin. lata. Pappus rufescens. Invol. squamæ nervo medio virescente, marginibus pallidis. (v. s. comm. à cl. Bojer.)

51. *C. VIRGATA* (DC. non Linn.), fruticosa erecta, ramis strictis virgatis teretibus pubescenti-hirtellis, foliis sessilibus linearilanceolatis acutis serratis pubescentibus subrubs punctulatis, corymbo irregulari composito, capitulis ad apices pedunculorum congestis subsessilibus, invol. squamis rufo-scariosis acutis, fl. disci circ. 5. — ½ in Madagascar detexit cl. Bojer. Baccharis virgata Bojer! in litt. 1835. Frutex 4-5-pedalis. Folia circ. 20 lin. longa, 4 lin. lata. Achænia compressa glabra. Flores mihi filiformes truncati, ex Bojer flavi marginales sæpè ligulati. An ideò Nidorellæ fortè species? (v. s. comm. à cl. inv.)

52. *C. FLEXILIS*, fruticosa glabra subscandens, ramis teretibus flexilibus, foliis sessilibus oblongo-lanceolatis basi attenuatis apice acuminatis integerrimis, paniculâ elongatâ, capitulis ad apices ramulorum congestis subsessilibus parvis, invol. squamis glabris scariosis rufescentibus acutis. ½ in sylvis Bèfouroun dictis ins. Madagascar prov. Bé-tani-mena super arbores subscandens. Baccharis Bojer! in litt. 1835. Flores flavi. Capitula parva in spec. juniora nec ideò satis ritè observata. Pappus rufus brevis. Folia 3 poll. longa, 8 lin. lata.

53. *C. HIRTELLA*, herbacea subscandens, ramis densè hirtellis, foliis petiolulatis lanceolatis utrinquè acuminatis remotè serratis utrinquè minutissimè scabridis, paniculis parvis axillaribus folio brevioribus, invol. squamis lanceolatis acuminatis scariosis rufescentibus glabris. ①? in sylvis ins. Madagascar cum *C. flexili* legit cl. Bojer. Folia (petiolo 2-3 lin. incluso) 3 poll. longa, 8-9 lin. lata. Capitula parva. Flores juniores non ritè exam. (v. s. comm. à cl. Bojer.)

- 54? *C. AGERATOIDES*, suffruticosa erecta ramosa infernè glabrata, ramis densè hispidis cylindricis, foliis petiolatis ovatis acutis grossè dentatis utrinquè hirtellis, petiolis densè hispidis, capitulis ad apices pedunculorum 7-8 congestis, invol. squamis glabris linearibus acuminatis, flor. fœmineis pauciseriatis. ½ in campis et agris cultis prov. Emirnæ insul. Madagascar legit cl. Bojer. Baccharis ageratum referens Bojer! An flores (mihi in specim. juniore non satis noti) fortè cœrulei? An à genere removenda et fortè ad Plucheam refer. ? (v. s. comm. à cl. Bojer.)

55. *C. LINEARELOBA*, caule erecto basi fruticuloso, ramis longè et densè hispidis, foliis parcius hispidis linearibus pinnatipartitis lobis 2-3-jugis distantibus linearibus integerrimis, summis indivisis, capitulis racemoso-glomeratis brevissimè pedicellatis, invol. squamis linearibus acuminatis dorso puberulis, fl. disci paucis. ½ in insulis Madagascar, Mauriti et Borboniæ, præcipuè in agris cultis legit cl. Bojer. Flor. fœm. innumeri ut centrales pappo 1-seriali longo subrufescente donati. Recept. foveolatum. Achænia parva compressa glabra. (v. s. ex Madag. comm. à cl. Bojer.)

56. *C. CALOCEPHALA* (Bory! voy. isl. afr. 2. p. 395), suffruticosa tota hirsuto-hispida, foliis sessilibus oblongis integerrimis aut hinc indè subserratis, pedunculis folia superantibus 1-cephalis, invol. squamis biserialibus acuminatis dorso hirsutis, fructibus compressis obovatis, pappo rufo-colorato flosculis longiore. ½ in insulâ Borboniæ (Comm.! Bont!), in montanis prope locum dict. Bras-de-Ponteau (Bory!). Erigeron rutilum var. Poir. suppl. 5. p. 494. Dimorphanthès. bidentata Cass. dict. 13. p. 256, et 37. p. 481. Suffrutex vix palmaris. Folia erecta 8-10 lin. longa, 2-3 lata, basi subattenuata utrinquè hirsuto-hispida. Achænia glabriuscula compressa. Rec. brevissimè fimbriatæ. (v. s.)

57. *C. ASPERA* (Bory! voy. isl. afr. 3. 174), suffruticosa tota hirsuto-hispida, foliis sessilibus semiaimplexicaulibus oblongo-lanceolatis grossè serratis trinerviis, capitulis ad apices ramorum 4-5 aggregatis breviter pedicellatis, invol. squamis acuminatis dorso hirsutis, achæniis compressis oblongis glabris, pappo rufo flosculis longiore. ½ in insulâ Borboniæ montanis (Comm.! Bory!). Conyza melastomoides Lam. dict. 2. p. 90. Erigeron scabrum Pers. ench. 2. p. 432. n. 35 nec n. 23. Erigeron Jussæi Spreng. syst. 3. p. 519. Valdè affinis *C. ca-*

locephalæ et *C. lithospermifoliæ*. Achænia compressa. Recept. brevissimè fimbriiferam. (v. s.)

58. *C. LITHOSPERMIFOLIA* (Lam. dict. 2. p. 90), suffruticulosa pubescenti-hirsutula, foliis sessilibus semi-amplexicaulibus oblongis apice serratis basi trinerviis, ramis subvirgatis apice aphyllis 1-cephalis, invol. squamis acuminatis dorso puberulis, achæniis compressis glabriusculis, pappo rufo corollis longiore. $\frac{5}{8}$ in summis montibus insulæ Borboniæ (Bory! Bont!). *Elphegea hirta* Cass. dict. 14. p. 364 ex syn. Valdè affinis *C. asperæ* et *C. calocephalæ* sed distincta videtur. Recept. brevissimè fimbriiferum. (v. s.)

59. *C. SERICEA* (Bory! voy. isl. afr. 2. p. 376 non Ait.), suffruticulosa tota villis adpressis sericeis cinerea, foliis sessilibus oblongis basi ferè angustioribus mucronatis integerrimis, pedicellis folio paulò longioribus 1-cephalis ad apicem circ. 3 approximatis, invol. squamis dorso sericeo-villosis subacuminatis, pappo rufo corollis vix longiore. $\frac{3}{8}$ in insulâ Borboniâ legit cl. Bory et mecum comm. Hæc et seq. commixtæ adsunt in h. Desf. sub nom. autographo *Eriogon rutilans* Poir.! suppl. Ferè media inter *C. calocephalam* et *C. argenteam*. (v. s.)

60. *C. ARGENTEA* (Lam. dict. 2. p. 87), suffruticosa tota villis adpressis sericeis argentea, foliis ovatis semiamplexicaulibus integerrimis mucronulatis, capitulis ad apices ramorum subglomeratis breviter pedicellatis, invol. squamis sericeo-villosis acuminatis, pappo rufo flosculis subæquali. $\frac{5}{8}$ in insulâ Borboniâ collibus aridis. Folia obscurè trinervia 9 lin. longa, 6 lin. lata. Achænia compressa glabra. (v. s.)

61. *C. ANCHUSÆFOLIA* (Poir. suppl. 2. p. 339), suffruticosa procera hirsuta, foliis ovali-oblongis utrinquè acuminatis in petiolum breviter attenuatis apice serratis penninerviis, corymbo paniculato terminali polycephalo foliis breviorè, pedicellis capitulo æqualibus, invol. squamis extus glabriusculis, achæniis oblongis glabris, pappo corollæ longitudine subrufescente. $\frac{5}{8}$ in ins. Borboniâ (Bory! Bont!). Conyza verbascofolia Bory! voy. isl. afr. 2. p. 377 non Willd. Folia 5 poll. longa, 15 lin. lata. Flores rubescentes. Achænia compressa glabra. (v. s.)

62. *C. AMYGDALINA* (Lam. dict. 2. p. 90), fruticosa villis longis adpressis sericeis subcanescentis, foliis petiolatis oblongo-lanceolatis utrinquè acuminatis dentata-serratis penninerviis suprà subviridibus subtus sericeo-argenteis, pedunculis axillaribus folio multò brevioribus capitula 3-4 congesta gerentibus, invol. squamis dorso subhirsutis acuminatis, pappo subrufescente corollæ longitudine. $\frac{5}{8}$ in insulâ Borboniâ prope Cafrorum planitiem. Folia cum petiolo semipollicari 4 poll. longa, pollicem lata. Achænium ignotum et idèò genus dubium. (v. s. comm. à cl. Bory.)

Sect. III. TUBIFERA.

Recept. fimbriiferum, fimbriis elongatis. Flores fœm. tubulosi cylindracei apice brevissimè 3-dentati discum superantes. — Species Borbonica.

3. *C. LAURIFOLIA* (Lam. dict. 2. p. 89), fruticosa pube brevissimâ subferrugineâ pubescens, foliis oblongo-lanceolatis in petiolum attenuatis subintegerrimis penninerviis, corymbo paniculato amplo, ramis elongatis, pedicellis ultimis vix capitulo longioribus, invol. squamis extus pubescentibus margine membranaceis, pappo rufescente corollis paulò breviorè. $\frac{5}{8}$ in insulâ Borboniæ sylvis (Bory! Bont!). *Fimbrillaria tubifera* Cass. dict. 17. p. 55. Folia 4-5 poll. longa, pollicem lata. Flor. fœm. pluriseriales tubulosi elongati 3-dentati stylo non exserto, masc. pasci anth. semiexsertis basi muticis. Achænia ex statu immat. sericeo-villosa compressa. Pappus rufus. Recept. fimbriis longis acutis onustum. (v. s.)

Sect. IV. FIMBRILLARIA. — *Fimbrillaria Cass. dict.* 17. p. 54.

Recept. totum fimbrilliferum, fimbrillis in centro longioribus sæpè setiformibus. — Species Capensæ.

64. *C. PODOCEPHALA*, caule imâ basi suffruticoso multicauli subadscendente, ramis hispido-scabris, foliis obovatis basi cuneatis integris apice grossè dentatis sparse scabro-hirtellis, summis oblongis agutis, ramis apice nudis 1-cephalis parçè hispidis, invol. squamis lineari-subulatis glabrisculis disci longitudine. $\frac{3}{4}$ in Africa australi ad Zw. Stoorm et Witbergen alt. 4-500 ped. legit cl. Drege! Recept. alveolatam brevissimè fimbrilliferam. Cor. masculæ glabræ numero fœm. subæquales. (v. s.)

ß. demissa, caulibus pluribus demissis folisque magis hispidis. — ad Nieuwewelde legit cl. Drege. (v. s.)

65. *C. PINNATILORATA*, caule herbaceo erecto vix puberulo, foliis radicalibus petiolatis, caulinis sessilibus semiamplexicaulis, omnibus glabrisculis pinnatipartitis, lobis lanceolatis subdentatis in apice caulis integerrimis margine et secus nervos scabris, corymbo paniculato laxo terminali, pedunculis pubescentibus, achæniis glabris. $\frac{1}{2}$ ad Cap. Bonæ-Spei post Thunb. legitur cl. Burckell (cat. geog. n. 4233!), Belanger (n. 80), Drege, Ecklon et Zeyh. *Erigeron pinnatum* Linn. f. suppl. 368. Thunb. prod. 153. fl. cap. 666. *Baccharis leucanthemifolia* Burm.! herb. et ideò ex aphan. *Xeranthemum leucanthemifolium* Burm. prod. 26? *Herba sesquipedalis*. Fol. rad. lobi ferè ovati sinuati obtusi. *Achænia compressa* ut in *Erigeronte*. (v. s.)

ß. hispida, caule magis hispido, foliis vix ad medium incisæ, capitulis paucioribus majoribus. $\frac{1}{2}$ in Witbergen alt. 3-400 ped. legit cl. Drege. (v. s.)

66. *C. PINNATIFIDA* (Less. syn. 204 non Roxb.), herbacea tota pilis patulis strigosis hirta, foliis caulinis pinnatipartitis lobis lanceolato-linearibus hinc inde dentibus lobulise acutis distantibus incisæ, capitulis 1-2 ad apices ramorum, achæniis puberulis. $\frac{1}{2}$ ad Cap. Bonæ-Spei (Drege!) in collibus extra urbem (Th.), in planitie prope Zeekuhwallei (Ecklon! in h. Dunant), in Stellenbosch (Eckl.). *Erigeron pinnatifidum* Thunb. fl. cap. 666. (v. s.)

67. *C. OBSCURA*, herbacæ erecta villis confertis subrigidis hirsuta, foliis obovato-oblongis sessilibus basi attenuatis subintegris semiamplexicaulis apice pinnatifidis, lobis obtusis aut in parte caulis super. acutis, corymbis laxiusculis, achæniis subpilosisculis. $\frac{1}{2}$? in Africa australi. Drege! Eckl. et Zeyh. (v. s.)

ß. calvescens, foliis infer. hirsutis, super. cauleque scabris, lobis acutis margine subrevolutis. $\frac{1}{2}$? in Lange-Kloof legit cl. Drege. An spec. propr.? (v. s.)

68. *C. INCISA* (Ait. kew. 3. p. 184), caule herbaceo? erecto puberulo, foliis petiolatis ovatis aut subcordatis subpiloso-viscosis dentato-incisæ basi subauriculatis, corymbis terminalibus laxiusculis, achæniis glabris. $\frac{1}{2}$ $\frac{3}{4}$ ad Cap. Bonæ-Spei. *Erigeron incisum* Th. fl. cap. 666. *Conyza incisa* et *Erigeron incisum* Willd. sp. 3. p. 1945 et 1962. *Baccharis ulmifolia* Burm.! prod. fl. cap. 26. *Achænia glabra*, compressa ut in *Erigeronte*. Recept. breviter fimbrilliferum.

a. stipulacea, petiolo supra medium appendice utrinquè aucto, limbo terminali subcordato. — ad Zneurebergen Drege! *Conyza stipulacea* Desf. cat. h. Paris. 1829. p. 164. excl. patr. *Dimorphanthus stipulacea* Cass. dict. 25. p. 94. excl. patr. (v. s.)

ß. hirta (Less. syn. 204), caulibus folisque magis hirsutis petiolo exappendiculato, limbo ovali. — ad Cap. Bonæ-Spei (Boiss.! herb.), in Tullagh, Uitenhage, Clan-William (Eckl. et Zeyh.) in Paarl (Drege!) *Erigeron hirtus* Th. fl. cap. 665. (v. s.)

69. *C. SCABRIDA*, tota pubescenti-scabra, caule subherbaceo erecto, foliis penninerviis lanceolatis acuminatis dentatis, inf. petiolatis, super. sessilibus brevissimè

auriculatis semiamplexicaulibus, corymbo terminali composito. § 24 ad Cap. Bonæ-Spei (Drege!) in distr. Uitenhage (Eckl. et Zeyh.). An *Erigeron dentatum* Burm.! prod. 26? Media inter *C. incisam* à quâ differt foliis lanceolatis 1½-2 poll. longis, 4 lin. latis, et *C. ivæfolia* à quâ distinguitur foliis pubescenti-scabris nec glabris penninerviis nec trinerviis. Achænia compressa. Rec. in ambitu breviter alveolatum medio setis elongatis onustum. (v. s.)

β. *angustifolia*, foliis linearibus remotius dentatis. — Borch.! n. 382. Eckl. ! n. 771. (v. s.)

70. *C. IVÆFOLIA* (Less. in Linnæa 1831. p. 138), fruticosa, ramis teretibus pubescentibus, foliis lanceolatis acuminatis in petiolum attenuatis grossè serratis basi trinerviis glabris, corymbo terminali composito. § ad Cap. Bonæ-Spei in collibus ad occid. montis Tabularis (Th.), ad Dutois-Kloof (Drege!). *Baccharis ivæfolia* Linn. sp. 1204. excl. syn. et patria. Berg. cap. 275. Thunb. fl. cap. 666. Burm.! prod. 26. *Fimbrillaria baccharoides* Cass. dict. 17. p. 54. Pluk. amalth. t. 454. f. 10. pessima et dubia. Barth. act. hafn. 2. t. 27. Linnæus, qui citat phrasin Pluken. ubi dicitur Africana, asserit suam stirpem ex Virginia et Peruv. ortam, sed fortè sub eo nom. *Baccharidem angustifoliam* designavit. Nostra certè Capensis ex spec. spontaneo Drege à me suprâ citato et altero Eckloniano in herb. Dnnant servato, hortensibus simillimis. Recept. fimbriillis in setis ferè desinentibus onustum. (v. s. sp. et v. c.)

71. *C. POSILLA* (Houttuyn! hist. nat. x. bl. 618. t. 69. f. 1), hispido-villosa cinerea herbacea ramosa, foliis superis sessilibus ovalibus mucrone apiculatis integerrimis aut ad apicem mucronibus 1-2 auctis basi trinerviis, pedicellis 1-floris folii longitudine in racemum laxum dispositis, inv. squamis linearibus acuminatis puberis demum reflexis. — ad Cap. Bonæ-Spei. Frustulum tantum vidi sed meo sensu distinctissimum. (v. s. in h. Delessert.)

† *Conyza auctorum non satis notæ pleræque verisimiliter è genere excludendæ.*

* *Gerontogea.*

72. *C. 2 ARABICA* (Willd. sp. 3. p. 1949), fruticosa stricta pubescens supernè tomentosa, foliis decurrentibus linearibus integerrimis tomentosis, capitulis pedicellatis paniculatis, invol. squamis setaceis. § in Arabiâ felici. *Erigeron decurrens* Vahl symb. 1. p. 72. An *Pluchea*, an *Blumea* spec.?
73. *C. DONGOLENSIS* (in Caill. Delil. pl. afr. voy. mer. p. 43), ramis villosis, foliis sessilibus oblongis dentatis sublyratis basi incis vel pinnatifidis, segmentis acutis. — circa Dongolah in Nubiâ. Flor. fructusque ignoti et ideò genus dubium.
74. *C. ? CRISPATA* (Vahl symb. 1. p. 71), fruticosa glabra, foliis decurrentibus lanceolatis serrato-dentatis, alis dentato-sinuatis, pedunculis 1-cephalis. § in Arabiâ. *C. caule alato* Forsk. cat. pl. ar. p. 119. An *Pluchea*? An *Blumea* sp.?
75. *C. ? FOLIOLOSA* (Willd. sp. 3. p. 1919, non Wall.), pubescens, caule herbaceo ramosissimo diffuso, foliis oblongis subintegerrimis subamplexicaulibus, pedunculis alaribus 1-cephalis demum elongatis folio sextuplò longioribus. — in Indiâ or. Con. foliosa Pers. ench. n. 3. An *Conyza*? an *Blumea*? an *Pluchea*?
76. *C. ? HETEROPHYLLA* (Lam. dict. 2. p. 84), caule striato herbaceo ramoso, foliis petiolatis cinereo-villosis aliis subcordatis aliis rotundioribus, paniculis terminalibus. — In Indiâ or. (Sonn.), in agro Malabarico (Rheed. mal. 10. t. 64). An *Vernonia* species cinereæ affinis?
77. *C. INCANA* (Willd. sp. 3. p. 1937), fruticosa tomentosa, foliis linearibus basi attenuatis subdentatis tomentosis, corymbo terminali. § in Arabiâ felici, *Erigeron incanum* Vahl symb. 1. p. 72. Invol. squamæ lineares. Pappus ferrugineus. An *Blumea*?
78. *C. ? LINIFOLIA* (Burm. prod. fl. cap. 26 non Linn.), vid. *Blumea* sp. ex capitulo à me olim dissecto, sed vix nota!
79. *C. ? LOUREIRI* (Poir. suppl. 2. p. 339), caule suffruticoso erecto simplici, foliis

- lanceolatis integerrimis breviter petiolatis subtus tomentosis, pedunculis axillaribus polycephalis, invol. cylindrico, recept. nudo. ζ prope Cantonem Sinarum. Baccharis Sinensis Lour. coch. 2. p. 604. Conyza Chinensis Pers. ench. n. 16 nec n. 21. An Conyza? an Blumea?
80. *C. MOLUCCANA* (Gaud. voy. p. 465), glabra, ramis suffrutescentibus sulcatis, foliis oblongis acuminatis basi obtusis argutè et remotè denticulatis subtus pallidioribus, paniculis terminalibus foliatis, involucribus pubescentibus. ζ in ins. Moluccis ad Rawak. An Blumea sp.?
81. *C. ORIENTALIS* (Willd. sp. 3. p. 1931), herbacea hispido-scabra supernè ramosa, foliis serratis, infer. obovatis petiolatis, superioribus lanceolatis sessilibus, capitulis congesto-corymbosis terminalibus, invol. squamis subulatis. \mathcal{Z} in Armeniâ. *C. orient.* Asteris attici folio flore luteo Tournef. cor. 35.
82. *C. PUBIGERA* (Linn. mant. 113), caule herbaceo sparsè piloso, foliis oblongis subdentatis subpetiolatis, pedunculis lanatis subdicephalis. — in Ind. orient. Sonchus volubilis Rumph. amboin. 5. p. 299. t. 103. f. 2, è Moluccis ortus et ad suam speciem à Linnæo relatus vix descriptioni respondet et ex icone videtur potius Senecioinea?
83. *C. STRICTA* (Willd. sp. 3. p. 1922), caule herbaceo erecto paniculato, ramis erectis, foliis lineari-lanceolatis basi attenuatis integerrimis pilosis obtusis cum acumine, corymbis fastigiatis. (1) in Indiâ orientali. Pappus rufescens. An Conyza? an Blumea?
84. *C. THYRSOIDEA* (Pers. ench. 2. p. 426. n. 8), herbacea, caule villosa, foliis pubescentibus obovatis sinuato-serratis infernè sublyratis, racemis foliosis terminalibus. — in Senegaliâ. Cæt. ign.
- ** *Americanae.*
85. *C. CAYENNENSIS* (Spreng. syst. 3. p. 512), herbacea, foliis longè petiolatis subcordato-ovatis acuminatis glabriusculis venosis obtusè serratis, paniculâ terminali, capitulis subsessilibus. — in Cayennâ ex Martin in herb. Willd. Cæt. ign. \dagger .
86. *C. CONICA* (Poir. suppl. 2. p. 341), fruticosa glabra, foliis subsessilibus coriaceis ovato-subrotundatis integris suprà lucidis subtus pallidis, capitulis cymosis subsessilibus, invol. conici squamis acutis, pappo intense rufo. ζ patr. et cæt. car. ign.
87. *C. DISTICHA* (Spreng. syst. 3. p. 509), fruticosa, foliis sessilibus oblongis dentatis subcoriaceis ramisque villosiusculis, spicis distichis foliatis terminalibus. ζ ad flumen magnum Brasiliæ. Cæt. ign. An Vernoniæ spec.? \dagger .
88. *C. GRAMINIFOLIA* (Spreng. syst. 3. p. 515), herbacea, caule glaberrimo simpliciori, foliis basi vaginantibus linearibus acuminatis elongatis striatis, pedunculis terminalibus corymbosis squamatis, invol. squamis lanceolatis cuspidatis. — in Monte-Video. Cæt. ign.
89. *C. LAXA* (Spreng. syst. 3. p. 513), herbacea, foliis glabris petiolatis oppositis ovato-lanceolatis utrinquè attenuatis serratis margine scabris tripinerviis, corymbis axillaribus terminalibusque oligocephalis, invol. squamis laxis linearibus. — in Brasiliâ. An Baccharidea? an Eupatoriacea?
90. *C. LUCENS* (Poir. suppl. 2. p. 342), fruticosa tortuosa, ramis teretibus tomentosis, foliis petiolatis ovatis integris suprà lucidis subtus incano-tomentosis, capitulis terminalibus agglomeratis, invol. squamis linearibus. ζ in Sancto Domingo. *C. Domingensis* Willd. herb. ex Spreng. syst. 3. p. 508. Cæt. ign. \dagger .
91. *C. MACROPHYLLA* (Spreng. syst. 3. p. 513), herbacea, foliis amplexicaulibus longissimis lanceolatis glabris sinuato-dentatis, paniculâ corymbosâ, invol. squamis lineari-lanceolatis. — in Monte-Video. Cæt. ign. \dagger .
92. *C. MONTEVIDENSIS* (Spreng. syst. 3. p. 510), suffrutescens glabra, foliis linearibus acuminatis elongatis serratis, capitulis corymbosis terminalibus. ζ in Monte Video. An Baccharidis spec.? \dagger .
93. *C. OBLONGIFOLIA* (Spreng. syst. 3. p. 512), herbacea, caule simplici stricto

superne aphyllto, foliis glaberrimis subintegerrimis oblongis obtusis sessilibus triplinerviis, inf. oppositis, paniculâ corymbosâ pubescente, invol. squamis oblongis obtusis. — in Brasiliâ ad Rio-Grande. Cæt. ign. †.

94. *C. PANAMENSIS* (Willd. sp. 3. p. 1942), fruticosa, ramis glabris striatis, foliis sessilibus ovato-lanceolatis scabris integerrimis triplinerviis reticulato-venosis, paniculâ terminali, invol. squamis oblongis. ♀ in Panamâ. Ex descr. valde manca videtur potius *Baccharis*?

95. *C. PUNCTATA* (Willd. sp. 3. p. 1937), fruticosa, ramis striatis glabris, foliis linearibus acuminatis basi attenuatis impresso-punctatis, pedunculis longissimis 1-cephalis. ♀ in Chili. Invol. squamæ lanceolatæ acutæ. Pappus rufus. Cæt. ign. An fortè *Baccharidis* spec.?

96. *C. RUFINULA* (Spreng. men. entd. 2. p. 141. syst. 3. p. 513), caule herbaceo erecto piloso, foliis brevissimè petiolatis oblongo-lanceolatis acutis utrinquè pubescentibus integerrimis, paniculâ pyramidalî. — in Brasiliâ. Folia sub lente denticulato-ciliata. Involucrum imbricatum glabrum squamis inter. elongatis coloratis. Pappus rufescens. Flores non observ. †.

97. *C. SQUMATA* (Spreng. syst. 3. p. 515), herbacea, caule ramosissimo scabro, foliis sessilibus linearibus acuminatis integerrimis, pedunculis squamatis 1-cephalis, invol. squamis lineari-lanceolatis. — in Monte-Video. Cæt. ign.

98. *C. TRIDENTATA* (Poir. suppl. 2. p. 342), fruticosa glaberrima, foliis confertis linearibus basi attenuatis subpetiolatis apice subtridentatis obtusis, capitulis parvis sessilibus. ♀ in Sancto-Domingo. Cæt. ign. An *Baccharidis* species? †.

*** *Patriæ non satis notæ.*

99. *C. CILIARIS* (Spreng. nov. prov. 14. syst. 3. p. 509), fruticosa, foliis spathulato-lanceolatis subsessilibus glabris ciliatis, pedunculis terminalibus solitariis elongatis 1-cephalis villosis. ♀ Patria et cæt. car. ign.

100. *C. GLOMERATA* (Link enum. h. Berol. 2. p. 322), caule herbaceo alato, foliis latè lanceolatis obtusis basi attenuatis serrulatis pubescentibus scabris, capitulis glomeratis bracteis obvallatis. — Patr. ign. An *Blumea*, *Pluchea* seu *Monentelis* spec.?

101. *C. LAXIFLORA* (Desf. cat. h. Par. 1830. p. 400), caule herbaceo villosio paniculato, foliis lanceolatis ciliatis, infer. remotè serratis rameis sup. lineari-lanceolatis integerrimis, capitulis paniculatis hemisphæricis, invol. squamis subulatis. ① Patr. et cæt. car. ign. †.

102. *C. PROCUMBENS* (Balb. cat. h. Taur. p. 28), caule procumbente ramoso, foliis spathulatis subtus tomentoso-niveis margine revolutis, pedunculis nudis moncephalis. — Patr. ign. *Conyza chrysocomoides* Ten. non Willd. ex Balthis. *Conyza prostrata* Ten. cat. h. Neap. 1819. p. 75?

103. *C. TORTUOSA* (Linn. hort. Cliff. 405), caule fruticoso tortuoso, foliis ovato-oblongis integerrimis, racemis reflexis. ♀ in Verâ-Cruce (Linn.), Brasiliâ (Spreng.), Madagascar (Vall. act. par. 1719. p. 396). Species mihi omninò obscura! An fortè *Vernonia*? †.

†† *Conyze exclusæ.*

C. ABBREVIATA Wall. = *Vernonia abbreviata*.

C. ACUMINATA Wall. = *Vernonia acuminata*.

C. ADNATA H. B. K. = *Pluchea subdecurrens*.

C. ALATA Roxb. = *Blumea alata*.

C. ALATA Baug. = *Inula bifrons*.

C. ALATA var. *β.* Wall. = *Blumea pterolonta*.

C. ALBICANS Wall. = *Vernonia albicans*.

C. ALOPECUROIDES Lam. = *Pterocanlon alopecuroides*.

- C. AMPLECTENS Wall. = *Blumea amplexens*.
- C. AMPLEXICAULIS Lam. = *Blumea amplexens*.
- C. AMPLEXICAULIS Wall. = *Blumea oligocephala*.
- C. AMPLEXICAULIS Michx. = *Pluchea bifrons*.
- C. ANAGALLIDIFOLIA Wall. = *Blumea anagallidifolia*.
- C. ANGUSTIFOLIA hort. Par. = *Baccharis Pingreei*.
- C. ANGUSTIFOLIA Lam. = *Monarhenus salicifolia* β .
- C. ANTHELMINTHICA L. = *Vernonia anthelmintica*.
- C. APPENDICULATA Blum. = *Pluchea appendiculata*.
- C. APPENDICULATA Lam. = *Senecio appendiculatus*.
- C. ARBOREA Wall. = *Vernonia arborea*.
- C. ARBORESCENS L. = *Vernonia arborecens*.
- C. ARBUTIFOLIA Lam. = *Baccharis arbutifolia*.
- C. ARGENTEA Wall. = *Inula argentea*.
- C. AROMATICA Wall. = *Blumea aromatica*.
- C. ARTICULATA Lam. = *Baccharis articulata*.
- C. ASPERA Wall. = *Decaneurum asperum*.
- C. ASTEROIDES Desf. = *Pluchea subdecurrens*.
- C. ASTEROIDES Lin. = *Sericocarpus comynoides*.
- C. ASTEROIDES Wall. = *Chrysopsis gossypina*.
- C. ATTENUATA Wall. = *Vernonia attenuata*.
- C. AURICULATA Mœnch. = *Phagnalon rapestræ?*
- C. AURITA L. F. = *Blumea aurita*.
- C. AXILLARIS Lam. = *Blumea axillaris*.
- C. BALSAMICA Wall. = *Psidium balsamica*.
- C. BALSAMIFERA L. = *Pluchea balsamifera*.
- C. BARBATA Wall. = *Blumea barbata*.
- C. BELLIDIFOLIA Wall. = *Vernonia bellidifolia*.
- C. BIFOLIATA Linn. = *Blumea bifoliata*.
- C. BIFOLIATA Cham. = *Blumea Chamissoniana*.
- C. BIFOLIATA Wall. = *Blumea Wightiana* β .
- C. BIFOLIATA Walt. = *Sericocarpus tortifolius*.
- C. BIFRONS Linn. = *Pluchea bifrons*.
- C. BLANDA Wall. = *Vernonia blanda*.
- C. BRACTEOLATA Wall. = *Vernonia bracteolata*.
- C. BRYOIDES Lam. = *Baccharis bryoides*.
- C. BUXIFOLIA Lam. = *Baccharis buxifolia*.
- C. CALYCINA Cav. = *Phagnalon calycinum*.
- C. CAMPHORATA Ell. = *Pluchea camphorata*.
- C. CANDIDA Linn. = *Inula candida* α .
- C. CANDIDA Willd. = *Inula candida* β .
- C. CANESCENS Bory. = *Schizogyne sericea*.
- C. CANESCENS Linn. f. = *Webbia pinifolia*.
- C. CAROLINENSIS Jacq. = *Pluchea odorata*.
- C. CERNUA Wall. = *Blumea alata* β .
- C. CHENOPODIFOLIA Lam. = *Gynura Mauritiana*.
- C. CHINENSIS Linn. = *Blumea Chinensis*.

- C. CHINENSIS Lam. = *Cyanopsis pubescens*.
 C. CHRYSOCOMOIDES Desf. atl. = *Nolletia chrysocomoides*.
 C. CHRYSOCOMOIDES Desf. arb. = *Felicia angustifolia*.
 C. CINEREA Linn. = *Vernonia cinerea*.
 C. CINEREA var. 2 Wall. = *Vernonia physalifolia*.
 C. CINERARIOIDES Bojer. = *Rochonia cinerarioides*.
 C. CINERASCENS Wall. = *Vernonia cinerea* s.
 C. CINTERÆFOLIA Willd. herb. = *Vernonia odoratissima*.
 C. COMMUNIS Wall. = *Blumea hieracifolia*.
 C. CORONOPUS Lam. = *Sarcanthemum Coronopus*.
 C. CORYMBOSA Roxb. = *Pluchea Indica*.
 C. CORTESII H. B. K. = *Pluchea Cortesii*.
 C. CUNEATA Bojer. = *Rochonia cuneata*.
 C. CUNEIFOLIA Lam. = *Baccharis cuneifolia*.
 C. CUNEIFOLIA Wall. = *Blumea cuneifolia*.
 C. CUNEIFOLIA Rensch. = *Solidago sparia*.
 C. CUPRESSIFORMIS Lam. = *Lepidophyllum cupressiforme*.
 C. DAMPIERI Cunn. = *Eurybia Dampieri*.
 C. DENSIFLORA Wall. = *Plumea densiflora*.
 C. DICHOTOMA Linn. = *Jasonia dichotoma*.
 C. DIOSCORIDIS Rauw. = *Bluchea Dioscoridis*.
 C. DISCOLOR Bory. = *Senecio appendiculatus*.
 C. DIVARICATA Spr. = *Vernonia diffusa*.
 C. DIVERGENS Wall. = *Vernonia subsessilis*.
 C. ELEGANTULA Wall. = *Vernonia bellidifolia*.
 C. ELÆAGNIFOLIA Wall. = *Vernonia elæagnifolia*.
 C. ELONGATA Wall. = *Blumea elongata*.
 C. ERICOIDES Lam. = *Vernonia ericoides*.
 C. EXCISA Wall. = *Blumea excisa*.
 C. EXTENSA Wall. = *Vernonia extensa*.
 C. FASCICULATA Wall. n. 3017. = *Blumea flavida*.
 C. FASCICULATA Wall. n. 3094. = *Blumea fasciculata*.
 C. FOETIDA Willd. = *Erigeron foetidum*.
 C. FOETIDA Lam. = *Pluchea Marylandica*.
 C. FOLIOLOSA Wall. n. 3030. = *Pluchea Martabanica*.
 C. FOLIOLOSA Wall. n. 3281. = *Pulicaria foliolosa*.
 C. FRUTICOSA Linn. = *Vernonia fruticosa*.
 C. GEMINIFLORA Ten. = *Phagnalon rupestre*.
 C. GENISTELLOIDES Lam. = *Baccharis genistelloides*.
 C. GLOMERATA Wall. = *Blumea glomerata*.
 C. GLOMERATA C. Wall. = *Blumea leptoclada*.
 C. GLUTINOSA Lam. = *Psiadia glutinosa*.
 C. GNAPHALODES H. B. K. = *Lænnecia gnaphalodes*.
 C. GRACILIS Wall. = *Blumea gracilis*.
 C. GRANDIS Wall. = *Pluchea grandis*.
 C. GRAVEOLENS Steud. = *Inula graveolens*.
 C. GRAVEOLENS Wall. = *Erigeron hispidum*.

- C. GUINEENSIS Willd. = *Blumea Guineensis*.
 C. GUMMIFERA Roxb. = *Commidendron gummiferum*.
 C. HALIMIFOLIA Desf. = *Baccharis halimifolia*.
 C. HAMILTONI Wall. = *Blumea Hamiltoni*.
 C. HELIOTROPIFOLIA Lam. = *Monarrhenus rufescens*.
 C. HEYNEANA Wall. = *Blumea Heyneana*.
 C. HIERACIFOLIA Spreng. = *Blumea hieracifolia*.
 C. HIRSUTA Linn. f. = *Pluchea hirsuta*.
 C. HIRSUTA Comm. = *Cylindrocline Commersonii*.
 C. HIRSUTA Wall. = *Erigeron sublystratum*.
 C. HOLOSERICA Wall. = *Blumea holosericea*.
 C. HYMENOPHYLLA Wall. = *Blumea hymenophylla*.
 C. JAVANICA Blum. = *Blumea lacera* J.
 C. INCANA Heyn. = *Vernonia physalifolia*.
 C. INCANA Wall. = *Vernonia abbreviata* var. ?
 C. INDICA Blum. = *Pluchea indica*.
 C. INDICA Wall. = *Decaneurum Dendigulense*.
 C. INTERMEDIA Lag. = *Phagnalon rupestre* β.
 C. INULOIDES Ait. = *Jasonia dichotoma*.
 C. LACERA Burm. = *Blumea lacera*.
 C. LACINIATA Roxb. = *Blumea laciniata*.
 C. LACINIATA Ham. = *Blumea crepidifolia*.
 C. LACTUCÆFOLIA Wall. = *Blumea lactucæfolia*.
 C. LANCEOLATA Wall. n. 3059. = *Vernonia divergens*.
 C. LANCEOLATA Wall. n. 2991. = *Berthelotia lanceolata* β.
 C. LIMONIFOLIA Sibth. = *Inula candida* β.
 C. LINEARIFOLIA Lam. = *Psidium linearifolia*.
 C. LINIFOLIA Linn. = *Sericocarpus solidaginoides*.
 C. LINIFOLIA Wall. = *Vernonia leptophylla*.
 C. LOBATA Linn. = *Neurolæna lobata*.
 C. LOBATA Juss. = *Ditrichum macrophyllum*.
 C. LONGICAULIS Wall. = *Eupatorium longicaule*.
 C. LONGIFOLIA Heyn. = *Blumea longifolia*.
 C. LUCIDA Spreng. = *Vernonia splendens*.
 C. LYCOPODIOIDES Lam. = *Eriothrix lycopodioides*.
 C. MACROPHYLLA Wall. = *Blumea macrophylla*.
 C. MACROSTACHYA Wall. = *Blumea macrostachya*.
 C. MADAGASCARIENSIS Lam. = *Psidium Madagascariensis*.
 C. MAGELLANICA Lam. = *Baccharis Magellanica*.
 C. MANILLENSIS Less. = *Blumea Manillensis*.
 C. MARYLANDICA Michx. = *Pluchea Marylandica*.
 C. MARYLANDICA Pursh. = *Pluchea foetida*.
 C. MEMBRANACEA Wall. = *Blumea membranacea*.
 C. MOLLIIS Willd. = *Vernonia cinerea*.
 C. MULTIFLORA Wall. = *Vernonia multiflora*.
 C. MURALIS Ham. = *Blumea muralis*.

- C. MUSRA Ham. = *Blumea Musra*.
 C. NUTANS Hain. = *Blumea commutata*.
 C. OBLIQUA Willd. = *Blumea amplexens*.
 C. OBOVATA Wall. = *Blumea obovata*.
 C. ODORATA Linn. = *Pluchea odorata*.
 C. OLAEFOLIA Lam. = *Linostyris villosa*.
 C. OVATA Wall. = *Blumea conyzoides*.
 C. OXYDONTA Wall. = *Blumea oxydonta*.
 C. PANICULATA Wall. = *Blumea phyllostachya*.
 C. PARVIFOLIA Wall. = *Blumea parvifolia*.
 C. PATULA Ait. = *Cyanopsis pubescens*.
 C. PAUCIFLORA Willd. = *Vernonia pauciflora*.
 C. PINIFOLIA Lam. = *Webbia pinifolia*.
 C. PINIFOLIA Bory. = *Faujasia pinifolia*.
 C. PLATENSIS Spreng. = *Vernonia Platensis*.
 C. PLATYLEPIS Spreng. = *Decaenium asperum*.
 C. POPULIFOLIA Lam. = *Distephanus populifolius*.
 C. POMÆ Sieb. = *Cylindrocline Commersonii*.
 C. PORTORICENSIS Bert. = *Vernonia Berteriana*.
 C. PROCERA Wall. = *Blumea procera*.
 C. PROLIFERA Lam. ? Blum. = *Microglossum proliferum*.
 C. PUMILA Sibth. = *Phagnalon pumilum*.
 C. PUNCTULATA Wall. = *Cyanopsis pubescens*.
 C. PUNDUANA Wall. = *Vernonia Punduana*.
 C. PUNGENS Lam. = *Iphiaea spinosa*.
 C. PURPURASCENS Sw. = *Pluchea purpurascens*.
 C. PYCNOSTACHYA Michx. = *Pterocaulon pycnostachyum*.
 C. PYGMÆA Sieb. = *Phagnalon pusillum*.
 C. RACEMOSA Wall. = *Blumea racemosa*.
 C. RAMIFLORA Spreng. = *Vernonia umbellata*.
 C. REDOLENS Willd. = *Monenteles redolens*.
 C. RIGIDA Sw. = *Vernonia rigida*.
 C. RIPARIA Blum. = *Pluchea riparia*?
 C. RIPARIA H. B. K. = *Tessaria mucronata*.
 C. ROBUSTA Roxb. = *Commidendron robustum*.
 C. RUGOSA Vahl = *Pterocaulon spicatum*.
 C. RUGOSA Ait. = *Commidendron rugosum*.
 C. RUNCINATA α Wall. = *Blumea runcinata*.
 C. RUNCINATA β Wall. = *Blumea cinerascens*.
 C. RUPESTRIS L. = *Phagnalon rupestre*.
 C. RUTILANS Poir. = *Decaenium Senegalense*.
 C. SAGITTALIS Lam. = *Baccharis sagittalis*.
 C. SALICIFOLIUS Lam. = *Monarrhenus salicifolius*.
 C. SALIGNA Wall. *Vernonia saligna*.
 C. SALSOLOIDES Turcz. = *Inula ammophila* β.
 C. SALVIFOLIA Bory. = *Blumea salvifolia*.

- C. SAXATILIS Linn. = Phagnalon saxatile.
- C. SAXATILIS Ten. = Phagnalon rupestre.
- C. SAXATILIS Sieb. = Phagnalon rupestre β .
- C. SCANDENS Wall. = Vernonia scandens.
- C. SEMIVESTITA Wall. = Blumea semivestita.
- C. SENECIOIDES Ham. = Blumea senecioides.
- C. SENEGALENSIS Willd. = Blumea Senegalensis.
- C. SENEGALENSIS Sieb. = Blumea Guineensis β .
- C. SERICEA Ait. = Schizogyne sericea.
- C. SERICEA Wall. = Blumea spinellosa.
- C. SERRATA Bory. = Senecio salicifolius.
- C. SERRULATA Lam. = Baccharis serrulata.
- C. SESSIFOLIA Blum. = Pluchea sessifolia.
- C. SESSILIFOLIA Wall. = Pluchea Wallichiana.
- C. SESILIFOLIA Sallab. = Distephanus populifolius.
- C. SICULA Willd. = Jasonia Sicula.
- C. SONCHIFOLIA Ham. = Blumea sonchifolia.
- C. SORDIDA Lin. = Phagnalon sordidum.
- C. SPICATA Lam. = Pterocaulon apicatum.
- C. SPINULOSA Spr. = Sommelfeltia spinulosa.
- C. SPLENDENS Spr. = Vernonia splendens.
- C. SQUAMOSA Bory. = Senecio squamosus.
- C. SQUARROSA Lin. = Inula conyzæ.
- C. SQUARROSA Wall. = Blumea myriocephala.
- C. STELLATA Spreng. = Vernonia oppositifolia.
- C. STIPULATA Vahl = Lihum Jussiei.
- C. SUBCAPITATA Wall. = Blumea subcapitata.
- C. SUBSIMPLEX Wall. = Blumea subsimplex.
- C. SUBSESSILIS Wall. = Vernonia subsessilis.
- C. SYLVATICA Blum. = Pluchea sylvatica.
- C. TENAXA Wall. = Blumea oxydonta β .
- C. TENORIT Spreng. = Phagnalon rupestre.
- C. THYOIDES Lam. = Baccharis thyoides.
- C. TOMENTOSA Forsk. = Phagnalon rupestre.
- C. TOMENTOSA Wall. = Pluchea tomentosa.
- C. TRICUNEATA Willd. = Baccharis cuneata.
- C. TRINERVIS Lam. = Baccharis trinervis.
- C. TRISULCA Ham. = Blumea trisulca.
- C. TUBEROSA Lam. = Jasonia tuberosa.
- C. ULIGINOSA Pers. = Pluchea bifrons.
- C. VAGANS Wall. = Vernonia vagans.
- C. VERBASCIFOLIA Willd. = Inula candida α .
- C. VILLOSA Willd. = Blumea aurita β .
- C. VILLOSA Wall. = Blumea arenaria.
- C. VIRENS A. B. D. Wall. = Blumea lapsanoides.
- C. VIRENS C. Wall. = Blumea virens.
- C. VIRENS E. Wall. = Blumea paucifolia.

- C. VIRGATA Linn. = *Pterocaulon virgatum*.
- C. VISCOSA Hortul. = *Inula viscosa*.
- C. VISCOSULA Wall. = *Blumea viscosula*.
- C. VOLKAMERIEFOLIA Wall. = *Vernonia volkameriæfolia*.
- C. VOLUBILIS Wall. = *Microglossum proliferum*.
- C. VULGARIS Lam. = *Inula conyza*.
- C. WIGHTIANA Wall. n. 3028. = *Monosis Wightiana*.
- C. WIGHTIANA Wall. n. 3093. = *Blumea Wightiana*.

CCXIII. PHAGNALON Cass. bull. philom. 1819. p. 174. dict. 19. p. 118, 23. p. 561. et 39. p. 400. Less. syn. 204. — *Conyza* et *Gnaphalii* sp. auct.

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, marginalibus pluriserialibus tenuissimis filiformibus fœmineis sæpè abortientibus, centralibus 5-dentatis hermaphroditis fertilibus. Recept. nudum planum. Involucrum imbricatum, squamis plùs minùs scariosis. Achænium teretiusculum erostre adpressè villosum. Pappus 1-serialis paucisetus conformis piloso-setaceus subacaber. — Fruticuli regiones Mediterraneas aut Canarienses habitantes, ramis tomentosis teretibus. Folia sparsa linearia aut lanceolata integra aut subdentata subtùs tomentosa sæpiùs margine revoluta. Capitula ramos sæpiùs apice aphyllis terminantia ebracteata sæpè solitaria. Flores lutei aut lutescentes.

1. P. SORDIDUM, caule basi fruticoso, ramis tomentosis ultimis aphyllis apice capitula 1-3 sessilia gerentibus, foliis linearibus integerrimis margine revolutis utrinquè tomentosis, invol. ovato-cylindracei squamis ovatis scariosis obtusiusculis. ζ ad rupes et muros Galliæ mediterraneæ, Italiæ, Mauritanis, Hispaniæ. *Conyza sordida* Linn. mant. 466. DC. fl. fr. 4. p. 140. *Gnaphalium sordidum* Linn. sp. 1193. *Gnaphalium conyzoides* Lam. fl. fr. 2. p. 63. *Phagnalon tricephalon* Cass. dict. 39. p. 401. — Barr. ic. t. 277 et 368. Invol. rufescens glaberrimum. Cor. flavidæ. (v. v. et s.)
2. P. UMBELLIFORME, caule basi fruticoso, ramis tomentosis apice ramulos plures 1-cephalos subumbellatos gracillimos interdum umbellato-ramulosos gerentibus, foliis integerrimis radicalibus lineari-oblongis planis caulinis linearibus margine revolutis subtomentosis, invol. oblongi squamis angustis acuminatis apice rufo-scariosis. ζ in rupestribus insulæ Palmæ arch. Canar. legerunt cl. Webb et Berthelot. Affine P. sordido sed evidenter distinctum! (v. s. in h. Webb et Berth.)
3. P. SAXATILE (Cass. bull. philom. 1819. p. 174), caule basi suffruticoso ramossimo, ramis tomentosis apice productis in pedunculos elongatos aphyllous 1-cephalos, foliis infer. oblongo-lanceolatis subdentatis super. linearibus margine revolutis suprà araneosis subtùs tomentosis, invol. campanulati glabri squamis sublinearibus acuminatis apice scariosis. ζ in rupibus et muris regionis Mediterraneæ à Cretâ ad insul. Canarienses, à Græciâ ad Lusitaniam. *Conyza saxatilis* Linn. sp. 1206. DC. fl. fr. 4. p. 140. *Phagn. subdentatum* Cass. dict. 39. p. 400. — Barr. ic. t. 426. Variat foliis caulinis planis aut margine revolutis, subdentatis aut integerrimis plùs minùs tomentosis. (v. v. et s.)
 β . *intermedium*, foliis subadnato-amplexicaulibus, invol. squamis angustioribus minùs acutis sed non obtusis. ζ in ins. Scio (Oliv.!), in muris ins. Crète (Sieb.!), in Catalonia (Pourr.!) et in Aragonia (Asso!). *Conyza saxatilis* Sieb. Asso! *Conyza intermedia* Lag. elench. p. 28. n. 358. *Phagnalon Lagasce* Cass. dict. 39. p. 401. An species propria? (v. s.)
4. P. RUPESTRE, caule basi suffruticoso procumbente, ramis tomentosis, pedunculis elongatis solitariis geminisve monocephalis aphyllis floccosis glabrisve, foliis basi adnato-amplexicaulibus oblongo-linearibus integris subdentatisve

sæpè undulatis suprà araneosis subtùs tomentosis, invol. subcampanulati glabri squamis rigidis obtusis planiusculis. ζ in rupibus calcareis Syriæ, Mauritaniae, Arabiae, Aegypti, Siciliae, Capreae, littoris Neapolitani, Sardiniae, insul. Olbiensium, Hispaniae ad Malagam et in ins. Canariensium. *Conyza tomentosa* Forsk. æg. p. 148. *Conyza rupestris* Linn. mant. 113. ex syn. *Conyza tomentosa* Shaw. sp. n. 176. ic. pess.? *Conyza saxatilis* Ten. fl. neap. 2. p. 213. Bové! pl. exs. n. 437 non Willd. *Conyza geminiflora* Ten. cat. h. Neap. 1819. p. 75. syll. p. 427. Link enum. h. Berol. 2. p. 323. *Conyza Tenorii* Spreng. ex Ten. *Phagnalon Tenorii* Presl. fl. sic p. xxix. *Phagnalon spathulatum* Cass. dict. 39. p. 401? Species à plurimis nunc cum *Ph. saxatili* nunc cum *Ph. calycino* mixta sed distinctissima foliis basi subdecurrente amplexicaulibus subhastatis, caule basi decumbente, invol. squamis nec denticulatis, nec undulatis nec membranaceo-scariosis. (v. s.)

5. *P. CALYCINUM*, caule basi suffruticoso, ramis tomentosis, pedunculis elongatis subtomentosis aphyllis 1-cephalis, foliis sessilibus oblongo-linearibus integris serratisve, super. margine revolutis, subtùs plùs minùs tomentosis, invol. campanulati glabri squamis obtusis subserratis membranaceo-scariosis subundulatis. ζ in rupibus insulae Maderae legit cl. Lowe et circa Mogador cl. Bronsonet. *Conyza calycina* Cav. annal. scienc. nat. v. 4. (1801) p. 87, ex Mogador orta. *Phagnalon Bennetti* Low. in litt. *Conyza rupestris* Buch in hist. Mad. (v. s.)
6. *P. MITIDUM* (Fresen. pl. eg. p. 81. t. 4. f. 2), caule vix basi suffrutescente erectiusculo ramoso tomentoso, foliis linearibus basi subauritis semiamplexicaulibus margine revolutis acutis suprà araneosis subtùs cano-tomentosis, invol. lati campanulati squamis scariosis glabris nervo medio fusco donatis, ext. ovatis obtusis intimis linearibus acutis. ζ in valle Arbain (Rupp. ex Fres.), in Monte Sinai (Bové!). Arabia dicitur *Honet-el-Naghi* *Conyza* n. 120 Bové! pl. exs. Flor. junio. Species ex invol. distinctissima. Pappus 4-5-setus albiissimus. Capitula majora quàm pler. sp. generis. (v. s. comm. à cl. Bové.)
7. *P. PUMILUM*, caule ad collum subtuberoso fruticoso cæterùm decumbente brevissimo, foliis obovatis in petiolum attenuatis integerrimis subtùs tomentosis, pedunculo vix foliis longiore aphylo 1-cephalo subtomentoso, invol. pauciserialis squamis acuminatis exterioribus dorso tomentosis intimis elongatis subscariosis. ζ in cæcunine montium Sphacioticorum Cretae. *Conyza pumila* Sibth. et Sm. prod. fl. Gr. p. 173. fl. Græc. (ined.) t. 863. *Conyza pygmaea* Sieb. pl. cret. exs. Suffrutex certè omnium minimus, vix pollicem longus. (v. s.)

CCXIV. CHIONOLÆNA.

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, radii tenuissimis 2-3-dentatis vel truncatis foemineis pluriserialibus, disci circ. 15 bisexualibus abortu sterilibus 5-dentatis. Invol. squamæ pluriserialis lineares siccae albæ glabræ demùm deciduæ. Recept. nudum punctatum persistens. Antherae ecaudatae. Stylus radii exsertus bifidus tenuis, disci indivisus filiformis minutè hispidulus. Achænium gracile teretiusculum villosum. Pappus 1-serialis setosus. — Suffrutex Brasiliensis arborem minutissimam referens, trunco nudo, ramis foliosis apice sericeo-canis mox glabris. Folia alterna conferta patula linearia acuta suprà glabra subtùs sericeo-cana enervia. Capitula ramos terminantia sessilia solitaria. Invol. ferè Helichrysi. Flores circ. 35 flavi? — Ex caract. accedit ad *Conyzam* et *Phagnalon* sed habitus valdè peculiaris.

1. *C. ARBUSCULA*. ζ in Brasiliæ provinciâ Minarum-Generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 298!). Frutex vix pedalis dendromorphus. Folia ferè acerosa, 5-6 lin. longa semilineam lata. (v. s.)

CCXV. ELACHOTHAMNOS.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus filiformibus fœmineis (fortè sterilibus), disci paucioribus tubulosis 5-dentatis hermaphroditis (fortè abortu masculis). Invol. imbricatum, squamis linearibus acutis 3-4-seriatis. Recept. nudum. Anth. Styl. Achænia teretiuscula. Pappus radii pluriserialis setiformis, setis inter se conformibus; disci biserialis, squamellis exter. paleaceis acutis, inter. paucis setiformibus. — Frutex Novo-Hollandicus glaber ramosus erectus. Folia sparsa teretiuscula acutissima carnosa erecta integerrima. Rami apice squamellati 1-cephali. — Ex capitulis junioribus characterem florum non ritè eruere quivi, sed pappo heterogeneo genus satis distinctum videtur. Nomen ex *ελαχος*, parvus, et *δαμνος*, frutex.

1. F. CUNNINGHAMI. ♂ in humidis depressis ad flumen Lachlan Nov.-Holland. julio flor. legit cl. All. Cunningham. Senecio othonnæoides Cunn. in litt. 1835. Frutex bipedalis Mesembryanthemi facie. Caulis albicans. Folia 7-10 lin. longa, acutiuscula lata. (v. s. comm. à cl. inv.)

Div. IV. BACCHARIDÆ. — Baccharidæ veræ *Cass. tabl. op. phyt.* 3. p. 68. — Baccharidæ *Less. syn.* 204 non p. 200.

Capitula (ut in Eutarchonantheis) dioica.

CCXVI. POLYPAPPUS *Less. in linnæa* 1819. p. 314. 1831. p. 149. *syn.* p. 204.

Capitula homogama dioica. Cor. masculæ et fœmineæ tubulosæ, ♂ 5-dentatæ, ♀ filiformes. Rachis plana ebracteolata. Invol. campanulatum imbricatum. Achænia teretiuscula. Pappus cum maribus tum fœmineis pluriserialis, setis plurimis apice clavellatis. — Fruticuli Americani ramosissimi. Ramuli pubescentes adulti glabri. Folia alterna sessilia oblonga subcuneata triplinervia. Capitula ramos terminantia solitaria in subdiv. majuscula. Cor. ex sicco purpurascens.

1. P. TRIPLINERVIS (*Less. syn.* 204), foliis apice serratis 3-5 notatis, achæniisque glabris. ♂ in Brasiliâ legit cl. Sellow. Vernonia triplinervia Spreng. *syst.* 3. p. 435. Folia pollicem longa, 3 lin. lata. Capitulum 6-7 lin. diam. Inv. squamæ lanceolatæ acutæ adpressæ minutè ciliolatæ. Pappus rufescens. (v. s. ♂ comm. à Mus. reg. Berol.)
2. P. PARVIFOLIUS, foliis lineari-oblongis integerrimis junioribus subtus achæniisque puberis. ♂ in Mexico ex h. Hænke. Frustula mascula imperfecta tantum video, sed habitus omnino prioris. Fol. sup. 4 lin. longa, lin. lata. Capitula 4-5 lin. longa. Cor. fœm. paulò crassiores et evidentius dentatæ. Indiv. ♂ ign. (v. s. ♀ in h. Hænke.)

CCXVII. BACCHARIS *Linn. gen. n.* 949. *excl. spec. Michx. fl. bor. Am.* 2. p. 125. *Brown obs. comp. in trans. Linn. soc.* 12. p. 115. — *Molina Ruiz et Pav. prod.* p. III. t. 24. *syst.* p. 200. — *Baccharis* et *Molina Less. syn.* p. 204 et 205. — *Sergilus* et *Baccharis Gærtn.* — *Baccharis*, *Sergilus*, *Pingrea*, *Tursonia* et *Arrhæuachne Cass.* — *Stephananthus Lehm.*

Capitula multiflora dioica, cor. homogamis tubulosis. Recept. nudum aut in paucis speciebus subpaleaceum. Invol. subhemisphæricum aut oblongum pluriserialè imbricatum. ♂ Cor. ad faucem dilatatæ 5-fidæ. Anthæræ exsertæ ecaudatæ. Stylus plus minus abortivus. ♀ Cor. filiformes subtruncatæ, stylus bifidus exsertus. Anth. omnino nullæ. *Achæmium sæpius sulcato-costatum*. Pappus pilosus ♂ uniserialis sæ-

pius tortuosus aut subplumoso-penicillatus involucri longitudinem subæquans, ♀ uni- aut pluriserialis apice gracilis invol. longitudinem sæpius superans. — Frutices rarius herbæ. Species genuinæ omnes in Americâ spontaneæ sæpius glabræ et resinoso-viscosæ interdum variè villosæ. Folia alterna in paucissimis opposita, sæpius ex utroque latere in nervulum aut interdum in alam decurrentia integra aut dentata. Capitula variè disposita. Cor. sæpius albæ. — Species inter se valdè affines nec ullo modo in genera plura divellendæ.

§. 1. *Trinervatæ, nempe foliis tri- aut triplinerviis non imbricatis, nec cuneatis, ramis apteris.*

1. *B. SPECIOSA*, suffruticosa glabra, ramis teretibus, foliis in petiolum attenuatis ellipticis acuminatis integerrimis trinerviis et reticulato-venosis, corymb terminali amplo fastigiato composito polycephalo, invol. utriusque sexus campanulati squamis ovatis submucronatis, achæniis adpressè pilosisculis ♂ in montanis Guadalupe jnnio flor. legit cl. Perrottet. Frutex 6-8-pedalis, ramis crassis medullosis semilignosis. Folia 3-4 poll. longa, 15-18 lin. lata. Corymbi rami angulati. Pappus rufus, ♀ uniserialis. (v. s. commun. à cl. Perrottet.)
2. *B. NERVOSA*, fruticosa glabra, ramis virgatis teretibus, foliis petiolatis ovali-lanceolatis acuminatissimis trinerviis integerrimis membranaceis, corymbis inf. axillaribus super. aphyllis in paniculam terminalem dispositis, invol. ♂ campanulati squamis lanceolatis acutis apice ciliatis ♂ in insul. Trinitatis (Sieb.), et Guadalupe (Krauss!). Eupatorium nervosum Sieb. ! fl. trin. exs. 76. non Swartz. Folia 3 poll. longa, vix pollicem lata. Flor. fœm. ign. (v. s.)
3. *B. TRINERVIS* (Pers. ench. 2. p. 423. n. 7), suffruticulosa glabra ramosissima, ramis teretibus, ramulis striato-sulcatis, foliis petiolatis ovato-lanceolatis acutis pnt acuminatis integerrimis 3-nerviis, caulinis alternis, rameis oppositis, corymbis terminalibus laxis, invol. subcampanulati utriusque sexus squamis ovatis obtusiusculis, achæniis adpressè puberis. ♂ in Brasiliâ (Commun. Lhotsky!), prope Rio-Janeiro (Lund!). Conyza trinervis Lam. dict. 2. p. 85. Pappus rufus. Folia 1-2 poll. longa, 6-12 lin. lata. (v. s.)
4. *B. QUINQUENNERVIS*, suffruticosa glabra, ramis teretibus, foliis alternis petiolatis ovatis acutis quinquenerviis à medio ad apicem crenato-serrulatis, capitulis ♂ ad apices ramulorum 5-7-congestis, invol. squamis lineari-oblongis obtusis disco æqualibus. ♂ in fruticetis prov. Sancti-Pauli Brasiliæ legit cl. Lund. A trinervi et ab omnibus mihi notis foliis quinque seu ferè quintuplinerviis diversa. (v. s. ♂ comm. à cl. Lund.)
5. *B. RHIXIOIDES* (H. B. et Kunth nov. gen. am. 4. p. 66), fruticosa volubilis, ramis sulcatis pubescentibus, foliis petiolatis elliptico-oblongis brevissimè acuminatis basi acutis integerrimis trinerviis subtus secus nervos et in petiolo pubescentibus, corymbis axillaribus et terminalibus in paniculam dispositis, invol. campanulati squamis oblongis in ♂ subobtusis in ♀ subacutis, achæniis striatis puberulis. ♂ in Andibus Peruvianorum (H. B. K.), ad Caracas (Vargas!), in Brasiliâ (h. Deless.), ad fluvior. margines circa Rio-Janeiro (Lund! Gand!) et ni fallor in prov. Matto-Grosso. Affinis *B. trinervi* et cineræ. Frutex ex Kunth volubilis, ex Lund 2-5-pedalis. An plures hic species confusæ? (v. s.)
6. *B. CAMFORUM*, collo fruticoso, caulibus erectis suffruticosis simplicissimis teretibus vix sub lente pubescentibus, foliis sessilibus glabris coriaceis alternis ellipticis trinerviis à medio ad apicem argutè serratis parcè reticulatis, capitulis ♀ in corymbum congestum terminalem congestis oblongis, invol. squamis oblongo-linearibus obtusis, interioribus submembranaceis disco æqualibus. ♂ in campis editis prov. Sancti-Pauli Brasiliæ. legit cl. Lund. *Chrysocoma decussata* fl. flum. 8. t. 9? quæ foliis alternis gaudet. Habitus hujus stirpis et plurimarum in campis Brasiliæ obviis præcipuè ex cl. Lund tribuendas more

vulgatissimo igne destruendi herbas camporum, unde rhizoma induratur et quotannis ampliatur (v. s. ♀ comm. à cl. Lund.)

7. *B. CINEREA*, suffruticulosa? totâ pube brevi velutino-cinereascens, caule tereti, ramis divergentibus striatis, foliis petiolatis ovato-lanceolatis acuminatis integerrimis trinerviis suprâ glabriusculis alternis rameis hinc inde suboppositis, corymbo terminali laxo, invol. ovati utriusque sexûs squamis lanceolatis acutis, acheniis adpressè puberis. ♂? in Brasiliâ circa Bahiam leger. cl. Salzmänn et Blanchet. Affinis *B. trinervi* sed canescenti-cinerea nec glabra, invol. squamæ acutæ nec obtusæ, etc. (v. s.)
8. *B. TRICHOCLADA*, fruticosa, ramis pediculisque teretibus hirsutissimis, foliis vix petiolatis ovato-ellipticis subacuminatis mucrone calloso apiculatis integerrimis grossè trinerviis suprâ glabris subtus sparsè villosis, pedunculis secus ramulos axillaribus capitula 3-5 subsessilia gerentibus, invol. ♂ squamis lanceolatis acuminatis subciliatis glabriusculis. ♂ in Mexico ex herb. Hænke. Affinis *B. cinereæ* et *rhexioidi* sed ramis saltem junioribus villis longis densis vestitiis facilè differt. Flor. ♀ ign., ♂ non satis notî. (v. s. in h. Hænke. à cl. de Sternberg miss.)
9. *B. CINNAMOMIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 65), fruticosa glabra, ramulis angulatis, foliis petiolatis oblongis breviter acuminatis basi angustatis integerrimis subcoriaceis trinerviis suprâ nitidis, paniculis terminalibus ramosissimis, capitulis ♀ pedicellatis, invol. hemisphærici squamis scariosis acutis. ♂ in convallibus Quitensium? Folia 4 poll. longa, 18 lin. lata, petiolo 7-8 lin. Flores in invol. crebri, masculi ign. Affinis *B. trinervi* sed folia basi angustata nec rotundata.
10. *B. VENUSTA* (H. B. et Kunth nov. gen. am. 4. p. 66), fruticosa glabra, ramulis angulatis, foliis petiolatis oblongis breviter acuminatis basi acutis integerrimis membranaceis triplinerviis, paniculis axillaribus terminalibusque ramosis, capitulis ♀ pedicellatis, invol. hemisphærici squamis scariosis acutiusculis, achenio sulcato glabro. ♂ in Novâ Andalusîa juxta Caripe et in sylvis Orinocensibus. Folia 3 poll. longa, pollicem lata, petiolo 3-4 lin. Flor. ♂ ign. Affinis *B. trinervi*.
11. *B. OXYPHYLLA*, suffruticosa? glabra resinoso-subviscosa, ramis virgatis sulcato-angulatis, foliis petiolatis elongato-lanceolatis basi acutis apice acuminatissimis integerrimis triplinerviis, paniculâ ramosâ laxâ terminali, capitulis ♀ pedicellatis, invol. squamis lanceolatis acutis intim. linearibus, achenio subpuberulo. ♂? in Peruvîa legit cl. Poeppig (pl. exs. n. 24). Petioli 4 linear. Limbus fol. 3-3 $\frac{1}{2}$ poll. longus, 7 lin. latus. Nervi laterales duo suboppositi ad quartam nervi mediî partem orti, cæterum nulli. Flor. ♂ ign. (v. s.)
12. *B. DIVERGENS*, fruticosa apice viscosa, ramis teretibus floridis divergentibus, foliis brevissimè petiolatis lanceolatis utrinquè acutis subtripplinerviis integerrimis glabris summis pedicellisque subpubescentibus, corymbis compositis confertis, capitulis subfasciculatis, invol. ♀ squamis lanceolatis acutiusculis apice summo subbarbatis. ♂ in Mexico ex h. Hænke. Folia pollicem longa, 4-5 lin. lata suprâ viscoso-nitida. Achenia parva glabriuscula. Pappus 1-ser. pallidè rufescens (v. s. ♀ in h. Hænke à cl. de Sternberg comm.)
13. *B. DOUGLASII*, herbacea erecta glaberrima viscosa, caule simplici supernè angulato, foliis breviter petiolatis lanceolatis acutis integerrimis trinerviis utrinquè punctatis, corymbo ♂ composito fastigiato polycephalo subnudo, invol. campanulati squamis lanceolatis margine membranaceis acutiusculis ciliatis. ♀? in Californiâ legit cl. Douglas. Folia 2 poll. longa 5-6 lin. lata, super. angustiora. Pappus ♂ invol. æqualis albidus. (v. s. ♂ comm. à soc. hort. Lond.)
14. *B. VIMINEA*, fruticosa glabra, ramis teretibus, ramulis striato-nervosis, foliis breviter petiolatis oblongo-lanceolatis utrinquè acutis basi tenuiter trinerviis, aliis integerrimis, aliis ad apicem minutè denticulatis, corymbis subracemosis confertis ramulos terminantibus, capitulis ♂ pedicellatis, invol.

squamis ovali-lanceolatis acutis membranaceis ciliatis. $\frac{5}{8}$ in California legit cl. Douglas. Folia 12-14 lin. longa, 2-2 $\frac{1}{2}$ lata, primâ fronte uninervia imò sese enervia, ramul. infer. breviora ferè squamas gemmaceas referentia. Pappus σ albidus crispus vix invol. longior. Fl. \varnothing ign. (v. s. comm. ab hom. soc. hort. Lond.)

15. *B. HÆNKEI*, herbacea? erecta ramosa glabra viscosa, ramis etiam floridis teretiusculis, foliis breviter petiolatis ovali seu oblongo-lanceolatis acutis integerrimis trinerviis utrinque valde punctatis, corymbo composito fastigiato polycephalo, invol. campanulati squamis lanceolatis margine membranaceis acutiusculis ciliatis, achæniis glabris minimis. $\frac{2}{3}$ $\frac{5}{8}$ in Mexico circa Real del monte legit cl. Hænke. Valdè affinis *B. Douglasii* sed caulis ramosior apice teretiusculus, folia latiora magis punctata pappus rufus (v. s. σ et \varnothing in herb. Hænke à cl. de Sternberg misso.)
16. *B. FUNKHUIZII*, caule herbaceo? erecto tereti simplici pubero, foliis oblongo-linearibus acutis sessilibus basi trinerviis integerrimis aut remotè subdentatis pube sparsâ subcabridis, paniculâ \varnothing elongatâ angustâ subracemosa, ramulis axillaribus distantibus oligocephalis, pedicellis 1-3-cephalis, invol. squamis oblongo-lanceolatis subacutis apice membranaceis, achæniis pube minimâ pubescentibus. $\frac{2}{3}$? in Brasiliæ prov. Sancti-Pauli. Caulis bipedalis. Pappus rufus invol. duplò longior pluriserialis. Achæn. et invol. ferè *B. cordifoliae* sed cæt. diversiss. Pl. σ ign. (v. s. \varnothing in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 515 miss.)
17. *B. LONGICEPS* (Kunze! in Poepp. coll. 2. n. 104), suffruticosa glabra resinoso-viscosa, foliis linearì-lanceolatis utrinque attenuatis mucronato-acutis integerrimis aut parè serratis punctatis trinerviis, nervis lateralibus margini adproximatis, corymbo composito exserto fastigiato polycephalo aphyllò, capitulis \varnothing campanulatis, invol. squamis lanceolatis acuminatis glabris margine scariosis, achæniis glabris nervoso-striatis. $\frac{5}{8}$ in lapidosis inundatis Rio de Chili legit cl. Poeppig. Proximè accedit ad *B. glutinosam* et præsertim ad *B. parvifloram* et si hybridæ facillè admittendæ essent, ut utriusque proles mixta ferè videtur (v. s. ex coll. Poepp.)
18. *B. SALICIFOLIA* (Pers. ench. 2. p. 425. n. 35), fruticosa, foliis lanceolato-linearibus triplinerviis serratis, serraturis remotis, corymbis terminalibus. $\frac{5}{8}$ in Peruvias montibus altis ad Pillao et Acomayo tractus. Molina salicifolia R. et Pav. syst. p. 210. An, suad. Spreng., eadem ac *B. guidifolia*? sed caract. non suffic. †
19. *B. INEQUINOXA* (H. B. et Kunth nov. gen. am. 4. p. 63), fruticosa ramosissima glabra, ramulis sulcatis viscosis, foliis petiolatis linearì-lanceolatis acutis basi angustatis dentatis trinerviis membranaceis, paniculis terminalibus, capitulis σ pedicellatis 10-floris, invol. hemisphærici squamis scariosis acutis. $\frac{5}{8}$ in calidis prov. Bracamorensis juxta Passo de Matara. Fol. 2 $\frac{1}{4}$ poll. longa, 4 lin. lata, petiolo 3 lin. Flor. \varnothing ign.
20. *B. GIBBIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 61), fruticosa glabra, ramulis sulcatis, foliis subsessilibus linearì-lanceolatis subacuminatis apicem versus argutè serrulatis trinerviis subcoriaceis subtus glanduloso-punctatis, corymbis σ terminalibus, pedicellis bracteis subulatis gerentibus, invol. hemisphærici squamis lanceolatis acutis scariosis. $\frac{5}{8}$ in scopulis montis Chimborazo. Folia Daphne Gnidii, 13-15 lin. longa, sesquilineam lata.
21. *B. LANCEOLATA* (H. B. et Kunth nov. gen. am. 4. p. 63), arborea glabra junior viscosa, ramulis sulcatis, foliis petiolatis lanceolatis acutis basi angustatis superne remotè serratis trinerviis utrinque glanduloso-punctatis, corymbis terminalibus ramosissimis dichotomis, capitulis σ pedicellatis 15-20-floris, invol. hemisphærici squamis scariosis glabris acutis. $\frac{5}{8}$ in littore Oceani Pacifici juxta Santa Peruvianorum. Folia bipollicaria, 5 lin. lata. Flor. \varnothing ign.
22. *B. RACEMOSA*, fruticosa, foliis ovato-lanceolatis trinerviatis acutè serratis, racemis terminalibus. $\frac{5}{8}$ in sylvis et nemoribus Chili ad Nonguen, Carcama,

Palomares, Huapen et Halqui tractus. Molina racemosa R. et Pav. syst. 209. Vulgò cum pluribus aliis *Chiloe* nuncupatur. Cœt. ign. †.

23. B. *COERULESCENS*, suffruticosa erecta glabra, ramis teretibus striatulis, foliis petiolatis elongato-lanceolatis utrinquè attenuatis trinerviis utrinquè punctulatis, infer. ferè à basi grossè serratis, super. minùs serratis imò subintegerrimis, corymbo paniculato laxo, capitulis longè pedicellatis, invol. campanulati squamis lanceolatis acuminatis glaberrimis margine albis scariosis vix subfimbriatis, achæniis glabris. ♀ in Mexico circa Matamoros legit cl. Berlandier (pl. exs. n. 2343). Cor. ex albo cœrulescentes. Capitula 2-3 lin. diam. pallidè rufa. Pappus foemin. 1-serialis. (v. s. comm. à cl. invent.)
24. B. *SERRULATA* (Pers. ench. 2. p. 423. n. 8?), herbacea erecta glabra, caule basi tereti apice angulato, foliis petiolatis ovato-lanceolatis acuminatis serratis trinerviis minutissimè punctulatis, corymbo composito terminali fastigiato, invol. ♂ campanulati squamis lanceolatis acutiusculis. — in Brasiliæ prov. Rio-Grande. *Conyza serrulata* Lam. dict. 2. p. 85? Specim. mihi obvium à desc. Lamarkii tantum recedit in eo quod sit glabrum nec subglabrum, masc. nec. foem. (v. s. in h. Mus. reg. Par. à Mus. Bras. sub n. 798 miss.)
25. B. *BAUVISSETA*, herbacea glabra erecta ramosa, caule ramisque teretiusculis, foliis subsessilibus elongato-lineari-lanceolatis utrinquè acuminatis acutè serratis triplinerviis, rameis sub 1-nerviis, corymbo composito terminali subpaniculato, invol. ♀ squamis oblongo-linearibus margine subscariosis, ext. obtusis, int. angustioribus acutis pappum æquantibus, achæniis parvis glabriusculis. — in Brasiliæ prov. Rio-Grande. Affinis B. ptarmicæfoliæ sed folia impunctata et flores in invol. numerosiores circ. 80. Pappus ♀ albidus 1-serialis vix invol. longior undè primà fronte capitula ♀ videntur ♂. Masc. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 781 miss.)
26. B. *PHLOGOPARPA*, fruticosa glabra, ramis teretibus apice angulatis, foliis lineari-lanceolatis utrinquè attenuatis basi tenuiter trinerviis margine integrè apice 1-nervosis serratis subtus punctatis suprà subpunctulatis, corymbo composito terminali, capitulis ♀ pedicellatis, invol. squamis lineari-lanceolatis subacutis, achæniis elongatis striatis, pappo intensè rufo colorato. ♀ in Americà olim Hispanicà sed loc. ign. Folia et rami obscuri in sicco subrostri. Fl. ♀ in invol. 20-25. Achænia ferè 2 lin. longa. Pappus rigidus multisetus. Flor. ♂ ign. (v. s.)
27. B. *ALAMANI*, suffruticosa glabra viscosa, ramis teretibus apice subangulatis, foliis brevissimè petiolatis angustè lineari-lanceolatis triplinerviis utrinquè attenuatis apice subserratis ad utramque faciem punctatis, corymbo paniculato conferto, capitulis pedicellatis ovatis, invol. ♂ squamis lanceolatis acuminatis densè ciliatis laxis subscariosis pappo longioribus. ♀ in Mexico legit cl. L. Alaman. Capitula ferè B. longifoliæ, folia ferè B. cœrulescentis sed angustiora et magis punctata. (v. s. comm. à cl. inv.)
28. B. *LONGIFOLIA*, herbacea erecta glabra ad apicem viscosa, caule tereti, ramis subangulatis, foliis elongatis lineari-lanceolatis breviter petiolatis utrinquè attenuatis à medio ad apicem serratis basi triplinerviis, corymbo composito subpaniculato ferè nudo, invol. campanulati squamis utriusque sexus lanceolatis acuminatis membranaceis densè ciliatis, dorso achæniisque glabris. ♀ in Mexico circa urbem legit cl. Berlandier flor. augusto (pl. exs. n. 624 et 653). Folia impunctata interdum supernè sublepidota, 3-5 poll. longa, semipollicem lata. (v. s. ♂ et ♀ à cl. inv. comm.)
29. B. *MARGINALIS*, suffruticosa resinoso-subviscosa glabra, foliis lineari-lanceolatis utrinquè acutis integerrimis aut parcè serratis trinerviis, nervis laterali-bus margini adproximatis, corymbis compositis foliosis polycephalis, invol. campanulati squamis lanceolatis ciliato-erosis ♂ latioribus magis fimbriatis, ♀ angustioribus, achænio glabro. ♀ in versuris et ripis Chaucay et Cercado prov. Peruvix (R. et Pav.), ad Valparaíso Chilensium (Gaud.!) Molina parviflora R. et Pav. syst. 209? Baccharis parviflora Pers. ench. 2. p. 425. n. 47? non Polr. Folia 2 poll. longa, 1½-2 lin. lata. Capit. utriusque sexus 2 lin. diam.

An stirps Chilensis certè eadem ac Peruviana? (v. s. ♂ et ♀ comm. cl. Gaudichaud.)

30. *B. FRVILLEI*, frutescens pube minutissimâ subpulverulenta, ramis teretibus substriatis, foliis breviter petiolatis lanceolatis utrinquè attenuatis grossè serratis triplinerviis, capitulis ♂ in corymbos compositos terminales anaphyllos digestis 18-20-floris, invol. ovati squamis lanceolatis acutis stramineis. ♀ in Peruvizæ (Caud.), ripis (R. et Pav.), ad sinum Chorillo (Mac-Ræ!), et circa Limam (Abadia!). Conyza frutescens, etc. Fenill. per. 2. p. 750. t. 37. *B. striata* R. et Pav. syst. 207 excl. syn. ? *B. ivæfolia* Peruviana (nec Capensis) auct. Habitus omninò Conyze ivæfoliæ sed etiam genericè distincta. Flor. ♀ ign. (v. s.)
31. *B. BERRÆFOLIA*, suffruticosa, ramis striato-angulatis villosò-hirtis, foliis breviter petiolatis ellipticis triplinerviis confertè et argutissimè serratis secus nervos utrinquè hirtellis, corymbo subpaniculato laxo polyccephalo, invol. fl. ♂ glabro, squamis linearibus margine angustè hyalinis apice subciliatis. ♀ in Mexico? Species ex foliis distinctissima. Folia 14 lin. longa, 7 lin. lata, petiolo 2 lin. hispido. Accedit ad *B. multifloram* sed foliis triplinerviis differt. (v. s.)
32. *B. RIPARIA* (H. B. et Kunth nov. gen. am. 4. p. 65), arborea glabra, ramulis striatis, foliis petiolatis oblongo-lanceolatis subacuminatis basi in petiolum angustatis dentato-serratis triplinerviis suprâ nitidis, paniculis terminalibus ramosis corymbosis divaricatis, capitulis ♂ pedicellatis circ. 25-floris, invol. hemisphærici squamis scariosis acutis. ♀ in convallibus Quitensibus prope Guanote et Rio-Chambo (H. et B.) fortè circa Limam? (Née herb.). An var. *B. floribunda* ut vult Spreng. sed petioli 3-4 nec 6-7 lin. longi? An ♂ *B. polyantha* sed flores in invol. 25 nec 120! Fl. ♀ ign. (v. s.)
33. *B. POLYANTHA* (H. B. et Kunth nov. gen. am. 4. p. 64), fruticosa glabra, ramulis angulatis, foliis petiolatis lanceolatis acutis basi angustatis crenato-serratis triplinerviis suprâ nitidis, paniculis terminalibus ramosissimis corymbosis diffusis, capitulis ♀ breviter pedicellatis 120-130-floris, invol. hemisphærici squamis scariosis acutis. ♀ in Peruvizæ? (H. B.), in Brasiliâ prope Monte-Video? (Spreng). Folia 4 poll. longa, 10 lin. lata, petiolo 3-4 lin. Flor. ♂ ign.
34. *B. FLORIBUNDA* (H. B. et Kunth nov. gen. am. 4. p. 64. t. 325), arborea glabra, ramulis sulcatis, foliis petiolatis subovato-oblongis subacuminatis basi in petiolum angustatis dentato-serratis triplinerviis suprâ nitidis, paniculis terminalibus ramosissimis corymbosis, capitulis ♂ pedicellatis 40-floris, invol. hemisphærici squamis scariosis acutis. ♀ in Andibus Novo-Granatensium prope Ancalguar et ad flum. Yacanatacu. Affinis *B. polyanthæ* sed folia 2½ poll. longa et petioli 6-7 lin. Arbor 18-24 ped. Flor. ♀ ign. (v. s. in h. Hænk. à cl. de Sternberg miss.)
35. *R. CONYZOIDES*, herbacea glaberrima, ramis striatis, foliis breviter petiolatis approximatis rotundo-ellipticis triplinerviis basi integerrimis apice crenato-serratis, paniculis terminalibus subcymosis, invol. ♀ latè campanulati squamis exter. subrotundis obtusis, inter. linearibus apice subciliatis, achæniis cylindraceis costatis hispidiusculis pappo subæqualibus. ♀ — ad fretum Sanctæ-Catharinæ Brasiliæ. Molina conyzoides Less. in linnæa 1831. p. 139. Fl. ♂ ign.
36. *B. ANOMALA*, suffruticulosa ramosa, canle tereti, ramis pubescentibus, foliis petiolatis ovatis basi obtusis apice acutè dentato-serratis trinerviis suprâ sparèe puberis subtus dense villosis, paniculis ramos terminantibus laxis, invol. ♂ squamis oblongo-linearibus acutis, fl. masc. stylum exsertum gerentibus. — in Brasiliæ prov. Rio-Grande. Fortè scandens et primâ fronte Clematidem fere referens. Folia alterna, rarissimè opposita, cum petiolo sesquilineari 7-9 lin. longa, 4 lin. lata. Stigmata maris perfecta sed ovarium abortivum. Pappus rufus scaber. Ind. ♀ ign. (v. s. ♂ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 856 miss.)
37. *B. PRUNIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 63), fruticosa ramosissima glabra viscosa, ramulis subangulatis, foliis petiolatis oblongis acutis basi an-

gustatis serratis subcoriaceis triplinerviis anprà nitidis, corymbis terminalibus et axillaribus, capitulis fœm. pedicellatis circ. 250-floris, invol. hemisphærici squamis scariosis acutis. ♂ in Peruvîa? Folia 15-16 lin. longa, 6 lin. lata. Flor. masculi ign.

38. *B. SEMISERRATA*, fruticosa, ramis teretibus glabris, ramulis subangulatis apice hirsuto-canescens, foliis lineari-lanceolatis utrinquè attenuatis trinerviis à medio ad apicem serratis, junioribus utrinquè pube sericeâ villosis demùm omninò glabris, pedunculis axillaribus folio brevioribus 1-5-cephalis, invol. ♀ squamis lanceolatis acuminatis, achæniis striatis glabris. ♂ in Brasiliæ prov. Rio-de-Janeiro ad Serra dos Orgaos legit cl. Vauthier (pl. exs. n. 338). Folia 1½ poll. longa, 2 lin. lata, pubem sericeam mox deponentia. Pappus vix rufescens invol. duplò longior. (v. s.)

ℓ. ? *glabrata*, ramulis hirsutiusculis, foliis etiam junioribus subglabris. ♂ in Brasiliæ prov. Sancti-Pauli. Nervi fol. laterales validiores paulò altius orti. An spec. propria? (v. s. in h. Mus. reg. Par. à Mus. imp. Brás. miss.)

39. *B. MONTANA*, arborea, ramis teretibus glabris junioribus araneoso-puberulis, foliis oblongo-lanceolatis utrinquè acuminatis margine præsertim ad basin revolutis dentes acutos 2-3 vix apice gerentibus triplinerviis subtus junioribus subcanescentibus demùm subglabris, paniculâ masculâ subracemosâ parvâ terminali foliis breviorè, inv. squamis ovato-lanceolatis margine membranaceis. ♂ in montosis prov. Rio-de-Janeiro vulgaris. Flores albidos octob. præfert ex Lund. mss. Affinis *B. semiserrata* sed folia subintegerrima verè tripli- nec trinervia. Ind. fœm. mihi ign. (v. s.)

40. *B. OXYDONTA*, herbacea? glabra erecta, caule ramisque teretibus, foliis petiolatis lanceolatis basi acutis integris apice acuminatis acutè serratis triplinerviis reticulato-venosis, racemis compositis axillaribus in paniculam foliosam dispositis, invol. ♂ campanulati squamis lanceolatis acutiusculis. — in Brasiliæ prov. Minas-Geraes ad Villa-Ricca legit cl. Vauthier. Folia ferè 1½ 3 poll. longa, 9-11 lin. lata. Capit. ♂ 3 lin. diam., ♀ ign. An fortè mas *B. triplinervia* sed folia angustiora subulata reticulata, capitula majora. (v. s. ex coll. Vauth. n. 260.)

41. *B. TRIPLINERVIA*, herbacea glaberrima, caule tereti, foliis petiolatis ovato-lanceolatis basi acutis apice longè acuminatis argutè serratis triplinerviis membranaceis, paniculis axillaribus terminalibusque laxis, capitulis ♀ pedicellatis, invol. squamis linearibus subacutis margine membranaceis pappum subsequantibus. — in Brasiliæ prov. Minarum generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 271). Simillima *Conyzæ triplinervia* sed dioica, folia subtus impunctata argutius serrata, achænia teretiuscula, caulis vix striatus nec sulcato-angulatus. Ind. ♂ ign. (v. s.)

42. *B. PUNCTIGERA*, herbacea pubero-pilosa subviscosa, foliis petiolatis ovato-lanceolatis basi acutis apice longè acuminatis argutè serratis triplinerviis subtus glanduloso-punctatis, corymbo composito fastigiato ramoso polycephalo, capitulis ♀ pedicellatis, invol. squamis linearibus acuminatis glabris. — in Brasiliæ prov. Minarum generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 261). Habitus *Conyzæ triplinervia* et *Baccharis triplinervia*, ab utraque primâ fronte differt pube pilosâ caulis ramorum petiolorum imò foliorum, et notis indic. Ind. ♂ ign. (v. s.)

43. *B. LUNDII*, fruticosa glabra, ramis teretibus ramulis striatis, foliis longè petiolatis lanceolatis basi cuneato-attenuatis integerrimis cæterùm serratis acuminatis triplinerviis membranaceis, corymbo composito terminali fastigiato polycephalo, invol. ♂ et ♀ squamis lanceolatis acutis in ♀ paulò majoribus, achæniis glabris, pappo 1-seriali. ♂ in Brasiliâ circa Rio-de-Janeiro legit cl. Lund. An *B. polyantha* à Spreng. in Brasiliâ citata sed non Kunthiana? *Chrysocoma sancta fl. flum.* 8. t. 3? Petioli 9-10 lin. longi. Limbus folior. 4-5 poll. longus, 1½ latus. Capitula parva 15-20-flora. Affinis *B. oxydonta* sed capit. ♂ dimidiò minora. (v. s.)

44. *B. GLUTINOSA* (Pers. ench. 2. p. 425), suffruticosa glaberrima viscosa, foliis

lanceolatis coriaceis grossè serratis trinerviis et penninerviis basi attenuatis apice acutis, corymbo breviter pedunculato, capitulis ♂ campanulatis, invol. squamis ovali-lanceolatis margine eroso-fimbriatis. ♀ in Chili prope Talca-guano (Cham.) Valparaiso (Bridg.), in aridis et in ruderalis ad Concept. et Puchacay prov. (R. et P.). Molina viscosa R. et Pav. syst. 207. Baccharis glutinosa Hook. et Arn. bot. Beech. 1. p. 31. Spreng. syst. 3. p. 463 excl. syn. Less. in Linnaea 1831. p. 148 et 506. Chilca Fenill. per. 2. t. 37 excl. Tourn. syn. Habitus Conyzae ivæfoliæ. Nervi fol. laterales aequaliter à nervo medio et margine distantes. Fl. foem. ign. undè genus dubium. (v. s. ♂ ex itin. Née et comm. à cl. de Chamisso.)

45. *B. MEDULLOSA*, suffruticosa glabra, ramulis striatis, foliis elongato-lineari-lanceolatis sessilibus utrinquè attenuatis serratis basi tenuè trinerviis, corymbo masculo composito fastigiato polycephalo, invol. squamis lanceolato-linearibus acuminatis. ♀ in Brasiliæ prov. Rio-Grande. Pl. foem. ign. Caulis digiti crassitie medullâ amplâ donatus. Fol. 5 poll. longa, 5-6 lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 850 miss.)

46. *B. PUNCTULATA*, suffruticosa glabra, ramis teretibus apice angulatis, foliis breviter petiolatis oblongis basi cuneatis integris cæterum regulariter serratis acutis trinerviis subtiè minutè et opacè punctatis, racemis compositis parvis ramulos axillares terminantibus et paniculam foliosam constituentibus, invol. ♀ ovatis squamis lineari-oblongis obtusis margine submembranaceis ciliolatis, achæniis glabris. ♀ in Brasiliæ prov. Rio-Grande. Fol. 2 poll. longa, 5 lin. lata. Pappus vix subrufescens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 970 miss.)

47. *B. SORDESCENS*, suffruticosa, ramis sulcato-angulatis puberulis, foliis sessilibus oblongis basi attenuatis apice vix acutis trinerviis velutinis margine tenuissimè serrato-ciliatis, ramulis in paniculam dispositis capitula 3-7 subsessilibus gerentibus, invol. ♂ squamis pluriserialibus oblongis subobtusis ad margines densè et longè molliter ciliatis. ♀ in Mexico legit cl. Hænke. Species *B. hirtæ* subaffinis sed ab ea et ab omnibus ferè involucro distincta. Flor. ♀ ign. ♂ parum noti. (v. s. in h. Hænke. à cl. de Sternberg nuss.)

48. *B. HIRTA*, herbacea erecta subsimplex tota strigoso-hispida, canle tereti striato, foliis sessilibus cuneatis apice grossè 3-7-dentatis basi trinerviis nervoso-reticulatis, corymbo composito terminali subpaniculato, capitulis 1-3 ad apices pedicellorum, invol. ♀ squamis lanceolatis vix acutis margine membranaceis subciliatis dorso glabriusculis, achænio jun. subpuberulo adulto glabro. — in Brasiliæ prov. Rio-Grande. Herba bipedalis. Folia 2 poll. longa, 5-8 lin. lata. Dentes seu lobuli fol. inf. obtusi, sup. acuti, omnium mucronati. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 833 et 967 miss.)

49. *B. MYRICEFOLIA*, fruticosa, ramis teretibus glabris, ramulis teretibusculis pube molli cano-hirsutulis, foliis oblongis basi longè attenuatis subpetiolatis apice dentibus utrinquè 1-2 incisus aut integerrimis trinerviis, nervis lateralibus tenuissimis, junioribus floccoso-pubescentibus adultis glabris, pedunculis axillaribus 1-2-cephalis folio brevioribus, invol. ♀ squamis lanceolatis acutis, achænio striato glabro pappi longitudini ferè aequali. ♀ in Brasiliæ prov. Sancti-Pauli. Affinis *B. semiserrata* sed faciliè distincta. Capitula dimidiò minor longius pediculata. Pappus brevis intensè rufus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 484 miss.)

50. *B. SCANDENS* (Pers. ench. 2. p. 424. n. 33), fruticosa scandens, foliis ovato-lanceolatis dentatis bidentatisque triplinerviis, corymbis terminalibus. ♀ in Peruviz versuris et aridis per Chancay et Cercado provincias ubi vern. *Chilca*. Molina scandens R. et Pav. syst. p. 205. Triorgyalis. Cæt. ign.

§. 2. *Cuneifoliæ, nempe foliis obovatis cuneatisve uninerviis aut triplinerviis non imbricatis, ramis apteris.*

51. *B. MAGELLANICA* (Pers. ench. 2. p. 425. n. 23), fruticulosa demisso-cæspitosa glabra viscosa, ramulis angulatis, foliis sessilibus confertis coriaceis obovato-

cuneatis obtusis aliis integerrimis aliis apice obtusè tridentatis, capitulis solitariis ad apices ramulorum sessilibus, invol. ovatis squamis margine ciliato-fimbriatis, ♂ ovali-lanceolatis, ♀ angustius lanceolatis acuminatis, achæniis striatis glabris. ♂ ad fretum Magellanicum (Lam.), ad portum Egmont et Desseo (Née!), in insulis Maclovianis (Gaud. ! D'Urv. !). Conyza Magellanica Lam. dict. 2. p. 91. Baccharis tridentata Gaud. ! fl. mal. p. 15. D'Urv. ! fl. mal. non Vahl Differt à spec. Vahl ramis glabris nec subvillosis, foliis 2-3 lin. longis nec pollicaribus, etc. (v. s. ♂ et ♀.)

52. *B. CUNEIFOLIA*, fruticulosa demissa glabra viscosa ramosissima, ramulis subangulatis, foliis sessilibus confertis coriaceis obovato-cuneatis obtusis ad apicem obtusè repando-subdentatis, capitulis ad apices ramulorum sessilibus confertis. ♂ ad fretum Magellanicum (Lam.), ad portum Egmont (Née!) et in Brasilii (Vahl), ad Monte-Video (Lam.). Conyza cuneifolia Lam. dict. 2. (1786) p. 91. Baccharis sessiliflora Vahl symb. 3. (1794) p. 97. An stirps Monte-Vidensis eadem ac Magellanica? An mera *B. Magellanica* varietas? (v. s. ♂.)

53. *B. ALPINA* (H. B. et Kunth nov. gen. am. 4. p. 48), fruticosa procumbens glabra, foliis obovato-spathulatis obtusis integerrimis coriaceis aveniis, capitulis ♂ subterminalibus solitariis sessilibus, inv. hemisphærici squamis obtusis, intimis oblongo-linearibus, recept. convexo nudo. ♂ ad radicis montis Pichinchæ et Antisanæ Quitensium. Folia 3 lin. et paulò longiora. Pappus ♂ apice incrassatus. Flor. in invol. circ. 70. An à *B. buxifolia* jam innuente Spreng. satis differt? sed cl. Kunth qui herb. Juss. sub oculis habuit ab eâ sejunctit.

54. *B. ? HUMIFUSA* (H. B. et Kunth nov. gen. am. 4. p. 48. t. 322), fruticosa prostrata repens glabra, foliis obovato-spathulatis obtusis integerrimis coriaceis aveniis, capitulis subterminalibus solitariis subsessilibus, invol. ovatis squamis lanceolatis scariosis acutis, recept. conico paleis subulatis obsito, achænio striato. ♂ in radicibus montis Pichinchæ et Antisanæ Quitensium. Furseniæ sp. Cass. dict. 37. p. 480. Affinis *B. Magellanica*. Pappus ♂ apice clavellatus. Flores in inv. circ. 35. — *B. cæspitosa* Pers. ench. 2. p. 428. n. 38, quæ Molina cæspitosa R. et Pav. syst. p. 203 ex Peruviae montanis orta sed vix descripta, ab hac fortè non differt? (v. s.)

55. *B. PROSTRATA* (Pers. ench. 2. p. 424. n. 37), fruticosa prostrata, foliis obovatis denticulatis integrisque, pedunculis axillaribus 1-cephalis. ♂ in Peruviae Andibus alpiis ad Bombom, Huamalies, Canta et Pillao. Molina prostrata R. et Pav. syst. 204. Vernaculè *Pachatacya*. Decoctum in dysurià prodest. Cæt. ign.

56. *B. FRIGIDA* (H. B. et Kunth nov. gen. am. 4. p. 49), fruticosa ramosissima glabra, foliis sessilibus lanceolato-spathulatis acutis integerrimis coriaceis uninerviis suprâ viscoso-nitidis subtus punctatis, capitulis ♂ terminalibus solitariis sessilibus, invol. hemisphærici squamis membranaceis, exter. ovato-oblongis, int. linearibus. ♂ in frigidis Peruviae. Flor. in invol. ♂ 60-65 albi, ♀ ign.

57. *B. MICROPHYLLA*, fruticosa erecta? ramosissima glabra viscosa, ramis teretibus, foliis petiolatis ovatis aut obovato-cuneatis denticulatis, capitulis racemosis breviter pedicellatis 5-floris, invol. squamis lineari-oblongis obtusis, interioribus duplò longioribus, achænio glabiusculo. ♂ in Peruvia legit cl. Hænke. Folia petiolo incluso 4 lin. longa, 3 lin. circ. lata, dentibus parvis 7-9 denticulata. (v. s. in h. Hænk. à cl. de Sternberg comm.)

58. *B. POLYGALEFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 49), fruticosa glabra subviscosa, ramulis subangulatis, foliis breviter petiolatis lanceolato-spathulatis acutis integerrimis coriaceis uninerviis suprâ nitidis subtus subpunctatis, capitulis ♀ axillaribus sessilibus subspicatis, invol. hemisphærici squamis acutiusculis scariosis, achænio striato glabro. ♂ juxta Moran Mexicanorum et in Paramo et decl. Andium Popayauensium (H. et B.), in Monte-Video (Spreng.)? An plures hic confusæ?

59. *B. OBTUSIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 51), fruticosa glabra, ramulis angulatis subviscosis, foliis sessilibus obovatis apice rotundatis et subrepandis basi cuneatis coriaceis triplinerviis, capitulis ♀ terminalibus capitato-

congestis sessilibus, invol. squamis scariosis acutiusculis, acheniis costatis glabris. ♀ in frigidis prope Gonzanama Peruvianorum. Folia 8-10 lin. longa, 5-6 lata. Pappus fuscescens. Fl. masc. ign.

60. *B. ODORATA* (H. B. et Kunth nov. gen. am. 4. p. 52), fruticosa ramosissima glabra viscosa, ramulis subangulatis, foliis brevissimè petiolatis ellipticis obtusis basi subcuneatis integerrimis coriaceis aveniis viscido-nitidis, capitulis ♂ terminalibus conglomeratis sessilibus, invol. hemisphærici squamis acutiusculis. ♀ in montosis prope Pasto Novo-Granatensium (H. et B.) et fortè in Brasiliâ? (Spr.). Folia ferè Buxi semipoll. longa, odorem balsameum redolentia. Fl. fœm. ign.

61. *B. BARBERIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 57), fruticosa ramosissima glabra viscosa, ramis teretiusculis, foliis subsessilibus obovatis obtusis mucronatis basi cuneatis integerrimis coriaceis reticulato-venosis nitidis, corymbis ♀ terminalibus densis, invol. hemisphærici squamis acuminatis, acheniis costatis glabris. ♀ in altâ planitie Quitensi ad radices montis Rucu-Pichincha. Folia 15 lin. longa, 6 lin. lata. Pl. ♂ ign.

62. *B. SQUARROSA* (H. B. et Kunth nov. gen. am. 4. p. 67), fruticosa glabra, ramis (caulibusve?) sulcato-angulatis paniculato-virgatis apice 1-cephalis, foliis breviter petiolatis spatulato-lanceolatis acutis integerrimis subcarnosis subaveniis viscosis, inv. ♀ squamis lineari-oblongis squarrosè imbricatis apice subsimbriatis. ♀ in aridis montium Mexicanorum prope Guanaxuato, ad radices montis La Buffa. Folia 10-11 lin. longa, 2 lin. lata.

63. *B. FILULARIS*, fruticulosa ramosissima glabra viscosa, ramulis angulatis, foliis sessilibus obovato-cuneatis obtusis integris aut sæpius subdentatis 1-nerviis coriaceis, capitulis ad apices ramorum et ramulorum axillarum solitariis globosis foliis parvis bracteatis, invol. ♂ squamis oblongis subacutis margine membranaceis apice subsimbrillatis. ♀ in Californiâ legit cl. Douglas. Folia 4 lin. longa, 2-3 lin. lata, summa capitulo breviora. Capit. 2 lin. diam. Fl. ♀ ign. (v. s. ♂ ex h. soc. hort. Lond. comm. à cl. Bentham.)

64. *B. AXILLARIS*, fruticulosa glabra, ramis teretibus junioribus striatis, foliis sessilibus obovatis obtusis basi cuneatis subnerviis integerrimis aut hinc inde subdentatis patulis coriaceis, capitulis ad axillas foliorum solitariis pedicellatis et idè racemos foliatis constituentibus, invol. ♂ ovati squamis lanceolatis acutis, flor. circ. 6 in capit. ♀ in Brasiliæ prov. Sancti-Pauli. Folia vix 3 lin. longa et 1½ lata. Pedicelli folio paulò breviores. Ind. ♀ ign. (v. s. ♂ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 481 miss.)

β. *dentata*, foliis cuneatis omnibus apice 3-5-dentatis. ♀ in prov. Rio-Grande. Capit. ♂ var. α conformia. Capit. ♀ brevissimè pedicellata oblonga cylindracea var. γ omninò conformia. (v. s. cum var. α, ♂ sub n. 956, ♀ sub n. 953.)

γ. *acutiloba*, foliis dentibus profundis acutis 5-7-incisis. ♀ in Bras. prov. Sancti-Pauli. Similis var. β, sed foliis acutè incisis non apice tantum sed etiam ad margines dentato-lobatis. Capit. ♂ similia in n. 417 et 419 herb. Mus. imp. Bras. Capitula ♀ in n. 418 obvia sunt cylindracea duplò longiora brevius pedicellata, squamis lanceolato-linearibus elongatis acutis. Pappus rufescens pluriserialis. Achenium striatum, nitidum. An certè aliarum (quarum specim. ♂ tantum vidi) conspecifica? (v. s. cum α et β sub n. 417, 418, 419.)

65. *B. MICROPHYLLA* (H. B. et Kunth nov. gen. am. 4. p. 53), fruticosa ramosissima glabra viscidula, ramulis angulatis, foliis subsessilibus cuneatis apice tridentatis coriaceis uninerviis, capitulis ♀ axillaribus breviter pedunculatis, invol. squamis subsquarrosis acutiusculis. ♀ prope Santa-Fe de Bogota. Bacch. tridentatæ proxima. Pl. ♂ ign.

66. *B. PEDICELLATA*, fruticosa ramosissima glabra viscosa, ramis teretiusculis, foliis obovato-cuneatis sessilibus apicem versus paucidentatis coriaceis sub 3-nerviis, venis lateraliter tenuibus aut subnullis, pedicellis axillaribus subnullis

- striato-sulcatis 1-cephalis, invol. ♀ squamis ovali-lanceolatis acutis apice subciliatis. ♀ in Chili ex h. Haenke. Folia serè B. cuneifoliæ aut B. concavæ viscoso-nitida 8-9 lin. longa 4 lin. lata. Pedicelli bracteolâ 1-2 instructi 7-9 lin. longi, involucrique pallida. (v. s. in h. Haenk. à cl. de Sternberg miss.)
67. B. CUNEATA, fruticosa glabra, ramulis angulatis, foliis cuneiformibus apice obtusè 3-dentatis basi in petiolum brevissimum attenuatis, capitulis ad apices ramulorum congestis subsessilibus ovatis, invol. squamis obtusis margine scarioso albo subfimbriato, acheniis glabris. ♀ in Mexico. Erigeron tricuneatum. Lin. f. suppl. 368? et ideò forsàn Conyza tricuneata Willd. sp. 3. p. 1946 excl. Lam. syn. Baccharis tricuneata Pers. ench. 2. p. 424. n. 26. Foliis accedit ad B. glomerulifloram sed glomeruli terminales nec axillares. (v. s. comm. à cl. Alaman et Mairet.)
68. B. GLOMERULIFLORA (Pers. ench. 2. p. 423. n. 11), fruticosa glaberrima, ramulis angulatis, foliis lævibus cuneato-obovatis in petiolum brevem attenuatis apice obtusis grossè dentatis summis obovato-oblongis integerrimis, capitulis ad axillas foliorum sessilibus congestis, invol. ♂ campanulati squamis oblongis obtusis. ♀ in sylvis Amer. boreal. littoralis à Virginiâ ad Carolinam (Fras. Bosc.) Baccharis sessiliflora (Pursh) Michx. fl. 2. p. 135. Ell. sketch 2. p. 320 non Vahl. Pappus ♂ brevis, ♀ multò longior. (v. s. ♂.)
69. B. CONSANGUINEA, fruticosa ramosissima glabra junior subviscosa, ramulis decurrenti-angulatis, foliis sessilibus obovato-cuneatis obtusis dentibus acutis utrinquè 2-4 notatis rigidis, capitulis ad apices ramorum et ramulorum sessilibus rarò solitariis sæpius 2-4 aggregatis, invol. ♀ squamis oblongo-linearibus obtusis margine membranaceis subfimbriatis, intimis acutiusculis. ♀ in Californiâ legit cl. Douglas. Valdè accedit ad B. confertam, cuneifoliam, etc. Folia 6 lin. longa, 3 lin. lata in sicco suprâ sublepidota. Ind. ♂ ign. (v. s. ♀ comm. ex h. soc. hortie. Lond. à cl. Bentham.)
70. B. SUBDENTATA, fruticulosa glabra, ramulis subangulatis junioribus folisque subviscosis, foliis sessilibus obovatis ovalibusve coriaceis integerrimis antè hinc inde apice subdentatis imâ basi subtrinerviis, capitulis ♀ oblongis in spicas breves foliis intermixtas confertis, squamis oblongis acutis inter. longioribus pappum subæquantibus. ♀ in Brasilæ prov. Minas-geraes (Vauth. pl. exs. n. 339), et Sancti-Pauli (h. imp. Bras. n. 474). Folia 6-9 lin. longa, 2-4 lata. Ind. ♂ ign. (v. s. ♀.)
71. B. OBOVATA (DC. non Hook. et Arn.), fruticosa ramosissima glabra viscosa, ramulis subangulatis, foliis sessilibus obovatis obtusis basi cuneatis integerrimis apice dentibus 5-7 notatis uninerviis, capitulis ♂ ad apices ramorum 3-5 congestis sessilibus, invol. squamis oblongo-lanceolatis acutiusculis. ♀ in Peruvæ collibus aridis affatim ad Tarmæ oppidum, vernac. *Tacamacho* apell. Molina obovata R. et Pav. syst. p. 206 (v. s. ex h. Dombey comm. à Mus. reg. Par.)
72. B. SINUATA (H. B. et Kunth nov. gen. am. 4. p. 53), fruticosa ramosissima glabra viscosa, ramulis angulatis, foliis sessilibus subrhomboides basi cuneatis apicem versus sinuatis subcoriaceis, corymbis terminali subtrichotomo, inv. ♂ hemisphærici squamis obtusiusculis, recept. paleas subulatas gerente. ♀ in regione temperatâ Peruvæ inter Guancabamba et Sondorillo. Turseniæ sp. Cass. dict. 37. p. 478. Folia 1½ poll. longa, 15 lin. lata. Cor. ♂ albæ 35-45 in invol., fœm. ign.
73. B. PHILIPPENSIS (H. B. et Kunth nov. gen. am. 4. p. 55), fruticosa ramosissima glabra junior viscosa, ramulis angulatis, foliis brevissimè petiolatis obovato-cuneatis apice 3-5-dentatis coriaceis, capitulis terminalibus conglomeratis sessilibus, invol. ♂ squamis scariosis acutis. ♀ in Venezuelâ prope San-Felipe. Affinis dicitur B. tridentatæ. Planta ♀ ign. Cor. ♂ albæ 30 in invol.
74. B. HAMBATENSIS (H. B. et Kunth nov. gen. am. 4. p. 62), fruticosa glabra, ramulis sulcato-angulatis, foliis petiolatis lanceolato-oblongis acutis basi cuneatis superne dentatis subcoriaceis uninerviis reticulato-venosis, corymbis terminalibus subpaniculatis, invol. ♂ hemisphærici squamis scariosis obtu-

sinuatis. ζ in aridis temperatis juxtâ Hambato Quitensium. Cor. σ albæ circ. 3o in invol., ζ ign. Folia pollicaria, 4 lin. lata.

75. B. VAUTHIERI, fruticosa glabra subviscosa, ramulis decurrenti-angulatis, foliis oblongo-cuneatis basi attenuato-subpetiolatis apice obtusis subtridentatis nitidulis venulosis, capitulis σ ad axillas solitariis breviter pedicellatis folio dimidiò brevioribus ovato-globosis, invol. squamis exter. minimis, inter. oblongis subobtusis disci longitudine. ζ in Brasiliæ prov. Minarum general. ad Tejucco legit cl. Vauthier (pl. exs. n. 328). Accedit hinc ad B. tridentatam illinc ad pilularem. Fœm. ign. Folia pollicem longa. Capit. pedicello incluso 3-4 lin. longa. (v. s. comm. à cl. coll.)

76. B. TRIDENTATA (Vahl symb. 3. p. 98), fruticosa glabra, ramulis decurrenti-angulatis, foliis oblongis basi attenuatis apice tridentatis rarius integris uninerviis, capitulis ad axillas summas et ad apices ramorum congestis subsessilibus oblongis, invol. utriusque sexus oblongis pallidis subacutiusculis, achænio striato glabro. ζ in collibus aridis Brasiliæ (Vahl), circa Bahiam (Salzm.), in prov. Minarum general. prope Tejucco (Vauth. sine num.), et Rio-Grande (h. imp. Bras. sub n. 963). B. tridentata Pers. et Spreng. excl. syn. Descr. Vahlî non satis completa huic potius quàm cæteris convenit, sed specim. non vidi. (v. s.)

μ ? *pluridentata*, foliis ad apicem dentibus 5-7 acutis instructis. ζ circa Bahiam in dumetis legit cl. Blanchet. An fortè spec. propria ad B. oblongifolias referenda? (v. s.)

77. B. BREVIIFOLIA, fruticosa ramosissima glabra, ramulis subangulatis, foliis obovato-cuneatis obtusis apice 3-dentatis, summis integris, capitulis ad apices ramulorum spicato-congestis numerosissimis, invol. fl. ζ squamis oblongis pallidis subacutiusculis, achænio glabro striatulo. ζ in campis editis prov. Sancti-Pauli Brasiliæ legit cl. Lund. fl. nov. An var. B. tridentatæ, sed folia 3-4 nec 12 lin. longa, obovato-cuneata nec oblonga, capitula paulò minora multò magis conferta et numerosa (v. s. comm. à cl. Lund.)

78. B. RETICULARIA, fruticosa glabra subviscosa ramosa, ramulis striato-subangulatis, foliis obovatis basi cuneato-attenuatis subsessilibus apice obtusis et obtusè 3-5-dentatis utrinquè reticulo tenui nervulorum ornatis, capitulis ad apices ramulorum circ. 3 in axillis ultimis solitariis subsessilibus, invol. ζ 5-flori squamis pallidis oblongis vix acutis, achænio striato glabro. ζ in Brasiliæ prov. Minarum generalium ad Villa Rica legit cl. Vauthier (pl. exs. n. 329). Folia ferè Buxi sed obtusiora magis obovata et dentata. Species affinis B. pauciflorescens. (v. s. comm. à cl. coll.)

79. B. SALZMANNI, fruticosa ramosissima glabra viscosa, ramulis striato-angulatis, foliis obovato-cuneatis subsessilibus dentibus grossis 7-9 instructis suprâ nitidis nervoso-reticulatis penniveniis, capitulis ad apices ramulorum inter folia subcongesto-corymbosis subsessilibus, inv. σ squamis acutis pappo æqualibus, ζ oblongi squamis elongatis acuminatis pappo paulò brevioribus, achænio striato glabro. ζ in collibus aridis circa Bahiam legit Salzman. Pappus albidus. Folia 12-15 lin. longa, 6-7 lata. (v. s. comm. à cl. coll.)

80. C. PLATYPODA, fruticosa glabra resinoso-viscosa, ramis striato-angulatis, floridis ancipiti-compressis, foliis petiolatis obovato-cuneatis obtusis grosse dentatis coriaceis uninerviis venis hinc inde innumeris pennatis subreticulatis, capitulis σ ad apices ramorum florid. dense aggregato-corymbosis, invol. squamis ovato-oblongis obtusis. ζ in Brasiliâ (herb. L'Hér.) prov. Minarum General. ad Mariannam (Vauth! pl. exs. n. 283). Folia 3 poll. longa, 1-1 $\frac{1}{2}$ poll. lata. Capitulorum glomeruli cincti foliis floralibus nunc brevibus oblongis, nunc elongatis oblongo-spathulatis. (v. s.)

81. B. CONFERTA (H. B. et Kunth nov. gen. am. 4. p. 55), fruticosa ramosissima glabra apice subviscosa, ramulis decurrenti-angulatis, foliis sessilibus obovato-oblongis acutis basi cuneatis apicem versus utroque latere dente unico instructis subaveniis, capitulis ζ terminalibus confertis subsessilibus, invol. squamis lanceolatis, inter. acuminatis demùm squarrosis, achæniis striatis glabris. ζ in

Mexici temperatis prope Cuernavaca (H. et B.) et ad Real del Monte (Berl.) pl. exs. n. 397). Fl. masc. ign. Folia 5-7 lin. longa. Affinis videtur *B. tridentata* et *cuneata*. (v. s.)

82. *B. PTERONIODES* (H. B. et Kunth nov. gen. am. 4. p. 51), fruticosa ramosissima, ramulis breviter petiolatis oblongo-cuneatis obtusis mucronatis dentibus utrinque 2-3 acutis instructis 1-nerviis utrinque punctatis, capitulis ad apices ramulorum solitariis sessilibus, invol. ♀ squamis lanceolatis acuminatis. ♂ in Mexico inter Tampico et Real del Monte maio flor. legit. cl. Berlandier (pl. exs. n. 398). Pappus ♀ corollam superans invol. duplō longior pallidē rufescens. Folia 3-5 lin. longa. (v. s.)

83. *B. ALATERNOIDES* (H. B. et Kunth nov. gen. am. 4. p. 51), fruticosa ramosissima, ramulis angulatis pulverulento-viscosis, foliis petiolatis obovatis basi cuneatis obsolete et remote dentatis coriaceis subvenosis glabris subviscosis, capitulis ♂ terminalibus congestis pedunculatis, invol. hemisphærici squamis scariosis acutis subfimbriatis. ♂ in altā planitie Peruvianā ad Caxaniarica. Cor. in invol. ♂ 45-50, fem. ign. Folia 6-7 lin. longa, 4-4½ lata.

84. *B. PÖPPIGIANA*, fruticosa glabra viscosa ramosissima, foliis obovatis basi cuneatis subsessilibus apice obtusis repando-dentatis, capitulis pedicellatis ad apices ramorum paucis umbellatis, invol. ♂ campanulati squamis lanceolatis vix apice subciliatis, ♂ inter fragmenta rupium graniticarum inter la Guardia et los Ojos de Agua Chilensiam. *Baccharis alaternoides* Poepp. ! pl. chil. exs. 2. n. 102. non Kunth. Differt à spec. Kunthianā pedicellis folio longioribus nec petiolo aequalibus. Affinis *B. cuneifolia* sed magis demissa et viscosa, dentes obtusiores, etc. (v. s. ♂ ex coll. Poepp.)

85. *B. UMBELLIFORMIS*, fruticosa glabra, foliis obovatis obtusis cuneatis et sessilibus planis dentatis uninerviis epunctatis, capitulis pedicellatis ad apices ramorum umbellatis, invol. ♂ campanulati squamis lanceolatis margine scariosis dense ciliatis. ♂ in Chili? Poepp. pl. exs. 860. An *Baccharis obovata* Hook. et Arn. bot. Beech. 1. p. 30? sed non Molina obovata R. et Pav. — (v. s. ♂ ex coll. Poepp.)

ℓ. invol. ♀ squamis minùs ciliatis — Poepp. ! exs. n. 695. An fœmina var. α? An spec. alt.? (v. s. ex coll. Poepp.)

86. *B. RUPICOLA* (H. B. et Kunth nov. gen. am. 4. p. 52), subarbores glabra, ramulis subangulatis, foliis subsessilibus confertis ellipticis obtusiusculis basi subcuneatis apicem versus remote denticulatis reticulato-venosis subtus viscoso-punctulatis, capitulis ♂ terminalibus congesto-corymbosis, invol. hemisphærici squamis coriaceis obtusis. ♂ in monte altissimo Purace juxta urbem Popayan. Arbor 12-18 ped. Folia 6 lin. longa 3½ lata. Fl. ♀ ign.

87. *B. RESINOSA* (H. B. et Kunth nov. gen. am. 4. p. 52. t. 323), fruticosa ramosissima glabra, ramulis striato-angulatis resinoso-viscosis, foliis brevissimè petiolatis obovato-oblongis acutiusculis basi subcuneatis 3-5-dentatis coriaceis subaveniis viscoso-nitidis subtus punctatis, capitulis terminalibus conglomeratis, ♂ sessilibus, ♂ brevissimè pedunculatis, invol. ovati squamis acutiusculis, achænio sulcato glabro. ♂ in Americā meridionali. Folia 8-9 lin. longa, 4-4½ lata. (v. s. ?)

88. *B. ROTUNDIFOLIA* (Spreng. syst. 3. p. 465?), fruticosa, ramis teretibus, ramulis striato-angulatis subviscosis, foliis sessilibus obovato-subrotundis, apice dentibus 3-5 repandis trinerviis coriaceis subtus præcipuè albido-furfuraceis, capitulis ♂ ad apices ramorum congestis sessilibus, ♀ minùs congestis subspicatis, bracteis ♂ parvis obovatis, ♀ invol. subæqualibus tridentatis, invol. squamis ♂ ovato-lanceolatis acutis, ♀ magis elongatis vix acutis, achæniis striatis glabris. ♂ in Brasiliæ prov. Rio-Grande. Phrasis Sprengelii optimè indiv. masculo convenit. Folia 6-9 lin. diam. Pappus ♀ rufus longus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. ♂ sub. n. 964, ♀ sub. n. 951.)

89. *B. CONGESTA*, fruticosa ramosissima glabra viscosa, ramulis striato-angulatis, foliis sessilibus obovatis basi breviter cuneatis apice obtusis margine dentibus 2-3 obtusis utrinque notatis penninerviis rigidulis, capitulis ad apices ramorum

sessilibus congestis, invol. ♂ hemisphærici squamis ovali-oblongis obtusiusculis margine fimbriato-ciliatis. — ♀ ad Mexico et Real del Monte legit cl. Hænke. Accedit ad B. confertam et cuneifoliam. Fl. fem. ign. (v. s. in h. Hænke à cl. de Sternberg miss.)

90. *B. XALAPENSIS* (H. B. et Kunth nov. gen. am. 4. p. 56), fruticosa glabra, ramulis sulcato-angulatis, foliis subsessilibus obovatis apicem versus acutè 5-dentatis basi cuneatis coriaceis subvenosis, junioribus subviscosis, capitulis ♂ terminalibus capitato-congestis sessilibus, invol. subhemisphærici squamis ext. ovatis, int. oblongo-linearibus. ♀ in montibus Mexicanis prope Xalapa ubi *Escovilla* dicitur. Aff. *B. cuneifolia*. Ind. ♀ ign.

91. *B. HETEROPHYLLA* (H. B. et Kunth nov. gen. am. 4. p. 62), fruticosa ramosissima glabra viscosa, ramulis sulcato-angulatis, foliis petiolatis oblongo-lanceolatis apicem versus sinuato-dentatis basi cuneatis membranaceis, pedunculis axillaribus et terminalibus 2-4-cephalis, capitulis ♂ subsessilibus hemisphæricis; inv. squamis ext. ovatis, int. oblongis obtusiusculis. ♀ in convallibus Mexicanis prope Guanaxnato et fodinam Valencianæ (H. B.). An ad Monte-Video, ut vult Spreng. syst. 3. p. 464?

α. foliis obtusis apicem versus sinuato-dentatis.

β. foliis acutis apicem versus dente 1-2 instructis. An spec. dist.?

92. *B. HALIMIFOLIA*, fruticosa ramosissima, apice tenuissimè subpubescens, ramulis angulatis, foliis sessilibus obovato-cuneatis subacutis apice dentibus 5-7 subrepandis penninerviis nervis suprà elevatis reticulatis, capitulis ♂ ad apices ramorum et axillas fol. pedicellatis subcorymbosim dispositis, invol. squamis pallidis subnerviis intimis acuminatis margine hyalinis. ♀ in Brasiliâ circa Bahiam. Fem. ign. Folia accedunt ad *B. halimifoliam*. (v. s.)

93. *B. NERI*, fruticosa glabra viscosa, ramulis subangulatis, foliis sessilibus oblongo-subcuneatis obtusis uninerviis dentibus 3-7 notatis, capitulis ♂ ad apices ramorum 3-5 sessilibus congestis circ. 15-floris, invol. squamis ovali-oblongis obtusis, int. linearibus vix acutis. ♀ in Amer. merid. Cordillera del Portillo legit cl. Née. Folia $3\frac{1}{2}$ lin. longa, $1-1\frac{1}{2}$ lata, sæpè deflexa coriacea. Fl. ♀ ign. (v. s. ex itin. Née.)

94. *B. CONCAVA*, fruticosa ramosissima, ramulis angulatis pubero-subvelutinis, foliis latè obovato-cuneatis apice obtusè tri-rariùs quinque-dentatis sessilibus glabris crasso-coriaceis uninerviis summis circa capitula subconcavis, capitulis ad apices ramulorum 3-5 congestis subsessilibus, invol. campanulatis squamis margine scariosis oblongis in ♂ obtusiusculis, in ♀ inter. linearibus subacutis, achæniis compressis glabris. ♀ in fruticetis collium Chilensium ad Conceptionem (R. et Pav. Mac-Ræ), Valparaiso (Bert. Gaud.), Topocalma (C. Gay.). Vulgò *Gaultro* ex R. et Pav. *Guanchu* ex Bert. Molina *conca* R. et Pav. syst. 206. *Baccharis tridentata* Poepp. coll. chil. n. 211. *Baccharis resinosa* Hook. et Arn. bot. Beech. 1. p. 31. excl. in utr. syn. (v. s. ♂ et ♀.)

95. *B. INTERMEDIA*, fruticosa glabriuscula resinoso-subviscosa, foliis lineari-cuneatis basi attenuatis apice repando-dentatis margine subrevolutis, capitulis ad apices ramulorum congesto-corymbosis brevissimè pedicellatis, invol. ♀ squamis ovali-lanceolatis vix acutis margine scariosis, achænio glabro striato. ♀ in Chili circa Valparaiso (Gadd.), in fruticetis collium (Bert.). Media ferè inter *B. rosmarinifoliam* et *concam*. (v. s.)

96. *B. VAHLII*, fruticosa glabra, ramis teretibus, ramulis angulatis, foliis obovato-cuneiformibus retusis integerrimis penninerviis, capitulis plurimis sessilibus ad apices ramorum congestis ovatis, inv. ♂ squamis ovali-oblongis exter. obtusis inter. vix acutis. ♀ in insulis Caribæis nempe Montserrat (Vahl), Santo-Domingo (Poit.), Sancta-Cruce (Vest.). *Baccharis dioica* Vahl symb. 3. p. 93. t. 74. Spreng. syst. 3. p. 459 (excl. syn. Kunth). Lam. ill. t. 698. f. 3. Cl. Vahl dixit folia trinervia sed in meis specim. (verisim. authenticis) potius penninerviis video. Capitula juniora oblonga subviscosa. Nomen specifi. mutavi cum ex gener. caract. *Baccharides* omnes sint dioicæ. (v. s.)

97. *B. HALIMIFOLIA* (Linn. sp. 1204), arborea glabra sublepidoto-glancoæscens, ramulis angulatis, foliis obovatis superne grosse angulato-dentatis basi cuneatis in petiolum subattenuatis, summis lanceolatis integerrimis, capitulis ♂ subglobosis pedicellatis solitariis adgregatis, ♀ fasciculatis laxè subpaniculatis ovato-oblongis, invol. squamis, ♂ ovatis ♀ inter. elongatis. 5 in maritimis Amer. bor. à Marylandiâ ad Floridam (Pursh), in Carolinâ (Michx.), Georgiâ (h. Duby.), *Baccharis halimifolia* Duham. arbr. ed. 1. v. 1. t. 35. ed. 2. v. 1. t. 60. *B. cuneifolia* Mœnch. *Conyza halimifolia* Desf. — Pluk. t. 27. f. 2. Herm. parad. t. 225. (v. v. c. et s. sp.)

♀. *angustior*, foliis cuneato-lanceolatis apice 3-5-dentatis, summis lanceolatis integerrimis. 5 in insulâ Cubâ circa Havanam. (v. s. comm. à cl. Ramon de la Sagra.)

98. *B. RETUSA*, fruticosa glabra apice viscosa, ramis teretibus, ramulis subangulatis, foliis obovatis subcuneatis in petiolum attenuatis obtusissimis dentibus obtusis utrinque 7-8 subrepandis basi trinerviis, capitulis ♂ ad axillas foliorum super. oblongorum solitariis pedicellatis folio brevioribus racemos foliaceos constituentibus, invol. ovati squamis lanceolatis acutis. 3 in Brasiliæ prov. Sancti-Pauli. Rami flor. laterales breves. Folia caulina cum petiolo 4-5 lin. 2 poll. longa, 9-10 lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 479 miss.)

99. *B. ILLINITA*, fruticosa glabra viscosa, ramis apice striatis, foliis breviter petiolatis latè obovato-cuneatis obtusissimis dentibus 7-9 distantibus grossis notatis suprâ sæpè glutine viscoso illinitis et lucidis trinerviis, nervis laterali-bus ferè æquidistantibus inter nervum medium et margines, racemo subpani-culato terminali, bracteis lineari-filiformibus pedicello brevioribus, invol. ♀ squamis lanceolato-linearibus acutis, achæniis striatis glabris. 3 in Brasiliæ prov. Sancti-Pauli. Affinis *B. mesoneura* sed bracteis multò angustiores et breviores, etc. Achænia parva. Pappus rufus invol. paulò longior 1-serialis. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 502 miss.)

100. *B. MESONEURA*, fruticosa ramosa viscosa, ramulis angulatis, foliis obovato-cuneatis in petiolum attenuatis à medio ad apicem grossè dentatis trinerviis, nervis laterali-bus æquidistantibus à nervo longitudinali et margine, capitulis racemoso-subpaniculatis, foliis floralibus oblongis integris pedicello longioribus, invol. ♀ squamis lanceolatis subacutis, achænio striato ovato glabro. 3 in Brasiliæ prov. Sancti-Pauli. Folia 20 lin. longa, 10 lin. lata. Pappus rufus invol. paulò longior. Fl. masc. ign. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 501 miss.)

101. *B. CASSINOIDES*, fruticosa glabra, ramis teretibus striatis junioribus foliisque superne viscosis, foliis in petiolum attenuatis obovatis obtusis mucronatis in-tegerrimis coriaceis penninerviis subtus minutissimè punctulatis suprâ nitidis, corymbis terminalibus foliis intermixtis, invol. ♂ ovati squamis linearibus acutiusculis. 3 in Brasiliæ prov. Sancti-Pauli. Valdè similis *B. cassinefoliæ*. Ind. fœm. ign. (v. s. ♂ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 477 miss.)

102. *B. CASSINEFOLIA*, fruticosa glabra, ramis teretibus, ramulis striatis, foliis in petiolum attenuatis obovatis obtusis mucronatis integerrimis coriaceis pen-ninerviis subtus minutissimè punctulatis, corymbis terminalibus foliis inter-mixtis, pedicellis bracteolatis, invol. ♂ ovati, ♀ oblongi, squamis ovatis ob-longis, ext. subacutis inter. longioribus obtusis, achæniis glabris. 3 in Brasiliâ circa Rio-Janeiro (Lund!), Bahia (Blanch.). Pappus ♂ invol. cor. æqualis, ♀ duplò longior albus pluriserialis. Folia 1 ½-2 poll. longa, 6-10 lin. lata (v. s.)

103. *B. VISMIRIODES*, fruticosa, ramis teretiusculis junioribus pedunculisque rufo-hirsutulis, foliis in petiolum brevem attenuatis obovatis basi cuneatis integerrimis ciliatis subpenninerviis brevissimè mucronatis, paniculâ terminali foliolis parvis ovali-oblongis instructâ polycephalâ, invol. ♂ squamis ovalibus obtusis glabris margine submembranaceis pappum subæquantibus. 3 in Brasiliæ prov.

- Sancti-Pauli. Fl. form. ign. Capitula ♂ parva. Folia rufescentia 15-16 lin. longa, 6 lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 478 miss.)
104. *B. CALVESCENS*, fruticosa, ramis teretibus junioribus dense villosis adultis glabris, foliis in petiolum attenuatis elliptico-oblongis submucronatis integerrimis uninerviis utrinque junioribus sericeo-villosis demum supra glabris subtus canescenti-velutinis subtuberculatis, racemis ramulos terminantibus simplicibus, invol. squamis oblongis acutis subrufo-membranaceis, achæniis striatis glabris. ♂ in Brasiliæ prov. Minas-Geraes ad Mariannam legit cl. Vauthier (pl. exs. n. 332) et circa Bahiam cl. Blanchet (n. 1972 et 1973!). Pappus rufus 1-serialis deciduus Ind. ♂ ign. (v. s. ♂ et ♀.)
105. *B. PAUCIFLOSCULOSA*, fruticosa glabra viscosa ramosa, ramulis decurrenti-angulatis, foliis obovatis basi cuneatis sessilibus apice obtusis obtusè dentatis tenuiter triplinerviis, capitulis ad axillas super. sessilibus solitariis, summis confertis, in ♂ magis glomeratis, in ♀ magis spicatis, invol. 5-8 flori squatis oblongo-lanceolatis, ♂ acutis, in ♀ acuminatis, achæniis glabris striatis. ♂ in Brasiliæ prov. Sancti-Pauli et Miner. gener. Folia 4-6 lin. longa, 3 circ. lata, subtus vix ac ne vix subpunctata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 480 et 486 miss.)
- β. punctulata*, foliis subtus punctulatis. ♂ in prov. Rio-Grande. Indiv. masc. ignota. An fortè species propria? (v. s. cum var. *α* sub n. 959.)
106. *B. LEPTOCERUALA*, fruticosa glabra, ramulis decurrenti-angulatis, foliis oblongo-linearibus seu obovato-oblongis integerrimis seu paucidentatis, obtusis basi attenuatis uninerviis coriaceis, capitulis axillaribus sessilibus ad apices ramorum et ramulorum spicato-congestis oblongis, ♂ 8-floris, ♀ 4-floris, invol. squamis oblongo-lanceolatis acutis membranaceis, achæniis glabris, pappo etiam ♀ involucrium vix superante. ♂ in Brasiliæ prov. Rio-Grande (herb. imp. Bras. n. 950, 977 et 948) et Rio-Janeiro (Blanch.). Folia 10-20 lin. longa, 2-4 lata. (v. s.)
107. *B. COGNATA*, fruticosa glabra viscosa ramosa, ramulis decurrenti-angulatis, foliis obovatis basi cuneatis sessilibus apice obtusis obtusè dentatis tenuiter triplinerviis, capitulis ad axillas super. sessilibus congestis subterminalibus, invol. ovati squamis utriusque sexus ovato-lanceolatis subacutis, floribus in invol. 14-18, achæniis striatis glabris. ♂ in Brasiliæ prov. Rio-Grande. Nimis *B. pauciflosculosa* affinis, sed ex numero florum et squamarum ferè duplo videtur diversa. Folia in specim. cæt. simillimis subpunctata aut epunctata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 985, 946 et 975 miss.)
108. *B. PLATENSIS* (Spreng. syst. 3. p. 465), fruticosa glabra viscosa, ramis oppositis junioribus striato-angulatis, foliis oppositis oblongo-cuneatis apice dentatis coriaceis, infer. obovato-oblongis, mediis oblongis, summis linearibus subintegerrimis, capitulis sessilibus glomeratis secus ramos et ramulos interruptè spicatis, invol. ovati squamis ovalibus subacutis pappi longit. subæquantibus. ♂ in Brasiliâ merid. ad Montevideo (Spr.), in littore dicto Bandâ orientale (Bacle!), ad Rio-Grande (h. Mus. Bras.). Folia infer. 2 poll. longa, 6 lin. lata. Variat foliis latioribus trinerviis aut angustioribus uninerviis. An duæ species hic fortè confusæ? (v. s.)
109. *B. SUBOPPOSITA*, fruticosa glabra subviscosa, ramulis striato-angulatis foliisque suboppositis cuneatis apice obtusè 5-7 dentatis basi integerrimis subpetiolulatis trinerviis, nervis lateralibus margini parallelis, capitulis ad apices ramorum paniculato-congestis breviter pedicellatis, invol. ♀ ovati squamis ext. ovatis int. oblongis acutis pappo æqualibus. ♂ in Brasiliæ prov. Rio-Grande: Habitus nonnull. Eupatorii specierum ob folia sæpius opposita aut subopposita rariùs alterna, sed flores Baccharidis. Folia 10-12 lin. longa, 5-6 lata. Capitula parva. (v. s. ♀ in h. Mus. Par. à Mus. imp. Bras. sub n. 978 et verisim. 949 miss.)
110. *B. AFFINIS*, fruticosa ramosa glabra, ramulis striato-angulatis, foliis cuneatis obtusis basi breviter petiolulatis integris apice obtusè 5-7-dentatis basi triner-

viii nervis lateralibus margini parallelis medio cæt. penninervio, summis obovato-oblongis integris, capitulis racemosis subpaniculatis pedicellatis ovatis, squamis utriusque sexus ovato-oblongis subacutis, achæniis striatis glabris, pappo invol. superante. ♀ in Brasiliæ prov. Sancti-Pauli et Rio-Grande. Affinis B. suboppositæ sed folia de generis more alterna obtusiora, capitula duplò majora, etc. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. ♂ sub n. 487 et 398, ♀ sub n. 969 miss.)

111. B. CRENULATA (Spreng. syst. 3. p. 465), fruticosa, foliis oppositis alternisque longè petiolatis oblongis obtusiusculis basi attenuatis crenulatis coriaceis reticulato-venosis 3-nerviis glabris, capitulis interruptè spicatis subsessilibus, spicis paniculatis. ♀ in Brasiliâ. †.

112. B. HOOKERIANA, fruticosa, foliis fasciculatis cuneatis basi attenuatis sessilibus dentato-spinosis rigidis, capitulis solitariis ramulos versùs apicem canlis foliosos breves terminantibus et ità quasi spicato-racemosis. ♀ ad Coquimbo Chilensium. Baccharis mucronata Hook. et Arn. bot. Beech. p. 30, non Kunth. Cæt. ign.

§. 3. Discolores, nempe foliis subcuneatis obovatis oblongis linearibusve uninerviis rarius penninerviis suprâ virescentibus subtus densè cano-tomentosis aut villosis, ramis apteris.

113. B. AVICENNIFOLIA, fruticosa, ramis striatis glabris junioribus sulcato-angularis, foliis petiolatis obovato-oblongis obtusis basi attenuatis integerrimis penninerviis suprâ glabris subtus cano-tomentosis, racemis strictis subaphyllis, capitulis breviter pedicellatis ovatis, invol. squamis ovatis acutis. ♀ in Brasiliæ prov. Minarum Generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 276). Conyza racemosa Spreng. syst. 3. p. 508? Petioli 6 lin. Limbus fol. 4 poll. longus, 15-16 lin. latus. Capitula immatura mascula non satis nota, sed certè Baccharidis potius quàm Conyze. (v. s.)

114. B. LESSINGIANA, fruticosa, ramis teretibus, foliis petiolatis ellipticis basi obtusis apice acutis integerrimis suprâ junioribus araneosis adultis glabriusculis subtus cum petiolis ramis pedunculisque densè cano-hirsutis, paniculâ terminali, invol. ♂ squamis ovato-lanceolatis ciliatis, ext. dorso subhirsutis, int. sublinearibus pappo brevioribus. ♀ in Brasiliâ. Bacch. nov. sp. Less! mss. in Lund. pl. miss. Flores ♂ in invol. circ. 15, majusculi. Achænium immat. pilosiusculum. Pappus subcrispus rufescenti-subflavicans. Petioli semipoll. Limbus fol. 3-3 ½ poll. longus, 15 lin. latus. Fl. ♀ ign. (v. s. comm. à cl. Lund.)

115. B. TARCHONANTHOIDES, fruticosa, ramis teretibus pedunculisque molliter tomentosis, foliis subpetiolatis lanceolatis obtusis basi longè cuneatis integris à medio ad apicem grossè serratis coriaceis suprâ glaberrimis subtus tomentosis penninerviis reticulatis, paniculâ terminali hirsutâ ramosissimâ polycephalâ subaphyllâ, inv. ♂ campanulati squamis hirtellis oblongo-linearibus acutis. ♀ seu ♀ in Brasiliæ prov. Minarum General. legit cl. Vauthier (pl. exs. n. 275). Folia 2 ½-3 poll. longa, 8-10 lin. lata. Cor. ♂ et pappus crassiusculus invol. æquales. Ind. fœm. ign. (v. s. comm. à cl. inv.)

116. B. GRANDIFLORA (H. B. et Kunth nov. gen. am. 4. p. 58), fruticosa volubilis, ramulis teretiusculis pubescenti-tomentosis, foliis petiolatis oblongis acuminatis basi acutis argutè dentatis subrugosis suprâ glabris subtus fusco-tomentosis, corymbis terminalibus dichotomis, invol. ♀ subhemisphærici squamis lineari-lanceolatis acuminatis subfimbriatis, achænio sulcato scabriusculo. ♀ in planitie Quitensi juxta Mulalo. Folia 3 poll. longa, 10-11 lin. lata. Pl. ♂ ign.

117. B. MOLLIS (H. B. et Kunth nov. gen. am. 4. p. 58), fruticosa, ramis teretibus fuscescenti-villosis, foliis subsessilibus ovatis obtusis basi rotundatis crenatis suprâ pilosiusculis subtus molliter villosi-pubescentibus, dentibus mucronatis, capitulis in spicas axillares terminalesque digestis sessilibus conglomeratis,

invol. ♂ hemisphærici squamis lanceolatis acutis. ♀ circa urbem Quito. Folia sesquipollicaria. Pl. ♀ ign.

118. *B. UNCINELLA*, fruticosa ramosissima, ramis teretibus glabris junioribus hirsutis, foliis sessilibus confertis patulis coriaceis integerrimis obovato-oblongis supra glaberrimis subtus cano-tomentosis junioribus apice mucrone recurvo uncinulatis adultis obtusis, capitulis ♂ paucis ad apices ramulorum axillaribus subcongestis, invol. ovati squamis exter. dorso subhirsutis, inter. linearibus glabrinisculis. ♀ in Brasiliæ prov. Sancti-Pauli. Rami conferti corymbosi breves. Folia 3 lin. longa, $1\frac{1}{2}$ lata. Flor. fœm. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 470 miss.)

119. *B. GNAPHALIOIDES* (Spreng. syst. 3. p. 461), fruticosa, ramis teretibus albo-tomentosis, foliis linearibus utrinque obtusis patulis reflexive uninerviis integerrimis margine subrevolutis supra araneosis subtus albo-tomentosis, capitulis ♀ racemoso-capitatis breviter pedicellatis, invol. squamis ovatis subscariosis, acheniis glabris muricatis ♀ in Brasiliæ prov. Rio-Grande. Folia 5 lin. longa, lineam lata. Flor. in invol. ♀ circ. 20. Pappus rufus pluriserialis. Species affinis *B. velutina* sed in genere suo distinctissima. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 824 miss.)

120. *B. MELICHRYSOIDES*, fruticosa, ramis teretibus lanato-hirsutis, foliis sessilibus basi truncato-subcordatis lanceolatis acuminatis, summis ovato-lanceolatis, omnibus integerrimis planis supra pubescentibus subtus dense albo-tomentosis, capitulis laxè corymbosis pedicellatis, invol. campanulati squamis oblongo-lanceolatis hirsutis, in ♀ longioribus acuminatis, achenio subangulato glabro. ♀ in Brasiliæ prov. Rio-Grande et Sancti-Pauli. Pappus albus nitens. Folia 12-15 lin. longa, 3-4 lata. Spec. speciosa distinctissima. (v. s. ♂ et ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 811 et 492 miss.)

121. *B. VELUTINA*, fruticosa ramosa tota molliter velutina, ramulis teretibus, foliis sessilibus linearibus integerrimis 1-nerviis margine subrevolutis, capitulis ♂ ex axillis breviter pedicellatis racemosis pedicellis nudis, invol. squamis oblongis ♀ ex axillis longius pedicellatis, pedicellis bracteolatis, invol. squamis ovatis obtusissimis laxis rufo-velutinis floribus in ind. ♀ 5-6, acheniis glabris, corollâ rigidâ pappo pluriseriali. ♀ in Brasiliæ prov. Rio-Grande. An fortè *B. ochracea* Spreng. syst. 3. p. 460? Habitus ferè *Ericæ*. Folia 3-6 lin. longa, in ♂ acutâ, in ♀ obtusâ. Non satis constat an ind. ♂ et ♀ hic descripta ad unam eandemque speciem pertinet. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. ♂ sub n. 995, ♀ sub 819 miss.)

122. *B. ERIOCLADA*, fruticosa, ramis teretibus præsertim junioribus villosis hirsutis, foliis sessilibus ovatis basi obtusis apice mucrone acutis integerrimis aut hinc inde 1-2 dentatis coriaceis supra glabris subtus villosis subhirsutis, capitulis ♀ sessilibus axillaribus ad apices ramorum et ramulorum spicato-congestis, invol. squamis lanceolatis rufo-membranaceis glabris, achenio striato glabro. ♀ in Brasiliæ prov. Sancti-Pauli. Folia 6-7 lin. longa, 3 lin. lata. Pappus rufus invol. paulò longior. Fl. masc. ign. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 469 miss.)

123. *B. PHYLICÆFOLIA*, fruticosa, ramis teretibus junioribus canescentibus velutino-hirsutis, foliis sessilibus approximato-patentibus ovato-oblongis basi obtusis subcordatis, apice obtusiusculis submucronulatis margine vix subrevolutis supra glabris subtus cano-tomentosis, paniculâ subnudâ ramosissimâ apice subcorymbosâ, capitulis pedicellatis, invol. ♂ squamis oblongo-linearibus obtusis dorso lanato-hirsutis. ♀ in Brasiliæ prov. Sancti-Pauli. Accedit ad descr. *B. phyllicoidis*. Folia uninervia 6 lin. longa, $2\frac{1}{2}$ lata. Pappus ♂ crispus albidus vix invol. longior. Fl. ♀ ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 491 miss.)

124. *B. LEUCOPAPPA*, caule fruticoso ramoso juniore canescente demùm glabrato, ramis teretibus, foliis sessilibus alternis ovatis acuminatis integerrimis supra glabris subtus cano-tomentosis, marginibus subrevolutis, ramis floridis subpaniculatis capitula corymboso-conferta gerentibus, invol. squamis oblongo-

linearibus acutis velutinis subplanatis. $\frac{3}{4}$ in prov. Rio-Grande Brasiliæ. (h. Mus. imp. Bras. n. 330). Folia coriacea 6-8 lin. longa, 3-4 lata, adulta reflexa. Pappus albusissimus. (v. s. in h. Mus. reg. Par.)

125. B. PHYLLOIDES (H. B. et Kunth nov. gen. sm. 4 p. 49), fruticosa ramossissima, ramulis villosis-pubescentibus, foliis sessilibus approximato-patentibus ovato-oblongis obtusiusculis margine revolutis integerrimis coriaceis enerviis supra nitidis glabris, subtus villosis-pubescentibus, capitulis σ terminatibus capitato-conglomeratis sessilibus, invol. hemisphærici squamis obtusiusculis. $\frac{3}{4}$ in Peruviae vexillis Parami de Guarami. Cor. 13-14 albæ, \varnothing ign.

126. B. RADICANS, caulibus suffruticosis teretibus virentibus ascendentibus albidis subtomentosis basi radicanibus, foliis sessilibus linearibus patulis integerrimis acutis margine revolutis supra glabris subtus albidis-tomentosis adultis subglabratibus, corymbo, terminali subcomposito conferto 8-12-cephalo, invol. \varnothing squamis ovatis exter. obtusiusculis dorso villosis, inter. mucronato-acutis. $\frac{3}{4}$ in Brasiliæ prov. Rio-Grande. Species distinctissima. Folia 6-8 lin. longa, 1-1 $\frac{1}{2}$ lata, nervo medio supra exarato. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 906 miss.)

\S . 4. Oblongifoliæ, nempe foliis oblongis ovalibus linearibusve glabris aut rariis villosis uni- aut penninerviis non distichè imbricatis, ramis, apteris.

127. B. CAPRARIEFOLIA, fruticosa, ramis teretibus junioribus laxè et molliter villosis-hirsutis, foliis sessilibus ovalibus aut obovato-oblongis apice serraturis utrinquè 3-5 acutis notatis junioribus laxè villosis adultis glabriusculis, capitulis axillaribus sessilibus ad apices ramorum in spicam foliaceam subcontinuum digestis, invol. \varnothing squamis lanceolatis margine membranaceis subacutis, achænio glabro. $\frac{3}{4}$ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 966 et 968). Capitula masc. parva, fœm. duplò ferè majora. Pappus \varnothing rufus invol. paulò longior. Folia 7-10 lin. longa, 3-lin. lata. (v. s.)

128. B. ORGYALIS, fruticosa glabriuscula, ramulis subangulatis adultis glabris, foliis petiolatis elliptico-lanceolatis utrinquè acuminatis penninerviis subtus pallidioribus et punctulatis à medio ad apicem remotè denticulatis, corymbis axillaribus 5-7-cephalis in paniculam foliosam dispositis, capitulis \varnothing pedicellatis, inv. squamis int. elongatis acuminatis, achæniis glabris. $\frac{3}{4}$ in Brasiliæ Serra dos Orgaos legit cl. Lhostky septemb. flor. Fl. masc. ign. Pappus \varnothing albidus elongatus. Folia 3 poll. longa, pollicem lata. (v. s.)

129. B. MICRODONTA, fruticosa glabra apice viscosa, ramulis decurrenti-angulatis, foliis oblongo-lanceolatis in petiolum subattenuatis acutis minutè et acutè ad apicem serrulatis uninerviis aut subtrinerviis venis lateraliibus tenuissimis, capitulis σ ad axillas foliorum secus ramulos axillares solitariis sessilibus ovato-oblongis, inv. squamis lanceolatis. $\frac{3}{4}$ in Brasiliæ prov. Sancti-Pauli. Folia 15 lin. longa, 3 lata. Capitula pallida 2 lin. longa. Fœm. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 483 miss.)

130. B. MACRODONTA, fruticosa glabra viscosa, ramulis subangulatis, foliis petiolatis oblongis ovalibusve utrinquè acuminatis hinc indè à medio ad apicem grossè serratis penninerviis, venis inf. submajoribus et indè falsè triplinerviis, pedunculis axillaribus folio brevioribus capitula 5-7 pedicellata corymbosa gerentibus, invol. \varnothing squamis ovali-lanceolatis, int. linearibus, margine membranaceis, achæniis striatis glabris. $\frac{3}{4}$ in Brasiliæ prov. Sancti-Pauli. Valde accedit ad B. microdontam sed diversa satis videtur. Flor. masc. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 485 miss.)

131. B. PENTLANDII, fruticosa glabra viscosa, ramulis sulcatis, foliis petiolatis lanceolatis utrinquè acutis imà basi integris cæterum argutè serratis penninerviis reticulato-venosis, capitulis σ ad apices ramorum densè congestis inter folia suprema sessilibus, invol. hemisphærici squamis lanceolatis scariosis acutis subimbricatis. $\frac{3}{4}$ in Republ. Boliviana Amer. austr. legit cl. Pentland et me-

cum comm. Folia cum petiolo 4' lin. bipollicaria, 6-8 lin. lata. Flor. ♀ ign. (v. s.)

132. *B. BUDDLEIODES* (H. B. et Kunth nov. gen. am. 4. p. 64), fruticosa glabra, ramulis subangulatis, foliis petiolatis lanceolatis acuminatis basi angustatis serrulatis subcoriaceis reticulato-venosis, corymbis terminalibus diffusis, capitulis ♂ pedicellatis 70-80-floris, invol. hemisphærici squamis lanceolatis acutis subcoriaceis. ♀ in Peruvia? Folia 3 ½ poll. longa, 8 lin. lata, petiolo 8-9 lin. Fl. ♀ ign.
133. *B. BOGOTENSIS* (H. B. et Kunth nov. gen. am. 4. p. 61), fruticosa glabra, ramis virgatis sulcatis, foliis breviter petiolatis lanceolato-oblongis acutis argutè serrulatis coriaceis reticulato-venosis supra nitidis, paniculis ♂ terminalibus ramosis foliatis, invol. subhemisphærici squamis lanceolatis acutis subciliatis. ♀ in altà planitie Bogotensi juxta Santa-Fé, et inter Chuapa et Chipó. Folia 7-8 lin. longa, 2-2 ½ lata. Fl. ♀ ign.
134. *B. VOLUBILIS* (H. B. et Kunth nov. gen. am. 4. p. 57), fruticosa glabra volubilis, ramis striatis, foliis petiolatis ovato-oblongis breviter acuminatis basi subrotundatis argutè serrulatis rigidis nitidis, pedunculis axillaribus terminalibusque oligocephalis, invol. ♀ subhemisphærici squamis scariosis lanceolato-linearibus, achænio costato glabro. ♀ in frigidissimis Andium Peruv. ad radices Parani de Guaramo. Cor. 25-30 albæ, ♂ ign. Folia sesquipollic., 8 lin. lata.
135. *B. ADSCENDENS* (Pers. ench. 2. p. 424. n. 13), caule adscendente, foliis ovatis acuminatis profundè dentato-serratis. ♀ in Peruvia ubi dicitur *Chilco* ex herb. Juss. Cum *B. volubili* conjunxit Spreng. syst. 3. p. 463. sed distinetam servo quia cl. Kunth qui herb. Juss. vidit cum eà non imò confert. Cæter. ign. †.
136. *B. ARBUTIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 54), fruticosa glabra, ramulis angulatis, foliis sessilibus oblongis utrinquè acutis argutè serratis coriaceis nitidis, capitulis terminalibus corymboso-conglomeratis breviter pedunculatis, invol. squamis ovato-lanceolatis. ♀ in Peruvia (J. Juss.), in devexis montis Pichinchæ Quitensium (H. B.). *B. arbutifolia* Vahl symb. 3. (1794) p. 97? *Conyza arbutifolia* Lam. dict. 2. (1786) p. 92.
137. *B. TEINDALENSIS* (H. B. et Kunth nov. gen. am. 4. p. 54), fruticosa ramosissima glabra apice viscosa, ramulis angulatis, foliis breviter petiolatis elliptico-oblongis utrinquè acutis grossè dentatis subcoriaceis, capitulis terminalibus conglomeratis breviter pedunculatis, inv. hemisphærici squamis scariosis acutiusculis. ♀ in altissimis inter Teindala et Pasto Novo-Granatensium. Cor. albæ in utroque sexu 30-40. Folia 14-15 lin. longa, 7-8 lata.
138. *B. MACRANTHA* (H. B. et Kunth nov. gen. am. 4. p. 54), fruticosa ramosissima glabra viscosa, ramis teretibus, foliis brevissimè petiolatis oblongis utrinquè acutis remotè et obsolete dentatis subcoriaceis, capitulis ♀ terminalibus capitato-congestis breviter pedunculatis, invol. squamis scariosis acutiusculis. ♀ propè Mulado Quitensium ad radices montis Cotopaxi. Capit. magna flores circ. 75 invol. longiores gerentia. Folia 12-14 lin. longa, 4-5 lin. lata.
139. *B. CAPITATA* (H. B. et Kunth nov. gen. am. 4. p. 53), fruticosa ramosissima glabra, ramulis subangulatis viscosis, foliis petiolatis subelliptico-obtusiusculis basi acutis sinuato-dentatis subcoriaceis glabris viscosis, capitulis ♀ terminalibus capitato-congestis sessilibus, inv. subhemisphærici squamis scariosis obtusiusculis, achænio sulcato glabro. ♀ in America meridionali. Folia 10-11 lin. longa. Fl. ♂ ign.
140. *B. ELEGANS* (H. B. et Kunth nov. gen. am. p. 60. t. 324), fruticosa glabra, ramis sulcato-angulatis, foliis petiolatis ellipticis acutis basi rotundatis apicem versus argutè dentatis supra nitidis reticulato-venosis, corymbis terminalibus ramosissimis divaricatis foliatis, capitulis ♀ ovali-oblongis, inv. squamis lanceolatis acutis, achæniis glabris. ♀ in Mexico? Pedicelli parè bracteolati. Flor. masc. ign. Affinis videtur *B. petiolaris*.

141. *B. MICRANTHA* (H. B. et Kunth nov. gen. am. 4. p. 61), herbacea? caule tereti striato hirtio, foliis petiolatis oblongis acutis basi subrotundatis argutè dentatis subcoriaceis densè hirtis, paniculâ terminali ramosissimâ foliatâ patulâ, invol. ♂ hemisphærici squamis lanceolatis acutis subciliatis. — Propè Guanaxuato Mexicanorum. Cor. alba. Fl. fœm. ign.
142. *B. BRACTEOLATA*, suffruticosa subviscosa, ramis junioribus subhirsutis, foliis linearibus obtusiusculis basi attenuatis confertis punctulatis puberoglandulosis integerrimis aut sæpius denticulatis, paniculâ brevi, squamulis acutis secus ramulos bracteolatâ, invol. squamis linearibus acuminatis puberulo-ciliatis. ♀ patria ign. Prov. juxta Poir. ex itin. Commers. Specimen habeo olim sub nom. *Chrysomæ* in hort. Mousseaux lectum. — *Baccharis* Lam. ill. t. 698. f. 2. *Bacch. parviflora* Poir. suppl. 1. p. 556. non Pera. Capit. specim. mei ut Poirerii cassa et ideo genus dubium. (v. s.)
143. *B. MUCRONATA* (H. B. et Kunth nov. gen. am. 4. p. 60 non Hook.), herbacea?, caule tereti densissimè hispido-hirtio, foliis petiolatis oblongis utrinquè acutis mucronatis remotè et argutè denticulatis rigidis hirtis, paniculâ terminali ramosâ patulâ bracteâtâ, invol. ♂ squamis lanceolatis acutis ciliatis. — in Mexici montibus Guanaxuatensibus. Fl. fœm. ign.
144. *B. SERRATIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 59), fruticosa, ramulis subangulatis hirtio-pubescentibus, foliis brevissimè petiolatis ovato-ellipticis brevissimè acumminatis basi angustatis argutè serratis rigidis suprâ hispidopilosis subtus molliter pilosis, paniculis terminalibus ramosis patulis bracteatis, invol. ♂ squamis lanceolatis acutis apice ciliatis. ♀ in montibus Mexicanis propè Guanaxuato (H. B.). An etiam in Martinicâ ut vult Spreng. syst. 3. p. 463? Bractæ lanceolatæ hispidæ Fl. fœm. ign.
145. *B. HIRTELLA*, fruticosa ramosa, ramis teretibus junioribus pubescenti-subhirtellis, foliis breviter petiolatis ellipticis acuminatis penninerviis membranaceis suprâ sparse setulos subtus æcus nervos hispidulis margine remotè et callosè serratis, paniculis ramulos terminantibus, capitulis ♂ parvis, invol. squamis lanceolato-linearibus subciliatis vix acutis. ♀ inter Mexicanas herb. Hænkeani vidi. Fl. fœm. ign. Fortè affinis *B. serratifoliæ* H. B. K.? sed folia nec rigida nec brevissimè acuminata nec subtus molliter pilosa. (v. s. in h. Hænk. à cl. de Sternberg miss.)
146. *B. SESSILIFOLIA*, fruticosa, ramis striatis sæpius divaricatis, foliis sessilibus rigidis ovatis acutis basi obtusis triplinerviis reticulato-venosis serratis suprâ glabris, paniculâ pyramidato-corymbosâ, invol. campanulati squamis margine scariosis glabris in ♂ exter. obtusis, in ♀ exter. ellipticis in utroq. sexu inter. linearibus, achæniis puberulis. ♀ in Chili.
- ♀. *hebecleda*, ramis folisque subtus pubescenti-hirtellis. ♀ ad Talcaguana (Cham.!). *M. sessilifolia* Less.! in linnæâ 1831. p. 140. *Baccharis ripariæ* Poepp.! coll. chil. n. 209. (v. s. ♂.)
147. *B. PETIOLATA*, glabra herbacea erecta ramosa, ramis teretibus apice subglandulosis, foliis petiolatis ellipticis grossè dentatis corymbis pedunculatis nudis compositis, invol. ♂ campanulati squamis ovato-lanceolatis subacutis. — in Chili ex h. Hænke. Species distinctissima. Petioli sesqui- aut bilineares. Limbus folii ro lin. longus, 4-5 latus. Fl. fœm. ign. (v. s. in h. Hænke à cl. de Sternberg comm.)
148. *B. XEROPHORA*, suffruticosa erecta glabra, ramis ad collum ortis sterilibus pilosiusculis, foliis sessilibus lineari-sublanceolatis integerrimis uninerviis, caulinis glabris, rameis pilosiusculis, capitulis in racemum laxum subpaniculatum dispositis, invol. fl. ♀ squamis ovali-oblongis glabris subobtusis pappo dimidio brevioribus, achæniis pube densâ et globulis glandulosis onustis. ♀ in campis editis prov. Sancti-Pauli legit cl. Lund. Pappus rigidulus basi rufescens. Cor. ♀ teretes quasi truncatæ minutissimè 4-dentatæ. Styli basi breves parum exserti. Flor. ♂ ign. et ideo genus dubium. (v. s. comm. à cl. Lund.)

149. *B. SEMISERRATA*, fruticosa erecta ramosa glabra, ramis junioribus angulatis adultis glabris, foliis sessilibus lineari-lanceolatis acutis uninerviis inferne integerrimis supernè argutè serratis, capitulis 3-5 corymbosis pedicellatis, invol. ♂ hemisphaerici squamis lanceolatis acuminatis margine eroso-scariosis, fl. circ. 30. ♀ in Peruvia Andium vallibus legit cl. Hænke. Folia 1-1½ poll. longa, 2 lin. lata. (v. s. comm. in h. Hænke à cl. de Sternberg miss.)
150. *B. PTARMICIFOLIA*, suffruticosa glabra, ramis striato-sulcatis, foliis subsessilibus lineari-lanceolatis utrinquè acuminatis argutè serratis subtrinerviis subtus punctatis, paniculâ terminali laxâ polycephalâ foliosâ, invol. ♀ squamis oblongo-linearibus subobtusis margine membranaceis ciliatis. ♀ in Panamaide (Née! ex h. Thib.). In hortis Mexicanis legit. cl. Berlandier julio flor. (pl. exs. n. 572). Valde affinis *B. thesioidi* sed folia 2 poll. ferè longa, 3 lin. lata, subtus punctata basi subtrinervia. (v. s.)
151. *B. SULCATA*, fruticulosa ramosissima gracilis glabra, ramulis sulcato-angulatis, foliis sessilibus linearibus acuminatis à medio ad apicem serratis utrinquè impresso-punctatis, paniculâ terminali laxâ foliis intermixtâ, capitulis ♂ ad apices pedicellorum solitariis, invol. squamis lanceolatis acutis margine scariosis. ♀ in Mexico circa Villalpando legit cl. Mendèz. Species inter eas quas vidi distinctissima, fortè *B. thesioidi* affinis. Fl. foem. ign. (v. s. ♂ comm. à cl. Alaman.)
152. *B. THESIODES* (H. B. et Kunth nov. gen. am. p. 61), fruticulosa glabra. ramulis sulcato-angulatis, foliis brevissimè petiolatis lineari-lanceolatis basi acutis apice acuminatis argutè serrulatis (nerviis rigidolis, paniculâ terminali ramosâ foliatâ, invol. squamis lanceolatis oblongis subobtusis fimbriato-ciliatis, pappo ♂ setiformi, acheniis glabris. ♀ in Mexico (Mair!), circa urbem (Berl. pl. exs. n. 529), juxta Santa-Rosa in montanis (H. et B.). Fol. 10-11 lin. longa, 1-2 lata. Affinis *B. multifloræ*. (v. s. ♂ et ♀.)
153. *B. MULTIFLORA* (H. B. et Kunth nov. gen. am. 4. p. 59), suffruticosa, caulibus simplicibus striatis puberulo-hirtis, foliis brevissimè petiolatis oblongis acutis basi rotundatis argutè serratis suprâ glabris subtus præcipuè secus nervum et petiolum puberulo-hirtis, paniculâ terminali ramosissimâ subfoliatâ, invol. utriusque sexûs squamis lanceolatis acutis margine scariosis fimbriato-scariosis, pappo ♂ apice penicillato-plumoso. ♀ in Mexico (Mair!), circa urbem (Berl. pl. exs. n. 900), et ad Tianguillo et Toluca in frigidis montanis (H. B.). Spec. ♀ descr. cl. Kunth et mea sunt omnia ♂. (v. s.)
154. *B. MYRSINITES* (Pers. ench. 2. p. 424. n. 25), fruticosa ramosissima, ramulis angulatis, foliis lanceolatis confertis apice sæpius bidentatis, capitulis terminalibus sessilibus congestis globosis ovatisve, invol. squamis acutis apice subciliatis. ♀ in Santo-Domingo. *Conyza myrsinites* Lam. dict. 2. p. 92. Spreng. syst. 3. p. 465 excl. syn. Poir. Habitus *B. Magellanicæ* et *cuneifoliæ* ex Lam.
155. *B. CHILCO* (H. B. et Kunth nov. gen. am. 4. p. 56), fruticosa ramosissima glabra viscosa, ramulis angulatis, foliis brevissimè petiolatis lanceolato-oblongis obtusiusculis basi cuneatis apicem versus utroque latere dente 1-2 instructis subcoriaceis subaveniis, capitulis ♀ terminalibus congestis subsessilibus, invol. squamis scariosis acutiusculis. ♀ prope Santa-Anna Novo-Granatensium, ubi dicitur *Chilco*. Plures aliæ species ex R. et Pav. in Peruvâ nuncupantur *Chilca*. Fol. 7 lin. longa. Fl. ♂ ign.
156. *B. CHILQUILLA*, fruticosa junior pulverulento-puberula dein glabra subviscosa, ramulis teretiusculis, foliis linearibus elongatis sessilibus acutis hinc inde grossè dentato-serratis aut integerrimis subuninerviis, corymbo terminali polycephalo, invol. ♂ ovati squamis scariosis stramineis glaberrimis ovali-oblongis subobtusis. ♀ in dumetis et sylvaticis secus flumina ad Quillota Chilensium octob. flor. legit cl. Bértero. Vulgò *Chilquilla del Rio*. Folia 3-4 poll. longa sesquillin. lata. Affinis *B. marginali*. (v. s.)
157. *B. ROSMARINIFOLIA* (Hook. et Arn. bot. Beech. 1. p. 30), caule fruticoso ra-

moso dense folioso, foliis linearibus obtusiusculis crasso-coriaceis integerrimis margine subrevolutis ramulisque resinoso-glandulosis, corymbis terminalibus dense oligocephalis, invol. ovalis squamis elliptico-oblongis ext. obtusis, int. subacutis, achæniis glabris. ζ in collibus fruticetis sylvaticis et arvis ad Valparaiso (Bert.), Talcahuano (Cham.), Conceon (Poepp.), Rancagua (Bert.), Conceptionem (d'Urv.), Colchagua (C. Gay!) Chilensium. Vulgò *Romero*, *Romerillo*, *Romero de la tierra*. *Baccharis lingulata* Kunze! in Poepp. coll. 1. n. 212. Less. in linnæa 1831. p. 147 et 506. An ex nom. vulg. *B. linearis* R. et Pav. syst. p. 205 potius hic quàm sequenti referenda videtur? (v. s.)

ß. subsinuata, foliis plurimis obtusè sinuato-repandis. — Verisim. in Chili legit cl. Dombey. Capitula pedicellata umbellata omnino ut in var. α . (v. s. σ ex h. itin. Dombey comm. à Mus. reg. Par.)

158. *B. OBLONGIFOLIA* (Spreng. syst. 3. p. 460 excl. syn.), fruticosa, foliis oblongis integerrimis, corymbis terminalibus. ζ in sylvis circa Conceptionem Chilensium et ad Hualpen tractus. Molina oblongifolia R. et Pav. syst. 203. Cæc. ign. An fortè var. latifolia et integrifolia *B. Pingræ*? \dagger .

ß. glabriuscula, ramis foliisque subtus glabris aut glabriusculis. ζ in fruticetis collium circa Valparaiso. *Baccharis rigida* Hook. et Arn. bot. Beech. 2. p. 57. (v. s. σ et \mathcal{Q} .)

159. *B. PINGRÆA*, caule suffruticoso ramoso laxè folioso, foliis linearibus aut lineari-lanceolatis integerrimis serratisve acuminatis ramisque punctulatis, corymbis laxis polycephalis, invol. ovato-subglobosi squamis in σ oblongo-lanceolatis macronatis, in \mathcal{Q} inter. acuminatis, achæniis glabris. ζ in Chili (R. et Pav.), (Cham.), frequens ad ripas arenosas fluviorum prope Conceon (Poepp.), San-Yago (Bert.), Coquimbo (Gaud.) et Valparaiso (Macr.), et in California (Cham. ex Less.), in prov. Rio-Grande (h. imp. Bras.) et ad Buenos-Ayres (Bacle!). Molina linearis var. R. et Pav. syst. p. 205? Hook. et Arn. in Beech. bot. 1. p. 57. Molina linearis Less. in linnæa 1831. p. 139 et 505.

a. latifolia, foliis lineari-lanceolatis serratis rariùs subintegris imò integerrimis. *B. linearis* Poepp. coll. chil. II. n. 103. *Conyza angustifolia* hort. Par. 1828. *Baccharis angustifolia* Desf. cat. ed. 3. p. 163 non Michx. et excl. patriâ. *Pingræa angustifolia* Cass. dict. 41. p. 58. (v. s. σ ex Poepp. et hort. Par. \mathcal{Q} ex Don et Cham.)

ß. angustissima, foliis angustissimè linearibus integerrimis aut rariùs denticulatis. (v. s. σ ex Gaud. \mathcal{Q} ex Née et Macrae.)

160. *B. LINIFOLIA*, herbacea? erecta simplex glabra, caule tereti, foliis linearibus elongatis acutis margine sub lente scabro-serrulatis primâ fronte 1-nerviis, sub lente 3-nerviis, venis lateralibus tenuissimis, paniculâ subcorymbosâ confertâ, invol. σ squamis lanceolatis acutis vix apice subciliatis. — In Mexico ad Real del Monte ex h. Hænke. Habitus ferè *Rigetowizæ*. Flor. masc.? in invol. 15-16. Pappus rufus invol. longit. æqualis. Fl. fœm. ign. (v. s. in h. Hænke à cl. de Sternberg miss.)

161. *B. PANICULATA*, fruticosa subviscosa glabriuscula, foliis linearibus acuminatis integerrimis subtrinerviis acumine reflexo, paniculâ elongatâ foliosâ, invol. \mathcal{Q} cylindraceis squamis glabris margine scariosis oblongo-lanceolatis, intimis acutis, achæniis glaberrimis. ζ in Chili circa Valparaiso (Macrae!) et Coquimbo (Gaud.). Affinis *B. Pingræ* sed dispositione capitulorum et foliis apice incurvis faciliè differt. Fl. σ ign. Cor. \mathcal{Q} truncatæ apice subpuberulæ. (v. s. comm. ab hon. Soc. hort. Lond. et ab amic. Gaudichaud.)

162. *B. PAUCIDENTATA*, fruticosa ramosissima glabra, ramulis striato-angulatis, foliis sessilibus linearibus aut lineari-oblongis utrinquè acutis uninerviis aut imâ basi subtrinerviis integerrimis aut dense 1-2 utrinquè notatis, capitulis \mathcal{Q} ad axillas superiores subsessilibus et idè in spicas breves digestis, invol. squamis lanceolatis acutis margine membranaceis, floribus in invol. circ. 10, achæniis striatis glabris. ζ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. sub n. 823 et 840), et ad Buenos-Ayres? Affinis *B. tenuifoliæ*. Ind. masc. ign. *Folia* 9-12 lin. longa, 1-1½ lata. (v. s.)

163. *B. DRACUNCULIFOLIA*, fruticosa ramosa ad apicem subpubescens, ramulis angulatis, foliis sessilibus lineari-oblongis acutis utrinque punctatis uninerviis basi integerrimis apice dentibus acutis parvis utrinque 2-3 notatis, capitulis in axillis super. breviter pedicellatis in racemum foliosum digestis, invol. squamis lanceolatis acutis margine membranaceis, floribus in invol. 25-30, achæniis glabris. $\frac{5}{8}$ in Brasiliæ prov. Rio-Grande, ad Sanctam-Catharinam et Rio-Janeiro. Affinis *B. paucidentata*. Capitula σ^7 pauciora. Variat foliis integerrimis dentatisve. Folia 9-11 lin. longa, 2-2 $\frac{1}{2}$ lin. lata. (v. s. σ^7 et φ comm. à cl. Lund. et in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 817 et 819 miss.)
164. *B. CEREALOTIS*, fruticosa glabra, ramulis teretiusculis, foliis sessilibus ellipticis utrinque acutis basi integerrimis apice parçé et minutè serratis uninerviis coriaceis erectis subimbricatis, capitulis ad apices ramorum et ramulorum brevium confertissimis, foliis summis bracteatis, invol. σ^7 squamis oblongis ciliatis subacutis. $\frac{5}{8}$ in Brasiliæ prov. Sancti-Pauli. Folia 7 lin. longa, 2-2 $\frac{1}{2}$ lata. Capit. σ^7 4-5 lin. longa, φ ign. Pappus σ^7 rufus invol. paulò longior. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 476 miss.)
165. *B. LEPTOSPERMOIDES*, fruticosa glabra, ramis teretibus junioribus decurrenti-angulatis, foliis linearibus sessilibus utrinque attenuatis integerrimis aut hinc inde subdentulatis penninerviis venis subreticulatis inter venas subtus punctatis, capitulis σ^7 axillaribus solitariis breviter pedicellatis folio brevioribus, invol. squamis oblongis subobtusis pappum æquantibus. $\frac{5}{8}$ in Brasiliæ prov. Sancti-Pauli. Affinis *B. axillari*. Capitula in medio ramorum sita; rami apice foliosi nec floriferi. Folia 9-10 lin. longa, sesquilineam lata. Fl. cœm. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 482 miss.)
166. *B. QUITENSIS* (H. B. et Kunth nov. gen. am. 4. p. 57), fruticosa ramosissima glabra viscosa, ramis sulcato-angulatis, foliis petiolatis oblongis acuminatis basi acutis integerrimis reticulato-venosis coriaceis nitidis, corymbis φ terminalibus axillaribusque, invol. hemisphærici squamis scariosis acutis. $\frac{5}{8}$ in altâ planitie Quitensi ad radices montis Ruca-Pichincha. Folia 2-4 $\frac{1}{2}$ poll. longa, 3-11 lin. lata. Pl. σ^7 ign.
167. *B. BRACHYLÆNOIDES*, fruticosa glaberrima nitida, ramis teretibus, foliis breviter petiolatis lanceolatis utrinque acuminatis integerrimis penninerviis, patulis rigidis subtus pallidioribus, paniculâ subcorymbosâ laxâ ferè aphyllâ polycéphalâ, invol. σ^7 squamis oblongis acutis brevibus, φ squamis linearibus acuminatis elongatis, achæniis apice sparsè puberis. $\frac{5}{8}$ in Brasiliæ monte Cercovado angusto flor. legit cl. Lund. Species pulchra habitu *Brachylæna* neriifoliam referens. Fol. 4-5 poll. longa, 9-12 lin. lata. Pappus albus 1-serialis. (v. s. comm. à cl. Lund.)
168. *B. VENULOSA*, fruticosa ramosa erecta glabra, ramis teretibus, foliis in petiolum brevem attenuatis subcoriaceis lanceolatis utrinque acuminatis integerrimis subclucidis utrinque venuloso-subreticulatis medio 1-nerviis, paniculâ terminalis corymbosâ laxâ ramis ramulisque puberulis, invol. σ^7 squamis lanceolatis acutis, flor. in inv. circ. 15-20. $\frac{5}{8}$ in Peruvæ montibus Oronocensis legit cl. Hænke. Affinis *B. brachylænoidi*. Folia 3-4 poll. longa, 8-10 lin. lata. Pappus rufescens. Fl. φ ign. (v. s. in h. Hænke à cl. de Sternberg comm.)
169. *B. MITIDA* (Pers. ench. 2. p. 425. n. 41), fruticosa, foliis oblongo-lanceolatis integerrimis denticulatisque, capitulis axillaribus corymboso-racemosis. $\frac{5}{8}$ in Peruvæ sylvaticis altis versus Pillao et Panao tractus. Molina nitida R. et Pav. syst. p. 204. Cœt. ign.
170. *B. LIGUSTRINA*, fruticosa glabra, ramis teretibus apice angulatis, foliis oblongo-sublanceolatis sessilibus integerrimis apice mucrone calloso et denticulis sæpè 1-2 minimis instructis uninerviis venosis, paniculâ terminali subnudâ polycéphalâ, capitulis σ^7 pedicellatis, invol. squamis paucis lanceolatis acutis margine membranaceis pappum æquantibus. $\frac{5}{8}$ in Brasiliæ prov. Minas Generalim ad Serra do Frio legit cl. Vanthier (pl. exs. n. 288). Fl. cœm. ign. Folia sesquipoll. longa, 3 lin. lata. (v. s.)

171. *B. PRINOIDES* (H. B. et Kunth nov. gen. am. 4. p. 56), fruticosa ramosissima glabra viscosa, ramulis striatis, foliis petiolatis oblongis acutis basi angustatis integerrimis coriaceis nitidis, pedunculis ♀ axillaribus 5-7-cephalis, invol. squamis scariosis acutiusculis, achæniis sulcatis glabris. ♂ in calidis prov. Jaen de Bracamoros, in ripâ flum. Amazonum. Folia 2 poll. longa, 10 lin. lata, petiolo 4-5 lin. Cor. in inv. 30-35, ♂ ign.
172. *B. VERNONIODES*, fruticosa, ramis teretibus dense hirsuto-hispidis, foliis sessilibus elliptico-oblongis basi obtusis apice subacutis integerrimis penninerviis suprâ puberulis subtus villosulo-hirsutis, paniculis amplis foliis oblongis intermixtis hirsutis, capitulis ♀ pedicellatis, invol. squamis ext. ovalibus dorso hirsutulis inter. longioribus membranaceis glabris, achænio puberulo, pappo 1-seriali. ♂ in Brasiliæ prov. Minarum General. ad Mariannam legit cl. Vauthier (pl. exs. p. 265). Habitus ferè Vernoniæ puberulæ. Hirsuties rufa. Folia 2 poll. longa, 6-8 lin. lata. Pappus albus. Fl. masc. ign. (v. s.)
173. *B. ORONOCENSIS*, fruticosa erecta glabra, ramis teretibus vix apice angulatis, foliis petiolatis ellipticis utrinquè acuminatis subintegerrimis subcoriaceis penninerviis, racemis axillaribus vix petiolo longioribus, capitulis ♂ pedicellatis 10-12-floris, invol. squamis ovato-oblongis subacutis. ♂ in Peruvîa montanis Oronocensibus legit cl. Hænke. Folia cum petiolo 6 lin. 3 poll. longa, 12-15 lin. lata; nervi laterales 8-9; denticuli 1-2 ad apices foliorum sæpè videntur. Flor. ♀ ign. (v. s. in h. Hænke. ab ill. de Sternberg comm.)
174. *B. VACCINIODES* (H. B. et Kunth nov. gen. am. 4. p. 50), fruticosa glabra viscosa, ramulis angulatis, foliis sessilibus oblongis utrinquè acutis subintegerrimis coriaceis aveniis, capitulis ♀ terminalibus conglomeratis sessilibus, invol. hemisphærici squamis scariosis ext. ovatis, inter. oblongis linearibusve, achæniis sulcatis glabris. ♂ in Mexico temperato prope Moram. Folia ad apicem utrinquè subunidentata, 5-6 lin. longa, 2 ½ lata. Ind. masc. ign.
175. *B. BUXTIFOLIA* (Pers. ench. 2. p. 424, n. 24), fruticosa glabra, ramis virgatis angulatis tuberculatis, foliis subsessilibus confertis oblongo-ovatis integerrimis subretusis, capitulis lateralibus axillaribus solitariis sessilibus, invol. squamis ovali-oblongis apice subciliatis. ♂ in Peruvîa ubi ex Jos. Juss. mss. *Almisquillo* dicitur. *Conyza buxtifolia* Lam. dict. 2. p. 92. Folia 6-7 lin. longa. Pappus albidus.
176. *B. CORIDIFOLIA*, fruticosa ramosa glabra, ramulis teretiusculis, foliis sessilibus linearibus mucrone acutis margine ciliis rigidulis brevibus serrulato-scarbris, caulibus elongatis, ramis brevioribus, paniculâ elongatâ terminali, capitulis pedicellatis 5-8-floris, invol. squamis ovali-oblongis margine membranaceis obtusis, achæniis pubescentibus. ♂ in Brasiliæ prov. Rio-Grande et fortè Sancti-Pauli. Caulis basi repens. Panicula maris magis ramosa, capitulis parvis numerosioribus, ferè Artemisiæ. Inv. foem. duplò majus, squamis magis scariosis. Pappus rufus pluriserialis. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 810, 826, 837, 855 et fortè 510 miss.)
 ♀. invol. feminei squamis intimis magis oblongis subacutis. ♂ in prov. Rio-Grande. (v. s. cum var. α sub n. 994, 793, 858, 838 et 839.)
177. *B. MEGAPOTAMICA* (Spreng. syst. 3. p. 461), fruticosa glabra, ramulis subangulatis, foliis linearibus acutis integerrimis margine revolutis suprâ nitidis uninerviis eciliatis, capitulis ad axillas fol. sup. solitariis sessilibus in spicam dispositis, invol. ovati squamis lanceolatis acutis, achæniis glabris, pappo ♀ invol. non superante. ♂ in Brasiliæ prov. Rio-Grande et Sancti-Pauli. Spica nunc capitulis distantibus interrupta nunc subcontinua. (v. s. ♂ et ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 511, 816, 818, 847 miss.)
178. *B. REVOLUTA* (H. B. et Kunth nov. gen. am. 4. p. 50), fruticosa ramosissima, ramulis villosulo-pubescentibus, foliis brevissimè petiolatis lanceolato-linearibus integerrimis uninerviis margine revolutis subcoriaceis rigidis villosulo-pubescentibus, capitulis ♂ in corymbum terminalem dispositis, inv. hemisphærici squa-

- nus acuminatis scariosis. ♀ in Peruvia? Cor. ♂ 50-60 in inv. fœm., ign. Folia 8-9 lin. longa.
179. *B. LEDIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 50), fruticosa, ramulis cano-tomentosis, foliis sessilibus lanceolato-linearibus acutis integerrimis uninerviis supra incano-pubescentibus subtus cano-tomentosis, capitulis ♂ terminalibus congestis sessilibus, invol. squamis scariosis acuminatis. ♀ in Peruvia? Folia 14-15 lin. longa, 2-3 lata. Cor. masculæ albæ 35-40 in inv., fœm. ign.
180. *B. TENUIFOLIA*, fruticosa ramosissima glabra, ramulis subtriatis, foliis alternis et suboppositis linearibus obtusis basi subattenuatis sessilibus integerrimis margine (in sicco) subrevolutis, capitulis ad axillas summas sessilibus spicatis folio suo longioribus oblongis 5-floris, invol. squamis lanceolato-linearibus acutis, acheniis glabris. ♀ in Brasiliæ prov. Sancti-Pauli. Valdè accedit ad *B. leptophyllam*, sed foliis apice obtusis et sublatioribus viscositate ramisque nitidis et capitulis spicatis differt. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 512 miss.)
181. *B. LEPTOPHYLLA*, fruticosa ramosissima glabra, ramulis tenuibus striato-angulatis, foliis alternis et suboppositis sessilibus linearibus acutiusculis integerrimis 1-nerviis, capitulis ♀ ad apices pedicellorum 1-3 subracemoso-paniculatis, invol. oblongi squamis lanceolato-linearibus inter. elongatis, floribus in inv. 5-6, acheniis striatis glabris. ♀ in Brasiliæ prov. Rio-Grande. Proximè accedit ad *B. angustifoliam* et ad *tenuifoliam*. Folia 9-10 lin. longa, semilineam lata. Inv. 3 lin. longum. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 993 miss.)
182. *B. GRACILIS*, herbacea glabra, caulibus è cœlo plurimis gracillimis subandis striato-sulcatis, foliis sessilibus linearibus acutis tristriatis integerrimis remotis, capitulis paucis longè pedicellatis subcorymbosis, invol. squamis 2-3-seriatis lanceolatis acutis margine membranaceis subciliatis, acheniis glabris ferè invol. equalibus, pappo achenii longitud. paulò superante scabro. ♀ in Brasiliæ prov. Sancti-Pauli. Valdè accedit ad *B. juncea* sed caulis duplò tenuior et valdè sulcato-angulatus, capitula dimidio minima, pappus intense rufus ferè duplò brevior, achenia duplò longiora, etc. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 516 miss.)
183. *B. JUNCEA* (Desf. ! cat. h. Par. 1829. p. 163 excl. patr.), herbacea glabra subglaucescens, caulibus è collo plurimis teretibus junceis, foliis sessilibus linearibus subnerviis integerrimis distantibus, capitulis terminalibus sessilibus aut paucis corymbosis pedicellatis, invol. hemisphærici squamis lanceolatis rigidis acutis, intimis linearibus minoribus, acheniis glabris. ♀ in Brasiliâ meridionali ex Less. in linnæa 1831. p. 149, nec in Ægypto ut vult Lehmann, aut in Senegaliâ ut hab. hort. Par. *Stephananthus junceus* Lehm. cat. sem. h. Hamb. 1826. p. 18. ind. schol. 1828. p. 34. pug. pl. nov. p. 35. act. nov. nat. cur. v. 14. p. 2. p. 821 (descr. ind. masc.). *Arrhenachne juncea* Cass. ! dict. 52. p. 253 (descr. fœm. et masc.). Ad *Eupatorineas* ref. Lehmann et ad *Vernoniaceas* Cassini sed certè (ut jam rectè dixit Less.). *Baccharidis* species. Pappus ♀ pluriserialis pallidè rufescens angustè subpaleaceo-pilosus vix ac ne vix serrulatus. (v. v. ♀ cult. in hort. Par.)
184. *B. ANGUSTIFOLIA* (Michx. fl. bor. am. 2. p. 125), fruticosa glaberrima ramosissima, ramis teretiusculis, foliis angusto-linearibus integerrimis vix acutis, paniculâ ♂ laxâ oligocephalâ, ♀ corymbosâ compositâ polycephalâ, inv. ♂ ovati squamis lanceolatis subacutis, ♀ oblongi squamis acuminatis, acheniis striatis glabris. ♀ in maritimis Carolinæ et Floridæ (Michx. ! Fras. ! Ell. !) et ad ripas fl. Mississippi (Pursh). Frutex 6-10-pedalis. *Baccharis angustifolia* Ell. ! sketch. 2. p. 318 non Desf. Capitula circ. 20-flora. (v. s.)
185. *B. GENISTIFOLIA*, fruticosa ramosissima glabra, ramulis striatis, foliis distantibus paucis sessilibus linearibus obtusis integerrimis subnerviis, capitulis ♂ ad apices ramulorum 3-5 sessilibus spicato-congestis ebracteatis, invol. obovati squamis ovatis acutis margine membranaceis. ♀ in Brasiliæ prov. Rio-Grande. Habitus ferè *Genistæ* tinctoriæ. Folia 5 lin. longa, $\frac{1}{2}$ lin. lata. Ind.

foem. ign. Pappus ♂ invol. non superans. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 835 miss.)

§. 5. *Sergilæ, foliis minimis aut subnullis, ramis non alatis.*

186. B. SCOPARIA (Pers. ench. 2. p. 425. n. 44), fruticosa ramosissima glaberrima viscosula, ramulis gracilibus tetraquetris, foliis distantibus minimis lineari-oblongis obtusis integerrimis, capitulis ♂ terminalibus solitariis ovatis, invol. squamis ovatis. § in Jamaica australis montibus Liguansensibus. Calea scoparia Linn. syst. nat. 3. p. 234. Swartz! obs. 302. Chrysocoma scoparia Linn. amcen. 5. p. 381 et 404. Sergilus scoparius Gærtn. fr. 2. p. 409. t. 174. — Browne jam. 316. t. 34. f. 4. Vulgò *Mountain Broom tree* Anglis Jam. Spec. foem. non vidi et ergò genus dubium. (v. s. comm. à cl. Swartz et Bertero.)

187. B. APHYLLA, fruticosa ramosissima erecta glabra, foliis nullis aut ad squamulas minimas reductis, capitulis secus ramos sessilibus spicatis ebracteatis, invol. fl. ♀ squamis lineari-oblongis obtusis, pappo demum elongato apice crispo. § in campis editis prov. Sancti-Pauli Brasiliæ legit cl. Lund. Chrysocoma aphylla fl. flum. 8. t. 1. B. scopariæ affinis sed nullo modo viscosa, capitula multò majora et spicata, etc. Stirpem huic valde affinem ad Coquimbo Chiliensium lectam video in h. Lindl. An eadem? An species in posterum stabi- lienda? (v. s. comm. à cl. Lund.)

188. B. NUDA; fruticosa? dichotomè ramosa erecta glabra, foliis nullis, capitulis 4-5 secus ramos sessilibus spicatis ebracteatis, invol. ovali, pappo cor. ferè brevior. § in Brasiliæ prov. Fluminensi. Chrysocoma nuda fl. flum. 8. t. 50. Priori ex icone valde affinis. †.

§. 6. *Caulopteræ, nempe foliis oblongis linearibus aut subnullis, ramis alas foliaceas è foliorum basi utrinquè decurrentes geren- tibus.*

189. B. GAUDICHAUDIANA, fruticosa interruptè bialata, alis planis semi-ellipticis, foliis minimis obtusis, capitulis secus ramos ramulosve sessilibus spicato-panicu- latis, invol. squamis ext. parvis acutiusculis, inter. acuminatis, acheniis bre- vibus ellipticis glaberrimis. § in Brasiliâ ad Sanctam-Catharinam legit cl. Gau- dichaud. An Calacia sessilis fl. flum. 8. f. 73? Simillima B. articulata sed non glauca, articuli elliptici latiores nempe 7-8 lin. longi 3 lin. lati. (v. s. ♀ comm. à cl. Gandichaud et postea ♂ à cl. Lund.)

190. B. ARTICULATA (Pers. ench. 2. p. 425), fruticosa glauca interruptè bialata, alis planis angustis, foliis minimis obtusis, capitalis secus ramorum ramulo- rumve apices sessilibus spicatis, invol. campanulati squamis ext. obtusis, in- timis in foem. acuminatis, acheniis brevibus ellipticis glaberrimis. § in Brasiliâ meridion. propè Montevideo (Comm.! Sellow!) et in prov. Rio-Grande (h. Mus. Par. ex h. Mus. imp. Bras.). Conyza articulata Lam. dict. 2. p. 94. Molina articulata Less.! in linnæa 1831. p. 140. (v. s. ♂ et ♀.)

191. B. PARVIFOLIA, fruticosa interruptè bi- aut trialata, alis angustis planis, foliis ellipticis obtusis, capitulis distantibus lateralibus sessilibus ad apices ra- morum interruptè spicatis, invol. ♂ squamis ovali-oblongis vix acutis. § in Peruvia ad Avebrada de Cohrachillo ex h. Hænke. Affinis B. articulata sed faciliè distincta. Folia non squamiformia sed ovalia 2-2 ½ lin. longa. (v. s. in h. Hænke à cl. de Sternberg miss.)

192. B. CRISPA (Spreng. syst. 3. p. 466), fruticulosa trialata, alis crispis coria- ceis cartilagineo-marginatis, foliis ad squamulas minimas reductis, capitulis approximatis, invol. in ♂ cylindræci squamis acuminatis. § in Brasiliâ meridional! (Sellow!), prov. Rio-Grande (h. Mus. imp. Bras.). Molina crispa Less.! in linnæa 1831. p. 141. Alæ angustæ breves. Capit. 2 ½ lin. longa. Ind. foem. ign. (v. s.)

193. B. CANSTELLOIDES (Pers. ench. 2. p. 425 excl. syn. Maregr. ad Cereum

phyllanthum potius referend.), suffruticosa trialata, alis planis enerviis brevibus seu crebre interruptis, foliis minimis ad squamulas acutiusculas subternatas reductis, capitulis paucis (1-2) interrupte spicatis, invol. turbinati squamis omnibus acuminatis, rachide ex Less. nudâ et achæniis glabris cylindraceis sulcatis. ♀ in prov. Tacungâ Peruviae (Jos. de Jussieu!), propè Loxam et in Audibus Novo-Granatensium (H. et B.). *Conyza genistelloides* Lam.! dict. 2. p. 93. *Molina reticulata* Less. in linnæa 1831. p. 143 non R. et P. (v. s. ♂ comm. à cl. Jussieu.)

194. *B. venosa*, suffruticosa trialata, alis angustis venoso-reticulatis remotè interruptis, foliis ad squamulas minimas reductis, capitulis ad apices ramorum spicato-glomeratis, invol. campanulati squamis omnibus in ♂ acuminatis. ♀ in Peruviae argillosis collibus (R. et Pav.). *Molina venosa* R. et Pav. syst. 212. A *B. genistelloide* differt articulis longioribus et ferè angustioribus, alis venosis, superficie totâ viscosâ resinosa, et capitulis magis approximatis. (v. s. ex h. Dombey comm. à Mus. reg. Par.)

195. *B. microcephala*, fruticulosa trialata, alis planis angustissimis remotè interruptis, foliis minimis oblongis, capitulis interrupte spicatis solitariis aut adgregatis, invol. campanulati squamis in ♀ interioribus aentis, in ♂ omnibus obtusis, achæniis obovatis, costis minutè puberulis. ♀ in Brasiliâ (Sellow!) prov. Rio-Grande (h. Mus. reg. Par. ex h. imp. Bras.!). *Molina microcephala* Less.! in linnæa 1831. p. 142. Alæ caulinae punctulatæ sæpè 2-3 poll. longæ, 2-3 lin. latæ. Folia 1-2 lin. longa, ad apices ramorum omnino abortiva. Capit. parva. (v. s. comm. à Mus. reg. Berol.)

196. *B. sagittalis*, fruticulosa trialata, alis planis remotissimè interruptis, foliis lineari-lanceolatis acutis internodio multò brevioribus, capitulis in spicam terminalem continuam aut subinterruptam dispositis, invol. campanulato-ovatis squamis in ♂ omnibus obtusis, in ♀ extimis acutis intimis acuminatis, achæniis glabris. ♀ in Chili (Née!), propè Talcaguanho (Cham!), Valparaiso (Macrae!), Coquimbo (Gaud.), ad Conceptionem (R. et Pav.?) et in Cordilleræ excelsis (cl. Gay!). *Baccharis genistelloides* Poepp. coll. Chil. 1. n. 213 non Pers. ex Less. *Molina sagittalis* Less.! in linnæa 1831. p. 142. *Molina reticulata* R. et Pav. syst. p. 212? Folia 7-9 lin. longa. (v. s.)

♂. *Pappigii*, foliis infer. ovatis ovalibusve mucronatis, super. lanceolatis. ♀ in Chili. *Baccharis venosa*? seu *Baccharis tripterix* Poepp.! coll. Chil. 1. n. 210 diar. 46. (v. s.)

197. *B. phytumoides*, herbacea trialata glaberrima simplex seu ramosa, alis planis caule angustioribus, foliis internodio longioribus penninerviis oblongis basi attenuatis, capitulis glomeratis folio suo multò brevioribus in spicam interruptam dispositis, invol. campanulati in utroque sexu conformis squamis intimis acuminatis, achæniis obovatis secus costas minutè puberulis. ♀ in Brasiliâ meridionali (Sellow!). *Molina phytumoides* Less.! in linnæa 1831. p. 146. Folia 2-3 poll. longa, 4-9 lin. lata. (v. s.)

198. *B. pentaptera*, fruticulosa quinque-alata, alis planis remotè interruptis, foliis ellipticis obtusis internodio multò brevioribus, ramis apice subcorymbosis, capitulis spicatis adgregatis, invol. cylindracei squamis in ♂ ext. sub-acutis, inter. longioribus acutioribus. ♀ in Brasiliâ (Sellow!), prov. Sancti-Pauli (h. Mus. imp. Bras.). *Molina pentaptera* Less.! in linnæa 1831. p. 505. Fœm. non novi. (v. s. ♂ comm. à Mus. imp. Bras.)

199. *B. trimera*, suffruticosa trialata, alis planis reticulato-venosis hinc inde interruptis, foliis ad squamulas minimas acutas adpressas reductis, capitulis glomeratis interrupte spicatis, invol. campanulati squamis ext. acutis, intimis in ♀ acuminatis, in ♂ obtusiusculis aut acutis, achæniis oblongis glaberrimis. ♀ in Brasiliâ littorali (Sellow!), ad fretum Sanctæ-Catharinæ (Cham.), propè Rio-Grande (L. imp. Bras.), pr. Rio-Janeiro (Gaud.), Bahiam (Blanch.). *Molina trimera* Less.! in linnæa 1831. p. 141. An *Cacalia decurrens* fl. flum. 8. 1. 72? Alæ 3-4 lin. latæ 1-3 poll. longæ. (v. s.)

200. *B. CYLINDRICA*, herbacea trialata, alis planis venosis interruptis, foliis ad squamulas minimas reductis, capitulis spicato-glomeratis, invol. ♂ campanulato, in ♀ oblongo et cylindraceo, squamis acutis, ovariis glaberrimis. 2/ in Brasiliâ (Sellow!), prov. Rio-Grande (ex h. imp. Bras.). Molina cylindrica Less. in linnæâ 1831. p. 141. Differt à *B. trimerâ* capitulis valdè approximatis, ♀ oblongo-cylindraceis meianthis squamis eciliatis acutis nec intimis acuminatis. (v. s.)
201. *B. MILLEFLORA*, fruticosa ramosissima trialata, alis planis latis crebrè interruptis subreticulatis, foliis minimis vix ullis, capitulis spicato-glomeratis, invol. campanulati squamis in ♂ omnibus obtusis, in ♀ interioribus acutis, achæniis oblongo-ellipticis pube costarum minutâ. 3 in Brasiliâ legit cl. Sellow. Molina milleflora Less. in linnæâ 1831. p. 143. Alæ 4 lin. latæ, 1 1/2-2 poll. longæ. Recept. subnudum. Capitula numerosa. (v. s. comm. à Mus. reg. Berol.)
202. *B. MYRIOCEPHALA*, fruticosa ramosissima trialata, alis planis subreticulatis hinc indè interruptis, foliis ad squamulas minimas reductis, capitulis parè glomeratis interruptè spicatis, spicis numerosis paniculatis, invol. oblongi squamis in ♀ ext. brevibus obtusis, inter. elongatis apice mucronatis membranaceis, achæniis glabris. 3 in Brasiliæ prov. Minas Geraes legit cl. Vauthier (pl. exs. n. 265!) et in Bandâ orientali (Bacle!). Affinis *B. millefloræ*. Ind. masc. non vidi. (v. s.)
203. *B. JUNCIFORMIS*, herbacea? quinque-alata, alis planis latè linearibus reticulato-venosis, ramis apice angustissimè alatis, foliis ovali-acuminatis petioulatis internodio multò brevioribus, capitulis spicato-glomeratis, invol. ♀ campanulati squamis exter. obtusissimis intimis acutiusculis, ovariis oblongis minutè puberulis 2/ ? in Brasiliâ. Molina juncea Less. in linnæâ 1831. p. 145 et p. 505, sed non est *B. juncea* Desf. cat. h. Par. 1829. Folia 4-5 lin. longa. Masc. ign.
204. *B. POLYPTERA*, caule simplicissimo 7-8-alato, alis angustis valdè elongatis, foliis elongato-lanceolatis integerrimis acutis suprâ glaberrimis subtus cauleque inter alas densè et candidè tomentosis, spicâ interruptâ terminali valdè elongatâ, capitulis sessilibus fasciculatis, invol. ♂ tomentosi squamis ovalibus mucronatis. 3 in Brasiliæ prov. Rio-Grande. Species distinctissima! Inv. scem. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 746, 750 et 1066 miss.)

§. 7. *Distichæ, nempe foliis minimis distichè imbricatis carinatis ramos apteros obtegentibus.*

205. *B. FERRUGINEA* (Pers. ench. 2. p. 425. n. 48), fruticosa, ramis junioribus tomentosis foliis obtectis, foliis sessilibus semiamplexicanlibus carinatis approximatis bifariâ imbricatis extus glabriusculis intus tomentosis, capitulis (masculis) solitariis terminalibus solitariis. 3 in Peruvîæ montibus excelsis vulgò Cordilleras et Pnnas, ubi vern. *Matará* et *Palmito* appellatur ex R. et P. Molina ferruginea R. et Pav. syst. 211. Rami breves subcorymbosi. Folia angusta extus subrugulosa ferè linearia. Invol. squamæ apice membranacæ stramineæ subpatulæ. Pappus apice clavellatus. Valdè affinis *B. thyoidi*. An hujus fortè indiv. masculum? (v. s. ex h. Dombey à Mus. reg. Par. comm.)
- B. thyopsis*, foliis latioribus extus glaberrimis lævibus, invol. squamis apice scariosis sed non patulis. 3 in Amer. merid. cl. Bonpland olim comm. sub nomine *Conyzae* thyoidis sed in opere Kunthiano ommissa. Folia omninò *B. thyoidis* sed capitula terminalia mascula. (v. s.)
206. *B. THYOIDES* (Pers. ench. 2. p. 425. n. 49), fruticosa, ramis junioribus tomentosis foliis densè obtectis, foliis sessilibus semiamplexicanlibus carinatis approximatis bifariâ imbricatis extus glaberrimis intus lanatis, capitulis axillaribus sessilibus solitariis. 3 in Peruvîæ montibus primis legit Jos. de Jussieu! *Conyza thyoides* Lam. dict. 2. p. 90. ill. t. 697. f. 5. Molina incana

R. et Pav. syst. p. 211. Vulgò *Palmito* ex R. et Pav. Fl. masc. non vidl. (v. s. specim. ♀ ex J. Juss! Domb! Née! Abadia!)

207. B. CATAPHRACTA (Spreng. syst. 3. p. 462), ramis? distichè imbricatis conglutinatis glabris, foliis ovatis obtusiusculis rigidis, capitulis terminalibus sessilibus. — in Peru. *Conyza cataphracta* Willd. herb. Cæt. ign. †.

§. 8. *Lepidophyllæ*, foliis imbricatis squamæformibus, capitulis minimis aggregatis.

208. B.? LEPIDOPHYLLA, fruticosa ramosissima, ramis teretibus junioribus tomentosis, foliis confertissimis sessilibus ovatis integerrimis concavis glutinosis coriaceis, capitulis ♂? circ. 5 sessilibus ad apices ramorum ovatis glutinosis 1-3? floribus, invol. squamis ovalibus flavescentibus. ♀ in terrâ Van-Diemen prope Hobart-Town ad summitatem frigidam montis Wellington alt. 4100 ped. angl. supra mare legit cl. A. Cunningham. Genus novum *Baccharideum* Cunn! in litt. 1835. Habitus *Lepidophylli*, sed frutex dioicus et masculus tantum mihi cognitus. Videtur genus proprium sed characteres mihi non satis clari. (v. s. comm. & cl. Cunn.)

† Non satis notæ.

* *Gerontogæ*. (A genere alienæ.)

209. B.? ARBORÆA (Linn. mant. 284), fruticosa, foliis alternis petiolatis elliptico-lanceolatis integerrimis scabridis, corymbo terminali paniculato, invol. subglobosi squamis imbricatis subulatis, achæniis striatis. ♀ in Ind. or. insulæ Johannæ sylvis (Kœn. ex Linn.), in ins. Johannæ canal. Mozambici nec Ind. or. (Spreng.). B. arborescens Decaisn. nouv. Ann. Mus. h. n. Par. 3. p. 412. dicitur tripedalis et maxima ordinis! Car. flor. et fruct. valdè incompleti si *Baccharis* vera, sola esset species *Gerontogæ*. †.

210. B.? VILLOSA (Vahl symb. 3. p. 98), foliis lanceolatis petiolatis extrorsum serratis subtus ramisque villosis subtomentosis. — in monte Chadrâ Arabiæ fellicis. Ex patriâ et ex invol. squamis lineari-setaceis potius ad *Conyzam*, anadente Persoon, aut ad *Blumeam* referenda.

** *Americana*.

211. B.? ASSURNIS (H. B. et Kunth nov. gen. am. 4. p. 67), arborea, ramulis angulatis tomentoso-pubescentibus, foliis petiolatis oblongis acutis basi rotundatis grossè dentatis membranaceis snprâ glabris nitidis subtus incano-tomentosis, corymbis terminalibus ramosis foliatis, capitulis masculis? circ. 13-floris longè pedicellatis, invol. squamis lanceolatis acutis, recept. paleis paucis inter flores onusto. ♀ in frigidis montanis Quitensium in Paramo de Assuaye. Cor. flavæ ex Bonpl. albidæ ex Kunth. An stylus et ovarium certè abortiunt?

212. B.? BRYOIDES (Pers. ench. 2. p. 425. n. 43), fruticulosa ramosa prostrata, ramulis confertis erectis, foliis minimis confertis linearibus subtus incanis, capitulis 6-8-floris terminalibus solitariis sessilibus, invol. cylindrici squamis oblongis. ♀ ad fretum Magellanicum seu littus Patagonicum legit cl. Commerçon. *Conyza bryoides* Lam. dict. 2. p. 91. Cæt. ign. Genus dubium. †.

213. B.? CANESCENS (Spreng. syst. 3. p. 466), herbacea, foliis fasciculatis petiolatis lineari-lanceolatis acutis subdentatis incano-tomentosis, paniculâ terminali. — in Montevideo. †.

214. B. CORYMBOSA (Pers. ench. 2. p. 425. n. 40), fruticosa, ramis corymbosis, foliis oppositis lanceolatis integerrimis margine revolutis capitulis axillaribus solitariis. ♀ in Peruvîæ alpihus versus Pillao. Molina corymbosa R. et Pav. syst. 210. Ex Spreng. phrasis falsa et species eadem ac B. revoluta. †.

215. B. PENDENS (Pers. ench. 2. p. 424. n. 28), fruticosa scandens, foliis ovatis cordatisque cum acumine integerrimis subquinqüenerviis, capitulis racemosis.

- 5 β in Andium nemoribus ad Cuchero, Casape et Casapillo tractus. Molina dependens R. et Pav. syst. p. 202. Cæt. ign.
216. B. EMARGINATA (Pers. ench. 2. p. 424), fruticosa, foliis ovalibus emarginatis, capitulis subcorymbosis. β in Peruviae collibus copiose in Tarmæ prov. Molina emarginata R. et Pav. syst. p. 202. Cæt. ign.
217. B.? FULIGINÆA (H. B. et Kunth nov. gen. am. 4. p. 68), fruticosa, ramis angulatis pulverulento-pubescentibus, foliis petiolatis oblongis acutis basi rotundatis margine integerrimis et revolutis coriaceis supra glabris nitidis subtus fuligineo-tomentosis, corymbis terminalibus ramosis foliatis, capitulis σ longè pedicellatis 8-floris, invol. squamis oblongis acutis, recept. paleas paucissimas gerente. β in montanis frigidis Quitensium cum B. Assuensi cui simillima.
218. B. GLABRA (Spreng. syst. 3. p. 466), herbacea, foliis oppositis petiolatis oblongis acuminatis profundè serratis membranaceis utrinquè glaberrimis, corymbo terminali. — in Montevideo. \dagger .
219. B. LATIFOLIA (Pers. ench. 2. p. 424. n. 32), fruticosa, foliis lanceolatis triplinerviis argutè serratis, corymbis terminalibus nudis. β in Peruviae ruderalibus et campis prope Muna et Rondosubi vern. Chilca et Anchochuta dicta. Molina latifolia R. et Pav. syst. 208. Cæt. ign. \dagger .
220. B.? QUADRINERVA (Pers. ench. 2. p. 423. n. 6), foliis subovalibus obtusis mucronatis hinc indè serratis quadrinerviis subviscosis. — in Amer. merid. ex herb. Juss. Numerus nervorum par videtur à genere alienus. \dagger .
221. B. APPRESCENS (Spreng. syst. 3. p. 464), fruticosa, foliis confertis spatulato-lanceolatis linearibusque acutis utrinquè mucronato-bidentatis glabris opacis glanduloso-punctatis, racemis foliatis oligocephalis axillaribus. β in Montevideo. \dagger .
222. B. SCABRA (Pers. ench. 2. p. 424. n. 34), ramis scabris, foliis oppositis ovato-lanceolatis acutis trinerviis serratis, corymbis terminalibus. β in Peruv. Tarmæ prov. montibus. Molina scabra R. et Pav. syst. 210. Cæt. ign. \dagger .
223. B.? TOMENTOSA (Pers. ench. 2. p. 424. n. 30 non Thunb.), fruticosa, foliis oblongo-lanceolatis integerrimis tomentosis, capitulis subcorymbosis. β in Peruviae altis frigidis in Caxatambo prov. versus Cheuchin. Molina tomentosa R. et Pav. syst. p. 203. Cæt. ign.
224. B. UNIFLORA (Pers. ench. 2. p. 425. n. 46), fruticosa, foliis lanceolatis serrato-dentatis confertis sessilibus, pedunculis terminalibus squamosis 1-cephalis. β in Peruviae collibus aridis ad Tarma. Molina uniflora R. et Pav. syst. 208. Cæt. ign. \dagger .
225. B.? VENETA (H. B. et Kunth nov. gen. am. 4. p. 68), fruticulosa, ramulis pubescenti-hirtellis, foliis subsessilibus oblongo-lanceolatis acutis basi valdè angustatis apicem versus utroque latere 2-3-dentatis crassiusculis glabris glaucescentibus, corymbis terminalibus simplicibus, capitulis pedicellatis, floribus hermaphroditis, achæniis albo-sericeis. β in declivitate occid. montium prope Cuernavaca Mexicanorum. An fortè Chrysocomea?

N. B. Præter species hinc indè citatas ad Baccharidem referendæ videntur species sequentes sine descriptione in Floræ fluminensis vol. octavo delineatæ, nempe: Chrysocoma cruciata t. 10, Chrysocoma indivisa t. 12, Chrysocoma maritima t. 22, Chrysocoma serrata t. 23, Chrysocoma dentata t. 27, Chrysocoma punctulata t. 49, Cacia frutescens t. 64, Cacia trinervia t. 65, Cacia rosmarinus t. 67, etc.

\dagger Species exclusæ.

B. ABSINTHIODES Hook et Arn. = Tessaria absinthioides.

B. ADNATA H. B. K. = Pluchea subdecurrens.

B. ÆGYPTIACA Forsk. = Pluchea Dioscoridis.

B. ARGUTA Pers. = Psadia balsamica.

- B. *ARUNCA* Schrank. = *Conyza Gouani*.
 B. *ASTEROIDES* Bert. = *Tripolium conspicuum*.
 B. *BANKSIIFOLIA* Bert. = *Tessaria absinthioides*.
 B. *BRAILIENSIS* Linn. = *Vernonia scabra*.
 B. *CHILENSIS* Weinm. = *Euxenia grata*.
 B. *CUPRESSIFORMIS* Pers. = *Lepidophyllum cupressiforme*.
 B. *DIOSCORIDIS* Linn. = *Pluchea Dioscoridis*.
 B. *ERICOIDES* Pers. = *Vernonia ericoides*.
 B. *FORTIDA* Linn. = *Pluchea foetida*.
 B. *FORTIDA* Walt. = *Pluchea Marylandica*.
 B. *HALIMIFOLIA* Mench. = *Conyza ivæfolia*.
 B. *HIERACIFOLIA* Lam. = *Conyza Gouani*.
 B. *INDICA* Linn. = *Pluchea Indica*.
 B. *ILICIFOLIA* Lam. = *Brachylæna dentata*.
 B. *IVÆFOLIA* Linn. = *Conyza ivæfolia*.
 B. *LINEARIFOLIA* Pers. = *Psiadia linearifolia*.
 B. *LYCOPODIOIDES* Pers. = *Eriothrix lycopodioides*.
 B. *MADAGASCARIENSIS* Pers. = *Psiadia Madagascariensis*.
 B. *NERIIFOLIA* Linn. = *Brachylæna neriifolia*.
 B. *NITIDA* Wall. = *Blumea Chinensis*.
 B. *OVALIS* Pers. = *Pluchea ovalis*.
 B. *OVATA* Sieb. = *Pluchea ovalis*.
 B. *PTEROCAULON* Bojer. = *Monenteles Pterocaulon*.
 B. *RUGOSA* Willd. h. = *Vernonia patens*.
 B. *SALVIA* Lour. = *Pluchea balsamifera*.
 B. *SENEGALENSIS* Juss. = *Decaneurum Senegalense*.
 B. *TAIFURCATA* Trev. = *Conyza absinthifolia*.
 B. *VISCOSA* Lam. = *Psiadia integrifolia*.
 B. *VISCOSA* Walt. = *Pluchea bifrons*.

Subtrib. III. *TARCHONANTHÆ* Less. syn. p. 205 non 208.

Capitula aut dioica aut heterogama nunquam radiata, floribus ambitus fœmineis pluriserialibus tenuissimis, disci hermaphroditis aut masculis paucioribus majoribus, antheræ caudatæ. — Folia alterna.

Div. I. *EUTARCHONANTHÆ*. — *Tarchonanthæ* Less. syn. 208 non 205. — *Vernoniæ* *Tarchonanthæ* pleræque Cass. tabl. in op. phyt. 3. p. 76.

Capitula (ut in *Baccharis*) dioica.

CCKVIII. *BRACHYLÆNA* Brown. trans. Linn. soc. 12. p. 115. Less. syn. 208. — *Oligocarpa* Cass. dict. 36. p. 21. — *Tarchonanthi* et *Baccharidis* sp. auct.

Capitula multiflora dioica. Recept. angustum nudum (aut ex Cass. sparse bracteolatum sed bracteolas in specim. meis non vidi). Invol. squamæ imbricatæ siccæ corollis breviores. Corollæ tubulosæ inæqualiter 5-dentatæ. ♂ antheræ basi bisetæ concretæ. Stylus filiformis simplex. Ovarium hispidum abortivum parvè papposum. ♀ antheræ liberæ abortivæ. Stylus apice in ramulos breves latos divisus. Achæmium

glanduloso-pubescent. Pappus setis biserialibus scabridus. — Frutices aut arbusculæ Capenses. Folia alterna coriacea breviter petiolata integra aut dentata suprâ glabra subtus glabra aut tomentosa. Capitula in racemos subpaniculatos disposita. Cor. luteæ.

1. *B. NERIIFOLIA* (Brown. et Less. ll. cc.), foliis lanceolatis integerrimis utrinque glaberrimis latitudine sexies circ. longioribus brevissimè petiolatis, paniculis terminalibus polycephalis. ♀ ad Cap. Bonæ-Spei non infrequens. Burch.! n. 7260, 4350. Drege! Eckl.! et Zeyh. *Baccharis neriifolia* Lin. sp. 1204. Burm.! prod. *Oligocarpa neriifolia* Cass. dict. 36. p. 22. *Tarchonanthus lanceolatus* Willd. Spreng. *Tarchonanthus dentatus* Ecklon! coll. in h. Dunant. *Conyza neriifolia* L'Hér. ex Steud. Burch! cat. geog. n. 7260 ♂ et 4850 ♀. Panicula pube brevissimâ cinnamomeâ junior velutina, demum glabrata. Folia 3 poll. longa, 6-7 lin. lata. (v. s.)
2. *B. GRANDIFOLIA*, foliis petiolatis lanceolatis longè cuneatis apice sinuatis obtusis utrinque glabris latitudine triplò longioribus subtus subferrugineis, paniculæ terminalis ramis divaricatis laxis. ♀ in Africâ Capensi ad Zneenebergen alt. 200-300-ped. legit cl. Drege! Petioli semipollic. Limbus fol. 4-4½ poll. longus, ferè sesquipollic. latus. Invol. *B. neriifoliae* sed ferrugineo-subvelutina. (v. s. ♂.)
3. *B. ELLIPTICA* (Less. syn. 208), foliis ellipticis denticulatis subtus tomentosis breviter petiolatis, racemis axillaribus glabris. ♀ ad Cap. Bonæ-Spei (Burch. n. 3267 et 4450), ad Kliplat rivier et Fisch rivier (Drege!). *Tarchonanthus ellipticus* Thunb. fl. cap. 638. *Tarch. camphoratus* Houtt. in h. Deless. (v. s.)
4. *B. RACEMOSA* (Less. syn. 208), foliis ellipticis oblongisve mucronatis integerrimis aut subdentatis suprâ glaberrimis subtus tomentosis brevissimè petiolatis, racemis ramulos laterales terminantibus 4-6-cephalis, capitulis distinctè pedicellatis. ♀ ad Cap. Bonæ-Spei in sylvâ Kukâ (Th.), Uitenhage (Eckl.! n. 234, 126, 1414, 1326, 8 et 1597). *Tarchonanthus racemosus* Thunb. fl. cap. 638. *Baccharis cuneata* Spreng. in Zeyh.! pl. exs. cap. n. 194 ex h. Moric. huc verisim. referenda? — Burch! cat. n. 4450 ♂ et fortè 3267 ♀. An certè à priori diversa? Folia 7-11 lin. longa, 3-4 lin. lata. Invol. subtrineti squamæ multiseriales, exter. brevissimæ subacutæ, inter. obtusæ dorso puberulæ. (v. s.)
5. *B. DENTATA*, foliis cuneiformibus indivisis aut subtrilobis grossè dentatis et denticulatis suprâ glabris subtus adpressè tomentosis brevissimè petiolatis, ad margines revolutis, racemis ramos terminantibus subspicatis oligocephalis, ♀ ad Cap. Bonæ-Spei in colon. orient. district. Albany (Burchell! n. 3400) ad Zneenebergen (Drege!) et Cafferland (Eckl.). *Tarchonanthus dentatus* Thunb. fl. cap. 638? *Baccharis ilicifolia* Lam. dict. 1. p. 345? Folia pollicem longa, 4-7 lin. lata, inter se formis sat varia sed semper basi cuneata. Invol. squamæ acuminatæ (v. s. ♀.)
6. *♀? salicina*, foliis lanceolatis per totam ferè longitudinem serratis, paniculâ (masculâ) terminali polycephalâ. ♀ in territorio cesso (Eckl.!) ad Omtata et Omsamwubo (Drege! n. 5054 et 3659). Folia basi nunc cuneata nunc obtusa, apice indivisa aut fortè trifida, ad margines vix subrevoluta. An var. sexui masculo propria? An species propr.? (v. s. ♂.)
7. *B. DISCOLOR*, foliis petiolatis oblongo-obovatis obtusis repando-dentatis basi attenuatis suprâ glabris subtus petiolis ramulisque tomentosis, racemis simplicibus subramosisve axillaribus folio vix longioribus oligocephalis, invol. squamis pluriserialibus imbricatis dorso glabris margine araneoso-lanatis. ♀ ad Cap. Bonæ-Spei in Coloniz orient. distr. Uitenhage (Burchell! cat. n. 3751 Eckl.!), ad Omsamculo et Omtendo (Drege!), ad sinum Algoa (Forb.). Folia 3 poll. longa (incl. petiolo 4 lin.), pollicem lata. Achænia matura teretiuscula striata ferè glabra. (v. s.)

CCXIX. TARCHONANTHUS Linn. gen. n. 301. Gärtn. fr. 2. p. 392. t. 166. f. 12. Lam. ill. t. 671. Cass. dict. 52. p. 245. Less. syn. 208 non Vaill. — Tarchonanthi sp. Thunb. Willd.

Capitula sæpius multiflora dioica. Invol. marium squamis 5 uniseriatis ad medium coalitis, foemineorum squamis plurimis distinctis biserialibus. Recept. ebracteolatum hirsutum. Cor. tubuloso-campulata 5-dentata extus hirsuta viscosa, intus glabra. ♂ antheræ in tubum coalitæ exsertæ breviter appendiculatæ et longè caudatæ, caudis setosis, filam. glabris. Ovarium abortivum. Nectarium amplum callosum apice concavum intra corollam ovarium mentiens (1); stylus filiformis vix apice subbilobus. ♀ Stam. abortiva. Nectarium nullum. Stylus exsertus apice bilobus, lobis arcuato-revolutis. Achæmium lanæ hirsutissimum epapposum. — Arbusculæ Capenses. Folia alterna petiolata coriacea, supra reticulato-areolata demum glabra subtus tomentosa aut rarius ad axillas subsolitaria. — Species omnes ab Hottentotis *Mohanka* dicuntur ex Burch. trav.

Sect. I. EUTARCHONANTHUS.

Capitula foeminea 3-5-flora, invol. squamis latè ovatis donata. Achænia hirsutissima. — Folia indivisa.

1. *T. CAMPHORATUS* (Linn. sp. 1179), foliis oblongis obtusiusculis integerrimis aut subdenticulatis, adultis supra glabris reticulato-venosis subtus tomentosis, paniculâ terminali polycephalâ. ♂ ad Cap. Bonæ-Spei frequens Drege! Eckl. Thunb. fl. cap. 637. Berg. cap. 236. — Herm. lugd. b. 229 ic. Pluk. alm. t. 174. f. 1. Lam. ill. t. 671. f. 1. Folia 2-3 poll. longa, 6-10 lin. lata, basi plus minus cuneata. Frutex camphoræ odorem redolens. (v. v. c. ♂ et s. sp. ♀ et ♂.)
2. *T. LITAKUNENSIS*, foliis oblongis integerrimis acutiusculis adultis supra glabris tenuissimè reticulatis areolis impresso-porosis subtus tomentosis, paniculâ terminali polycephalâ. ♂ in Africâ australi extratropicâ inter flumen Makkwarin et urbem Litakun legit cl. Burchell et mecum sp. foem. fructiferum sub n. 2202 cat. geog. comm. Conf. Burch. tr. 2. p. 340. Idem ni fallor ad Omsamculo et Omtendo legit cl. Drege! *T. camphorato* similis sed folia 15 lin. longa, 3 lata et diversè reticulata. Paniculæ rami et ramuli multò tenuiores, albedo nec rufescenti-velutini. Specim. masc. mihi ign. (v. s.)
3. *T. OBOVATUS*, foliis obovatis obtusis basi cuneatis supra glabris reticulato-venosis subtus tomentosis, capitulis ad axillas foliorum solitariis pedicellatis. ♂ ad Cap. Bonæ-Spei in regione Trans-Gariepinâ propè Sensavan. Folia 8 lin. longa, 3-4 lin. lata. Pedicelli cum capitulo circ. folii longitudine. Capitula fructifera priori similia sed ad axillas solitaria. (v. s. comm. ♀ à cl. Burchell sub n. 2155.)
4. *T. ANGUSTISSIMUS*, foliis oblongo-linearibus integerrimis supra glabris vix reticulatis subtus tomentosis, capitulis solitariis. ♂ ad Cap. Bonæ-Spei in regionibus interioribus propè Karra-Poort (Burchell! n. 2691), ad Zneurebergen (Drege!), in Albany et Tambukiland (Eckl.!). Folia 7-8 lin. longa, 1 ½-2 lin. lata. Capitula parca et nimis juniora ut structuram discernere possim et ideo mihi ut jam cl. inventori genus omninò dubium. (v. s.)

N. B. T. minor Less. syn. 208 (de quo dicuntur: Folia breviora, capitulorum masculorum squamæ spatulato-obovata acutæ foliaceæ impellucidæ extus densè

(1) Quâ insolitâ nectarii grossificatione decepti fuerunt Bergins et Linnæus. Ipse olim hunc errorem, sed dubitans et theoriam veram simul indicans, admisi in disc. compos. ann. Mus. h. nat. Par. 16. p. 138.

tomentosæ et circiter sesquilineam longæ), fortè ad quandam ex duabus ultimis speciebus referendus, sed cùm capit. fœminea sola viderim, mihi obscurus remanet.

Sect. II. HENOTOGYNA.

Capitula fœminea uniflora, invol. squamis linearibus donata. Achænia parvè hirsuta. — Folia apice grossè trifida. — An genus?

5. *T. trilobus*, foliis cuneatis margine integris aut subsinuatis apice obtusè trifidis, lobo medio minore, suprâ glabris minutissimè reticulatis, subtus hirsutis subtomentosis penninerviis reticulatisque, racemis in paniculam subdispositis. \S in asperis Africæ austr.-orient ad Omtendo legit cl. Drege! Species distinctissima, ferè inter *Brachylænas* et *Tarchonanthos* veros ex habitu media. Folia rugosa petiolata 3-4 pollic. longa, $1-1\frac{1}{2}$ lata, lobis exceptis *Budleia* aut *Salvia* folia referentia. Flor. masculi omninò *Eutarchonanthi*, fœminei 1-flori; achænia epapposa ovata per anthesin saltem parvè hirsuta. (v. s.)

+ *Species exclusæ.*

T. DENTATUS Thunb. = *Brachylæna dentata*.

T. ELLIPTICUS Thunb. = *Brachylæna serrulata*.

T. ERICOIDES Linn. f. = *Eriocephalus glaber*.

T. GLABER Linn. f. = *Brachylæna dentata*.

T. LANCEOLATUS Thunb. = *Brachylæna neriifolia*.

T. RACEMOSUS Thunb. = *Brachylæna racemosa*.

Div. II. PLUCHEINÆ Cass. tabl. sub *Vernoniaceis* (1). — *Epaltidæ* Less. syn. 205.

Capitula heterogama seu monoica.

CCXX. *BLUMEA* DC. in ann. bot. Guill. 1833. v. 2. p. 514 (2). non Spreng. nec Nees (3). — *Erigeron* Don prod. fl. nep. p. 171. non Linn. — *Conyzæ* species pleræque Blum. bijdr. Wall. cat.

Capitulum multiflorum heterogamum, fl. radii multiserialibus fœmineis tenuissimè filiformibus truncatis aut sub-2-3-dentatis, disci paucis (5-25) masculis cylindricis 5-dentatis, fauce non aut vix dilatata. Recept. planum sæpius omninò nudum interdum tenuissimè fimbriiferum hirsutum. Invol. pauciseriale subimbricatum, squamis linearibus acuminatis. Antheræ basi tenuissimè caudatæ. Achæmium teretiusculum. Pappus 1-serialis, setis capillaribus vix subscabris. — Herbæ aut suffrutices. Species *Gerontogæ*, ferè omnes ex Indiâ orientali seu insulari seu præsertim continentali, paucissimæ Africanæ. Folia alterna sæpius cum caulibus variè villosa. Capitula paniculata aut laxè subcorymbosa. Flores flavi aut purpurascens. — Genus vastissimum à *Conyzæ* et

(1) Stylus enim florum disci non malè refert stylos *Vernoniacearum*, sed affinitates potius juxta *Conyzeas* hæc genera trahunt.

(2) Caractere emendato ex observationibus per litteras à cl. Decaisne communicatis.

(3) *Blumia* Nees est *Talauma*. *Blumia* Spreng. est *Reinwardtia* Blum., nomen evidentissimè à Sprengelio injustè mutatum et sanctè servandum cum *Reinwardtia* Spreng. sit. *Dufourea* Kunth. = *Prevostea* Choisy. = *Calycobolus* Willd., et *Reinwardtia* Dumort. = *Lini* species.

Phagnalo achæniis teretiusculis nec plano-compressis, ab Erigeronte ligulis nullis et achæniis teretiusculis, à Laennecii pappo 1-seriali distinctum, à Pluchæ tandem corollis masculis cylindricis, invol. squamis linearibus vix imbricatis, et inflorescentiâ plus minus paniculatâ diversum, dicavi cl. C. L. Blume qui plures hujus species ipse detexit et qui Botanicam Indicam strenuè promovit.

Ser. I. *APTERÆ, nempè foliis caulinis non decurrentibus.*

* Dissitifloræ, nempè foliis integris dentatisve, pedunculis solitariis monocephalis.

1. *B. AMPLECTENS* (DC. in Wight contrib. bot. ind. p. 13.), caule herbaceo piloso tereti divaricato-ramoso, foliis junioribus villosis adultis subglabratiss, foliis caulinis ovatis semiamplexicaulibus acutè dentatis, infimis hinc inde basi incis, pedunculis paucis subterminalibus folio paulò longioribus patentim villosis 1-cephalis, invol. squamis linearibus aristatis disco paulò longioribus. (1) in rudertis Indiæ Maderaspatanæ frequens, circa Negapatam (Wight!). *Erigeron obliquum* Linn. mant. 572? ex Russ. herb. et ideo *Conyza obliqua* Willd. sp. 3. p. 1930? *C. amplexicaulis* Lam. dict. 2. p. 85. Willd. l. c.? Folia var. subrotunda obtusa, ovata, ovali-oblonga acuta, plus minus serrata, semper semiamplexicaulia ut dicitur de *C. amplexicauli* et ex sicco obliqua ut de *C. obliqua*. Flor. fœm. pluriser. tenues stig. exsertis rubris, masc. circ. 15-20 anth. exsertis. *Conyza pumila* Pers. ench. 2. p. 427, huc ut varietas citata est fortè diversa sed vix nota. (v. s. in h. Cur. Ind. or. nunc soc. Linn. Lond. et in h. Wight.)
2. *B. ARENARIA* (DC. in Wight contrib. bot. ind. p. 13), basi glabruscula apice undiquè sericeo-villosissima, caule à basi ramoso diffuso tereti, foliis oblongis sessilibus mucrone callosio acutis integerrimis aut acutè denticulatis, pedunculis solitariis 1-cephalis hirsuto-lanatis axillaribus terminalibusque, invol. squamis linearibus acuminatis dorso hirsutis apice ciliatis, corollis masc. apice barbatis. (1) in arenosis Indiæ Maderaspatanæ propè Negapatam (Wight!). *Conyza villosa* herb. Mair. ex Wall. cat. n. 3105 non 3083 nec Willd. Radix longa gracilis. Flor. fœm. tenues stylo longè exserto bifido, masc. pauci. Rec. nudum. (v. s. in h. Wight et in h. hon. Cur. Ind. or. nunc Soc. Linn. Lond.)
3. *B. INTEGRIFOLIA*, herbacea erecta ramosissima infernè pilis confertis subglutinosus subhirtella, supernè pilis longioribus sparsis pilosula, foliis linearilanceolatis acutis integerrimis, summis linearibus, ramulis elongatis apice 1-cephalis nudis, invol. squamis dorso ad basin pilosiusculis cæterum glabrusculis linearibus demum reflexis. (1) in sylvis aridis Novæ-Hollandiæ bor.-occid. ad Port-Kearls lat. 14° legit cl. A. Cunningham septemb. flor. Herba semipedalis. Radix gracilis simplex. Flores flavi, masc. in specim. incompleto destructi et ideo genus subdubium. (v. s. comm. à cl. Cunn.)
4. *B. TENELLA* (DC. in Wight contrib. bot. ind. p. 13), caule herbaceo multiplici puberulo tereti parçè ramoso, foliis oblongis basi obtusè cordato-semiamplexicaulibus acutis distantèr acutè et callosio-dentatis pube parvâ scabridis, ramis acuminatis subintegerrimis, pedunculis terminalibus et oppositifoliis 1-cephalis puberis folio triplo longioribus, invol. squamis linearibus acuminatis disco ferè sublongioribus. (1) in insulâ Timor leger. Botanici expedit. Bandinæ, et in arenosis versus Negapatam (cl. Wight!). Decaisn. nonv. ann. Mus. h. Par. n. 3. p. 411. Fl. flavi, fœm. innumeris tenuissimis masc. 8-9, anth. inclusis. Valde affinis *B. oligocephalæ* et *B. amplexenti*. (v. s. comm. à Mus. reg. Par.)
5. *B. ANAGALLIDIFOLIA* (DC. l. c. p. 13), caulibus à collo multiplicibus diffusis ramosis teretibus apice puberulis, foliis caulinis ovali-oblongis sessilibus hinc inde acutè dentatis, rameis lanceolatis integerrimis puberulis, capitulis longè pedicellatis solitariis axillaribus terminalibus in paniculam foliosam laxam subcorymbosam dispositis, invol. squamis linearibus acutis glabrusculis disco paulò longioribus. (1) in Indiâ or. Maderaspatanâ ad Negapatam in humidis

frequens (Wight!), in Pegu (Belang!). *Conyza anagallidifolia* Wall. f. n. 3098. comp. n. 208. Recept. nudum punctulatum. Fl. fœm. innumeri tenuissimæ masc. 10-12 cor. apice subbarbata. *Achæmium glabrum*. An fortè *Conyza scabra* Linn. mant. 113? quæ vix nota!

6. *B. GLUTINOSA*, herba erecta paniculato-ramosa parçè pilosa tota glutine viscosa, ramis teretibus, foliis oblongis grossè et acutè dentatis basi attenuatis, capitulis pedicellatis subsolitariis, invol. squamis linearibus glabrisculis, floribus masculis circ. 3. — in aridis insulæ Madagascar circa pagum Majangay in sinu Bombatoe legit cl. Rojer. Cor. glabræ, fœminæ innumeræ. *Achænia glabra*. (v. s. comm. à cl. inv.)

7. *B. OLIGOCEPHALA* (DC. in Wight contr. bot. ind. p. 13), parçè puberula herbacea multicaulis subdiffusa, foliis infer. obovato-oblongis basi cuneatis, super. oblongis ovalibus sessilibus subsemiauriculatis omnibus apice remotè et acutè dentatis, pedunculus 1-3 ex apice caulium 1-cephalis villosis folio longioribus, invol. squamis linearibus acutis disco æquantibus. ① circa Madras. C. auriculatis Wight! in Wall. cat. n. 3104. comp. 214 non Lam. nec Michx. Herba vix semipedalis, affinis *B. bifoliatæ*, sed præter car. indic. fl. masc. 15-20, antheris exsertis. (v. s. in h. soc. Linn. Lond. et comm. à cl. Wight.)

8. *B. RUBIFLORA*, caule herbaceo erecto ramoso pilis mollibus patulis plus minus hispidis foliis sagittato-semiauriculatis ovali-oblongis acutis subintegerrimis aut acutè dentatis, ramulis floridis 1-cephalis subunifoliis pedicelliformibus, invol. squamis linearibus acuminatis flores æquantibus, cor. disci apice hirsutis. — in Indiâ Maderaspatanâ. *Erigeron asteroides* Roxb. ex Wall. cat. et herb. n. 2975. comp. n. 85. Styli fl. fœm. longè exserti! Variat ramis fere glabris aut molliter apice villosissimis foliorum dentibus nullis aut utrinquè 7-8. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

9. *B. BIFOLIATA* (DC. l. c. p. 14 sub *Conyzâ*), caule herbaceo adscendente tereti ramoso apice præsertim puberulo, foliis acutè dentatis, infer. ovatis basi longè attenuatis glabrisculis, super. sessilibus oblongis cuneatis puberis, pedunculis axillaribus supra basin sæpius folia duo subopposita gerentibus, pedicellis elongatis 1-cephalis, invol. squamis linearibus acutis disco subæqualibus. ① in Indiâ or. peninsulâ frequens in nemorosis humidis ex Wight! C. foliolosa Wight! in Wall. cat. n. 3092. comp. 202 non n. 3030 nec Willd. C. bracteolata Rottl. in h. Puer. Pluk. alm. t. 177. f. 1. Habitus varius et fortè spec. 2-3 sub hac confusa. In specim. diversis eodem loco collectis video folia nunc longè nunc breviter petiolata, super. et involucria plus minus villosa, fl. masc. indiv. 8-16, etc., in omnibus recept. nudum Cor. masc. flavidæ vix ac ne vix apice barbellatæ antheris subinclusis, achænia glabra. (v. s.)

** *Paniculatæ, nempe foliis dentatis laciniatisve, pedunculis nunc axillaribus polycephalis nunc verò in paniculam dispositis, invol. cinereo-pubescentibus.*

10. *B. BARBATA* (DC. l. c. p. 14. sub *Conyzâ*. Wall. cat. n. 3099. comp. 209), caule herbaceo erecto simplici tereti pilosiusculo, foliis lanceolatis basi attenuatis acutè et distanter serratis acutis supra sparsè puberis subtus densè et adpressè sericeo-villosis, ramis axillaribus foliosis folio brevioribus apice racemosis, summis corymbosis, invol. squamis linearibus acutis glabrisculis disci longitudine. ① in Indiâ peninsulâ legit cl. Wight. Herba pedalis. Folia 1 $\frac{1}{2}$ -2 poll. longa, 5-6 lin. lata. Recept. nudum. Fl. flavidi, fœm. innumeri stylo exserto bifido, masc. 5 cor. apice glabriscula. Achæn. glabra. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et in herb. Wight.)

11. *B. CHAMISSONIANA*, caule herbaceo erecto tereti apice pilis mollibus patulis hispidis, foliis obovatis obtusis basi cuneatis subsessilibus denticulatis utrinquè molliter pubescenti-villosis pedunculis axillaribus folio brevioribus oligocephalis, invol. squamis linearibus disco æqualibus pubescentibus. ① in Læsoniâ legit cl. de Chamisso. C. bifoliata Cham! et Less. in linnæa 1831. p. 135. (v. s. comm. à cl. Chamisso.)

12. *B. AXILLARIS*, caule tereti erecto herbaceo pubescenti-villoso, foliis petiolatis

- obovatis ovalibusve membranaceis argutè denticulatis penninerviis, pedunculis folio brevioribus foliola 1-2 gerentibus 7-8-cephalis, ultimis in paniculam dispositis, invol. squamis dorso hirsutis linearibus acuminatissimis. (1) in insulâ Mauritii ad ripas rivorum (Bout. Bojer!). *Conyza axillaris* Lam. dict. 2. p. 84. Petioli pollicares. Limbi fol. 2 $\frac{1}{2}$ poll. longi, 15-18 lin. lati basi subcuneati. Pappus albidus. Cor. purpurascens. Affinis *B. Wightianæ* et *Chamissonianæ*, sed foliis petiolatis ab utraq. faciliè differt. *C. axillaris* Willd. enum. 871. Horn. cat. suppl. 12, est certè diversa sed mihi ignota. (v. s.)
13. *B. MANILLENSIS*, caule herbaceo erecto tereti striato hirsuto suprâ parèe folioso, foliis semiamplexicaulis sparsè hirsutis obovatis sinuato-dentatis subtus pallidioribus, racemo terminali simplici, capitulis jun. subsessilibus, invol. squamis linearibus ciliatis squarrois. (1) in Luzoniâ legit el. de Chamisso. *Conyza Manillensis* Less. in linnæa 1831. p. 135. Ovaria hirsutiuscula. Affinis dicitur *B. Wightianæ* var. γ .
14. *B. WIGHTIANA* (DC. l. c. p. 14), caule herbaceo erecto tereti striato puberulo, foliis inf. petiolatis ovalibus basi cuneatis acutè et inæqualiter dentatis subobtusis, super. cuneatis ovali-oblongis, capitulis in paniculam racemosam interruptam terminalem digestis, invol. squamis linearibus acutis inter. glabrisculis disco paulò longioribus demum reflexis, cor. masculis extis tuberculoso-glandulosis. (1) in Indiâ Maderaspatanâ legit cl. Wight, ad Pondichéry (Belanger!), et in regione Birmanorum circa Prome (cl. Wallich). *Conyza Wightiana* Wall. cat. n. 3063. comp. 203. Recept. nudum. Fl. sem. innumeri tenues, masc. 8-12 ex sicco subrubentes. Achenia glabra. (v. s. comm. ab hon. Cur. Ind. or.)
- f. cuneiformis*, foliis ferè omnibus oblongo-cuneatis subacutis, capitulis thyrsoidè-confertis. In Indiâ Maderaspatanâ. *C. bifoliolata* Wall. cat. n. 3091. comp. 201 non Linn. (v. s. comm. cum *a*.)
- f. Madagascariensis*, floribus masc. 5-6. — in agris cultis ins. Madagascar legit cl. Bojer. Similis var. *a*. (v. s.)
15. *B. CUNNINGHAMI*, caule herbaceo erecto tereti paniculato-ramoso basi folisque glabris apice molliter villosis, foliis sessilibus oblongo-lanceolatis grossè et irregulariter inciso-dentatis lobulis dentibusve acutis rigidulis, capitulis paniculatis, invol. squamis linearibus acuminatis dorso puberis apice ciliato-barbatis demum reflexis. — in umbrosis ripis abruptis Novæ-Hollandiæ bor.-occid. ad Regent's-River latit. 15° legit cl. A. Cunningham. Herba circ. bipedalis. Flores flavidi, masc. pauci. (v. s. comm. à cl. inv.)
16. *B. MUARA*, tota cinerea molliter villosa, caule herbaceo erecto tereti ramoso, foliis ovali-oblongis aristato-serratis basi attenuatis, inf. petiolatis, infernè lyrato-appendiculatis lobatis angustis acutis nunc à limbo distinctis nunc cum eo confluentibus, paniculæ ramis erectis apice confertè corymbosis, invol. squamis linearibus vix acutis villosis disci longitud. cor. masc. apice barbatis. — in Indiâ orient. prope Hologung legit cl. Hamilton. *Conyza Muara* Ham! in Wall. cat. n. 3095. Rec. nudum. Fl. flavidi, masc. 20-25. Achenia glabra. (v. s. in h. hon. Cur. Ind. or. nunc soc. Lin. Lond.)
17. *B. LESCHENAUULTIANA*, caule erecto herbaceo tereti piloso, foliis ovalibus serratis puberulis, inf. obtusis basi attenuatis, sup. sessilibus acutis, ramulis floridis hirsutissimis oligocephalis infer. axillaribus folio brevioribus, super. in paniculam brevem digestis, invol. squamis linearibus hirsutis discum æquantibus. — in Zeylonâ et in Indiâ peninsulâ legit cl. Leschenault. Rec. subalveolare nudum. Fl. sem. innumeri stylo exserto bifido purpureo, masc. 10-11 glabri anth. semiexsertis. Achenia glabra. Pappus paucisetus albus. Valdè affinis *B. molli*. (v. s. comm. à cl. Leschenault.)
18. *B. LACTUCIFOLIA*, infernè sparsè pilis rigidulis scabra, supernè confertius et molliùs velutina, caule herbaceo erecto tereti apice ramoso, foliis infer. ovatis acuminatis grossè dentatis in petiolum longè attenuatis, mediis ad basin auriculato-cordatis, summis sessilibus oblongo-lanceolatis, paniculâ laxè patente,

ramis spicato-racemosis, capitulis pedicellatis dissitis, invol. squamis linearibus acutis disci longitudine. (1) in Ind. or. regione Birmannorum ad fl. Saluen supra Troglā legit cl. Wallich et nī fallor ad Sornogon cl. Hanke. *Conyza lactuæfolia* Wall.! cat. n. 3088. comp. n. 198. Fl. fœm. innumeri, stylo exserto bifido. masc. circ. 25-floris, lobis ciliato-hirtellis. (v. a. comm. ab hon. Cur. merc. angl. Ind. or.)

19. *B. LACERA* (DC. in Wight contr. p. 14), tota tomentosa aut pubescenti-hirsuta albida, caule herbaceo erecto tereti, foliis obovato-oblongis grossè dentatis sæpè basi incisis aut pinnatifidis, paniculā oblongā aut corymbosā confertā aut laxā, invol. squamis linearibus acuminatis flores æquantibus. — ad oryzetā Ind. or. continent. et insularis nec non in Chinā. Flor. flavi, fœm. innumeri, stylo bifido, masc. circ. 20. *Achænia* subangulata glabriuscula. Planta multiformis fortè dividenda, sed caract. non satis perspicui.

a. *Burmanni*, foliis lyrato-pinnatifidis, ramis paniculatis. — in Javā. C. lacera Burm. fl. ind. 180. t. 69. f. 1. Blum.! bijdr 897. (v. s.)

β. *Commersonii*, foliis obovatis grossè dentatis, capitulis confertis. — in Javā. Con. lacera Lam.! dict. 2. p. 83. (v. s.)

γ. *Wallichii*, foliis obovatis profundè et acutè dentatis, paniculā laxā. — in Silhet, Chinā, etc. Con. lacera Wall.! cat. et herb. n. 3045 et 3082. comp. n. 155 et 192. C. villosa Wall. ll. cc. n. 3083 et 193, ab hac specie non differre videtur. (v. s.)

δ. *Blumiei*, foliis obovatis acutè et exsertè serratis, paniculā laxā. — in Javā. *Conyza Javanica* Blum.! bijdr 897. (v. s.)

20. *B. LACINIATA*, caule herbaceo erecto ramoso tereti puberulo apice molliter et parcè piloso, foliis infer. pinnatipartitis utrinquè subscabris lobis utrinquè 4 oblongis ovalibus summis cum impari subconfluentibus, omnibus subrepandis acutè et inæqualiter dentatis, summis oblongis sessilibus hinc indè serratis, paniculā amplā polycephalā, ramis elongatis apice thyrsoides, capitulis pedicellatis, invol. squamis linearibus acuminatis disco longioribus. — In Indiā orient. *Conyza laciniata* Wall. cat. n. 3100. comp. n. 210. Recept. nudum. Fl. masc. 15-16 lobis apicè puberulis. *Achænia* pilis sparsis scabrida oblongo-teretia. (v. s. in herb. hon. Cur. Ind. or. nunc Soc. Lin. Lond.)

21. *B. CERNUA*, caule herbaceo erecto tereti subsimplici basi puberulo apice velutino-villoso, foliis obovatis basi attenuatis grossè et acutè dentatis suprā puberulis subtus villosis, infer. longè basi productis, summis oblongis utrinquè villosis, paniculā elongatæ ramis folio longioribus apice capitula glomerata subsessilia cernua gerentibus, invol. squamis linearibus acutis flores æquantibus. — in Indiā orientali ad Bhiratghur februario flor. legit cl. Hamilton. *Conyza nutans* Hamilt.! in Wall.! cat. n. 3080. comp. n. 190. non Blum. Folia infer. 6 poll. longa, 1 $\frac{1}{2}$ lata. Fl. fœm. innum. tennes, stylo exserto bifido, masc. circ. 7-8. Pappus albus. *Achænia* puberula. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

22. *B. ? LEUCANTHEMA*, cinereo-pubescent herbacea, foliis elliptico-oblongis acuminatis sessilibus denticulatis, corymbis compositis patulis polycephalis in paniculam amplam digestis, invol. tomentosi squamis lineari-lanceolatis aristatis discum subæquantibus. 2/ ad Bessaria pagum Nepaliæ. *Erigeron leucanthum* Don prod. fl. nep. p. 171. Folia inf. spithamea, 2 poll. lata. Fl. masc. paucissimi.

23. *B. TRICHOPIORA*, tota villis sericeis confertis hirsuto-villosa, caule herbaceo stricto erecto parcè ramoso, foliis obovatis acutè dentatis, infimis breviter petiolatis obtusis, caulinis basi cuneatis, ramulis floridis folio longioribus in paniculam angustam elongatam dispositis, invol. squamis linearibus disco longioribus. (1) in Nepaliā legit cl. Wallich. *Erigeron molle* Don prod. fl. nep. p. 172. sed. div. à *Conyza* molli Willd. An fortè *Bl. Wightianæ* var.? Recept. nudum Fl. purpurei fœm. innum. stylo exserto bifido, masc. 10-12 glabri. (v. s. comm. à cl. Wallich anno 1821.)

24. *B. DREGHANA*, caule herbaceo erecto apice præsertim molliter villosa, foliis

sessilibus obovatis dentatis, infer. basi longè attenuatis in parte attenuatà acutè inciso-dentatis suprâ pubescentibus subtus villosis, super. utrinquè adpressè sericeo-villosis, paniculâ contractâ sericeo-villosâ, invol. squamis linearibus disci longitudine. — in Africâ austr.-orient. ad Zw. Omata et Omsamwubo legit cl. Drage n. 5055. Habitu ferè *B. trichophoræ*. Invol. pubens post anthesin reflexum. Squamæ et styli purpurascens. Recept. punctulatum. Fl. masc. 8-10. (v. s.)

ß. gracilis, caule gracili simplici pubescente, foliis infer. basi integerimis. — cum var. *α* legit cl. Drage. (v. s.)

25. *B. PARVIFOLIA*, tota et præsertim ad apicem cinerea pubescenti-villosa, caulibus è collo plurimis erectis ramosis, foliis infer. obovatis basi attenuatis breviter petiolatis acutè dentatis, superioribus oblongis sessilibus vix dentatis, paniculâ ramosâ polycephalâ, invol. squamis exter. brevibus dorso subhirsutis inter. linearibus acuminatis flores subæquantibus demum reflexis. ① in Nepaliâ legit cl. Wallich. *Conyza parvifolia* Wall! cat. n. 3004. comp. n. 114. *Erigeron glomeratum* Spreng. in litt. ad Wall. Radix dura subsimplex. Caulis 8-10 poll. Folia pollicem longa, 6 lin. lata. Cor. purpureâ, fœm. innumeræ, stylo bifido, masc. 8-10. An *Bl. oxyodontæ* var. ? (v. s. comm. ab hon. Car. merc. angl. Ind. or.)

26. *B. TIMORENSIS* (DC. in Decaisne! nouv. ann. Mus. h. n. Par. 3. p. 411), caule erecto tereti apice ramoso puberulo basi subglabrato, foliis oblongis acutis puberulis irregulariter serratis, infimis basi attenuatis, mediis basi incisis auriculâ ntrinquè exsertâ sublineari semiamplexicanlibus, summis sessilibus subintegris, paniculâ corymbosâ ramis erectis, capitulis pedicellatis, invol. squamis linearibus acuminatis dorso puberis discum ferè superantibus. — in insulâ Timor leg. Baudiniani collectores. Recept. nudum. Cor. rubentes, fœm. innumeræ, masc. paucae subpubescentes sub oculo armato. Affinis *B. lactucæfolia*. (v. s. comm. à Mus. reg. Par.)

27. *B. TRIGONA*, tota glanduloso-pubera, caule herbaceo erecto infernè trigono superne tereti parçè ramosa, foliis petiolatis ovalibus acutis irreg. dentatis basi inciso-lyratis, summis subintegris acuminatis, paniculâ subracemosâ foliosâ strictâ, invol. squamis linearibus acutis glabrisculis, recept. glabro, achæniis teretiusculis glabris. — in Indiâ or. prov. boreali-occid. legit cl. Royle. Valde accedit ad *Bl. Timorensem* sed diversa videtur. Herba in siceo nigricans. An caulis casu tantum trigonus? Fl. fœm. innum., masc. circ. 5 breves. (v. s. comm. à cl. Royle.)

28. *B. CICHORIIFOLIA* (DC. in Decaisne! nouv. ann. Mus. h. n. Par. 3. p. 410), caule herbaceo erecto ramoso tereti striato puberulo in ramis floridis villos glandulis intermixtos gerentibus, foliis glabriusculis, inf. lyrato-pinnatilobatis infernè attenuatis lobis inf. parvis oblongis super. majoribus subtriangularibus confluentibus, omnibus irreg. mucronato-dentatis, fol. summis oblongis sessilibus subintegris, paniculâ amplâ divaricatâ, ramis apice racemiferis, invol. squamis linearibus ciliatis disco subæqualibus. — in insulâ Timor colleg. exp. Baudiniane collect. Recept. alveolare fimbriiflereo-velutinum. Fl. flavidis, fœm. innum. masc. 8-10 glabri. (v. s. comm. à Mus. reg. Par.)

29. *B. SÆSSILIFLORA* (Decaisne nouv. ann. Mus. h. n. Par. 3. p. 410), ramis herbaceis erectis ramulisque virgatis striatis puberulis, foliis subruncinatis v. lanceolato-oblongis irregulariter serratis sessilibus subauriculatis puberulis, capitulis axillaribus plerumque sessilibus 3-5-glomeratis, invol. squamis linearibus puberulis, recept. glabro subtuberculato. — in insulâ Timor. Proximè ad *B. cichoriifoliam* accedit.

30. *B. CREPIDIFOLIA*, tota tenuiter pubescenti-villosa subcinerea caule erecto striato-sulcato subramoso, foliis infer. basi attenuatis lyratis pinnatipartitisve, lobis inf. ovali-oblongis, summo majore obovato, foliis omnibus acutè et irregulariter dentatis, summis oblongis acuminatis, ramis apice capitula dense thyrsoidæa gerentibus in paniculam dispositis, invol. squamis linearibus acutis dorso pubescentibus disco paulò longioribus. — in Indiâ or. ad Nakh-Wari

februar. florentem legit cl. Hamilton. *C. laciniata* Ham. ! in Wall. cat. n. 3084. non Roxb. Recept. densè alveolare. Flores fœm. tennes glabri, stylo breviter exserto, masc. circ. 18-20 lobis ciliato-villosis. Achænia subpubescentia. (v. s. in h. Cur. Ind. or. nunc Sec. Lin. Lond.)

31. *B. GLANDULOSA* (DC. in Wight contrib. p. 14), tota brevissimè piloso-glandulosa, caule herbaceo erecto subtereti striato-nervoso, foliis caulinis breviter petiolatis ovatis subacuminatis irregulariter dentatis basi lyrato-incisis, ramis basi cuneatis obovato-oblongis subdentatis, capitulis ad apices ramorum paniculam foliosam constituentibus, pedicellis ramosis glandulosis folio brevioribus, invol. squamis linearibus acutis discum æquantibus. ① ? in Indiâ Madraspat. propè Negapatam legit cl. Wight. Herba supernè verisim. viscosa. Recept. convexum. Achænia glabra. (v. s. in h. Wight.)
32. *B. PHYLLOSTACHYA*, caule herbaceo erecto tereti basi glabriusculo apice cum ramis petiolisque pilis mollibus patulis hispido, foliis petiolatis lyratis pubescentibus, lobis infer. paucis parvis oblongo-linearibus acutis aut tridentatis, terminali maximo ovato-obtusulo basi sublobato acutè dentato, paniculâ confertâ, ramis brevibus folia obovata, ovaliave acutè dentata capitulis intermixta gerentibus, invol. squamis linearibus acutis glabris disco longioribus. — in Ind. orient. verisim. Madraspatanâ. *C. paniculata* herb. Madr. ex Wall. ! cat. n. 3090. comp. n. 200 non Willd. Flor. fœm. innumeri, masc. 5. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
33. *B. CINERASCENS*, tota cinereo-pubens, pube alterâ brevi confertâ et pilis longis mollibus patulis intermixtis apice confertioribus, caule herbaceo erecto tereti ramoso, foliis ovali-oblongis lanceolatisve, infer. irregulariter dentatis petiolatis, mediis basi attenuatis, summis integerrimis, omnibus exauriculatis, paniculâ ramosâ polycephalâ, ramis elongatis basi interruptè spicatis apice congesto-corymbosis, invol. squamis linearibus vix acutis dorso villosis disco longioribus. ① in Nepaliâ legit cl. Wallich. *C. runcinata* var. B. Wall. ! cat. n. 3087. comp. 197. Affinis *Bl. runcinata* ex florum structurâ et numero sed foliis pilisque certè distincta. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
34. *B. RUNCINATA*, caule herbaceo erecto tereti ramoso pubescenti-velutino, foliis inf. in petiolum basi auriculato-dilatatum productis oblongis runcinatis irreg. dentatis suprâ puberulis subtus et novellis villosis, paniculâ elongatâ, ramis capitula congesta interruptè spicatis apice gerentibus, invol. squamis linearibus acutis dorso villosis disco longioribus. — in Ind. or. ad Hurdwar prov. Delhi legit cl. Wallich. *Conyza runcinata* Wall. ! cat. n. 3087. A. comp. n. 197. A. Recept. alveolatum. Fl. fœm. tenues, stylo vix exserto, masc. 12-15. Achænia subpuberula. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
35. *B. ACUTATA* (DC. in Decaisn. ! nouv. ann. Mus. h. n. Par. 3. p. 409), tota (supernè imprimis) tenuiter velutina pubescens, caule herbaceo erecto simplici nervulis striato-angulato, foliis petiolatis pinnatis irregulariter paucilobatis, lobis oblongis acuminatis terminali majore ovali, omnibus serratis profundis acutis inæqualibus incisis, summis oblongis basi attenuatis subcæcatis, paniculâ contractâ terminali, invol. squamis linearibus pubescentibus acutis disco ferè longioribus. ① ? in insulâ Timor leger. Baudiniani peregrinatores. Recept. alveolare subhirsutum. Affinis *C. crepidifolia*. (v. s. comm. à Mus. reg. Par.)
- β. foliis infimis lyrato-lobatis, cæteris ovatis aut ovato-lanceolatis grossè et acutè dentatis. — in Timor cum priore. (v. s. comm. à Mus. reg. Par.)
36. *B. SONCHIFOLIA*, tota pubescenti-villosa subviscida, caule erecto simplici striato, foliis infer. petiolatis lyratis, lobis inf. paucis oblongis acutis, terminali maximo triangulari, omnibus grossè et inæqualiter dentatis, super. incisis dentatisve, paniculâ terminali elongatâ, pedunculis later. folio duplò longioribus invol. squamis linearibus acutis disco paulò longioribus. ① in Indiâ orient. ad Patna legit cl. Hamilton. *Conyza sonchifolia* Ham. ! in Wall. ! cat. n. 3085. comp. n. 195. Affinis *B. laciniata* sed recept. laeve punctatum,

- flor. fœm. stylo longius exserto prof. bifido, masc. 3-4 oor. brevius ciliatâ. Achænium hirtellum striatum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
37. B. SUBCAPITATA, tota cinereo-pubens apice et in surculis novellâ magis villosa, caule herbaceo erecto tereti simplici unico aut ad basin submultiplici, foliis petiolatis ovatis inæqualiter et acutè dentatis ad basin incisâ fere lyratis, capitulis ad apicem caulis in spicam ovato-oblongam congestis, invol. squamis linearibus acutis florum longitudine. ① in Nepaliâ legit cl. Wallich! Conyza capitata Wall. cat. n. 3056. comp. n. 166. Herba vix pedalis. Folia sesquipoll. longa 8-9 lin. lata. Flores pallidè flaventes, fœm. innumeri graciles stylis exsertis bifidis, masc. circ. 12 antheris non exsertis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
38. B. HAMILTONI, caulis pluribus solitariisve herbaceis erectis simplicibus lanuginosis parè foliosis, foliis in parte caulium infer. approximatis sessilibus oblongis cartilagineo-dentatis suprâ glabriusculis subulatis vix acutis, super. parvis ramosis, capitulis in apicem terminalem ovatam basè subinterruptam aggregatis, invol. squamis linearibus acutiusculis apice ciliatis subcariosis pallidè flavidis disco æqualibus. ② in Indiâ or. ad Kamaon, ad Sokenaghar, ad Koemari legit cl. Hamilton. Conyza necessaria Ham.! herb. Gnaphalium Hamiltoni Wall.! cat. n. 3938 comp. n. 48. Fl. fœm. tennes innum. stylo indiviso longè exserto, masc. 8-10. Achæn. glabra. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or. et à cl. Royle.)
- *** Senecionifloræ, nempe foliis dentatis incisive membranaceis subglabris, capitulis paniculatis, invol. glabris aristatis.
39. B. SENECIOIDES, caule herbaceo erecto tereti striatulo brevissimè subpuberalo, foliis petiolatis ovalibus subacutis basi longè cuneatis distanter callosè-dentatis membranaceis in petiolo et nervo subhirsutis, paniculâ laxâ terminali subnudâ, capitulis longè pedicellatis, invol. squamis linearibus acuminatis ciliatis disco subæqualibus. ① in Indiâ orient. loco dicto Nawabgunj legit cl. Hamilton. Conyza senecioides Hamilton in Wall.! n. 3102. comp. n. 212. Fl. flavi, fœm. tennes stylo bifido, masc. cylindrici circ. 15. Achænia glabra (v. s. in herb. hor. Cur. Ind. or. nunc Soc. Lin. Lond.)
40. B. FLAVA, tota glabra herbacea erecta, caule tereti lævi ramoso, foliis infimis obovatis basi attenuatis subserratis obtusis, caulinis cordato-semiamplexicaulis acuminatis grossè callosè-serratis, ramis floridis axillaribus racemum corymbosum confertum gerentibus in paniculam laxè dispositis, invol. squamis linearibus cuspidato-acuminatis disco circ. æqualibus. ② in Indiâ or. prov. Nepal, Silhet, Ava, Tavoy, ins. Penang, etc. Conyza fasciculata Wall.! cat. n. 3017. comp. n. 127. non n. 3094. comp. 204. Senecio fasciculatus et Baccharis amplexicaulis Wall.! herb. non Lam. Erigeron obovatus Spreng.! in litt. ad Wall.! Conyza obliqua Ham.! herb. Erigeron falcatum Don prod. 172? sed quid falcatus? Flores flavi, fœm. multiseriales tenuissimi truncati, stylo exserto bifido, masc. 10-12. Pappus albus. Ovarium glabrum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et à cl. Royle.)
- β. subnuda, caule simplici subaphyllo. Hab. cum var. α. et cum eâ comm. (v. s.)
41. B. VIRENS (DC. in Wight contrib. p. 14), caule herbaceo erecto tereti glabro ramoso, foliis rigidulis sessilibus utrinquè sparsè puberulis, caulinis elongatis basi cuneatis sessilibus trifidis, lobis lanceolatis mucronatis regulariter serratis, terminali longiore, ramis obovatis basi cuneatis parè serratis, paniculâ laxâ, pedicellis axillaribus folio longioribus 1-5-cephalis capitulis pedicellatis, invol. squamis int. linearibus disco longioribus mucronatis. — in Indiâ orient. legit cl. Russel, in arenosis maritimis ad Negapatam (Wight!). Conyza virens Wall.! cat. n. 3037. comp. n. 147. var. C. Erigeron virens Russ.! herb. Fl. purpurascentes, fœm. innumeri, masculi 5. Achænia puberula. (v. s. in herb. hon. Cur. merc. angl. nunc Soc. Lin. Lond.)

42. *B. LAPSAŃOIDES* (DC. l. c. p. 15), glabra, caule herbaceo erecto tereti ramoso, foliis petiolatis membranaceis, caulinis sinuato-repandis hinc indè mucronato-serrulatis, lobis mucronatis lateralibus subtriangularibus terminali ovato, rameis obovatis vix subserratis, paniculà laxà oligocephalà, ramis lateralibus foliosis, capitulis pedicellatis, invol. squamis linearibus mucronatis disco paulò longioribus. (I) in Indiæ orient. sylvis secus ripas flum. Saluen prov. Martabanis ad Trogia, Kogun, Moalmyne et Amherst leg. cl. Wallich, ad Conrallum in Carnaticà austr. *Conyza virens* A. B. D. Wall. ! cat. et herb. n. 3037. comp. 147. Flor. flavidi, fœm. innumeri, masc. 10-12. Invol. post anth. omninò reflexa. Rami floridi glabri aut pube glandulosà minime subviscosi. (v. s. com. ab hon. Cur. Ind. or. et in herb. Wight.)
43. *B. HYMENOPHYLLA*, grabriuscula, caule herbaceo erecto tereti ramoso, ramis floridis glanduloso-puberis, foliis petiolatis membranaceis mucronibus exsertis sparsè subserratis, infimis longius petiolatis basi sublyrato-incisis, super. ovatis ovalibusve, paniculà laxà, pedunculis folio longioribus, capitulis pedicellatis, invol. squamis linearibus flores non superantibus. (I) in Martabanis æcus flumen Saluen leg. cl. Wallich. *Conyza hymenophylla* Wall. ! cat. n. 3038. comp. n. 138. Valdè affinis *B. lapsanoidi*, sed folia minùs divisa, fl. masc. circ. 20 et fœm. pauciores. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
- ß. subpubescens*, foliis inferioribus ovalibus basi cuneatis, vix subrepandis nec basi sublyratis, ramis floridis pube parçà non glandulosà subpubescentibus. (I) in Martabanis ad Kogun. (v. s. comm. cum *α*.)
44. *B. PAUCIFOLIA*, caule herbaceo erecto tereti subsimplici undiquè minutissimè puberulo apice subviscido, foliis petiolatis obovato-oblongis basi cuneatis hinc indè exsertè serratis suprà adultis glabris subtùs et novellis utrinquè molliter pubescenti-villosis paniculà elongatà subaphyllà, invol. squamis linearibus acuminatis flores æquantibus. (I) in Indiæ orient. verisim. prov. Silhet. *Conyza virens* E. Wallich ! cat. et herb. n. 3037. comp. 147. excl. A.-D. Affinis *B. lapsanoidi* et *hymenophyllæ*. Invol. demùm omninò reflexa. Fl. flavidi, masc. 12-18. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
45. *B. NAPIFOLIA*, caule herbaceo erecto tereti basi subglabro apice pubescente subglanduloso, foliis pubescentibus petiolatis lyratis lobis dentatis lateralibus ovali-oblongis terminali subrotundo, paniculà parvâ oligophyllà, ramis parçè foliosis, invol. squamis linearibus acuminatis florum longitudine. — in Indià orient. ad Tavoy. *Conyza napifolia* Wall. ! cat. n. 3075. comp. n. 185 Herba pedalis. Folia sesquipoll. longa. Flor. flavi, fœm. pauciseriales truncati, stylo exserto bifido, masc. centrales circ. 30. An fortè genericè distincta? (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
46. *B. MURALIS*, infernè glabra medio puberula apice glanduloso-pilosa, caule herbaceo erecto simplici tereti, foliis in petiolum angustatis lyrato-pinnatifidatis, lobis grossè dentato-serratis inf. utrinquè 2-3 triangularibus oblongive super. confluentibus ultimo majore ovato, paniculà elongatà laxà, capitulis pedicellatis, invol. squamis linearibus acuminatis glabris subcoloratis flores paulò superantibus. (I) in Indià orient. ad Paingti (an fortè Painkhandi distr. Kamaon?). *Conyza muralis* Ham. ! in Wall. ! cat. n. 3077. comp. n. 177. Fol. infer. 5 poll. longa, 1 $\frac{1}{2}$ poll. lata. Inv. et fl. purpurasc. Fl. fœm. innumeri stylo bifido exserto, masc. circ. 5-7 breviores. Pappus albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
47. *B. TRISULCA*, caule herbaceo erecto lævi trisulco, foliis sessilibus hispidis dentatis, radicalibus cuneatis, caulinis oblongis amplexicanlibus, corymbis paniculatis oligocephalis, squamis invol. aristatis disco longioribus. (I) in Nepalià. *Erigeron trisulcus* Don prod. fl. nep. 171. *Conyza trisulca* Ham. ex Dou. Pappus nivens. Flor. masc. pauci.
48. *B. MEMBRANACEA*, caule herbaceo erecto simplici tereti infernè glabro cæterùm piloso, foliis membranaceis sparsè pilosis grossè serratis, infer. obovatis acuminatis basi longè cuneatis in parte cuneatà nunc integerrima nunc

inciso-lobata sublyrata, summis distantibus linearibus serratis, pedunculis axillaribus glanduloso-pilosis capitula subglomerata sessilia gerentibus interruptè spicatis in paniculam laxam dispositis, invol. squamis linearibus acuminatis disco æqualibus. ① in Ind. orient. ad Prome regni Avenis legit ocul. Wallich. *Conyza membranacea* Wall.! cat. n. 3019. comp. n. 129. Folia infer. pauca, 7-8 poll. longa, 2-2½ poll. lata, summa pedunculis paulò breviora. Flor. fœm. pluriseriales tenuissimi circ. 30-40, masc. circ. 25-30. Pappus albus tenuissimus parvus. (v. s. com. ab hon. Cur. merc. angl. Ind. or.)

49. *B. viscosula*, caule herbaceo erecto tereti subimplici apice puberulo-viscoso, foliis membranaceis denticulis exsertis, callosis serratis obovatis basi cuneatis subpetiolatis sæpe lyrato-incisis tenuissimè utrinquè puberulis, paniculâ laxâ, ramis subnudis apice capitula pauca sessilia subspicata gerentibus, invol. squamis linearibus acuminatis flores æquantibus in lineâ mediâ extus puberulis. ① in Indiæ orient. regno Avenis propè Taong-Dong novembri legit cl. Wallich. *Conyza viscosula* Wall.! cat. n. 3081. comp. n. 191. Herba pedalis. Fol. 4 poll. longa 1-½ lata. Capitula parvula 2 lin. lata. Flor. fœm. tennes truncati stylo brevissimo exserto, masc. 15-20. Pappus albus. (v. s. comm. ab hon. Cur. Ind. or.)

β. *hispidula*, caule a basi ad apicem patentim hispido, pilis supernè pluribus glanduloso-viscosis foliis subtus villosulo-hirsutis, invol. squamis flores superantibus dorso puberulis apice longè ciliatis. ① cum priori eodem loco et die lecta à cl. Wallich et tamen fortè species propria? (v. s. comm. cum α.)

50. *B. HEYNEANA* (DC. in Wight contrib. p. 15), caule herbaceo erecto ramossissimo subtilissimè striatulo et puberulo, foliis oblongis basi attenuatis apice obtusis hinc inde dentes exsertos acutos gerentibus utrinquè glabriusculis, infer. basi lyrato-auritis, paniculâ foliosâ laxâ, ramis folii longitudine apice subcorymbosis, invol. squamis linearibus acuminatis dorso subpuberulis disco paulò longioribus. — in Indiâ orient. legit cl. Heyne. *Conyza Heyneana* Wall.! cat. n. 3089. comp. n. 199. Con. aurita Heyne! herb. non Lin. f. Folia 2-3 poll. longa, pollicem lata, infima ignota. Fl. flavidi, fœm. innum. tennes stylo exserto, masc. circ. 12. Pappus albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et in herb. Wight.)

51. *B. CUNEIFOLIA*, pube brevissimâ subsabrida, caulibus herbaceis erectis teretibus striatis, foliis sessilibus cuneiformibus obtusis serraturis exsertis acutis mucroniformibus distantibus sæpè reflexis, paniculæ oligocephalæ ramis erectis capitula pauca congesta apice gerentibus, invol. squamis linearibus acutis discent æquantibus, fl. masc. numero fœmineos superantibus. — in Indiæ prov. Martaban ad ripas flum. Saluen propè Phanoë legit cl. Wallich. *Conyza cuneifolia* Wall.! cat. n. 3101. comp. n. 211. non Lam. nec Rausch. Folia 12-15 lin. longa, 5-6 lata. Recept. nudum. Flores flavidi, fœm. pauciseriales, masc. circ. 40-50. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

52. *B. SUBSIMPLEX*, caule imâ basi fruticuloso gracili glabro, ramis herbaceis teretibus elongatis subnudis pubescenti-villosis, foliis infer. obovatis petiolatis basi cuneatis rariter et acutè callosa-dentatis suprâ glabriusculis subtus pallidis pubescentibus, rameis minoribus petiolatis ad petiolum sæpè incisis, capitulis ad axillas supremas 2-3 brevè pedicellatis spicam interruptam ferè constituentibus, invol. squamis linearibus acuminatis dorso puberulis apice glabris arcuato-reflexis. — in Indiæ. or. prov. Silhet. *Conyza subsimplex* Wall.! cat. n. 3003. comp. n. 113. C. paniculata Willd. sp. 3. p. 1920 ex Spreng. in litt. ad Wallich sed descr. non ritè quadrat. Cor. ochroleucæ, fœm. tenuissimæ stylo simplici, masc. circ. 20. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

*** *Fasciculiform, nempè foliis dentatis, ramis elongatis capitula ad axillas foliorum floralium aggregata et interruptè spiciformia gerentibus.*

53. *B. RACEMOSA*, caule herbaceo tereti villosa aut demum glabrato, foliis ellipticis acutis basi attenuatis petiolatis inæqualiter serratis utrinquè sparse villosa, capitulis ad axillas foliorum superior. dense congestis sessilibus in apice interruptè formam longè digestis, invol. squamis linearibus acuminatis densa puberis demum stellato-patulis intus nitidis. ① in Indiâ orient. ad Nokote legit cl. Wallich. *Conyza racemosa* Wall.! cat. n. 3002. comp. n. 112. non Spreng. *C. stachyoides* Spreng. in litt. ad Wall. Cor. purpureæ, fœm. innumeræ stylo bifido, masc. 5-7. Pappus albidus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
54. *B. MACROSTACHYA*, tota densè hirsuta albida, caule herbaceo erecto tereti subsimplici, foliis lanceolatis utrinquè acuminatis acutè serratis, superioribus subsessilibus ex axillis ramos abbreviatis foliolosos densè floridos gerentibus, paniculâ elongatâ interruptè subspicatâ confertâ fasciculatâ, invol. squamis linearibus acutis, ext. hirsutis, interioribus apice coloratis. — in Ind. orient. ad Nokote Nepalesium martio flor. legit cl. Wallich. *Conyza macrostachya* Wall.! cat. n. 3053. comp. n. 143. *Baccharis auriculata* Wall.! olim in herb. Flor. purpurei, fœm. innum. tennes, masc. 10-12. *Achænia glabra*. Rec. plannum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
55. *B. HIERACIFOLIA* (DC. in Wight contrib. p. 15), tota hirsuta, caule herbaceo erecto tereti simplici, foliis calloso-dentatis, infer. obovatis obtusis in petiolum attenuatis, super. ovalibus oblongisve acutis sessilibus aut semiamplexicaulibus, capitulis sessilibus confertis in thyrsum ovato-oblongum digestis, invol. squamis linearibus acuminatis glabriusculis disco longioribus. ① in Indiâ orient. prov. Nepaliâ (Ham.), Silhet (Wall.) et penins. Carnaticâ (Wight!). *Conyza hieracifolia* Spreng. syst. 3. p. 514. *Erigeron hieracifolium* Don prod. fl. nep. 172. *Conyza communis* Wall.! cat. et herb. n. 3018. comp. n. 128. Herba pedalis. Fol. subtus magis villosa-hirsuta. Invol. apice subpurpurascens. Flor. fœm. innumeri tennes stylo exserto indiviso, masc. 5. Ovaria pubescentia. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et in herb. Wight.)
56. *B. HOLOSERICA*, tota villis sericeis longis vestita, caule herbaceo erecto tereti apice hirsuto subsimplici, foliis oblongis basi attenuatis acutis profundè parcè et acutè serratis adpressè sericeo-villosis, paniculâ elongatâ interruptâ, ramis baseos folia superantibus apice racemosis, sup. brevioribus congestè polycephalis, invol. squamis linearibus acuminatis puberis ciliatis disco ferè longioribus. — in Indiâ or. prov. Martabanâ ad ripas flum. Saluen propè Phanoë legit cl. Wall. *Conyza holosericea* Wall.! cat. n. 3103. comp. n. 213. *C. sericea* Wall.! cat. burm. n. 1892. Pars plant. inferior ignota. Flor. purpurascens, fœm. innumeri tennes, stylo exserto subsimplici, masc. 10-12 glabri, anth. exsertis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
57. *B. VIMINEA*, caule erecto ramoso, ramis elongatis teretibus pubescentibus, foliis lanceolato-oblongis basi attenuatis acuminatis grossè ad apicem et acutè serratis utrinquè pubescentibus, ramulis floridis axillaribus folio brevioribus foliola et capitula 7-8 gerentibus, invol. squamis linearibus pubescentibus disci longitudine. — in insulâ Timor. Recept. alveolare glabrum. Flor. fœm. purpurei innumeri stylo exserto bifido, masc. 5-7 glabri. *Achænia glabra*. (v. s. comm. à Mus. Reg. Par.)
58. *B. FASCICULATA*, herbacea erecta ramosa tota cinereo-villosa, fol. infer.... super. oblongo-lanceolatis basi attenuatis sessilibus supernè rariter serratis acutis, capitulis sessilibus interruptè spicatis nempè ad axillas foliorum rami inferiorum solitariis superiorum 2-3 congestis, invol. squamis linearibus acuminatis flores æquantibus, ovario glabro, pappo 6-8-seto. — in Indiâ or. insulâ Timor et regione Burmanorum propè Amherst. *Conyza fasciculata* Wall.! cat. n. 3094. comp. n. 204. *C. glomerata* Wall.! cat. Burm. n. 1409. (v. s.)

59. *B. GRACILIS*, infernè glabra supernè puberula, caule herbaceo striato erecto? ramoso, foliis oblongis basi attenuatis obtusis, inf...., mediis sinuato-incis, summis subserrulatis acutis, paniculâ ramosissimâ laxâ, invol. squamis subvillosis linearibus acutis flores subæquantibus. — in Indiâ orient. legit cl. Heyne. *Conyza gracilis* Heyne! in Wall.! cat. n. 3064. comp. n. 174. Pars infer. plantæ ignota. Folia sub paniculâ ramis ferè pollicem longa, 3-4 lin. lata. Flores pallidè flaventes, fœm. innum. tenues, masc. circ. 12. Achænia minima teretiuscula. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
60. *B. PURPUREA*, tota parvè villosa, caule erecto herbaceo simplici tereti, foliis ovatis acutis serratis basi integerrimis attenuatis, ramis plurimis subnudis paniculatim dispositis capitula sessilia spicata subconferta gerentibus, invol. squamis linearibus acutis, exter. brevibus cinereo-virescentibus, intimis coloratis flores paulò superantibus. ① in montibus seu collibus Indiæ or. circa Rajmahal legit cl. Hamilton. *Conyza purpurea* Ham.! in Wall.! cat. n. 3076. comp. n. 186. Radix exilis subsimplex. Herba pedalis. Involucra purpurascencia. Flor. fœm. innumeri tenues, masc. circ. 20. Pappus albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
61. *B. GLOMERATA* (DC. in Wight contrib. p. 15), tota villosa-subhirsuta cinerea, caule herbaceo erecto tereti ramoso, ramis in paniculam dispositis foliosis apice capitula sessilia glomerata interruptè spicata gerentibus, foliis lanceolatis serratis utrinquè acutis, invol. squamis multiserialibus, ext. minimis, intimis linearibus flores paulò superantibus, fl. masc. 30. ① in Indiâ orient. reg. Burmannorum ad Prome, Taong-Dong, Amherst, Moalmyne legit cl. Wallich, in arenosis maritimis circa Madras et Negapatam cl. Wight. *Conyza glomerata* Wall.! cat. n. 3055. comp. n. 165 excl. litt. C. *Conyza argentea* Spreng. in litt. ad Wall. non alior. Flor. fœm. vix masc. numerosiores truncati, stigm. bifido purpureo, masc. 5-fidi antheris luteis exsertis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
62. *B. LEPTOCLADA*, infernè glabrata supernè cinereo-villosula, caule herbaceo erecto tereti ramoso, ramis virgatis paniculatis capitula sessilia solitaria aut apice glomerata interruptè spicata gerentibus, foliis glabrisculis obovato-oblongis basi longè cuneatis ferè petiolatis apice grossè et acutè serratis, invol. squamis linearibus acuminatis florum circ. longitudine, fl. masc. circ. 12-15. ① in Nepaliâ legit cl. Wallich. *Conyza glomerata* var. C. Wall.! cat. et herb. n. 3055. comp. n. 165. Valdè accedit ad *B. glomeratam* sed pars infer. ferè glabra, invol. squamæ pauciores, fl. masc. dimidiò numerosiores, antheræ inclusæ. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
63. *B. PERROTETIANA*, tota cinereo-velutina, caulibus herbaceis erectis subsimplicibus teretibus, foliis oblongis acutis serratis basi attenuatis, racemo spiciformi terminali oligocephalo, invol. squamis linearibus puberulis apice subcoloratis acutis disco paulò longioribus. — in Africæ occid. arenosis locis inundatis propè flum. Senegal et inter fl. Gambie et peninsulam promontorii Viridis mense martio flor. legit cl. Perrottet. Flor. purpurei, fœm. innumeri, masc. 15-20. Achænia teretiuscula glabra. (v. s. comm. à cl. Perrottet.)
64. *B. SOLIDAGINOIDES*, caule herbaceo erecto tereti subsimplici molliter villosa, foliis lanceolatis sessilibus basi attenuatis suprâ pubescentibus subtus molliter villosa-canescens serratis, capitulis aggregatis in racemum interruptum subfoliosum subspicatum dispositis, squamis lineari-subulatis acutis villosis apice laxiusculis, flor. fœm. innumeris, masc. 4-5. ②? in Sierra-Leonâ. *Gnaphalium solidaginoides* Poir.! suppl. 2. p. 805. excl. patr. Species à Poirétia descr. in h. Desf. ex specim. quod ipse cl. et amico magistro olim communicavi, ex herb. L'Hér. orto. (v. s.)
- *** Oxyodontæ, nempe basi glabræ apice hirsutæ, foliis obovatis dentes duros acutos subaristatos gerentibus, capitulis fasciculatis aut subpaniculatis.
65. *B. SPINELLOSA*, caule herbaceo tereti basi pilosiusculo apicè ramisque cinereo-villosissimo, foliis infer. obovatis acutis longè cuneato-attenuatis sparse pilosis

inæqualiter et grossè serrato-spinulosis, summis oblongo-linearibus utrinquè sericeo-villosissimis æqualiter spinuloso-serratis, ramis floridis paniculatis folia longè superantibus apice capitula sessilia conferta spicata gerentibus.

①? in Indiâ orient. ad Prome regni Avenis legit cl. Wallich. *Conyza sericea* Wall. cat. et herb. n. 3021. comp. n. 131 non Ait. nec Bory. Folia inf. 6-poll. longa, a poll. lata. Rami florif. infer. 6 pollic. Flor. fœm. pluriser. tenuissimi stylo indiviso, masc. circ. 20. Pappus albus tenuis deciduus parvus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

66. *B. BELANGERIANA*, caule herbaceo erecto à basi ramoso basi sparso hispido apice molliter villosa, foliis infimis obovatis subobtusis in petiolum brevem attenuatis, summis subsessilibus, omnibus argutè serratis suprà sparso pilosis subtus densè sericeo-villosis, capitulis ad axillas superiores congestis ferè interruptè spicatis, invol. squamis linearibus glabriusculis apice ciliatis. ① ad littus Malabaricum legit cl. Belanger. Affinis *B. oxyodonta* sed erecta, multò major. Recept. angustum nudum. Flor. fœm. in genere pauci, masc. 8-10. Antheris exsertis. *Achænia* teretia glabriuscula. (v. s. comm. à cl. inv.)

67. *B. ERANTHA* (DC. in Wight contrib. p. 15), caule herbaceo erecto tereti basi glabro apice ramoso et mollissimè hispido, foliis membranaceis caulinis sessilibus ovalibus acutè et exsertè apiculato-dentatis molliter pubescentibus, summis cordato-amplexicaulibus, pedunculis axill. et terminalibus 1-cephalis solitariis aut paucis aggregatis demum valdè elongatis, invol. squamis linearibus ciliatis acuminatis, cor. fœm. et masc. apice barbatis. ①? in Indiâ verisim. *Maderaspatanâ* legit cl. Wight. Herba pallidè virens. Fol. inf. ignota caulina 3 poll. longa, 15-16 lin. lata. Invol. demum reflexum. Rec. glabrum. Flor. masc. 25-30. (v. s. in h. Wight.)

68. *B. OXYDONTA* (DC. in Wight contrib. p. 15), herbacea villis longis sericeis laxè villosa, caulibus pluribus simplicibus ramosive diffusis, foliis argutè et grossè serratis subtus magis sericeo-villosis, infimis obovatis petiolatis, superioribus ellipticis subsessilibus, capitulis paucis ad apices ramorum racemoso-corymbosis subsessilibus, invol. squamis linearibus acuminatis disco paulò longioribus. ① in Indiâ orient. ad Prome regni Avenis legit cl. Wallich, circa Pondichéry (Belanger!) et ad littora maris versus Negapatam frequens (Wight). *Conyza oxydonta* Wall. cat. n. 3015. comp. n. 125. *Erigeron graminosus* Russ.! herb. Folia infima 2-2½ poll. longa, 9-10 lin. lata, caulina 9-12 lin. longa, 4-5 lata. Flor. fœm. pluriseriales, masc. circ. 15-20. Pappus albidus parvus tenuis deciduus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

β. *tenera*, tota magis sericeo-villosa, foliis summis vix serratis, paniculâ laxiore polycephalâ. ① in Indiâ oriental. ad Onde, Holasannj et in Nepaliâ leger. cl. Hamilton et Wallich. *Conyza tenera* B. et C. Wall. cat. et herb. n. 3033. comp. n. 133. *Conyza* bovina Ham.! herb. *Erigeron cuspidatum* Spreng. in litt. ad Wall. (v. s. comm. cum α.)

γ. *glabrata*, tota glabriuscula laxa, capitulis distantibus longius pedicellatis. — in Indiæ prov. boreali-occid. *Conyza tenera* A. Wall. cat. et herb. n. 3033. comp. n. 133. (comm. cum var. α et β à cl. Royle.)

***** Semivestitæ, nempe infernè *glabrata* supernè *velutina*, foliis oblongis acuminatis subintegris, pappo albo.

a. Receptaculo hirsuto.

69. *B. RIPARIA*, glabra, caule fruticoso scandente tereti, foliis breviter petiolatis ellipticis breviter acuminatis glanduloso-serratis rigidis subtus subpallidioribus, paniculæ oblongæ ramis velutinis folio longioribus 1-3-cephalis, capitulis pedicellatis, invol. squamis lanceolato-linearibus acutis florum disco paulò brevioribus. ② in insulâ Javâ ad ripas fluminum legit cl. Blume. *Conyza riparia* Blume! bijdr. 899 non Kunth. Flor. masc. circ. 20 lobis valdè hirtis. Recept. hirsutum. (v. s. comm. ab ill. inv.)

70. *B. CHINENSIS*, caule herbaceo erecto tereti glabro apice cum pedunculis in-

volucrisque pubescenti-velutino, foliis breviter petiolatis ovalibus apice angustè acuminatis utrinquè glabris dentibus distantibus callosis acutis exsertis hinc inde instructis, paniculâ laxâ, invol. squamis linearibus, exter. brevissimis, intimis multò longioribus, recept. villosissimo, acheniis teretiusculis subcostatis puberulis. — in Chinâ (Linn.), in Javâ sylvis montosis (Blume!), et insulâ Penang (Wallich!). Baccharis nitida Wall.! cat. et herb. n. 3043. comp. n. 133. Conyza Chinensis Linn. sp. 1208 (excl. Rumph. syn.). Blume! bijdr. 898. Pers. ench. n. 21 nec n. 16. Eupatorium reflexum Linn. ex Steud. Affinis B. semivestitæ nec ullo modo genericè separanda. Differt foliis etiâ superioribus petiolo donatis, paniculâ minùs velutinâ, invol. squamis acutioribus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

71. B. SEMIVESTITA, caule herbaceo tereti erecto apice cum ramis floridis densè velutino-villoso, foliis elliptico-oblongis acuminatis basi attenuatis subsessilibus grossè serratis utrinquè glabris, paniculâ terminalis ramis inferioribus distantibus, capitulis ad apicem ramulorum sessilibus densè congestis, invol. squamis linearibus acutis dorso pubescentibus flores æquantibus, recept. villosissimo. in Nepaliâ legit cl. Wallich. Conyza semivestita Wall.! cat. n. 2996 A. comp. n. 106 A. Folia 6 poll. longa, 2 poll. lata. Fl. fœm. innumeri, masc. pauci. Ovarium puberulum subcylindraceum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

72. B. PROCERA, caule herbaceo erecto tereti paniculato-ramoso infernè glabro apice ramisque lanuginoso-villosis, foliis caulinis glabris sessilibus ovali-lanceolatis acuminatis subserratis, paniculâ amplâ divaricatâ, capitulis ad apices ramulorum aggregatis, invol. squamis dorso pubescenti-hirsutis linearibus flores paulò superantibus, receptaculo villoso. — in Indiâ or. prov. Silhet. Conyza procera Wall.! cat. n. 3050. comp. n. 140 non Desf. Folia 5 poll. longa, 2 poll. lata, infima ignota. Fl. purpurei, fœm. innum. stylo bifido, masc. 12-15. Achenia puberula. Pappus albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

b. Receptaculo glabro.

73. B. MYRIOCEPHALA, caule herbaceo erecto tereti infernè cum foliis glabro supernè cum paniculâ ramis puberulo-hirtis, foliis oblongis acuminatis basi longè in petiolum attenuatis subserrulatis, paniculâ ramosissimâ subaphyllâ, capitulis secus ramulos laterales sessilibus et terminalibus, invol. squamis linearibus acuminatis recurvo-patentibus dorso puberulis disco paulò longioribus. — in Indiâ or. prov. Silhet legit amic. Wallich. Conyza squarrosa Wall.! cat. et herb. n. 3025. comp. 135 non Linn. Herba elata spectabilis. Folia 8-12 poll. longa, 1-2 poll. lata. Rami inf. ex axillis orti villosi oligocephali, super. elongati ramosissimi. Flor. fœm. innumeri truncati stylo bifido, masc. 5-9. Achenia teretiuscula. Pappus pallidè rufescens. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

74. B. SPECTABILIS, ferè tota glabra ramulis subhirsutis, caule herbaceo erecto paniculato ramoso tereti, foliis utrinquè lævibus ovali-lanceolatis basi longè attenuatis apice acuminatis hinc inde subincisis remotè et calloso-serratis suprâ lucidis, paniculâ amplissimâ laxâ, ramis velutino-hirtellis lateralibus folio longioribus polycephalis, invol. squamis linearibus acuminatis dorso glabrisculis apice ciliatis flores ferè superantibus, recept. alveolato glabro. — in peninsulâ Indiâ orient. legit cl. Wight (pl. exs. 2. 16). Media omninò inter B. myriocephalam et elongatam, sed paniculâ laxissimâ polycephalâ ab utrâque præter car. indic. diversa. Folia 8-9 poll. longa, 1 $\frac{1}{2}$ -2 lata. An var. B. myriocephalæ? sed capitula duplò majora quam B. myriocephalæ, invol. squamæ non squarrosæ. (v. s. comm. à cl. Wight.)

75. B. ELONGATA, basi glabriuscula supernè pubero-velutina subsericea, caule herbaceo erecto simplici tereti, foliis suprâ scabridis subtus pubescentibus lanceolatis calloso-serratis, inferioribus basi attenuatis et hinc inde subincisis, superioribus vix basi attenuatis apice acutioribus, paniculâ elongatâ ramis plaribus brevibus oligocephalis, invol. squamis linearibus acuminatis dorso

puberulis apice ciliatis flores paulò superantibus demùm reflexis, recept. alveolato glabro. — in Indiâ orient. insulâ Penang legit cl. Wallich. Conyza elongata Wall.! cat. n. 3078. comp. n. 188. Herba bipedalis. Flores albidî, fœm. innumeri tenues stylo exserto apice bifido, masc. circ. 12. Pappus albus. Achenia glabra teretiuscula. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

96. B. OBOVATA, caule herbaceo erecto tereti simplici molliter hirsuto, foliis obovatis acuminatis basi cuneatis apice serrulatis suprà sparse pilosis subtus cinereo-villosis, pedunculis paucis molliter hispidissimis folio longioribus 1-2-cephalis in paniculam laxam oligocephalam dispositis, invol. squamis linearibus acuminatis disco paulò longioribus, recept. glabro. ① in Nepaliâ inferiore legit cl. Wallich. Conyza obovata Wall.! cat. n. 3022. comp. n. 132. Herba sesquipedalis. Folia 5 poll. longa, 2 poll. lata. Capitula diam. semipollic. Fl. purpurascens, fœm. innumeri truncati tenues stylo bifido, masc. circ. 20. Pappus albus deciduus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

***** *Macrophylla, nempe foliis ovalibus oblongive subintegerrimis amplis, capitulis paniculatis, pappo rufescente.*

77. B. MACROPHYLLA, caule suffruticoso, foliis ovato-aut oblongo-lanceolatis acuminatis basi in petiolum angustatis glanduloso-serratis suprà puberulis subtus griseo-tomentosis, paniculâ divaricatâ laxâ, ramis nudis apice subracemosis, invol. squamis linearibus acuminatis glabriusculis flores æquantibus. ③ in insulâ Javâ montanis ad rivulos legit cl. Blume. Conyza macrophylla Blume! bijdr. 896. Folia pedem et ultrâ longa, 5 poll. lata. Recept. alveolatum. Flor. fœm. innumeri stylo exserto bifido, masc. pauci. Pappus pallidè rufescens. Achenia teretiuscula striata pallida subpuberula. (v. s. comm. à cl. Blume.)

78. B. AROMATICA, caule herbaceo erecto tereti ramoso pube minimâ confertâ subglandulosâ præsertim supernè vestito, foliis ovatis acuminatis basi longè secus nervum petioliformem productis utrinquè puberulis, paniculâ laxâ polyccephalâ, invol. squamis exter. brevibus subrecurvîs dorso puberulis, inter. elongato-linearibus acuminatis glabris flores paulò superantibus. — in Ind. or. inter vallem sensu stricto Nepaliâ et Noakote legit cl. Wallich martio forè defloratam. Conyza aromatica Wall.! cat. n. 3054. comp. n. 164. Erigeron Cochinchinensæ Spreng. in litt. ad Wallich non Pers. Folia 5-6 pollic. longa, 1-2 lata. Pappus rufus. Flor. fœm. innumeri filiformes, masc. circ. 10. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

79. B. DENSIFLORA, caule herbaceo erecto ramisque molliter lanuginosis, foliis elliptico-oblongis acuminatis basi attenuatis hinc indè acutè inciso-dentatis suprà scabridis subtus lanuginosis nonnullis basi sublyratis, paniculâ ramosissimâ confertissimâ, invol. squamis linearibus acuminatis glabriusculis corollas subæquantibus, flor. fœm. innumeri, masc. circ. 5. — in Indiâ orient. ad Pundua legit Fr. de Silva. Conyza aromatica Wall.! cat. n. 2997. comp. n. 107. Conyza amygdalina Spreng. in litt. non Lam. Folia ad paniculæ ramos linearî-lanceolata rariè serrata ramis breviora, caulina infer. 6 poll. longa, 2 poll. lata. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

80. B. EXCISA, caule herbaceo erecto tereti ramoso velutino-pubescente, foliis elliptico-oblongis dentatis basi incisîs sublyratis suprà puberulis subtus lanuginosis, paniculâ amplâ ramosissimâ, invol. squamis linearibus disco longioribus hirsutis subacutis falcatis recurvis. — in Indiâ Malaicæ insulâ Penang legit cl. Wallich. Conyza excisa Wall.! cat. n. 3011. comp. n. 122. Affinis B. densifloræ sed panicula magis laxa. Fol. infer. basi longè attenuata. Fl. fœm. innumeri, masc. 5-6. Pappus rufus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

81. B. LONGIFOLIA, caule herbaceo erecto tereti simplici infernè cum foliis glabro supernè cum paniculâ ramis pubero aut villosis, foliis oblongis utrinquè acuminatis serratis, paniculâ elongatâ polyccephalâ, ramis subsimplicibus, capitulis lateralibus subsessilibus et terminalibus subsessilibus, invol. squamis linearibus acuminatis glabris flores æquantibus. — in Indiâ orient. legit cl. Heyne. Co-

- nyza densiflora* Heyne in Wall.! cat. n. 3026. comp. 126. Folia 8-9 poll. longa, 1½-2 poll. lata. Panícula pedem et ultra longa. Inv. squamis longiores 4 lin. Pappus rufescens. Achænia striato-angulata velutina. Fl. fœm. innumeri tenuissimi stylo bifido, masc. circ. 5. Cor. extus ad apicem parce glandulosa. (v. s. comm. ab hon. Curia merc. angl. Ind. or.)
82. *B. SYLVATICA*, caule herbaceo simplici, foliis breviter petiolatis oblongo-lanceolatis utrinquè acuminatis suprâ basin argutè serrulatis et ad costam infrâ subvelutinis, racemis simplicibus paniculatis nutantibus, invol. squamis linearibus acutis exter. brevibus pubescentibus int. disco paulò longioribus. — in insulæ Javæ montanis Megamendung. Con. sylvatica Blume! bijdr. 898. Pappus rufescens. Flor. fœm. innumeri tenues stylo exserto bifido, masc. 6-7 lobis hirtellis anth. et interdum stylo exsertis. (v. s. comm. à cl. Blume et Belanger.)
83. *B. SASSILIFOLIA*, caule herbaceo erecto apice confertè puberulo, foliis sessilibus oblongo-lanceolatis utrinquè acuminatis superne serratis discoloribus suprâ scabris infrâ rugosis et arachnoideo-puberis, racemis compositis laxè paniculatis, invol. squamis linearibus, exter. brevibus puberulis, int. glabris disci longitudine. — in ins. Javæ altis montis Burangrang. *Conyza sessilifolia* Blume! bijdr. p. 899. Pappus subrufescens. Fl. fœm. tenues stylo longè exserto bifido, masc. 5 antheris exsertis lobis hispidulis. (v. s. comm. à cl. Blume.)

***** Appendiculatæ, nempe foliis petiolatis quasi lyratis seu petiolis appendices foliaceas vel lobos distinctos gerentibus.

84. *B. BALSAMIFERA*, caule basi suffruticoso, ramis teretibus lanato-villosis, foliis oblongo-aut elliptico-lanceolatis duplicato-dentatis rugoso-venosis suprâ villosis subtus sericeo-lanatis, petiolis lobis lineari-lanceolatis appendiculatis, corymbo subpaniculato divaricato, invol. squamis linearibus acutis hirsutis. § 2 in Indiâ orient. continentali et insulari, Luzoniâ, Javâ, Cochinchinâ. *Conyza balsamifera* Linn. sp. 1208. Blume! bijdr. 895. Wall.! cat. et herb. n. 2998. comp. 108. *Conyza vestita* Wall.! herb. 1824. *Pluchea balsamifera* Less. syn. 1831. p. 150. An *Erigeron sericeum* Retz obs. 5. p. 28? *C. odorata* Rumph. amb. 6. t. 24. f. 1. non Linn. *Baccharis Salvia* Lour. cochinch. 2. p. 603. Variat foliis suprâ nunc villosissimis nunc subvillosis ferè glabratiss, dentibus inæqualibus aut æqualibus acutis embohtusive, petiolis interdum nudis sæpius appendiculatis undè folia stricto sensu lyrata. Pappus rufescens. Antheræ caudatæ. Achæniûm maturum glabrum striato-sulcatum (v. s.)
85. *B. GRANDIS*, caule herbaceo erecto angulato tenuiter pubescente, foliis ovalibus acuminatis suprâ glabris subtus pubescentibus serratis petiolatis, petiolo lobulos 5-6 distantes lineares acutos gerentibus. — in Indiâ orient. ad Tavoy Burmannorum legit cl. Wallich sinè floribus. *Conyza grandis* Wall.! cat. n. 3051. comp. 161. *Conyza falcigera* Wall.! cat. Burm. n. 2048. Valdè affinis *B. balsamiferæ* et ex hac analogiâ ad Blumeam relata. Folia cum petiolo 9 poll. longa, 3 poll. lata. (v. s. sine fl.)
86. *B. APPENDICULATA*, caule suffruticoso, foliis oblongo-lanceolatis argutè serrulatis subtomentosis et subrugosis, petiolis appendiculatis, racemis compositis paniculatis. § in insulâ Javâ ad fluviorum ripas circa Buitenzorg. *Conyza appendiculata* Blume bijdr. p. 895 non Lam. *Baccharis gratissima* Blume! in litt. 1831? Florum non satis evolutorum struct. non examinare quivi.

Ser. II. CAULOPTERÆ, nempe foliis secus caulem ramosve decurrentibus.

* Alatæ, nempe alis integerrimis.

87. *B. VERONIOIDES*, suffruticulosa tota densè velutino-hirsuta, foliis ovato-lanceolatis acutè et callosè denticulatis in alas integerrimas desinentibus, paniculæ terminalis ramis axillaribus 2-3-cephalis folium paulò superantibus, invol. squamis imbricatis linearibus acuminatis subsquarrosis, exter. dorso

hirsutulis, intimis glabris scariosis longioribus. ζ in Indiâ or. *Vernonia Wightiana* Wall. cat. n. 2923. comp. 33 sed in h. Wight non reperi et ideo errorem quemdam timeo. Diversissima à *Conyza Wightiana* quæ *Blumea nostra Wightiana*. Fl. purpurei, margin. innumeri fœminei, centrales 10-12 masculi stylo incluso verisim. ovario abortivo. Achænium teres tenue. (v. s. in h. Cur. Ind. or. nunc soc. Linn. Lond.)

88. *B. OLOPTERA*, tota pubescenti-subglandulosa, caule basi suffruticulosos, foliis oblongis rariter denticulatis acutis basi productis in alas longas latas integerrimas, capitulis ad apices ramulorum solitariis, invol. squamis multiseriatis basi erectis apice subulatis squarrosos-reflexis. ζ in Africâ ad flumen Gambie mense martio legit cl. Perrottet, in campis recenter igne devastatis. Fl. fœm. innumeri tenuissimi truncati, masc. exacte cylindrici. Anth. basi caudatæ aut vix ac ne vix subdentatæ. Achænium teres villosus. Recept. glabrum. Pappus flosculis longior. (v. s. comm. ab indef. Perrottet.)

89. *B. SALVIFOLIA*, herbacea simplex molliter villosa-tomentosa, foliis lanceolato-oblongis subintegerrimis acutis basi semimplexicanibus decurrentibus, ramis lateralibus 2-5-cephalis in paniculam elongatam dispositis, capitulis nutantibus, invol. squamis ext. brevibus foliaceis squarrosis, intimis elongatis acuminatis glabris. η in insulâ Bourboniæ ubi ab incolis dicitur *Sauge*. *Conyza salvifolia* Bory voy. isl. afr. 3. p. 176. Folia trita odorata, loco Tabaci adhibentur. Rec. nudum. Fl. fœm. innumeri filiform. stylo exs. masc. circ. 12-15. Achænia teretiuscula glabra. (v. s.)

90. *B. ALATA*, caule herbaceo erecto ramoso foliisque pube brevi subrufâ pubescenti-velutinis, foliis elliptico-oblongis dentatis secus caulem in alas integerrimas longè productis, pedunculis axillaribus mono-aut oligocephalis racemoso-paniculatis, capitulis suberectis, squamis ext. lanceolatis foliaceis squarrosis subpuberis, inter. linearibus scariosis flores æquantibus. — in Indiâ or. legit cl. Heyne. *Conyza alata* Roxb. cat. h. calc. p. 61. *Conyza alata* var. *A.* Wall. cat. n. 3039. comp. 149. *Erigeron alatum* Don prod. 171. *Vernonia alata* Heyne! herb. Fl. purpurei, masc. 10-12. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

β . *cernua* (Wall. cat. et herb. n. 3012. comp. n. 122 sub *Conyza*), caule herbaceo erecto ramoso foliisque pube brevi subrufâ pubescenti-velutinis, foliis oblongis acuminatis denticulatis secus caulem in alas integerrimas longè productis, pedunculis axillaribus mono-aut oligocephalis racemoso-paniculatis recurvis, capitulis cernuis, invol. squamis ext. lanceolatis foliaceis puberis, inter. longis nitidis scariosis apice recurvis demum patentibus. — in Nepaliâ, ad Kamaon, Sirmore. Pubes fortè glandulosa. Invol. radians Carlinæ more. Fl. fœm. innumeri stylo bifido brevè exserto, masc. 12-18. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

γ *Natalensis*, alis integerrimis in parte caulis infer. subundulatis, capitulis cernuis, invol. multiseriatis magis squarrosis. — in Africâ austr.-orient. ad Port-Natal legit cl. Drege, n. 4996. An species propr.? Fl. masc. circ. 12 anth. ecandatis. Cor. apice summo subscabris. (v. s.)

91. *B. GARIEPINA*, suffruticulosa ramosissima tota canescenti-villosa, foliis oblongo-linearibus integris subacutis in alas integras productis, capitulis pedicellatis subpaniculatis, invol. squamis lineari-subulatis rectis disco subæqualibus, intimis non scariosis. ζ in Africæ austr. regione *Gariépina* legit. cl. Drege, n. 2722. Folia 8-9 lin. longa, $1\frac{1}{2}$ -2 lata. Fl. masc. 10. Anth. tenuissimæ caudatæ? Achænia tenuia vix pubescentia. (v. s.)

* * Aurisæ, nempe alis incisus aut interruptis.

92. *B. PTERODONTA* (DC. in Wight contrib. p. 15), caule herbaceo tereti ramoso vix subpuberulo apice viscoso, foliis elliptico-oblongis glabris subserratis in alam profundè et acutè dentatam ferè incisam longè productis, ramis foliosis subpaniculatis apice mono-aut oligocephalis, pedicellis nudis, invol. squamis ext. oblongis foliaceis brevibus, intimis scariosis linearibus acutis flores paulò superantibus. — in Indiæ or. littore arenoso maris Coromandeliani frequens

ex Wight! herb. nec ab eâ differre videntur specimina à cl. Bojer in ins. Madagascar ad urbem Tananarivou lecta; alæ caulinae in iis variant nunc grosse nunc minutissime serratae. *Conyza alata* B. Wall.! cat. n. 3039. comp. 149. *Ethulia divergens* et *Inula indica* Russ.! herb. Valde accedit ad *B. alata* sed decurrentis fol. inciso-dentatis facile differt. (v. s.)

93. *B. aurita* (DC. in Wight contrib. p. 16), hirsuto-villosa, caule erecto tereti ramoso, foliis lyrato-pinnatifidis, lobis oblongis acutis dentatis, terminali majore, infer. basi decurrentibus secus caulem 1-2-auritis, paniculâ terminali laxâ, capitulis pedicellatis erectis, invol. squamis lanceolatis acuminatis, ext. dorso hirsutis, inter. subscariosis disco paulò longioribus. ① in Indiâ or. peninsula (Leach.!), frequens in agro Maderaspatao et Coromandeliano (Wight!), ad Surjaghora et Tukanaghur (Ham.!), in Pegu (Belang.!), etc. Con. aurita Linn. suppl. 367? Roxb.! Ham.! in Wall. cat. n. 3069 et 3086. Ex Linn. pars infima plantæ glabra, sed mihi ignota. Recept. nudum. Flor. masc. 5-8. Achænia glaberrima. Styli fl. fem. exserti. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

β. *Ægyptiaca*, caule folisque hirsutis subsericeis, capitulis subminoribus. — in Ægypto ad Hermonthium legit cl. Sieber, et in Sennaar cl. Acerbi. *Erigeron Chinense* Sieb.! fl. exs. non Jacq. An *Conyza villosa* Willd. sp. 3. p. 1929? (v. s.)

γ? *Berteriana*, tota piloso-subhirsuta vix subcinerea, foliis infimis latè obovatis. ① Hanc stirpem ex insulâ Americanâ Sanctæ-Marthæ ad Balbisiolum olim misit Bertero, sed ibi fortè culta aut introducta? An *B. aurita* aut *B. Senegalensis* var.? (v. s.)

94. *B. GUINEENSIS*, tota puberula subglandulosa, foliis oblongis acutis dentatis basi incisâ infernè secus caulem 1-2-auritis, paniculâ terminali laxâ, capitulis breviter pedicellatis erectis, invol. squamis lanceolatis acuminatis, ext. dorso puberulis, intim. subscariosis disco paulò longioribus. ① in Guineâ (Willd. Thonn.), in Senegaliz arenosis apricis humidis (Perr.!). *Conyza Guineensis* Willd. sp. 3. p. 1930. *Erigeron stipulatum* Schum. guin. p. 385. *Conyza argentea* Perr.! et Lepr.! ined. Nimis *B. aurita* affinis et fortè varietas? (v. s.)

β. *foliolosa*, squamis foliaceis linearibus innumeris secus pedicellos. — *Conyza Senegalensis* Sieb. fl. exs. n. 42. non Willd. Potius monstrum quàm varietas vera. (v. s.)

95. *B. SENEGALENSIS*, tota puberula subglandulosa, foliis obovatis obtusis dentatis basi attenuatis et incisâ subdecurrenti-auritis, paniculâ confertâ terminali, capitulis pedicellatis, invol. squamis lanceolatis acuminatis, exter. brevibus puberulis, int. subscariosis disco vix longioribus. ① in Senegaliâ olim legit Ronssillon, ad Gambiam recentius Don (h. Lindl.!). *Conyza Senegalensis* Willd. sp. 3. p. 1928. (v. s.)

CCXXI. *PLUCHEA* Cass. bull. philom. 1817. p. 31. non Arrab. fl. flum. (1835). — *Pluchea* et *Chlænobolus* Cass. dict. 42. p. 1. — *Conyza* et *Baccharidis* sp. auct. — *Pluchea* sp. Less. syn. 207.

Capitulum multiflorum heterogamum, fl. ambitus multiserialibus foemineis filiformibus truncatis aut 2-3-dentatis, centralibus paucis masculis aut subhermaphrodito-sterilibus 5-dentatis glabris. Recept. planum epaleaceum nudum aut hirsuto-fimbriliferum. Invol. pluri-seriale imbricatum. Antheræ basi bicaudatæ. Achæmium erostre cylindraceum aut sulcato-angulatum. Pappus uniserialis, setis filiformibus vix scabris. — Herbæ aut suffrutices in Americæ, Asiæ imò Africæ partibus calidis dispersi. Folia alterna, varia in speciebus diversis, sæpe pubescenti-glandulosa. Capitula corymbosa pedicellata. — Genus multifforme, in posterum fortè dividendum, à *Conyza* et *Blumea* antheris caudatis diversum. — Placus Lour. huc fortè revocandus, sed non satis notus.

Sect. 1. *STYLIMNUS*. — *Stylimnus* et *Gynema* Rafin. *jour. phys.* 1819. p. 100 et *ann. nat.* 1820. p. 15. — *Leptogyne* Ell. *sketch.* 2. (1824). p. 322. non Less.

Recept. nudum.

§. 1. *Foliis decurrentibus*. — *Species Americanæ et Gerontogææ*.

1. *P. MACROCEPHALA*, herbacea erecta simplex pilosa, caule striato, foliis longè decurrentibus, infer. ovali-oblongis serratis basi longè attenuatis, super. lanceolatis linearibusve integerrimis, capitalis 7-8 in corymbum terminalem dispositis, pedicellis patentim hirsutis, involacris tomentoso-lanatis. — in Brasiliæ prov. Rio-Grande (h. imp. Bras. n. 747). Herba $1\frac{1}{2}$ -2-pedalis. Capit. 9-10 lin. diam. Pappus copiosus subrufescens. Anth. longè caudatæ. (v. s. in h. Mus. reg. Par.)
2. *P. QUITOC*, herbacea, foliis sessilibus longè in alas foliaceas decurrentibus oblongis lanceolatisve callosio-denticulatis, corymbo composito subfastigiato, invol. squamis lanceolatis subacutis. ① in Brasiliæ campis arenoso-paludosis (Lund!), circa Bahiam (Blanch. Salzm.), Rio-Janeiro (Gaud.), in littore dicto Banda orientalis (Baile!), in prov. Rio-Grande, Mato-Grosso. Dicitur vern. *Quitoc* (Lund), *Quitogue* (Blanch.), *Quitoco* (h. Mus. Bras.). Flor. nov. dec. Gnaphalium squaevoles Arrab. fl. flum. 8. t. 100. Herba odore suavi aromatico, sapore amaro subcamphorato. Fl. invol. innumeri, marginales pallidè flavescentes, centrales violacei. Variat superficie glabrâ aut plus minùs pubescente, imò in vaginâ foliorum infer. canescenti-subvelutinâ; foliis oblongis lanceolatis aut subovatis acutis, rariùs obtusis, plus minùs dentatis; alis caulinis integris aut dentatis. Ex alis caulinis accedit ad *Pterocaulon* sed ex habitu toto et caract. capituli certè *Pluchea* pertinet. (v. s.)
3. *P. SUBDECURVENS* (Cass. dict. 42. p. 4), caule fruticoso erecto tereti supernè præsertim puberulo, foliis in alam subintegram obtusam brevem decurrentibus lanceolato-linearibus acutis à medio ad apicem serratis snprâ scabrisunculis subtus viscoso-punctulatis, corymbo terminali conferto polyccephalo, capitulis pedicellatis, invol. squamis lanceolatis acuminatis disco subæqualibus glabriunculis. ③ in Mexici salsis inter Istla et Acamistla (H. B. K.). *Conyza* asteroides Desf. 1 cat. h. par. 1815. p. 114. non Linn. nec Walt. *Conyza* adnata H. B. et Kunth nov. gen. am. 4. p. 74. *Baccharis* adnata Willd. enum. 870. Flor. lutei, form. pluriseriales truncati stylo exserto, masc. 5-10. Achænia glabra. (v. v. c. in h. Par.)
4. *P. OVALIS*, tota cinereo-puberula, caule suffruticoso ramoso tereti, foliis ovalibus ovatisve dentatis basi productis in alam inæqualiter dentatam, capitulis corymbosis. ⑤ in Senegaliâ olim leger. Geoffroy de Villeneuve, Rousillon et Sieber, in siccis ad Promont. Viride recentius Perrottet. *Baccharis ovalis* Pers. ench. 2. p. 424. n. 14. *Baccharis ovata* Sieb. fl. seneg. exs. n. 61. Invol. pallidi squamæ inter. apice subobtusæ ciliatæ. (v. s.)

§. 2. *Antheris brevius caudatis, foliis sessilibus non decurrentibus aut petiolatis*. — *Species Gerontogææ*.

5. *P. DIOSCORIDIS*, tota cinereo-puberula, caule herbaceo basi suffruticulosæ tereti apice ramoso, foliis sessilibus ovalibus acutis serratis basi attenuatis inciso-sublobulatis, lobulis stipulæ-formibus, capitulis corymbosis, invol. squamis acuminatis. ⑥ in Ægypto ad Alexandriam, Kahiram, Damiettam; in Nubiâ (Delil.) et in Syriâ? (J. Bauh.). *Baccharis Dioscoridis* Linn. amœn. ac. 4. p. 289. *Conyza Dioscoridis* Rauw. it. t. 54. *Baccharis Ægyptiaca* Forsk. (v. s. et v.)
6. *P. TOMENTOSA* (DC. in Wight contrib. p. 16), fruticosa, ramis teretibus ramulis et foliis utrinquè pube brevi vix tomentosâ griseo-velutinis, foliis obovatis ovalibusve cordato-seniamplexicaulibus dentatis, pedunculis ex axillis

- superioribus folio longioribus, capitula pedicellata conferta gerentibus, in paniculam confertam brevem dispositis, invol. squamis oblongo-linearibus acutis disco paulò brevioribus. ζ in Indiâ or. circa Mysore legit cl. Heyne. Conyza tomentosa Wall.! cat. n. 3031. comp. n. 141. Wight! herb. non Mill. nec Forsk. Eupatorium Mysorens Heyne! herb. Folia 12-15 lin. longa, 9-10 lin. lata. Flor. fem. innum. stylo vix exserto, masc. 5. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
7. *P. INDICA* (Less.! in linnæa 1831. p. 150), caule fruticoso tereti supernè corymboso ramoso, infernè glabro, ramulis pedicellis foliisque novellis minutè pulverulento-puberulis, foliis obovatis breviter petiolatis distanter et acutè serratis, corymbis ramos terminantibus confertis, invol. ovatis squamis ext. ovatis dorso puberulis, intimis linearibus acuminatis glabris. ζ in Ind. or. ad ripas Gangis, prope Chitagong et Penang (Wall.!), in Luzoniâ (Cham.!), Javâ (Bl.!); Timor (h. Mus. Par.), Borneo? (h. Merat). Baccharis indica Linn. sp. 1205. Willd. sp. 3. p. 1916. Conyza indica Blume! herb. Conyza corymbosa Roxb. mas. Wall.! cat. et herb. n. 3009. comp. n. 119 non Mill. — Breyn. cent. 1. 70. Flor. fem. innumeri, masc. 3-5. Antheræ breviter caudatæ. (v. s.)
8. *P. WALLICHIANA*, caule fruticuloso tereti ramoso foliis pedunculisque pube brevissimâ subpulverulentâ scabridis, foliis ovalibus obovatisve acutiusculis obtusisve serrulatis integerrimisve sessilibus, corymbis terminalibus trichotomia, capitulis breviter pedicellatis ovali-oblongis, invol. squamis lanceolatis acutis florum longitudine. ζ in Indiâ orient. ad Srinaghur. Conyza sessilifolia Wall.! cat. et herb. n. 3029. comp. n. 139 non Blume. Folia 1-2 poll. longa, 6-10 lin. lata. Fl. fem. innumeri tennes truncati stylo bifido, masc. circ. 5. Antheræ basi brevissimè caudatæ. Valdè habitu accedit ad Pluch. Indicam. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
9. *P. FOLIOLOSA*, glabra, caule fruticoso tereti ramoso, foliis ellipticis subacutis grossè et acutè paucidentatis breviter petiolatis, summis basi cuneatis sessilibus, corymbis terminalibus compositis confertis, invol. squamis inf. latè ovatis, super. linearibus acuminatis florum longitudine. ζ ad ripas fluminum Saluen et Caluen prov. Martabanæ Ind. or. (Wall.!) et in Pegu (Belang!). Conyza foliolosa Wall.! cat. n. 3030. comp. n. 140 non Willd. Eupatorium foliosum Wall. cat. Burm. n. 1565. Habitus Pluchæ Indicæ et Wallichianæ. Anth. brevissimè caudatæ. Flor. fem. innumeri tennes stylo bifido, masc. circ. 5. Ovaria glabrinsecula. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
- §. 3. *Antheris longiùs caudatis, foliis non decurrentibus sessilibus petiolatisve. — Species Americanæ.*
10. *P. OBLONGIFOLIA*, herbacea subfusco-pubescent subglutinosa, foliis sessilibus basi obtusè truncatis oblongis acutis serratis venoso-reticulatis, corymbo terminali subcomposito, invol. squamis dorso puberis glandulosis lanceolatis acuminatis disco longioribus. \mathcal{L} in Brasiliæ paludosis ad Rio-Janeiro et Bahiâ legit cl. Blanchet (n. 71. 72 et 1035!) et in prov. Sancti-Pauli cl. Lund! Affinis *P. bifronti*, sed notis indicatis facillè distincta. (v. s.)
11. *P. BIFRONS*, herbacea subglutinosa, foliis amplexicaulibus cordatis ovali-lanceolatis serratis nervoso-reticulatis, corymbo composito conferto, invol. squamis lanceolatis acutis dorso puberis subglandulosis. \mathcal{L} in humidis uliginosis et aggeribus Carolinæ. Conyza bifrons Linn. sp. 1207. Ell. sketch 2. p. 322. Conyza amplexicaulis Michx. fl. bor. am. 2. p. 126 non Willd. Baccharis viscosa Walt. fl. car. p. 202. Conyza uliginosa Pers. ench. 2. p. 427. — Plukn. alm. 1. 87. f. 4. Cor. sordidè purpurascens ut congeneres, nec pallidè flavæ ut vult Pursh. Per hiemem in horis matutinis, si cælum serenum, fibræ cristallinæ pollicem longæ è basi caulis oriuntur! (Ell. l. c.). (v. s.)
12. *P. MARYLANDICA* (DC. non Cass.), herbacea minutè pubescens subglutinosa, foliis sessilibus lanceolatis acutis subserratis membranaceis uninerviis, corymbo composito fastigiato polycephalo subaphyllo, capitulis pedicellatis, invol.

squamis lanceolatis acutis florum disco brevioribus. ① in humidis et inundatis Amer. bor. à Noveboraco ad Carolinam (Pursh fl. 523), Novâ-Cesareâ (Torr.!). *Conyza Marylandica* Michx. fl. 2. p. 126. Ell. sketch 2. p. 321. *Baccharis foetida* Walt. car. 202? *Stylimnus maritimus* Raf. in litt. 1819. Sæpe in hortis vagat sub nom. *Erig. camphorati*. (v. s.)

13. *P. FORTIDA*, herbacea glabrescens, foliis petiolatis ovali-lanceolatis utrinque acuminatis grossè serratis penninerviis membranaceis, corymbo fastigiato subpaniculato, capitulis pedicellatis, inv. squamis lanceolatis acutis dorso subpubescentibus eciliatis. ② in Virginiâ (Linn.), in Kentucky (Raf.). *Baccharis foetida* Linn. sp. 1205. *Gynema dentata* Raf. in litt. 1819. *Gynema viscida* Raf. ann. nat. 1820. p. 15. *Conyza marylandica* Pursh fl. 523? *Pluchea marylandica* Cass. dict. 42. p. 2 non DC. Dill. elth. t. 89. f. 105. (v. s.)
14. *P. CAMPHORATA*, herbacea subpubescens, foliis petiolatis ovatis seu ovato-lanceolatis acutis subrepando-dentatis membranaceis penninerviis subtus granulata resinosa gerentibus, corymbo composito fastigiato, capitulis pedicellatis, invol. squamis membranaceis ciliatis ovato-lanceolatis acutis dorso pubescenti-glandulosi. ① circa Bostoniam (Greene!), in Pensylvaniâ nec in Carolinâ et Georgiâ (Ell.). *Erigeron camphoratum* Linn. sp. 1212. Willd. sp. 3. p. 1960. *Conyza camphorata* Ell. sketch 2. p. 321 non Pursh? *Pluchea petiolata* Cass. dict. 42. p. 2. *Gynema balsamica* Raf. (v. s.)
15. *P. PURPURASCENS*, herbacea pubescens, foliis petiolatis lanceolato-ovatis acutis serratis subtomentosis, corymbo terminali composito, capitulis pedicellatis ovato-globosis, invol. squamis ext. ovatis ferrugineo-pubescentibus, int. ovato-lanceolatis patentibus. ① in subhumidis et depressis Jamaicae. *Conyza purpurascens* Swartz fl. ind. occ. 1342. — Sloan. hist. 1. p. 152. f. 1 malè. Brown. jam. 318. Affinis dicitur *P. odorata*, sed herbacea serratifolia, etc.
16. *P. GLABRATA*, herbacea glabra aut vix subpubescens, foliis petiolatis lanceolatis basi longè acuminatis apice acutis serratis penninerviis, corymbis compositis laxis, capitulis pedicellatis subrotundis, invol. squamis ovato-lanceolatis acutis. ① in Guadalupâ legit olim cl. Badier et recentius Perrottet ad Pointe-à-Pitre. An var. glabra *P. purpurascens*? An *Conyza odorata* Guadalupensis Spreng. syst. 3. p. 507 ad hanc aut ad *P. odoratam* referenda? (v. s.)
17. *P. ODORATA* (Cass. dict. 42. p. 3), fruticosa pubescenti-velutina, foliis petiolatis ovalibus oblongisve utrinque attenuatis, integerrimis subtus subcanescentibus, corymbo composito fastigiato polycephalo, capitulis pedicellatis, invol. squamis ext. ovatis dorso pubescentibus, inter. linearibus acuminatis scariosis glabresculis. ② in Americâ merid. (Linn.), Mexico ad Tampico de Tamaulipas (Berl. pl. exs. n. 3) et in distr. Oaxacamo (Andr. pl. exs. n. 275), in Guadalupâ (Badier!). *Conyza odorata* Linn. sp. 1208. — Plum. ed. Burm. t. 97 (fol. separ.), sed nunquam vidi folia inciso-dentata. Folia tacta odorem subsuavem exhibent. Ab hac nullo modo differre videtur *Conyza Carolinensis* Jacq. ic. rar. 3. t. 585. coll. 2. p. 271. Willd. sp. 2. p. 271, in insulis Bahamensibus spontanea, sed verisim. excl. habit. in Carolinâ et Floridâ. (v. v. c. et s. sp.)
18. *P. CORTESII*, suffruticosa ramosissima canescenti-hirta, foliis petiolatis oblongis acutis basi angustatis integerrimis canescentibus, corymbis terminalibus ramosis, capitulis breviter pedicellatis, inv. squamis imbricatis oblongo-lanceolatis apice subulatis. ③ prope villam Ferd. Cortesii et urbem Cuernavaca Mexicanorum. *Conyza Cortesii* H. B. et Kunth nov. gen. am. 4. p. 75. Folia 3 poll. longa, 10 lin. lata. Flor. disci circ. 20. Affinis dicitur *Pluch. odorata* et ad *Plucheam* idèò referenda? quod in specim. Kunthiano confirmavit ocul. Decaisne.
19. *P. CHINGOYO*, fruticosa, ramis hirtis pubescentibus, foliis subrotundo-ellipticis utrinque obtusis integerrimis suprâ hirtis pubescentibus subtus molliter pubescentibus canescentibus, corymbis terminalibus ramosis, invol. hemisphaerici squamis glabris linearibus apice subulatis. ④ in locis planis prope urbem Santa Peruvianorum ubi dicitur *Chingoyo*. *Conyza Chingoyo* H. B. et Kunth

nov. gen. am. 4. p. 76. t. 328. Ex habitu ad Plucheam referenda videtur, cl. Decaisne reverâ ex specim. Kunthiano hanc suspicionem confirmavit.

N. B. Placus tomentosus Lour. fl. coch. 2. p. 606. Placus lævis Lour. fl. coch. 2. p. 606. Gynema balsamica Raf. fl. lud. 63. Gynema? argentea Raf. fl. lud. 63 et Gynema? parviflora Raf. fl. lud. 64 sunt species fortè huic sectioni addendæ sed omninò ignotæ.

Sect. II. HEBEPHORA.

Recept. hirsutum. Caules herbacei. Folia sessilia non decurrentia. — Species Asiaticæ, mediâ inter Plucheam veram et Pterocaulon, à priori recept. hirsuto, à poster. invol. persistente et inflor. corymbosâ diversæ.

20. *P. HIRUTA* (Less. in linnæâ 1831. p. 150), caule herbaceo erecto tereti hirsuto, foliis sessilibus lanceolatis basi acuminatis apice subserratis suprâ lucidis asperis subtus subhirsutis, capitulis spicato-glomerulatis, invol. cylindrici squamis linearibus acuminatis hirsuto-ciliatis, recept. hirsuto-fimbrillifero, corollis glabris, antheris longè caudatis. — in Chinâ (Torr. ex Linn.), in Cochinchinâ? (Lour.), in Luzoniâ (Cham. ex Less.). *Conyza hirsuta* Linn. sp. 1209. An plantæ omnes hic recensitæ omniâ eadem?

21. *P. SCABRIDA*, suffruticosa, ramis teretibus confertè pulverulento-scabris, foliis subsessilibus basi attenuatis obovatis grossè et irregulariter dentatis utrinquè sparè pulverulento-scabris, corymbo terminali composito, capitulis ad apices ramorum sessilibus fasciculatis, invol. ovati squamis ext. ovalibus, inter. linearibus glabris apice subciliatis. ♂ ad Luzoniam legit cl. Hænke. Proxima *P. hirsutæ* Less. sed foliorum formâ et capitulorum dispositione diversa. Recept. hirsutum. Cor. masc. glabræ apice tuberculatæ. (v. s. in h. Hænke à cl. de Sternberg miss.)

† Species exclusa.

P. PYCNOSTACHYA Less. = *Pterocaulon pycnostachyum*.

CCXXII. *PTEROCAULON* Ell. sketch 2 (1824). p. 323. — *Chlænobolus* Cass. 49. (1827) p. 337. — *Pluchea* sp. Less. — *Conyza* sp. Lam.

Capitulum multiflorum, fl. marginalibus fœmineis filiformibus 3-dentatis pluriserialibus. Recept. tenuissimè fimbrilliferum hirsutum, rariùs nudum. Invol. oblongi squamæ imbricatæ pluriseriales adpressè aut apice summo vix squarrosæ caducissimæ! Antheræ basi caudas subulatas gerentes subexsertæ. Achænia angulata adpressè pilosa. Pappi setæ filiformes pluriseriales inter se æquales involucri longiores. — Herbæ interdum basi suffruticosæ Americanæ. Rhizoma subtuberosum sæpè lignosum. Caulis tomentosus alis sex foliaceis ornatus. Folia alternâ lanceolata, integrâ aut denticulata, suprâ glabra aut juniora araneosa, subtus densè tomentosa, utrinquè producta in alas longas integras. Capitula sessilia glomerata in spicas thyrsosve spicatos digesta.

Sect. I. CHLÆNOBOLUS.

Recept. hirsutum. Folia subtus, caules et pagina alarum cum inferiore folii contigua densè et cano-tomentosa.

1. *P. PYCNOSTACHYUM* (Ell. sketch 2. p. 324), caule herbaceo simplici, foliis lanceolatis subundulatis minutè et remotè denticulatis suprâ glabris, capitulis in spicam densam continuam confertis, invol. squamis extus sericeo-tomentosis apice squarrosis. ♀ in arenosis siccis campis et sylvis Carolinæ et Floridæ (Pursh, excl. syn. Cav.). *Conyza pycnostachya* Michx. fl. 2. p. 126. *Gnaphalium undulatum* Walt. fl. car. 203. *Chlænobolus pycnostachyos* Cass. dict. 49. p. 338. *Pluchea pycnostachya* Less. Radix tuberosa vulgò Black-Root dicta *placera vetera sanat.* (v. s.)

2. *P. alopecuroides*, basi suffrutescens simplex, foliis ellipticis irregulariter et calloso-crenato-serratis supra junioribus arachnoides adultis glabris, capitulis in thyrsium spiciformem densum aggregatis, infer. hinc inde interruptis, invol. squamis dense tomentosis apice acutis glabris squarrosis. ζ in sabulosis aridis Jamaica australis (Sw.), Guadalupæ (Bert.!) et Martinicæ (h. Duby!). *Conyza alopecuroides* var. α . Lam. dict. 2. p. 93. *Conyza alopecuroides* Swartz fl. ind. occ. 1344. *Chlænobolus alopecuroides* Cass. dict. 49. p. 339. (v. s.)

β . *polystachyum*, caule apice ramoso, spicis thyrsoides plurimis. ζ in insul. Caribæis? (v. s. comm. à cl. Lambert.)

3. *P. spicatum*, suffrutescens simplex ramosumve, foliis oblongo-aut elliptico-lanceolatis acutis irregulariter dentatis subrugosis capitulis thyrsoides-spicatis nempe in glomerulos spicatis aggregatis, invol. squamis sericeo-tomentosis squarrosis, inter. subglabris. ζ in Brasiliâ circa Rio-Janeiro et Bahiâ. An fortè *Conyza alopecuroides* Kunth prope Cumanam lecta huc referenda?

α . *brachystachyum*, thyrsos spicato ovato denso continuo. ζ frequens circa Rio-Janeiro et in prov. Sancti-Pauli (Lund!). *Conyza rugosa* Vahl symb. 1. p. 71. *Chlænobolus rugosus* Cass. dict. 49. p. 341. Habitus *P. alopecuroides* et forsan species duas conjungens. (v. s.)

β . *cylindricum*, thyrsos cylindrico elongato continuo. ζ in Amer. merid. (Lam. Cav.). *Conyza spicata* Lam. dict. 2. p. 93. Cav. ic. 1. t. 12. *Chlænobolus spicatus* Cass. l. c. (v. s.)

4. *P. interruptum*, suffrutescens, foliis oblongo-lanceolatis irregulariter dentatis, capitulis fasciculatis sessilibus, fasciculis in thyrsium spicatum cylindrico-elongatum interruptum digestis, invol. squamis linearibus, ext. dorso tomentosis subacutis, inter. elongatis acuminatis apice glabris. ζ \mathcal{U} in Brasiliâ circa Bahiam (Salzm.!) Rio-Janeiro (Gandich.!), prov. Rio-Grande (h. mus. Bras!)

α . *polystachyum*, ramis plurimis elongatis apice thyrsoides-spicatis. (v. s.)

β . *monostachyum*, caulibus simplicibus apice thyrsoides-spicatis. An species propria ex inv. brevioribus et thyrsos minus interrupto. (v. s.)

5. *P. angustifolium*, herbaceum gracile ramosum, foliis linearibus acutis integerrimis, capitulis sessilibus in spicam subrotundam brevem aphyllam confertissimis, invol. squamis inter. linearibus acuminatis glabris, flor. hermaphrod. 2-3. \mathcal{U} in Brasiliæ prov. Sancti-Pauli. Folia sesquipoll. longa, 2 lin. lata, supra juniora araneosa, adulta glabra. Capitula 15-20 oblonga in glomerulum sphericum conferta. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 431 miss.)

6. *P. virgatum*, herbaceum simplex, foliis lineari-lanceolatis vix serrulatis acutis, capitulis glomeratis, glomerulis spicato-thyrsoides valde remotis, invol. squamis dense tomentosis mucrone acuto glabro. ζ \mathcal{U} ? in insulis Caribæis nempe Jamaica (Sloan.), Portorico et Sancti-Thomasii (Bert.!), nec in Carolinâ. *Conyza virgata* Lin. sp. 1206 Lam. dict. 2. p. 93. *Gnaphalium virgatum* Lin. amcn. 5. p. 405. *Chlænobolus virgatus* Cass. dict. 49. p. 340. — Plum. ed. Burm. t. 98. f. 2. Sloan. hist. 1. p. 206. t. 152. f. 5. (v. s.)

Sect. II. PTEROCAULOPSIS.

Recept. nudum nec hirsutum. Invol. squamæ subæquales vix imbricatæ glabræ et folia subtus non cano-tomentosa. — Herba basi suffruticosa foliis Pluchææ, inflorescentiâ Pterocauli, fortè cum Monentelis sectione primâ genus constituens.

7. *P. polystachyum*, suffruticosum erectum puberulo-hirtellum, foliis lanceolatis seu oblongo-lanceolatis acutis dentato-serratis reticulato-nervosis in alas foliaceas dentatas aut subintegras secus caulem decurrentibus, ramis floridis plurimis paniculato-corymbosis, capitulis oblongis sessilibus fasciculatis secus rhachin thyrsoides-spicatis. invol. squamis linearibus acuminatis glabris. ζ in Brasiliæ prov. Rio-Grande. Cor. ferè Pluchææ, sed folia decurrentia et apicæ thyrs-

soides Pterocauli. Fl. disci 2-3, form. innumeri, omnes longitudine involucri. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub. n. 861 miss.)

CCKXIII. MONENTELES Labill. sert. caled. p. 43. Cass. dict. 53. p. 236. — Tessariæ sp. Less. syn. 207.

Capitulum pluriflorum, fl. marginalibus 2-3-seriatis fœmineis filiformibus apice subdentatis, disci 1-5 hermaphroditis seu styli abortu masculis subcylindricis 5-dentatis extus subglandulosus. Recept. nudum planum angustum. Involucri oblongi squamæ biseriatae persistentes subæquales. Antheræ disci caudis subulatis donatæ. Styl. disci et radii apice bifidus. Achænia oblonga subcompressa parcè villosa. Pappus 1-serialis, setis filiformibus imâ basi subconcretis vix invol. longioribus. — Herba Filagineæ ramosæ. Folia oblonga seu lanceolata denticulata, in alas secus caulem longas integerrimas producta. Capitula sessilia, ad axillas foliorum floralium fasciculata, in spicas interruptas disposita. — Genus ad Pterocaulon proximè characteribus et habitu accedens, sed satis diversum videtur invol. persistente non imbricato, recept. nudo, floribus disci paucioribus imò solitariis, ovario subcompresso et pappo 1-seriali.

§. 1. Floribus disci 3-5. — An potiùs Pterocauli sectio?

1. M. PTEROCAULON, basi suffruticosus totus cinereo-villosus, caule erecto ramossissimo, foliis oblongo-linearibus distanter denticulatis secus ramos in alas longas foliaceas integerrimas productis, capitulis ad axillas supremas et apices ramorum sessilibus congesto-fasciculatis. § in agris circa urbem Tananarivou metrop. prov. Emirnæ insul. Madagascar legit cl. Bojer. Baccharis Pterocaulon Bojer! in lit. 1833 cum icon. Habitus omninò Monentalis. An fortè Conyza decurrens Lin. sp. 1206 eadem aut affinis? (v. s. comm. à cl. inv.)

§. 2. Flore disci solitario.

* Capitulis in glomerulos spicatum dispositos aggregatis.

2. M. SPICATUS (Labill. ! sert. nov. cal. 43. t. 43), caulibus herbaceis dense tomentosis alatis, foliis longè decurrentibus oblongis serrulatis suprâ araneosis subtus dense lanuginosis, capitulis in spicas elongatas dispositis, squamis invol. omnibus acuminatis, exter. brevioribus lanatis. ¶ in Novâ-Caledoniâ (Labill. !) et in Indiæ montibus regni Avenis ad Taong-dong, Paghamew, alibique ad ripas fl. Irrawaddy (Wall. !). M. spicatus Cassini dict. 53. p. 237. Gnaphalium cylindrostachyum Wall. ! cat. n. 3931. comp. n. 41. Gn. polycephalum Wall. ! cat. biru. n. 655. Recept. nudum. (v. s.)
3. M. REDOLENS, caule lignoso ramoso toto alato, foliis longè decurrentibus lanceolatis integerrimis suprâ pubescentibus subtus leviter cano-tomentosis, capitulis in spicas terminales glomeratis. § in Novâ-Caledoniâ (Forst. !) et in Luzoniâ (Cham. !) Gnaphalium redolens Forst. prod. n. 533. Conyza redolens Willd. sp. 3. p. 1951. Tessaria redolens Less. in linnæa 6. p. 151.

** Capitulis in glomerulum globosum dispositis.

4. M. GLOBIFRONS, caulibus suffruticulosus, foliis lanceolatis acutis reticulato-nervosis utrinquè pubescenti-hirsutis in alas angustissimas productis, glomerulis globosis terminalibus, invol. squamis exter. brevibus hirsutis apice maculatis subtruncatis, inter. elongatis linearibus acutis glabris, flor. fœm. capituli cujusque 10-12. § in paludosis sylvaticis ad fluvium Brisbane in sinu Moreton legit cl. A. Cunningham. Affinis M. sphacelato. (v. s. comm. à cl. Cunningham.)
5. M. INTERMEDIUS, caulibus suffruticulosus hirsutus, foliis elliptico-lanceolatis

mucronato-acutis reticulato-venosis utrinque hirsutis in alas latiusculas integerrimas longè productis, glomerulis terminalibus globosis, invol. squamis ext. hirsutis apice ustulatis, inter. elongatis glabris, floribus fœm. circ. 10. 5 in arenosis littoribus ad Caput-Cleveland Novæ-Cambiæ aust. inter tropicos. Species media inter *M. globiferum* et *M. sphacelatum*, à priore alis caulinis duplò triplòve latioribus et longioribus, superficie magis hirsutâ, foliis bracteantibus majoribus, à sequente foliis acutis et fl. fœmineis paucioribus diversa. (v. s. comm. à cl. Cunningham.)

6. *M. SPHACELATUS* (Labill. sert. n. cal. 43. t. 44), caulibus herbaceis alatis, foliis longè decurrentibus oblongis obtusis, capitulis in glomerulos laterales terminalesve globosos aggregatis, invol. squamis exterioribus ad apicem ustulatis. 2? in Novâ-Caledoniâ. *Monenteles sphacelatus* Cass. dict. 53. p. 237. Flos centr. ex. Cass. fertilis, fl. fœm. circ. 25. Recept. pilos lanatos paucos gerens.
7. *M? SPHÆRANTHOIDES*, caule fruticuloso glabro, foliis lineari-lanceolatis elongatis margine subrevolutis rugosis penninerviis denticulatis in alas dentatas decurrentibus glabriusculis, pedunculis pubescenti-hirtellis subscabris, glomerulis globosis terminalibus. 5 in insulâ Enderby Archipelagi Dampieriani ad oram bor. occid. Australiæ legit cl. A. Cunningham. *Sphæranthoides* Cunningham! in litt. Frondescencia fere *Sphæranthi*, capitulum *Monentelis*, sed in specim. equidem malè conservato achænia omninò calva! An pappus deciduus, an verè nullus? Invol. squamæ ext. breves hirsutæ, inter. elongatæ glabræ. Species distinctissima! Genus dubium! (v. s. comm. à cl. peregr.)

CCXXIV. *TESSARIA* Ruiz et Pavon prod. p. 112. t. 24. syst. p. 213 Cass. dict. 53. p. 233. — *Tessaria* sp. et *Gynheteria* Less. syn. 207 et 208. — *Gynheteria* Willd. mag. nat. Berl. 1807. p. 140. — *Gynheteria* Spreng. neu. entd. 2. p. 135.

Capitulum multiflorum, fl. marginalibus pluriserialibus tenuissimis truncatis seu 2-3-dentatis fœmineis, centralibus 1-7 majoribus hermaphroditis aut styli abortu masculis, cor. apice dilatatâ 5-fidâ. Recept. tenuiter fimbrilliferum hirsutum. Invol. oblongi squamæ imbricatæ scariosæ, ext. breves persistentes ciliatæ, inter. elongatæ acutæ deciduæ subintegerrimæ. Antheræ basi brevissimè caudatæ. Stylus disci indivisus undiquè subhispidulus. Fl. fœm. exsertus glaber bifidus. Achænia brevia teretiuscula glabra. Pappus 1-serialis, setis filiformibus. — Frutices Americæ australis incolæ, foliis ramulisque adpressè cano-villosis. Folia alterna ovali-oblonga aut lanceolata integra aut in iisdem indiv. plus minus dentata. Corymbi terminales ramosi. Cor. disci purpureæ, fœmineæ pallidæ.

§. 1. *Flore hermaphrodito-sterili unico centrali.*

1. *T. LEGITIMA*, foliis ramulisque hirsuto-canescens ovali- aut subobovato-oblongis vix acutis, invol. squamis ext. ciliis subhirsutis fimbratis. 5 in Peruvîâ ubi dicitur *Pazaro-Bobo*. *Tessaria* Cass. dict. 53. p. 233. Variat foliis integerrimis aut sinuato-subdentatis: ergò certè videtur *T. integrifolia* R. et Pav. syst. p. 213 et fortè etiam *T. dentata* R. et Pav. p. 214. *Gynheteria* incana Willd. mag. nat. berl. 1807. p. 140? Spreng. neu. entd. 2. p. 135. t. 1. f. 5-11 ex icone, sed excl. patr. et descr. part. (v. s. ex itin. Dombey, Née et Gaudichaud.)
2. *T. MUCRONATA*, ramis glabriusculis, foliis adpressè serieo-incanis ellipticis basi cuneato-acuminatis apice mucronato-acutis, aliis integerrimis, aliis apice paucidentatis, invol. squamis glabris margine vix ac ne vix subciliatis. 5 seu 5 ad Baranquilla Novo-Granatensium propè ostia fl. Magdalenæ legit cl. Bertero. *Gynheteria* incana è Magdalenâ Spreng. syst. 3. p. 482 (excl. syn. et aliis habit.) ex specim. ab ipso auct. ad Balbisium misso! An fortè *Conyza*

riparia Magdalenensis H. B. et Kunth nov. gen. am. p. 4. 76? (ex quâ cl. Cassini genus suum Phalacromesum dict. 53. p. 235. 57. p. 339. 60. p. 586 constituit) sed certè nostræ flos centralis papposus nec nudus. (v. s. comm. à cl. Balbis et Bertero.)

§. 2. *Floribus disci 3-8 hermaphroditis seu abortu masculis.*

3. T. *AMAREUA*, ramulis foliisque adpressissimè pulverulento-canis, foliis lanceolatis subobtusis basi in petiolum attenuatis sinuato-paucidentatis, invol. squamis exter. ovato-rotundis ciliatis, inter. linearibus scabris glabris, disco 3-floro. ♀ in America merid. fortè in Peruvia. Habeo ex itin. Neæano sub nom. vernaculo *Pajro-Bovo*, vix à nom. primæ generis speciei distincto.
4. T. *ABSINTHIODES*, ramis foliisque adpressissimè canis subargenteis, foliis lanceolatis utrinquè acuminatis, nunc integerrimis, nunc dentibus grossis acutis hinc indè incisis, subtus nervoso-reticulatis, invol. squamis glabris acutis, disco 7-8 floro. ♀ in Chili ad Valparaíso, Coquimbo, et Conceptionem secus flumina. Vulgò ex Bert. dicitur *Bréa*. Baccharis absinthioides Hook. et Arn. bot. Beech. p. 57. Baccharis banksiæfolia Bert. herh. Gynheteria incana Chilensis Spreng. syst. 3. p. 482. (v. s. comm. à cl. Bertero, D'Urville, Gaudich.)

† *Species exclusa.*

T. *REDOLENS* Less. = *Monenteles redolens*.

CCXXV. MONARRHENUS Cass. bull. philom. 1817. febr. dict. 32. p. 433. 53. p. 236. — Mahometa DC. 1802. mss. — Conyzæ sp. Lam. — Tessariæ sp. Less.

Capitulum pluriflorum heterogamum, fl. omnibus tubulosis, ambitus plurimis subbiseriatis filiformibus 3-4-lobis fœmineis, disci solitario 5-dentato hermaphrodito aut ovarii abortione masculino. Recept. nudum planum parvum. Invol. ferè campanulatum floribus paulò brevius imbricatum, squamis oblongis obtusis apice summo subtomentosis. Cor. disci extus subglandulosa. Antheræ breviter caudatæ longè appendiculatæ. Stylus disci ramis inclusis brevibus, fl. fœm. longis exsertis filiformibus. Achænia fl. fœm. oblonga cylindracea supernè subhispidula callo basilari crasso. Pappus setis 1-serialibus scabris. — Frutices Mauritanii. Folia alterna linearia aut sublanceolata integerrima, suprà scabrida, subtus tomentosa aut sericea. Corymbus compositus. Capitula ad apices pedunculorum conferta subsessilia. Flores lutei. — Genus affine Tessariæ et Monenteli, ob flor. ♂ solitarium inter plurimos ♀ olim à me Mahometa dictum.

1. M. *SALICIFOLIUS* (Cass. l. c.), foliis breviter petiolatis lanceolato-linearibus subtus sericeo-tomentosis penninerviis. ♀ in insulis Mauritiæ ad montem Pollicis, et Borboniæ ad montem Bernica (Bout!). Conyza salicifolia var. α. Lam. dict. 2. p. 89. Bory! voy. isl. afr. 2. p. 28. Folia omnino Salicis viminalis, 2-3 poll. longa, 4 lin. lata. Vulgò *Bois de Senil* et *Bois de Chenille* dicitur. (v. s.)
2. M. *PINIFOLIUS* (Cass. ll. cc.), foliis angustè linearibus sessilibus margine subrevolutis uninerviis subtus tomentosis. ♀ in insulâ Borboniæ. Conyza salicifolia var. β. Lam. dict. 2. p. 89. Conyza salicifolia angustifolia Pers. ench. 2. p. 428. Vulgò *Bois de Senil de Bourbon*. Folia 2-3 poll. longa, semi-lin. lata. Priori valdè affinis sed distinctus videtur et nulla specimina intermedia observantur. (v. s.)
3. M. *RUFSCENS*, foliis sessilibus lineari-lanceolatis planis penninerviis utrinquè ramulisque pube rufâ minimâ ferrugineis integerrimis aut hinc indè argutè et minutissimè serratis. ♀ in insulis Mauritanis nempe in insulâ Borboniæ (Comm.) in insulâ Mauritiæ (Bout. in h. Lindl.). Conyza heliotropifolia Lam. dict. 2. p. 90? (v. s. in h. Lindl.)

CCXXXVI. CYLINDROCLINE *Cass. bull. philom. 1817. jan. dict. 12. p. 318. Less. syn. 207.* — *Lepidopogon Tausch in florâ 1829. p. 2. p. 37. et 1831. p. 1. p. 224.*

Capitulum pluriflorum heterogamum, flor. omnibus tubulosis, marginalibus in parte recept. spiciformi seu inter involucri squamas sitis pluriserialibus foemineis filiformibus, disci in parte recept. terminali sitis paucis abortu styli subsimplicis masculis. Recept. totum bracteo-latum et spiciforme apice subconcauum. Invol. oblongum, squamis secus pedicelli apicem spicæ more dispositis imbricatis oblongis, infer. sterilibus apice ad dorsum tomentosis, superioribus in axillâ floriferis fimbriato-ciliatis. Achæmium oblongo-cylindraceum basi attenuatum subarcuatum 8-costatum pilosiusculum. Pappus 1-serialis, paleis axi corneis pectinato-barbellatâ. — Fruticulus Mauritanus ramis crassis basi glabris inter folia dense et longè sericeo-hirsutis. Folia ad apices ramorum conferta rosulata integra ovalia basi attenuata et vix idè subspathularia juniora, suprâ et demùm ad basin sericeo-hirsuta, ad apicem breviter strigilloso-sabra, subtus albo-tomentosa ad nervos et basin sericeo-hirsuta. Corymbus compositus confertus, ramis longè sericeo-hirsutis aphyllis folia vix æquantibus. — Est quasi *Evax* arborescens!

1. *C. COMMERSONII* (Cass. II. cc.). 3 in insulâ Mauritiâ montibus propè montes Pollicis et *Piton de la rivière noire* dictis (Boj.!). *Lepidopogon gnaphalodes Tausch in florâ 1831. Conyza hirsuta Commers. herb. Conyza Sieb. fl. maur. exs. 2. n. 334. Lepidopogon Ponæ Tausch in florâ 1829. c. 2. p. 37 et 1831. c. 1. p. 224. Flores cœruleæ! Species valdè singularis identitatem squamarum involucri et palearum receptaculi optimè demonstrans! (v. s.)*

CCXXXVII *EVAX* *Gærtn. fr. 2. p. 393. t. 165. Lam. ill. t. 694. Pers. ench. n. 1842.* — *Filago Willd. sp. 3. p. 2387. Cass. dict. 17. p. 2.* — *Gnaphalium Paill. act. ac. Par. 1719. p. 314.* — *Filaginis sp. Linn. — Micropi sp. Desf.*

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, radii multiserialibus foemineis gracilibus, disci paucis masculis 4-crenatis. Receptaculum elongatum subulatum, apice inter fl. masc. nudum, paleolas scariosas acuminatas inter. fl. foem. gerens. Invol. 1-2-seriale, squamis adpressis scariosis ovalibus aristato-acuminatis. Achænia (radii) obcompressa obovalia glabra calva. — Herba Europæ australis et Orientis indigenæ, minimæ, subcaules, tomento plus minù copioso canescentes. Folia alterna circa capitulorum glomerulum rosulata. Capitula plurima conferta subumbellata, medio majore.

Sect. I. *EUEVAX.*

Invol. squamæ et recept. paleæ planæ scariosæ abruptè aristato-acuminatæ. Capitula dense umbellata, bracteis foliaceis inter se nullis, foliis floralibus circa umbellam radiantibus.

1. *E. PYGMÆA* (Pers. ench. 2. p. 422), foliis floralibus obovatis obtusis. (1) in apricis et rupestribus maritimis Europæ mediterraneæ. Lam. ill. t. 694. f. 1. *Filago pygmæa* Linn. sp. 1311. *Filago acanthis* All. ped. n. 620. *Gnaphalium pygmæum* Lam. dict. 2. p. 761. *Evax umbellata* Gærtn. l. c. *Micropus pygmæus* Desf. atl. 2. p. 307. — Barr. ic. t. 127. n. iv. Variat caule simplicissimè verè pygmæo subnullo et caule à basi ramoso multicipiti longiore ferè pollicari. (v. et s.)
2. *E. EXIGUA*, foliis floralibus oblongis acutis vix glomerulo longioribus, invol.

squamis exter. dorso lanatis. ① in apricis et rupestribus Hispaniæ circa Madritum (Lag.!), in Siciliâ meridionali (Guss.!), in archipelagi insulis (Sibth.), in rupibus vulcanicis novæ insulæ Camini prope Theram (D'Urv.), et in agro Byzantino (Cast.!). etiam habeo inter plantas itin. Brousson. sed locus non indicatus. Filago exigua Sibth. fl. græc. t. 922. ined. ex prod. 2. p. 207. Micropus exiguus D'Urv. enum. p. 109. n. 802. Filago pygmæa Cav. ic. 1. p. 23. t. 36. non Linn. Variat ut prior caule simplici humillimo et à basi multicauli imò ramosissimo 2-4 poll. longo. Folia valdè cano-lanata. Omnino media inter priorem et sequentem. (v. s.)

3. E. ASTERICIFLORA (Pers. ench. 2. p. 422), foliis floralibus oblongis acutis radiantibus capitulorum glomerulo multò longioribus. ① in apricis et rupestribus Hispaniæ (Lam. ex Vahl's specim.), Sardiniae (Moris!), Siciliæ (Guss.!), Calabriæ (Duf.), et Orientis (Labill.) Lam. ill. t. 694. f. 2. Gnaphalium astericiflorum Lam. dict. 2. p. 760. — Moris. ox. s. 7. t. 11. f. 5. Variat etiam caule simplici 2-pollic. et ramoso, sæpiùs ad apicem ramosiore, 3-4 poll. longo. Folia glomerulum circumdantia 9-10 lin. longa. (v. s.)

Sect. II. PSEUDEVAX.

Invol. squamæ exter. lanatæ, inter. et recept. palæ oblongæ chartacæ obtusæ. Capitula sparsa aut congesta, bracteis foliaceis intermixtis et foliis floralibus irregulariter circa glomerulum dispositis. — Habitus Filaginis.

4. E. DISCOLOR, foliis obovato-lanceolatis, floralibus vix involucrium superantibus, capitulis congestis lanâ floccosâ basi circumdatis. ① in Siciliæ montibus Nebrodensibus (Guss.!). Filago discolor Guss. in litt. 1831 et 1834. Herba 1-2-pollicaris à basi ramosa. (v. s. comm. à cl. Guss.)
5. E. MARROTTICA, foliis lineari-oblongis, floralibus vix capitulo longioribus, capitulis axillaribus unilateralibus imbricatis. ① prope Alexandriam ad lacum Mareotidem. Filago Mareotica Delil. fl. eg. 130. t. 47. f. 2. Micropus Mareoticus Spreng. syst. 3. p. 499. Habitus ferè Filaginis montanæ sed caract. omnino Evacis.
6. E. PROLIFERA (Nutt. in litt. 1825), caule solitario erecto apice capitula inter folia glomerata gerente, ramis subglomerulo ortis ipsis apice capitula glomerulata gerentibus, foliis lineari-oblongis, floralibus conferto-subverticillatis capitula superantibus ① in Americâ bor. ad Red-River detexit cl. Nuttall. Herba stricta 2 $\frac{1}{2}$ -3 poll. longa tota villosa-canescens. (v. s. comm. à cl. Nutt.)
7. E. MULTICAULIS, caulibus è collo plurimis subdiffusis erectiusculis ramosis, foliis oblongis obtusis laxè ramisque araneoso-lanatis, summis circa capitula lanata glomerata conferto-verticillatis, invol. squamis obtusis membranaceis. ① in Mexico circa lacum Sancti-Nicolai in sinu Spiritûs-Sancti et ni fallor in campis prov. Texas ad Fernando de Bexar legit cl. Berlandier (pl. exs. n. 1958 et 2109). Herba 4-6 poll. longa. (v. s.)

+ Species dubia.

- E. VERA (Raf. herb. p. 24), in Texas et Louisianâ. An huic generi adnumeranda?

++ Species exclusæ.

- E. ERICOIDES Schrank = *Eolopes tenuifolia* et *Helichrysum teretifolium*.
 E. INVOLUCRATA Schrank = *Petalacte coronata*.
 E. INDICA Ham. = *Filago pygmæa*.
 E. SPATHULATA Less. = *Gnaphalium bracteatum*.

CCXXVIII. MICROPSIS.

Capitulum pluriflorum heterogamum, fl. omnibus tubulosis, exter. 2-3-serialibus fœmineis, inter. circ. 3 masculis. Invol. squamæ 2-3-se-

riales apice membranaceæ oblongæ subcutæ glabræ. Recept. angustum nudum. Achænia villosa oblonga pappo parvo squamelloso superata. — Herba Chilensis habitu Evacis seu Micropi, undè nomen.

1. *M. NANA*. (1) in pascuis declivibus collium prope Quillota Chilensium legit cl. Bertero! (pl. exs. n. 1146), flor. septemb. Herba multicaulis vix pollicem longa. Folia alterna obovata adpressè cano-villosa integerrima obtusa mucronata, super. congesta subverticillata in axillis capitula parva sessilia foventia. (v. s.)

CCXXXIX. *MICROPUS* Linn. gen. n. 996. Gærtn. fr. 2. p. 389. t. 164. Cass. dict. 31. p. 39. Less. syn. 205. — *Gnaphalodes* Tourn. inst. 439. t. 261. Adans. fam. 2. p. 118. — *Micropi* sp. Desf.

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, radii circ. 5-7 uniserialibus fœmineis filiformibus, disci 5-7 masculis 5-dentatis. Invol. biseriale, squamis flores et fructus radii involventibus. Recept. ebracteolatum angustum. Achænia radii lateraliter compressa obovata glabra calva squamis inv. involuta et cum eo decidua. — Herbæ Europæ, lanato-tomentosæ, pusillæ, Filaginis s. *Gnaphalii* facie. Folia alterna. Capitula glomerata.

Sect. I. ACANTHOLÆNA.

Invol. squamæ achænia involvèntes dorso aculeis rigidis echinatæ.

1. *M. SUPINUS* (Linn. sp. 1313), totus adpressè sericeo-lanatus, caulibus diffusis erectisve, foliis spatulatis obtusis, capitulis axillaribus vix aggregatis, invol. squamis dorso echinatis. (1) in aridis inundatisque dessiccatis apricis Europæ Mediterraneæ ab Hispaniâ ad Persiam, nec non in Oriente et Mauritanîâ. Cav. ic. 1. t. 35. Lam. ill. t. 694. f. 1. Filago supina Lam. fl. fr. 2. p. 60. *Gnaphalodes* dentata Moench meth. 568. — Pluk. alm. t. 187. f. 6. Variat caule solitario erecto, aut plurimis diffusis imò procumbentibus. Fructus squamis inv. muricati involuti, ferè triangulares, non malè *Melampodii* achænia referunt. (v. v. et s.)

Sect. II. BOMBYCILÆNA.

Invol. squamæ achænia involvèntes dorso inermes lanatæ.

2. *M. ERECTUS* (Linn. sp. 1313), totus niveo-lanatus, caulibus erectis aut subdiffusis, foliis oblongo-lanceolatis obtusiusculis, capitulis axillaribus terminalibusque aggregatis, invol. squamis dorso convexis parè lanatis. (1) in apricis sterilibus rupestribus et inundatis exsiccatis Europæ Mediterraneæ et mediæ ab Hispaniâ ad Persiam, nec non in Oriente et Mauritanîâ. Lam. ill. t. 694. f. 2. Loeff. itin. 166. t. 1. f. 3. Lag. nov. gen. et sp. p. 32. n. 401. Variat ut prior caule sæpius erecto simplici aut multicauli, rariùs multicapiti supino (*Filago* multicaulis Lam. fl. fr. 2. p. 59). (v. v. et s.)
3. *M. BOMBYCINUS* (Lag. nov. gen. et sp. p. 32. n. 400), totus niveo-lanatus, caulibus erectis aut subdiffusis, foliis lineari-oblongis obtusis, capitulis congestis axillaribus et terminalibus, invol. squamis dorso convexis lanâ floccoso-tumidâ (nec adpressâ) obductis. (1) in collibus aridis Hispaniæ circa Madritum (Lag.), Tudelam (Duf.), et Galliæ australis in Occitaniâ totâ. *Micropus* erectus Guss. pl. rar. p. 345? Cass. dict. 31. p. 40? — Clns. hist. 1. p. 329. f. 2? Variat ut duæ sp. priores caule erecto subsimplici apice subdichotomo. (*M. discolor* Pers. ench. 2. p. 422), aut multicauli supino vel adscendente (Barr. ic. t. 296). Prioris fortè varietas? (v. v. et s.)
4. *M. GLOBIFERUS* (Bert. herb. n. 433), totus niveo-tomentosus, caulibus à basi ramosis diffusis, foliis oblongo-linearibus, floralibus latioribus obtusis, capitulis terminalibus lateralibusque, invol. squamis planiusculis inermibus to-

mentosis obtusissimis. ① in Chili ad Rancagua legit cl. Bertero. Squamæ recept. involutæ obovato-compressæ tomentosæ achænium foveantes et globum incanum cum invol. constituentes. (v. s.)

5. *M. MINIMUS*, totus niveo-tomentosus, caule brevissimo erectiusculo parçè ramoso, foliis ovali-oblongis obtusis, capitulis ad apices ramorum confertis. ① in Mexico ad Monterey januar. fl. legit cl. Berlandier (pl. exs. n. 1861). Priori affinis sed multò minor. Species quoad caract. fl. et fr. recognoscenda. (v. s.)

† *Species exclusæ.*

M. XXIGUUS Spreng. = *Evax exigua*.

M. MAREOTICUS Spreng. = *Evax Mareotica*.

M. PYGMÆUS Desf. = *Evax pygmæa*.

GCXXX. *EPALTES* Cass. bull. philom. 1818. p. 139. dict. 15. p. 6. 37. p. 339. Less. syn. 206. — *Ethulia* Gartin. fr. 2. p. 389. t. 164. non Cass. — *Ethulia* sp. Linn.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus fœmineis, disci paucioribus androgynis sterilibus. Invol. bi- aut pluriserialia imbricatum campanulatum. Recept. nudum convexum. Cor. radii filiformes tenuissimæ vix apice dentatæ, disci obconicæ 5-dentatæ extus parçè glandulosæ. Antheræ inclusæ. Stylus radii exsertus ramis revolutis, disci breviter exsertus indivisus. Achænium radii obovato-subangulatum glabrum epapposum, disci abortivum. — Herbæ ramosæ. Folia alterna oblonga aut obovata integra aut vix dentata sessilia aut longè decurrentia. Pedunculi oppositifolii. Capitula solitaria aut gemina pedicellata subpaniculata. — Genus affine *Denekia* et *Dichrocephala*.

* *Foliis decurrentibus.*

1. *E. DIVARICATA* (Cass. ll. cc.), caule herbaceo erecto divaricato-ramosissimo, foliis lineari-oblongis basi attenuatis longè decurrentibus remotè denticulatis, pedunculis folio brevioribus subbracteatis, invol. squamis acuminatissimis disco longioribus. ① frequens in oryzetis et agris Indiæ orient. *Ethulia divaricata* Linn. mant. 110. Burm. ind. 176. t. 58. f. 1. Wall. cat. et herb. n. 98. A et B. — Pluk. alm. t. 21. f. 4. Variat omninò glabra aut hinc indè versus apicem sublanuginosa. Invol. albida subscariosa rariùs subpurpurascens. (v. s.)
2. *E. LINEARIFOLIA*, caule herbaceo erecto apice corymboso-ramoso, foliis linearibus angustè decurrentibus subintegerrimis, pedunculis folio longioribus bracteolatis, invol. squamis acuminatissimis disco longioribus. ① in Ind. orientali circa Prome regni Avenis. *Ethulia divaricata* var. C. Wall. cat. et herb. n. 98. C. Priori simillima et fortè varietas? Involucrum in omnibus specim. mihi obviis subpurpurascens. Folia infer. sesquipollicaria vix ultra lineam lata. Pedunculi ferè pollicares bracteolas 3-5 acutissimas patulas gerentes. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
3. *E. PYGMÆA*, caule herbaceo nano ramoso, foliis lineari-oblongis subdenticulatis angustissimè decurrentibus, pedunculis ebracteatis folia vix superantibus, involucri squamis adpressis vix acutis flores non superantibus. ① in Indiæ orientalis agro Carnatico legit cl. Belanger. Herba sesquipoll. glabra. (v. s. comm. à cl. inv.)
4. *E. BRASILIENSIS*, caule herbaceo erecto ramoso, foliis oblongo-lanceolatis acuminatis dentato-serratis in alas integerrimas latas foliaceas secus caulem et ramos decurrentibus, pedunculis ramosis subulatis, corymbis folio brevioribus, invol. squamis ovalibus obtusis flores non superantibus. ① in humidis circa Bahiam legit cl. Salzmann. Habitus ferè *Pluchea* Quiloc, sed caract. genericis

et speciatim pappo nullo faciliè distinguenda. Cor. ex Salzm. lutescentes. (v. s. comm. à cl. Salzm.)

5. *E. MEXICANA* (Less. in linnæâ 1830. p. 147), caule infernè lignoso ramoso, foliis oblongo-obovatis denticulatis decurrentibus, pedunculis folio brevioribus, invol. squamis ovalibus acutis adpressis flores non superantibus. 3 in humidis prope Vera-Cruz. Cor. fl. femineorum stylum includunt. Herba glabra.

** *Foliis non decurrentibus.*

6. *E. AUSTRALIS* (Less. in linnæâ 5. p. 148. syn. 206), foliis oblongis parçè dentatis sessilibus non decurrentibus, pedunculis 1-3-cephalis, invol. squamis adpressis. — in Novâ-Hollandiâ (Less.), in paludosis ad ripas flum. Lachlan in Nov.-Holl. interiore (Cunn.!), flor. junio. Capitula globosa dimidiò quàm *E. divaricata* minora. (v. s.)
7. *E. HIRSUTA* (Less. syn. 206), tota hirsuta procumbens, foliis non decurrentibus, invol. squamis arcè imbricatis, capitulis solitariis vel geminis. — in Indiâ orientali. Cœt. ign.
8. *E. LITTORALIS* (Less. in linnæâ 1831. p. 151), procumbens strigosa, foliis spathulatis serrato-dentatis, capitulis pedicellatis solitariis binatisve, invol. scariosis nudis. — in littoribus marinis Indiæ orientalis. *Artemisia littoralis* Retz obs. 5. p. 28. Cœt. ign.

CCXXXI. *DENEKIA* Thunb. prod. fl. cap. in præf. nov. gen. 14. p. 177. fl. cap. 665. Willd. sp. 3. p. 1912. Cass. dict. 13. p. 65. Less. syn. 206. — Sellow sp. Spreng.

Capitulum multiflorum heterogamum, fl. radii pluriseriatis foemineis disco æqualibus bilabiatis, labiis exiguis æqualibus integerrimis, disci tubulosis 5-dentatis masculis. Recept. nudum planum. Invol. campanulati squamæ biserialis disco subæquales. Achænium sessile erostre. Pappus radii nullus, disci palea apice palmatisecta et aliis nonnullis brevioribus integris sæpius constans. — Herba Capensis erecta debilis. Folia alterna semiamplexicaulia oblonga undulata, subtus araneoso-tomentosa, suprâ glabra, mucrone apiculata, integra aut superiora sinuato-dentata. Capitula paniculato-corymbosa subconferta parva. Flores albi. Invol. glanduloso-hirtella.

1. *D. CAPENSIS* (Thunb. ll. cc.), foliis subtus canescenti-tomentosis hinc inde acutè dentatis. — ad Cap. Bonæ-Spei in aquosis prope Langekloof (Th.), ad Coloniam distr. orient. (Burch. ! cat. geog. n. 5062), ad Fisch-Rivier et Zwartz. Omtata (Drege! n. 5071 et 3776), in territorio cesso (Eckl. ! n. 1395). *Selloa Capensis* Spreng. syst. 3. p. 496. (v. s. comm. à cl. Burch. et à cl. Thunberg olim ad L'Hérit. sub nomine *Gnaphalii* miss.)

ℓ. *latifolia*, foliis radicalibus elliptico-oblongis petiolatis, caule superne glanduloso-hirtello basi longè radicante. — ad Port-Natal et Omtata legit cl. Drege. (v. s.)

γ. *minor*, foliis lineari-lanceolatis subintegerrimis, caule exili araneoso-tomentoso. 2 ad Stormberg alt. 4-500 ped. et circa Basche legit cl. Drege. Herba 6-7 pollic. Folia radic. angusta. (v. s.)

2. *D. GLABRATA*, foliis utrinquè viridibus, glabris aut pube subglanduloso-sparâ minutâ subinstructis, integris aut vix dentatis. 2 in Africâ australi ad Kipplat-Rivier, Zneureberge, Dumpf (Drege!), in Uitenhage et Tambukiland (Eckl. !). An prioris adhuc varietas? (v. s.)

Subtrib. IV. *INULEÆ* Cass. ann. sc. nat. 1829. p. 20. Less. syn. 197.

Capitula nunquàm dioica, rariùs homogama discoidea, sæpissimè heterogama radiata, nempe fl. foemineis ferè semper ligulatis. Recept. (excepto

Rhanterio) epaleaceum. Antherae caudatæ. — Folia alterna. Capitula in glomerulum non aggregata. Ligulæ ubi adsunt disco concolores. — Species omnes Gerontogæ.

DIV. I. EUINULÆ.

Capitula pluri- aut multiflora in glomerulum non conereta.

CCXXXII. RHANTERIUM Desf. *fl. atl.* 2. p. 291. t. 240. *Cass. dict.* 45. p. 312. *Less. syn.* 199.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus foemineis ligulatis, disci tubulosis hermaphr. 5-dentatis. Invol. ovatum, squamis basi concavis imbricatis productis in appendicem rigidam uncinato-reflexam. Recept. bracteolatum, bracteolis involucrique squamis internis flores subamplectantibus. Antherae apice acutè appendiculatæ, basi caudatæ. Achænia ovali-subpentagona glabra erostria, radii epapposa, disci coronata paleis 5 caducis basi integerrimis utrinquè ad apicem densè barbellatis. — Suffrutex Mauritanicus rigidus ramosus parçè foliosus, ramis tomentosis apice 1-cephalis. Folia sparsa lanceolata argutè serrata glabriuscula. Flores lutei. — Genus recept. paleaceo ad Bupthameas accedit.

1. R. SUAVEOLENS (Desf. l. c.). ♂ in arenis maritimis prope Sfax regni Tuncani. Folia trita suaveolentia. (v. s. comm. à cl. Desfontaines.)

CCXXXIII. INULA Gært. *fr.* 2. p. 449. f. 170. *Less. syn.* 198. — Inulæ et Conyze sp. *Lin.* — Inula, Limbarda, Corvisartia et Conyza *Cass.* — Conyza *Don prod. fl. nep.* 176 non *Lin.*

Capitulum multiflorum heterogamum, fl. radii 1-serialibus foemineis abortu interdum substerilibus, sæpius ligulatis, rarius subtubulosis 3-fidis, disci hermaphroditis tubulosis 5-dentatis. Invol. imbricatum pluriseriale. Recept. planum aut subconvexum nudum. Antherae basi bisetæ. Achænium erostre teretiusculum aut in I. Helenio tetragonum. Pappus conformis 1-serialis, setis capillaribus subscabris. — Herba gerontogæ pleræque Europæ aut Asiaticæ, perennes, rarius biennes aut annuæ. Folia caulina alterna, sæpè amplexicaulia, indivisa integerrima aut serrata. Capitula ad apices pedunculorum solitaria sæpè corymbosa flava.

Sect. 1. CORVISARTIA *Merat fl. par. ed.* 2. v. 2. p. 261. *Cass. dict.* 10. p. 572.

Invol. squamæ ext. latè ovatæ foliaceæ, inter. obovato-spathulatæ obtusæ. Achænium glaberrimum, in I. Helenio tetragonum, in I. Royleanâ fortè teretiusculum. — Si in utrâque specie achænium tetragonum Corvisartia ut genus habitu consentiente admittendum, si achænia diversa I. Royleana optimè jungit I. Helenium cum cæteris Inulis.

1. I. HELENIUM (Linn. sp. 1236), caule erecto, foliis dentatis subtus velutino-tomentosis acutis, radicalibus ovatis in petiolum longè attenuatis, caulinis semialexicaulibus, pedunculis paucis 1-cephalis ad apicem corymbosis. ♀ in Angliâ, Belgio, Galliâ septentr. et orientali, Germaniâ. fl. dan. t. 728. Gært. *fr.* 2. t. 170. Aster Helenium Scop. carn. n. 1078. Aster officialis All. n. 705. Corvisartia Helenium *Merat et Cass. ll. ec.* Helenium *Cam. epit.* 35. Fuchs hist. 242. Radix magna amara et aromatica, *Enula Campana* offic. dicæ (v. v. et s.)

2. *I. ROYLEANA*, tota molliter villosa subtomentosa, caule erecto simplici r-cephalo, foliis ovatis subdenticulatis, caulinis infer. in petiolum alatum basi obtuse auriculato-dilatatum semialexicaulem productis, summis paucis sessilibus cordato-amplexicaulibus, acheniis teretiusculis? $\frac{1}{2}$ in Indiæ prov. boreali-occid. detexit cl. Royle. Species insignis Inula Helenio habitu, involucri, ligulis numerosissimis angustis et pappo rigidioris proxima, sed foliis etiam caulinis plerisque longè petiolatis mediisque subpanduræformibus distincta. Achenia ex immaturo statu vid. teretiuscula nec tetragona. (v. s. comm. à cl. Royle.)

Sect. II. BUBONIUM. — Enula *Duby bot. gall. 1. p. 267* (sed Enula veterum est *Corvisartia*.)

Squamæ lineares aut lanceolatae, apice appendiculo parvo foliaceo auctæ, exter. subbreiores.

3. *I. CONYZA*, caule erecto apice ramoso subcorymboso pubescenti-subtomentoso, foliis ovato-lanceolatis subacutis subintegerrimis supra subpubescentibus subtus tenuè tomentosis, radicalibus in petiolum attenuatis, caulinis sessilibus, invol. squamis linearibus apice foliaceis patulo-recurvis, fl. radii foemineis trifidis subligulatis. $\frac{1}{2}$ ad sepes, colles, prata sicca præsertim in calcareis totius ferè Europæ usque ad Byzantium, Tauriam et Persiam. *Conyza squarrosa* Linn. sp. 1205. fl. dan. t. 622. engl. bot. t. 1195. Cass. dict. 10. p. 306. Lam. ill. t. 697. f. 1. *Conyza vulgaris* Lam. fl. fr. p. 73. *Inula squarrosa* Bernh. non Linn. *Erigeron squarrosus* Clairv. herb. val. 243. *Inula* sp. Less. syn. 198. — Cam. epit. 61a. Herba sordida obscura trita foetida. Invol. è virescente purpurascens. Cor. flavæ, marginales ad latus internum profundius fissæ et ideo ad ligulas subaccedentes. (v. v.)
4. *I. SUAVIOLENS* (Jacq. hort. vind. 3. p. 29. t. 51), caule erecto simplici apice subcorymboso, foliis ellipticis subvillosis, inferioribus in petiolum longè attenuatis remotè serrulatis, caulinis basi breviter attenuatis integerrimis, capitulis 5-7 pedunculatis, invol. campanulati squamis apice foliaceis subrecurvis, ligulis linearibus involucri longioribus. $\frac{1}{2}$ in Italiæ littoribus circa Monfalcone (Suffr. ex Reichenb. fl. exc. a. p. 237), fortè in Gallo-Provinciâ ex Lam. Murr. syst. p. 766. Wild. sp. 3. p. 2099. DC. fl. fr. n. 3144. Herba trita potius gravè quàm suavè olens. Valdè affinis *I. Conyza* et hujus varietas ligulata videretur nisi capitula majora. (v. s. c. in hort. Taurin. et Cels.)
5. *I. CANDIDA* (Cass. dict. 23. p. 554), tota densè candido-tomentosa, foliis ovatis radicalibus longè, caulinis breviter petiolatis, pedunculis terminalibus et axillaribus r-cephalis, invol. squamis oblongo-linearibus obtusis. $\frac{1}{2}$ in rupestribus apricis Cretæ, Sami, Dalmatiæ, Istriæ.
- a. *verbascifolia*, foliis subcrenato-dentatis acutiusculis, capitulis radiatis, ligulis elongatis. $\frac{1}{2}$ *Conyza verbascifolia* Willd. sp. 3. p. 1924 non Bory. Con. candida Linn. sp. 1208. Sibth. fl. græc. t. 861 ex Sm. prod. 2. p. 173. — Bocc. sic. t. 31. f. 2. (v. s. ex Dalmatiâ.)
- β. *limonifolia*, foliis integerrimis obtusis, capitulis flosculosis aut ligulas involucri breviores gerentibus. $\frac{1}{2}$ *C. limonifolia* Sibth. fl. græc. t. 865 ex Sm. prod. 2. p. 174. *C. candida* Will. sp. 3. p. 1923 non Lour. (1) Barr. ic. t. 217. (v. s. ex Atticâ, Cretâ et Samo.)
6. *I. TRAPSOIDES*, tota tomentoso-lanata, caule erecto apice corymboso-ramoso, foliis serratis acuminatis radicalibus petiolatis ovali-oblongis, caulinis adnatis longè decurrentibus, capitulis corymbosis confertis, invol. squamis linearibus. $\frac{1}{2}$ in herbidis ad Caucasum. Etiam si nonnulli dixerint nihil esse clarius ac identitas synonym. infra citatorum, fortè 3 varietates sequentes ut species habendæ :
- a. *Sprengelii*, tomento denso crasso, foliis caulinis brevius decurrentibus,

(1) *Conyza candida* Lour. coch. 2. p. 605 est verisimiliter species omnino diversa, sed nondum satis nota.

capitulis paucioribus latioribus, involucri tomentoso-lanati squamis obtusis, ligulis involucri discumque longè superantibus. — *Inula thapsoides* Spreng. ! pug. 1. p. 58. Cat. sem. h. hal. 16. Media ferè inter *I. candidam* var. α et var. seqq. (v. s. comm. à cl. Spreng. et v. c. in h. Monsp.)

6. *Poiretii*, tomento parciore vix albido, foliis caulinis longius decurrentibus, corymbis numerosioribus confertis, invol. ovati squamis linearibus acutis glabriusculis, ligulis vix involuero longioribus. — *Inula verbascifolia* Poir. suppl. 3. p. 154. Horn. h. hafn. supp. 99. *Inula thapsoides* Reichenb. icon. t. 347. (v. s. et v. c. in hort. Par.)

7. *Biebersteinii*, tomento denso-erasso, foliis caulinis longè decurrentibus, corymbis numerosis confertis, invol. ovato-oblongi squamis linearibus rectis acutis vix dorso villosis, ligulis nullis aut paucissimis brevissimis trifidis. — *Conyza thapsoides* Bieb. fl. taur. et suppl. n. 1716. (v. s. comm. à cl. Steven et v. c. in hort. Monsp.)

7. *I. POLYGONATA*, ramis teretibus villosis apice tomentosis foliorum decurrentiis integerrimis longis alatis, foliis elliptico-oblongis utrinquè attenuatis dentatis suprâ asperis subtus et summis utrinquè densè tomentosis, corymbo terminali, capitulis subsessilibus paucis bracteis tomentosis circumdatis, invol. squamis ext. oblongis densè tomentosis obtusis, intimis lanceolatis acuminatis scariosis glabris, ligulis nullis. 5? in Indiâ or. ad Prome in regno Avenis legit cl. Wallich. *Conyza polygonata* Wall. ! cat. n. 3010. comp. n. 120. Caulis fortè basi frutescens? Rami pedales et ultrâ crassitie pennæ anserinæ. Foliorum superiorum dentes callosi. Flores ex statu misero specim. non ritè vidi et hic retuli speciem ex analogiâ cum *In. candidâ* et *thapsoides*. (v. s. comm. ab amp. Cur. merc. angl. Ind. or.)

8. *I. BIFRONS* (Linn. sp. 1236), caule apice ramoso corymboso-subdichotomo viscoso, foliis glabris ovali-oblongis aliis cordato-amplexicaulibus aliis decurrentibus, basi dentatis apice integerrimis mucronatis, capitulis confertè corymbosis subsessilibus, invol. squamis oblongo-linearibus subsquarrosis, ligulis brevissimis. ② in vallibus et humidis Pyrenæorum (Herm.), Arvernæ, Occitanie, Galloprovinciæ, Pedemontii, Liguriæ, Galliciæ (Bess.), Italiæ. DC. fl. fr. 4. p. 155. Reichenb. icon. 4. n. 527. t. 346. Inv. glabra Bess. cat. h. Cram. ex Reich. — Herm. par. t. 127. Bocc. mus. 1. t. 122. Garid. Aix. t. 23. — *Conyza alata* Baumg. ex Reichenb. (sed in fl. trans. non reperi). *Aster bifrons* All. ped. n. 714. *Inula glomeriflora* Lam. fl. fr. 2. p. 150. *Conyza bifrons* Gou. hort. 436 non Linn. (v. s.)

9. *I. OCLUS-CHRISTI* (Linn. sp. 1237), caule erecto apice parcè ramoso-corymboso molliter villosa aut lanata, foliis oblongo-lanceolatis integerrimis utrinquè villosis-sericeis, radicalibus in petiolum attenuatis, caulinis basi obtusâ amplexicaulibus, capitulis 3-5 longiusculè pedunculatis, invol. squamis oblongo-linearibus dorso hirsutis, ligulis invol. duplò longioribus. 2? in collium lateribus graminosis apricis Austriæ, Hungariæ, Dalmatiæ. Willd. sp. 3. p. 2090. — Clus. hist. 2. p. 20. f. 2. Plantam ex Austriâ loco Clusiano lectam (quæ typus speciei) à Jacquinio illustratam et à filio suo mihi missam comparavi cum Caucasicâ nec ullum discrimen specificum vidi. — *Inul. montanam* fl. taur. n. 1746 huc refert cl. Steven. (v. s.)

6. *campestris*, tota parcius villosa, foliis superioribus non tantum cordato-amplexicaulibus sed etiâ subdecurrentibus. 2? in Podoliâ (Bess.), Tauriâ (Bieb. Stev.!) et Persiâ (Szow.!). *I. Oclus-Christi* Bieb. fl. taur. n. 1744. *I. campestris* Bess. enum. Volh. p. 33 et 108. Recept. in hac et in var. α pube minutissimâ breviter puberulum. (v. s.)

10. *I. HELENIOIDES* (DC. fl. fr. suppl. (1815) p. 470), caule erecto hirsuto supernè corymboso-ramoso, foliis ovali-oblongis subhirsutis subdentatis, radicalibus petiolatis, superioribus basi obtusâ amplexicaulibus, capitulis 3-5-foliis floralibus bracteatibus, invol. campanulati squamis linearibus hirsutis, ligulis invol. duplò longioribus, recept. ad medium setuloso. 2? in agris asperis Pyrenæorum

orientalium damissorum legi ad pagos Finestret, Ria, Billoc, etc. et in montibus Corbarienais. Inula Oculus-Christi Lapeyr. abr. 522 non Linn. (v. v. sp.)

β. *flaculosa*, ligulis nullis. (v. s. c.)

11. *I. HIRTA* (Linn. sp. 1239), caule erecto subramoso piloso-hirto, foliis sessilibus lanceolato-oblongis rigidis integerrimis secus nervos et margines hirtis, capitulis solitariis aut subternis corymbosis, invol. squamis hirtis ciliatis linearilanceolatis, exterioribus interiores superantibus, acheniis glabris. ½ in pratis asperis et montosis Galliæ imò prope Parisios, Pedemontii, Appennini Genueusis et Etrusci, Sabaudia, Helvetiæ, Carniolæ, Thuringiæ, Austriæ, Tauriæ, Sibiria, Dahuriæ, Jacq. fl. austr. t. 358. DC. fl. fr. 4. p. 151. Inula montana Poll. pal. Aster hirtus Scop. carn. n. 1082. t. 58. Clus. hist. 2. p. 14. f. 2. Variet 1-3-cephala. (v. v. et s.)

12. *I. VAILLANTII* (Vill. dauph. 3. p. 216), caule erecto ramosissimo subcorymboso folisque subtus pube brevi velutino-cinerea, foliis oblongo-lanceolatis suberratis, caulibus sessilibus, infimis basi subattenuatis, invol. squamis ext. oblongo-linearibus subfoliaceis, intimis linearibus glabris, ligulis invol. duplò longioribus, acheniis glabris. ½ in dumetis et pratis montosis Helvetiæ orientalis, Sabaudia, Delphinatus et Pedemontii. Fl. fr. n. 315a. Gaud. fl. helv. 5. p. 325. Aster Vaillantii All. fl. ped. n. 719. In. cinerea Lam. dict. 3. p. 259. Inula ramosissima Clairv. mau. herb. 244. — Hall. helv. n. 73. t. 2. opt. Herba Melissam ferè olet. (v. v. et s.)

13. *I. SEMIAMPLEXICAULIS* (Renter! mem. soc. Gen. 7. p. 169. ic.), caule erecto apice corymboso pubescenti-subviloso, foliis semiaplexicaulibus oblongo-lanceolatis acutis calloso-serratis suprà glabris subtus pubescenti-villosis, invol. squamis ext. oblongo-lanceolatis patulis foliaceis, inter. linearibus glabris, ligulis invol. duplò longioribus, acheniis glabris. ½ olim legi in Pedemontii agro Caneensi et cum *I. Vaillantii* præeunte Allionio (n. 719?) confudi, recentius in sylva La Batte prope Genevam cum *I. Vaillantii* et *I. salicina* legit oral. Renter. Utriusque fortè hybrida? Differt ab *I. salicina* foliis dorso cauleque villosis, herbà Polii odore donatà, foliis frequentius serratis. Distinguitur ab *I. Vaillantii* foliis amplexicaulibus minùs villosotomentosis paulò rigidioribus, odore subdiverso, ligulis paulò longioribus. (v. v.)

14. *I. SALICINA* (Linn. sp. 1238), glabra, caule erecto apice corymboso-ramoso, foliis semiaplexicaulibus lanceolatis margine ciliato-scabris apice subrecurvis, ramis 1-cephalis, invol. campanulati squamis ovato-lanceolatis serrulato-scabris apice subreflexis, acheniis glabris. ½ in pratis asperis dumetis et sylvis Gallia, Helvetiæ, Italiæ, Germaniæ, Daniæ, etc. Fl. dan. t. 786. Fl. fr. 4. p. 151. Gaud. fl. helv. 5. p. 324. Aster salicinus All. ped. n. 709. — Clus. hist. 2. p. 15. f. 1. Mor. ox. s. 7. t. 21. f. 10. Herba inodora. (v. v. sp.)

β. *lanceifolia*, foliis ovato-lanceolatis, rigidioribus margine magis scabridis. ½ Inula aspera Poir. suppl. 3. p. 154. (v. v. c.)

γ. *lanceifolia*, caule hirto, capitulis minoribus. ½ Inula lanceifolia Wenderoth in florà 1829. erg. p. 1. p. 26. An potiùs *I. squarrosa* seu *spiræifolia* var.?

15. *I. ENSIFOLIA* (Linn. sp. 1240), caule erecto glabro apice araneoso simplici 1-5-cephalo, foliis sessilibus linearibus calloso-acuminatis multinerviis glabris, capituli squamis lanceolatis erectis foliosis subtus araneoso-tomentosis. ½ in collibus et asperis Pedemontii, Italiæ, Carniolæ, Baviaræ, Austriæ, Tauriæ, Caneensi. Jacq. fl. austr. t. 162. Biebert. fl. taur. n. 1753. D'Urv. enum. n. 811. Aster ensifolius Scop. carn. n. 1084. — Bocc. mus. 1. p. 31. t. 18. Variet caule 1-cephalo aut polycephalo sed trea varietates species mihi non satis notas distinguit cl. Wenderoth in florà 1829. p. 1. p. 26. (v. s.)

16. *I. SQUARROSA* (Linn. sp. 1240 non Bernh.), caule erecto vix ramoso scabriusculo, foliis oblongis acuminatis serratis utrinquè et margine plùs minùs scabridis reticulato-venosis, capitulis ad apices ramulorum solitariis vix corymbosis, invol. campanulati squamis ovatis apice patulo-reflexis margine scabridis.

¶ in sylvis petrosis Gallie, Helvetiæ et Germaniæ australis, Italiæ, etc. Bert. em. ital. p. 192 et Gaud. fl. helv. 5. p. 321. var. æ. DC. fl. fr. n. 3148. Aster corymbosus Moench. non Ait. In. spiræifolia Lam. dict. 3. p. 258. — Fink. alp. t. 16. f. 1. (v. v. sp.)

β. *latifolia*, foliis ovato-lanceolatis subscabris. ¶ in Delphinatu prope Grandanopolim. Inula germanica Vill. Dauph. 3. p. 219. Lam. dict. 3. p. 258. DC. fl. fr. n. 3149 non Linn. (v. s.)

17. I. *spiræifolia* (Linn. sp. 1238 non Lam.), puberulo-scabra, caule erecto simplici apice summo confertè corymboso, foliis sessilibus oblongis cartilagineo-denticulatis et mucronatis rigidis reticulatis; invol. squamis ovatis glabris obtusiusculis squarrosis. ¶ in Pedemontio, Carniolâ calidiore, Austriâ littorali, Podoliâ, Tauriâ, Caucaso, Persiâ. Inula Bubonium Murr. syst. 767. Jacq. fl. austr. app. t. 19. Bieb. fl. taur. n. 1751. In. salicina Pall. ind. taur. n. 154 ex Bieb. Aster Bubonium Scop. carn. n. 1083. t. 58. Aster squarrosus All. pedem. n. 708. excl. syn. ex cl. Moretti qui primus synonym. hujus speciei elucida vit. — Aster IV Clus. hist. 2. p. 14. Bubonium Tab. hist. 716. Inula spiræifolia et Bubonium Willd. sp. 3. p. 2094 et 2097. (v. s.)

β. *hybrida* (Bauing. fl. trans. 3. p. 132), capitulis subternis majoribus. ¶ in montosis Podoliæ et Transylvaniæ. Fortè hæc var. I. spiræifolia et I. squarrosam Linn. præcæntibus cl. Lamarck Duby et Gaudin in unam colligit. (v. s.)

18. I. *micranthos* (Poir. dict. suppl. 3. p. 153), caule erecto simplici pubescente, foliis sessilibus lanceolatis integerrimis suprâ subscabris subtus villosis subserricis, capitulis corymbosis confertis, invol. squamis linearibus acutis dorso puberulis, achæniis villosis. ¶ in Armeniâ (Tourn.), in collibus Bulgariæ circa Sisepolim (D'Urv.) et in herbidis apricis Tauriæ (Bieb.). Inula orientalis Willd. sp. 3. (1800) p. 2100, non Lam. (1789). I. germanica Bieb. fl. cauc. n. 1752 non Linn. ex D'Urv. enum. 815. Aster armenius salicis folio flore luteo minime umbellato Tourn. cor. 36. Affinis I. Germanicæ sed quidquid dix. Spreng. diversa foliis subtus serricis et præsertim invol. subcampanulato nec oblongo, achæniis villosis, etc. (v. s. comm. à cl. D'Urv.)

19. I. *germanica* (Linn. sp. 1239), caulibus erectis simplicibus apice corymbosis, foliis cordato-semiamplexicaulibus ovali-lanceolatis integerrimis suprâ nudiusculis subtus pubescentibus, capitulis confertis breviter pedicellatis oblongis, invol. squamis adpressis apiculo subreflexo, ligulis disco parum longioribus, achæniis glabris. ¶ in herbidis apricis Saxonie, Thuringiæ! Bessarabiæ!, in Tauriâ ad Odessam, etc. fortè in Sibirîâ (sic icon Gmel. fl. sib. 2. t. 78. f. 2. huc pertinet?). — Jacq. fl. austr. t. 134. opt. I. Germanica D'Urv. enum. xi. I. media Bieb. suppl. p. 676. Petit enum. pl. pal. 94. — Moris. ox. sv 7. t. 19. f. 26. pess. — I. Germanica florist. Gall. est diversa. (v. v.)

20. I. *media* (Bieb. suppl. 576), pilosiuscula, caule apice corymbum compositum glomeratum gerente, foliis sessilibus cordato-lanceolatis, infimis subdentatis, ligulis invol. parum longioribus, invol. subtomentosis. ¶ in Poloniâ australi legit cl. Besser ex Bieb. Media inter I. salicinam et Germanicam. Corymbi rami sub-1-cephali capitulo æquales aut breviores. †

21. I. *caspica* (Blum. in Ledeb. ind. sem. h. dorp. 1822. p. 10), caule herbaceo tereti stricto simplici glabro, foliis elongatis lineari-lanceolatis sessilibus basi obtusè subauritis glabris integerrimis margine scabris, corymbi compositi ramis villosis, invol. squamis linearibus, ext. subfoliaceis rigidè ciliato-scabris, ligulis numerosis angustis. ¶ ad mare Caspium in salsis prope Astrachan. Folia 4 poll. longa, 3-4 lin. lata. (v. s. comm. à cl. Fischer.)

22. I. *britannica* (DC. fl. fr. 4. p. 149), caule erecto villosa apice corymboso 3-5-cephalo, foliis longè lanceolatis remotè denticulatis subtus villosis, infimis basi attenuatis, superioribus dilatatis semiamplexicaulibus, invol. squamis linearibus laxis achæniisque hirsutis. ¶ in pratis secus vias et fossas, in Galliâ ferè totâ, Germaniâ, Daniâ, Italiâ, Græciâ (Sibth.), Persiâ (Szow.), Bulgariâ (D'Urv.), Sibirîâ, sed non in insulis Britannici! Inula Britannica Linn.

- sp. 1237. Schkuhr handb. t. 247. Fl. Dan. t. 413. Aster Britannicus All. pedem. n. 712. In. hirta Poll. pal. — Moris ox. s. 7. t. 19. f. 8. — Britanica Dalech. hist. 1083 unde nomen specificum à plurimis orthographiâ erroneâ corruptum *Βριτανικα*, Diosc. ex Dalech. Variat foliis integerrimis aut semiseratis, caule 1-9-cephalo, capitulis minoribus ubi numerosi, majoribus ubi pauciores adsunt, foliis præsertim superioribus latioribus quæ I. Oëtteliana Reich. fl. exc. 2. p. 237. (v. v. et s.)
- β. *comosa*, invol. squamis in folia disco longiora abeuntibus. I. *comosa* Lam. fl. fr. 2. p. 147. (v. s. olim.)
- γ. *dichotoma*, pedunculis bifidis dichotomisve, capitulis minoribus, invol. squamis apice patulo-reflexis. — in Oriente nempe in *Egypto*? ex Zucc., in Persiæ prov. Aderbeidschan ad rivulorum margines circa Khoi (Szow.! ex sp. Fisch.) et ad mare Caspium prope Astrachan et Hinkenberg? ex Ledeb. Inula dichotoma Zucc. obs. bot. n. 88. Inula Britanica anthod. minor. Fisch. et Mey. pl. Szow. ined. (v. s.)
23. I. *MONTANA* (Linn. sp. 1241), caule erecto 1-cephalo villosa, foliis lanceolatis integerrimis villosis, radicalibus in petiolum attenuatis, invol. squamis ext. oblongo-lanceolatis obtusiusculis extus villosis, inter. linearibus acutis, achæniis hirtis. ½ in sterilibus montanis apricis Hispaniæ, Siciliæ, Galliæ australis, Pedemontii, Helvetiæ et Germaniæ australis, Istriæ et Tauriæ. DC. fl. fr. 4. p. 154. — Garid. Aix. t. 10. — Lob. ic. 350. f. 2. — Aster montanus All. ped. n. 706. In. odora D'Urv. enum. n. 812. excl. syn. Pulicaria calycina Presl. del. prag. p. 96 et Inula calycina Presl fl. sic. 1. p. xxx huc pertin. monente cl. Gussone, Multum refert Pulicariam odorem sed facili differt pappo uno nec biseriali. Variat caule solitario aut multicapiti, 1-cephalo aut rarius apice ramoso 1-3-cephalo. (v. v. et s.)
24. I. *GRANDIFLORA* (Willd. sp. 3. p. 2098), caule simplici 1-cephalo apice villosa, foliis sessilibus oblongis parè et sparsè pubero-villosis serratis, serraturis apice glandulosis, invol. squamis lineari-subulatis fulvo-hirsutis. ½ in alpestribus Caucasii Iberici (Stev.!), et ad Guriel Persiæ (Szow.!), Bieb. fl. taur. et suppl. n. 1748. Inula alpina Adam in Web. et Mohr cat. 1. p. 68. In. glandulosa ÷ Sims. bot. mag. n. 1907. (v. s. comm. à cl. Steven.)
25. I. *GLANDULOSA* (Willd. sp. 3. (1800) p. 2097 non Lam. 1789), caule simplicissimo 1-cephalo patentum et molliter hirtum, foliis sessilibus ovali-oblongis utrinquè villosis hirtis glandulas secus marginem sessiles dentiformes gerentibus, invol. squamis lineari-subulatis nigricantibus hirsutis. ½ in alpestribus Caucasii. Bieb. fl. taur. et suppl. n. 1749. Bot. mag. t. 1907. bot. reg. t. 334. I. orientalis Lam. dict. 3. (1789) p. 255. non Willd. (1800). Capitulum radio excepto ferè sesquipoll. diam. Ligulæ lineares angustæ pollicem longæ. Valde affinis I. grandifloræ. Nomen Lamarckii arbitrariò à Willd. mutatum, imò cum altero confusum, stricto jure servandum esset. (v. s. comm. à cl. Steven.)
26. I. *GRAVEOLENS* (Desf. cat. ed. 2. p. 121), piloso-viscosa ramosissima, foliis inf. oblongo-lanceolatis subdentatis, sup. linearibus integris, ligulis minimis vix invol. æquantibus. (†) in apricis Europæ australis ad Constantinopolim, agrum Trajanum, Græciam, Italiam, Galliam australem et occid. Hispaniam, etc. Cass. dict. 23. p. 553. Erigeron graveolens Linn. sp. 1210. Solidago graveolens Lam. fl. fr. 2. p. 145. DC. fl. fr. n. 3162. — Barr. ic. t. 370. Caulis vetustus sæpè indurato-suffrutescens. Pappi setæ numerosiores quàm in cæteris. (v. v. et s.)
- β. *frutescens*, caule humili fruticoso cæspitoso ramosissimo, foliis linearibus brevibus obtusiusculis integerrimis, ligulis fortè nullis. ½ ad Alexandriam Egypti. (v. s. comm. à cl. Martins.)
27. I. *LATIFOLIA*, caule simplici basi tereti glabriusculo suffruticuloso, apice striato-sulcato setis hispido, foliis sessilibus latè ovalibus acutis basi longè cuneatis remotè serratis sparsè setosis obliquè penninerviis, capitulis paucis terminalibus, invol. squamis ext. linearibus subfoliaceis hirtis interioribus paulò brevioribus, ligulis invol. superantibus, achæniis villosis. ½ ¾ in Indiâ orient.

- legit cl. Royle (n. 113. 2). Pedalis aut sesquipedalis. Folia circ. 4 poll. longa, 2 poll. lata, membranacea, serraturis callosis exsertis. (v. s. comm. à cl. Royle.)
28. *I. AFRICANA* (Lam. ! dict. 3. p. 256), caule herbaceo tenui vix pilosiusculo ramoso, foliis amplexicaulibus subscabridis subtus glanduloso-punctatis, inferioribus oblongis acutis serratis, summis lanceolatis brevioribus subintegerrimis, ramis apice 1-cephalis aliis aphyllis aliis parce foliolosis, invol. squamis lineari-setaceis. — ad Cap. Bonæ-Spei ex Sonnerat in h. Lam. et ad Galam Africæ (Cass.). *Iphiona punctata* Cass. dict. 23. p. 610. *Inula punctata* Less. syn. 198. Folia 6-7 lin. longa, 2-3 lin. lata. Capitula subglobosa 3-4 lin. diam. pallide flava. Ligulæ breves. Habitus Pulicariæ sed pappus 1-serialis, Vicoæ sed fl. radii papposi! (v. s.)

Sect. III. CAPPA.

Invol. squamæ multiseriales obtusiusculæ regulariter imbricatæ, ext. brevissimis, int. gradatim longioribus. Ligulæ paucæ parvæ aut subnullæ, fl. radii nempe 2-3-dentatis ferè tubulosis (sed uniserialibus fœmineis ut in *Inulâ*). — Suffrutices Indici.

29. *I. EUPATORIODES*, caule fruticoso, ramis teretibus striatis rufo-velutinis, foliis ellipticis acuminatis basi attenuatis breviter petiolatis rariter et minutè serratis suprâ sparse scabris subtus rufo-cinereo-pubescentibus, paniculâ thyrsoidæ polycephalâ compositâ, ramis inf. axillaribus, invol. squamis oblongis obtusis subhirsutis, ligulis paucis inv. paulò longioribus, achæniis villosis. § in montibus prov. Silhet et fortè in Nepaliâ Ind. or. leger. cl. Bruce et Wallich. *Conyza eupatorioides* Wall. ! cat. et herb. n. 2993. comp. n. 103. Anth. breviter caudatæ. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)
30. *I. OBLONGA*, fruticosa, ramis teretibus apice præsertim sericeo-villosis, foliis brevissimè petiolatis oblongis utrinquè attenuatis dentato-serratis suprâ scabris subtus sericeis pubescentibusve, corymbo terminali congesto oligocephalo, capitulis vix pedicellatis, invol. squamis linearibus acutis velutino-hirsutis disco brevioribus, fl. fœm. paucis circ. 2-serialibus. § in Indiæ orient. montanis. Cor. purpureæ. Pappus apice crassius barbellaris subclavellatus. Ovarium teres pubescenti-sericeum. Anth. caudatæ.
- α. *argentea*, foliis subtus densè sericeis albis, à medio ad apicem serratis. § ad Taong-Dong, Srinaghur, Kamroop et Charcool in Deyra-Dhoon legit cl. Wallich. *Conyza argentea* A. B. C. Wall. ! cat. et herb. n. 3014. comp. 124 non Lam. *Conyza oblonga* Wall. ! herb. (v. s. comm. ab ampl. Cur. merc. Angl. Ind. or.)
- β. *pubescens*, foliis tenuioribus subtus pubescentibus vix subsericeis ferè à basi ad apicem serratis. § in demissis Nepaliæ inferioris. C. *argentea* D. Wall. ! cat. l. c. An species propria? (v. s. comm. cum α.)
31. *I. CAPPA*, caule fruticoso, ramis teretibus albedo-velutinis, foliis elliptico-oblongis acuminatis basi attenuatis breviter petiolatis suprâ scabris subtus tomento sericeo niveo lanuginosis, paniculâ compositâ terminali polycephalâ, ramis niveis, inf. axillaribus folio multò brevioribus, invol. squamis linearibus subacutis albo-velutinis, ligulis subnullis, achæniis villosis. § in Nepaliâ superiore. *Conyza Cappa* Ham. in Don prod. 176. *Conyza lanuginosa* Wall. ! cat. et herb. n. 2992. comp. n. 102. *Eupatorium lanatum* Spreng. in litt. ad Wall. Frutex elegans. Pappus apice scaber. Fl. radii 1-ser. tubulosi 3-dentati aut subligulati sed cæteros non superantes. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)
32. *I. PSEUDO-CAPPA*, caule fruticoso, ramis teretibus scabridis villosis, foliis ellipticis vix acutis basi subobtusis sessilibus subintegerrimis suprâ sparse scabris subtus sericeo-lanato-villosis albidis, paniculâ terminali compositâ polycephalâ, ramis inf. axillaribus folia subabortiva triplò superantibus, inv. squamis lineari-oblongis vix subhirsutis, ligulis subnullis, achæniis villosis. § in Chinâ legit cl. Staunton. Valde affinis *I. Cappa*. Folia infer. non vidè sed super. sessilia ubi in *I. Cappa* petiolatâ. (v. s. comm. à cl. Lambert.)

33. *I. xanthoxona*, caule suffruticoso erecto tereti dense hirsuto, foliis subsessilibus oblongo-lanceolatis subintegerrimis suprâ hispido-velutinis subtus dense albo-lanuginosis, corymbo terminali conferto 8-9-cephalo, invol. ovati squamis linearibus acutis multiserialibus dorso tomentosis, ligulis subnullis, achæniis villosis. $\bar{\zeta}$ in Nepaliâ inferiore legit cl. Wallich. *Conyza eriophora* Wall. ! cat. n. 3013. comp. 123. Fl. radii. 1-ser. fœm. tubulosi tenues, cæst. herm. stigmat. longis exsertis. Pappus 1-serialis. Affinis *I. Cappæ*. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)

Sect. IV. LIMBARDA. — *Limbarda Adans. fam. 2. p. 125. Cass. dict. 23. p. 565. et 26. p. 437. Reichenbach fl. exc. 2. p. 237.*

Invol. squamæ omnino exappendiculatæ adpressæ acuminatæ pauciseriales, inter. non multò exteriores superantibus. Achænium villosum teretiusculum.

34. *I. CRITHMOIDES* (Linn. sp. 1240), fruticulosa glaberrima, foliis linearibus obtusis carnosis integerrimis aut apice trifidis. $\bar{\zeta}$ ad rupes maritimas secus Oceanum ab Angliâ ad Lusitaniam et secus mare Mediterraneum ferè totum. DC. fl. fr. 4. p. 154. Engl. bot. t. 68. *Inula crithmifolia* Willd. sp. 3. p. 2101. Senecio succulentus Forsk. descr. æg. ar. 149. Senecio crithmifolius Scop. carn. n. 1066. *Limbarida tricuspidis* Cass. dict. 26. p. 438. *Inula Canariensis* Mill. dict. n. 12? excl. patr. — Lob. ic. t. 395. f. 2. (v. v.)

35. *I. AMMOPHILA* (Bunge! enum. mong. ex litt.), glabrescens, caulibus suffruticulosus diffusis gracilibus, foliis lanceolato-linearibus sessilibus, infer. subcordatis integerrimis crassiusculis patentibus, capitulis solitariis terminalibus, invol. squamis adpressis acutis pallidis, fl. radii angustè ligulatis. $\bar{\zeta}$ in sabulosis Chinæ borealis legit cl. Bunge. Folia 2-3 lin. longa subvillosa, infer. breviora latiora, sup. angustiora paulò longiora. (v. s. comm. à cl. Bunge.)

ß. salsoloides, glanduloso-subviscosa, foliis evidentius cordatis, fl. radii tubulosi. $\bar{\zeta}$ in sabulosis Mongoliæ Chinensis circa Chadatu detexit cl. Turczaninow. *Conyza salsoloides* Turcz. ! pl. chin. et mong. dec. n. 20. Certè *Inula* ex antheris basi bisetis et *I. ammophila* var. *discoidea* videtur etiamsi viscosa et foliis paulò latioribus donata. (v. s. comm. à cl. Turczaninow.)

36. *I. VISCOSA* (Ait. kew. ed. 1. v. 3. p. 223), caule erecto piloso-viscoso apice paniculato-ramoso, foliis lanceolatis serratis utrinquè glanduloso-pilosis sessilibus cordato-auriculatis, auriculis reflexis, ramis lateralibus foliosis oligocephalis, invol. squamis linearibus. $\bar{\zeta}$ in apricis Teneriffæ (Cham.!), ina. Archipelagi (Beaupré!), agri Hierosolymitani (Bové!), Hispaniæ, Galliæ australis, Italiæ mediter., Mauritanie. Desf. fl. atl. 2. p. 274. DC. fl. fr. 4. p. 173; *Erigeron viscosus* Linn. sp. 1209. Jacq. hort. vind. t. 165. *Solidago viscosa* Lam. fl. fr. 2. p. 144. — Clus. hist. 2. p. 20. f. 1. Bocc. sic. t. 7. f. B? Variat foliis 3-9 lin. latis. (v. v. et s.)

37. *I. BARBATA* (Wall. ! cat. et herb. n. 71), caulibus erectis subsimplicibus basi glabris apice sparse pilosis, foliis oblongo-lanceolatis acuminatis basi vix attenuatis semiamplexicaulibus serrulatis ciliato-barbatis, capitulis sessilibus subsolitariis foliis cinctis, invol. squamis lanceolato-linearibus foliaceis longè barbato-ciliatis. $\bar{\zeta}$ ad Srinaghur legit cl. Blindworth. Herba pedalis et sesquiped. Radix fasciculata multicaulis. Capitula ampla ferè bupthalmoides. Ligulæ longissimæ lineares stylo brevissimo subabortivè glabræ pollicem longè semin. lin. latæ. Pappus 1-serialis capillaris albus. Achænium parvum subpuberulum. (v. s. comm. ab hon. Aul. merc. angl. Ind. or. et à cl. Royle.)

38. *I. VESTITA* (Wall. ! cat. et herb. n. 72), caule erecto ramoso apice præsertim cum foliis villosissimo, foliis semiamplexicaulibus lanceolatis caulinis basi obtusis apice acuminatis irregulariter serratis utrinquè villosis, pedunculis asphyllis molliter hirsutissimis, invol. squamis lineari-subulatis scabridis ligulis linearibus barbatis. $\bar{\zeta}$? ad Kamaon et Srinaghur legit rev. Blindworth. Herba pedalis à basi ramosa. Ligulæ stylo subabortivo fortè steriles. Pappus albus

capillaris 1-ser. Achenium parvum teretiusculum glabrum. Hieracium hispidum Don prod. fl. nep. 165 ex Spreng. in litt. An rectè? (v. s. comm. ab hèn. Aul. merc. angl. Ind. or. et à cl. Royle.)

β. minor, caule digitali aut subpalmari simplici oligocephalo. — ad Deab Wall. ! l. c. var. B. (v. s. comm. cum var. α.)

39. I. ? NERVOSA (Wall. ! cat. et herb. n. 70), caule erecto villosa subsimplici, foliis ovali-lanceolatis basi attenuatis apice acuminatis penninerviis utrinque villosis scabridis serraturis callosis distantibus, capitulis paucis corymbosis terminalibus, pappi setis apice rigidis barbellatis. ½ in Nepaliâ legit cl. Wallich. Herba pedalis, caule tereti aut subcompresso. Invol. campanulati squamæ lanceolato-lineares exteriores subfoliaceæ dorso scabridæ. Ligulæ lineares angustæ glabræ. Achenium villosum teretiusculum. Pappus 1-serialis basi villis achenii quasi calyculatus rigidus basi subinteger, apice barbellis setæ. (v. s. comm. ab hèn. Aul. angl. Ind. or.)

40. I. ACUMINATA, caule erecto ramoso teretiusculo puberulo, foliis caulinis glabris integerrimis basi dilatatis subauriculatis cordato-semiamplexicaulibus apice lanceolatis acuminatis, capitulis ad apices ramorum solitariis, invol. squamis linearibus foliaceis laxiusculis, acheniis glabris. ½ in Indiâ prov. boreali-occid. legit. cl. Royle. Paulò accedit ad I. spiræifoliam et I. Germanicam sed foliis ad basin dilatatis et ad apicem acuminatis faciliè differt. (v. s. comm. à cl. Royle.)

41. I. JAPONICA (Th. fl. jap. 318), caule erecto villosa tereti, foliis sessilibus subamplexicaulibus oblongo-lanceolatis serratis villosis, pedunculis axillaribus monocephalis subpaniculatis tomentosis, invol. squamis linearibus subulatis villosis. ½ in Japoniâ. Banks ic. Kämpf. t. 30. Ben Fuku Kämpf. ante. 877. — Affinis videtur I. nervosæ Wall. sed differt foliis caulinis basi nec attenuatis nec subpetiolatis. Pappus indescr.

† Species non satis notæ.

42. I. QUADRIDENTATA (Lag. nov. gen. et sp. 30. n. 374), herbacea villosa-viscosa, foliis lanceolatis integris, floralibus linearibus, squamis invol. exterioribus laxis, corollulis 4-dentatis. (1) in arenosis ruderalis cultis Hispaniæ ab urbe Alonensi ad Murciam usque. An vera Inulæ species? An fortè Francoeuræ sp.?

43. I. PROVINCIALIS (Linn. sp. 1241), caule erecto 1-cephalo, foliis subserratis subulis tomentosis, radicalibus ovatis, caulinis paucis lanceolatis sessilibus. ½ in Collendaneis Galloprovinciæ. I. Provincialis Gon. videtur Seneciò uniflorus, sed species Linneæana omninò dubia! An fortè Pulicariæ odora var.?

44. I. MOLLIS (Bernhardi in Link enum. h. ber. a. p. 335), caule erecto tenui-tomentoso, foliis serratis, radicalibus in petiolum attenuatis, caulinis lanceolatis acutis sessilibus hirtis, invol. squamis lanceolatis hirtis extimis reflexis. ½ Patr. ign. †.

45. I. ? PRIMULÆFOLIA (Lam. dict. 3. p. 261), caule erecto herbaceo polycephalo, foliis spatulatis crenatis subulis incanis. — In pratis montosis Sancti-Domithi. — Plum. et Burm. t. 40. f. 1. Verisim. è genere excludenda, sed hodièrnia ignota.

†† Species exclusæ.

I. ACRIS Bernh. = Erigeron acre.

I. ÆSTUANS Linn. = Verbesina? æstuans.

I. APPENDICULATA B. Wall. = Vicia appendiculata.

I. APPENDICULATA C. Wall. = Vicia Indica γ.

I. ARABICA Linn. = Pulicaria Arabica.

I. ARGENTEA Pers. = Chrysopsis argentea.

I. AROMATICA Linn. = Printzia aromatica.

I. CANADENSIS Bernh. = Erigeron Canadensis.

- I. CAPENSIS Spreng. = *Pulicaria Capensis*.
- I. CAUCASICA Pers. = *Telekia cordifolia*.
- I. CERNUA Berg. = *Printzia Bergii*.
- I. CHRYSOCHOMOIDES Poir. = *Jasonia Sicula*.
- I. CÆRULEA Linn. = *Printzia Bergii*.
- I. CONYSEA Lam. = *Pulicaria dysenterica*.
- I. CONYZOIDES Desf. = *Pulicaria odora*.
- I. CRISPA Desf. = *Francoëuria crispâ*.
- I. CRISPA Sieb. = *Pulicaria undulata*.
- I. CYLINDRICA With. = *Pulicaria vulgaris*.
- I. DENTATA Sibth. = *Pulicaria dentata*.
- I. DIVARICATA Nutt. = *Heterotheca Lamarckii*.
- I. DUBIA Thunb. = *Aster Japonicus*.
- I. DYSENTERICA Linn. = *Pulicaria dysenterica*.
- I. FALCATA Pursh. = *Chrysopsis falcata*.
- I. FETIDA Linn. = *Erigeron foetidum*.
- I. GLANDULOSA Lam. = *Chrysopsis Mariana*.
- I. GLUTINOSA Pers. = *Grindelia glutinosa*.
- I. GNAPHALODES Vent. = *Strabonia gnaphalodes*.
- I. GOSSEYINA Michx. = *Chrysopsis gossypina*.
- I. GRAMINIFOLIA Michx. = *Chrysopsis graminifolia*.
- I. INCISA Lam. = *Pulicaria incisa*.
- I. INDICA Linn. = *Vicoa Indica*.
- I. LANATA Banks. = *Chrysopsis gossypina*.
- I. MACROPHYLLA hortul. = *Telekia cordifolia*.
- I. MARIANA Linn. = *Chrysopsis Mariana*.
- I. MONTANA Biv. = *Pulicaria odora*.
- I. ODORA Linn. = *Pulicaria odora*.
- I. PALUDOSA Link. = *Pulicaria dentata*?
- I. PINIFOLIA Linn. = *Senecio pinifolius*.
- I. PUNCTATA Muhl. = *Heterotheca scabra*.
- I. PULICARIA Brot. = *Pulicaria dentata*.
- I. PULICARIA Linn. = *Pulicaria vulgaris*.
- I. PULICARIA d'Urv. = *Pulicaria dysenterica*.
- I. QUADRIFIDA Ham. = *Francoëuria crispâ* β .
- I. REPENS Fisch. = *Pulicaria uliginosa*.
- I. SATURNIA Spreng. = *Lorentea satureioides*.
- I. SATUREIOIDES Mill. = *Lorentea satureioides*.
- I. SAXATILIS Lam. = *Jasonia glutinosa*.
- I. SCABRA Nutt. = *Heterotheca scabra*.
- I. SCABRA Wall. = *Vicoa appendiculata*.
- I. SERRATA Pers. = *Grindelia inuloides*.
- I. STOLONIFERA Gay. = *Aplopappus stoloniferus*.
- I. SUBAXILLARIS Lam. = *Heterotheca Lamarckii*.
- I. TRICHOPHYLLA Nutt. = *Chrysopsis trichophylla*.
- I. TRIKIS Linn. = *Trixis frutescens*.
- I. TUBEROSA Lam. = *Jasonia tuberosa*.

I. ULIGINOSA Stev. = *Pulicaria uliginosa*.

I. UNDULATA Linn. = *Pulicaria undulata*.

I. VILLOSA Vahl. = *Pulicaria Arabica*.

CCXXXIV. SCHIZOGYNE Cass. dict. 56. p. 23. Less. syn. 123. — *Conyzæ* et *Chrysocomæ* sp. auct. — *Gnaphalii* sp. Poir.

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, marginalibus paucis 1-seriatis tenuibus bidentatis fœmineis, cæteris 5-dentatis hermaphroditis. Recept. nudum angustum. Invol. oblongo-obconicum imbricatum, squamis colorato-scariosis oblongis acutis serrulatis glabris ciliolatis. Anthæræ basi breviter bisetæ. Achænium oblongo-cylindraceum glabriusculum. Pappus 1-serialis setosus scabridus. — Frutex Canariensis ramosissimus. Folia sparsa sessilia linearia obtusa integerrima. Corymbi terminales subpaniculati polycephali. Pedicelli squamulosi. Invol. citrina. Flores intensius lutei. — Genus differt à *Conyzâ* et *Phagnalo* floribus fœmineis uni- nec multiserialibus, à *Chrysocomâ* floribus non omnibus hermaphroditis. Ex antheris bisetosus-caudatis accedere videtur ad *Inulam* sed faciliè differt capitulis non radiatis, involucre imbricato, etc.

1. S. SERICEA, ramis foliisque tomento albo subsericeo undiquè obtectis. ♂ in insulis Canar. Teneriffâ, Palmâ, magnâ Canariâ legerunt et mecum comm. cl. Collignon, Ledru, Broussonet, Bory, Chr. Smith, Webb, et Berthelot, fortè etiâ ex h. Puer. in insulis Azoricis sed non in ins. Mauritiî aut ad Cap. Bonæ-Spei quidquid dixerit cl. Cassini. *Conyza sericea* Ait. kew. 3. p. 183. L'Hér. descr. mss. Link. Buch! can. 165. *Chrysocomâ sericea* Linn. f. suppl. 360. *Conyza tenuifolia* Salisb. prod. 194. *Conyza canescens* Bory! in litt. Sch. obtusifolia Cass. l. c. ips. test. in h. Merat. *Gnaphalium multiflorum* Poir. suppl. 2. p. 803 (ex specim. Ledru) et idè *Gnaph. Teneriffæ* Spreng. syst. 3. p. 468. *Gnaphalium compressum* Poir. suppl. 2. p. 804 (ex specim. Brousson.) ipso auctore teste et autogr. in h. Desf. Variat involucri glaberrimis aut extis subaraneosis. Cortex sapore acri contra odontalgiam ab incolis usitatus ex Linn. (v. s.)
2. S. GLABERRIMA, ramis foliisque glaberrimis. ♂ in insul. Canariensibus verisim. Teneriffâ. *Conyza Chrysocomæ linosyridi* affinis Webb et Berth. herb. Priori simillima et fortè varietas. Differt tamen glabritie perfectâ, foliis paulò brevioribus, corymbis oligocephalis. (v. s. in h. Webb. et Berth.)

CCXXXV. VARTHEMIA.

Capitulum multiflorum homogamum, fl. omnibus tubulosis 5-dentatis hermaphroditis. Recept. alveolatum. Invol. obovatum imbricatum, squamis adpressis acutiusculis gradatim longioribus, intimis disco æqualibus. Anthæræ basi bisetæ. Stigmata brevia inclusa. Achænium compressum oblongum pubescens. Pappus 1-serialis, setis subscabridis. — Suffrutex Persicus erectus ramosissimus glaberrimus. Folia sparsa pauca linearia aut oblongo-linearia integerrima patulo-deflexa. Capitula ad apices ramorum solitaria. Flores lutei. — Ex capitulis homogamis ad *Chrysocomam*, ex antheris basi bisetis ad *Inulam* accedit. — Genus dicavi veteri peregrinatori Lud. de Varthemo, qui ineunte seculo sexdecimo Persiam et Orientem ferè totum peragravit.

1. V. PERSICA. ♂ in Persiâ prope Teheran leger. cl. Olivier et Brugnière et recentius cl. Belanger. (v. s.)

CCXXXVI. *VICOA* Cass. ann. sc. nat. 1829. p. 33. dict. 60. p. 594. Less. syn. 199. Inulæ sp. Linn.

Capitulum multiflorum heterogamum, fl. radii angustè ligulatis fœmineis uniserialibus, disci tubulosis 5-dentatis hermaphr. Invol. imbricatum, squamis lineari-subulatis. Recept. ebracteolatum areolatum plano-subconvexum. Antheræ basi setoso-appendiculatæ. Achænium erostre. Pappus radii nullus, disci capillaris 1-serialis. — *Herbæ Indicæ* annuæ ramosæ erectæ. Folia alterna sagittato-auriculata semiamplexicaulia lenceolata aut linearia integra aut dentata. Rami parçè foliosi apice 1-cephali. Capitula flava.

1. *V. AURICULATA* (Cass. ll. cc.), foliis subdentatis oblongis acutis basi auriculatis, ligulis disco subæqualibus. (I) in Ceylonâ.

2. *V. INDICA* (DC. in Wight contrib. bot. ind. p. 10), foliis basi auriculatis lanceolatis acuminatis serratis subintegerrimisve utrinquè plus minùs puberulis, ligulis disco duplò longioribus. (I) per totam Indiæ orientalis peninsulam frequens. Capitula 3 lin. diam. (v. s.)

a. *serrata*, foliis per totam longitudinem serratis. Inula Indica Willd. sp. 3. p. 2092. Wight! herb. Inula Indica var. F. Wall.! cat. n. 74. Folia basi latè dilatato-auriculata in parte dilatata præsertim dentata.

β. *subintegra*, foliis subintegerrimis aut hinc indè serratis. — Inula Indica Linn. sp. 1237. Wall. cat. n. 74. — *Doronicum calcaratum* Roxb. fl. ind. 3. p. 434 ex Wight. Burm. ceyl. 124. t. 55. f. 2. Pluk. alm. t. 149. n. 4. f. 3. — *Aster Indicus* Willd. sp. 3. p. 2041 non Linn. ex Wight contrib. bot. ind. p. 10.

γ. *attenuata*, foliis subintegerrimis inferioribus basi angustatis ferè petiolatis nec auriculatis. — Inula appendiculata var. C. Wall.! cat. n. 76 lecta ad Prome.

δ. *aspera*, foliis subintegerrimis omnibus auriculatis suprâ sparsè et rigidè setosis, subtùs puberulis. Inula Indica var. G. Wall.! herb.

3. *V. AURITA*, foliis subintegerrimis omnibus basi latè et obtusè auriculatis angustis suprâ densè setoso-echinatis subtùs pubero-subtomentosis, ligulis disco paulò longioribus. — Inula auriculata Wall.! herb. n. 75. *Erigeron linearis* Spreng. in litt. Valdè accedit ad *V. Indicam* sed ex pubescentiâ subtomentosâ vid. diversa. (v. s. commun. ab hon. Cur. merc. angl. Ind. or.)

4. *V. APPENDICULATA*, foliis linearibus basi auriculatis acuminatis subintegerrimis margine revolutis utrinquè minutissimè et densè puberulis, ligulis disco duplò longioribus. (I) in Indiâ orientali ad ripas excelsas flum. Yenangham. Inula scabra Wall.! cat. h. bot. Inula appendiculata Wall.! cat. et herb. n. 76. var. B. Folia 3 poll. longa, 2 lin. lata. Capitula radio excepto 6-7 lin. diam. et ideò priori dopla. (v. s. commun. ab hon. Aul. angl. Ind. or.)

CCXXXVII. *PENTANEMA* Cass. bull. philom. 1818. p. 75. dict. 23. p. 565. 38. p. 373. Less. syn. 198.

Capitulum multiflorum heterogamum, fl. radii 1-ser. fœmineis angustè ligulatis puberis apice 3-dentatis, fl. disci tubulosis hermaphr. 5-dentatis. Recept. ebracteolatum subconvexum. Invol. subglobosi squamæ lineares imbricatæ villis longis mollibus hispidæ. Achænium erostre hispidulum. Pappus 1-serialis, setis 5 capillaribus. — *Herba Orientalis* annua erecta gracilis molliter hispida. Rami divaricati rubentes. Folia alterna obovata aut oblonga obtusa integerrima parçè villosa membranacea. Capitula pedunculata oppositifolia subglobosa flava. Nomen ex πέντε, quinque, et ἵμα, filum, et ergò neutri nec fœminiini generis.

1. *P. DIVARICATUM* (Cass. ll. cc.). ① in Oriente sine loci design. legit Æl. Michaux, et postea inter Alep et Bagdad cl. Olivier et Bruguère. Capitulum omnino Pulloarie trichoccephalæ sed pappus diversus. (v. s.)

CCXXXVIII. *FRANCOEURIA* Cass. dict. 34. p. 44. v. 38. p. 374. Less. syn. 198. — *Duchesnia* Cass. dict. 13. p. 545 non Smith et exal. ear. — *Inula* sp. Pers. — *Asteris* sp. Forsk.

Capitulum multiflorum heterogamum, fl. radii ligulatis breviusculis 1-ser. fœmineis, disci tubulosis 4-dentatis hermaphr. Recept. nudum planum. Invol. campanulatum imbricatum, squamis lineari-subulatis. Antheræ basi setoso-appendiculatæ. Achæmium erostre. Pappus caducus, setis 1-serialibus infernè subscabris supernè barbellato-subplumosis basi conferruminatis in anulum brevissimum setulis paleolisve minimis coronatum. — Suffruticulus Africanus erectus. Rami teretes 1-cephali divergentes. Folia alterna semiamplexicaulia oblonga dentata. Capitula flava.

1. *F. CRISPA* (Cass. ll. cc.), foliis margine undulato-crispatis. 5 et sæpè ①. In arenosis apricis Senegaliæ et Egypti. *Helenium Egyptiacum*, etc. Vaill. act. par. 1720. p. 304. n. 24. *Aster crispus* Forsk. æg. ar. descr. p. 150. *I. crispa* Pers. ench. 2. 450. excl. syn. Vent. Delille! fl. eg. p. 25. t. 45. f. 2. *Duchesnia crispa* Cass. dict. 13. p. 545. *Conyza undulata* Poir. suppl. 2. p. 339? Variat subherbacea aut fruticosa, tota subglabrata aut densè tomentoso-alba, foliis angustioribus latioribus semper margine crispatis. (v. s.)

Æ. *Indica*, foliis margine planiusculis. — ad Bar in prov. Bahar prope ripas Gangis legit cl. Hamilton. *Inula quadrifida* Ham. herb. Wall. cat. n. 73. (v. s. comm. ab hon. Aulâ merc. angl. Ind. or.)

CCXXXIX. *IPHIONA*. — *Iphionæ* sp. Cass. — *Chrysocomæ* sp. Forsk. — *Conyza* sp. Lam. — *Stæbelinæ* sp. Vahl.

Capitulum multiflorum homogamum, fl. omnibus hermaphroditis tubo stricto sensu subnullo, fauce longâ cylindricâ, dentibus 5. Recept. alveolatum angustum. Invol. imbricatum obovato-oblongum, squamis adpressis acuminatis seu ovalibus mucronatis. Antheræ basi breviter setoso-caudatæ. Filam. imæ cor. insertæ! Achæmium subcylindraceum striato-sulcatum parçè hispidum. Pappus rigidus multiserialis subscaber, setis exter. brevioribus, intimis gradatim longioribus corollam æquantibus. — Fruticuli Ægyptiaci glabri ramosissimi. Folia sparsa pauca subulata rigida pungentia basi auriculis 1-2 pungentibus instructa. Capitula ad apices ramorum subnudorum solitaria lutea. Pappus rufescens.

1. *I. JUNIPERIFOLIA* (Cass. dict. 23. p. 610), tota glabra, foliis apice subulato-pungentibus, inv. squamis obtusis mucronatis. 5 in Egypto prope Cahiram inter urbem et Soneys, versus Ramalieh, etc. (Delil! Acerbi! Radd!), in valle Hebran (Rupp.), in Arabiæ desertis circa montem Sinai (Bové!). Arabicè *Daffy* in Egypto, *Daffara* in Arabia dicta. *Conyza Memphisica*, etc. Vaill. act. par. 1719. p. 301. *Chrysocoma mucronata* Forsk. descr. 147. Fres. beitr. p. 77. *Stæbelina spinosa* Vahl symb. 1. p. 69. *Conyza pungens* Lam. dict. 2. p. 86. *Chrysocoma spinosa* Delil. fl. eg. p. 128. t. 46. f. 3. *Conyza pungens* et *Stæbelina spinosa* Pers. *Serratula spinosa* Poir. Iph. dubia Cass. bull. philom. 1817. oct. Jasoniæ sp.? Less. syn. 195. Folia variant basi uni- aut bispinosa et rariùs in medio adhuc 1-spinosa, summa semper squamiformia integerrima. (v. s. comm. à cl. Acerbi et Raddi.)

2. *I. SCABRA* (DC. in ann. sc. nat. 1834. bot. p. 263), tota pube brevi glandulosa

scabrâ, foliis subulato-pungentibus, invol. squamis linearilanceolatis glabris. ζ in Ægypti deserto circa Suez et in Arabiâ circa Tor mense junio flor. legit cl. Bové. *Chrysocoma*? Bové! pl. exs. n. 104. *Arabis* dicitur *Saffara* ex Bové. Priori habitu similis sed folia et capitula majora. Caulis ramosissimus. Folia linearilanceolata pungentia utrinquè ad basin dentibus 1-3 spinosis aucta. *Achænia* villosissima. (v. s. comm. à cl. Bové.)

† *Species exclusa.*

I. PUNCTATA Cass. = *Iqula* Africana.

CCXL. JASONIA. — *Jasonia Chiliadenus* et *Allagopappus* Cass. dict. 24. (1822) 34 (1825) et 56 (1828). — *Jasonia Allagopappus* et *Donia* sp. Less. syn. 197. — *Inula* et *Erigerontis* sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis uniserialibus aut homogamum fl. omnibus tubulosis 5-dent. hermaphrod. Recept. nudum subalveolatum planum. Invol. imbricatum disco circ. æquale, squamis linearibus acutis. Antheræ basi setis caudatæ. *Achænia* villosum oblongo-cylindraceum basi attenuatum. Pappus biserialis utrinquè setis filiformibus barbellatis, exterioribus brevissimis, inter. corollam ferè æquantibus. — *Herbæ* aut suffrutices humiles in regione Mediterraneâ aut Canariensi indigenæ, sæpius pilosi aut glandulosi ramosi, ramis patentibus apice 1-cephalis. Folia alterna sessilia oblonga aut linearia integra aut dentata. Flores lutei. — *An* genus in quatuor divellendum?

Sect. I. CHILIADENUS Cass. dict. 34. (1825) p. 34. — *Myriadenus* Cass. dict. 23. p. 565 non Desv. — *Orsina Bertol. giorn. h. nat.* 1829. v. 1. p. 362.

Capitula discoidea homogama. *Achænia* basi villosa apice glandulosa. Pappus inter. multi- (10-15) setosus. — *Caulis* herbacei.

1. *J. GLUTINOSA*, herbacea erecta glutinoso-pubera, foliis linearilanceolatis integerrimis sæpè obliquè tortis, ramis ad apicem usquè foliosis. \mathcal{U} in apricis rupestribus Galloprovinciæ, agri Ruscinonensis (Benth.), Catalanniæ (Pourr.), Siciliæ, Lopedusæ et Melitæ (Guss.). *Erigeron glutinosus* Linn. mant. 112. *Inula saxatilis* Lam. fl. fr. 2. p. 153. DC. fl. fr. ed. 3. v. 4. p. 154. *Chrysocoma camphorata* Rob. et Cass. in litt. *Chrysocoma saxatilis* DC. fl. fr. suppl. p. 468. *Chiliadenus camphoratus* Cass. dict. 34. p. 35. *Orsina camphorata* Bertol. giorn. sc. nat. Bol. v. 1. 1829, p. 362. opt. ex Guss. — *Bocc. sic.* t. 13. f. 4: et idè *Inula foetida* Zeraph. fl. melit. prod. p. 25. ex Bertol. — *Barr. ic.* t. 158. (v. s.)

Sect. II. EUJASONIA. — *Jasonia* Cass. dict. 24 (1822). p. 200.

Capitula radiata aut subdiscoidea. *Achænia* teretiuscula undiquè villosa. Pappus interior multi- (10-20) setus.

2. *J. TUBEROSA*, suffruticulosa erecta, radice præmorsâ incrassato-tuberosâ ligulosâ, foliis linearibus aut sublinearibus integerrimis, ramis parè foliosis ligulis involucri longioribus. \mathcal{U} ζ in apricis Galliæ mediterraneæ, Hispaniæ, Syriæ, etc. *Erigeron tuberosus* Linn. sp. 1212. Willd. sp. 3. p. 1961. *Inula tuberosa* Lam. dict. 3. p. 260. DC. fl. fr. ed. 3. v. 4. p. 153. *Jasonia radiata* Cass. dict. 24. p. 201. *Aster punctatus* Lapeyr. abr. pyr. p. 513. excl. syn. ex Benth. — *Lobel ic.* t. 250. f. 3. Variat canle vix palmari et culta pedali et sesquipedali. (v. v.)

3. *J. SIGUIA* (DC. in ann. sc. nat. 1834. bot. p. 261), herbacea erecta subvillosa.

radice gracili, foliis radicalibus lanceolatis dentatis, caulibus linearibus basi auriculatis semiamplexicaulibus integerrimis ramis parce foliosis. ☉ ? 2/ in paludosis maritimis arenosis Cretæ, Siciliæ, Hetruriæ, Galliæ et Hispaniæ mediterraneæ, Mauritanie et ad montem Sinai (Bové l. n. 118). Caulis 1-1 1/2 pedalis ramosus, purpurascens. Rami paniculato-corymbosi. Pappus biserialis, serie extima brevissima.

α. *discoidea*, floribus marginalibus subtubulosis aut in ligulam involucri brevioris fissis. — *Erigeron Siculum* Linn. sp. 1210. *Conyza Sicula* Willd. sp. 3. p. 1931. DC. fl. fr. 4. p. 129. Poir. suppl. 2. p. 358. *Jasonia discoidea* Cass. dict. 24. p. 201. et 39. p. 406. *Inula chrysocomoides* Poir. voy. 2. p. 239. Lam. dict. 3. p. 261. ill. t. 680. f. 3. Desf. atl. 2. p. 275. *Tubilium Siculum* Choisy! in h. DC. ined. — Boiss. sic. t. 31. f. 4. Magn. monsp. 76. ic. Pluk. alm. t. 168. f. 2. Species ab auctoribus sæpè confusa et characteres radii reverà variantes et fallaces. An *Dimorphanthès Sicula* Cass. dict. 13. p. 255. huc revocanda sed excl. car. gener.? (v. v. et s.)

β. *radiata*, floribus radii in ligulam involucri longiorem productis. — in Galliâ Narbonensi et Mauritanâ Tingitanâ. DC. fl. fr. 4. p. 129. Salzmann! pl. exs. *Aster pyrenaicus* Pourr.! in h. L'Hér. excl. syn. Ligulæ luteæ ferè similis priori. (v. s.)

Sect. III. *ALLAGOPAPPUS* Cass. dict. 56. p. 21. Less. syn. 197.

Capitulum discoideum homogamum. Achænium angulatum ad angulos villosum. Pappus inter. pauci- (5-7) setus, setis filiformis scabris, exter. brevis paucisetus. — Caulis fruticosi. Folia glandulosa. An genus proprium?

4. *J. DICHOTOMA*, fruticosa subdichotomè ramosa glabrescens, foliis linearibus basi attenuatis apice serratis, ramis floriferis squamas plurimas lineari-subulatas gerentibus, ligulis nullis. 5 in insulis Canariis, Palmâ, Gran-Canariâ et frequentior in insulâ Teneriffâ ad rupes Barancorum legerunt Ledru! Chr. Smith! Broussonet! Webb et Berthelot! nec in insulâ Mauritiâ ut incautè dixit Cass. ex h. Merat. *Conyza dichotoma* Linn. f. suppl. 359. *Chrysocoma dichotoma* Jacq. ic. rar. 1. t. 171. *Conyza inuloides* Ait. kew. 3. p. 182. *Conyza Canariensis* Willd. sp. 3. p. 1937. Link in Buch can. 148 et 165. *Allagopappus dichotomus* Cass. l. c. exc. patr. Antheris basi, jam teste Antonio, bisetis à Conyzâ et Chrysocomâ differt. Pappo biseriali extimo minimo plurisetoso ad Eojasoniam accedit. Folia utrinquè multiglandulosa. (v. s.)

Sect. IV. *DONIOIDES*. Doniæ sp. Less.

Capitulum radiatum, ligulis fœmineis. Achænium subangulatum hispidum. Pappus interior 7-10-setus, setis crassis ciliato-scabris, exterior plurisetus brevissimus.

5. *J. LÆVIGATA*, fruticosa glabra lævis, foliis oblongo-lanceolatis subcoriaceis integerrimis aut apice grossè serratis, pedunculis subaphyllis. 5 in ins. Teneriffâ ad Caput Teno rarior Broussonet! Webb et Berthelot! Folia basi subtrinerivia 1-2 poll. longa, 7-12 lin. lata. Inv. squamæ foliaceæ disco ferè longiores. *Euphthalmum lævigatum* Brouss. herb. Willd. mag. nat. ges. Berl. 1807. p. 138. *Donia Canariensis* Less. syn. 199. (v. s.)

CCXLI. *PULICARIA*. — Inulæ sp. Linn. — *Pulicaria* et *Tubilium* Cass. Less.

Capitulum multiflorum heterogamum, fl. radii fœmineis 1-serialibus ferè semper ligulatis, disci tubulosis 5-dent. hermaphr. Recept. nudum areolatum planiusculum. Invol. laxè imbricatum pauciseriale, squamis linearibus. Achænium erostre puberulum teretiusculum nec compressum. Pappus biserialis, serie exteriori coroniformi dentatâ brevissimâ, inter. setis 10-20 scabridis. — Herbæ pleræque Europææ nonnullæ Africanæ, in uliginosis præsertim nascentes, villosæ odoratæ.

erectæ ramosæ. Folia caulina cordato-sagittata integra aut dentata lanceolata. Pedunculi apice 1-cephali. Capitula flava.

Sect. 1. *PULICARIA*. — *Pulicaria* Gærtn. fr. 2. p. 461. t. 173. Cass. dict. 23. p. 565 et 44. p. 93. Less. syn. 197.

Capitula radiata, fl. angustè ligulatis.

1. *P. TRICHOCEPHALA*, caule demisso à basi ramosissimo cæspitoso, foliis lineari-sublanceolatis integerrimis acutis sessilibus pubescentibus, capitulis parvis foliis bracteatis lateralibus subsessilibus et terminalibus subglobosis, invol. squamis linearibus villosissimis, radii ligulis disco paulò longioribus, pappo ext. dentato, inter. 10-seto. (1) in Oriente sed locus propr. ignotus fortè Syria. Valdè accedit ad *P. Arabicam* sed differt præcipuè involuero villis longis albis hirsutissimo ferè ut in *Pentanemate*. Herba vix digiti longitudine. Radix perpendicularis subsimplex. (v. a.)
2. *P. ARABICA* (Cass. dict. 44. p. 94), caule erecto divaricato-ramoso, foliis oblongis cordato-semiamplexicaulibus mucrone acutis utrinquè plùs minùs pubescenti-villosis integerrimis, pedunculis axillaribus terminalibusque, invol. squamis lineari-subulatis puberulis, radii ligulis disco paulò longioribus, pappo ext. dentato, inter. 10-seto. (2) in Arabiâ? (Forst. ex Linn.), in cultis montosis ad Sinâi (Bové), Ægypte (Delil. Coqueb.), Crêtâ (Oliv. et Br.), Mauritanîâ (Desf. Salsm.), Andalusiâ (Hæussel). *Inula arabica* Linn. mant. 114? (exol. syn. Pluk. et ideò patr. in Indiâ indic.) Desf. fl. atl. 2. p. 276. *Inula villosa* Vahl ex Horn. h. hafn. 2. p. 822 non Nutt. *Pul. villosa* Link enum. h. berol. 2. p. 336. Variat caule decumbente aut subdiffuso. Capit. 5 lin. diam. Valdè dubium mihi videtur stirpem in Arabiâ habitare. Forskahl in florâ suâ Arabicâ hanc non enumerat. Descr. Linnæi est etiam valdè manca. (v. s.)
3. *P. DESERTORUM* (DC. ann. sc. nat. 1834. bot. p. 263), caulibus erectis ramosis pubescenti-villosis, foliis semiamplexicaulibus oblongis grossè et obtusè dentatis undulato-crispis villosis, pedunculis 1-cephalis, capitulis depressis hemisphæricis, invol. squamis linearibus acuminatis, ligulis invol. paulò longioribus, pappo ext. brevissimo subintegerrimo, inter. 10-seto. (3) in Arabiæ desertis circa montem Sinâi legit cl. Bové mense junio flor. Arabis *Desdas* dicitur. *Inula Arabica* Bové! pl. exs. n. 109. Ex loco natali est verè Arabica sed differt à stirpe sub hoc nomine adnødum admissâ et fortè in Arabiâ nunquam repertâ. (v. s.)
4. *P. VULGARIS* (Gærtn. fr. 2. p. 461. t. 173), caule paniculato erecto, foliis semiamplexicaulibus oblongis acutis undulatis subintegerrimis villosis, pedunculis 1-cephalis oppositifoliis, capitulis hemisphæricis, radii ligulis brevissimis, pappo ext. dentato, inter. circ. 15-seto. (4) in uliginosis viis et plateis Europæ temperatæ et australis usque ad Caucasum (Bieb. Stev.), et ad Sibiriam Altai-cam propè Semipalatinsk (C. A. Mey.), in Teneriffâ fortè introducta (Brouss.). *Inula Pulicaria* Linn. sp. 1238. Fl. dan. t. 613. Engl. bot. t. 1196. *Aster pulicarius* Scop. caru. n. 1080. *Inula cylindrica* With. Hnds. et *Inula uliginosa* Sibth. ex Sm. fl. brit. 892. *Conyza minima* Dod. pempt. 52. Variat capitulis flosculosis ex Sm. fl. brit. Variat præcipuè invol. in region. frigidioribus glabrescentibus aut molliter hirsutis in reg. calidioribus. An fortè species duæ hic adhuc confusæ? (v. v. et s.)
5. *P. ULLIGINOSA* (Stev. in litt.), caulibus erectis apice tantùm paniculato-ramosis, foliis caulinis cordato-aureolatis semiamplexicaulibus lanceolatis acutis subserratis subtùs præsertim velutino-tomentosis, capitulis pedunculatis terminalibus et axillaribus, invol. squamis lineari-subulatis, radii ligulis disco duplò ferè longioribus, pappo ext. dentato, int. 15-seto. (5) in Tauriâ meridionali ad fossas aquosas legit cl. Steven. Eandem ni fallor habeo lectam ad ripas Jordanis et prope Beyrouth in Bové pl. exs. n. 441? *P. dysenterica* Bieb.

suppl. p. 574 ex Stev. *Inula repens* Fisch. ex Trevir. act. nat. cur. 13. p. 203. Ferè media inter *P. vulgarem* et *dysentericam*. (v. s. comm. à cl. Steven.)

6. *P. ANGUSTIFOLIA*, tota subcinereo-pubescent, caule basi suffruticuloso ramoso, foliis sessilibus oblongo-linearibus integerrimis aut hinc inde dentatis, pedunculis subnudis 1-cephalis folio triplò longioribus, invol. squamis foliaceis, ligulis invol. longioribus, pappo ext. denticulato, inter. 10-13-seto. ♂ ♀ in Indiæ prov. boreali-occident. legit cl. Royle. Accedit ad *P. uliginosam* sed foliis sessilibus nec cordato-auriculatis capitulis longius pedunculatis et invol. squamis latioribus faciliè differt. (v. s. comm. à cl. Royle.)
7. *P. DYSENTERICA* (Gærtn. fr. 2. p. 462), caule paniculato erecto, foliis caulinis aurito-amplexicaulibus ovali-oblongis subintegerrimis callosis-acutis subtus incano-tomentosis, pedunculis axillaribus et terminalibus 1-cephalis, squamis invol. apice setaceo-cuspidatis, radii ligulis disco duplò triplòve longioribus, pappo ext. dentato, inter. circa 20-seto. ♀ in Europæ ferè totius uliginosis, fossis, humidis. Cass. dict. 44. p. 93. *Inula dysenterica* Linn. sp. 1237. DC. fl. fr. 4. p. 149. Fl. dan. t. 410. Engl. bot. t. 1115. Hayn. arzn. 6. t. 46. Schknh. hand. t. 247. ic. fr. *Aster dysentericus* All. pedem. n. 711. *Inula conyzæa* Lam. fl. fr. 2. p. 149. *Aster undulus* Moench. *Inula pulicaria* D'Urv. ! enim. n. 816. excl. syn. Variat foliis longioribus coloribus. An In. planifolia Link ab hac satis differt? — Fuchs hist. 436. ic. (v. v.)

β. *breviradiata*, ligulis invol. paulò longioribus aut eo subæqualibus. ♀ circa Genevam ad pagum Vezenaz obs. Reuter. (v. s.)

8. *P. UNDULATA*, albo-lanuginosa, caule erecto paniculato, foliis cordato-auriculatis semiamplexicaulibus lanceolatis undulato-crispis dentatis, pedunculis squamiferis terminalibus et axillaribus, invol. squamis linearibus acutis subcaehris, radii ligulis disco paulò longioribus, pappo ext. dentato, inter. 15-20-seto. (1) (2) in desertis Ægypti. *Inula undulata* Linn. mant. 115. Lam. dict. 3. p. 256. Delil. fl. eg. 126. t. 46. f. 1. excl. var. β. *Helenium Ægyptiacum* tomentosum, etc., Vaill. act. Par. 1720. p. 305. *Inula crispa* Sieb. ! herb. Ægypt. I. odora Forsk. fl. æg. p. 150? (v. s.)

9. *P. INCISA*, pubero-villosa, caule erecto paniculato, foliis basi dilatata cordato-semiamplexicaulibus oblongo-linearibus acutis plus minus incisè dentatiave planiusculis, pedunculis apice subaphyllis terminalibus et axillaribus, invol. squamis linearibus acutis glabriusculis, radii ligulis disco paulò longioribus, pappo ext. dentato, inter. 15-seto. (1) in Senegalià propè St.-Louis legerunt cl. Roussillon et Perrottet. *Inula incisa* Lam. ! dict. 3. p. 256. In. undulata β Pers. Delil. Valdè affinis I. undulata sed nec albo-lanuginosa nec undulatifolia. (v. s.)

10. *P. ODORA* (Reich. fl. exc. 2. n. 1532), caule erecto subsimplici oligocephalo, foliis villosis remotè denticulatis, radicalibus petiolatis ovatis, caulinis auriculato-amplexicaulibus oblongis lanceolatis, pedunculis subternis axillaribus et terminalibus, radii ligulis disco duplò longioribus, pappo ext. 10-dentato, inter. 10-seto. ♀ in apricis agri Ruscinonensis (Duby bot. gall. 1. p. 268). Galloprovinciæ, Liguriæ, Istriæ (Reich.), Græciæ (Sibth.; h. Acad. mon. 1). Aleppi, Constantinopoli, Etruriæ, agri Neapolitani (Ten. syll. p. 464), Siciliæ (Guss. !), Ischiæ (Schouw. !), Mauritanix, Corsicæ, nec in Burgundiâ ut falso indicavit Durande. *Inula odora* Linn. sp. 1236. *Aster odoratus* All. ped. n. 713. *Inula dumetorum* Salz. ! herb. — Barr. ic. t. 1145. bona. An fortè huc referenda *Pulicaria calycina* Presl. del. 96, quæ *Inula montana* Bivon. rar. cent. 1. p. 34. non Linn. et *Inula calycina* Spreng. syst. 3. p. 521 et I. conyzoides Desf. cor. Tourn. p. 49. t. 38? quæ *Aster Creticus conyzoides* flore magno *Asphodeli* radice Tourn. cor. 36. (v. v.)

11. *P. CAPENSIS*, caule erecto basi simplici apice corymboso folisque præsertim superioribus utrinquè densè sericeo-villosis, foliis cordato-amplexicaulibus mucrone calloso apiculatis et calloso-serrulatis, invol. squamis linearibus acuminatis laxè imbricatis pedunculisque villosissimis, ligulis linearibus involucre paulò longioribus, achæniis villosis, pappo ext. 10-dentato inter. 15-seto.

24 ad Cap. Bonæ-Spei ad fossas juxta montem Diaboli (Eckl. ! it.), in distr. Worcester (Eckl. !), ad Kleindrakenstein (Drege !), ad Port-Natal (Drege !), et in coloniæ distr. orient. (Burch. ! cat. geog. n. 4407). Herba 1-2-pedalis. Capitula 5-6 lin. diam. Variat foliis caulibusve plus minus sericeo-tomentosis et incanis. An fortè daz species distinguendæ ? Species simillima *P. inuloidi* sed radiata. (v. s.)

12. *P. ERIGEROIDES*, caule erecto basi simplici apice subpaniculato pubescente, foliis semiamplexicaulibus oblongo-lanceolatis remotè denticulatis aut subintegerrimis minutissimè puberulis, invol. squamis lineari-subulatis subscabridis, ligulis angustis parvis, pappo exter. 10-dentato, achæniis vix pilosiusculis. 24 ? in Africâ Capensî Burch. ! cat. n. 4408 et 1949. Eckl. ! Drege ! *Erigeron scabrum* Thunb. fl. cap. 665. Affinis *P. vulgari* et *P. Capensî* sed ab utrâque sat diversa. (v. s.)

13. *P. DENTATA*, caule erecto ramosissimo, foliis amplexicaulibus acuminatis dentatis, capitulis hemisphæricis, radii ligulis longitudine involucri. 24 in insulâ Crêtâ. *Inula dentata* Sibth. fl. græc. t. 874 (ined.) ex Smith prod. 2. p. 181. *Inula pulicaria* ß Linn. sp. 1238. syn. dub. An fortè *In. paludosa* Link. seu *In. Pulicaria* Brot. huc referenda ? †.

Sect. II. *TUBILIUM* Cass. bull. philom. 1817. p. 153. dict. 23. p. 565. v. 56. p. 19. Less. syn. 197.

Capitulum discoideum, fl. radii 3-5-fidis stamina effreta sæpè gerentibus. *Achna glabruscula*. — Sectio cum *Pulicariâ* conjungenda eodem jure ac *Conyzâ squarrosa* cum *Inulâ*.

14. *P. INULOIDES*, caule herbaceo erecto puberulo, foliis semiamplexicaulibus obtusè auriculatis lineari-oblongis integerrimis aut dentato-subcincis vix pubescentibus, ramis 1-cephalis paucis subcorymbosis, invol. squamis lineari-subulatis villosis, fl. radii irregulariter tubulosis cæteros paulò asperantibus. 24 ad Mogador (nec in Teneriffâ) legit cl. Broussonet ! *Erigeron inuloides* Poir. ! suppl. 5. p. 464. *Tubilium angustifolium* Cass. ! dict. 56. p. 20. Simillima *P. Capensî* et etiâ ad *Conyzam* Gonani habitu accedens. Fol. 2 poll. longa, 2 lin. lata. (v. s. comm. à cl. Broussonet.)

15. *P. FOLIOLOSA*, tota cinereo-villosa, caulibus herbaceis ramosissimis apice molliter hispidis, foliis oblongis acutis utrinquè puberulis, inf. basi attenuatis subdentatis, super. integris basi auriculatis semiamplexicaulibus, invol. squamis lineari-subulatis villosis, fl. radii tubulosis tenuissimis cæteris ferè brevioribus, pappo ext. coroniformi dentato, inter. 5-10-seto. 24 ? in Indiâ orient. prope Patna legit cl. Hamilton, et in prov. bor.-occident. cl. Royle. *Conyza foliolosa* Ham. ! in Wall. cat. n. 3281. comp. 391 non 3092 nec Willd. Anth. setas ex cor. minutie non vidi sed ex habitu et pappo hic refero speciem. Cor. apice subglandulosa. (v. s. in h. Cur. Ind. nunc Soc. Linn. Lond.)

16. *P. HYDRABADENSIS* (Lindl. ! in litt. 1835), caule herbaceo erecto tereti ramoso subpubero, foliis lanceolatis puberulis, infer. obtusis dentatis basi in petiolum attenuatis, super. sessilibus angustis acutis subintegris, ramis parce foliosis apice monocephalis, invol. squamis linearibus acuminatis, exter. foliaceis dorso puberulis, inter. siccis glabris. ① in Indiâ orient., ad Hyderabad (ex nom.) et ad muros Seringapatam (h. madr. !). *Aster Hydrabadensis* Wall. ! cat. n. 2973. comp. n. 83. Recept. latum eleganter punctatum aut brevè alveolatum. Pappus interior rufus rigidulus deciduus coronâ albidâ cinctus. Cor. tubulosæ brevissimè 5-dentatæ. Anth. tenuissimè caudatæ. (v. s. comm. ab hon. Aul. merc. angl. Ind. or.)

† *Species exclusæ.*

P. ANNUA Gært. = *Stenactis annua*.

P. CALYCINA Presl. = *Inula montana*.

CCXLII. STRABONIA.

Capitulum multiflorum heterogamum, fl. radii subligulatis difformibus minimis 1-serialibus fœmineis, disci tubulosi 5-dentatis hermaphroditi. Recept. planum nudum. Invol. squamæ lineares pauciseriales disco dimidiò breviores. Antheræ basi bisetæ. Achænium compressum villosum apice obtusum basi subangustatum. Pappus collo vix notando ab achæni apice distinctus duplex, serie ext. brevissimâ dentatâ subcoroniformi, int. setis pluribus densè et crassè barbellatis 1-serialibus. — Herba Persica multicaulis erecta tota lanugine albâ molli densè vestita, pedicellis involucriisque subglabris pubescentibusve. Folia obovato-spathulata obtusissima undulata semiamplexicaulia. Capitula ad apices ramulorum solitaria. Flores flavi. — Genus achænio compresso nec teretiusculo ab Inuleis plerisque diversum. Dicavi memorie cl. Strabonis inter veteres geographi insignis.

1. *S. GNAPHALODES*. (1) ex Vent. sed 2/2 vid. ex radice crassâ tereti subligulâ. In Persiâ inter Teheran et Isbahan ex herb. Oliv. et Brug. prope Tegrich ad radices montium Albourghs ex semin. ad Cels missis. Innula gnaphalodes Vent. pl. rar. jard. Cels n. et t. 75. (v. s. ex h. Oliv. et Brug.)

CCXLIII. PEGOLETTIA Cass. dict. 38. p. 230. Less. syn. 200.

Capitulum multiflorum homogamum. Recept. ebracteolatum punctatum planum. Invol. 2-3-seriale imbricatum floribus brevius, squamis lineari-subulatis. Cor. tubulosæ 5-fidæ subringentes. Antheræ basi longè bicaudatæ. Stylus Inulearum. Achænium cylindraceum sulcatum subangulatum erostre. Pappus conformis biserialis, paleis seriei ext. brevibus basi paleaceis, inter. elongatis piliformibus barbellatis ferè plumosis. — Fruticuli vel herbæ Africani. Folia alterna punctata elliptica aut linearia. Capitula ad apices ramulorum solitaria. Flores flavi.

1. *P. SENEGALENSIS* (Cass. dict. 38. p. 230), herbacea erecta tota villis brevibus hispida subglutinosa, foliis sessilibus lineari-sublanceolatis acutis vix apice subdentatis, pappo biseriali, inter. subplumoso, ext. brevi paleaceo barbato. (1) in arenosis secus flumen Senegal in regione Walo, ad Dagana? legit indef. Perrottet. Herba pedalis paniculato-ramosa. Radix subsimplex. Flores flavi senescentes subpurpurei. (v. s.)
2. *P. OXYODONTA*, herbacea, basi glabra supernè puberulo-glandulosa viscosa, foliis ovali-lanceolatis utrinquè acuminatis dentibus utrinquè 2-3 grossis acutis exsertis callosis, invol. squamis acuminatis, floribus in invol. 35-40, pappo biseriali exteriore brevissimo parco obtuso. — in Africâ austr. ad flum. Gariep, in Klein-Namaqualand, legit cl. Drege. Folia 2 poll. longa, 7 lin. lata. Achænium teres sulcatum hinc indè pilosiusculum. (v. s.)
3. *P. ACUMINATA*, suffruticosa versus apices glandulosa, foliis oblongis integris utrinquè acuminatis, invol. squamis longè acuminatis apice laxis, pappo biseriali ext. breviori. 3 in Africâ australi ultrâ flum. Gariep legit cl. Burchell, n. 2396. Ramuli et folia puberulo-glandulosa. Achænium glabratum striatum teretiusculum. (v. s.)
4. *P. POLYGALÆFOLIA* (Less. syn. 200), fruticosa glabra, ramis tortuosis (aut ex Thunb. retrofractis), foliis petiolatis obovatis ovalibusve acutis integris, invol. squamis acuminatis, pappo biseriali ext. brevi subpaleaceo. 3 ad Cap. Bonæ Spei (Th. Licht.), ad Klein-Fischrivier et Zwarteberge (Drege!). Eupatorium retrofractum Thunb. fl. cap. 628. Vernonia polygalæfolia Licht. naa. ex Less.
5. *P. SACCHARIDIFOLIA* (Less. syn. 200), fruticosa, foliis petiolatis ovalibus obo-

vatis viscidis grossè dentatis, invol. squamis acutis, circ. 2-3-serialibus pappo pluriseriali, exter. vix distincto. ♀ ad Cap. Bonæ-Spei (Krebs! pl. exs. n. 191), ad Fischervier et Zneureberge (Drege!) Pteronia dentata Spreng. in h. Moric. Valde priori ex habitu affinis, sed viscosa, folia dentata et pappi series externa vix ab inter. distincta. (v. s.)

6. P. ? TENELLA, herbacea erecta tenuis sparsè hispida, foliis elliptico-oblongis utrinquè attenuatis integerrimis, ramis apice longè aphyllis pubescentibus x-cephalis, invol. squamis biserialibus acuminatis, pappo biseriali, exteriorè brevissimo. — in Africâ austr. legit cl. Drege, n. 5159, sine loci design. Flores non vidi unde genus subdubium. Recept. nudum punctatum. Achænium teres substriatum. Pappi setæ crassæ nivæ. (v. s.)

CCKLIV. GEIGERIA *Griesselich in linnæa* 1830. p. 411. — *Geigera Less. syn.* 199. — *Zeyhera Spreng. f. diss.* 1828 non *Mart.* 1826. — *Dizonium Willd. herb. ex Schlecht.*

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci tubulosi regulariter 5-dent. hermaphr. Recept. conicum bracteolatum piloso-fimbriiferum. Invol. imbricatum, squamis ext. bractæformibus foliaceis laxis, cæteris adpressis. Achænium subtrigonum erostre basi attenuatum pilis vestitum. Pappi paleæ 8-12 biseriales obovatæ apice obtusæ et ex Griess. subincisæ. — Herba humilima Capensis, radice seu rhizomate lignoso simplici, caule brevissimo subdichotomo. Folia linearia conferta integerrima. Capitula fasciculata sessilia. Flores lutei.

1. G. AFRICANA (Griess. l. c.). ♀ ad Cap. Bonæ-Spei. Zeyher! coll. exs. (1828) n. 147 in h. Moric. Eckl. pl. exs. n. 650. *Zeyhera acaulis Spreng. f. l. c. Dizonium longifolium Willd. herb. ex Less.* (v. s.)

DIV. II. CÆSULIÆ.

Capitula 1-flora in glomerulum aggregata, involucri proprio bivalvi demum cum achænio concreto. — Locus in serie naturali dubius.

CCKLV. CÆSULIA *Roxb. corom.* 1. p. 64. t. 93. *Cass. dict.* 6. suppl. p. 4. *Brown trans. lin. soc.* 12. p. 94. *Less. syn.* p. 200. — *Meyeræ sp. Don. Spreng.*

Capitula plurima uniflora adgregata in glomerulum seu capitulum generale. Involucrum proprium valvis 2 carinatis membranaceis inter se basi et demum cum ovario concretis tunc pappum Bidentis more bipartitum simulantibus. Cor. tubulosa, fauce ovatâ, lobis 5 patulis. Styli rami filiformes divergentes glabri. Antheræ rigidulæ, basi setis lævibus caudatæ. Achænium compressum epapposum quasi bialatum ex involucri valvis concretis. — Herba Indica glabra adscendens. Folia alterna lanceolato-linearia utrinquè acuminata distanter serrata, basi ad florum originem dilatata semiamplexicaulia. Ramuli florales axillares brevissimi; folia floralia 2-3 colorata orbiculata capitulum generale adjectis bracteis paucis parvis cingentia. Flores pallide violacei aut albi à centro ad peripheriam evoluti. — Ad Heliantheas et deindè ad Vernoneas dubitanter refert Cass.; Less. ad Asteroideas Inuleas. Stylus enim non vid. Vernoniaceæ et antheræ Inulearum, sed habitus et infloresc. valde diversi. Habitus ferè Cryphiospermi sed alternifolia.

1. C. AXILLARIS (Roxb. l. c. et in fl. ind. 3. p. 447). ♀ in locis subhumidis *Coromandelis*, in orysetis prope Tanjore (Wight!) et in paludosis Nepalice inferioris regione subsylvaticâ Turraye. Wallich! cat. n. 2183. comp. 143.

Andr. bot. rep: t. 431. *Meyera orientalis* Don prod. fl. nep. 180. *Melananthera orientalis* Spreng. in litt. ad Wall. 1830. *Folia* 3 pollices longa, 3 lin. lata. (v. s. comm. ab hon. Aulâ angl. Ind. or.)

Species exclusæ.

C. RADICANS Willd. = *Cryphospermum repens*.

C. SCANDENS Poepp. = *Wulfia scandens*.

Subtrib. V. *BUPHTHALMÆ*. — Inulæ *Bupthalmæ* §. 1. *Cass. dict.* 23. p. 566. — *Bupthalmæ* Less. *syn.* p. 209 (*adm. Ceruana*). — *Bupthalmi* Linn. *spec. Gerontogæ*.

Capitula multiflora heterogama (aut in Anvilleâ homogama), fl. marginalibus scæmineis sæpius ligulatis, in Ceruanâ tubulosus 3-fidis. Anth. nunc longè nunc breviter caudatæ nunc ecaudatæ (in *Lepidochætâ* et *Ceruanâ*). Recept. totum paleaceum, paleis floris discum subsequantibus. Achænium erostre aut brevissimè rostratum. Pappus coroniformis lacerus aut pilosodentatus. — An genera infrâ indicata pro sect. tantum habenda? — Herbæ aut suffrutices. Folia in omnibus ritè cognitis alterna integra aut dentata. Capitula lutea. — Species omnes *Gerontogæ*.

CCXLVI. *BUPHTHALMUM* Neck. *elem.* 1. p. 20. *Cass. dict.* 34. p. 276. Less. *syn.* 209. — *Bustia* sp. *Adans.* — *Asteroidis* sp. *Tourn.*

Radius 1-serialis, ligulis latiusculis; corollarum omnium tubus teres internè angustatus. Antheræ obsoletissimè caudatæ. Achænium radii triquetrum angustè trialatum, disci plano-compressum margine interiori unialatum. Pappus scarosus coroniformis pilosodentatus conformis. — Herbæ perennes glabriusculæ. Folia lineari-lanceolata subintegra. Capitula terminalia solitaria. Invol. squamæ pauciseriales longè acuminatæ. — Genus habitu *Inulas* referens.

1. *B. SALICIFOLIUM* (DC. fl. fr. 4. p. 218), caule herbaceo, foliis lineari-sen oblongo-lanceolatis subdentatis, achæniis radii apice obtusis subtruncatis. 2½ in Europâ ferè totâ ad radices montium. Jacq. fl. austr. t. 370. *B. grandiflorum* Schkuhr handb. t. 257. *B. grandiflorum* et *salicifolium* Linn. sp. 1275. Lam. dict. 1. p. 516. Reichenb. fl. exc. 2. p. 236. Gaud. fl. helv. 5. p. 380. Variat caule simplici aut ramoso, subvillosa aut glabra, ramis floridis erectis aut subpatulis, foliis acuminatis aut rariùs obtusis integerrimis aut dentatis, subvillosis aut glabris, latioribus et angustioribus, basi subsemiamplexicaulibus aut subangustatis, sed hæ variationes adeò in specim. variis combinantur ut species auctorum vix inter varietates veras recipiendæ videantur. (v. v.)
2. *B. INULOIDES* (Moris! sard. app. 1828), caule basi suffruticuloso, ramis foliis involucrique pube brevî hirtis viscosis, foliis infer. ovatis grossè dentatis petiolatis summis lanceolato-linearibus integerrimis sessilibus, achæniis radii supernè in cornu duo acuminata productis. 2½ 3 inter fissuras rupium calcar. insulæ Tavolara prope Sardiniam. Pappus irregulariter coroniformis inæqualis dentato-ciliatus. Paleæ minùs acuminatæ quàm prioris. (v. s. comm. à cl. Moris.)
3. *B. ANGUSTIFOLIUM* (Pursh fl. 2. p. 564), foliis alternis linearibus supernè latioribus integerrimis glabris, invol. squamis acutè lanceolatis. — in *Georgiâ* et *Floridâ* ex h. Banks in quo Pursh descr. Elliotio et aliis bot. amer. ign. An *Mnesitheon album* Raf. fl. Ind. 68? ex Spreng.

† *Species non satis notæ.*

4. *B.? LONGIPES* (Cass. bull. philom. 1822. p. 144. dict. 34. p. 277), caule herbaceo

repente, foliis alternis petiolatis cordiformibus obtusis trinerviis inæqualiter crenatis, pedunculis elongatis oppositifoliis 1-cephalis, 1-bracteatis. — In ins. Madagascar. Certè ex Cass. priorum congener sed habitus diversissimus et fructus indesciptions dubitationem suadent.

5. B.? MELISSOIDES (Poir. suppl. 1. p. 747), caule herbaceo aspero, foliis oppositis longè petiolatis cordato-ovatis obtusè crenatis asperis, pedunculo longissimo 1-cephalo. — Patr. et car. pleriq. ignot. An *Helianthea*? fortè *Wedelia* sp.?
6. B.? OLERACEUM (Lour. fl. coch. 2. p. 618), caule herbaceo erecto lævi, foliis oppositis lineari-lanceolatis inæqualiter dentatis recurvis glabris, capitulis solitariis, invol. squamis acutis membranulâ laterali connexis. — Cultum in hortis Cochinchinæ et Chinæ. — Discus flavus. Radius albus. Recept. paleæ minimæ. Toto cœlo differt à B. graveolente ad quod Willd. refert et imò à genere omnino alienum videtur.
7. B.? RAMOSUM (Forsk.æg. ar. descr. 151), caule herbaceo hispido, foliis oppositis lanceolatis scabris enerviis denticulatis subciliatis, capitulis terminalibus pedunculatis solitariis, invol. squamis ext. foliaceis radio parum brevioribus. — In Arabiâ felici. Vahl symb. 2. p. 92. Ligulæ circ. 8 spice tridentatæ purpureo-nervosæ.
8. B.? SAGITTATUM (Pursh fl. bor. am. 2. p. 564), tomentosum, foliis radicalibus longissimè petiolatis oblongis sagittatis integerrimis subtrinerviis, caulinis oblongis in petiolum attenuatis, caule subtricephalo, invol. squamis exter. disco longioribus. 2ℓ in Amer. bor. montibus dictis Rocky-Mountains. Capitula magna lutea (Pursh). Non est *Bupthalmum* (Nutt.), sed quid? †.

B. MELITENSE Ræusch. }
B. ORTEGÆ Spreng. } nomine tantum nota.

†† Species exclusæ.

- B. AQUATICUM Linn. = *Asteriscus aquaticus*.
 B. ARABICUM Delil. = *Anvillea Garcini*.
 B. ARBORESCENS Linn. = *Borrichia arborescens*.
 B. ASPERRIMUM Spr. = *Anomostephium bupthalmoides*.
 B. ASPERUM Horn.? = *Wedelia hispida*.
 B. ASTEROIDEUM Viv. = *Pallenis spinosa*.
 B. AUSTRALIS Spreng. = *Wollastonia aristata*.
 B. BONARIENSE Pers. = *Acicarpa spathulata*?
 B. CAPENSE Linn. = *Oederia prolifera*.
 B. CANUM L'Hér. herb. = *Borrichia argentea*.
 B. CHRYSANTHEMOIDES Poir. = *Asteriscus imbricatus*.
 B. CORDIFOLIUM W. et Kit. = *Telekia cordifolia*.
 B. CRASSIFOLIUM Raf. = *Asteriscus maritimus*.
 B. DIFFUSUM Vahl mss. = *Eclipta procumbens* β.
 B. DURUM Linn. = *Osteospermum spinosum*.
 B. ELEGANS Brouss. = *Asteriscus imbricatus*.
 B. FLOSCULOSUM Vent. = *Anvillea Garcini*.
 B. FRUTESCENS Linn. = *Borrichia frutescens*.
 B. GARCINI Burm. = *Anvillea Garcini*.
 B. GRAVEOLENS Vahl = *Asteriscus graveolens*.
 B. HELIANTHOIDES Linn. = *Heliopsis lævis*.
 B. HETEROPHYLLUM Willd. = *Wedelia pedunculosa*?

- B. *HELIANTROIDES* Forst. = *Wollastonia aristata*.
 B. *IMBRICATUM* Cav. = *Asteriscus imbricatus*.
 B. *INCANUM* Mill. = *Borrichia frutescens*.
 B. *LÆVIGATUM* Brouss. = *Jasonia lævigata*.
 B. *LINEARE* Willd. = *Borrichia frutescens* β?
 B. *MARITIMUM* Linn. = *Asteriscus maritimus*.
 B. *ODORUM* Schousb. = *Asteriscus odoratus*.
 B. *PERSICUM* Garc. = *Anvillea Garcini*.
 B. *PERUVIANUM* Lam. = *Borrichia Peruviana*.
 B. *PRATENSE* Vahl = *Ceruana pratensis*.
 B. *REPENS* Lam. = *Wedelia carnosa*.
 B. *RAMTILLA* Ham. = *Guizotia oleifera*.
 B. *SCABRUM* Cav. = *Wedelia hispida*.
 B. *SERICEUM* Linn. f. = *Asteriscus sericeus*.
 B. *SESSILE* Mill. = *Asteriscus maritimus*.
 B. *SPECIOSISSIMUM* Linn. = *Telekia speciosissima*.
 B. *SPECIOSUM* Schreb. = *Telekia cordifolia*.
 B. *SPINOSUM* Linn. = *Pallenia spinosa*.
 B. *STENOPHYLLUM* Link. = *Asteriscus sericeus* γ.
 B. *STRIGOSUM* Spreng. = *Wedelia pedunculosa*?
 B. *TRIFOLIUM* Poepp. = *Oyedæa buphthalmoides*.
 B. *UNIFLORUM* Forst. = *Wollastonia Forsteriana*.

CCXLVII. *TELEKIA* Baumg. *fl. trans.* 3. (1816) p. 149, *malè*. Less. *syn.* 209.
 — *Molpadia* Cass. *bull. philom.* 1818. *dict.* 32. p. 400. — *Inulæ* sp. *Bieb.*
Pers. — *Asteroidis* sp. *Tourn.*

Radius 1-serialis ligulis longis angustis; corollarum omnium tubus teres infernè sensim angustatus. Anth. basi caudis elongatis. Achænium disci et radii lineare elongatum multicostatum exalatum triquetro-obcompressum. Pappus coroniformis denticulatus conformis subcartilagineus (nec plumosus ut vult Baumg.). — Herbæ proceræ habitu *Inulæ Helenii*.

1. *T. CORDIFOLIA* (Kit. pl. rar. hung. 2. p. 117. t. 113. sub *Buphth.*), caule erecto pubescente, foliis subtus pubescentibus inferioribus petiolatis cordatis duplicato-callososerratis, superioribus ovatis sessilibus simpliciter serratis. ½ in pomariis sylvaticis et pratis montosis Banatùs, Croatiae, Transylvaniæ, Tauriæ, Cappadociæ, Persiæ ad Guriel (Szow!), *Bieb. fl. taur. n.* 1796. *Telekia speciosa* Baumg. *fl. trans.* 3. p. 150. *Molpadia suaveolens* Cass. *dict.* 32. p. 401. *Inula Caucasica* Pers. *ench.* 2. p. 450. *Buphthalmum speciosum* Schreb. *dec. p.* 11. t. 6, *Asteroides orientalis petasitidis folio*, etc. *Tourn. cor.* 51. *inst. t.* 487. Capitulum *Inulæ Helenii*. Invol. squamæ ovatæ mucronatæ. Pappus brevissimus. Herba trita suaveolens. (v. s.)
2. *T. SPECIOSISSIMA*, caule erecto sparsè piloso, foliis cordato-amplexicaulibus ovalibus acutis serratis rigidulis glabris nervo medio subtus subvillosis. ½ in vallibus montium Brixienisium (Ard.), ad rupes circa lacum Comensem (Com!), et in provinciis vicinis ad Corni di Canzo (Duby!), in valle Sasana (Thom!), etc. *Buphthalmum speciosissimum* Ard. *specim.* 1. p. 26. t. 1. Linn. *mant.* 117. nec 517. *Reichenb. icon.* 4. p. 37. t. 349. *Gaud. fl. helv.* 5. p. 382. Invol. squamæ elongatæ acuminatæ. Pappus paulò magis elongatus quàm prioris. (v. s.)

CXXLVIII. *ASTERISCUS* Moench meth. 592. Less. syn. 209. — *Nauplius* Cass. dict. 34. p. 272. — *Asterisci* sp. Tourn.. — *Asteroidis* sp. Vaill. — *Odonostigma* Neck. ex Less.

Radius 1-serialis, ligulis cuneiformibus apice tridentatis, tubo brevi basi biarticulato, cor. disci exalata tubo inferne incrassato. Anth. basi longè caudatæ. Achænium triquetro-obcompressum exalatum conforme. Pappus conformis coroniformis irregulariter dentatus. — Herbæ basi sæpè suffruticosæ ramosæ regiones mediterraneas habitantes plus minùs hirsutæ. Folia obovata oblonga aut linearia integerrima. Capitula terminalia sæpiùs ramis lateraliter ortis et elongatis lateraliter facta. Invol. squamæ ext. foliaceæ inermes obtusæ disco multò longiores.

1. *A. MARITIMUS* (Moench l. c.), caule irregulariter ramoso, ramis apice 1-cephalis nunc foliosis nunc denudatis, foliis oblongo-spathulatis villosis, invol. bracteis spathulatis. 2/ in rupibus maritimis Hispaniæ, Galliæ australis, Italiæ, Aleppi, ins. Mediterran., Mauritanæ. *Bupthalmum maritimum* Linn. sp. 1274. *Nauplius maritimus* Cass. dict. 34. p. 274. *Bupthalmum crassifolium* Raf. speech. 1. p. 191. ex descr. et sp. à cl. Guss. misso. — Barr. ic. t. 251. Bocc. mus. t. 129. Clus. hist. 2. p. 13. Variat caule basi suffruticoso et herbaceo, supino et suberecto, ramis brevibus foliosis (Bupht. sessile Mill.) aut elongatis parè foliosis post anthesin nudis. Invol. squamæ ext. spathulatæ post anthesin deciduæ, int. tunc adpressæ in discum depressum. (v. v. et s.)
2. *A. AQUATICUS* (Moench l. c.), caule erecto di- aut trichotomo, capitulis sessilibus axillaribus et terminalibus, foliis oblongis obtusis villosis aut glaberrimis. (1) in maritimis aquaticis regionis Mediterraneæ à Canariis ad Cretam, à Lusitaniâ ad Byzantium. *Bupthalmum maritimum* Linn. sp. 1274. Gærtn. fr. t. 169. DC. fl. fr. 4. p. 217. Schkuhr handb. t. 237. *Nauplius aquaticus* Cass. dict. 34. p. 273. — Seba thes. 1. t. 29. f. 7. Zan. hist. t. 24. Breyn. cent. t. 77. Interdum caulis induratur. (v. v. et s.)
3. *A. GRAVEOLENS*, caule erecto, ramis sub capitulis ortis elongatis proliferis, foliis oblongis basi cordato-amplexicaulibus incisus dentatisve villosis, capitulis axillaribus brevissimè pedicellatis et terminalibus, invol. bracteis oblongo-linearibus acutis. 2/ ? in Egypti desertis circa Kahiram, necnon circa Suez, Sinai (Bové!). *Bupthalmum graveolens* Forsk. descr. 151. Vahl symb. 1. p. 74. t. 19. Sieb. herb. Egypt. Delil. ill. Capitulum post anthesin subglobosum. (v. s.)
4. *A. SERICEUS*, caule fruticoso ramoso, foliis cuneatis linearibusve integerrimis villosis-sericeis, capitulis terminalibus et lateralibus sessilibus, bracteis linearibus, rec. bracteolis acuminatis. 5 in ins. Canariis. *Bupthalmum sericeum* Linn. f. suppl. 379. Folia alterna nec opposita.
 - a. *latifolius*, foliis spathulato-oblongis densè villosis-sericeis, bracteis radio brevioribus lineari-subspathulatis, invol. squamis linearibus sericeo-hirsutissimis. 3 in insulâ Fuertaventurâ. B. sericeum L'Hér. herb. et mus. Ait. kew. 3. p. 245. Sims. bot. mag. t. 1836. opt. Capit. cum radio diam. sesquipollicari. (v. s.)
 - ß. *intermedius*, foliis spathulato-oblongis adpressius villosis-sericeis bracteis oblongo-subspathulatis radio subæqualibus, invol. squamis linearibus pubescenti-sericeis. 3 in ins. Lancerotte. Bupht. sericeum Link in Buch. ! can. 150. (v. s.)
 - γ. *stenophyllum*, foliis linearibus acutis villosis-sericeis, caule ramosissimo depresso, bracteis folio conformibus, invol. squamis adpressissimis. 3 in campis maritimis ins. Magnæ-Canariæ. Bupht. stenophyllum Link in Buch. ! can. 150. Gravè olet ex Chr. Smith. Capit. cum radio 9-10 lin. diam. (v. s.)
5. *A. ODORUS*, caule suffruticoso erecto trichotomo, foliis lineari-lanceolatis

acutis integerrimis pubescenti-subsericeis, capitulis terminalibus et axillaribus subsessilibus, bracteis plurimis folio conformibus, rec. bracteolis obtusè truncatis apice ciliato-villosis. \S in arenosis imperii Maroccani (Schousb.), prope Mogador (Brouss.), Bupthalmum odorum Schousb. mar. 2. p. 199. t. 7. Prioris var. γ affinis sed bracteolis obtusis faciliè differt. (v. s.)

6. *A. IMBRICATUS*, glaber atrovirens, caule fruticoso ramoso, foliis in petiolum attenuatis spatulatis obtusis argutè et grossè serratis, capitulis terminalibus solitariis, bracteis cuneiformibus apice 3-dentatis. \S in agro Mogadoriensi detexit cl. Broussonet. Bupthalmum imbricatum Cav. annal. scienc. nat. 1801. v. 4. p. 94. Bupth. elegans Willd. mag. naturl. ges. Berl. 1807. p. 138. Buph. chrysanthemoides Poir. suppl. v. 1. (1810) p. 747 excl. patr. Nomina 3 certè synonyma cum tres descr. è specim. Broussonetianis concinnatè! (v. s. comm. à cl. Brouss.)

CCXLIX. *PALLENIS* Cass. dict. 37. p. 275. Less. syn. 210. — *Athalmum* Neck. elem. 1. p. 20? — *Asterisci* sp. Tourn. — *Asteroidis* sp. Vaill.

Radius biserialis. Cor. radii angustè ligulata apice tridentata tubo crasso utrinquè alato, disci tubo crasso hinc marginato-alato. Achæmium obcompressum orbiculatum alatum; disci subtriquetrum villosiusculum. Pappus denticulato-ciliaris, radii dimidiatus externus, disci coroniformis. — Inv. squamæ exter. foliæ patentès apice spinosæ.

1. *P. SPINOSA* (Cass. l. c.), caule erecto hirsuto, foliis obovato- aut. oblongo-lanceolatis, invol. squamis ext. patulis spinescentibus. ① in asperis et subhumidis apricis, ad margines agrorum Europæ australis et Africæ borealis, à Teneriffa ad Ægyptum, ab Armorica Nannetensi ad Constantinopolim. Bupthalmum spinosum Linn. sp. 1274. DC. fl. fr. 4. p. 217. Schkuhr handb. t. 258. Bupth. asteroidum Viv. fl. lyb. sp. p. 57. t. 25. f. 2. — Blackw. herb. t. 272. Clos. hist. 2. p. 13. f. 1. Variat caule plus minus hirsuto simplici 1-cephalo aut corymboso, capitulis centralibus sessilibus cæteris longè pedunculatis, foliis integerrimis aut denticulatis. *B. aureum* Salzm. pl. exs. circa Tanger lectum videtur var. foliis latioribus caule et involucri basi hirsutissimis. (v. v. sp.)

CCL. ANVILLEA.

Capitulum multiflorum homogamum, fl. omnibus tubulosis 5-dentatis hermaphroditis. Recept. convexi paleæ basi oblongæ convexæ, apice ext. in acumen, interiores in setam productæ. Involucri campanulati squamæ ext. seu bractæ foliæ spatulatæ patulæ, inter. biserialis adpressæ imbricatæ. Anthæræ basi ecaudatæ (Vent.). Achæmium tetragonum durum pappo coroniformi brevi integro coronatum. Herba, erecta aut multicaulis diffusa præsertim ad summitates velutino-canescens, caule ad basin aut rhizomate fruticoso. Folia obovata spatulatave basi attenuata grossè et obtusè dentata. Capitula sessilia terminalia, ramis sæpè sub capitulis ortis quasi lateralia. Flores aurei. — Genus à cæteris Bupthalmeis distinctum floribus omnibus tubulosis hermaphr. et ideo ut ex antheris ecaudatis ad Ceruanam vergens. — Dicavi cl. d'Anville (Joh. Bapt. Bourguignon d'Anville), summo geographo Parisiensi qui Orientalium regionum situs labore indefesso sagacissimè exploravit.

1. *A. GARCINI*. \S in Oriente non longè à Bagdad leger. cl. Olivier et Bruguière! in Arabia-Petræa (ex syn. Delil.) et in Persiâ olim legit Garcin (ex Burm.). Bupthalmum Persicum Garc. ex Burm. Bupthalmum Garcini Burm. fl. ind. t. 60. f. 1. Anthemis Garcini Burm. fl. ind. p. 183. Bupthalmum Arabicum

Delil. fragm. fl. arab. petr. p. 14. t. 4. ex Guillemain qui specim. vidit. Buphthalmum floeculosum Vent. h. Cels n. et t. 25. (v. s. ex herb. Oliv. et Brug.)

CCLI. CERUANA Forsk. *æg. descr.* 153. *Cass. dict.* 8. p. 22. *Less. syn.* 202. — Buphthalmi sp. Vahl.

Capitulum multiflorum heterogamum discoideum; flor. radii pluri-seriales fœminei graciles 3-dentati, disci hermaphr. 5-dentati. Recept. convexum totum bracteolatum. Invol. campanulato-hemisphæricum bracteis foliaceis extus cinctum erectum subbiseriale. Antheræ ecaudatæ. Achæmium oblongo-obovatum bicostatum compressum, callo apicali rostrum brevissimum et annuliforme sistente. Pappus brevissimus coroniformis piloso-dentatus. — Herbæ rarius fruticuli hirsuti Africani. Folia alterna subsemiamplexicaulia grossè dentata aut incispinnatifida. Capitula cymosa, nempe pedunculi 1-3 infrà capitulum terminale quodque nascentes et elongati in pseudo-laterales conversi. Flores lutei. — Genus habitu Buphthalmo affine sed discoideum, Grangæ sed recept. bracteolatum.

1. C. PRATENSIS (Forsk. fl. æg. p. LXXIV), caule herbaceo erecto hispido, foliis obovato-cuneatis dentatis aut pinnatilobatis, bracteis capitulo longioribus lanosolato-linearibus. ① in Ægypto ad ripas Nili. Buphthalmum pratense Vahl symb. 1. p. 75. Delil! fl. eg. p. 132. t. 48. f. 2. Sieb. herb. eg. exs. (v. s.)
2. C. SENEGALENSIS, caulibus herbaceis erectis hispidissimis, foliis obovato-cuneatis grossè dentatis, bracteis ovatis obtusis capitulo longitudine. ①? in Senegaliâ ad Podor legit indef. Perrottet mense 7^{br} flor. Caulis à basi multiceps. Radix simplex crassitie pennæ columbinæ. Herba tota hispidissima. An fortè ex patriâ C. rotundifolia Cass. dict. 41. p. 123? sed forma foliorum non ritè cum nostrâ congruit. (v. s. comm. à cl. Perrottet.)
3. C. FRUTICOSA (Less. syn. 202), caule procumbente lignoso ramosissimo, foliis oblongo-ellipticis subsemiamplexicaulibus integerrimis seu pinnatifido-dentatis, ramulis confertis 1-cephalis. 3 in Ægypto prope Alexandriam. Facies Grangæ Maderaspatanæ.

Subtrib. VI. ECLIPTEÆ Less. syn. 210. — Helianthearum gen. Cass.

Capitula multiflora heterogama, fl. radii fœmineis (excl. Chryphiospermo) ligulatis, disci hermaphroditis. Antheræ ecaudatæ. Recept. paleaceum. Pappus nullus seu aristatus nunquam setosus. Folia opposita. — Subtribus Heliantheis proxima sed stigmati structurâ ad Asteroideas potiùs referenda.

CCLII. BORRICHIA Adans. fam. 2. (1763) p. 130. — Diomedea Cass. bull. philom. 1815. p. 175 et 1817. p. 70. dict. 13. p. 283. Kunth nov. gen. 4. p. 212. Less. syn. 240. non Zoolog. — Odontospermum Neck. elem. 1. (1791) p. 20? — Buphthalmi sp. Linn.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis 1-serialibus, disci hermaphrod. tubulosi 5-fidis. Involucrum hemisphæricum, squamis imbricatis, ext. foliaceis acutis, int. obtusissimis margine membranaceis. Recept. planum paleis lanceolatis onustum. Cor. faux parçè dilatata. Styli rami longi per totam ferè longitudinem hispidi acuti. Antheræ nigricantes. Achænia cuneata compresso-angulata corticata apice pappo brevi coroniformi dentato coronata, disci subdifformia subcalva. — Frutices suffruticesve Americani maritimi. Folia opposita oblonga integerrima coriacea medianibus utrinquè li-

gulâ triangulari adpressâ (stipulâ auct.) subconnata. Capitula terminalia pedicellata. Flores lutei. — *Borrichia* est nomen vetustius, caractere pro tempore rectè stabilitum, dictum cl. Borrich. *Diomedea* est nomen multò recentius et jam veteribus et nunc zoologis avem designat.

1. *B. ARBORESCENS*, glabra, foliis lanceolatis mucronato-acutis basi angustatis integerrimis, invol. squamis ovatis ext. subacutis adpressis, paleis spathulatis obtusis. ζ 5 in insulis Cayo, Flamingo et Bonito, ad littus australe insulæ Cubæ (H. B.), in insulis Bahamensibus (Linn.), et Bermudianis (Pluk.). *Bupthalmum arborescens* Linn. sp. 1273. *Diomedea indentata* Cass. dict. 13. p. 284. *Diomedea glabrata* H. B. et Kunth nov. gen. am. p. 213. — Dill. elth. 1. t. 38. f. 43. Plum. amer. ed. Burm. t. 16. f. 2. (v. s.)

2. *B. FRUTESCENS*, adpressè sericeo-canescens, foliis lanceolatis obovatisve obtusis basi attenuatis utrinquè grossè et acutè ad basin 1-2-dentatis, rameis sæpius edentulis, invol. squamis ext. apice recurvo-aquarosis, paleis acuminatis indurato-spinosis. ζ 5 in Jamaica (Sloane), in maritimis Americæ borealis à Virginia ad Floridam (Pursh!), in Mexico (Berl. ! Pl. exs. n. 1963). *Bupthalmum frutescens* Linn. sp. 1273. *Bupthalmum incanum* Mill. *Diomedea bidentata* Cass. dict. 13. p. 283. — Dill. elth. 1. t. 38. f. 44. Catesb. car. 1. t. 93. Plum. et Burm. t. 107. f. 1. (v. s.)

β . *angustifolia*, foliis lineari-lanceolatis medio aut basi obtusè et acutè dentatis. ζ in Mexico ad Tampico (Berl. ! pl. exs. n. 126). An *Bupthalmum lineare* Willd. sp. 3. p. 2260 ad hanc aut ad sequentem referendum? (v. s.)

3. *B. PERUVIANA*, foliis suprà glabriusculis subtùs sericeo-canescens linearilanceolatis submucronatis medio denticulis subobtusis 1-2 sæpius notatis. ζ in Peruvia (Domb. ex h. L'Hér. !). *Bupthalmum Peruvianum* Lam. dict. 1. p. 515. (v. s. sine capit.)

4. *B. ARGENTEA*, sericeo-lanata argentea, foliis lanceolatis mucronatis integerrimis, involucri squamis extimis ovatis sericeis vix mucronatis adpressis, int. glabris ciliatis amplis, paleis acutis flosculis ferè brevioribus. ζ in Peruvia (Domb. ex h. L'Hér. !), in maritimis insulæ Cubæ prope Havanam (La Sagra) et Guanavacoa (H. et B.), in ins. Magni-Kaimanis prope Jamaicam (Don ex Lindl.). *Diomedea argentea* H. B. et Kunth nov. gen. am. 4. p. 213. Lindl. trans. hort. soc. Lond. 7. p. 50. report 1827. p. 7. *Bupthalmum canum* L'Hér. ! herb. et descr. mas. Cum priore in hortis interdum confusa sed involucrio diversissima. (v. s.)

5. *B. DEMISSA*, caule demisso ramoso glabro, foliis strigilloso-pubescentibus confertis subspathulatis basi longè attenuatis obtusis integerrimis, pedunculis terminalibus, invol. squamis ovatis adpressis post anthesin conniventibus, paleis acutis flosculo brevioribus confertis et alveolas ferè constituentibus. ζ circa Carthagenam Amer. austr. legit cl. Gaudichaud. (v. s.)

CCLIII ECLIPTA Linn. mant. (1767) p. 157. Gærtn. fr. 2. p. 441. t. 169. Cass. dict. 14. p. 231. 46. p. 399. et 410. Less. syn. 212. — *Micrelum* Forsk. descr. p. 152. — *Eupatoriophalacron* Vaill. act. Par. 1720. excl. sp. Adans. fam. 2. p. 130.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus foemineis angustissimè et breviter ligulatis, disci tubulosi 4-dentati hermaphr. Recept. è plano convexum, bracteolis lineari-filiformibus apice ciliatis achænia æquantibus instructum. Inv. biseriale, squamis 10-12 ovato-lanceolatis acuminatis. Achænia radii triquetra, disci compressa ad latera muricato-tuberculosa, apice epapposa mutica aut denticulis 1-3 minimis apiculata, juniora ad apicem puberula. — Herbæ æquinoc-tiales plus minus scabræ aut hirsutæ ramosæ erectæ aut prostratæ. Folia opposita integra aut serrata penninerviâ. Pedunculi axillares soli-

tarii gemini aut terni 1-cephali. Flores albi. Succus ad colorem nigrum inficiendum adhibendus. — Genus ex Cass. affine hinc Bupthalamo illinc Melantheræ.

Sect. I. EUECLIPTA. — Ecliptæ sp. auct.

Achaenia omnia muricato-tuberculosa calva aut vix ac ne vix aristulata, radii triquetra, disci modicè compressa oblonga.

1. *E. RARECTA* (Linn. mant. 286? Willd. sp. 3. p. 2217), caule erecto adpressè strigoso, foliis oblongo-lanceolatis utrinquè acuminatis, pedicellis solitariis geminive capitulo quintuplò longioribus. ① per orbis totius regiones calidas in graminosis humidis argillosis inundatis ferè vulgata; habeo ex Chinâ, Cochinchinâ, Ind. or., Ægypto, Senegaliâ, agro Ovariensi, Africâ austr.-orient. prope Port-Natal (Drege!) et ex insulis Cubâ, Trinitatis, Mauritiî, etc.; indic. etiâ in Moluccis, Javâ, Asiâ minore, Surinamo, Virginiâ, Floridâ, Novâ-Granatâ, Brasiliâ, Sancto-Domingo, sinu Mullgrave Amer. bor.-occ., etc. *Verbesina alba* Linn. sp. 1272. *Cotula alba* Linn. syst. nat. 2. p. 564. *Microlium asteroides* Forak. descr. 152. *Ecl. adpressa* Moench meth. — Dill. elth. f. 137. Rumph. amb. 6. p. 18. f. 1. Pluk. alm. t. 109. f. 1. Folia utrinquè adpressè strigosa, integra parè aut intensè serrata. Fortè species Linneana est sequens sed cum omnes auctores hanc sub nomine *E. erectæ* habeant, etiâ admisit. (v. v. et s.)

β. *diffusa*, caule à basi ramosissimo diffuso aut adscendente pedunculis vix capitulo duplò longioribus. ① in Senegaliæ regione Walo cum var. *erectâ* in argillosis inundatis legit cl. Perrotet. (v. s. comm. à cl. inv.)

2. *E. LONGIFOLIA* (Schrad! in litt. 1832), caule erecto adpressè strigoso, foliis oblongo-lanceolatis acuminatis basi obtusis semialexicanilibus ad latera reflexis pedunculis 1-3 capitulo quintuplò longioribus. ① in insulis Caribæis, Guadalupâ; fortè in Porto-Ricco, Domingo et Martinicâ si *Bellis ramosa* Jacq. amer. 216. t. 129 et idè *E. punctata* Linn. mant. 2. p. 286 huc ritè referendæ? *E. erecta* Linn. mant. 286? (v. s.)
3. *E. ZIPPELIANA* (Blum. I bijdr. 914), caule erecto apice patentim hirsuto, foliis ovali-lanceolatis undulatis utrinquè attenuatis, pedunculis 1-4 capitulo vix duplò longioribus. ① in graminosis paludosis Bataviæ. Pedunculi hirsuti. Folia utrinquè adpressè strigosa serrata. (v. s. comm. à cl. Blume.)
4. *E. PROSTRATA* (Linn. mant. 286), caule prostrato aut adscendente adpressè strigoso ramulis subhirsutis, foliis ovali- aut oblongo-lanceolatis basi attenuatis subserratis subundulatis scabris, pedunculis 1-2 capitulo duplò triplòve longioribus. ① in humidis Indiæ orientalis ut vid. frequens et in insulâ Maderâ ad rivulos indigena facta ex rev. Lowe *Verbesina prostrata* Linn. sp. 2. p. 1272. *Cotula prostrata* Linn. syst. 2. p. 564. *Microlium Tolak* Forak. descr. 153. — Dill. elth. f. 138. Pluk. alm. t. 118. f. 5. (v. s.)
- β. *undulata*, ramis apice strigosis patentibus hirsutis, foliis ovatis, pedunculis vix capitulo longioribus. ① in Indiâ orientali. *Ecl. undulata* Willd. sp. 3. p. 2219. Hanc et veram prostratam conjungo ex pluribus specimenibus à cl. Wallichio sub n. 319 et 320, et à cl. Royle comm. nec inter se meo sensu separatu dignis. (v. s.)
5. *E. PARVIFLORA* (Wall. cat. et herb. n. 323), caule prostrato-repente adpressè etiâ in ramulis strigoso, foliis oblongo-lanceolatis utrinquè acuminatis subintegerrimis subscabridis, pedunculis 1-2 capitulo vix duplò longioribus. ① in Indiâ orientalis humidis legerunt cl. Heyne et Wight. Hic etiâ pertinere videtur *E. prostrata* D et *E. Wall.* cat. et herb. comp. n. 319 et *Ecl. prostrata* à cl. Chamisso in Luzoniâ lecta. Folia ferè *E. erectæ*, habitus *E. prostratæ*, caulibus longè radicantibus ab utrâque distincta. (v. s.)
6. *E. TERNALIS* (Bunge! enum. pl. chin. bor. n. 224), caule erecto simplici aut divaricato-ramoso adpressè strigoso apice subhiépido, foliis oblongis utrinquè

- attenuatis subintegerrimis adpressè strigosis, pedunculis 1-2 capitulo vix longioribus, achæniis angulis crassioribus granulatis. ① ad thermas prope Tanschan Chinæ borealis legit cl. Bunge fl. maio. (v. s. comm. à cl. inv.)
7. *E. PROCUMBENS* (Michx. fl. bor. am. 2. p. 129), caule procumbente adpressè strigoso, foliis longo-lanceolatis utrinquè attenuatis rariter serratis utrinquè adpressè scabris, pedunculis 1-2 capitulo duplò circ. longioribus, inv. squamis oblongo-lanceolatis acuminatis. ① in humidis Carolinæ (Bosc!). Ell. sketch 2. p. 403. Grangea lanceolata Poir. suppl. 2. p. 826. Rami ab Ell. dicuntur radicantes. Invol. squamæ radio longiores. (v. s. comm. à cl. Bosc.)
- E. patula*, foliis latioribus ovali-lanceolatis subtriplinerviis. ① Patr. ign. Ecl. patula Schrad. in litt. 1832. Bupthalmum diffusum Vahl herb. ex Puer. herb. (v. s.)
8. *E. BRACHYPODA* (Michx. fl. bor. am. 2. p. 130?), caule divaricato-prostrato adpressè strigoso, ramis apice pedunculisque hirsutis, foliis longo-lanceolatis utrinquè acuminatis rariter serratis utrinquè adpressè scabris, pedunculis 1-2 capitulo paulò longioribus, inv. squamis ovatis cuspidato-acuminatis. ① in Louisiana prope Novam-Aureliam et in Carolinâ (Michx.) fortè in Mexico littorali ad Tampico (Berl. n. 23?). Amellus Carolinianus Walt. fl. car. p. 213 ex Mich. Flores dixit 5-fidos cl. Michaux et in nostrâ 4-fidi! sed ipse Elliot nunquam 5-fidos vidit. An species dum hic confusæ? an caract. varians? (v. s.)

Sect. II. PLATECLIPTA.

Achaenia lævia ant partim muricata aristis 2-3 coronata, lateraliter in alam latam complanata, radii subtrigona, disci compressissima subrotunda. — Herbæ glabriusculæ, foliis basi triplinerviis.

9. *E. LANCEOLATA*, erecta glabra, foliis oblongo-lanceolatis integerrimis, pedicellis terminalibus et è dichotomiis ortis folio longioribus apice adpressè pubescentibus, invol. squamis 10 lanceolatis inter se subæqualibus, achæniis medio dorso muriculatis. ②? in Brasiliæ prov. Rio-Grande. Caulis purpurascens pedalis sesquipedalisve. Folia 12-15 lin. longa, 4-5 lata internodiis breviora. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1007.)
10. *E. ELLIPTICA*, ascendens glabra, foliis elliptico-oblongis basi attenuatis integerrimis, pedicellis terminalibus folio paulò longioribus adpressè pubescentibus, invol. squamis 5 ext. majoribus oblongis, 5 inter. submembranaceis dimidiò minoribus, achæniis non aut vix tuberculosi. ② in Brasiliæ prov. Rio-Grande. Folia vix pollicem longa 4 lin. lata internodiis sæpè longiora. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. n. 1024.)

+ Species vix notæ.

11. *E. PALUSTRIS* (Forst. prodr.), erectiuscula glabra foliis lanceolatis subintegerrimis, capitulis pedunculatis. — in insulis maris Pacifici. An Meyeræ seu Enhydræ sp.? †.
12. *E. PUSILLA*, basi repens, caule simpliciter foliis petiolatis ovalibus acutis glabris subenticulatis 3-nerviis, capitulis axillaribus pedicellatis solitariis. — in Porto-Ricco. Cor. flavidæ seu purpurascens. Verbesina pusilla Poir. dict. 8. p. 459. †.
13. *E. GILIATA* (Raf. lud. fl. p. 69), ex foliis alternis à genere videtur aliena nec satis nota.

++ Species exclusæ.

- E. HUMILIS* Kunth. = Sabazia humilis.
E. INTEGRIFOLIA Spreng. = Bellis integrifolia.
E. LATIFOLIA Linn. f. = Blainvillea latifolia.
E. SCABRIUSCULA Wall. = Wollastonia scabriuscula.
E. SESILIS Sw. = Enhydra sessilis.

CCLIV. BLAINVILLEA *Cass. journ. phys.* 1823. p. 216. *dict.* 29. p. 493. 47. p. 90. 59. p. 138. *Less. syn.* 212. — *Ucacea Cass. dict.* 27. p. 9 *excl. syn. ad.* — *Verbesinæ et Ecliptæ sp. auct.*

Capitulum pluriflorum heterogamum, fl. radii paucis 1-seriatis fœmineis in ligulam latam brevem 3-dentatam irregularem expansis, disci tubulosis 5-dentatis hermaphr. Invol. ovato-cylindraceum 1-2-seriale, squamis erectis exter. ovali-oblongis foliaceis, int. paleiformibus. Recept. planum angustum, paleis flores amplectentibus concavis submembranaceis apice dentatis. Stylus disci inclusus, ramis brevibus ferè Bupthalmi. Achænia radii triquetra glabriuscula in angulis hispidula aristas 3 persistentes rigidas ciliatas apice gerentibus, disci lateral. compressa biaristata rarius 3-4-arist. — *Herbæ Æquinoctiales Gerontogæ erectæ scabræ aut hirsutæ. Folia opposita petiolata triplinervia ovato-rhombea aut lanceolata dentata. Pedunculi ex axillis seu aliis ramorum 1-3, monocephali nudi. Flores albi.*

1. B. RHOMBOIDEA (*Cass.!* *dict.* 29. p. 493), foliis rhomboideis acuminatis irregulariter dentatis utrinquè tomentoso-hirsutis, ramis inf. oppositis summis dichotomis, pedicellis petiolo multò longioribus, achæniis aristis crassis triquetris à basi ad apicem barbato-plumosis. ① in *Brasiliâ* (*Comm.!*) circa *Rio-Janeiro* (*Lund.!* *Gaud.!*) et *Bahia* (*Blanch.!* *Salzm.!*). *Verbesina dichotoma* *Murr. comm. Goett.* 1779. p. 15. t. 4 opt. sed *excl. patr.* Stirps in hortis botanicis non infrequens, sequenti similis sed certè diversa. (v. s. ap. et v. c.)

β. *lanceolata*, foliis angustioribus oblongo-lanceolatis. — *Verbesina lanceolata* *Poir.!* *dict.* 8. p. 461. (v. s. c.)

2. B. LATIFOLIA (*DC.* in *Wight contrib.* p. 17), foliis rhomboideis acuminatis grossè dentatis utrinquè sparse pubescenti-hirtellis, ramis petiolisque hirtellis oppositis summis dichotomis, pedunculis petiolo plerumquè brevioribus, achæniis aristis basi barbatis apice nudis sæpè subabortivis. ① in *Indiâ orient.* ad montes *Dendigul Malabaricæ* legit *cl. Wight* et ad *Munghir. Eclipta latifolia* *Linn. f. suppl.* 378 et *Linn.!* *herb. ex Wight!* *herb. Verbesina dichotoma* *C et E Wall.!* *cat. et herb.* n. 314. *Verbesina Lavenia* *Roxb. fl. ind.* 3. p. 442 non *Linn. ex Wight.* Ad priorem valdè accedit sed pubero-hispidula nec velutino-hirsuta, folia latiora duplò longius petiolata papyraceo-membranacea, superiora ferè lanceolata. (v. s.)

β. *angustifolia*, foliis angustioribus oblongo aut vix ovato-lanceolatis. — in *Indiâ orient.*, *Ceylonâ*, etc. *Verbesina dichotoma* *Wall.!* *cat. et herb.* n. 314 *A. C. D.* (v. s.)

3. B. GAYANA (*Cass.!* *dict.* 47. p. 90), foliis oppositis et alternis longè petiolatis ovato-lanceolatis dentatis suprà scabris subtus pubescentibus, pedunculis è dichotomiis ramorum ortis aut oppositifoliis petiolo duplò longioribus, invol. squamis int. striatis, vix exter. subvillosis. ① in *Senegaliâ* *ex semin.* ad *cl. Gay* missis. Ligulæ dicuntur biligulatæ, in specim. meo deciduæ. (v. s. comm. à *cl. Gay.*)
4. B. PRIEURANA, foliis oppositis breviter petiolatis oblongo-lanceolatis acuminatis integerrimis aut subserratis utrinquè strigoso-hispidis subtus pallidioribus, pedunculis axillaribus et alaribus petiolo triplò et ultrà longioribus, invol. squamis linearibus dorso strigosis intimis levibus non striatis. ① in *Senegaliâ* superiore ad montes *Bakel* legit *cl. Leprieur.* Flores non vidi. Achænia omnia in specim. meo oblonga subcuneata utrinquè convexa biangulata apice biaristata juniora præsertim villosa submaculata. An fortè *Wedelia Africana* *Pers. ench.?* p. 490 huc refer.? (v. s. comm. à *cl. inv.*)
5. B. BIARISTATA, foliis oppositis longè petiolatis ovatis acutis serratis utrinquè sparse et adpressè pilosis subtus pallidioribus, pedunculis alaribus petiolo paulò longioribus, inv. squamis 5 oblongis acutis, achæniis omnibus biaristatis. ① in

Brasilæ prov. Rio-Grande. Ligulæ parvæ fœminææ. Capitula parva pauciflora. Petioli 3-9 lin. longi. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 873 miss.)

CCLV. SALMEA DC. cat. h. Monsp. (1813) p. 140. Brown. trans. linn. soc. 13. p. 112. Cass. dict. 47. p. 87. Less. syn. 212 non Cav. nec Willd. (1). — Hopkirkia Spreng. nov. prov. (1818) p. 23. — Bidentis sp. Linn.

Capitulum homogamum. Inv. imbricatum. Recept. conicum paleaceum, paleis convolutis flores amplectentibus. Cor. tubulosæ, fauce vix dilatata, lobis 5 subrevolutis. Antheræ sagittatæ (Br.). Styli basi subbulbosi rami breves crassi extus subhispidi suprâ papuloso-puberi obtusi. Achænia obcompressa linearia aptera, marginibus ciliatis apice in aristas 2 persistentes productis. — Frutices erecti aut subscandentes Americani et præs. Caribæi. Rami teretes. Folia opposita petiolata ovata acuta penninervia dentata aut subintegerrima. Capitula paniculato-corymbosa. Flores albidi. — Genus affine Dahliz ex stigmate, sed valdè affinè Spilantho ex receptaculo achænio et paleis. Valdè dubius hæreo de loco huic generi in ordine naturali tribuendo, aut inter Eclipteas aut inter Verbesineas.

1. S. EUPATORIA (DC. cat. h. monsp. p. 140 cum ic. ined.), caulibus plurimis suberectis glabris apice pubescentibus, foliis ovatis acuminatis hinc indè dentatis glabris, aristis æqualibus, corollis rectis. ♀ in ins. Caribæis. Bidens scandens Swartz obs. 927? Salmea grandiceps Cass. dict. 47. p. 88? Hopkirkia eupatoria Spreng. syst. 3. p. 443. Jacq. f. ecl. v. 2. t. 139. ined. Eupatorium Houstonis et Eup. Gonani hortul. non Linn. Rami divaricati. Pedunculi puberulo-hirsuti divaricati, inf. sæpius alterni paniculati, sup. oppositi corymbosi. (v. s. sp. et v. c.)

β. *intermedia*, ramis ramulisque glaberrimis, foliis basi cuneatis repandodentatis glabris. ♀ in Jamaicâ! An fortè species propria exactè inter S. eupatoriam et scandentem media. (v. s.)

2. S. SCANDENS (DC. l. c.), caule scandenti ramisque etiâ ad apicem lævibus, foliis ovatis acuminatis subintegerrimis glabris, aristis æqualibus, corollis rectis. ♀ in ins. Caribæis et sp. in Jamaicâ. Bidens scandens Linn. sp. 1167. hort. cliff. 399. Sims bot. mag. t. 2062. Salmea scandens Brown l. c. Bidens frutescens Mill. dict. n. 7. Hopkirkia scandens Spreng. syst. 3. p. 443. Calea Amellus Linn. sp. 1179? ex Spreng. et suad. Willd. sp. 3. p. 1795. Salmia parviceps Cass. dict. 47. p. 88? Isocarpha Amellus Spreng. in h. Balb. (v. s. sp.)

3. S. HIRSUTA (DC. l. c.), caule scandente striato, ramis hirsutis, foliis ovatis basi subcordatis apice acuminatis integerrimis suprâ pubescentibus subtus villosis-hirsutis, aristis inæqualibus, corollis rectis. ♀ in Jamaicâ. Bidens hirsuta Swartz fl. ind. oec. 3. p. 1295. Salmea hirsuta Brown l. c. Hopkirkia hirsuta et Eupatorium vestitum Spreng. syst. 3. p. 444 et 416. Salmea oppositiceps Cass. dict. 47. p. 89? (v. s. comm. à cl. Bertero.)

4. S. ? CURVIFLORA (Brown trans. soc. lin. 13. p. 112), tubo corollæ insigniter extrorsum curvo, aristis inæqualibus, interiori alata, stigmate revoluta. ♂ Patr. et cæt. car. ign.

N. B. Hopkirkia fruticulosa Spreng. syst. 3. p. 444, capitulo radiato et pluribus aliis caracteribus à genere diversa, est Armania fruticulosa Bert.

(1) Salmea Cav. est Sansevieria. Salmia Willd. est Carludovica.

triplo longioribus. ① in planitie Bogotensi ad urbem Santa-Fé (H. et B.), forte in Mexico ad Zacatlipam (Berl.). (v. s.)

4. *S. serrata*, foliis ovatis acutis regulariter dentato-serratis breviter petiolatis, summis sessilibus ovato-lanceolatis ciliatis, invol. ext. squamis interiore duplo longioribus. ① in Chili legit cl. Chamisso et in America austr. cl. Poeppig. *S. cordifolia* Poepp. diar. 463. herb. n. 55. Differre videtur à priore petiolo semipollic. nec 3-pollic. limbo ovato nec cordato, inv. ext. squamis brevioribus. (v. s.)
5. *S. isERICA* (Willd. sp. 3. p. 220), foliis deltoideo-triangularibus grosse et irregulariter dentatis in petiolo cuneato-decurrentibus, summis ovato-lanceolatis, invol. ext. squamis interiores paulo superantibus. ① in Caucasi Iberici sylvis (Stev.), Tauriæ merid. vineis (Bieb. Mey. l. enum. n. 595) et circa pagos Mediæ (Buxb.). *S. Orientalis* Bieb. fl. taur. n. 179 r. — Buxb. cent. 3. p. 29. t. 52. (v. s.)
6. *S. MICROCEPHALA*, foliis breviter petiolatis oblongo-lanceolatis hinc inde serratis utrinque sparse puberulis, involucri ext. squamis lineari-subspathulatis glanduloso-ciliatis vix capitulo longioribus. — in Novâ-Hollandiâ ex herb. Lambert. Caulis dichotomos gracilis sparse puberulus. Petioli 2 lin. Folia fere sesquipollic., 5-6 lin. lata. Capit. 3-4 lin. diam. Ligulæ paucæ obovatæ 3-dentatæ. (v. s. comm. à cl. Lambert.)
7. *S. GRACILIS*, foliis breviter petiolatis oblongo-lanceolatis subintegerrimis utrinque sparse puberulis, invol. ext. squamis linearibus apice subspathulatis glanduloso-ciliatis capitulo duplo longioribus. — in Novâ-Hollandiâ ex herb. Lambert. Similis *S. microcephalæ* et forte mera varietas. (v. s. comm. à cl. Lambert.)

Seet. II. *TRIMERANTHES* Cass. dict. 49. p. 114. 59. p. 237. —
Schkuhria Moench meth. 566, non Roth.

Fl. radii difformes breviter et irregulariter ligulati 2-3-dentati, disci 3-dentati 3-andri. Achænia collo omnino destituta.

8. *S. FLOSCULOSA* (L'Hér. l. stirp. nov. p. 37. t. 19), foliis sessilibus ovatis dentatis, invol. ext. squamis interiore duplo longiore. ① in Peruvia. *Schkuhria dichotoma* Moench meth. 566. *Trimeranthes dichotoma* Cass. dict. 49. p. 115. (v. v. et s.)

CCLIX. *SABAZIA* Cass. dict. 46. p. 480. v. 55. p. 264 et 273. Less. syn. 211. —
Ecliptæ sp. Kunth.

Capitulum multiflorum heterogamum, fl. radii foemineis 1-seriatis ligulatis, ligulis cuneatis grosse tridentatis 6-nerviis invol. longioribus, disci hermaphr. tubulosi 5-dentatis. Recept. conicum, paleis linearibus seu setiformibus glabris achænia fere superantibus, intimis sæpe post anthesin ad apicem recept. persistentibus. Invol. semiglobosum biseriale 7-8-phyllum. Cor. extus pilosiuscula. Achænia conformia pentagona striata cuneata apice omnino calva nec apiculata. — Herbae Mexicanæ ramosæ. Folia opposita petiolata oblonga dentata glabra. Pedunculi alares aut plerique terminales elongati 1-cephali. Ligulæ albidæ extus purpurascentes. Genus ab *Ecliptâ* differt ligulis latè cuneatis, tubulis 5-dentatis, fructibus striatis nec tuberculatis 5-gonis omnibus nec 2-3-gonis dissimilibus apice omnino calvis et muticis. Habitat Galinsogæ sed achænium calvum.

§. 1. Recept. conico, achæniis glabris.

1. *S. HUMILIS* (Cass. dict. 46. p. 481), erectiuscula ramis pedunculisque pilosohirsutis, ligulis profunde trifidis. ① in Novâ-Hispaniâ prope Ario et Pazcuaro. *Eclipta humilis* H. B. et Kunth nov. gen. am. 4. p. 264. t. 394.
2. *S. SARMENTOSA* (Less. in linnæa 1830 p. 148), caule sarmentoso, foliis ovatis

acuminatis teneris triplinerviis pilosiusculis anticè remotè serratis, invol. squamis obtusis, inter. ciliatis majoribus. — in Mexico ad Serro-Colorado.

3. *S. MICROGLOSSA*, caule erecto pedunculisque glabris, foliis ovato-lanceolatis acuminatis teneris ferè à basi ad apicem remotè serratis, invol. squamis obtusis, inter. vix ciliatis, ligulis minimis. ① in montanis circa Mexico ad S.-Augustinum legit Berlandier (pl. exs. n. 733). (v. s.)

ß. puberula, caule ad apicem pedicellisque pubescentibus. ① circa Mexico in montanis. (Berl.! pl. exs. n. 910). (v. s.)

§. 2. *Recept. convexo, achæniis hispidulis.*

4. *S.?* *URTICÆFOLIA*, hispida, foliis ovatis acuminatis grossè dentatis, invol. squamis exter. piloso-glandulosis, recept. convexo, achæniis hispidulis. ② in agro Quitensi inter Mulalo et Pensache. *Viborgia urticæfolia* H. B. et Kunth nov. gen. am. 4. p. 257. t. 389. An *Calydermis* spec. Less. syn. 245? *Habitus* *Galinsogæ* sed achænium calvum. Differt à *Calyderme* capitulis radiatis.

† *Species exclusæ.*

S. OCCIDENTALIS Lin. = *Verbesina phaetusa*.

S. OCCIDENTALIS Walt. = *Verbesina Siegesbeckia*.

S. PORTORICCENSIS Bert. = *Ogiera Portoriccensis*.

CCLX. *CRYPHIOSPERMUM* Beauv. fl. ovar. 2. p. 24. t. 74. (excl. descr. ferè tota et icon. stigm.). Cass. dict. 12. p. 78. Less. syn. 210. — *Wahlenbergia* Schum. pl. guin. 387. non Schrad. — *Enhydra* sp. Cass. — *Meyera* sp. Brown. — *Cæsulia* sp. Willd.

Capitulum multiflorum heterogamum, fl. omnibus intra rec. paleas complicatas omnino involutis, ext. pluriserialibus fœmineis, centr. hermaphroditis stigmatibus alteratione subabortivis. Invol. 4-phyllum, squamis foliaceis, 2 oppositis majoribus. Recept. hemisphæricum paleaceum, paleis coriaceis subglandulosis apice pilos articulos paucos gerentibus. Cor. basi latiores corneæ apice attenuatæ, limbo brevissimo, radii subligulato 3-4 dentato, disci 5-dentato campanulato. Stigmata breviter extus arcuata obtusa. Achænium obcompressum basi attenuatum erostre omnino calvum paleâ inclusum. — Herba Africana aquatica radicans glabra oppositifolia *Enhydra* omnino referens, dichotoma procumbens. Folia integerrima oblonga remota subserrata. Capitula axillaria solitaria sessilia.

1. *C. REPENS* (Beauv.! l. c.). — in Africâ æquinoctiali ad ripas flumin. Formosæ (Beauv.!), in Guineâ (Thonn.). Less. in linnæa 1831. p. 154. *Cæsulia* radicans Willd. sp. 3. p. 1797. *Wahlenbergia glomerata* Schum. guin. 387. *Enhydra cæsulioides* Cass. dict. 14. p. 554. *Meyera* Guineensis Spreng. syst. 3. p. 602. (v. s.)

Trib. IV. *SENECIONIDEÆ* Less. syn. p. 218.

Stylus apice cylindræus, in flor. hermaphroditis supernè bifidus, ramis longiusculis linearibus apice penicillatis, nunc truncatis, nunc ultra penicillum in conum brevem aut appendiculum elongatum angustum et hispidulum productis. Stigmatibus series latiusculæ et prominulæ usque ad penicillum seu originem coni seu appendicis extensæ. Corolla disci pellucida regularis. Pollen globosum echinulatum.

CONSPECTUS TRIBUS.

Subtrib. I. MELAMPODINEÆ. Flores omnes unisexuales nec ulli hermaphroditi, masculi et foeminei in diversis plantis (dioici), aut in earumdem plantarum capitulis diversis (heterocephali), aut in iisdem capitulis (monoici). Antheræ ecaudatæ. Recept. plerumquæ paleaceum. Pappus nunquàm setosus.

Div. I. EUXENIÆ. Capitula dioica multiflora discoidea. Achænia ecorticata.

Div. II. MILLENIÆ. Capitula monoica pauciflora, fl. radii paucis ligulatis aut trifidis. Achænia ecorticata. Recept. sæpius epaleaceum.

Div. III. SILPHIÆ. Capitula monoica multiflora, fl. radii pluribus ligulatis. Achænia ecorticata. Recept. totum aut saltèm margine paleaceum.

Div. IV. MELAMPODIÆ. Capitula monoica multiflora radiata. Achænia corticata.

Div. V. AMBROSIEÆ. Capitula heterocephala, nempe alia ♂, alia ♀. Antheræ marium liberæ. Cor. tubulosæ, interdum in ♂ nullæ.

Div. VI. IVEÆ. Capitula monoica. Cor. tubulosæ aut rariùs in fl. ♀ nullæ. Antheræ liberæ.

Div. VII. PARTHENIÆ. Capitula monoica radiata. Achænia radii obcompressa aut rariùs trigona. Ligulæ tardè deciduæ.

Subtrib. II. HELIANTHEÆ. Capitula sæpiùs heterogama radiata aut homogama discoidea. Recept. totum aut margine paleaceum. Cor. fl. herm. lobi crassi. Pappus nullus coroniformis aut aristatus. Antheræ nigricantes ecaudatæ. — Folia sæpiùs opposita.

Div. I. HELIOPSIDEÆ. Capitula heterogama ligulis foemineis aut discoidea. Achænia grossa cortice crasso sæpiùs adhærente instructa, nunquàm obcompressa. Pappus nullus, coroniformis, aut breviter aristulatus.

Div. II. RUDBECKIÆ. Capitula heterogama ligulis neutris aut sterilibus. Achænia erostria. Pappus nullus aut coroniformis.

Div. III. COREOPSIDEÆ. Capitula heterogama, ligulis neutris rarisimè discoidea. Achænia erostria. Pappus aristis 2-4 et interdum etiàm squamellis constans.

Div. IV. BIDENTIDEÆ. Capitula heterogama ligulis neutris, rarisimè discoidea. Achænia rostrata, rariùs calva, sæpiùs aristis 1-4 superata.

Div. V. VERBESINEÆ. Capitula heterogama ligulis foemineis, rariùs discoidea. Achænia cortice tenui donata, compressa aut obcompressa, calva, aut aristas tot quot anguli, aut setas rigidas fragiles gerentia.

Subtrib. III. FLAVERIÆ. Capitula uni- aut pauciflora dense aggregata heterogama, fl. fœmineis nunc solitariis nunc pluriserialibus, aut heterogama cum prioribus mixta. Achænia calva. Recept. in capit. paucifloris epaleaceum, in multifloris paleaceum. — Folia opposita.

Subtrib. IV. TAGETINÆ. Capitula heterogama fl. radii fœmineis aut discoidea homogama. Invol. sæpius 1-seriale aut gamophyllum. Recept. epaleaceum. Achænium erostre basi attenuatum. Pappus aristatus, rarius setosus. — Folia opposita rarius alterna. et involucria grossè pellucido-glandulosa.

Div. I. TAGETÆ. Capitula sæpius radiata. Cor. disci regulares. Styli rami cono superati. Pappus aristatus.

Div. II. POROPHYLLÆ. Capitula discoidea. Cor. disci inæqualiter fissæ. Styli rami in appendicem subulatam producti. Pappus setosus pluriserialis.

Subtrib. V. HELENIÆ. Capitula sæpius heterogama. Achænium erostre. Pappus squamellis pluribus 1-serialibus scariosis ferè semper liberis constans, rarissimè calvus. — Folia alterna rarius opposita.

Div. I. GAILLARDIÆ. Receptaculum epaleaceum.

Subdiv. 1. EUGAILLARDIÆ. Ligulis neutris.

Subdiv. 2. EUHELENIÆ. Ligulis fœmineis aut nullis.

Div. II. GALINSOGÆ. Receptaculum paleaceum. Involucri squamæ achænia radii non involventes.

Subdiv. 1. EUGALINSOGÆ. Ligulis fœmineis seu nullis.

Subdiv. 2. SPHENOGYNEÆ. Ligulis neutris.

Div. III? MADIÆ. Recept. margine aut ferè totum paleaceum. Invol. squamæ 1-2-seriales convolutæ aut complicatæ achænia radii semper calva includentes.

Subtrib. VI. ANTHEMIDÆ. Capitula sæpissimè heterogama, ligulis fœmineis seu neutris. Antheræ ecaudatæ. Styli rami truncati barbati rarissimè cono superati. Achænium angulatum teres aut in radio obcompressum. Pappus nullus aut coroniformis, rarius squamellis aut pilis capillaribus constans. — Folia sæpissimè alterna.

Div. I. EUANTHEMIDÆ. Receptaculum paleaceum. Capitula radiata.

Div. II. CHRYSANTHEMÆ. Recept. epaleaceum. Capitula radiata.

Div. III. COTULÆ. Recept. epaleaceum. Capitula discoidea homogama aut heterogama.

Div. IV. ATHANASIEÆ. Receptaculum paleaceum. Capitula homogama, cor. omnibus teretibus.

Div. V. ARTEMISIEÆ. Recept. epaleaceum. Capitula discoidea homo- aut heterogama. Cor. disci teretes stylum bifidum gerentes.

- Div. VI. HIPPIÆ. Capitula monoica discoidea. Recept. epaleaceum. Cor. teretes. Stylus disci simplex clavatus.
- Div. VII. ERIOCEPHALÆ. Capitula monoica, fl. foemineis paucis ligulatis tubulosisve. Stylus fl. disci simplex clavato-truncatus. Receptaculum paleaceum.
- Subtrib. VII. GNAPHALIÆ. Capitula homogama aut heterogama, discoidea rarissimè radiata, interdum 1-flora. Antheræ caudatæ. Achænia pappo piloso seu setaceo coronata, rarissimè calva. — Folia sæpius alterna.
- Div. I. ANGIANTHÆ. Capitula pauciflora in glomerulum invol. generali cinctum densè aggregata.
- Div. II. CASSINIÆ. Capitula non glomerata. Recept. totum aut margine paleaceum.
- Div. III. HELICHRYSÆ. Capitula non glomerata. Recept. epaleaceum.
- Div. IV. SERIPHIÆ. Capitula 1-flora, nunc libera, nunc glomerata et invol. generali cincta.
- Div. V. ANTENNARIÆ. Capitula dioica aut subdioica non glomerata. Styli fl. masc. simplices. Recept. epaleaceum aut margine tantum paleaceum.
- Div. VI. LEYSSERÆ. Capitula multiflora radiata, ligulis foemineis. Recept. epaleaceum. Pappus squamellis setisve constans.
- Div. VII. RELHANIÆ. Capitula multi- rariùs pauciflora homogama aut radiata. Recept. epaleaceum. Pappus nullus aut coroniformis.
- Subtrib. VIII. SENECTIONIÆ. Capitula homo- aut heterogama, discoidea aut radiata. Antheræ ecaudatæ. Achænia pappo piloso aut setaceo coronata, exteriora rarissimè calva. — Folia alterna.
- Div. I. NEUROLINÆ. Receptaculum paleaceum.
- Div. II. ERECHTITÆ. Recept. epaleaceum. Capitulum discoideum heterogamum, flor. radii tubulosis foemineis.
- Div. III. EUSENECTIONIÆ. Recept. epaleaceum. Capitulum aut homogamum discoideum, aut radiatum. Cotyledones planæ.
- Div. IV. BALBISIÆ. Recept. epaleaceum. Capitulum discoideum homogamum aut radiatum subdioicum. Cotyledones involutæ.
- Subtrib. I. MELAMPODINÆ. — Melampodiæ et Ambrosiæ *Less.* 213 et 218. — Ambrosiæ, Milleriæ et Baltimoreæ *Cass. opusc.* 3. p. 60 et 61. — Polygamia necessaria *Linn.*
- Flores omnes unisexuales, nec ulli hermaphroditi, in iisdem aut diversis plantis, et in eadem stirpe in iisdem aut diversis capitulis siti, nempe monoici aut dioici. Antheræ ecaudatæ. Recept. ferè semper paleaceum. Pappus nullus, subcoroniformis aut aristatus, nunquam setosus. Styli rami pilis collectoribus sæpius deficientibus sæpius subdeformati, aut more Senecionidearum apice barbati.

Div. 1. EUXENIÆ. — Asteroideæ Melampodicæ Euxeniæ Less.
syn. 216.

Capitula dioica, multiflora, discoidea. Achænia ecorticata.

CCLXI. EUXENIA Cham. hort. ber. (1820). p. 75. t. 16. (excl. syn.) Cass. dict. 35. p. 446. 50. p. 472. Less. syn. 216 non Nees et Mart. — Ogiera Spreng. syst. 3. p. 674 non Cass. — Podanthus Lag. nov. gen. (1816). p. 24. Cass. dict. 46. p. 404. Spreng. syst. 3. p. 456. Less. syn. 419 non Haw. (1812).

Capitula dioica globosa homogama discoidea. Invol. 1-seriale patulum demum reflexum, squamis oblongo-lanceolatis 10 liberis æqualibus aut 2 oppositis sæpè majoribus. Recept. convexum paleis oblongis subobtusis deciduis onustum. Cor. tubo angusto, fauce campanulata extus hirsuta et glandulosa, limbo 5-dentato. ♂ anth. ecaudatæ parum cohærentes (nec liberæ) nigricantes. Stylus inclusus, ramis brevibus crassis obtusis dorso hispidulis. Ovar. abortivum. ♀ anth. abortivæ liberæ. Styli rami crassi obtusi exserti. Achænium obpyramidatum basi attenuatum tetragonum calvum aut rarius pappo subauriculiformi coronatum glabrum apice pilis minutissimis circa areolam subinstructum. — Frutices Chilenses dichotomi ramosissimi, ramis teretibus junioribus adpressè pubescentibus. Folia opposita breviter petiolata ovata aut ovali-lanceolata acuta serrata triplinervia utrinquè subscabrida. Pedunculi 1-3 è dichotomis orti 1-cephali. Flores flavi.

1. E. GRATA (Cham. hort. ber. p. 75. t. 16), foliis latè ovatis in petiolum non decurrentibus. ♂ in rupestribus apricis Chilensibus (Domb.!), nempè ad Talcaguano (Cham.), Conception (Beech.), in Brasiliâ? (Nees et Mart. Beitr. in nov. act. nat. cur. 12. p. 7). Cass. 50. p. 472. Hook. et Arn. bot. Beech. 1. p. 57. Ogiera triplinervia Spreng. syst. 3. p. 674 non Cass. Baccharis Chilensis Weium. syll. pl. nov. 2. p. 21 ex Less. Podanthus ovatifolius Lag. nov. gen. p. 24. n. 322. Vulgò Chilensibus dicitur *Palo-Negro* ex Lag. (v. s.)
2. E. MITQUI, foliis ovali-lanceolatis basi longè cuneatis apice acuminatis medio grossè serratis. ♂ in Chili circa Valparaiso (Macr.! Gaud.!), in sylvaticis collium ad Quillota et Rancagua (Bert.!), Podanthus Mitqui Lindl. in litt. 1832. Loud. hort. brit. p. 488. Ogiera triplinervia Bert. herb. non Cass. Vulgò Chilensibus dicitur *Mitri* ex Bert. et verosim. *Mitqui* ex nom. spec. Lindleyano. Priori affinis sed foliorum formâ sat distincta. (v. s.)

β. *subintegerrima*, foliis subintegerrimis. ♂ in Chili. Græmia aromatica Poepp. pl. exs. n. 208. diar. n. 30 non Hook. Folia minora ferè omnia integerrima, hinc indè subdenticulata. An fortè species propria? (v. s.)

† *Species exclusa.*

E. RADIATA Nees. = *Gymnosoma eusenoides*.

CCLXII. PETROBIUM R. Brown trans. Linn. soc. 12. p. 113. — Laxmannia Forst. nov. gen. n. et t. 47 non Schreb. nec R. Br. nec Fisch (1). — Spilanthi spec. Forst. — Drimiphyllum Burch mss.

Capitula pluri- (10-12) flora dioica. Recept. paleis membranaceis invol. squamas referentibus inter flores sitis. Invol. subcampanulatum biseriale oligophyllum floribus brevius, squamis ovali-oblongis glabris. Cor. tubus teres vix aristis achæni longior, limbo 4-partito, lobis ovali-

(1) Laxmannia Schreb. (1791) est Cyminosina. Laxmannia Fisch. (1812) est Stictogonum sect. Gei. Laxmannia Brown (1810) est genus inter Asphodelas. Stricto jure genus Forsteri servandum et novum genus ad Brownii nomen fingendum esset, sed juris prioritatem hic neglexi nè augeatur confusio.

oblongis. ♂ antheræ exsertæ appendiculatæ. Stigmata acuta hispidula. ♀ stam. distincta. Stigmata acuta recurva. Achænium lineare plano-compressum, ad latus utrinquè medio nervigerum, sparsè hirtellum, apice productum in aristas 2 rigidulas ex angulis ortas margine strigillis erectis subciliatis. Interdum arista tertia longior adest. — Arbor triorgyalis Heleniana. Ramuli alterni. Folia opposita petiolata latè obovata seu ovalia, anticè serrata, suprà glabra, subtus sparsè subpuberula. Panícula (ex Burch.) terminalis oligocephala ramulis oppositis. Capitula longè pedicellata erecta albida.

1. *P. ARBOREUM* (Brown l. c.), 5 in insulâ Stæ-Helenæ ubi Anglis dicitur *White-Wood*, *Cabbage-Tree*. *Bidens arborea* (fœm.) et *Spilanthes tetrandra* (masc.?) Roxb. in *Beata*. app. p. 301 et 325. *Spilanthes arboreus* G. Forst. comm. goett. 9. p. 66. *Laxmannia arborea* Forst. nov. gen. t. 47. *Spilanthes pseudogummifera* Forst. in h. Mus. Par. *Drimyphyllum Helenianum* Burch. in litt. (v. s. specim. fœm. comm. à cl. Burch.)

CCLXIII. *ASTEMMA* Less. *syn.* 216. — *Monactidis* spec. Kunth.

Capitula 10–15-flora homogama discoidea dioica. Invol. tubuloso-campanulatum, squamis linearibus obtusis subimbricatis. Recept. paleis squamas referentibus membranaceis ciliatis. Cor. tubulosa dentibus 5 revolutis, fœmineæ stam. sterilia gerentes. Achænium lineare calvum breviter rostratum stipitatumque. — Arbor Americæ æquinoct. Rami angulati puberuli. Folia alterna subdenticulata subtriplinervia. Panícula fastigiata terminalis ramosissima foliata. Corollæ albæ ex Kunth, flavæ ex Less. — Genus valdè dubium.

1. *A. DUBIUM* (Less. l. c.), 5 in valle Tarquensi regni Quitensis. *Monactis dubia* H. B. et Kunth nov. gen. am. 4. p. 287.

Div. II. MILLERIÆ. — *Helianthæ* Milleriæ veræ et Baltimoreæ *Cass. opusc.* 60. — *Asteroidæ* *Melampodiæ* Milleriæ et *Senecionidæ* *Ambrosiæ* Less. *syn.* 2–15 et 219.

Capitula monoica, fl. nempe in iisdem capitulis fœmineis sæpè paucis ligulatis aut tubulosis subtrifidis, masculis centralibus tubulosis. Achænia ecorticata nempe cum squamis inv. seu paleis rec. non concreta exalata, calva aut rariùs pappo coroniformi parvo superata. Recept. interdum epaleaceum. Antheræ concretæ.

- CCLXIV. *ELVIRA*. — *Elvira* Cass. dict. (1824) 30. p. 67 et *Meratia* Cass. l. c. p. 65 non *Lois*. — *Delilia Spreng. bull. philom.* 1823. apr. non Kunth. — *Eugamelia fl. mexic. ic. ined.* Milleriæ sp. L.

Capitulum monoicum biflorum, flore altero fœmineo ferè ligulato, altero masculo tubuloso 5-fido. Invol. 2–4-phyllum squamis oppositis, altero majore orbiculato basi cordatâ subcrenato, altero minore ovali, 3°–4°ve ubi adsunt oblongæ acutæ. Recept. minimum epaleaceum. Cor. fœminea longè tubulosa; ligulâ brevi integerrimâ? Anth. ♂ vix exsertæ. Styl. abortivus breviter bifidus. Styl. ♀ apice bifidus. Achænium plano-obcompressum ovale involucri immutato tectum nec involutum, calvum. — Herba Americana annua trichotoma. Folia opposita petiolata ovata aut oblonga acuminata serrulata triplinervia pubescenti-subscabrida. Pedicelli in axillis supremis aggregati 1-cephali subumbellati. — Descr. Sprengelii pessima; ex eâ sagaciss. Cassinius finxit *Meratiam* suam cujus *descriptio* multò præcellit descr. autoptæ sed adhuc tamen subincorrecta. Nomen et descr. fl. Mex. optima, sed inedita et ideò *Elviræ* no-

men admittendum, etiamsi sensûs saltem cogniti expers. An capit. interdûm 3-florum, ut patet ex descr. Linnæi et Spreng.?

1. *E. biflora*. (1) in agro Campechiano (Mart.), in umbrosis circa Bahiam (Salzm.), Novâ-Andalusiâ prope Cumanacoa (H. et B.), insulâ Cubâ juxta Havanam (Lasagr.), ad Santa-Martha (Bert.) et in Mexici ditone Oaxacâ circa prædium Tapanala (Alam.). *Milleria* Mart. cent. t. 47. f. 1. *Linn.* hort. Cliff. t. 25 *Milleria biflora* Linn. sp. 1301 *Gærtn.* fr. 2. p. 423. t. 168. *Lam.* ill. t. 710 f. 1. *Delilia Berteri* Spreng. bull. phil. 1823 apr. syst. 3. p. 674. *Elvira Martyni* et *Meratia Sprengelii* Cass. H. cc. (v. s.)

CCLXV. *MILLERIA* Cass. dict. 30 (1824). p. 67. 59. p. 235. Less. syn. 215. *Milleriæ* sp. Linn. Mart. Gærtn.

Capitulum monoicum pauciflorum, fl. foemineo ligulato, cæteris masculis tubulosis 5-dent. Invol. 3-5-phyllum, altero majore cucullato. Recept. epaleaceum. Ligulæ profundè trifidæ. Stylus ♂ integerrimus, ♀ bifidus. Achænium nitidum plano-obcompressum calvum involucri demum carnosio indurato atque irregulariter gibbo omnino involutum, areolâ laterali. — Herba Americana annua trichotoma subvillosa. Folia opposita petiolata ovato-rhomboidæ tenerè membranacea triplinervia dentata. Stipula ovata sæpè reflexa utrinquè inter folia ut in Rubiaceis. Pedunculi terminales dichotomi, pedicelli è dichotomiis et apice ramorum. Cor. luteæ.

1. *M. quinqueflora* (Linn. sp. 1301), caule subtetragono pubescente, foliis latè ovatis subcordatis abruptè acuminatis serratis infernè in petiolum longè decurrentibus. (1) in Panamâ, Verâ-Cruce, Mexico (Linn. Cav.), in valle Sancti-Jacobi (H. et B.), in Chilapæ circuitibus (fl. mex. ic. ined.). *Gærtn.* fr. 2. p. 423. t. 168. f. 5. *Lam.* ill. t. 710. f. 2. *M. dichotoma* Cav. ic. 1. p. 58. t. 82. — Mart. cent. t. 41. Canlia obtusè tetragonus variat glaber pubescens aut villosus. *Florum capituli* 5-9 (v. s.)

β. *maculata*, foliis maculatis profundius serratis Mart. cent. t. 47. f. 2. — *M. maculata* Mill. dict. n. 2.

2. *M. glandulosa*, caule tereti pilis glandulosis asperulo, foliis ovalibus acuminatis serraturis callosis minimis donatis, in petiolum latè alatum attenuatis. (1)? in Mexico legit cl. Berlandier (pl. ex. m. 955!). Petiolus seu nervus medius infra foliū dilatationem latius, superne glaber, ad marginem utrinquè serie pilorum quasi sursum ciliatus. Pedicelli fructu ferè breviores. Capitulum minimum. Fructus grossus obtusus irregulariter rugosus constans squamis 2 coriaceis subinæqualibus, achænio obovato levè longitudinaliter striatulo apice mucronulato calvo. Cæter. ut in ic. cit. *Gærtn.* (v. s.)

† Species exclusæ.

M. alba Hortul. — *Fougerouxia alba*.

M. angustifolia Cav. = *Flaveria angustifolia*.

M. biflora Linn. = *Elvira biflora*.

M. contrayerva Lam. = *Flaveria Contrayerva*.

CCLXVI. *RIENCOURTIA* Cass. bull. philom. 1818. p. 76. dict. 45. p. 466. Less. syn. 215. — *Petrantha* Poit. ms.

Capitulum monoicum 4-7-florum discoideum heterogamum, fl. disci 3-6 masculis, radii unico foemineo tubuloso. Invol. oblongum disco brevius, squamis biserialibus in utrâque serie 4 æqualibus adpressis coriaceis. Recept. parvum nudum. ♂ Cor. disci fauce ampliâ, limbo 4-5-dentato, lobis fasciculo pilorum superatis. Anth. nigricantes. Sty-

lus symplex exsertus. ♀ Cor. radii cylindrica 3-dentata. Styli rami duo. Achænium obcompressum ovale s. orbiculatum calvum. — Herbæ erectæ ramosæ Australi-Americanæ. Folia opposita petiolulata lanceolata 3-nervia pilosa subdentata. Capitula spicata vel glomerata ad apices ramorum supernè aphyllorum. — Conjux cl. Cassinii orta è gente de Riencourt.

1. *R. SPICULIFERA* (Cass. l. II. cc.), capitulis in spicam dispositis distichis ad bractearum axillas sessilibus, rachi hispidâ, bracteis ciliatis. — in Americâ austr. ? Corollæ masculæ 5-dentatæ. (v. s. specim incomp. in h. Juss.)
2. *R. GLOMERATA* (Cass. dict. 43. p. 370. 45. p. 468), capitulis plurimis in glomerulos terminales solitarios subglobosos collectis bracteatis, bracteis apice piloso-hispidis. — in Guianâ gallicâ. *Tetrantha suaveolens* Poit. ined.

CCCLXVII. LATREILLEA.

Capitulum monoicum multiflorum, fl. radii 3-4 fœmineis, disci masculis. Recept. subspiciforme. Invol. squamæ 1-3-seriales, exteriores breves potiùs bracteæ dicendæ, inter. majores concavæ. Rec. paleæ latè ovales suborbiculatæ ciliatæ inter flores sitæ. Cor. radii breves tubulosæ subtruncatæ pilos articulos extus gerentes, disci 5-fidæ glabræ. Antheræ concretæ ecaudatæ. Stylus radii bifidus, ramis 2 crassius glabris acuminatis intus sulcatis, disci simplex inclusus abortivus. Achænia radii grossa glabra obcompressa, disci abortiva. — Herba Brasiliensis habitu paululùm *Chromolaenæ*. Rhizoma lignosum. Caulis herbaceus basi striatus glaber apice puberulus aut hispidus. Folia opposita sessilia oblonga, infer. glabra, super. hirtella. Corymbus breviter trifidus hirtus, capitulis circ. 3-5 ovatis lupulaceis. — Dixi in memoriam cl. Petr. Andr. Latreille entomologi peritissimi, de geographiâ etiâ benè meriti et omnibus naturæ curiosis morum amenitate cari.

1. *L. SERRATA*, caule dichotomo basi glabro apice puberulo, foliis oblongo-lanceolatis à medio ad apicem grossè et remotè calloso-serratis. ½ in campis editis prov. Sancti-Pauli novemb. flor. legit cl. entomologus Lund. Folia 3 polli. longa, 4-6 lin. lata subtriplinervia. Capitula nivea. Invol. veri squamæ basi ovatæ apice acuminatæ. Rec. paleæ apices latæ fimbriatæ basi angustatæ. (v. s. comm. à cl. Lund.)
2. *L. INTEGRIFOLIA*, caule simplici basi glabro apice hispido, foliis lanceolatis integerrimis. ½ in Brasiliæ prov. Sancti-Pauli ex h. Mus. imp. Bras. n. 447 ! Invol. squamæ obtusissimæ dorso hispidæ. (v. s. in h. Mus. reg. Par.)

CCCLXVIII. ICHTYOTHERE Mart. arzn. pfl. p. 27.

Capitulum monoicum, floribus omnibus tubulosis, radii fœmineis subquinis fertilibus, corollulæ limbo in fimbriis tenuissimas laciniato; disci plurimis 5-fidis. Involucrum minimum cupuliforme persistens. Recept. paleaceum, paleis singulis florem singulum fœventibus. Antheræ basi intus denticulo auctæ. Stylus radii bifidus, disci sterilis. Achænia intra paleas extimas concavas grandefactas obovata absque pappo. — Herba perennis, foliis oppositis, petiolis breviter vaginantibus limbis argutè serratis. Capitula florum pedunculata terminalia aggregata, rachi post grossificationem per dichotomiam excrescente, involucri persistentibus tunc grandefactis. — Affinis *Baillieria Cacosmiæ* et *Melampodio* ex Mart.

1. *I. CUNABI* (Mart. l. c.). — in Brasiliæ prov. Rio-Negro. Pisces inebriat undè nom. gener.

CCLXIX. CLIBADIUM *Linn. mant.* 161 *non Neck.* — Clibadium, *Oswalda et Baillieria Cass.* 59. p. 324. — *Baillieria et Oswald Less. syn.* 215. — *Baillieria Aubl.* — *Trixis Swartz, Pers. non P. Brown.*

Capitulum monoicum pluriflorum discoideum, fl. radii paucis tubulosis 5-fidis hermaphroditis sterilibus. Invol. ovatum, squamis conniventibus ovatis concavis, interioribus achænia radii in parte concava semifoventibus. Recept. planum in disco omnino epaleaceum, paleas seu invol. squamas inter. in radio interdum biseriali gerens. Stylus radii bifidus, ramis divaricatis subrevolutis glabriusculis subobtusis, disci ex *Cass.* indivisus vix subbifidus apice clavatus hispidulus. Achænia radii grossa obovata intus obcompressa extus convexa epapposa apice pilis articulatis villosa, disci abortiva gracilia linearia apice villosissima. — Herbarum seu suffrutices Americani. Rami teretes sæpius subvillosi. Folia opposita petiolata ovata aut lanceolata serrata plerumque aspera. Capitula ovata albida parva in paniculam sæpius corymboso-fastigiatam disposita.

Sect. I. EUCLIBADIUM. — Clibadium *Linn. l. c. Cass. dict.* 9. p. 395. 29. p. 176 et 181. 59. p. 324.

Achænia drupacea succo viscoso scatentia, radii uniserialia. Paleæ recept. nullæ.

1. C. SURINAMENSE (*Linn. mant.* 294), foliis petiolatis ovatis acuminatis acute crenatis scabris, pedunculis oppositis. — Surinami. C. foetidum *Alam. ex Linn.* Invol. violaceum. Cor. alba. — Species ab hodiernis ign. Habui ex Amer. sub hoc nom. fragmentum sine flore, cum descriptione non malè conveniens. Rami et petioli hirsutuli. Petioli 10-12 lin. longi. Folia 4 poll. longa, 2 $\frac{1}{2}$ poll. lata, argutè serrata, suprà scabrida, juniora villosa demum scabrida. (v. s.?)

Sect. II. OSWALDA *Cass. dict.* 59. p. 319 et 322. *Less. syn.* 215. — *Baillieria Aubl.* — *Baillieria* sp. *Spreng.* — *Trixis* sp. *Sw. Pers.*

Achænia sicca villosa, radii uniserialia obcompressa obovata dorso convexa, disci gracilia linearia apice hispida. Paleæ recept. nullæ.

2. C. ROTUNDFOLIUM, foliis brevissimè petiolatis ovato-subrotundis basi triplinerviis utrinque subscabridis, paniculà corymboso-fastigiatà. ζ in Brasilià circa Bahiam legerunt cl. Salzmann, Blanchet, etc., in prov. Sancti-Pauli (h. Mus. Bras. n. 466). Petioli 3-4 lin. longi. Limb. fol. 3 poll. longus, circit. 2 poll. latus. Capitula ovato-globosa. Pedunculi hirsuti. Inv. squamæ pubescentes, inter. glabræ multistriatæ acutæ. (v. s.)

β . Foliis ovatis acutis. — in prov. Sancti-Pauli (Lund!), et prov. Minarum Generalium (Gaudichaud!). (v. s.)

3. C. PERUVIANUM (*Pœpp.*! pl. exs. n. 35. diar. n. 1611), foliis petiolatis ovali-lanceolatis longè acuminatis basi cuneatis adpressè serrulatis triplinerviis suprà scabris subtus ramis petiolisque hirsutis, paniculà corymboso-fastigiatà. ζ in Peruvia legit cl. Pœppig. Petioli 4-5 lin. longi. Limb. fol. 4 poll. longus, 10-12 lin. latus. Capit. ovato-globosa. Inv. squamæ ext. dorso puberæ, intimæ glabræ striatæ apice puberæ acutæ. (v. s.)

4. C. TRINITATIS, foliis petiolatis ovali-lanceolatis basi longè cuneatis apice acuminatis acutè serrulatis subtrinerviis, suprà secus nervos et vix inter nervulos scabridis, subtus ramisque hirtello-subhirsutis, paniculà ovatà. \mathcal{U} ζ in insulà Trinitatis legit cl. Sieber fl. trin. pl. exs. n. 71! Petioli 6-7 lin. longi. Limb. fol. 5 poll. longus, 1 $\frac{1}{2}$ latus. Capitula ovato-globosa parva. Panicula non fastigiata, foliis subbrevior. Inv. squamæ paucæ extus subpuberæ apice crosso subdenticulatæ. (v. s.)

5. *C. CARACASANUM*, foliis petiolatis ovato-lanceolatis basi breviter cuneatis apice acuminatis argutè serratis triplinerviis suprâ scabris subtus molliter puberohirsutis, ramis piloso-scabris, paniculâ corymboso-fastigiatâ. 2½ 3 circa Caracas legit cl. Vargas. Petioli 5 lin. longi. Limb. fol. 3 poll. longus, 15-16 lin. latus. Capitula alba subglobosa vix pubera. Achenia disci abortiva à basi hirsutissima. (v. s. comm. à cl. Vargas.)
6. *C. ASPERUM*, foliis petiolatis ovato-lanceolatis basi breviter cuneatis apice acuminatis argutè serrulatis triplinerviis utrinquè scabris, ramis adpressè puberulis, paniculâ corymboso-fastigiatâ. 2½ 3 in Cayennâ (Aubl. Pers.), prope Tioselo Mexic. (Schiede). Baillieria aspera Aubl. guian. 2. p. 805. t. 317. Trixis aspera Pers. ench. 2. p. 491. Trixis scabra Swartz fl. Ind. occ. 1379. Oswaldia baillieroides Cass. dict. 59. p. 322. Less. in linnæa 1834. p. 267. Petioli 5 lin. longi. Limb. fol. 3 poll. longus, æquipoll. latus. Capitula subglobosa extus subpubera. Achenia disci abortiva apice tantum hispida. Desor. Cass. optima: Aubl. manca, sed ex patriâ et iconè huc referenda. Herba amara odore pisces inebrians. — An Baillieria sylvestris Aubl. l. c. p. 307 etiamsi foliis angustioribus satis diversa? (v. s.)
7. *C. VARGASII*, foliis petiolatis ovali-lanceolatis basi cuneatis apice acuminatis argutè serrulatis triplinerviis suprâ vix subscabris subtus vix subhirsutis, ramis apice adpressè puberis, paniculâ ovatâ non fastigiatâ. 3 circa Caracas legit cl. Vargas. Petioli 4 lin. longi. Folia 2 poll. longa, 7 lin. lata. Capitula extus pubera. (v. s.)
8. *C. HAVANENSE*, foliis petiolatis ovali-lanceolatis utrinquè longè acuminatis medio grossè et remotè serratis triplinerviis utrinquè setis sparsis paucis subscabris, ramis apice adpressè puberis, paniculâ laxiusculâ non fastigiatâ. 3 in insulâ Cubâ prope Havanam legit cl. de La Ossa. Petioli 5-6 lin. longi. Limb. fol. 5-6 poll. longus, 18-20 lin. latus. Capitula parva. (v. s.)
9. *C. TEREBINTHINACEUM*, foliis petiolatis ovali-lanceolatis basi cuneatis apice acuminatis triplinerviis puberulo-scabris subtus hirsutiusculis, ramis petiolisque brevissimè subhirsutis, paniculâ subcorymbosâ. 3 in insulis Caribæis, in Jamaica occident., nempe vallibus saxosis (Sw.). Trixis terebinthinacea Swartz! fl. ind. occ. 3. p. 1375. t. 26. Baillieria terebinthinacea Spreng. syst. 3. p. 624. Petioli 7-8 lin. longi. Limb. fol. 4 poll. longus, 15 lin. latus. In ipso specimine recept. vidi nudum. (v. s.)
- β. *Badieri*, foliis glabrioribus grossius serratis. 3 in Guadalupâ legit cl. Badier. (v. s.)

Sect. III. *TRIXIDIUM*. — Baillieria Cass. et Less. ex car. — Trixidis sp. Swartz.

Achenia radii sicca apice villosa biserialia, ext. squamis, inter. paleis squamas omnino referentibus extus semi-involuta, disci abortiva epaleacea.

10. *C. EROSUM*, foliis petiolatis latè ovatis basi cuneatis breviter acuminatis grossè et irregulariter inciso-serratis penninerviis rugosis scabris subtus ramisque subhirsutis, paniculâ corymbosâ. 3 in insulis Caribæis Sancti-Christophori et Dominicæ (Sw.), Trinitatis (Sieb. fl. trin. n. 319), Guadalupæ (Badier!). Trixis erosa Swartz fl. ind. occident. 1377. Baillieria erosa Spreng. syst. 3. p. 624. Petioli 7-9 lin. longi. Limb. fol. 5 poll. longus, 3 poll. latus. Capitula latiores quam priorum præsertim post anthesin, evidenter in radio paleacea, in disco epaleacea. (v. s.)

† *Species non satis notæ.*

11. *C. P. COMMELINOIDES*, herbaceum, foliis oppositis cordato-lanceolatis integerrimis nervosis utrinquè glabris, capitulis terminalibus subsessilibus, squamis receptaculi obtusissimis. — in Brasiliâ. Rolandra terminalis Spreng. syst. 3. p. 673. Baillieria commelinoides Less. syn. 215.

12. C. PEDUNCULOSUM, herbaceum, ramis gracilibus apice summitate promissè nudis 2-3-cephalis, foliis lanceolatis serratis 5-nerviis hispidis. — in Cayennâ. Trixis pedunculosa Rich. act. soc. h. n. Par. 1792. p. 112. Pers. ench. 2. p. 491. Baillieria pedunculosa Spreng. syst. 3. p. 625.
13. C. BARBASCO, fruticosa, foliis oppositis acutis basi valdè angustatis remotè et obsoletè serrulatis suprâ pilosiusculis subtus adpresso-pubescentibus, corymbis semilibus trichotomis, recept. paleis oblongis singulis flore masculo duplò brevioribus, ovario compresso? dentibus 2 minutis apice instructo. 5 in ripâ flum. Tuamini et Tima prope Javita in missionibus Orinoci. Baillieria Barbasco H. B. et Kunth nov. gen. am. 4. p. 189. Vulgò dicitur *Barbasco*. An hujus generis?
14. C. MERIIFOLIUM, arboreum, ramis albo-lanatis, foliis sparsis verticillato-congestis lanceolatis integerrimis coriaceis suprâ glabris subtus albo-lanatis, corymbis terminalibus villosissimis, invol. squamis interioribus lineari-lanceolatis, recept. paleaceo convexo-conico, fl. fœmineis ligulatis, achænio margine villoso coronato. 3 in monte Sillâ de Caracas ubi dicitur *Incienso*. Baillieria meriifolia H. B. et Kunth nov. gen. am. 4. p. 189. Trixis meriifolia Humb. rel. hist. 1. p. 645. A genere certè removenda! An Melampodii sp.?

CCLXX. UNXIA Linn. f. suppl. 56. Juss. gen. 186. Less. syn. 220 non Kunth.

Capitulum monoicum pluriflorum, fl. radii 5 ligulatis fœmineis parvis ovatis, disci 5 masculis tubulosis 5-dentatis. Invol. subrotundum, squamis 5 ovatis. Recept. nudum? Achænia radii ovata dura (Linn. f.) compressa lævissima calva, areolâ laterali (Less.). — Herbæ dichotomæ Guianenses. Folia opposita hirsuta oblongo-obovata acuminata basi rotundata 5-nervia villosa. Capitula axillaria breviter pedunculata. — An Clibadio affinis? †.

1. U. CAMPHORATA (Linn. f. suppl. 368), foliis lanceolatis, capitalis 10-floris. — in locis arenosis Surinam. Odor gravis camphoratus. *Camphorplant Surinam* Holland.
2. U. HIRSUTA (Rich. act. soc. Par. 1. p. 105), foliis oblongis subcordatis, capitulis multifloris. (1) Cayennæ. Pers. ench. 2. p. 492.

†† Species exclusæ.

- U. ACHILLÆOIDES Spr. = Villanova achillæoides.
- U. ALTERNIFOLIA Spr. = Villanova alternifolia.
- U. ANEMONIFOLIA H. B. K. = Villanova anemonifolia.
- U. ANTHEMIFOLIA Bert. = Apalus anthemifolius.
- U. DISSECTA Hook. = Villanova dissecta.
- U. PRATENSI H. B. K. = Chlamisperma pratense.

CCLXXI. APALUS.

Capitulum monoicum, fl. radii circ. 5-7 uniseriatis fœmineis ligulatis ovali-oblongis integris, disci circ. 8-10 masculis glabris tubo tereti limbo campanulato 5-dentato. Invol. 5-7-phyllum uniseriale, squamis ovatis. Recept. nudum angustum. Styl. radii bifidus, ramis teretibus extrorsum arcuatis. Achænia radii oblonga erostria calva sulcato-angulata pube glandulosâ brevi crassâ velutino-hirtellâ, disco epigyno. — Herba Chilensis tenella annua erecta glabra versus apicem subvillosa. Folia alterna semimplexicaulia, basi linearia petioliiformia integra apice pinnatipartita, lobis linearibus oppositis integris obtusis utrinquè 2-4. Rami apice pedicelliiformes 1-cephali. Cor. disci et radii

flavæ. — Differt ab *Unxiâ* achæniis nec compressis nec areolâ laterali donatis; à Villanovâ et *Chlamiispermate* disco masculo; ab omnibus achæniis glanduloso-puberis et foliis alternis. — Nomen ex ἀπαλις, tener.

1. *A. ANTHEMIFOLIUS*. (1) In pascuis sterilibus circa Rancagua, Valparaiso et ad montem La Leona Chilensium legit cl. Bertero. *Unxia anthemifolia* Bert. l herb. n. 726.

CCLXXII. PRONACRON *Cass. dict. 43. p. 370. Less. syn. 214.*

Capitulum monoicum multiflorum, fl. radii 5 uniseriatis foemineis ligulatis, disci circ. 12 masculis tubulosis. Invol. subglobosum biseriale disco æquale, squamis 7, duabus ext. rotundatis villosis submajoribus, 5 int. glabris membranaceis. Recept. planiusculum, paleis paucis inæqualibus subulatis. Cor. radii tubo glanduloso, ligulâ ellipticâ subintegrá. Achænia glabra, disci gracilia, radii lateraliter plano-compressa sublenticularia crassa lævia truncata calva. — Herba ramosissima Guianensis villosa. Folia opposita lanceolata subdentata. Capitula pedunculata terminalia inter ramulos novellos orta. Flores flavi.

1. *P. RAMOSISSIMUM* (Cass. l. c.). (1) ? in Guianâ gallicâ.

CCLXXIII. AIOLOTHECA.

Capitulum monoicum multiflorum, fl. radii circ. 5-6 ligulatis foemineis, disci tubulosis 5-dent. masculis. Invol. biseriale; squamis exter. oblongis erectis 8-10 disco brevioribus, inter. paleas referentibus et achænia radii semi-includentibus. Recept. parvum convexum, paleis membranaceis cuneatis apice dentatis acutis achænia involv. Ligulæ breves obtusæ 3-dentatæ. Achænia radii villosa subtrigona epapposa villis subcoronata inter squamas int. et paleas ext. inclusa, disci gracilia glaberrima calva. — Herba Mexicana incana apice corymboso-ramosa. Caulis teres basi glabratus. Folia alterna sessilia oblonga obtusa sinuata. Capitula corymboso-fastigiata. Cor. luteæ. — Habitus herbæ affinis *Parthenio* incano sed fructus diversus et flores lutei. — Nomen ex αἰολος, varius, et θῆκη, theca.

1. *A. PARTHENIOIDES*. — in Mexici ditione Sancti-Ludovici de Potosi decembr. flor. legit cl. Berlandier (pl. exs. n. 1341). Herba circ. pedalis. (v. s.)

CCLXXIV. TRIGONOSPERMUM *Less. syn. 214. et in linnæâ 1834. p. 267.*

Capitulum monoicum minutum pluriflorum, fl. radii 3 foemineis ligulatis profundè 3-4-lobis parvis, disci crebrioribus masculis tubulosis. Invol. 6-8-phyllum, squamis linearibus acutis 1-nerviis, alternis majoribus rotundis. Recept. paleis scariosis difformibus, ext. majoribus erosis, int. linearibus. Stylus ♂ integerrimus inclusus abortivus, ♀ bifidus. Achænia disci 0, radii triangularia squamâ int. et palcâ ext. cincta lucida calva. — Herba annua Mexicana erecta dichotoma. Folia opposita triplinervia rhombeo-ovata breviter acuminata usquè ad basin petioli cuneato-decurrentia utrinquè hirsutiuscula. Paniculæ cymosæ. Rami glanduloso-hirsuti. Capitula lutea.

1. *T. ADENOSTEMOIDES* (Less. l. c.), capitulis minimis subdiscoideis, ligulis nempe vix discum superantibus. (1) in Mexico circa Jalapan. Herba ultrâ-orgyalis. Capitula parva. Cor. masc. 4-dentatæ. Folia 6 poll. longa, 3 lata.

2. *T. MELAMPODIODES*, capitulis evidenter radiatis, ligulis nempe disco triplò circ. longioribus. — in Mexico ad montem San-Felipe prope Oaxacam legit cl. Andrieux (pl. ex. n. 320). Caulis subhirsutus. Folia 3 poll. longa, sesquipoll. lata. Paniculae rami pilos parcos et glandulas turbinatas gerentes. Capitula magn. ferè Siegesbeckiæ. Ligulae 3-5 elongatæ apice irregulariter lobato-dentatæ, discique cor. aurantio-flavæ. Achænia trigona nigra. (v. s. ex coll. Andr.)

CCLXXV. XENISMIA.

Capitulum monoicum, fl. radii 5-7 fœmineis ligulatis, disci 10-12 tubulosis 5-dentatis masculis. Invol. 1-serialis squamæ oblongæ. Recept. epaleaceum. Ligulae obtusæ basi ciliatæ. Anth..... Styl..... Achænia disci 0, radii crassa glabra aculeis rigidis crassis undiquè echinata calva. Semen crassiusculum oblongum basi attenuatum. — Herba Australi-Africana annua subpuberula multicaulis erecta. Radix exilis. Folia alterna oblongo-cuneata apice dentata basi attenuata. Capitula ad ramorum apices pedicellata solitaria parvula. — Nomen (ex ξενισμός, peregrinitas) alludit ad habitationem plantæ quæ sola totius divisionis in Africâ versatur.

1. *X. ACANTHOSPERMA*. (1) in Africâ austr.-orient. legit cl. Drege! Herba 5-6 poll. Habitu Acanthospermum humile paulò referens. (v. s.)

CCLXXVI. SCOLOSPERMUM Less. in *linnæa* 1830. p. 152. t. 2. f. 19-31. syn. p. 219.

Capitulum pluriflorum monoicum, fl. radii fœmineis ligulatis, disci masculis tubulosis obconicis 5-dentatis. Invol. triserialis squamæ in seriebus 2 ext. septenæ subæquales. Recept. paleaceum. Stylus fl. ♂ indivisus exsertus teres obtusiusculus apice hirsutiusculus. Achænia calva triquetra versus apicem tuberculata in squamis invol. intimis capsulatum inclusa. Embryo triquet. Cotyledones conduplicatæ. — Herba Mexicana erecta dichotoma Fougereuxiæ facie. Folia opposita petiolata ovato-lanceolata acuminata serrata triplinervia. Capitula in dichotomiis axillaria et terminalia pedunculata lutea.

1. *S. FOUGEROUXIAE*. — in Mexico ad Santa-Fé (Schied.) et ad Panamam (Hænk.). *S. Baltimoroides* Less. ll. oc. Petioli breves hispidi. Folia utrinquè adpressè villosa. (v. s. in h. Hænk. à cl. de Sternberg miss.)

CCLXXVII. FOUGEROUXIA. — Baltimore et Fougéria *Manch meth.* 592 et suppl. 243. *Cass. dict.* 17. p. 283. 46. p. 399 et 412. — Baltimore *Linn. mant.* 288. *gen. n.* 1323. *Gartn. fr.* 2. p. 443. t. 169. *Lam. ill.* t. 709. — Niebuhria *Scop. non. DC.*

Capitulum pluriflorum monoicum, fl. radii 5 ligulatis fœmineis, disci masculis tubulosis 5-dent. Invol. 2-serialis squamæ adpressæ lanceolatæ, ext. foliaceæ. Recept. planum, paleis amplectentibus membranaceis apice ciliato-fimbriatis. Cor. disci extus hirsutiusculæ. Achænia radii juniora subtrigona obcompressa pappo coroniformi dentato deciduo (1), adulta trigona omnino calva lævia, disci abortiva elongata compressa papulosa. — Herbae annuæ erectæ Americanæ. Caulis obtusè tetra-

(1) Quo pappi delapsu plurimi decepti fuerunt, et præ aliis Mœnchius qui plantam adhuc pappiferam sub nom. Baltimore et pappo dejecto sub nom. Fougéria descripsit.

gonus subscaber. Folia opposita petiolata ovata acuta serrata trinervia subscabrida. Capitula pedicellata è dichotomiis orta aut rarissimè in iisdem indiv. oppositifolia. Flores pallidè flavi aut albi? — Nomen *Baltimoræ* rejiciendum! cum planta ad *Baltimoram* non erescat, admisi cum mutatione parçâ etymologiæ conformi nomen *Mœnchii* in honorem cl. Fougereux de Bondaroy Gaillardæ monographi botan. et agronomi periti sæc. XVIII.

1. *F. RECTA*, foliis pubescentibus serratis, pedicellis elongatis. (1) in Mexico (*Less.*) fortè circa *Veram-Crucem* nec certè circa *Baltimoram* (*Pursh. Nutt.*). *Baltimora recta* *Linn.* mant. 288 excl. *Pluk. syn.* et idèd patriâ. *Schkuhr. handb.* 3. s. 261 b. opt. *Baltimora trinervata* *Mœnch meth.* 592, et *Fougeria tetragona* *Mœnch suppl.* 243. *Moris. oxon.* 5. 7. t. 13. f. 16. (v. v. cult.)
2. *F. ALBA*, foliis subtomentosis subserratis, capitulis subsessilibus. — olim in horto *Trianonensi* culta ex specim. herb. *Rich.* sub nom. *Milleriæ albae*. *Baltimora alba* *Pers. ench.* 2. p. 489.

CCLXXVIII. *CHRYSOGONUM* *Linn. gen. n.* 988. *Gærtn. fr.* 2. p. 436. t. 174. *Lam. ill.* t. 713. *Less. syn.* 220. — *Chrysogonum* et *Diotostephus* *Cass.*

Capitulum multiflorum monoicum, fl. radii circ. 5 uniseriatis ligulatis fœmineis, disci masculis tubulosis 5-dentatis obtusis. Recept. planum, squamis linearibus obtusis. Cor. disci glabrinseulo, tubo brevissimo, fauce cylindricâ. Achænia radii obcompressa vix subtrigona glabra pappo coroniformi persistenti coronata, disci abortiva linearia elongata pappo coroniformi dentato. — Herbæ perennes Boreali-Americanae plus minus hirsutæ humiles. Caules è collo plurimi, alii floridi, alii stoloniformes. Folia alia radicalia, alia secus caules opposita petiolata ovata crenata. Capitula solitaria breviter pedicellata. Flores lutei.

1. *C. VIRGINIANUM* (*Linn. sp.* 1303), petiolis limbo longioribus, foliis demùm glabris, pappo radii coroniformi ad latus interius fissio apice crenato. 2/ in Virginia? (*Linn.*), in Carolina superiore (*B. Greene!*). *Gærtn. fr.* 2. p. 473. t. 174. f. 8. *Pluk. alm.* 100. t. 83. f. 4 (excl. altero *Pluk. syn.* à *Linn.* et *Willd.* citato). *Lam. ill.* t. 713. f. B. (v. s.)
2. *C. DIOTOSTEPHUS*, hirsutissimum, petiolis limbo subbrevioribus, pappo radii biaruito. 2/ in siccis pinguibus Carolinae. *Diotostephus repens* *Cass. dict.* 48. p. 543. *Chrys. Virginianum* *Michx. amer.* 2. p. 472. *Ell. sketch* 2. p. 472. — *Lam. ill.* t. 713. f. A. Achænia radii ante squamas invol. interiores sita et cum floribus disci circit. 4. 5. à recept. cum squamâ ipsâ separata. (v. s.)

+ *Species vix nota.*

C. DICHOTOMUM (*Vahl in Vest St.-Cr.* p. 303. sine descr.). — in insulâ Sanctæ-Crucis.

++ *Species exclusa.*

C. PERUVIANUM *Linn.* = *Zinnia pauciflora*.

Div. III. *SILPHIÆ.* — Asteroides *Melampodiæ* *Silphiæ* *Less. syn.* 213.

Capitula monoica multiflora, fl. fœmineis pluribus (3-10) ligulatis. Achænia eocorticata calva aut biaristata. Recept. paleaceum, paleis squamisve nunquàm cum achæniis concretis.

CCLXXIX. *GUARDIOLA* *H. Bonpl. pl. eq. 1. p. 144. t. 41. H. B. et Kunth nov. gen. am. 4. p. 247. Cass. dict. 20. p. 12. 59. p. 319. Less. syn. p. 419. — Guardi-
ola Steud. nom. 304.*

Capitulum pluriflorum heterogamum, fl. radii 3-5 ligulatis fœmineis, disci 10-11 masculis tubulosis 5-dentatis. Invol. 3-phyllum, squamis ovalibus erectis. Recept. planum angustum paleaceum. Ligularum tubus longus basi hispidus, cor. radii tubus longior glaber. Achænia disci compressa (Kunth) seu obcompressa (Cass.) lævia calva, disci abortiva. — Herba Mexicana glabra trichotoma. Folia opposita longè petiolata ovato-lanceolata acuminata serrata basi subcuneata. Capitula terna pedicellata corymbosa terminalia. Discus flavus, antheris nigri-
cantibus, radius albus brevis 3-dentatus. — Certè *Helianthea* sensu Cassiniano. An *Milleriis*, an *Silphicis* adjungenda?

1. *G. MEXICANA* (H. et B. l. c. t. 41). (1) in aridis Mexici iuter Ario-Aguasarco et montem Jorullo.

CCLXXX. *HIDALGOA* *Less. in linnæa 1831. p. 406. syn. 213. non Lallav.*

Capitulum monoicum multiflorum, fl. radii 5 uniseriatis fœmineis ligulatis, radii tubulosis abortu masculis. Inv. cylindræo-campanulatum. biserialè disco brevius, squamis integerrimis obtusis. Recept. planum, paleis membranaceis linearibus. Stylus disci simplicissimus clavatus, radii bifidus ramis semi-lanceolatis. Achænia disci lineare, radii valde compressum læve ellipticum angustè bialatum apice continuatum in pappum brevissimum bicornem. — Herba scandens Mexicana. Folia alterna petiolata pedati- seu ternati-secta, segmentis lateralibus plus minus bilobis. Capitula solitaria longè pedunculata aurantiaca.

1. *H. LESSINGII*. — in Mexico temperato, in sylvis Misantlæ, in Cuesta-Grande de Yala-Cingo. *H. ternata* Less. l. c. non Lallav. Ex descr. auctorum differt à stirpe *Lallavei* (mihi ut Sprengelio *Melampodio* adjungenda), genericè fructu nec drupaceo nec apice cuncillato; specificè foliis alternis nec oppositis, pedati- nec strictè ternati-sectis, segmentis lateralibus bilobis.

+ *Species exclusa.*

H. TERNATA Lallav. = *Melampodium Hidalgo*.

CCLXXXI. *SILPHIUM* *Linn. gen. n. 1334. Cass. dict. 59. p. 319. et 34. Less. syn. 213.*

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis, ligulis 1-serialibus, fructibus pluriserialibus, fl. disci tubulosis 5-dentatis abortu styli masculis. Invol. campanulatum imbricatum, squamis basi adpressis strictis apice foliaceis laxis. Recept. subconvexum paleaceum. Cor. disci tubo brevissimo, fauce cylindræa lobis, superne puberulo-hirtis. Stylus indivisus apice hispidus. Achænia radii plano-obcompressa bialata emarginata dentibus aristisve 2 nunc liberis nunc cum alâ utrinquè concretis coronata. — Herbæ Boreali-Americane sæpius elatæ interdum gummiifere, caule tereti aut tetragono, foliis alternis verticillatis aut oppositis sæpè connatis. Capitula pedicellata. Flores lutei. — *Silphium* veterum erat etiam herba gummiifera inter Umbelliferas quærenda.

§. 1. *Tereticaulia, caulibus teretibus seu obtusè subhexagonis, foliis alternis aut ternato-verticillatis, achæniis apice biaristatis aut bidentatis.*

* *Foliis pinnatilobatis alternis.*

1. *S. LACINIATUM* (Linn. sp. 1301), caule tereti supernè hispido, foliis alternis petiolatis pinnatipartitis scabris amplexicaulibus, lobis acutis indivisis aut dentato-sinuatis, invol. squamis hispidulis basi ciliatis latè ovatis in appendicem lanceolatam acuminatam productis. \mathcal{L} ad ripas flum. Amer. bor. præsertim ad Mississipi (Pursh), in Georgia occident. et ad montes Alleghany (Ell.), Linn. f. dec. p. 5. t. 3. Ell. sketch 2. p. 46r. Jacq. f. ecl. 1. t. 90. *S. spicatum* Poir. suppl. 5. p. 157. Herba 1-2-orgyalis. Capitulum amplum. Achænia radii pluralia, aristis 2 cum alâ confluentibus. (v. v. c.)
2. *S. GUMMIFERUM* (Ell. sketch 2. p. 46o. in adn.), caule valdè hispido et scabro, foliis sinuato-pinnatifidis subtus subhispidis, lobis acutis incisus, invol. squamis ovatis acuminatis margine hispidis. \mathcal{L} in pratis Alabamæ prope Georgiam occid. Herba 2-3-pedalis valdè *S. laciniato* affinis. Caulis gummi theribinthaceum copiosum fundit. Capitula majora quàm in omnibus aliis speciebus. Aristæ subulatæ.
3. *S. PINNATIFIDUM* (Ell. sketch 2. p. 46a), caule etiam ad apicem glabriusculo, foliis sinuato-pinnatifidis subscabris subtus parcè pilosis, invol. squamis glabris obtusis, ext. rotundis, int. ovalibus, ligulis invol. circ. æqualibus. \mathcal{L} in pratis Alabamæ et Georgiae occid. Herba 4-6-pedalis. Fol. lobi sæpius acuti. An *S. pinnatifidum* Nees et. sem. h. Bonn. 1831 omninò idem?
4. *S. COMPOSITUM* (Michx. fl. bor. am. 2. p. 145), caule tereti lævi, foliis alternis petiolatis sinuato-pinnatifidis lobis obtusis secus petiolum et margines limbi vix ac ne vix pilosisculis, invol. squamis ovalibus obtusis glabris. \mathcal{L} in sylvis glareosis maritimis à Virginia ad Carolinam (Pursh), in pinetis siccis Carolinae merid. (Fras.! Bose!). *S. laciniatum* Walt. car. 217. *S. sinuatum* Banks herb. ex Pursh. *S. pseudo-laciniatum* Fras. in herb. L'Hér. Radix ex Fraser crassa venenata. Capitula parva paniculata. (v. s.)
5. *S. SUBACAULE* (Nutt. in Sillim. am. journ. 5. 1822. p. 301.), scabriusculum, caule brevissimo, pedunculis longissimis, foliis alternis sessilibus subsinuato-pinnatifidis dentatis, invol. squamis dilatatis rhomboideis.—in Florida orientali. Caulis vix 2-3-poll. Folia 2-3 poll. longa exactè ut in Senecione alternè sinuata. Pedunc. palmares.

** *Foliis indivisis alternis.*

6. *S. TEREBINTHACEUM* (Linn. suppl. 383), caule tereti striato glabro, foliis alternis, infer. petiolatis ovatis subcordatis serratis scabridis, sup. ovatis, paniculâ laxâ nudiusculâ, invol. squamis glabris ovatis obtusis. \mathcal{L} in montibus Carolinae (Ell.) et Louisianae (Pursh). Jacq. hort. vind. 1. t. 43. Gärtn. fr. 2. p. 445. t. 171. Herba 4-5-pedalis ferè tota glabra. Denticuli 2 ad apicem achæniæ breves ab alâ disjuncti. Variat, ex Ell. sketch 2. p. 463, foliis radicalibus reniformibus subtus scaberrimis et foliis lobato-angulatis subtus ferè glabris. Prima var. est verisimil. *S. reniforme* Raf. med. fl. 2. p. 283 et ultima fortè *S. elatum* Pursh fl. bor. am. 2. p. 579 ex h. Banks in Carolina indicatum.
7. *S. PUMILUM* (Michx. fl. bor. am. 2. p. 146), caule adscendente tereti tomentoso, foliis alternis petiolatis cordato-ovatis dentatis supra pubescentibus subtus petiolisque subtomentosis, capitulis paucis longè pedicellatis, invol. squamis latè ovatis obtusis velutinis. \mathcal{L} in Florida (Michx.).

β. *tomentosum* (Nutt. gen. am. 2. p. 183), caule petiolis folisque subtus densè albo-tomentosis. \mathcal{L} in Georgia (Bartr.), pinetis siccis Carolinae (Fras.! Ell.). *S. tomentosum* Pursh fl. 2. p. 579. *S. pumilum* Ell. sketch 2. p. 469. Achænia adpressè villosa, vix alata, juniora setas 2 deciduas gerentia demum mutica apice hispidula. (v. s.)

8. *S. ASTERISCUS* (Linn. sp. 1302), caule tereti hispido, foliis inf. oppositis

grossè serratis, super. alternis sessilibus oblongis seu ovali-lanceolatis hirtis serratis integrisve, capitulis paucis solitariisve, invol. squamis ovatis ciliatis, exter. acutis. \mathcal{L} in pratis montanis (Pursh) et arenosis siccis (Ell.). Virginie et Carolinæ. Dill. elth. 42. t. 37. f. 42. S. reticulatum Mœnch. Achænia radii latè alata apice callis à obtusis ab alâ disjunctis coronata. (v. s. sp. et v. c.)

β . *lævicaule*, caule lævi, invol. squamis dorso glabris margine ciliatis apice subobtusis, achæniis dentibus paulò magis conspicuis. \mathcal{L} in Georgiâ occident. Silph. Asteriscus β Nutt. gen. 2. p. 183. \bar{S} . scabrum Walt. car. 217. S. dentatum Ell. sketch 2. p. 468. (v. s.)

9. S. QUERCIFOLIUM, caule tereti hispido densè folioso, foliis sparsis petiolatis elliptico-sublanceolatis obtusiusculis obtusè sinuato-repandis penninerviis supra strigilloso-scaberrimis subtùs puberis, petiolis ciliatis, capitulis corymbosis, invol. squamis ovali-oblongis obtusis glabris subciliatis apice patentis-squarrosis. \mathcal{L} patr. ign. Vidi vivum anno 1807, in horto Burdigalensi nec unquam in hortis inveni. Differt à S. Asterico foliis petiolatis; à S. trifoliato et ternato caule hispido et foliis alternis; ab omnibus foliis repando-sinuatis.

10. S. RETICULATUM (Pursh fl. bor. am. 2. p. 579, caule...., foliis alternis ovato-lanceolatis serratis obtusiusculis villosiusculis. \mathcal{L} in Floridâ. Cæt. ign. Species Nuttallio dubia. †.

*** *Folii indivisi verticillatis.*

11. S. TRIFOLIATUM (Linn. spec. 1302), caule tereti subhexagono lævi, foliis ternato-verticillatis ovatis dentatis scabris, infer. longè, super. brevissimè petiolatis, petiolis subnudis, paniculâ 3-chotomâ, invol. squamis ovali-oblongis subacutis vix ciliatis, ligulis 15-18. \mathcal{L} in montosis Virginie, Carolinæ (Pursh) et Georgiæ (Ell. sketch 2. p. 466). S. ternatum Desf. cat. h. Par. ed. 3. p. 403 non Retz. — Moris. ox. s. 6. t. 3. f. 68. Achænia glabra latè alata apice emarginata dente calloso intra marginaturam notata. Habui sub nom. admissio ex hort. Hafn. et ex Lugd. Bat. tempore Royeni collectum. (v. v. c.)

12. S. TERNATUM (Retz in Willd. sp. 3. p. 2333), caule tereti lævi, foliis ternato-verticillatis breviter petiolatis lanceolatis acuminatis serratis scabris, petiolis ciliatis, paniculâ subdichotomâ, invol. squamis ovali-oblongis obtusiusculis ciliatis, ligulis 8-9. \mathcal{L} in Amer. bor. (Retz.), montanis Carolinæ et Georgiæ (Ell. sketch 2. p. 467). S. ternifolium Michx. fl. bor. am. 2. p. 146. S. trifoliatum Desf. hort. Par. ed. 3. p. 182. Hook. bot. mag. t. 3355. non Retz. Linnæus Michaux et plures alii S. ternatum et trifoliatum verisim. confuderunt. S. ternatum præter alias notas differt achænio apice subciliato, aristas 2 obtusas cum aliis concretas gerente. (v. v. c.)

β . *atropurpureum* (Nutt. gen. am. 2. p. 183), caule tereti lævi atropurpureo, foliis quaterno-verticillatis sparsisve lanceolatis scabris, petiolis ciliatis. \mathcal{L} in Amer. bor. Carolinâ et Georgiâ (Pursh fl. 2. p. 575). \bar{S} . atropurpureum Retz in Willd. sp. 3. p. 2334.

- §. 2. Tetragona, caule tetragono apice interdum teretiusculo, foliis oppositis, superioribus sæpè connatis, achæniis exaristatis, aristis nempe cum alâ utrinquè omninò concretis.

13. S. PERFOLIATUM (Linn. sp. 1301), caule tetragono lævi, foliis oppositis ovato-deltoides grossè serratis glabriusculis, inf. petiolatis, sup. connato-perfoliatis, invol. squamis latè ovatis obtusis tenuè ciliatis, ext. subsquarrosis, ligulis 15-20. \mathcal{L} in Amer. bor. à Pennsylvania ad Carolinam in montibus Alleghany (Pursh). Cass. dict. 59. p. 324. Hook. bot. mag. t. 3354. S. tetragonum Mœnch. Achænia glaberrima latè alata emarginata exaristata, lobis obtusis, aristis nempe cum alâ omninò et strictissimè sub nervi marginalis specie concretis. (v. v. c.)

β . *conjunctum*, invol. squamis ext. 4 expansis et cætera superantibus. S. conjunctum Willd. enum. p. 633 ex Pursh.

14. *S. HORNEMANNI* (Schrad. hort. Goett. sem. 1829 et in litt. 1832), caule tetragono infernè hispido, foliis oppositis subtus subscabris latè ovatis acutis grossè serratis, infer. petiolatis petiolis sæpius alatis, superioribus connatis-perfoliatis, invol. squamis ovatis subobtusis tenuè ciliatis, ligulis circ. 20. $\frac{1}{2}$ in Amer. bor. *S. Asteriscus* Hornem. h. Hafn. ex Schrad. *S. connatum* Desf. hort. Par. cat. ed. 3. p. 182 non Linn. *S. perfoliatum* Poir. suppl. 5. p. 157? *Achænia* glaberrima alata apice acutè biloba, more *S. perfoliati* exaristata. Species media inter *S. perfoliatum* et *S. connatum* et cum utroque sæpè confusa. (v. s. comm. à cl Schrad. et v. c.)
15. *S. CONNATUM* (Linn. mant. 574), caule basi tetragono apice tereti hispidissimo, foliis oppositis latè connatis-perfoliatis ovato-oblongis serratis scabris, invol. squamis ovatis obtusiusculis subsquarrosis, ligulis circ. 12. $\frac{1}{2}$ in Amer. boreali. *S. scabrum* Moench. Ligulæ 7-8-nerviæ latiores quàm *S. perfoliati*. *Achænia* non vidi. (v. s. cult. in hort. Hafn.)
16. *S. SCABERRIMUM* (Ell. sketch 2. p. 466), caule subangulato, angulis superius scabris, foliis breviter petiolatis interdum connatis ovatis acuminatis serratis rigidis utrinquè scaberrimis, capitulis subcorymbosis, invol. squamis ovatis ciliatis, ligulis 12-14. $\frac{1}{2}$ in Amer. bor. Georgiâ. *Achænia* orbiculata alata profundè emarginata.
17. *S. INTEGRIFOLIUM* (Michx. fl. bor. am. 2. p. 146), caule tetragono aspero, foliis oppositis sessilibus oblongis integerrimis scabris, capitulis paucis breviter pedunculatis, invol. squamis oblongo-ovatis glabris ciliatis, ligulis circ. 14. $\frac{1}{2}$ in regione Illinoensium (Michx.), ad montes Alleghany (Pursh), in montanis Carolinæ (Ell. sketch 2. p. 465.)
18. *S. LEVIGATUM* (Pursh fl. bor. am. 2. p. 578), caule tetragono sulcato glabro simplici, foliis oppositis sessilibus ovatis acuminatis tenuissimè serratis basi subcordatis utrinquè glabris, capitulis densè corymbosis, invol. squamis ovatis ciliatis. $\frac{1}{2}$ in Georgiâ inter Savannah et Louis-Ville (Ensl.) et in distr. occident. (Ell. sketch 2. p. 466.)
19. *S. ERYTHROCAULON* (Bernh. ex Spreng. syst. 3. p. 630), caule tetragono glabro, foliis oppositis cordato-lanceolatis acutis inæqualiter dentatis scaberrimis, petiolis alatis perfoliatis, invol. squamis oblongis acutiusculis. $\frac{1}{2}$ in America boreali.

† *Species vix nota.*

S. TRACHOPUS (Raf. herb. p. 24). — In America boreali.

†† *Species exclusæ.*

S. SOLIDAGINOIDES Linn. = *Heliopsis lævis*.

S. TETRAGONOTHECA Gærtn. = *Tetragonotheca helianthoides*.

S. TRILOBATUM Linn. = *Wedelia carnosa*.

CCLXXXII. *POLYMNIA* Linn. gen. n. 987. Gærtn. fr. 2. p. 429. t. 174. f. 2. Less. syn. 213. — *Polymnia* et *Alymnia* Neck. elem. 1. p. 31. — *Polymnia* et *Polymniastrum* Lam. ill. t. 711 et 712. Cass. dict. 59. p. 246 et 247.

Capitulum multiflorum monoicum, fl. radii foemineis ligulatis 1-seriatis, disci tubulosi 5-dentati abortu styli masculis. Invol. duplex, ext. squamis 5 patulis magnis foliaceis ovato-lanceolatis, int. squamis pluribus brevioribus (sed disco longioribus) adpressis achænia amplectentibus. Recept. planum paleaceum. Stylus disci supernè pubescens apice bifidus. Achænia glabra, disci abortiva teretia, radii obovata crassa subcompressa lævia exalata calva. — Herbæ Americanæ erectæ. Folia alterna seu opposita aspera semiamplexicaulia subcorymbosa. Flores lutei, disco sæpè fusco-purpurascens.

Sect. I. UVEDALIA. — Polymnia Neck. Lam. Cass. II. cc.

Ligulæ planæ elongatæ involucrium superantes.

1. P. UVEDALIA (Linn. spec. 1303), caule sulcato glabriusculo, foliis oppositis utrinque sparse puberis sinuato-lobatis subtrilobis in petiolum decurrentibus, ligulis 10 lineari-lanceolatis tridentatis. \mathcal{L} in pascuis montanis Amer. bor. à Virginîâ ad Carolinam. Michx. fl. 2. p. 147. Ell. sketch 2. p. 471. Mill. dict. n. 1. Lam. ill. t. 711. f. 2. bona. Willd. sp. 3. p. 2335. excl. syn. Cav. — Pluk. t. 83. f. 3. Moris. ox. s. 6. t. 7. f. 55. (v. s.)
2. P. MACULATA (Cav. ic. 3. p. 14. t. 227), caule sulcato plus minus villosa sæpius maculato, foliis oppositis junioribus molliter villosis, inf. deltoideo-concave sinuato-angulatis, supremis rhombeis subrepandis, ligulis circ. 20 lanceolatis tridentatis. invol. squamis exter. latè ovatis. \mathcal{L} in Mexico (Cav.), in valle Mexicanâ (Berl. pl. exs. n. 719), in montibus Aihualtampec (fl. Mexic. ined.). P. Uvedalia hort. Paris. 1818 et hort. Lugd. B. tempore Royenii. (v. v. c. et s. sp.)
3. P. ASPERA (Mart. herb.), caule teretiusculo scabro apice breviter hirsuto, foliis oppositis suprâ aspero-scabris subtus molliter velutinis subcanescentibus hastato-trifidis, lobis acuminatis suberoso-dentatis, lateralibus subbifidis ad basin acutè auriculatis, invol. squamis exter. lineari-oblongis disco duplò ferè longioribus. \mathcal{L} in Mexico. Enata anno 1829 in hort. Monac. seminibus ab hortul. Kerl. adportatis. Species aff. P. maculatæ sed folia basi amplè et acutè auriculata. Ex capitulis junioribus caract. non eruerè potui. An P. scabra Zuccar. pl. nov. 1. p. 313 eadem? Ex cl. auct. capitula dicuntur discoidea nec radiata! (v. s. in h. Mart.)
4. P. RIPARIA (H. B. et Kunth nov. gen. am. 4. p. 282), caule compresso fistuloso glabro, foliis oppositis ovatis sinuato-angulatis subcordatis basi in petiolum decurrentibus dentatis hispidulis, ligulis 12-15 elongatis 3-dentatis. \mathcal{L} ad ripam fl. Magdalenæ inter Cascajo et Angustura de Carare.
5. P. GLABRATA, caule tereti subangulato glabro ad apices scabriusculo, foliis oppositis subconnatis ovato-lanceolatis acuminatissimis basi in petiolum alatum decurrentibus subsinuato-dentatis seu integerrimis, adultis glabris, junioribus subasperulis, corymbo composito laxo, invol. squamis exterioribus latè ovalibus. \mathcal{L} in Chili legit cl. Hænke. Folia 9 poll. longa, 3 poll. lata. (v. s. in h. Hænke à cl. de Sternberg comm.)
 β . *angustifolia*, foliis oblongo-lanceolatis subintegerrimis. \mathcal{L} in Cordilleris Chilensium et fortè Peruvianorum legit Hænke. Folia 4 poll. longa, pollicem lata. (v. s. comm. cum α .)

Sect. II. ALYMNIA. — Alymnia Neck. elem. 1. p. 31. — Polymniastrum Lam. ill. t. 712.

Ligulæ involucrio breviores obovatæ aut lineares.

6. P. CANADENSIS (Linn. sp. 1303), caule tereti puberulo-glanduloso, foliis oppositis alternisve ovato-triangularibus sublobatis acutis petiolatis sæpè basi auriculatis, infimis sublyratis, invol. squamis ovatis acuminatis ciliatis, ext. paulò majoribus, achæniis obovatis subcompressis, ligulis minutissimis trifidis. \mathcal{L} in montanis Americæ borealis à Canadâ ad Virginiam (Pursh), in Carolinâ (Ell. sketch 2. p. 471), in Kentucky (Raf.!), ad saltus Niagaræ (B. Greene!), Linn. amæn. ac. 5. p. 15. t. 1. f. 5. Lam. ill. t. 711. f. 1. Michx. fl. bor. am. 2. p. 147. (v. s.)
7. P. VARIABILIS (Poir. dict. 5. p. 505), caule tereti puberulo-glanduloso, foliis oppositis ovato-triangularibus sublobatis in petiolum cuneato-decurrentibus subpuberis, invol. squamis subæqualibus subacutis dorso puberis eciliatis, ligulis minutis subintegris. — patr. ign. Polymniastrum Lam. ill. t. 712. Capitulum

primâ fronte discoideum sed verè radiatum ligulis minutissimis. An à priori certè distincta? (v. s. c. in hort. Par.)

8. *P. SIBIENSIS* DC., caule tereti trichotomo apice puberulo, foliis oppositis sessilibus ovato-deltoides basi latè cuneatis apice acuminatis subdentatis indivisis suprà glabriusculis subtùs pubescenti-villosis, summis lanceolatis subintegerrimis, invol. squamis ext. 5 oblongo-linearibus subpuberulis, int. summo dorso pilos apice glandulosos gerentibus, ligulis parvis subintegris. — in Brasiliâ circa Rio-Janeiro legit cl. Gaudichaud. Species distinctissima et primâ fronte à genere diversa, sed à prioribus non separanda. Stylus disci minimus subnullus. Recept. post. anthesin elevatum supernè plano-subconcauum et Berlandieræ recept. ferè referens (v. s. comm. à cl. inv.)
9. *P. SILPHIOIDES*, caule tereti puberulo-glanduloso, foliis oppositis alternisve membranaceis puberulis triplinerviis grossè dentatis apice subtrilobis, basi in petiolum alatum irregulariter dentatum attenuatis, prope basin auriculato-dilatatis, inv. squamis ext. ovatis acutis dorso villosis, achæniis obovatis subcompressis, ligulis linearibus integris. — in Brasiliâ prov. Rio-Grande (h. Mus. imp. Bras. n. 885!). Affinis priori sed folia basi dilatata, capitula duplò majora.

† *Species non satis nota aut dubia.*

10. *P. CAROLINIANA* (Poir. dict. 5. p. 505), caule tereti pubescente, foliis alternis indivisis crenatis subtùs albidis, invol. squamis ext. majoribus, int. 4 minoribus. — in Carolinâ. †.

†† *Species exclusæ.*

- P. ABYSSINICA* Linn. = Guizotia oleifera.
P. CARNOSA Linn. = Didelta carnosa.
P. FRONDOSA Bruc. = Guizotia oleifera.
P. PERFOLIATA Poir. = Melampodium perfoliatum.
P. REVOLUTA Quorumd. ex Steud. = Wedelia perfoliata.
P. SPINOSA Linn. = Didelta spinosa.
P. TETRAGONOTHECA Linn. = Tetragonotheca helianthoides.
P. WEDELIA Linn. = Wedelia frutescens.

CCLXXXIII. *ESPELETIA* Mutis in Humb. et Bonpl. pl. equin. 2. p. 11. t. 70-72.
 H. B. et Kunth nov. gen. am. 4. p. 280.

Capitulum multiflorum monoicum, fl. radii numerosissimis 1-seriatis ligulatis fœmineis, disci tubulosi 5-dentati bisexualibus abortu masculis. Invol. campanulatum imbricatum polyphyllum. Recept. planiusculum paleaceum, paleis membranaceis obtusis. Antheræ exsertæ. Stylus radii bifidus gracilis, disci simplex. Achænia obovata subangulata calva, disci verisim. abortiva. — Herbæ æquinoct. Americanæ densè lanato-tomentosæ resinosæ perennes. Folia opposita integra. Capitula ramos terminantia corymbosa lutea. — Genus caract. videtur affine Silphio sed habitus diversus. Antheræ ex icone breviter caudatæ, sed de hoc caract. cum habitu discrepante silent descriptiones.

1. *E. GRANDIFLORA* (H. et Bonpl. l. c. t. 70), rufescenti-lanata, foliis radicalibus lanceolatis subtùs lineatis, corymbo simplici laxo. ♀ in Amer. austr. locis subfrigidis prope Santa-Fé de Bogota et in Andibus Quinduensium. A Bogotensibus dicitur *Fraxylejon*. Fundit resinam *trementhina* dictam. Achænia omninò glabra. Ligulæ circ. 60 lineares apice 3-dentatæ.
2. *E. ARGENTEA* (H. et Bonpl. l. c. t. 71), caule densissimè argenteo-piloso, foliis suprà strigosis subtùs molliter tomentosis, corymbo ramoso subfastigiato. ♀ in Novæ-Granatæ frigidis juxta urbem Zypaquira. Capitula dicuntur radiata, sed ligulæ indescriptæ nec in icone obviæ. Stylus radii ad

basin more insolito moniliformi-incrassatus. Achænia omnino glabra. Cor. ad tubi apicem pilosiusculæ.

3. *E. corymbosa* (H. et Bonpl. l. c. p. 16. t. 72), incano-tomentosa, foliis coriaceis, radicalibus supernè lato-lanceolatis subtùs venosis tomentosis, capitulis subcorymbosis confertis. 2/ in frigidis Andiam Novo-Granatensium juxta Almaguer. Èsp. rigida H. et B. pl. equin. t. 72. Ligulæ circ. 20 invol. paulò longiores. Cor. in tubo glabræ. Achæniû apice barbatum.

CCLXXXIV. BERLANDIERA.

Capitulum multiflorum monoicum, fl. radii 1-serialibus ligulatis foemineis ad axillas squamarum invol. interiorum sitis, fl. disci masculis tubulosis 5-dentatis ad axillas palcarum recept. exteriorum sitis, centralibus abortivis. Recept. turbinatum obconicum supernè truncatum. Invol. 3-seriale foliaceum, squamis ext. circ. 3 oblongis, mediis obovatis majoribus acutiusculis circ. 4-5, intimis maximis subrhombeis subdentatis circ. 6-8. Paleæ recept. foliaceæ, exteriores binæ floriferæ ad basin cujusque squamæ recept. intimæ ortæ cucullatæ apice dilatata obtusæ, aliæ ex margine et ex disco terminali recept. ortæ minores cuculliformes florem ad filum reductum aut nullum gerentes. Achænia radii obcompressa obovato-orbiculata exalata coriacea pappo brevissimo subaristato coronata. Cotyledones subcarnosæ invol. squamis parallelæ. — Suffrutex Texano-Mexicanus. Rami teretes hirsuti. Folia alterna sessilia cordata ovato-sublanceolata crenata, suprâ strigilloso-villosa, subtùs canescenti-villosa. Capitula longè pedicellata ad apices ramorum 3-5 corymbosa. Invol. et recept. paleæ pubescentes. Flores lutei. — Aff. videtur hinc Melampodio, illinc Polymniæ. — Genus dixi in honorem cl. Berlandier qui Mexicanas plagas peragravit et plantas plurimas his regionibus proprias collegit.

1. *B. TEXANA*. 5 in sylvis Mexicanæ provinciæ Texas legit cl. Berlandier. (Pl. exs. n. 2044.) (v. s.)

Div. IV. MELAMPODIEÆ. — Asteroideæ Melampodieæ veræ. Less. syn. 217.

Capitula monoica multiflora, fl. foemineis pluribus ligulatis. Achænia corticata (1) calva. Recept. paleaceum, paleis plùs minùs cum achæniis concretis. — Cassini hanc et priorem divisionem sub nom. Helianthæ Milleriæ Melampodieæ junxit. (Opusc. 3. p. 60.)

- CCLXXXV. MELAMPODIUM Linn. gen. n. 989. Brown trans. hort. soc. 12. p. 104. H. B. et Kunth nov. gen. am. 4. p. 272. Less. syn. 217. — Melampodium, Dysodium et Alcina Pers. — Melampodium, Zarabella et Alcina Cass.

Capitulum multiflorum monoicum, fl. radii 5-10 uniseriatis foemineis ligulatis, disci tubulosis masculis. Invol. duplex, squamis ext. 5, rariùs 3, planis patulis, int. tot quot flores radii et eos involventibus. Recept. convexo- aut subulato-conicum paleaceum, paleis diaphanis deciduis. Achænia disci abortiva, radii obovata subcurvata lævia calva intra squamas inv. interiores plùs minùs inclusa et ideò sæpiùs

(1) *Corticata* dixi achænia squamis inv. paleisve recept. involuta et cum iis adeò concreta ut massa unica ex achænio et squamâ coadunatis evadat. Sed *involuta* dixi cum squamis paleisve sunt inclusa nec concreta. Ergò achænia Chrysogoni et plur. aliorum generum à cl. Lessingio corticata dicta hoc nomine non designavi.

externè tuberculosa aut rugosa et apice sæpè coronata. — Herbæ aut suffruticuli Americani. Caules dichotomi. Folia opposita. Pedicelli inter dichotomias orti 1-cephali. Flores flavi, ligulis in unicâ spec. albis.

Sect. 1. *EUMELAMPODIUM*. — *Melampodium* Linn. l. c. *Gærtn. fr. 2. p. 436. t. 169. Lag. nov. gen. et sp. p. 32. Cass. dict. 59. p. 237.*

Invol. squamæ interiores achænia involventes, extûs tuberculose seu muricatæ nervoso-costatæ, supernè productæ in cucullum nunc truncatum nunc in aristam uncinatam desinentem.

1. *M. DIFFUSUM* (Cass. dict. 59 (1829). p. 238), caule erecto-diffuso di- seu tri-chotomo, foliis oblongis utrinquè acutis integerrimis suprâ pubero-scabris subtûs longè sericeo-villosis, pedunculis è dichotomiis ortis 1-cephalis, invol. squamis 5, ext. obovato-cuneatis mucronatis achænia involv., basi nervoso-striatis seriem tuberculorum ad latus internum gerentibus, cucullatis in acumen uncinatum productis. (1) in insulâ Luzoniâ ad Manillam (Cham!), *M. Manillense* Less. in linnæâ 1830. p. 155. t. 2. f. G. (Achænium). Affine *M. Americano*. Discus oblongus. Folia ex Less. interdum variant pinnatifida. (v. s.)
2. *M. SERICEUM* (Lag. elench. h. madr. 1805. nov. gen. (1816) p. 32. n. 404 non Kunth), caule erecto diffuso di-trichotomo, foliis suprâ scabridis subtûs sericeo-villosis, infer. trifidis, sup. integris lineari-lanceolatis, capitulis in dichotomiis et apicibus breviter pedicellatis, inv. squamis ext. oblongis villosissimis, int. achænia involv. undiquè muricatis apice cucullatis in acumen extûs uncinatum productis. (1) in Mexico (Lag.) prov. Leoninæ ad occid. urbis Guanaxuato (Mendès). *M. diffuso* valdè affinis, sed pedicellis vix invol. longioribus bracteis foliaceis binis angustis sub involuero et achæniis valdè muricatis differt. (v. s.)
3. *M. RAMOSISSIMUM*, caule suffruticuloso ramosissimo glabro, foliis linearibus subadpressè pubescentibus, aliis integerrimis, aliis hinc indè obtusè dentato-lobatis sinuatisve, pedicellis folio longioribus 1-cephalis, invol. squamis ext. ovalibus extûs pubescentibus, inter. achænia involventibus basi tuberculosi apice expansi in cucullum latum dorso in acumen uncinatum desinentem. 5 in Mexici prov. Texas legit cl. Berlandier (pl. exs. n. 2017). Ligulæ oblongo-lineares parvæ. (v. s.)
4. *M. LINEARILOBUM*, caule erecto diffusè ramoso, foliis pinnatipartitis lobis utrinquè 2-4 linearibus suprâ strigoso-pilosis subtûs sericeo-villosis, summis paucis integerrimis linearibus, pedicellis elongatis 1-cephalis, invol. squamis exter. latè ovalibus basi villosissimis apice membranâ marginatis inter. achænia involv. basi rugoso-tuberculatis apice cucullatis, cucullo margine intûs revolutò dorso brevissimè apiculato. (1) in Mexico ad Guilo-tepec et Las Bacas. Ligulæ latè obovatæ. Cucullus interdum exapiculatus interdum apiculo recurvo. Facilè lobis fol. linearibus ab icone Houston. typo *M. Americano* differt. (v. s. comm. à cl. Alaman.)
5. *M. AMERICANUM* (Linn. sp. 1303), caule procumbente, foliis integris seu in lobos lato-lanceolatos utrinquè 2-3-fidis suprâ strigoso-hispidis subtûs junioribus sericeo-hirtis, pedicellis elongatis hispidis 1-cephalis, ligulis 10 oblongis invol. duplò longioribus integris, inv. squamis ext. latè ovatis, inter. achænia obtegentibus costatis auriculato-tuberculatis, apice cucullatis, cucullo extûs aspero marginibus inflexo exappendiculato. — in Mexico ad Vera-Cruz (Houst. reliq. p. 9. t. 21) et ad Santa-Fé in savanis (Less. in linnæâ 1830. p. 149). *Gærtn. fr. 2. p. 436. t. 169. f. 8.*
6. *M. CINEREUM*, caule diffuso villosò ramoso, foliis oblongo-linearibus obtusis utrinquè cinereo-pubescentibus paucis integris plerisque obtusè sinuato-repan-dis, pedicellis 1-cephalis folio paulò brevioribus, ligulis invol. longioribus apice emarginatis, invol. squamis ext. ovatis subacutis dorso villosis, inter. achænia invol. basi muricato-tuberculatis apice cucullatis, cucullo amplo ex-

aristato extus lævi, margine integerrimo intus revoluto. (1) in Mexico ad San-Fernando oct. fl. legit Berlandier (pl. exs. n. 2243). Flores ex eo albi, in sicco albid. (v. s.)

7. *M. KUNTHIANUM*, caule fruticoso, ramis junioribus villosa-pubescentibus, foliis lanceolato-linearibus integerrimis supra sericeo-pubescentibus, subtus lanuginoso-sericeis argenteis, pedicellis 1-cephalis folio longioribus, invol. squamis ext. obovato-cuneatis acutis extus sericeis, int. achænia involv. basi costatis muricato-tuberculatis apice cucullatis, cucullo vix apiculato marginibus vix inflexis. 3 in Mexico inter. ad Tehuillotepec et Tasco. Mel. sericeum H. B. et Kunth nov. gen. am. (1820), p. 272. t. 398 non Lag. Flores surantiaco-flavi; ligulæ invol. triplò longiores. Capitulum 8-9 lin. diam. Caules et pedunculi ex icone hispidi (v. s. sine fl.)
8. *M. LANCEOLATUM*, caule fruticoso basi glabro ad apices adpressè villosa, foliis lanceolatis acuminatis integerrimis supra parcè strigillosis demum glabris subtus sericeo-villosis argenteis, pedicellis 1-cephalis vix folio longioribus, invol. squamis ovatis dorso villosis, int. achænia involv. nervuloso-costatis et tuberculatis apice cucullo exaristato instructis. 3 in itiner. suis legit Née sed locus ignotus. An Manillense? an potiùs Americanum? Species facillè distincta. Capitula 4 lin. diam. Discus elevatus ut in Spilanthe. Fl. flavi. Ligulæ inv. duplò longiores ovales. (v. s. ex herb. Thib.)

Sect. II. *ZARABELLIA*. — *Dysodium* Rich. in Pers. ench. 2. p. 489 non Cav. — *Zarabellia* Cass. dict. 59. p. 240 non Neck.

Invol. squamæ inter. achænia radii involventes, rugosæ aut tuberculatæ, supernè aut truncatæ aut productæ in mucrones dentesve 1-3 antiocè subhiantes et achæni apicem denudatum monstrantes. — Sect. inter priorem et tertiam media, ad *Eumelampodium* M. heterophyllo, ad *Alcinam* M. divaricato ex habitu accedens.

9. *M. HETEROPHYLLUM* (Lag. nov. gen. et sp. p. 33. n. 405), caule fruticoso erecto piloso, foliis lanceolatis in iisdem aut diversis individuis indivisis trifidisque, supra puberulo-scabridis, subtus sericeo-villosis argenteis, pedicellis 1-cephalis folio longioribus, invol. squamis ext. ovato-rhombeis dorso villosis, int. achænia involv. nervuloso-subcostatis parcè tuberculatis in mucrones 1-3 minutos subproductis. 3 in Mexico (Lag.), in pratis Papantlæ (Schied.), ad colles prope Tantoyma et Tampico de Tamaulipas (Berl. pl. exs. n. 2161 et 2281). Less. in linnæa 1831. p. 407. Herba ferè M. sericei seu lanceolati sed cucullus ad apicem achæni nullus. (v. s.)
10. *M. ANGUSTIFOLIUM*, caule fruticoso erecto puberulo-hirtello, foliis linearibus integerrimis acutis supra puberulo-hirtellis subtus sericeo-villosis argenteis, pedicellis folio duplò longioribus, invol. squamis ext. ovatis acutis dorso villosis, inter. achænia involv. apice truncatis vix mucronatis. 3 in Peruviae montanis, ad Oronocum (Hænk. herb.). Hæc et præcedens species habitum sect. primæ optimè exhibent sed cucullus nullus. Fortè caract. paulò immutato ad *Eumelampodium* referendæ. (v. s. in h. Hænke à cl. Sternberg miss.)
11. *M. GRACILE* (Less. in linnæa 1831. p. 407), caule herbaceo erecto trichotomo, foliis semiamplexicaulis supra asperissimis junioribusque strigosohispidis subtus asperis pallidioribus et glabridisculis elliptico- seu obovato-rhombeis integerrimis seu hastato-panduratis posticè sinuato-dentatis, sup. triangularibus integerrimis acuminatis basi obtusissimis, invol. squamis ext. 3. ligulis 5, fructibus trapezoideo-obovatis apice nudis glabris lævibus obsolete reticulato-venosis. (1) in pratis Papantlæ Mexicanorum. Simile *M. paludoso* sed multò tenerius.
12. *M. OBLONGIFOLIUM*, caule herbaceo erecto di- et trichotomo sparse puberulo-hirtello, foliis sessilibus oblongis acuminatis integerrimis aut basi interdum subdilato-denticulatis utrinquè sparse et rariter strigillosis, pedicellis folio longioribus, invol. squamis ext. 3 lanceolatis, ligulis 5 bifidis, fructibus obo-

- vatis apice nudis dorso reticulato-rugulosis. (1) in Mexico ad Tantoyuca legit Berlandier (pl. exs. n. 2153). Capitula parva lutea. Affine *M. gracili*, sed folia nec rhombea nec formis varia nec nullâ basi obtusissima. (v. s.)
13. *M. MICROCEPHALUM* (Less. in *linnæa* 1834. p. 268), herbaceum gracile hirsutiusculum, foliis oblongo-rhomboideis basi perangustâ sessilibus, pedicellis folio longioribus, invol. squamis ext. 5 curvilineo-acuminatis, achæniis curvato-obpyramidalis apice truncatis et nudis dorso tuberculatis. (1) in Mexici cultis prope Hacienda de la Laguna. Proximè dicitur *M. divaricatum*.
14. *M. HISPIDUM* (H. B. et Kunth nov. gen. am. 4. p. 273 t. 399), herbaceum erectum, ramis oppositis hispidis, foliis sessilibus subconnatis linearis-lanceolatis integerrimis suprâ hispido-scarbis subtus pubescentibus, pedicellis folio duplò triplòve brevioribus, invol. squamis ext. 5 ellipticis obtusis dorso villosis, int. achænia involv. costatis spinuloso-tuberculatis apice truncatis, ligulis bifidis brevibus. (1) in Mexico inter Pascuaro et Valladolid de Mechoacan. Ligula flavidæ.
15. *M. LONGIFOLIUM* (Cav. ! hort. madr.), caule herbaceo erecto sparsè puberulo-hirtello, foliis oblongis subrhombeis semiamplexicaulibus obtusè cordato-auretis apice subacutis integerrimis utrinquè sparsè puberulis, pedicellis folio brevioribus, invol. squamis ext. 5 lanceolatis subacuminatis dorso puberis, int. achænia involv. compressis rugosis apice muticis aut dorso brevissimè mucronatis. (1) in Mexico (Cav.). Willd. enum. p. 934. Brouss. hort. monsp. 1806. (v. v. c.)
16. *M. RHOMBOIDEUM*, caule herbaceo erecto bifariam piloso, foliis ovato-oblongis basi aurito-semiamplexicaulibus à medio ad basin angustatis apice subacutis vix dentatis utrinquè puberulis, pedicellis è dichotomiis ortis folio brevioribus gracilibus, invol. squamis ext. 5 lanceolatis acuminatis, inter. achænia involv. rugosis dorso ad apicem in spinam erectam productis. (1) in Amer. calid. sed locus ign. Zarabellia rhomboidea Cass. dict. 59. p. 240. Dysodium divaricatum hort. par. 1819! An *M. Berterianum* Spreng. syst. 3. p. 619? sed inter plantas Berterianas non reperio. Affine *M. longifolio*, sed differt caulibus bifariam nec sparsè pilosis fructibusque longè spinosis. (v. v. c.)
17. *M. DIVARICATUM*, caule herbaceo dichotomo bifariam piloso, foliis ovatis acuminatis basi cuneatis grossè dentatis suprâ scarbis subtus hispidulis, pedicellis è dichotomiis ortis glabriusculis folio sæpius longioribus, invol. squamis ext. 5 obovatis obtusis basi subconcretis, int. achænia obtegentibus compressis rugosis apice truncatis extus subdentatis, ligulis brevibus 3-dentatis. (1) in Amer. paludosis ad sinum Darien (H. et B.), circa Sanctam-Martham (Rich. Bert.!), in Mexico (Lag. Karv.!). Dysodium divaricatum Rich. (1807), in Pers. ench. 2. p. 489. Colla in Ac. Taur. 1827. p. 323 cum icon. Melampodium paludosum H. B. et Kunth nov. gen. am. (1830) 4. p. 215. Less. in *linnæa* 1830. p. 149. Alcina ovatifolia Lag. nov. gen. et sp. (1816) n. 403. Jacq. f. ! eclog. 1. p. 115. t. 78. Melamp. ovatifolium Reich. exot. (1827) p. 30. t. 42, sed canlis undiquè pilosis! Wedelia ovatifolia Willd. enum. suppl. 61. Wedelia minor Horuem. Hafn. hort. p. 855. Alcina minor Cass. dict. 59. (1829) p. 243. Dysodium radiatum Desf. ! cat. h. par. 1830. p. 182. (v. v. c. et s. sp.)

Sect. III. *ALCINA*. — *Alcina Cav. ic. p. 10. t. 15. Cass. dict. 59. p. 242.* — *Wedeliæ sp. Willd.* — *Alcinia Steud.* — *Camutia Bonat.*

Invol. squamæ int. achænia radii involventes ovali-oblongæ læves apice tubercula minima 2-4 gerentes et inter eos foramine minimo vix apertæ. Achænium striatum.

18. *M. PERFOLIATUM* (H. B. et Kunth nov. gen. am. 4. p. 274), herbaceum erectum dichotomum glaberrimum, foliis basi appendiculatis in pixidem connatis in petiolum alatum angustatis latè ovato-rhombeis triplinerviis subdentatis, pedicellis è dichotomiis ortis folio paulò brevioribus, invol. squamis ext. ovali-oblongis capitulo duplò longioribus, int. achænia involv. ovatis lævibus

apice summo tubercula 3-5 minima gerentibus. ① in Mexico (Cav. Alam.), inter Valladolid et Pánuco (H. B.), in valle Tolucanâ (Berl. pl. exs. n. 1134). *Alcina perfoliata* Cav. ic. 1. p. 10. t. 15. Lag. nov. gen. p. 32. *Wedelia perfoliata* Willd. sp. 3. p. 2335. *Polymnia perfoliata* Poir. dict. 5. p. 506. *Melamp. connatum* fl. mex. ic. ined. *Camutia perfoliata* Bonat. ex Steud. (v. s.)

† *Species non satis notæ.*

19. M. ? *DOMBEYANUM*, caule herbaceo erecto basi glabrato medio sparse apice conferte hirtis, foliis sessilibus basi breviter attenuatis ovatis vix acutis grosse dentatis glabrisculis, junioribus subtus villosis et glandulosis, capitulis in dichotomis summis subsessilibus, invol. squamis ext. ovali-oblongis obtusiusculis..... ① ? in Amer. austr. legit Dombey sed locus propr. ign. Ob capitula nimis juniora caract. gener. non ertere quivi. Habitus ferè M. paludosus. (v. s. comm. à Mus. reg. Par.)

20. M. ? *RUDERALE* (Swartz fl. ind. occ. 1372), caule herbaceo erecto tetragono pubescente, foliis petiolatis ovato-lanceolatis acutis serratis trinerviis hispidis, pedicellis axillaribus petiolo brevioribus, capitulis subnatis discoidis. ① in ruderatis Jamaicæ australioris. A genere videtur removenda. Inv. squamæ 5 oblongæ erectæ. Fl. 10-11 hermaphr., exteriores tantum fertiles. Fructus ovato-oblongi tetragoni, ad angulos obtusè muricati, calyculo minuto cylindrico superati. Paleæ acuminatæ flosculos cingentes. (Sw.)

21. M. ? *HIDALGOA*, caule volubili, foliis oppositis longè petiolatis ternatisectis, segmentis ovatis crassiusculis glabris serratis, invol. biseriali, ligulis 5, achænio drupaceo ovato, initio apice auriculato, maturo forcipato. — in densissimarum sylvarum marginibus arborescens scandens prope flumen Blanco non longè à San-José del Corral Mexicanorum. *Hidalgoa ternata* Lallav. et Lex. nov. veg. mex. 1. p. 15 non Less. Ligulæ coccineo-aurantiaceæ. Discus flavus. Conf. *Hidalgoa* Lessingii.

†† *Species exclusæ.*

M. *AUSTRALE* Linn. = *Acanthospermum xanthioides*.

M. *BARANQUILLÆ* Spreng. = *Sclerocarpus Africanus*.

M. *HUMILE* Sw. = *Acanthospermum humile*.

CCLXXXVI. *ACANTHOSPERMUM* Schrank pl. rar. hort. monac. v. 2. (1819) t. 53. — *Centrospermum* Kunth nov. gen. am. (1820) p. 271. t. 397 (non Spreng.). Cass. dict. 59. p. 235 et 244 (excl. syn. Spreng.). Less. syn. 217. — *Echinodium* Poir. mss. ex Cass. — *Melampodii* sp. Læfl. Linn. Brown.

Capitulum pluriflorum heterogamum, fl. radii 1-serialibus fœmineis ligulato-cucullatis, disci masculis tubulosis 5-dentatis. Invol. 1-seriale, squamis 5 ellipticis concavis. Recept. planum paleaceum, paleis exter. ovaria involventibus extus aculeis herbaceis uncinatis onustis, inter. concavis apice dentatis fl. masculos fœventibus. Anthæræ ecaudatæ appendiculatæ. Styli rami recurvati apice glabri. Achænia paleis aculeatis arcuè involuta compressa obtusa basi subattenuata epapposa. — Herbæ Americanæ ramosæ demissæ aut diffusæ trichotomæ. Folia opposita subtus punctata. Capitula solitaria ad apices aut dichotomias ramorum per anthesin parva. Cor. luteæ.

Sect. 1. *XANTHIOIDES*.

Paleæ exteriores seu achæniorum tegumenta longitudinaliter striato-sulcata, secus nervulos aculeos herbaceos uncinatos gerentia, apice obtusa nec in cornua producta, fructus *Euxanthii* referentia.

1. A. *XANTHIOIDES*, caule diffuso, ramis pubescentibus, foliis petiolatis subintegris aut dentatis glabrisculis, fructibus apice inermibus. ① in Amer. austr.

- a. obtusifolium*, foliis ovalibus obovatisve obtusis minùs serratis, ramis pubescentibus. — in Cumanà? (Loefl.), in Brasiliæ prov. Rio-Grande (h. Mus. Bras. n. 1069). *Melampodium australe* Loefl. itin. 268. Linn. sp. 1303. (v. s. ex Forsyth sine loci design.)
- β. acutifolium*, foliis ovatis acutis serratis, ramis pubescenti-hirtis. — inter Cumana et Bordonas. *Centrospermum xanthioides* H. B. et Kunth nov. gen. am. 4. p. 271. t. 397.
- γ. glabratum*, foliis ovalibus utrinquè acutis serratis glabris, ramis pubescentibus. — in Brasilià (Sell.), circa Rio-Janeiro (Martius! Lund! Gaudichaud!). *Acanthospermum Brasilum* Mart. in litt. (v. s.)
2. *A. hirsutum*, caule diffuso, ramis hirsutis, foliis petiolatis latè ovalibus obovatisve obtusis subdentatis secus nervos subtùs hirsutis. ① in Brasilià circa Bahiam leger. cl. Salzman, Blanchet et Lhotsky, et ad Mariannam cl. Vauthier (pl. exs. n. 563). *Acanthospermum Brasilum* Schrank pl. rar. h. monac. 2. p. 53. An prioris adhuc varietas? (v. s.)

Sect. II. CERATOCHLÆNA.

Paleæ exter. seu achæniorum integumenta trigona esulcata in cornua duo elongata producta.

3. *A. HUMILE*, caule erecto subhirsuto, foliis sublyratis inciso-dentatis. ① in ruderatis et cultis Jamaicæ (Sw.), Sancti-Domingi (Bert.); ubi dicitur *Semæcontra-Marron*. *Melampodium humile* Swartz! prod. 114. flor. 1370. Moris ill. pl. rar. h. taur. p. 19. act. Acad. Taur. 36. p. 193. *Centrospermum humile* Less. syn. 217. (v. s.)
4. *A. HISPIDUM*, caule erecto hispido, foliis ovatis basi cuneatis sessilibus à medio subdentatis membranaceis subhispidis. ① in Brasiliæ sabulosis maritimis circa Bahiam legit cl. Salzman. Pili longi albi patentes articulati. Fructus exactè prioris paulò magis echinatus. (v. s. comm. à cl. Salzmann.)

Div. v. AMBROSIEÆ. — Ambrosiæ archetypæ *Cass. dict.* 29. p. 176. *opusc.* 3. p. 61. — Senecionidæ Ambrosiæ *Less. syn.* 218 *excl. lva.*

Capitula in iisdem stirpibus, alia mascula, alia fœminea. Invol. ♂ gamophylla multiflora, ♀ uni- aut pauciflora. Cor. tubulosæ breviter 5-dentatæ aut interdum in fl. ♀ nullæ. Anthere ecaudatæ approximatæ, nec verè cœnetæ. Achænia calva.

CCLXXXVII. XANTHIUM. *Tourn. inst.* t. 252. *Linn. gen.* n. 1056. *Gærtn. fr.* 2. p. 418. t. 164. *Cass. dict.* 25. p. 195. *Less. syn.* p. 219.

Capitula monoica homogama. ♂ Involucrum subglobosum multiflorum squamis liberis 1-seriatis. Recept. cylindraceum paleaceum. Corollæ clavatæ pilosiusculæ breviter 5-lobæ. Stam. filam. corollæ vix adnata, antheræ liberæ ecaudatæ. Stigm. 2 concretæ. ♀ Flores 2 intra invol. hamato-aculeatum rostris 1-2 superatum inclusi. Cor. filiformes. Stam. o. Stigm. 2 linearia divergentia ad latera papillifera. Achænia compressa singula in loculis singulis involucri demum indurati et bilocularis nidulantia. Embryo oblongus. — Hærbæ sporadicæ ramosæ annuæ. Folia alterna variè incisa. Capitula irregulariter glomerato-spicata, spicâ terminali supernè masculâ infernè fœmineâ. — Genus inter Compositas valdè abnorme! ad Urticeas à Ventenatio, ad Cucurbitaceas à Reichenbachio rejectum!

Sect. I. EUXANTHIUM. — *Xanthium Diosc.*

Involucri fructigeri rostrajduo apice plūs minūs uncinato-incurvata rariūs recta. — Spinæ ad basin foliorum nullæ. Folia cordata in sinu cuneata triloba dentata.

1. *X. MACROCARPUM* (DC. fl. fr. suppl. p. 356), involucro fructigero ovali-oblongo inter aculeos et ad basin rostrorum hispidissimo, rostris apice uncinatis. ① in vineis et cultis Galliae australis et Italiae, nec non in America boreali, Mexico, agro Bonariensi et verisim. Peruviano (Domb.) *X. orientale* Linn. f. dec. 33. t. 17 (excl. syn. et patr.). Gartn. fr. 2. t. 164. Cav. icon. 3. t. 221 (mala seu dubia). *X. echinatum* Murr. comm. goett. 1784 cum ic.? Willd. sp. 4. p. 374? Balb. et Nocc. fl. Ticin. 2. p. 184. t. 17. *X. Italicum* Moretti! dec. 5. p. 8. n. 50. Reichenb. icon. 4. p. 22. t. 323. *X. maculatum* Rafin. in Sillim. journ. vol. 1. p. 149 ex Torr.! in litt. *X. Brasilicum* Arrab. fl. flum. 10. t. 23. Invol. fruct. latitudine duplò longius. (v. v. et s.)

β. *glabratum*, involucro inter aculeos et in rostris glaberrimo. ① in Carolina. *X. Americanum* Walt. car. p. 231. *X. Cauadense* Linn. sp. 1400? Mill. dict. n. 2. — Dill. elth. 2. p. 432. t. 321 excl. fig. 13 et 14. An ab hoc satis differt *X. occidentale* Bertol. lucubr. 1822. p. 38 à cl. Bertero in Sancto-Domingo lectum? (v. s. comm. à cl. Bosc et à cl. Bertero sine fr.)

2. *X. INDICUM* (Roxb.! cat. calc. 67), involucro fructigero ovali inter aculeos et ad basin rostrorum pubescente, rostris apice uncinatis. ① in Ægypto (Coqneb.), in ruderatis Indiae orientalis frequens, verisim. etiam in China. *X. orientale* Linn. sp. 1400 excl. syn. *X. Chinense* Mill. dict. n. 5? *X. Indicum* Wall.! cat. et herb. n. 291. Wight. herb. *X. strumarium* Delil. ill. fl. eg. n. 891? Proximè accedit ad sequens sed paulò majus et fructus rostris diversum DC. in Wight. contrib. p. 17. (v. s.)
3. *X. STRUMARIUM* (Linn. sp. 1400), involucro fructigero ovali inter aculeos et ad basin rostrorum pubescente, rostris rectis, foliis basi cordatis 3-5-nerviis 3-5-lobis grossè dentato-incisis. ① ad vias, agros, vineas, rudera totius Europæ frequens, etiam in Sibiria, in Africa boreali, imò in America verisim. introd. Fl. dan. t. 270. Lam. ill. t. 765. f. 1. Engl. bot. t. 2544. Schkuhr. handb. t. 291. *X. vulgare* Lam. fl. fr. 2. p. 56. — *Xanthium* Fuchs. hist. 579. Blackw. herb. t. 444. Dill. elth. t. 321. f. 13 et 14. (v. v. et s.)
4. *X. INÆQUILATERUM*, involucro fructigero ovali inter aculeos et ad basin rostrorum pubescente, rostris rectis, foliis ovatis angulato-trilobis repando-serratis basi inæquali attenuatis, summis penninerviis. ① in ruderatis ins. Javæ prope Buitenzorg legit cl. Blume. *X. orientale* Blume! bijdr. p. 915 non allor. Fructus minor quàm *X. strumarii*, cæterum similis. Species foliis basi inæqualibus cuneatis insignis. (v. s. comm. à cl. invent.)

Sect. II. ACANTHOXANTHIUM.

Involucri fructigeri rostrum sæpius unicum semper rectum. — Spinæ validæ ad basin foliorum. Folia nunquàm cordata, basi cuneata.

5. *X. SPINOSUM* (Linn. sp. 1400), spinis tripartitis, foliis trilobis suprà minutè puberulis secus nervos canis subtus canescentibus. ① ad agros et secus vias Galliae præsertim australis, Italiae, Austriae littoralis, Hungariae, Podoliae (Bess.), et introd. in America hinc inde. Lam. ill. t. 665. f. 4. Moris. ox. s. 15. t. 2. f. 3. Spinæ flavescentes graciles pollicem longæ, parte indivisâ vix bilineari. (v. v. et s.)

β. *brachyacanthum*, spinis brevioribus. In specimine Brasiliano mihi obvia ex comp. Bras. n. 867, spinæ multò breviores quàm in Europæis. An discrimen fortè constans? Conf. *X. spinosum* Arrab. fl. flum. 10. t. 24. (v. s. in h. Mus. reg. Par.)

6. *X. CATHARTICUM* (H. B. et Kunth nov. gen. am. 4. p. 274), spinis ternatis, foliis pinnatifido-lacinatis suprà strigoso-hispidis subtus cano-tomentosis. (Linn. agro Quitensi prope Chillo et Quito. Herba cathartica Caxamarone.)

7. *X. PARVIFOLIUM*, spinis solitariis simplicibus, foliis oblongis subintegerrimis mucrone spinuloso terminatis utrinque glabrescentibus. (T) ? Patr. ign. verisim. Amer. austr. Spinæ crassæ 3-4 lin. longæ. Folia 4-5 lin. longa, $1\frac{1}{2}$ -2 lin. lata. (v. s. comun. ab am. Delessert.)

† *Species vix nota.*

8. *X. ? HOMOTHALAMUM* (Spreng. neu. entd. 1. p. 259), caule inermi, involucri androgynis pentaphyllis, foliis obovato-subrotundis. — in Brasiliâ. Species à genere videtur aliena, et postea ab auctore ipso ommissa! †.

†† *Species exclusa.*

X. FRUTICOSUM Linn. = *Franseria artemisioides*.

CCLXXXVIII. *FRANSERIA* Cav. ic. 2. p. 78. Willd. sp. 4. p. 378. Cass. dict. 17. p. 364. Less. syn. 219.

Capitula monoica homogama. ♂ Invol. hemisphæricum 8-12-dentatum seu squamis 8-12 uniserial. concretis. Recept. paleis filiformibus inter flores. Flores 15-20. Cor. 4-fida. Stam. imæ corollæ inserta, vix huic adnata. Antheræ liberæ. Pistilli rudimentum. ♀ Invol. ovatum extus aculeis uncinatos aut spinescentes gerens et in cornua cava 3-4 apice desinens, 1-florum aut in loculamenta 3-4 uniflora divisum. Cor. nulla aut stylo adpressissima. Anth. o. Stylus filiformis intra cornua productus bifidus. Stigm. apice obtusa. Achænium calvum in invol. nidulans. — Suffrutices aut herbæ in Americâ crescentes. Folia alterna grossè dentato-lobata uni- aut bipinnatisecta. Capitula spicata, macula superiora, feminea inferiora sæpius aggregato-spiculata. — Genus medium iter *Xanthium* et *Ambrosiam*.

Sect. I. XANTHIOPSIS.

Invol. fructigera aculeis mollibus uncinatis hispida.

1. *F. ARTEMISIOIDES* (Willd. sp. 4. p. 374. hort. berol. 1. t. 2), fruticosa, foliis bipinnatifido-dentatis, petiolis alatis, limbo supra virescente subtus albidopubescente. ♂ in Peruvîâ. *Xanthium fruticosum* Linn. f. suppl. 418. *Ambrosia arborescens* Lam. dict. 1. p. 127. Invol. masc. circ. 12-dentatum, foem. ex Willd. biflorum. (v. s.)
2. *F. AMBROSIOIDES* (Cav. ic. 2. p. 79. t. 200. excl. syn.), fruticosa, foliis oblongo-lanceolatis dentatis basi cordatis truncatis inæqualibus utrinque hirsutis viridibus, petiolo sæpè appendiculato cæterum aptero. ♂ in Mexico. fl. mex. ic. ined. Invol. masc. 8-fidum. Cor. more Compositarum 5-nervis! Inv. femineum intus 3-4-loc. 3-4-florum. (v. v. c. in hort. Monsp.)

Sect. II. ACANTHOLÆNA.

Invol. fructigera aculeis spinescentibus echinata.

3. *F. CHAMISSONIS* (Less. in linnæâ 1831. p. 507), subherbacea flexuoso-decumbens, foliis rotundo-ellipticis utrinque hirsuto-incanis longè petiolatis. — in Californiâ. Hinc cl. Lessing. refert var. duas diversissimas in posterum fortè separandas.

a. *malvæfolia*, basi lignosâ, foliis integris plicatis crenatis hirsuto-subniveis.

β. *bipinnatisecta*, canescens, caule herbaceo inani, foliis bipinnatisectis, segmentis divergentibus oppositis, segmentulis minutis obtusis. (v. s. olim specim. in h. Par. culta ex sem. Californicis à cl. Collignon lectis et sp. sioc. ex Dougl. itin. ab hon. soc. hort. Lond. comm.)

CCLXXXIX. *AMBROSIA* Tourn. inst. t. 252. Linn. gen. n. 1057. Gaertn. fr. 2. p. 417. t. 164. Cass. dict. 25. p. 203. Less. syn. 219.

Capitula in eadem plantâ alia fœminea ad basin spicæ, alia mascula ad apicem, omnia homogama. ♂ Capitula pluriflora. Invol. squamis uniserialibus ferè omnino in cupulam concretis. Recept. epaleaceum. Cor. tubulosa brevis. Stam. corollæ non adhærentia. ♀ Capitula plura 1-flora aggregata, invol. communi cincta, bracteolis interpositis instructa. Corolla nulla. Styli rami elongati è collo involucelli exserti. Achænium ovatum intrâ invol. persistens incurvatnm et sæpius in dentes aut cornua excrescens. — Herbæ interdum suffruticosæ, aliæ gerontogæ, aliæ Americanæ. Folia infer. opposita, super. alterna, pinnatilobata trilobata aut indivisa. Rami desinentes in spicam basi compositam fœmineam, apice masculam simplicem. — Genus quoad caracteres genericos difficilè intelligendum et quoad specierum discrimina, fructibus sæpius indescriptis, non satis elaboratum.

* *Foliis uni- aut sæpius bipinnatifidis.*

1. *A. MARITIMA* (Linn. sp. 1481), tota villosissima cinereo-canescens, caule divaricato villosulo-hispido, foliis bipinnatipartitis, lobis apice obtusis hinc inde utrinquè 1-dentatis, invol. masculo breviter hispido campanulato 15-20-floro, fœm. fructigero ovato cornubus 4-5 rectis echinato. (I) in herbidis ad ripas maris et fluviorum in Hispaniâ circa Taragonam (Delaroche!), ad Niceam (All.), Venetiam (Moric.), in Hetruriâ, agro Neapolitano (Ten.), Cretâ (Oliv.), Ægypto (Coqueb. ! Bové!), Sennaar (Acerb. !), Cappadociâ (Linn.), etc. Schkuhr handb. t. 292, sed cornua inv. fruct. omitta. A. villosissima Forsk. descr. eg. p. 160. Dod. pempt. 35. ic. A. artemisiæfolia et A. elatior Bess. prim. fl. gal. 2. p. 389 et 390 mihi ignotæ, videntur ex descript. meræ *A. maritimæ* variet. ? (v. s.)
2. *A. SENEGALENSIS*, tota cinereo-pubescent, summitatibus villosis, caule ramoso subvillosulo, foliis bipinnatipartitis, lobis acutis dentatis inter dentes subserrulatis, inv. masculis campanulatis breviter hispidis 15-20-floris, fœm. ovatis cornubus 4-5 rectis echinatis. (I) secus ripas fl. Senegal locis humidis argillosis leger. cl. Bacle et Perrottet. An *A. maritimæ* var. ? sed pubes multò minor et lobi fol. longiores acuti denticulati discrimen in genere adeò intricato fortè constituunt. (v. s.)
3. *A. CRITHMIFOLIA*, caule basi suffruticuloso supernè confertè velutino, foliis oppositis petiolatis crassiusculis bipinnatifidis, lobis angustis obtusis utrinquè pube adpressâ puberis, spicis paucis, terminali multò longiore, fructibus junioribus villosis inermibus. ♀ ♂ in ins. Cubâ circa Havanam legit cl. Ferrero. Ambr. maritima Ferr. ! in litt. 1821. Inter Americanas magis enim quàm alteræ species *A. maritimam* refert ? (v. s.)
4. *A. ABROTANIFOLIA* (Wall. ! cat. n. 3234. comp. 344), minutissimè pubescens, caule basi suffruticoso ramosissimo, foliis pinnatifidis basi cuneatis petiolatis, lobis oblongis acutis hinc inde dentatis subtrifidis, summis linearibus integerimis, racemis brevibus foliosis, invol. masc. vix puberulo 5-7-floro, fœm. ovato vix corniculato. ♂ culta in. h. bot. calc. et ex Wall. vix non Indica. Eandem sine loci design. video inter plantas Lundii Brasilianas, et circa Bahiam à cl. Blanchet (pl. exs. n. 972) lectam. Facilè ab omnibus differt capitulis masculis minimis. (v. s.)
5. *A. FRUTICOSA*, caule fruticoso erecto ramoso, foliis bipinnatipartitis, lobis lineari-sublanceolatis acutis integris aut lobulis hinc inde incisus suprâ glabrisculis subtus adpressè pubescentibus, racemis terminalibus elongatis inter capitula mascula ebracteatis, fructibus echinato-spinosis. ♂ in locis umbrosis Mexici in distr. Tamaulipas prope Reynosa april. flor. legit cl. Berlandier (pl. exs. n. 2112 !). Variat caule ramisque setis strigillosis sparè hispidis aut vix

hispida subpubescens, capitulis masculis minoribus aut paulò majoribus. (v. s.)

β. *intermedia*, caule suffruticoso supernè hispido, foliis bipinnatilobatis lobis acutis, racemis elongatis confertis. ♂ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 869!). Media inter *A. fruticosam* et *A. crithmifoliam*, priori caule et foliis, posteriori inflorescentiâ similis. An spec. propria? (v. s. in h. Mus. reg. Par.)

6. *A. ELATIOR* (Linn. sp. 1401), caule virgato vix pubescente, foliis bipinnatifidis glabriusculis, petiolis ciliatis, racemis paniculatis, lateralibus abbreviatis, fructibus apice 6-cornutis. ① in pascuis fertilibus à Canadâ ad Carolinam (Pursh), in Georgiâ circa Savannah (h. Duby!). Willd. sp. 4. p. 376. Ell. sketch 2. p. 477. Affinis *A. artemisiæfoliæ*, sed foliis caulibusque glabriusculis, racemis lateralibus caule brevioribus et petiolis longè ciliatis differt. (v. s.)
7. *A. POLYSTACHYA*, tota pube confertâ velutina, caule sulcato, foliis bipinnatilobatis, lobis elongatis inciso-dentatis acuminatis, racemis numerosissimis in paniculam amplam dispositis, fructibus inermibus. — in Brasiliæ prov. Sancti-Pauli locis cultis humidis nov. et dec. flor. legit cl. Lund; etiâ video in h. Mus. imp. Bras. sub n. 460! Affinis hinc *Franseriæ artemisioidi*, illinc *Ambr. artemisiæfoliæ*; à priore diversa caule herbaceo, petiolis teretibus nec alatis et caract. genericis; à posteriore caule sulcato multò elatiore, foliorum lobis multò magis acuminatis, rachi inter lobos lobulatâ, racemis numerosis, fructu inermi, etc. An *A. maritima* Arrab. fl. flum. 10. t. 26 non Linn.? (v. s. comm. à cl. Lund.)
8. *A. ARTEMISIÆFOLIA* (Linn. sp. 1401), caule tereti pubescente, foliis bipinnatifidis subcanescentibus, summis pinnatifidis, ramis fastigiatis, racemis terminalibus ternis, fructûs spinis brevibus acutis. ① in agris Amer. bor. à Canadâ ad Carolinam Ell. sketch 2. p. 477. Lam. ill. t. 765. f. 1. *A. absinthifolia* Michx. fl. bor. am. 2. p. 183. Fortè etiâ hab. in Mexico (fl. mex. ic. ined.) circa urbem (Berl. ! pl. exs. n. 573) et ad Havanam (Ferr.!), sed ex sicco de identitate specim. dubius hæreo. — Hern. mex. p. 446. f. 1? (v. s.)
9. *A. PSILOSTACHYA*, caule herbaceo erecto tereti paniculato-ramoso foliisque scabridis, foliis subsessilibus pinnatipartitis lobis lineari-lanceolatis acuminatis hinc indè pinnatim incisis, racemis plurimis gracilibus inter capitula mascula remota ebracteatis, fructibus infra apicem tuberculis obtusis 5-6 onustis apice conico elongato. — in Mexico inter San-Fernando et Matamoros legit cl. Berlandier (pl. exs. n. 2280!). Fructus plurimi parvi in invol. fœm. Capitula masc. pubero-scabrida parva. (v. s.)
10. *A. CONFERTIFLORA*, caule herbaceo erecto sulcato paniculato-ramoso, foliis petiolatis pinnatisectis suprâ glabriusculis subtùs in petiolo puberulo-scabris, segmentis pinnatipartitis linearibus elongatis integris aut subincisis, racemis plurimis elongatis confertifloris inter. capitula masc. ebracteatis. — in Mexico ad Matamoros legit cl. Berlandier (pl. exs. n. 2297!). Affinis *A. psilostachyæ*, sed foliis petiolatis pinnatisectis et capitulis masc. confertis diversa. Fructus ex statu juniore non satis noti. (v. s.)
11. *A. ACANTHOCARPA* (Hook. fl. bor. am. 1. p. 309), valdè ramosa, setis asperis rigidis hispidis, foliis bipinnatifidis lobis linearibus, racemis terminalibus, fructibus spinis rigidis subulatis longis patentibus undiquè tectis. — in Amer. bor. ad ripas fluminum sœcus Saskatchewan, Red-River, Columbia.
12. *A. PERUVIANA* (Willd. sp. 4. p. 377), caule erecto virgato glabriusculo, foliis infer. bipinnatifidis, interm. pinnatifidis, summis oblongo lanceolatis indivisis, racemis terminalibus solitariis. ② in Peru (Poepp. ! pl. exs. n. 17), ad sinum Chorillo (Macrae!), in Mexico (Alam.!), ad Real del Monte (Hænke!), An stirps Mexicana omninò eadem ac Peruviana? Huc refer. videtur *A. humilis* Arrab. fl. flum. 10. t. 25? (v. s.)
13. *A. HETEROPHYLLA* (Muhl. in Willd. sp. 4. p. 377), caule pubescente aut villosulo paniculato, foliis caulinis pinnatifidis subdentatis petiolatis, rameis summis lanceolatis sessilibus, petiolis longè ciliatis, racemis terminalibus sub-

solitarius, fructu sub apice spinis 5-6 acutis instructo. ① ad ripas fluminum circa Philadelphiam (Pursh). (v. s.)

14. *A. MICROCEPHALA*, glabrescens, foliis superioribus alternis longè petiolatis pinnatilobatis sinuatisve obtusis, lobis inæqualibus integris obtusissimis, foliis summis indivisis subspathulatis obtusis. — in Cayennâ legit olim cl. Patris. Invol. mascula nutantia orbiculata integerrima dimidiò minora quàm in omnibus aliis. Fl. fœmin. juniores tantum vidi. (v. s.)
15. *A. PANICULATA* (Michx. fl. bor. am. 2. p. 183), caule pubescente ramosissimo supernè paniculato, foliis utrinquè viridibus bipinnatifidis, lobis lanceolatis, fructibus aggregatis pusillis globoso-obovatis subinermibus. ① in cultis à Canadâ ad Floridam necnon ad Nootka Am. bor. occid. (Hook.). Willd. sp. 4. p. 376. Ell. sketch 2. p. 478. Iva monophylla Walt. car. 232. — Plak. alm. t. 10. f. 5. — Ambr. pusilla Rafin. in litt. hujus videtur var. nana. (v. s.)
16. *A. TOMENTOSA* (Nutt. gen. am. 2. p. 186), caule humili, foliis bipinnatifidis subtus albis et tomentosis, spicis solitariis. ½ in Louisianâ super. ad ripas flum. Missouri, rara, 1-2 pedalis.
17. *A. HISPIDA* (Pursh fl. suppl. p. 743), undiquè canescenti-hispida, foliis bipinnatifidis, lobis incisis, racemis terminalibus subpaniculatis ① in Carolinâ meridionali ex herb. Sherard sed Elliottio et Nuttallio ignota.
18. *A. TENUIFOLIA* (Spreng. syst. 3. p. 851), caule hispido, foliis bipinnatifidis subtus canescentibus, lobis remotis linearibus acutis subdentatis, racemis solitariis. — in Montevideo.

** *foliis trilobatis.*

19. *A. CUMANENSIS* (H. B. et Kunth nov. gen. am. 4. p. 276), ramis hirsutis, foliis tripartitis pubescentibus subtus canescentibus, lobis pinnatilobatis, intermedio trifido, fructibus 4-5 spinosis. ½ in humidis prope Cumanam Novæ Andalusie.
20. *A. TRIFIDA* (Linn. sp. 1401), caule erecto piloso aspero, foliis etiâ super. oppositis pilosis et scabris trifidis, lobis ovali-lanceolatis acuminatis serratis, petiolo angustè alato ciliato, fructu infra apicem tuberculis 6 subspinosis instructo apice conico. ① ad ripas flum. Amer. bor. à Canadâ ad Georgiam. Ell. sketch 2. p. 476. — Moris. ox. s. 6. t. 1. f. 4. Herba 4-6-pedalis. Cor. albæ. (v. v. c. et s. sp.)
21. *A. APTERA*, junior sparsè setulosa, adulta tuberculis aspera, caule striato, foliis trifidis, lobis ovali-lanceolatis acuminatis serratis, petiolo tereti aptero, fructu infra apicem tuberculis 5-6 subspinosis instructo apice conico. ① in Mexico circa Bejar locis inundatis aug. flor. legit cl. Berlandier (pl. exs. n. 1778!). Herba 3-6-pedalis ad A. trifidam proximè accedens, sed petiolo faciliè distincta. Lobi interd. 5, duobus exter. minoribus. (v. s.)

*** *foliis indivisis.*

22. *A. BIDENTATA* (Michx. fl. bor. am. 2. p. 182), hirsutissima, foliis arcuè sessilibus indivisis oblongis supra basin latecentem utrinquè unidentatis, fructibus tetragonis infra apicem quadrispinosis. ① in regione Illinoensium (Michx.); ad partes occident. montium Alleghany (Pursh).
23. *A. INTEGRIFOLIA* (Muhl. in Willd. sp. 3. p. 275), caule...., foliis oppositis sessilibus ovatis acuminatis basi ciliatis, racemis terminalibus plerumquè ternis. ① ad ripas paludum Pensylvaniæ et Virginie. Pursh fl. 2. p. 530. (v. s.)

† *Species exclusa.*

A. ARBORESCENS Linn. = *Franseria artemisioides*.

Div. VI. IVEÆ. — Ambrosiæ IVEÆ Cass. dict. 29. p. 175. —
Ambrosiæ spurix Cass. opusc. 3. p. 61.

Flores ♂ et ♀ in iisdem capitulis. Cor. omnes tubulosæ 5-dentatæ aut inter-

dum in ♀ nulla. ♂ Anthæræ liberæ approximatæ. Styli apice incrassati. ♀ Styli bipartiti aut simplices elongati ferè subulati hispiduli.

CCXC. PINILLOSIA *Ossa in litt. descr. et ic.*

Flores monoici ♂ et ♀ in iisdem capitulis: fœminei 2 apicali ante invol. squamas 2 majores siti; masculi 2 cum iis alterni corollâ tubulosâ 5-fidâ donati. Invol. 4-phyllum squamis cruciatim oppositis 3-nerviis imâ basi subconcretis constans, bracteolis 2-linearibus brevibus extus stipatum. Recept. planum fortè omninò epaleaceum. Cor. ♂ tubo tereti, fauce apertâ, lobis 5 ovatis papilloso-puberis. Anth. exsertæ ovato-oblongæ, utrinquè obtusæ approximatæ nec concretæ. Stylus antheras paulò superans apice bilobus incrassatus glaber. Ovarium abortivum. Cor. ♀ nulla. Styli rami à basi ferè liberi elongati graciles apice subhispidi. Achænia obovata subcuneata, juniora præter apicem mucroniformem instructa pappo 4-aristato, aristis crassis retrorsum aculeolatis apice capitellatis, adulta quadricornuta cornubus conicis nudis patulis et nectariò mucroniformi terminata. Embryo crassus, cotyl. obovatis extus convexis. — Herba Caribæa repens glaberrima, Tetranthi facie. Folia opposita orbiculata longè petiolata integra. Pedunculi axillares petiolis longiores nudi 1-cephali. Flores albi. — Plantam antè diem supremum misit b. Ant. de La Ossa rogans ut si genus ex eâ extruendum, dicatum fuerit ill. Claudio Martinez de Pinillos insulæ Cubæ gubernatori, scientiæ fautori!

1. P. TETRANTHOIDES. ? in insulâ Sancti-Domingi legit cl. Bertero et in Cubâ circa Havannâ cl. de La Ossa et de La Sagra. Tetranthus Berterii Spreng.! syst. 3. p. 459. Herba elegans ferè Linnæam referens sed multò minor. (v. s.)

CCXCI. TETRANTHUS *Swartz prod. 116. fl. ind. occid. 3. p. 1885. t. 27. Cass. dict. 57. p. 343. Less. syn. 420.*

Flores monoici, ♂ et ♀ in iisdem capitulis; fœminei 2 ante squamas inv. majores tubulosi 5-dentati; masculi 2 ante squamas minores etiâ tubulosi 5-dentati. Invol. 4-phyllum squamis ovalibus obtusis 2 sub ext. paulò majoribus constans, bracteolis 4-5 linearibus ciliatis acutis stipatum. Recept. planum paleis convolutis amplex. Cor. tubo gracili, fauce infundibuliformi, lobis oblongis subpuberulis subinæqualibus. ♂ Anthæræ oblongæ apice breviter mucronatæ approximatæ nec concretæ. ♀ Styli rami longi graciles exserti apice divergentes hispiduli. Achæmium oblongum substriatum intra paleam involutum. Pappus brevis coroniformis ciliolatus. — Herbæ Domingenses repentes. Folia opposita petiolata ovata mucronulata. Pedunculi axillares nudi foliis longiores 1-cephali. Flores albidii.

1. T. LITTORALIS (Swartz! l. c.), caule petiolisque glaberrimis, pedunculis adpressè subpubescentibus, foliis ovato-orbiculatis basi hinc indè cordatis. (I) ad ripas fluviorum Hispaniolæ. (v. s.)
2. T. MINUTUS (Spreng. neu. entd. 3. p. 40), caule petiolisque hirsutis, pedunculis basi laxè hirsutis cæterum adpressè pubescentibus, foliis ovatis nupquàm cordatis. (I) in Sancto-Domingo legit cl. Bertero. T. littoralis Spreng.! syst. 3. p. 459 non Sw. Paleæ distinctè ciliatæ. Herba tota priore paulò major. (v. s.)

† *Species exclusa.*

T. BERTERII Spreng. = Pinillosia tetraanthoidea.

CCXCH. IVA Linn. gen. n. 1059. Gartn. fr. 2. p. 394. t. 164. Cass. dict. 24. p. 43. 59. p. 176. Less. syn. 219 non Riv. — Denira Adans. fam. 2. p. 118. — Tarchonanthes sp. Vaill. — Parthenii sp. Linn. ed. 1737. p. 265.

Flores monoici, ♂ et ♀ in iisdem capitulis, foemineis paucis 1-ser. in ambitu tubulosis seu campanulatis, masc. pluribus in disco tubulosis 5-dentatis. Invol. sæpius 3-5-phyllum campanulatum, squamis ovatis subuniseriatis, rarius imbricatum squamis 3-4-seriatis. Recept. planum paleis lineari aut lineari-spathulatis. Styli fl. foem. longè exserti subulati subhispidi, fl. masc. breviores apice incrassati. Achænia disci abortiva, radii obcompressa calva cornubus nec tuberculis onusta. — Herba aut frutices ex Americæ ripis orti. Folia opposita aut summa alterna trinervia serrata aut integra. Capitula ad fol. super. axillas solitaria seu terna, breviter pedicellata, deflexa, semiglobosa, racemos terminales foliaceos constituentia. Cor. albidæ. Anth. luteæ. — Habitus Artemisiæ sed capitulis monoicis et fructu facilè diversa.

§. 1. *Involucris 3-5-phyllis subuniseriatis.*

1. I. XANTHIFOLIA (Nutt. gen. am. 2. p. 185), herbacea, foliis oppositis petiolatis cordato-ovatis acuminatis duplò serratis sericeo-villosis subtus canescentibus, spicis paniculatis nudis, invol. squamis 5 ovato-lanceolatis acuminatis. ① in aridis circa Fort Mandan ad ripas flum. Missouri. Herba 5-6-pedalis. Folia ferè Xanthii strumarii.

2. I. CILIATA (Willd. sp. 3. p. 2386), caule herbaceo hirsuto, foliis oppositis petiolatis lanceolato-ovatis rariter serratis, petiolis longè ciliatis, racemo confertissimo, bracteis ovato-lanceolatis acuminatis longissimè ciliatis, capitulis axillaribus deflexis. ① in Amer. bor. pratis in dition. Kentuckensium et Illinoisensium. Pursh fl. 2. p. 580. Iva annua Michx. fl. bor. am. 2. p. 184 non Linn.

β. *latifolia*, foliis latè ovatis remotè dentatis trinerviis utrinquè petiolisque pubescentibus, summis alternis lanceolatis basi petiolisque ciliatis, bracteis ovatis acuminatissimis squamisquè involucri obtusis ciliatis. — in Mexico circa Bejar legit cl. Berlandier (pl. exs. n. 18641). An à specie Illinoisensi diversa? (v. s.)

3. I. AXILLARIS (Pursh fl. bor. am. 2. suppl. p. 743), herbacea subpilosa, foliis plerumquè oppositis lineari-lanceolatis integerrimis, capitulis axillaribus solitariis nutantibus, invol. simpliciusculi squamis ovatis. ① in Louisiana super. (Pursh), in aridis et salinis ripis fl. Missouri (Nutt.), ad flum. Saskatchewan et Columbia (Hænk.). — Herba 8-9 poll. Flores albi (Pursh). Folia cuneato-oblonga obtusa ex Nutt. gen. am. 2. p. 185. — An species ab Hookero fl. bor. am. 2. p. 309. t. 107 egregiè descripta et delineata, ipso auct. dubitante, omninò eadem ac Purshiana?

4. I. ANGUSTIFOLIA (Nutt. in litt. 1825), herba tota adpressè pubera, foliis sessilibus linearibus aut lanceolato-linearibus, infer. subserratis oppositis, super. alternis integerrimis, capitulis axillaribus subsessilibus deflexis, invol. obtusis dorso subpuberis. ① in Amer. bor. territorio Arkansasano detexit et mecum comm. cl. Nuttall. Capitula multò minora quàm I. annuæ, magis approximata et spicam foliaceam continuam constituentia. Fruct. mat. non vidi. (v. s.)

5. I. ASPERIFOLIA (Less. in linnæa 1830. p. 151), caule herbaceo adscendente, foliis lanceolato-linearibus acutis strigosis integerrimis, infer. oppositis, capitulis alternis cernuis, invol. 3-gono denticulato gamophyllo. — in pascuis prope Vera-Cruz.

6. I. ANNUA (Linn. sp. 1402), caule herbaceo striato petiolisque pubescentibus, foliis oppositis petiolatis ovali-lanceolatis utrinquè acutis grossè serratis trinerviis scabridis, summis integerrimis, capitulis axillaribus cernuis pedicellatis

invol. squamis ovalibus dorso pubescentibus. ① in America merid. (Linn.), sed locus propr. incert. Linn. hort. ups. 285. Schmid. icon. p. 61. t. 16. Variat foliis ternato-verticillatis. (v. s.)

7. *I. FRUTESCENS* (Linn. amoen. ac. 3. p. 25), fruticosa, foliis oppositis lanceolatis subpetiolatis acutis profundè serratis subscabris, summis integerrimis, capitis axillaribus deflexis vix pedicellatis depresso-globosis, invol. 5-phyllo obtusissimo. ③ in salsuginosis maritimis à Novâ-Angliâ ad Floridam (Pursh), in Novâ-Cæsareâ (Torr.!). Ell. sketch 2. p. 474. Gærtn. fr. 2. t. 164. Lam. ill. t. 166. f. 2. — Pluk. alm. t. 27. f. 1. vulgò dicitur *Marsh-Elder* ex Nutt. Variat foliis ternato-verticillatis. (v. v. c. et s. sp.)
8. *I. CHRISANTHIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 276), frutescens pubescenti-canescens, foliis oppositis petiolatis lanceolatis integerrimis glaucescentibus triplinerviis, capitulis ad axillas subternis cernuis, invol. squamis 3-5 orbiculatis, rec. paleis obtusis, ext. obovatis, int. lineari-spathulatis ③ in ins. Cubæ humidis juxta Havanam.

§. 2. *Involucris imbricatis 6-9-phyllis.*

9. *I. IMBRICATA* (Walt. car. 232), suffrutescens seu herbacea glabra, foliis sessilibus lineari-lanceolatis cuneatis carnosius, infer. oppositis trinerviis denticulatis, super. alternis integerrimis, invol. imbricatis 3-4-seriatis, recept. paleis spathulatis. ② ③ ad ripas arenosas maritimas Carolinæ et Georgiæ. Michx. fl. bor. am. 2. p. 184. Pursh fl. 2. p. 580. Nutt. gen. 2. p. 185. Ell. sketch 2. p. 475. desc. opt. Inv. squama 6-9. Cor. fœm. tennes, stigm. simplicibus elongatis. Cor. fl. fœm. interdum ad latus internum fissæ et ideo ferè ad ligulatas et ad div. seq. vergentes. (v. s. sp.)

+ *Species exclusa.*

- I. MONOPHYLLA* Walt. = *Ambrosia paniculata*.

CCXCIII. EUPHROSYNÆ.

Capitula monoica, fl. radii fœmineis apetalis 1-seriatis, disci masculis tubulosis 5-dentatis. Invol. 1-2-seriale patens squamis obovatis margine submembranaceis. Recept. planiusculum, paleis obovatis membranaceis, exter. majoribus, intimis multò minoribus. ♂ Auth. distinctæ. Cor. breves, tubo vix ullo. ♀ Styli rami teretes glabri obtusi. Flos centralis omnino abortivus filiformis apice penicilliformis. Achænia obcompressa; margine calloso cincta, lævia, inter squamas et paleas exter. sita, apice omnino calva. — Herba Mexicana glabra erecta, habitu nunc Ambrosiam nunc Parthenium referens. Folia alterna petiolata pinnatipartita, lobis oblongis pinnatilobatis sinuatis, lobulis secus rachiam intermixtis parvis sinuato-dentatis. Pili pauci subtus secus nervos primarios rigiduli. Capitula in paniculam ramosissimam disposita breviter pedicellata. Flores albid. Achæmium Parthenii, sed cum squamis vicinis non adhærens. — Ex fœminis nudis Euphrosynen dixi alludens ad nuditatem Gratiarum.

1. *E. PARTHENIFOLIA*. ② ? ① ? in Mexico circa urbem legit cl. Berlandier (pl. exs. n. 587). *Gynnostyles parthenifolia* Moc. fl. mex. icon. ined. Folia 5 lin. longa, variè pinnatilobata. (v. s.)

DIV. VII. PARTHENIÆ. — Senecionidæ Ambrosiæ Partheniæ Less.
syn. 220.

Fl. ♂ et ♀ in iisdem capitulis, fl. radii 1-seriatis fœmineis ligulatis, ligulis subcoriaceis persistentibus aut tardè deciduis, fl. disci masculis tubulosis 5-dentatis. Recept. sæpius paleaceum. Achænia radii obcompressa margine calloso cincta aut rarius trigona.

CCXCIV. CONIOTHELE.

Capitulum multiflorum monoicum, ligulis 7-10 uniserialibus fœmineis, fl. disci pluribus masculis tubulosis fauce ampliata 5-dentata. Invol. subcampanulati squamæ 7-10 subuniseriales oblongæ. Recept. epaleaceum convexum. Achænia... teretia calva, papillis minutis confertis pulverulento-velutinis, disci omnino nulla. — Herba Californica glaberrima erecta tenuis annua subsimplex. Folia alterna pinnatipartita, rachi lobisque remotis 2-3-jugis linearibus. Capitula terminalia pedicellata solitaria. Flores pallide flavi. Inter Parthenieas accedit ad Leptosynen sed ab hac et à cæteris differt recept. nudo. — Nomen ex *Novia*, pulveris, et *ἄνθη*, papilla, alludit ad fructum pulverulento-pappiliosum.

1. C. CALIFORNICA. ① in Novâ-Californiâ legit cl. Douglas. Semen hujus stirpis sub microscopio observavit cl. Duby. Papillæ aquâ imbibitæ elongantur, rumpuntur et ex iis oriuntur filamenta tennia non articulata molecularis intus donata quæ seminis quasi integumentum villosum conficiunt. (v. s. comm. ab hon. Soc. hort. Lond.)

CCXCV. LEPTOSYNE.

Capitulum multiflorum, fl. radii ligulatis fœmineis, disci tubulosi 5-dentatis bisexualibus abortu sterilibus? Invol. biserialia, squamis exter. seu bracteis paucis linearibus foliaceis disco sublongioribus, inter. ellipticis vix acutis submembranaceis disco æqualibus. Recept. paleis membranaceis lanceolatis persistentibus onustum. Ligulæ latè ovato-subrotundæ. Cor. tubi fauce obconicâ basi subbarbellatâ. Anth. ecaudatæ. Stylus bifidus, radii vix exsertus obtusus, disci inclusus ramis cono hispido lato brevi capitellatis. Achænia (ex flore) obcompressa, calva seu pappo minimo coronato superata, margine subalata, glabra. — Herba Californica glabra annua tenuis. Radix simplex gracilis. Folia subradicalia erecta angustissimè linearia, alia integerrima, alia lobis parvis linearibus pinnatipartita. Caulis teres gracilis nudus 1-cephalus foliis triplò quadruplò longior. Flores flavi. — Genus fructibus maturis ignotis subdubium. An, floribus disci verè hermaphroditis admissis, potiùs Anthemideum? sed Anthemideæ paleaceæ nullæ in Americâ hucusque notæ? — Nomen ex *λεπτοσύνη*, tenuitas, gracilitas.

1. L. DOUGLASII. ① in Californiâ legit cl. Douglas. Herba strictè erecta 6-8-pollic. (v. s. comm. ab hon. Soc. hort. Lond.)

CCXCVI. PARTHENIUM Linn. gen. n. 1058. Gærtn. fr. 2. p. 429. t. 168. Cass. dict. 38. p. 14. Less. syn. 319. — Partheniastrum Nissol. act. ac. Par. 1711. — Hysterothorus Vaill. act. ac. Par. 1720. — Argyrochæta Cav. ic. 4. p. 54. — Villanova Ort. non Lag. — Trichospermum Beauv. ined. ex Cass.

Capitulum multiflorum heterogamum, fl. radii 5 uniserialis fœmineis ligulatis, disci tubulosi 5-dentati abortu styli masculis. Invol. hemisphæricum biserialia, squamis ext. ovatis, int. suborbiculatis. Recept. conicum cylindraceumve, paleis semiamplect. membranaceis apice latioribus onustum. Fl. ♂ stam. imò tubo inserta, anth. juniores vix confertæ. Stylus indivisus. ♀ Ligulæ obcordatæ breves. Styli rami semiteretes obtusi. Achænia obcompressa lævia, cincta margine calloso, utrinquè basi squamis contiguis adhærente et demùm ab ovario secedente. Pappus bisquamellatus, squamellis aristæformibus seu suborbi-

culatis. — Herbæ seu suffrutices Americani hirsuto-canescens. Folia alterna. Capitula albidia fastigiato-paniculata.

Sect. I. PARTHENIASTRUM.

Pappi squamellæ tennes brevissimæ imò nullæ. — Folia indivisa.

1. *P. PAUCIFLOSUM* (Less. in Hassk 1830. p. 152), caule fruticoso, foliis petiolatis triangularibus acutis basi subcoarctatis aut in petiolum decurrentibus scabris dentatis, invol. squamis sub-3-seriatis obtusissimis. ♀ in Mexico ad Plan del Rio.
2. *P. TOMENTOSUM*, caule fruticoso, foliis petiolatis ovatis acutiusculis in petiolum non decurrentibus grossè dentatis suprâ molliter velutinis subtus petiolis ramisque albo-tomentosis, paniculâ corymbosâ polycephalâ confertâ, invol. squamis biseriatis, inter. obtusissimis, achænio epapposo. ♀ in Mexico inter Oaxacam et Mitlam legit cl. Andrienx (pl. exs. n. 281). Habitus in sicco paulò refert icon. 44. ser. 2. Sweet br. fl. gard. ad genus omninò diversum referendam. (v. s.)
3. *P. INTEGRIFOLIUM* (Linn. sp. 1402), caule herbaceo, foliis ovalibus asperis inæqualiter dentatis aut basi subincisis, infer. in petiolum decurrentibus, super. amplexicaulibus, invol. squamis exter. subacutis. ♀ in montosis Virginie (Pursh), Carolinæ (Fras.!) et Georgiæ (Ell.). Willd. hort. ber. t. 4. — Dill. elth. 302. t. 225. f. 292. (v. v. c. et s. sp.)

Sect. II. PARTHENICHÆTA.

Pappus biaristatus, aristis longis rigidis. — Folia obtusè sinuato-pinnatifida.

4. *P. RAMOSISSIMUM*, caule fruticoso ramosissimo, foliis obovatis obtusè sinuatis incano-lanatis, invol. squamis obtusissimis, achæniis puberis obovatis, aristis 2 rectis elongatis ♀ in Mexici ditione Sancti-Ludovici de Potosi legit Berlandier (pl. exs. n. 13421). Achænia fortè juniora non marginata nervo solubili observavi, cæterùm *P. incano* valdè simile. An genus diversum. (v. s.)
5. *P. INCANUM* (H. B. et Kunth nov. gen. am. 4. p. 260. t. 391), herbaceum totum albidoincanum, foliis obtusè sinuato-pinnatifidis, lobis inf. inciso-dentatis, terminali trilobo, invol. squamis omnibus obtusissimis, achæniis aristis recurvo-subdivergentibus ♂ in montibus circa urbem Mexici. (Berl.!) pl. exs. n. 63a.) (v. s.)

Sect. III. ARGYROCHÆTA. — *Argyrochæta* Cav.

Pappi squamellæ ovali-oblongæ obtusæ membranaceæ. Folia bipinnatifida.

6. *P. HYSTEROPHORUS* (Linn. sp. 1402), herbaceum puberulum basi subhispidum apice subincanum, foliis variè bipinnatifidis, invol. squamis acutis. ♂ in glareosis Amer. calid. frequens, nempe circa Mexico (Berl. pl. exs. n. 42), in prov. Texas (n. 1575), prov. Leoninâ (Mendez!), ins. Cubâ (h. Madr.), Trininitatis (Sieb.!), Jamaicâ (h. L'Hér.!), Sancti-Thomasii (Wyd.!), Martinicâ, (Willd.), etc., nunc in ins. Mauritiæ introd. (Bout.!). *Argyrochæta bipinnatifida* Cav. ic. 4. p. 54. t. 378. Villanova bipinnatifida Ort. dec. 48. t. 6. — Nissol. act. par. 1711. p. 423. t. 13. ♂. 2. (v. v. c. et s. p.)

† *Species exclusa.*

P. LUTRUM Spreng. = *Guizotia oleifera*.

CCXCVII. MENDEZIA.

Capitulum multiflorum radiatum, fl. radii paucis 1-serialibus fœmineis ligulatis, disci tubulosis 5-dentatis hermaphr. abortivis. Invol. squamæ paucæ 2-3-seriales ovales obtusæ imbricatæ. Recept. parvum

conicum, paleis membranaceis oblongis complicatis. Cor. radii tardè deciduæ ellipticæ, radii tenuis glaberrimæ. Styli disci rami apice summo hirtelli. Achænia radii obcompresso-plana elliptico-cuneata apice emarginata breviter biaristata utrinquè glabra margine subtuberculata, tuberculis pilos fasciculatos gerentibus; disci compressa abortiva. — Herba Mexicana annua trichotoma, caule tereti pubescente. Folia opposita oblongo-linearia integra puberula. Capitula terminalia solitaria bicolora, disco nempe intensè croceo ligulis pallidis. — Genus affine Verbesinæ ex achænio biaristato, sed multò affinius Parthenio et Tragocerati, priori characteribus, posteriori habitu. Dixi in honorem cl. Mendez botanici Mexicani, qui, mediante amiciss. L. Alaman, hanc et plurimas alias ab ipso detectas humanissimè communicavit.

1. *M. bicolor*. (1) in Mexici prov. Leoninâ ad occid. urbis Guanaxato legit cl. Mendez. (v. s.)

CCXCVIII. TRAGOCERAS Less. syn. 220. — Tragoceros H. B. et Kunth nov. gen. am. 4. p. 248. t. 385.

Capitulum 11-14 florum heterogamum, fl. radii 6-8-ligulatis fœmineis, disci 5-6 tubulosi masculis 5-dentatis. Invol. ovatum, squamis 5-7 imbricatis membranaceis obtusis. Recept. minutum paleaceum. Styli rami capillacei patentes. Achænium disci abortivum, radii oblongo-ovatum, calvum apice truncatum, corollâ persistente magnâ coronatum. — Herbæ Mexicanæ ramosissimæ graciles humiles annuæ. Folia opposita integra. Capitula solitaria ramos supernè aphyllis coronantia.

Sect. 1. EUTRAGOCERAS.

Achænia radii costis extus crenatis callosis destituta.

1. *T. zinnioides* (H. B. et Kunth l. c.), ligulis apice in cornua duo ferè subulata recurva fissis, achæniis utrinquè glabris subunicostatis. (1) in aridis calsis Mexici inter Aria et Montem Jorullo. Cor. disci violacæ, radii albidæ.
2. *T. microglossum*, ligulis indivisis concavis achænio duplò brevioribus extus villosis, achæniis intus 3-striatis apice utrinquè villosis. (1) in Mexici prov. Leoninâ ultra Guanaxato legit cl. Mendez. Cor. disci pallidæ, radii vircescentes. (v. s.)

Sect. II. SKIRROGLYPHIS.

Achænia radii costis 3 callosis, mediâ minore, lateralibus crenato-tuberculatis instructa.

3. *T. flavicomum*, ligulis indivisis 2-3-dentatis planis achænio vix brevioribus, supra subtilissimè velutinis, subtus sparsè glandulosi et puberulo-hirtelli, achæniis intus glabriusculis 1-costatis extus callosis 3-costatis, costâ mediâ minimâ, lateralibus crenato-tuberculatis. (1) in Mexico circa Tehuantepec, Chignitan et Ixtaltepec. Fl. jul. aug. Caulis purpurascens. Ligulæ supernè pallidè luteæ, subtus subvirescentes. (v. s. comm. à cl. L. Alaman.)
4. *T. schiedeanum* (Less. in linnæâ 1834. p. 269), ligulis emarginatis concavis reflexis subtus pubescentibus achænio multò brevioribus, achæniis intus pubescentibus costis integerrimis extus glabris calloso-3-costatis, costâ mediâ angustâ, lateralibus crenato-tuberculatis. (1) in Mexico ad Malpays de Nanligo. Ligulæ albidæ. Ex descr. conjungere videtur ligulas *T. microglossi* et achænia *T. flavicomi*.

† Species exclusa.

T. frueckeanifolium Spreng. = *Isostigma peucedanifolium*.

Subtrib. II. *HELICANTHÆE* Less. syn. p. 221. — *Helianthæ* sect. 2, 3 et 4 Cass. opusc. 3. p. 57. excl. gen. paucis.

Capitula sæpius heterogama, fl. radii ligulatis seu foemineis, disco hermaphr., rarissimè homogama discoidea, nunquam monoica seu dioica. Recept. paleaceum aut rarissimè ad centrum subnudum. Cor. fl. hermaphr. lobi crassi sæpius supernè papilloso. Achænia sæpius tetragona, nunc teretia aut obcompressa aut compressa, alia calva, alia pappo coroniformi aristato aut partim paleaceo, nunquam nec verè setoso nec omninò paleaceo superata. Anthera nigricantes ecaudatæ. Styli fl. herm. rami divergentes nunc truncati apice penicillati, nunc sæpius cono acuto aut appendiculo lineari terminati. — Folia sæpius opposita.

Div. I. *HELIOPSIDÆE*. — *Wedeliæ* et *Heliopsidæe* Less. syn. 221 et 223. — *Helianthæ* *Heliopsidæe* et *Rudbeckiearum* pars Cass. opusc. 3. p. 59.

Capitula heterogama, fl. radii foemineis, aut rariùs homogama discoidea. Achænia sæpè cortice crasso donata obovata angulata aut compressa, nunquam obcompressa, calva aut pappo coroniformi aut setis paucis rigidulis irregularibus coronata. — Cortex achæni (calycis tubus) nunc à pericarpio vero subseparabilis, nunc arcuè adhærens.

CCXCIX. *PHILACTIS* Schrad. ind. sem. hort. Gætt. 1831 et diss. mss. cum icon. 1832.

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis circ. 8-10 uniser., disci tubulosis 5-dentatis hermaphroditis sterilibus. Invol. campanulatum imbricatum, squamis biserialibus adpressis. Recept. elongato-conicum scrobiculatum, paleis oblongis acutis complicatis. Cor. radii cum ovario continuæ oblongæ amplæ obtusæ 5-dentatæ, disci extùs parcè glandulosæ. Anth. disci ecaudatæ. Styli radii exserti bipartiti, disci sæpius inclusi, ramis cono brevi superatis. Achænia radii trigona obpyramidata fertilia glabra coronata, extùs ligulâ persistente, intùs aristâ crassâ acutâ subciliatâ, disci compresso-subtetragona ex sicco abortiva subhirtella coronata aristis 4 cruciatim inæqualibus scabris imâ basi subconcretis ex angulis ortis. — Herba basi suffruticosa Mexicana dichotoma. Rami teretes, juniores villosi. Folia opposita petiolata latè ovata acuminata triplinervia serrulata utrinquè adpressè villosa. Pedicelli inter dichotomias orti 1-cephali foliis breviores. Ligulæ flavæ demùm subrufæ. Discus luteus. — Genus inter *Tragoceratem* et *Zinniam* medium.

1. *P. ZINNIODES* (Schrad.! l. c.) 2. 5 in Mexico. Petioli 4-lin. Limbus fol. 2 poll. longus, sesquipollic. latus. (v. s. comm. à cl. Schrad.)

CCC. *ZINNIA* Linn. gen. n. 974. Gærtn. fr. 2. p. 459. t. 172. Cass. dict. 59. p. 315. Don trans. lin. soc. 16. p. 2. p. 155. Less. syn. 224. — *Crassina* Scop. — *Lejica* Hill.

Capitulum radiatum, fl. radii foemineis ligulatis, disci tubulosis 5-dentatis hermaphr. Invol. imbricatum, squamis ovato-rotundis nigro marginatis. Recept. conicum cylindraceumve, paleis oblongis complicatis disci flores involventibus. Ligulæ coriaceæ reticulatæ obovatæ aut oblongæ cum achæni apice continuæ persistentes. Cor. disci super ovarium articulatæ, supernè pilis corollinis velutina. Achænia corticata

membranacea subalata, radii subtrigona calva, disci ext. subtrigona, inter. compresso-plana, apice ex angulo interiore uniaristata aut biaristata (aristâ inter. longiore), aut bidentata seu mutica. Cotyled. in achæniis trigonis involucro parallelæ, in achæniis compressis invol. contrariæ. — Herbæ Americanæ annuæ, ramis foliisque oppositis. Folia 3-5-nervia integra. Capitula ramos terminantia solitaria, elegantia, diù persistentia. — Cl. Cassini dubitat an ligula sit verus pappus radii et cl. Don hanc sententiam confirmat.

§. 1. *Achæniis disci ad latus interioris uniaristatis aut rariis subbinis valdè inæqualibus, paleis receptaculi integerrimis obtusis.*

1. *Z. TENUIFLORA* (Jacq. ic. rar. 3. t. 590. coll. 5. p. 153), caule erecto vix pubescente, foliis brevissimè subpetiolatis cordato-lanceolatis, pedunculis longissimis cylindricis, invol. oblongi squamis adpressis vix marginatis, ligulis lineari-oblongis sæpius apice revolutis, recept. paleis integerrimis, achæniis disci uniaristatis. (I) in Mexico (Cav.), ad Tlapujahua (Keerl.!) prope Cinapequara, Valladolid, Guanajuato, etc. (H. et B.). Sims bot. mag. t. 555. *Z. revoluta* Cav. ic. 3. t. 251. Ligulæ suprâ coccineæ, 7-8 lin. longæ, vix 2 lin. latæ, integræ aut apice bidentatæ. (v. v. c. et s. sp.)
2. *Z. BICUSPIS* (ic. fl. mex. ined.), caule erecto hirtello, foliis breviter petiolatis lanceolatis villosis, pedunculis folio brevioribus cylindricis, invol. ovalis squamis adpressis, recept. paleis integerrimis, ligulis lineari-lanceolatis apice acutè bifidis, achæniis disci uniaristatis. (I) in Mexico. An var. *Z. tenuifloræ*, sed ligulis suprâ sordidè albido-virescentibus et notis suprâ indicatis satis distincta videtur. (v. ic. pict.)
3. *Z. LEPTOPODA*, caule erecto confertè pubescente, foliis sessilibus ovali-oblongis, pedunculis cylindricis folio æqualibus longioribusve, invol. obovatis squamis adpressis, ligulis obovatis emarginatis, recept. paleis integerrimis obtusis, achæniis disci 1-aristatis. (I) in Mexico, ad Tlacolola in Oaxacanâ ditione (Andr. fl. exs. n. 314), et in montibus circa urb. Mex. (Berl.!) pl. exs. n. 686). *Z. pauciflora* in fl. mex. ined. non Linn. Affinis *Z. multifloræ* sed pedunculis nec cavis nec incrassatis nec striatis et capitulo minore differt. Ligulæ suprâ purpureæ aut rubræ. (v. s.)
4. *Z. MULTIFLORA* (Linn. sp. 1269. Linn. f. dec. 23. t. 12), caule erecto ramoso subhirtello, foliis vix subpetiolatis ovato-lanceolatis, pedunculis folia superantibus apice (præsertim centrali) obconicis cavis striatis, invol. campanulati squamis adpressis, ligulis obovatis obtusis emarginatisve, recept. paleis integerrimis obtusis, achæniis disci 1-aristatis. (I) in Mexici pratis juxta Valladolid et Pazcuaro (H. et B.), in Louisianâ (Willd.), in Brasiliâ circa Bahiam (Blanch.!) n. 191, ad littora maritima ins. Mauritiî (Met.!), sed verisim. culta aut introducta. An & *Z. pauciflorâ* satis distincta?
a. rubra, ligulis supernè rubris coccineisve. Jacq. obs. 3. p. 40. (v. v. c.)
β. flava, ligulis supernè flavis luteisve. H. B. et Kunth nov. gen. am. 4. p. 250. (v. s. in Carol. introd. ex Bosc.)
5. *Z. VERTICILLATA* (Andr. bot. rep. 3. t. 189), caule erecto sparsè hirsuto, foliis oblongo-lanceolatis, nunc irregulariter verticillato-congestis, nunc spiraliter subdigestis, pedunculis brevibus obconicis cavis striatis, invol. campanulati squamis adpressis, ligulis obovatis apice emarginatis sæpius 2-3-seriatis, paleis recept. integerrimis subacutis, achæniis disci 1-aristatis. (I) in Mexico. *Z. hybrida* olim in hort. Par. *Z. verticillata* Schkuhr handb. 3. t. 252 evidenter monastrum sed verisim. speciei nondum in statu naturali descriptæ. Caules angulato-sulcati sæpè quasi fasciati. Capitula fortè è pluribus facta. Fl. disci 5-10-andri, styli 3-5-fidi. Conf. Don l. c. (v. v. c.)
6. *Z. PAUCIFLORA* (Linn. sp. 1269), caule erecto hirsuto, foliis sessilibus cordato-lanceolatis subamplexicaulibus, pedunculo terminali striato obconico cavo

brevi, lateralibus longioribus tenuioribus, invol. squamis adpressis, recept. paleis integris, achæniis subbiaristatis, aristâ ext. brevissimâ aut nullâ. ① in Peruvia. Lam. ill. t. 685. f. 1. Rudbecka fol. oppos. Zinn. h. Gætt. 409. t. 1. Chrysogonum Peruvianum Lin. sp. ed. 1. p. 920 excl. syn. Zinnia lutea Gærtn. fr. 2. p. 459. t. 172. Bidens Mill. dict. t. 64. Lejica Hill. exot. 29. Crassina Scop. diss. Ligulæ flavæ obovatæ apice obtusæ aut 1-2-dentatæ. (v. v.)

§. 2. *Achæniis disci apice biaristatis bidentatisve aut muticis, recept. paleis integerrimis acutis.*

7. *Z. HYBRIDA* (Sims bot. mag. t. 2123), caule erecto pubescente, foliis cordatis sublancoelatis sessilibus amplexicaulibus, pedunculo terminali solitario brevi obconico cavo, invol. squamis adpressis, recept. paleis acutis integris, achæniis bidentato-aristatis. ① in Amer. merid. ex h. Pavon. *Z. grandiflora* hortul. Affinis *Z. paucifloræ* sed differt caule pubescente nec hirsuto, fl. disci obscure purpureis, ligulis suprà obscure-coccineis nec flavis, pedunculo non striato. Si nomen justo impositum, hybrida ex *Z. pauciflorâ* et *Z. elegante*? (v. v. c.)

8. *Z. ANGUSTIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 251), caule erecto pubescente sparse hirtello, foliis sessilibus lanceolatis hispido-scabris, pedunculis folia subæquantibus cylindricis, invol. squamis squarrosis, ligulis ellipticis, recept. paleis integerrimis angustis acutis, achæniis disci emarginato-bidentatis muticis lævibus ciliatis. ① in Mexico prope Guanajuato. Ligulæ suprà flavæ; discus valde convexus, ligulis paulò intensius flavus. (v. s. sp. comm. à cl. Mendez et Alaman.)

9. *Z. MARITIMA* (H. B. et Kunth nov. gen. am. 4. p. 251), caule procumbente ramosissimo lævi apice subhispidulo, foliis breviter petiolatis oblongis obtusis hispido-scabris, pedunculis striatis superne incrassatis, invol. hemisphærici squamis adpressis, recept. paleis lanceolatis integerrimis, achæniis disci apice bidentatis. ① in arenosis maritimis prope Acapulco.

§. 3. *Achæniis disci muticis aut subbidentatis, recept. paleis acuminatis apice serrato-cristatis.*

10. *Z. ELEGANS* (Jacq. coll. 3. p. 152), caule erecto hirtio, foliis sessilibus amplexicaulibus cordato-ovatis, pedunculis solitariis folio longioribus cylindraceis, invol. squamis ovatis obtusis, superioribus nigro marginatis, recept. paleis serrato-cristatis, achæniis disci muticis vix subbidentatis, ligulis obovato-subrotundis. ① in Mexico (Cav. Karv.), ad montes Tixtlæ (fl. mex. ined.), circa Cuernavaca (Berl. n. 975). Planta in hortis seminibus varians, an fortè in species duas ex involucrio lævi aut squarroso divellenda?

A. invol. squamis adpressis aut intimis vix subsquarrosis.

a. *violacea*, palearum apicibus ligulisque suprà purpureo-violaceis. *Z. violacea* Cav. ic. 1. p. 57. t. 81. Andr. bot. rep. 1. t. 55. *Z. elegans* Jacq. ic. rar. 3. t. 589. Sims bot. mag. t. 527. (v. v. c.)

β. *alba*, palearum apicibus ligulisque suprà albis. (v. v. c.)

B. invol. squamis omnibus squarrosis.

γ. *purpurascens*, palearum apicibus ligulisque suprà purpureo-violaceis. (v. v. c.)

δ. *coccinea*, palearum apicibus ligulisque suprà splendide coccineis. — *Z. violacea coccinea* Lindl. bot. reg. t. 1294. (v. v. c. et s. sp.)

† *Species exclusa.*

7. *BIDENS* Retz. = *Glossogyne pinnatifida*.

CCCI. *HELICTA* Cass. bull. philom. nov. 1818. dict. 20. p. 461. 46. p. 399 et 406 non Less.

Capitulum multiflorum heterogamum, fl. radii ligulatis 1-seriatis fœmineis, disci hermaphr. tubulosis 5-dentatis. Invol. campanulatum biseriale disco longius, squamis ext. 5 spathulatis apice foliaceis, inter. minoribus oblongis. Recept. convexum, paleis amplectentibus. Ligulæ ad basin usquæ fissæ nec basi tubulosæ. Cor. disci tubo nullo et ideò stam. sublibera. Antheræ nigricantes in appendice terminali tubercula grossa gerentes. Achænia conformia erostria compressa obovata oblonga, supernè hispidula, margine calloso utrinquè cincta. Pappus sessilis coroniformis brevis cartilagineus dentatus. — Frutex (patriâ ignotâ) sarmentosus, ramis teretibus junioribus hispidulis. Folia opposita subconnata ovalia obtusa remotè subdentata crassiuscula lucida. Pedicelli solitarii terminales et axillares. Flores lutei.

1. *H. SARMENTOSA* (Cass. l. c.). ♂ Patr. ign. diù in hort. Par. sub nomine *Verbesinæ muticæ* culta.

† *Species exclusæ.*

H. ALTERNIFOLIA Less. = *Epallage salicifolia*.

H. MADAGASCARIENSIS Less. = *Epallage dentata*.

CCCL. *ALARÇONIA*.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis 1-seriatis, disci tubulosis 5-dentatis hermaphr. Invol. 2-3-seriale campanulatum, squamis laxis foliaceis hirsutis oblongis disco æqualibus longioribusve, intimis minoribus paleas referentibus. Recept. planum, paleis flores subæquantibus et semiamplect. Ligulæ amplæ apice 3-dentatæ. Cor. disci tubo brevi coriaceo, fauce cylindricâ longâ 5-nerviâ, dentibus vix extûs apice subbarbatis. Styli rami fl. ♀ glabri, fl. ♂ longi densissimè hispidi acuti, exserti revoluti. Achænia prismatica crassa elongata, pappo calyciformi vario. — Herbæ perennes villosæ Californicæ Inulæ Helenii faciem ferentes. Caulis teres simplex hirsutus. Folia alterna oblonga aut elliptico-oblonga, inferiora basi attenuata, super. semiamplexicaulia, integerrima. Capitula ampla solitaria terminalia. Flores flavi. Genus habitu spectabile. An *Wedeliæ*, an *Helenicis* affine? — Dixi in honorem peregrinatoris Ferdin. Alarçon, qui primus aut inter primos Californiam adiit anno 1540.

1. *A. HELENIODES*, caule folisque molliter velutinis, radicalibus petiolatis amplis elliptico-oblongis utrinquè acutis, invol. squamis disco longioribus, achænio glabro dentibus 5 lanceolatis acutis sericeo-ciliatis crassis coronato. ♀ in Californiâ legit cl. Douglas. Folia radicalia pedem longa, 2½ pollic. lata. Ligulæ ferè sesquipollic. (v. s. comm. ab hon. soc. hort. London.)

2. *A. ANGUSTIFOLIA*, caule hirsuto, foliis oblongo-lanceolatis, infer. subglabris, super. vix subhirsutis, invol. squamis disco æqualibus longè hirsutis, achænio apice villoso, nunc truncato, nunc in aristam setosam scabram elongatam producto. ♀ in Californiâ legit cl. Douglas. Recepti mixtam cum præcedente cui habitu sat similibus, sed caract. carpici adeò dissimiles ut facile ex hac specie genus constituendum esset. (v. s. comm. cum priore.)

CCCHII. WEDELIA *Jacq. amer.* 217. t. 130. *Rich. in Pers. ench.* 2. p. 490 non *Adans.* — Wedelia, Stemmodontia et verisim. *Trichostemma Cass. dict.* 46. p. 407-409. — *Wedeliæ sp. Less. syn.* 222.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis foemineis, disci hermaphr. tubulosis 5-dentatis. Invol. 2-3-seriale, squamis ext. foliaceis, intimis membranaceis. Recept. subconvexum paleaceum. Styli fl. hermaphr. rami cono superati. Achænia obovata aut compressa erostria, superata pappo coroni- seu calyciformi sæpius substipitato squamellis concretis ciliatisque constante. — Suffrutices aut herbæ. Species pleræque Americænæ, rariùs Asiaticæ aut Australasicæ. Folia opposita breviter petiolata serrata rariùs trifida. Pedicelli terminales sæpius solitarii, sæpè ex dichotomiis ramorum, rariùs ex axillis foliorum, orti. Corollæ luteæ.

Sect. I. CYATHOPHORA.

Achænia omnia exalata, apice calyculum dentatum gerentia, nec in tubercula callosa producta.

* *Herbaceæ, foliis ad medium trilobis aut subtrilobis.*

1. W. CARNOSA (*Rich. in Pers. ench.* 2. p. 490), caule herbaceo repente glabro, foliis sessilibus subcarnosis obovatis basi cuneatis subtrilobis repando-subdentatis glabris, pedunculis axillaribus 1-cephalis, folio brevioribus aut paulò longioribus adpresso-villosis, invol. squamis biserialis, ext. foliaceis oblongis obtusis disco æqualibus dorso subpuberis, int. minoribus membranaceis glabris, pappo coroniformi irregul. denticulato. 2. in maritimis insul. Caribæarum, ad Sanctum-Domingum (*Poir.*), Jamaicam (*Sloan.*), etc. — *Cass. dict.* 46. p. 409. *Plum. ed. Burm.* t. 107. f. 2. *Silphium trilobatum Linn. sp.* 1302. *Buphthalmum repens Lam. dict.* 1. p. 515. *Sloan. jam.* t. 155. f. 1. Ligulæ 10-12 apice dentatæ. (v. s.)

β. *triloba* (*Rich. l. c.*), parçè pilosa, foliis manifestè trilobis. — Habeo ex ins. Cubæ fol. lobis ferè acuminatis et ex Guadalupâ fol. lobis obtusis. An diversæ? (v. s.)

γ. *aspera* (*Rich. l. c.*), foliis asperiusculis obsolete trifidis. — in Guadalupâ legit Bertero. (v. s.)

2. W. PALUDOSA, caule herbaceo adscendente? glabro subanguloso, foliis subpuberulis sessilibus oblongo-lanceolatis, basi angustatis obtusis, apice acuminatis, longè supra basin triplinerviis remotè serratis, serraturis duabus mediis longioribus, pedicellis axillaribus folio triplò longioribus pubescentibus, invol. squamis oblongis subacutis foliaceis disco paulò longioribus, intimis paleisque brevioribus scariosis, achæniis elongati pappo calyciformi denticulato. ① ad margines paludum in Brasiliâ circa Rio-de-Janeiro legit cl. Lund. Flores aurei. Ligulæ apice acutè 2-3-dentatæ. (v. s.)

β. *vialis*, caule hinc indè pilosinaculo magis demisso, foliis breviter et acutè trilobis, pedicellis villosis, invol. squamis paulò latioribus ciliatis. — ad viarum margines circa Bahiam legit cl. Salzmänn. An species propria, an eadem in solo sicciore nata? (v. s.)

γ. *latifolia*, caule glabriusculo, foliis latè ovatis profundè trilobis subserratis. — circa Bahiam legit cl. Blanchet. Hinc fortè pertinet *Acmella Brasiliensis Spreng. syst.* 3. p. 592, sive *Buphthalmum strigosum Spreng. neu. entd.* 2. p. 140, sed nihil asserere queo cum *Acmella Spreng.* septem speciebus ad septem genera diversa referendis constet. (v. s.)

3. W. GRACILIS (*Rich. in Pers. ench.* 2. p. 490), caule herbaceo repente subfiliformi, foliis anguloso-lobatis pubescentibus membranaceis, achæniis turgidis subtrigonis muricatis. — in Antiquâ et Sancto-Domingo, in siccis et montosis. An ex ipso auct. var. W. carnosæ?

** Herbaceæ, foliis serratis.

4. *W. HISPIDA* (H. B. et Kunth nov. gen. am. 4. p. 215. t. 71), herbacea, ramis erectis hispidis, foliis lanceolatis aut subovato-lanceolatis basi cuneatis apice acuminatis medio inciso-dentatis subtriplinerviis utrinquè hirtis, pedunculis elongatis 1-cephalis è dichotomiis ortis, invol. squamis ext. oblongo-lanceolatis hirtis disco æqualibus, int. glabrinusculis subæqualibus, achænio compresso-tetragono, pappo coronif. potius calyculiformi ciliato denticulato. $\frac{1}{2}$? in Mexici valle Tenochitl (H. et B.), Ker bot. reg. t. 543. Desf. cat. h. par. ed. 3. p. 185. *W. calendulacea* Rich. in Pers. ench. 2. p. 490 non Less. *Stemmodontia scaberrima* Cass. dict. 46. p. 407. *Bupthalmum scabrum* Cav. ic. 2. p. 53. t. 167 vix bona, sed sub hoc nomine semina et specim. ex hort. Madr. habui. An fortè *Bupthalmum asperum* Horn. h. Hafn. suppl. p. 102? Flores suberocci ad quod alludit nomen Rich. Achænia sparse puberula maculata utrinquè ad basin disco rufo calloso ornata. Pappus hinc indè in setulam productus. (v. s. et v.)
5. *W. MACRODONTA*, caule herbaceo ramoso tortuoso hispido, foliis sessilibus ovato-lanceolatis acutis grossè dentatis utrinquè hispidis, pedicellis è dichotomiis et apicibus ortis 1-cephalis hispidissimis aphyllis, invol. squamis foliaceis scabris acutis disco circ. æqualibus, ligulis plurimis vix invol. longioribus. $\frac{1}{2}$ in campis editis prov. Sancti-Pauli Brasilie ad Taubalé, novemb. flor. legit cl. Lund. Capitula multiflora, ligulis 15-20. (v. s. comm. à cl. Lund.)
6. *W. LUNDII*, caule herbaceo erecto trichotomo gracili setis hispido, foliis sessilibus lanceolatis utrinquè acuminatis utrinquè hispidis serraturis paucis minimis apice donatis, pedicellis è dichotomiis ortis elongatis aphyllis 1-cephalis, invol. squamis ext. foliaceis disco longioribus, pappo calyciformi ciliato subaristulato. $\frac{1}{2}$ in campis editis prov. Sancti-Pauli, ad Mugi flum. legit cl. Lund. Affinis *W. hispide*, sed habitu multò graciliori et foliis vix subserratis præcipuè differt. Ligulæ apice dentatæ 8-9 lin. longæ. (v. s. comm. à cl. Lund.)
7. *W. VAUTHIERI*, caule subherbaceo erecto villis hirtello subsimplici, foliis ovalibus sessilibus acutiusculis basi trinerviis subtus reticulatis utrinquè sparse setuloso-scabris, capitulo terminali solitario pedicello hirsuto instructo, invol. squamis ovalibus obtusis dorso scabris disci circ. longitudine. $\frac{1}{2}$ in Brasiliæ prov. Minarum-Generalium legit cl. Vauthier (pl. exs. n. 319!). Affinis hinc *W. Lundii*, illinc *W. radiosæ*. Rhizoma fruticosum. Ligulæ 8-9 lin. longæ, apice dentatæ, invol. duplò et ultrà longiores. (v. s. comm. à cl. Vauthier.)
8. *W. AFRICANA* (Pers. ench. 2. p. 490), herbacea, foliis breviter petiolatis lanceolatis serratis acuminatis trinerviis cauleque pubescentibus, invol. squamis ext. obtusis dorso puberis, int. subacutis glabrinusculis paulò longioribus, ligulis circ. 20 apice bidentatis, achæniis calyculo subsessili denticulato. $\frac{1}{2}$ in Africæ æquinoctialis regno Owariensi. Beauv. fl. owar. 2. p. 18. t. 69.
9. *W. CALENDULACEA* (Less. syn. p. 222 non Rich.), foliis oblongo-lanceolatis basi attenuatis utrinquè strigoso-pilosis apice rariter serratis, pedunculis 1-cephalis solitariis axillaribus folio triplò longioribus, invol. squamis ext. oblongis subacutis disco longioribus, achæniis calyculo substipitato denticulato. — in Indiâ orient. pascuis humidis frequens, in Silhet, Bengalo, agro Maderaspatano, Zeylonâ, etc. DC. in Wight contrib. p. 17. *Verbesina calendulacea* Linn. sp. 1222. Willd. sp. 3. p. 2226. Wall. cat. et herb. n. 315. Lam. ill. t. 686. f. 1. *W. Benghalensis* Rich. in Pers. ench. 2. p. 10? *Jageria calendulacea* Spreng. syst. 3. p. 500. — Burm. zeyl. 52. t. 22. f. 1. Rheed. mal. 10. t. 42? mal. Variat foliis obtusis aut sæpius acuminatis, integerrimis plus minùs serratis aut incisissimis, imò rariùs, ferè trilobis. Achænia crassa obtusè tetragona rugoso-muricata margine 4-5-dentato crasso obtuso persistente coronata. (v. s.)
10. *W. URTICÆFOLIA* (DC. in Wight contrib. p. 18), herbacea subscandens, foliis petiolatis ovato-lanceolatis grossè et subinæqualiter serratis utrinquè strigosiuspidis basi brevè apice longè acuminatis et sæpè incurvis, pedunculis solitariis 1-cephalis, invol. squamis 10 biseriatis acuminatis dorso pubero-strigosis, recept.

paleis acuminatissimis, achæniis calyculo brevè denticulato superatis. — in Indiâ orient. Fortè species 2 seu 3 distinguendæ, nempe:

- a. *Wallichii*, pedunculis folio brevioribus, foliis utrinquè longè acuminatis membranaceis parcius involucriisque strigosis. — in Nepaliâ. Weddellia Wallichii Less. in Kunnâ 1831. p. 162. (v. s. comp. à cl. Wallich. anno 1831.)
 - β. *Wightii*, pedunculis folii circ. longitudinem æquantibus, foliis basi breviter acuminatis parè strigosis. — in collibus Indiæ australis, ad Dendigul. *Verbesina urticifolia* Wight! herb. *Verbesina biflora* var. Wall! cat. n. 317. litt. B. et fortè D. (v. s.)
 - γ. *scaberrima*, pedunculis folio longioribus, foliis breviter basi acuminatis magis ovatis utrinquè caulibus involucriisque scaberrimis. — in regno Ava, ad Prome et in ins. Javâ (Bl.). Culta flore pleno evadit. *Verbesina urticæfolia* Blume! hijdr. gr. 1. *Verbesina scaberrima* Wall! cat. n. 311. An species propria? (v. s.)
11. *W. CUNNINGHAMI*, herbacea erecta, caule subasperulo, foliis petiolatis oblongo-lanceolatis basi acutis apice acuminatis serrulatis utrinquè puberulis, pedicellis axillaribus terminalibusque 1-cephalis folio brevioribus, invol. squamis 6 biserialis, ext. lanceolatis acutis longioribus puberis, achæniis obcompressis calyculo brevè denticulato superatis. — in rupestribus insularum Goulburn ad oram borealem Australiæ mart. flor. legit cl. A. Cunningham. Affinis *W. urticæfoliæ* var. sed involucri oligophyllis foliisque multò angustioribus facillè differt. (v. s. comm. à cl. inv.)
 12. *W. TRICHOSTEPHIA*, caule herbaceo erecto tereti hispido, foliis breviter petiolatis ovali-oblongis remotè serratis triplinerviis utrinquè hispidis, pedicellis solitariis terminalibus villosis, invol. squamis ovato-lanceolatis foliaceis disco paulò longioribus, achæniis tetragonis pilorum zonâ superatis. — in Brasiliâ. *Trichostemma hispidum* Cass. dict. 46. p. 409. *Trichostephus* Cass. dict. 60. p. 618. *Trichostephium* Cass. opusc. 3. p. 59.
 13. *W. MOLLIS* (Rich. in Pers ench. 2. p. 490), herbacea, foliis petiolatis cordato-lanceolatis crenatis membranaceis subtus molliter pubescentibus. 2/ in Americâ merid. Caulis pubescens. Folia obsoletè 2?-nervia, sed imprimis subtus valdè venosa. †.
 14. *W. SUBVELUTINA*, caule herbaceo erecto tereti villosa, foliis vix petiolatis angustè lanceolatis basi obtusis apice acuminatis parvè et remotè serrulatis, supra ad basin triplinerviis, subtus reticulato-venosis velutino-villosis, supra scabridis demum glabris, pedicellis solitariis brevissimis, invol. squamis ovato-lanceolatis, achæniis calyculo minimo denticulato-ciliato. 2/ in Brasiliæ prov. Sancti-Pauli (h. imp. Bras. n. 408!), in paludosis (Lund!). Folia 3 poll. longa, 5 lin. lata, petiolo 2 lin. longo. (v. s.)
 15. *W. PUPERULA*, caule herbaceo erecto tereti striato apice puberulo, foliis sessilibus ovatis vix acutiusculis dentatis utrinquè tenuè puberulis subtus propè basin triplinerviis nervoso-reticulatis, pedicellis terminalibus ternis vix capituli longitudine, invol. squamis ovatis disco brevioribus, achænio subpubente calyculo multidentato coronato. 2/ in Brasiliæ prov. Minarum Generalium ad Sabarra legit cl. Vauthier (pl. exs. n. 327!). Folia sesquipollicem longa, ferè pollicem lata. Ligulæ ovali-oblongæ obtusæ. (v. s.)
 16. *W. EHREBERGII* (Schlecht. in Linnæ 1831. p. 729), caule herbaceo tereti strigoso-piloso inâ basi repente, foliis petiolulatis triplinerviis oblongo-ellipticis acutis serrato-crenatis hirsuto-hispidis suprâ asperrimis, capitalis pedicellatis terminalibus, invol. 10-phylli squamis ext. apiculatis extus hirsutis, paleis ciliatis, ligulis circ. 12 truncatis sub-3-crenatis, achænio obovato-tetragono, pappo brevissimo coroniformi. — in Sancto-Domingo ad urbem Cap Français.
- *** *Fruticosa aut suffruticosa, foliis serratis dentatisve.*
17. *W. RETICULATA*, caule suffruticuloso sulcato adpressè puberulo subaspero, foliis breviter petiolatis ovato-lanceolatis basi obtusis apice acuminatis cæser-

ruulatis rigidulis suprà strigilloso-asperis subtus velutino-pubescentibus subtriplinerviis nervoso-reticulatis, pedicellis folio brevioribus solitariis adpressè villosis, invol. squamis ovalibus disco subbrevioribus, ext. dorso puberulis, achæni calyculo villosulo, radii margine ciliato. \S in Porto-Ricco legit cl. Bertero, et verisimiliter in aliis insul. Caribæis. W. Jacquin Rich. in Pers. ench. 2. p. 490. excl. syn. W. frutescens Spreng. l. in h. Balb. non Jacq. Ligulæ oblongo-lineares obtusæ invol. duplò longiores. (v. s.)

18. W. RADIOSA (Ker bot. reg. t. 610), caule basi suffruticnoso tereti asperulo, foliis petiolatis ovato-sen oblongo-lanceolatis acutis serrulatis utrinquè puberulis asperulis, pedunculis è dichotomià et apice ramorum 1-3 monocephalis folio longioribus, invol. squamis 3-seriatis, ext. ovatis puberis ciliatis foliaceis, intimis oblongo-linearibus membranaceis, pappo coroniformi ciliato-serrulato. \S in Brasiliâ. Ligulæ flavæ obtusæ vix 3-dentatæ invol. duplò longiores. Achænia juniora obcompressa disci ciliata, radii hirsutula, matura mihi ignota. (v. s. ex hort. Angl.)

19. W. CARACASANA, caule suffruticoso tereti subscabrido, ramulis pedicellis petiolisque hispidulis, foliis petiolatis ovato-lanceolatis basi cuneatis apice longè acuminatis serratis triplinerviis utrinquè sparsè strigosis subtus pallidioribus, pedicellis terminalibus 1-cephalis subternis, invol. squamis ext. ovali-lanceolatis disco longioribus dorso scabris, ligulis elongatis subintegris, achæni pubescentis calyculo serrulato-ciliato. \S in Amer. austr. propè Caracas legit cl. Vargas. Folia ferè W. acuminatæ, sed distinctius triplinervia, pedunculi terni longiores. Cor. intensè luteæ. (v. s.)

20. W. ACUMINATA, caule fruticnoso, ramis teretibus pedunculisque strigoso-scabris, foliis petiolatis ovato-lanceolatis basi cuneatis apice longè acuminatis serratis utrinquè strigilloso-asperis subtriplinerviis, pedicellis solitariis 1-cephalis folio brevioribus, invol. squamis ext. lanceolatis acuminatis strigillosis radium subæquantibus, ligulis angustis acutè bidentatis, achæni pubescentis calyculo substipitato denticulato. \S in ins. Cubâ juxta Havannam legit cl. Ant. de la Ossa. Folia 3 poll. longa, 12-15 lin. lata, interdum basi inæqualia. Petiolus junior ciliatus. (v. s.)

21. W. FRUTESCENS (Jacq. amer. p. 217. t. 130), caule fruticoso scandente glabro, ramis teretibus, foliis breviter petiolatis ovatis acutis leviter serratis utrinquè scabris, pedicellis terminalibus solitariis vix capituli longitudine, invol. squamis ext. subovatis laxis, ligulis obcordatis semibilibis, achæni obovati calyculo 10-dentato. \S in fruticetis et sylvaticis circa Carthagenam. Polymnia Wedelia Linn. mant. 118.

22. W. AMBIGUA, caule fruticoso tereti glabro, ramulis petiolisque hispidissimis, foliis breviter petiolatis ellipticis utrinquè acuminatis subserratis, suprà sparsè strigillosis, subtus secus nervos pubero-scabris, vix triplinerviis, pedicellis 1-cephalis hispidis folio paulò brevioribus, invol. squamis ext. lanceolatis scabris, int. paulò longioribus ovatis obtusissimis glabris, ligulis acutè bidentatis, achæni calyculo subdenticulato. \S in Guadalupâ legit cl. Bertero. W. calycina Spreng. l. syst. 3. p. 580. ex ipso in h. Balb. sed excl. syn. etiã Sprengeliano! Achæni calyculus nigricans. (v. s.)

23. W. LANCEOLATA, caule fruticoso tereti nervulis elevatis striato, ramis oppositis piloso-scabris, foliis breviter petiolatis lanceolatis basi brevè cuneatis apice acuminatis serrulatis suprà strigilloso-scabris subtus puberulis triplinerviis, pedicellis solitariis hispidulis folio paulò longioribus, invol. squamis ext. foliaceis lanceolatis acutis, int. obtusis subbrevioribus ovali-oblongis glabris, achænio radii pubescente calyculo substipitato inæqualiter denticulato. \S in Porto-Ricco legit cl. Ledru. Ligulæ invol. paulò longiores oblongo-lineares obtusæ vix subdentatæ. Folia supernè ad orig. pilorum albo-punctata. Affinis W. frutescenti, sed invol. squamæ acuminatæ nec obtusæ. (v. s.)

24. W. AFFINIS, caule fruticoso dichotomo tereti glabro, ramulis hispidulis, foliis breviter petiolatis lanceolatis basi cuneatis apice acuminatis subserrulatis

- utrinquè parè puberulis vix subtriplinerviis, pedicellis solitariis 1-cephalis hispidulis vix folio longioribus, invol. squamis foliaceis lineari-lanceolatis disco paulò longioribus, ligulis acutè bifidis vix invol. longioribus, achæniis subpuberuli calyculo substipitato vix subdentato. ♀ in insulâ Sancti-Thomasii legit cl. Wydler (pl. exs. n. 51). W. calycina Rich. in Pers. ench. 2. p. 490? Valde affinis W. lanceolata. (v. s.)
25. W. *PULCHELLA* (H. B. et Kunth nov. gen. am. 4. p. 214), caule fruticoso procumbente, ramis teretibus adpressè pilosis, foliis petiolatis ovatis acutis basi cuneatis serratis triplinerviis utrinquè piloso-strigosis, pedicellis 1-2 terminalibus elongatis pubescentibus, invol. squamis ext. obovatis aut subrotundo-ovatis acutis adpressè hispidulis, inter. glabris, ligulis circ. 7 bifidis, achæniis pubescentis calyculo fimbriato. ♀ in humidis propè Cumana ubi dicitur *Mirasol*.
26. W. *ACAPULENSIS* (H. B. et Kunth nov. gen. am. 4. p. 215), caule fruticoso procumbente, ramis striatis junioribus piloso-hispidis, foliis petiolatis ovato-oblongis acuminatis basi acutis remotè serrulatis triplinerviis suprà hispidoscabris subtùs piloso-strigosis, pedicellis solitariis 1-cephalis folio brevioribus pilosiusculis, invol. squamis ext. lanceolatis aut ovato-lanceolatis hispidis ligulis apice bifidis, achænio puberulo calyculo substipitato fimbriato-ciliato ♀ propè Acapulco Mexicanorum ad littus Oceani Pacifici (H. et B.), et in ins. Sancti-Thomasii ex Schlecht. in linnæa 1831. p. 727. An W. calycina Rich. in Pers. 2. p. 490? ex Schlecht.
27. W. *LATIFOLIA*, caule fruticoso tereti, ramis scabridis, foliis petiolatis latè ovatis basi subobtusis apice acuminatis remotè subserratis utrinquè sparsè et adpressè puberis triplinerviis, petiolis ciliatis, pedicellis 1-cephalis capituli longitudine, invol. squamis ext. ovato-lanceolatis acuminatis dorso villososcabris radium subæquantibus, ligulis latis acutè bidentatis, achænio calyculo substipitato denticulato-ciliato. ♀ in Amer. austr. Terrâ firmâ legit cl. Bertero. *Polytmia australis* Spreng. ! in h. Balb., sed in ejus syst. non reperire quivi. Cor. flavo-croceæ ! (v. s.)
28. W. *CRENATA* (Rich. in Pers. ench. 2. p. 490), suffrutescens procumbens, foliis petiolatis oblongis inæqualiter crenatis obtusis membranaceis subrugosis, ovarii glandulosis, pappo obsoleto. ♀ in Guadalupe arenosis. Ligulæ obtusæ 3-dentatæ. †.
29. W. *SERRATA* (Rich. in Pers. ench. 2. p. 490), suffruticosa, foliis longiusculè petiolatis subcoriaceis lævibus lucidis, serraturis magnis, ligulis tridentatis, achæniis calyculo margine excepto glabro. ♀ ad Sanctum-Domingum. Folia ferè illicina. †.
30. W. *CRUCIANA* (Rich. in Pers. ench. 2. p. 490), frutescens tota molliusculè villosa, foliis petiolatis ovatis acutè dentatis, pedunculis longissimis, achæniis pubentis variegati calyculo stipitato multipartito. ♀ in insulâ Sanctæ-Crucis ad sepes arborum viarumque W. Caribæa Spreng. syst. 3. p. 580. †.
31. W. *PARVIFLORA* (Rich. in Pers. ench. 2. p. 490), frutescens, foliis sessilibus ovato-lanceolatis acutissimis integris asperrimis rugosis, pedunculis brevibus, ligulis semibifidis, achæniis pubentis calyculo subintegro seu levissimo crenulato. ♀ in campestribus Guadalupe. Ex foliis sessilibus ab omnibus differre videtur. †.

Sect. II. CALLOPHORA.

Achænia omnia exalata apice producta in tubercula duo callosa obtusa inter quæ pappus brevissimus coroniformis subserrulato-ciliatus sedet.

32. W. *GAUDICHAUDI*, caule suffruticuloso tereti, ramis pedicellisque hispidis, foliis subsessilibus ellipticis utrinquè acuminatis vix apice serrulatis subtriplinerviis, suprà strigillosa, subtùs præsertim secus nervos puberulis, pedicellis elongatis 1-cephalis terminalibus seu è dichotomiis, invol. squamis biseriatis oblongis villososubhirtis disco ferè longioribus, ligulis obtusis achæniis apice

obtusè bicallosis et in centro pappum sessilem minimum subserrulato-ciliatum gerentibus. 3 in Peruvia legit cl. et am. Gaudichand. Habitus *W. Caracasanae* et *W. acuminatae* sed achænium diversissimum. An genus proprium? (v. s.)

Sect. III. ACTINOPTERA.

Achænia radii alâ angustâ non membranaceâ marginata, disci exalata, omnia in pappum cyathiformem membranaceum obliquum dentatum producta.

33. *W. HELIANTHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 216. t. 372), caule herbaceo dichotomo, ramis teretibus piloso-pubescentibus, foliis breviter petiolatis ovato-oblongis acuminatis remotè serratis triplinerviis suprà hispidulo-scabris subtùs pubescenti-pilosis, pedicellis terminalibus elongatis solitariis invol. squamis subæqualibus oblongo-lanceolatis hispidis, ligulis 15-25? bidentatis, achæniis maculatis calyculo obliquo laciniato-fimbriato coronatis. 2 in collibus siccis inter Alausi et Ticsan Quitensium. Achænia disci immarginata, radii alâ utrinquè cincta, quâ notâ à genere *Weddiæ* differt.

Sect. IV? AGLOSSA.

Capitulum homogamum discoideum. Involucrum 1-seriale 5-phyllum. Achænia ad angulos tuberculosa. — An certè congener?

34. *W. DISCOIDEA* (Schlect. in linnæâ 1831. p. 728), caule herbaceo prostrato subhirsuto, foliis oppositis petiolulatis triplinerviis ovatis obtusis serrato-crenatis scabris utrinquè hirsutis, capitulis ad apices ramorum 1-3 breviter pedunculatis discoideis, involucri 1-serialis squamis 5-ovatis obtusis, paleis acuminatis apice longè ciliatis, achæniis ad angulos tuberculatis. — in insulâ Sancti-Thomasii.

† *Species exclusæ.*

W. ARISTATA Less. = *Wollastonia strigulosa*.

W. AUREA Hook. = *Verbesina aurea*.

W. CHAMISSONIS Less. = *Wollastonia canescens*.

W. FORSTERIANA Endl. = *Wollastonia Forsteriana*.

W. MINOR Spreng. = *Melampodium divaricatum*.

W. OVATIFOLIA Willd. = *Melampodium divaricatum*.

W. PERFOLIATA Willd. = *Melampodium perfoliatum*.

CCCIV. *JÆGERIA* H. B. et Kunth nov. gen. am. 4. p. 277. t. 400. Cass. dict. 24. p. 125. Less. syn. 223. non *Ræusch*. (1)

Capitulum multiflorum radiatum, ligulis fœmineis 1-serialibus, disci hermaphr. 5-dentatis. Invol. campanulatum oligophyllum æquale, squamis florum fœmineorum basin et demùm achænia involventibus. Recept. parvum conicum, paleis carinatis diaphanis persistentibus. Styli rami recurvati. Achænia compressa 3-4-gona oblonga lævia apice calva aut pappo coroniformi minimo superata disco epigyno perexigu. — Herbæ Americanæ annuæ pilosiusculæ erectæ simplices aut oppositè ramosæ. Folia opposita subsessilia aut breviter petiolata ovalia subserrata. Capitula parva pedicellata.

1. *J. MNIOIDES* (H. B. et Kunth l. c. t. 400), caule simplici puberulo 1-2-cephalo, invol. squamis ligulisque. 5 (1) in temperatis prope Ario Mexicanorum. Flor. lutei. Achænia calva.

(1) *Jægera* *Ræusch* genus monandrum sine descript. omnino ignotum.

2. *J. PARVIFLORA*, caule subsimplici oligocephalo sparsè pubero apice hirsuto, foliis in petiolum ciliatum attenuatis subintegris suprà sparsè pilosis subtùs glabris, infer. ovalibus sublanceolatis, ligulis paucis minimis. ① in Brasiliæ prov. Sancti-Pauli. Affinis *J. hirtæ*. Fructus junior refert iconem Kunth. Ligulæ ex sicco vid. albæ aut albidæ. (v. s. in h. Mus. Par. à Mus. imp. Bras. sub n. 433 miss.)
3. *J. HIRTA* (Less. syn. 223), caule oppositè ramoso hirtò polycephalo, foliis ovalibus, invol. squamis ligulisque circ. 10 ligulis obovatis exiguis. ① in America (Lag.), in aquosis prope Jalapam (Less.), in Brasiliâ (h. Mus. Berol. I.), *Aemella hirta* Lag. nov. gen. et sp. 3o. n. 383. Less. I in linnæa 1830. p. 163. Variat 3-poll. ad 3-ped. (v. s.)
4. *J. REPENS*, caule ramoso ascendente ad nodos radicante infernè glabrato superne confertè et breviter hirsuto, foliis ovatis ovalibusve subacutis trinerviis subintegerrimis parè puberulis, ligulis circ. 10 invol. longitudine. 2/2 ? in Brasiliæ prov. Rio de Janeiro ad Serra dos Orgaos legit cl. Vauthier (pl. exs. n. 3231). Flores flavi. (v. s.)
5. *J. MACROCEPHALA* (Less. in linnæa 1834. p. 270), caule ramoso oligocephalo, foliis plerisque oblongis, ligulis oblongo-ellipticis. ① prope Jalapam Mexic. Capitula longè pedunculata solitaria minora quàm in *Erigeronte alpino*.

† *Species exclusæ.*

J. ABYSSINICA Spreng. = *Guizotia oleifera*.

J. CALENDULACEA Spreng. = *Wedelia calendulacea*.

CCCV. *LIPOTRICHE* R. Brown in trans. Linn. soc. 12. p. 118. in adn. non Less.

Capitulum multiflorum radiatum, fl. radii ligulatis 3-dentatis fœmineis 1-serialibus, disci hermaphr. tubulosis. Invol. duplici serie imbricatum subæquale. Recept. convexum, paleis foliaceis distinctis. Styli disci rami appendice acuta hispidula. Achænia subuniformia turbinata, pappo setaceo caduco uniseriali 8-10-seto denticulato coronato. — *Herbæ* in Africâ æquinociali spontaneæ. Folia opposita indivisa. Pedunculi terminales terni. Flores flavi. Genus à cl. auctore Melantheræ proximè accedit, affine etiam *Wedeliæ Ecliptæ* et *Diomedæ* dicitur.

1. L. BROWNII. — in Africâ æquin. ad ripas fluminis Congo legit infeliciss. Chr. Smith. Cæt. ign.

† *Species exclusæ.*

L. AUSTRALIS Less. = *Lipochæta lobata*.

L. GYMNOLOMOIDES Less. = *Perimenium gymmolomoides*.

CCCVI. *MELANTHERA* Rohr in Kiobek. nat. hist. selsk. 2. (1792) p. 1. p. 213. Cass. journ. phys. 1823. mai. p. 207. dict. 29. p. 483. 59. p. 158. — *Melanthera* Rich. in Michx. flor. bor. am. 2. (1803) p. 106. Brown. trans. soc. Linn. 14. p. 117. Less. syn. 222. — *Amellus* P. Brown. jam. 317 non Linn. — *Bidentis* sp. Linn. — *Calex* sp. Jacq.

Capitulum multiflorum homogamum. Invol. disco breviori squamæ biserialibus subæquales. Recept. convexum paleis flores amplectentibus persistentibus onustum. Cor. tubo brevi, limbo 5-fido, lobis suprà papillosis. Achænia compresso-subtetragona basi incrassata apice truncata. Pappi setæ 2-10 rigidulæ caducæ irregulariter dispositæ. — *Herbæ* rariùs suffrutescentes Americanæ ramosæ. Caules subtetragoni.

Folia opposita, rarius ternato-verticillata, petiolata, triplinervia, ovato-delloidea aut triloba, dentato-serrata, scabra aut subcanescentia. Capitula longè pedunculata 1-3 terminalia. Cor. albæ. Anth. nigricantes. — Nomen *Melantheræ* vetustius et constructum ex typis veteribus consecratis nempè *Melanthium*, *Melanthemum*, etc.

1. *M. HASTATA* (Rich. in Michx. l. c.), foliis oppositis hastato-trilobis scabris, invol. squamis lanceolatis, recept. paleis longè acuminatis rigidis. \mathcal{L} in siccis Carolinæ (Bosc! Fras! etc.) et in Cubâ prope Havanam (La Sagra!). Ell. sketch 2. p. 314. *Bidens nivea* Linn. sp. 1167 (excl. primò Dill. syn.)
 - a. *lobata* (Pursh fl. bor. am. 2. p. 519), foliis profundè trilobatis. — *Melanthera trilobata* Cass. dict. 29. p. 485. — Dill. elth. t. 47. f. 55. *Athanasia hastata* Walt. car. 201? (v. s.)
 - ß. *pandurata* (Pursh l. c.), foliis leviter lobatis panduræformibus. — *Melanthera panduræformis* Cass. dict. 29. p. 485. *Bidens nivea* Walt. car. 201? — Dill. elth. t. 46. f. 54. (v. s.)
2. *M. DELTOIDEA* (Rich. l. c. in adnot. sub *Melantherâ*), foliis oppositis indivisis ovato-delloideis aut subcordatis capescenti-scabris, invol. squamis ovali-oblongis, recept. paleis obtusis breviter mucronatis subfoliaceis. \mathcal{L} seu \mathfrak{J} in loco natali; sæpius ① in hortis Eur. Hab. in insulis Caribæis, Jamaicâ (Br. Sw.), Martinicâ (h. Duby!), Santo-Domingo (Bert.), Cubâ (La Sagra!), Novâ-Granatâ ad fl. Magdalenæ (H. et B.). *Melanthera urticæfolia* Cass. dict. 29. p. 484. *Melanthera* Linnæi Kunth nov. gen. 4. p. 199. *Bidens nivea* Linn. sp. 1167 var. α excl. syn. Dill. Swartz! obs. bot. 296. *Calea aspera* Jacq. ic. rar. t. 583. Amellus P. Browne jâm. 217. — Dill. elth. p. 55. t. 47. f. 55. n. 3. (v. v. c. et s. sp.)
3. *M. OXYLEPSIS*, foliis oppositis ovato-delloideis acuminatis dentatis utrinquè scabris subtus nervoso-reticulatis pallidis, involneri squamis receptaculique paleis acuminatis. \mathcal{L} ad Gnyquil legit cl. Hænke. Affinis *M. deltoideæ* sed paleis valdè acuminatis et capitulis paulò minoribus præcipuè differt. *Achænia* truncatis setis fragilibus numero variis interdum nullis (aut deciduis) coronata. (v. s. in h. Hænke.)
4. *M. CORYMBOSEA* (Spreng. syst. 3. p. 445? excl. syn.), foliis ternato-verticillatis ovato-lanceolatis acuminatis serratis utrinquè scabris subcanescentibus, capitulis pedunculatis laxè corymbosis, invol. squamis ovatis, recept. paleis linearicuneatis vix acutis. \mathfrak{J} ? in Santo-Domingo legit cl. Bertero, fortè etiàm in Guadalupe et Antiguâ ex Spreng. neu. entd. 2. p. 135 sed ipse in vol. 3. p. 40 ejusd. libri Antiguam omisit? An var. prioris? an folia constanter verticillata? Folia cum petiolo 15-16 lin. longa, 5-6 lin. lata. (v. s. ex herb. Bert. comm. à cl. Balb. qui etiàm ad cl. Spreng. misit.)

† *Species exclusæ.*

M. GLANDULOSA Wall. = *Wollastonia scabriuscula*.

M. ORIENTALIS Spreng. = *Casulia axillaris*.

CCCVII. OGIERA Cass. dict. 35. p. 445. 43. p. 371. *opusc. phyt.* 3. p. 61. — Wedeliæ sp. Less. syn. 222.

Capitulum circ. 8-florum homogamum discoideum. Invol. biserialis circ. 10-phylli squamæ ext. foliaceæ majores, inter. paleas referentes. Recept. parvum planum paleaceum. Cor. 5-dentatæ. Styli rami in appendicem hispidam producti. Achænia oblonga gracilia subtetragona apice in collum brevissimum attenuata calva. — Herbæ piloso-hispidæ Americanæ. Folia opposita petiolata ovalia subdentata triplinervia subtus plius minùs glandulosa. Capitula parva pedicellata. Cor. luteæ. — Cassinius prope Siegesbeckiam, Lessingius inter Wedeliæ collocat. Styli rami, in paucis floribus equidè visì, potiùs ad *Heliopeides* per-

tinent, sed à Wedeliâ genus facile differt capitulis discoideis et achæniis castrato calvo.

1. *O. TRIPLINERVIS* (Cass. dict. 35. p. 445), foliis utrinquè hispidis subtus glandulosis, achæniis tuberculatis, capitulis solitariis. — in Guianâ?
 - β. *leiocarpa*, adscendens, foliis utrinquè pilosiusculis, achæniis lævibus, capitulis subadgregatis. — in Guianâ gallicâ. Ogiera leiocarpa Cass. dict. 43. p. 371. Chalarium Poit. ined. (v. s. specim. à cl. Ad. Brongniart miss.)
 - γ. *Portoriccensis*, prostrata, foliis utrinquè pilosiusculis, pedicellis ad ramorum dichotomias 2-4 subadgregatis, achæniis immaturis lævibus, maturis tuberculosis. (L?) in insulâ Porto-Ricco legit cl. Bertero. Siegesbeckia Portoriccensis Bert. herb. (v. s.)

† *Species exclusæ.*

- O. TRIPLINERVIS* Bert. = *Euxenia Mitiqui*.
- O. TRIPLINERVIS* Spreng. = *Euxenia grata*.

CCCVIII. *MONACTIS* H. B. et Kunth nov. gen. am. 4. p. 286 (excl. sp. 2). Less. syn. 211.

Capitulum 5-11-florum heterogamum, fl. radii unico ligulato foemineo, disci tubulosis hermaphrod. 5-dentatis. Invol. cylindraceum subimbricatum, squamis lanceolato-linearibus. Recept. planum, paleis squamas referentibus sed carinatis. Stigma disci exsertum lobis revolutis. Achænia compressa, breviter rostrata et stipitata, calva glabra. — Arbor Aust.-Americana glabriuscula. Rami angulati subflexuosi. Folia alterna petiolata ovato-oblonga subdenticulata triplinervia. Panicula terminalis fastigiata ramosissima foliata. Flores flavi. Squamæ bracteolæque ex Less. demum spiraliter tortæ.

1. *M. FLAVERIÆ*. β in prov. Bracamorensi inter urbem Jaen et ripam fl. Amazonum. *Monactia Flaverioides* H. B. et Kunth nov. gen. am. 4. p. 286. t. 403. *Phætusa australis* Spreng. syst. 3. p. 591.

CCCIX. *WOLLASTONIA* DC. in Decaisn. nouv. ann. Mus. h. n. Par. 3. p. 414. — *Wedeliæ* sp. Less. — *Verbesinæ* sp. Blum.

Capitulum multiflorum heterogamum, fl. radii 1-seriatis ligulatis foemineis, disci hermaphr. tubulosi 5-dentatis. Invol. 2-3-seriale, squamis ext. oblongis foliaceis, internis minoribus membranaceis. Recept. planum aut subconvexum paleaceum. Cor. super ovarium articulata. Styli fl. herm. rami cono superati. Achænia crassa, obovata turbinata aut compressa, apice umbilicata, nunc nuda, sæpius aristas 1-5 tenues rigidas fragiles deciduas gerentia. — Suffrutices aut rariùs herbæ. Species ex Indiâ orient., Moluccis aut Australiâ ortæ, sæpius hispidæ, *Wedeliæ* facie. Folia opposita breviter petiolata ovata acuta triplinervia dentata. Pedicelli sæpius ad apices ramorum terni 1-cephali. Flores lutei. — Genus medium inter *Wedeliam* et *Heliopsidem*, ab utrâque achæniis structurâ et pappo diversum. — Dixi in memoriam cl. chemici anglî Wollaston de scientiis physicis et naturalibus optimè meriti et ex morum suavitate semper lugendi.

§. 1. *Wollastoniæ veræ, gerontogæ, radiatæ.*

1. *W. BIFLORA*, foliis petiolatis ovatis basi breviter apice longè acuminatis acutè serratis suprâ sparse puberulo-scabris subtus glabrisculis, pedunculis 1-3 mono-

cephalis, uno terminali, 1-2 ex axillis supremis ortis, invol. squamis hiseriatu elongato-lanceolatis, achænio calvo seu 1-aristato. — in Indiâ orient. ad Malabarâ, Martabaniam, Sangore, etc., frequens. *Verbesina biflora* Linn. sp. 1272. Willd. sp. 3. p. 226. Wall. cat. n. 317. excl. specim. paucis. *Wedelia biflora* DC. in Wight contrib. p. 18. *Acmella biflora* Spreng. syst. 3. p. 59r. Rheed. mal. 10. t. 40 bona. Invol. squamæ oblongo-lineares acutæ foliaceæ dorso puberulæ. Recept. paleæ similes. Achænia crassa obtusè 3-4-gona basi attenuata 3-4-sulcata tuberculato-rugosa, vertice lato pubero, areolâ minimâ nunc aristâ ad latus inter. instructâ nunc nudâ. Certè ad Wollastoniam, nec ad Adenostemma ut vult Lessing (p. 156), nec ad *Acmellam* ut Spreng., pertinet. (v. s. comm. ab. hon. Cæt. merc. angl. Ind. or.)

2. *W. ASPERRIMA* (Decaisne nouv. ann. Mus. h. n. Par. 3. p. 414), herbacea, ramis teretibus subtrigosis, foliis petiolatis lanceolato-attenuatis acutis serratis basi acutis trinerviis utrinquè asperimis, capitulis solitariis, invol. squamis lanceolatis, bracteis acutè acuminatis, achæniis obovatis apice truncatis angulatis tenuissimè punctulatis v. tuberculatis, pappo subuniseto. — in insulâ Timor.
3. *W. SCABRIUSCULA* (DC. in Decaisn. n. ann. mus. 3. p. 414), caule herbaceo sulcato-angulato glabresculo, foliis petiolatis ovato-lanceolatis acuminatis basi cuneatis supra basin triplinerviis grossè serratis suprâ sparè strigilloeis subtitis adpressè puberulis, corymbis terminalibus compositis trichotomis, invol. squamis biserialis ovali-oblongis acutis, achæniis crassis apice umbilicatis exaristatis. — in insulâ Timor leger. itin. Bandiniani botanici, in Indiâ orient. ad Penang, Singapore, Silhet, Rangoon cl. Wallich, et in insulis Mariannâ cl. Hænke. *Eclipta scabruscula* Wall. cat. n. 3212. comp. 322. *Melananthera glandulosa* Wall. herb. Meyera capitata Spreng. syst. 3. p. 601 ex litt. ad Wall. sed excl. syn. *Verbesina dichotoma* Heyn. herb. *Verbesina biflora* Blum. bijdr. p. 911. *W.* aristatæ et glabræ proxima, sed folia angustiora basi sensim cuneata supernè pilis delapsis albo punctata, capitula numerosiora compositè corymbosa et achænia calva, sed in 2 prioribus aristæ adèd variables ut vix earum absentiam tutò admittam. (v. s. comm. à Mus. reg. Par. et ab hon. Cur. Ind. or. angl.)
4. *W. MONTANA*, subscandens, foliis breviter petiolatis ovatis acuminatis grossè serratis utrinquè sparè striguloso-scabris, pedunculis 1-3 folio longioribus, invol. squamis 5 uniseriatis ovali-lanceolatis acutis, achæniis 1-aristatis vix angulosis. — in montanis insulæ Javæ. *Verbesina montana* Blum. bijdr. 911. Caulis vix puberulus. Paleolæ recept. acuminatæ apice serrulatæ. Achænia apice minutè puberula ferè obovata. (v. s. comm. à cl. Blum.)
5. *W. MOLUCCANA*, suffruticosa di-trichotoma, foliis breviter petiolatis ovatis acuminatis raritè serratis basi sub-5-nerviis utrinquè ramulisque scabrusculis, pedunculis confertis subumbellatis scabrido-subhirsutis folio brevioribus, invol. 4-5-phylo, squamis ovatis paleisque margine ad apicem hirsutis. 3 in insulis Moluccanis. *Verbesina Moluccana* Blum. bijdr. p. 912. (v. s. comm. à cl. Lambert.)
6. *W. ZANZIBARENSIS*, caule herbaceo erecto glabresculo, foliis petiolatis ovato-subrhombis apice acuminatis trinerviis grossè serratis utrinquè setulas sparsas adpressas gerentibus, capitulis subternis longiusculè pedicellatis, invol. squamis et recept. paleis oblongo-linearibus subfoliaceis persistentibus cor. disci brevioribus, achæniis obovatis glabris obtusis, junioribus apice hirsutulis, adultis calvis. — in ins. Zanzibar legit cl. Bojer. *Anthemiopsis macrophylla* Bojer. in litt. 1836. Styli rami inclusi apice hispidi cono acutissimo. (v. s. comm. à cl. inv.)
7. *W. CANESCENS*, herbacea scandens? hirsuto-canescens, foliis ovato seu elliptico-lanceolatis serratis acuminatis hirsuto-canescens, paniculis subcorymbosis, invol. biserialis squamis hirsutis acuminatis, achæniis muticis. — in insulâ Guahan. *Verbesina canescens* Gandieh. in Freyc. voy. p. 183. *Wedelia Chamissonis* Less. in Linnæâ 1831. p. 161.

8. *W. ELONGATA*, caule herbaceo erecto tereti ramoso glabriusculo, foliis petiolatis oblongo-lanceolatis acutis serratis utrinquè setis adpressis subscabris, pedicellis ternis folio duplò longioribus 1-cephalis, invol. squamis ovali-oblongis dorso subscabridis disco æqualibus, achæniis obovatis glabris calvis. — in insulâ Madagascar legit cl. Bojer. Anthemiopsis elongata Bojer. ! in litt. 1835. Foliolorum juga remota internodio breviora. Recept. palææ submembranaceæ. Achænia epapposa collo minutissimo terminata nec omninò truncata. (v. s. comm. à cl. Bojer.)
9. *W. STRIGULOSA* (DC. in Decaisn. nouv. ann. mus. 3. p. 414), foliis longè petiolatis trinerviis aut propè basin triplinerviis ovatis acuminatis grossè serratis, suprâ sparsè, subtus in nervis et petiolis striguloso-pubescentibus, pedunculis ad apicem ramorum 2-3 subalternis, invol. squamis biserialibus ovato-lanceolatis acuminatis dorso striatulis puberulis. — in insulis Rawak, Waigiu, Pisang, etc. Moluccarum. Verbesina strigulosa Gandich ! in Freyc. voy. bot. p. 463. Wedelia aristata Less. ! in linnæa 1831. p. 160. Bupthalthum helianthoides Forst. prod. 57. nom L'Hér. Bupthalthum australe Spreng. syst. 3. p. 605 ? ex Less. Achænia obtusè obpyramidata 3-et 4-gona obtusa subpuberula aristis 1-3 deciduis coronata. Recept. palææ obovato-oblongæ mucronatæ. (v. s. comm. à cl. Gandichand et Less.)
10. *W. GLABRATA*, caule herbaceo glabro, foliis longè petiolatis altè triplinerviis ovatis subacuminatis remotè dentatis, suprâ sparsè, subtus secus nervos pilis parvis substrigillosis paucis puberis, pedunculis ternis terminalibus, invol. squamis biserialis ovali-oblongis acutis, achæniis 1-aristatis nauticisve. — in insulâ Timor leger. Bot. Bauidiane exped. Affinis *W. aristata* sed car. ind. faciliè differt. (v. s. comm. à Mus. reg. Par.)
11. *W. INSULARIS*, caule basi suffruticoso adscendente, ramis vix apice pubescentibus, foliis petiolatis ovato-lanceolatis basi acutis trinerviis apice acuminatis subserratis utrinquè petiolisque pubescentibus, pedunculis è dichotomiis ortis apice folia 2 ferè linearia et capitula 3 pedicellata gerentibus, invol. brevius polyphylli squamis ovali-oblongis acutis, fructibus exaristatis. 3 in sterilibus saxosis insularum ad oram borealem Australasie apr. flor. legit cl. A. Cunningham. Affinis ex habitu *W. scabrusculæ*, ex calyce *W. Forsteriana*. (v. s. comm. à cl. inv.)
12. *W. PROCUMBENS*, caule procumbente basi suffruticuloso, ramis adscendentibus scabris, foliis sessilibus ovato-lanceolatis seu lanceolatis acuminatis rariter et grossè dentatis utrinquè strigillosis asperrimis trinerviis, capitulis pedicellatis ex dichotomiis solitariis, invol. biserialis squamis substrigillosis exter. foliaceis oblongis disco longioribus, achænio subtrigono glabro coronulâ trisetâ brevissimâ superato. 3 2 in littoribus insularum ad oram borealem Australasie apr. flor. legit cl. A. Cunningham. Folia pilis albis et post eorum delapsum maculis albis ut in Borragineis notata. Capitula pauciflora. Ligulæ vix invol. longiores. (v. s. comm. à cl. inv.)
13. *W. FORSTERIANA*, caule suffruticoso, foliis petiolatis oblongo-lanceolatis acutis grossè serratis suprâ strigosis subtus alido-villosis, pedunculis axillaribus terminalibusque 1-cephalis. 3 in insulâ Norfolk (Forst.), locis apicis graminosis ad oram meridionalem insulæ (Cunn. !), Bupthalthum uniflorum Forst. prod. 541. Spreng. syst. 3. p. 605. Wedelia Forsteriana Radd. fl. norf. p. 51. n. 102. Ex Spreng. folia quintuplinervia sed in specim. meo evid. triplinervia. Invol. squamis 15-20 bi- seu triseriatis constans. Folia specim. mei ovali-lanceolata dicenda. Fructum ex specim. juniore non judicare queo: (v. s. comm. à cl. Cunningham.)

§. 2. *Wollastonia dubia*, *Americana*, *discoidea*.

14. *W. LANTANOIDES*, caule fruticuloso, ramis striato-sulcatis subasperulis, foliis petiolatis latè ovatis basi obtusis apice vix acutis crenatis, propè basin triplinerviis, utrinquè subpuberulis, subtus nervoso-reticulatis, pedunculis terminalibus ternis 1-cephalis, capitalis discoideis, invol. squamis ellipticis acutis

- disco brevioribus, achæniis obovatis calvis aut setas 1-3 fragiles caducas apice gerentibus. $\bar{3}$ in insulis Caribæis legit cl. West. Bupthalmum latifolio affine Puer. ! herb. et h. Vahl. An Wedeliæ discoidæ Less. syn. 222 affinis aut fortè eadem? Cor. flavæ. Petioli pollicares. Limbus fol. 2 poll. longus et latus. (v. s.)
15. F. ? PROSTRATA, caulibus herbaceis procumbentibus adpressè pubescentibus, foliis oppositis obovatis obtusis basi in petiolum attenuatis integris glabrisculis, pedicellis axillaribus ascendéntibus 1-cephala folio triplò longioribus, invol. sub-1-serialis squamis elliptico-oblongis, achæniis crassis subtrigona compressive calvis aut aristam brevem gerentibus, recept. paleis lineari-subulatis. \mathcal{U} in Brasiliæ prov. Rio-Grande. Herba habitu ferè Veroniam Allionii, fortè genus proprium constituens, sed floribus omninò in specim. deficientibus hic provisoriè adjuncta. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1073 miss.)

CCCX. TILESIA Meyer esseq. p. 252.

Capitulum multiflorum homogamum seu heterogamum, fl. radii ligulatis fœmineis bifidis, disci tubulosis 5-dentatis hermaphr. Invol. imbricatum 2-3-seriale, squamis coriaceis ovatis apice squarrosis. Recept. hemisphæricum, paleis squamas referentibus flores amplectentibus. Styli disci et fortè radii rami setacei plumosi. Achænia obpyramidata tetragona immarginata calva, integumento externo (calycis tubo) separabili. — Planta scandens Guianensis scabra. Folia opposita petiolata ovata serrato-crenata utriusquè scaberrima. Capitula 1-3 ad apices pedunculorum congesta. Cor. crocæ.

1. T. CAPITATA (Mey. l. c.). — in Guianâ batavâ ad Essequebo. Meyera capitata Spreng. syst. 3. p. 601. ex syn.

CCCXI. PASCALIA Ort. dec. 4. p. 39. t. 4. Willd. sp. 3. p. 2228. Cass. dict. 46. p. 406. Less. syn. 223.

Capitulum multiflorum radiatum, fl. radii ligulatis fœmineis 1-serialibus, disci hermaphr. tubulosis 5-dentatis. Invol. disco æquale, squamis subbiserialis linearibus subfoliaceis. Recept. subconvexum paleaceum, paleis lanceolatis semiamplectentibus. Achænium radii triquetrum squamis inv. non inclusum, disci tetraquetrum obovatum paleis semi-inclusum (utrumque maturum ex Ort. drupaceum carnosum). Pappus coroniformis brevissimus irregulariter dentatus. — Herba Mexicana scabriuscula parcè ramosa strobilorum Pini odorem spirans (Ort). Folia opposita lineari-lanceolata subtriplinervia integerrima, inferiora irregulariter dentata. Capitula terminalia solitaria. Cor. lutæ.

1. P. GLAUCA (Ort. l. c.). \mathcal{U} in Chili non longè ab oppido Chillan. (v. v. c.)

+ Species vix nota.

- P. GLANDULOSA Fisch. ex Steud. nom. ph. 59a.

++ Species exclusa.

- P. BACCATA Spreng. = Wulfia membranifolia.

CCCXII. RUMFORDIA.

Capitulum radiatum, ligulis multinerviis fœmineis circ. 10, tubulis hermaphroditis 5-dentatis. Invol. duplex, exter. squamæ 5 patulæ foliaceæ ovali-oblongæ disco breviores, inter. circ. 10 complicatæ ovatæ mucronatæ, dorso glanduloso-puberæ, achænia radii involventes. Recept. plano-convexum, paleis squamas inv. inter. referentibus. Cor.

omnes basi puberæ super ovarium articulatæ. Stylus radii longus apice bifidus glabrinsculus, disci exsertus ramis semiteretibus apice puberulis cono brevissimo subsuperatis. Achænia obovato-compressa glabra calva conformia. — Frutex Mexicanus glaber. Rami teretes. Folia opposita ampla sessilia latè ovalia, basi attenuata, apice breviter acuminata, ad quartam partem longitudinis triplinervia, margine basi integerrima apice serrulata. Panícula oppositè ramosissima polycephala bracteis oblongis intermixta. Capitula pedicellata. Ligulæ pallidè flavæ, discus obscuris luteus. — Caract. ferè Heliopsidis, sed habitus diversissimus. Dicavi in memoriam cl. physici Benj. Thomson comitis de Rumford qui experimentis circa calorem et lucem et etiàm circa lignorum naturam momentosis inter hodiernos spectabilis remanet.

1. R. FLORIBUNDA. 3 in Mexico detexit cl. de Karwinski. Folia super. 8 poll. longa, 5-6 poll. lata. (v. s. in h. acad. reg. Monacensis.)

CCCXIII. HELIOPSIS Pers. ench. 2. p. 473. Dunal mem. Mus. h. n. Par. 5. p. 54. Less. syn. 223. — *Heliopsis* et *Kallias* Cass.

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis 1-serialibus, disci hermaphr. tubulosi 5-dentatis. Invol. bi-seu pluriseriale, squamis ext. sæpius foliaceis, cæt. subimbricatis. Recept. conicum, paleis oblongo-lanceolatis achænia amplectentibus. Achænia angulato-compressa glabra omninò calva, disco epigyno parvo. — Herbae Americanæ erectæ perennes oppositifoliæ. Folia dentata. Capitula solitaria pedunculata. Cor. flavæ. — An ab hac satis differt *Helepta* Raf. neogen. 1825. p. 3?

Sect. I. EUHELIPSIS. — *Heliopsis* Cass. dict. 24. p. 333.

Corollæ omnes glabræ, ligulæ super ovarium articulatæ. Achænia lævia. Invol. 2-3-seriale. — Herbae Boreali-Americanæ. Folia serrata.

1. H. LÆVIS (Pers. ench. 2. p. 473), caulibus glaberrimis, foliis glabris ovato-acuminatis serratis, invol. squamis externis lanceolatis patulis apice subseratis. 2 in Amer. bor. à Pensylvaniâ ad Carolinam (Pursh), in fluminum ripis umbrosis rupestribus (Dun.). Dun! mem. mus. 5. p. 55. Hook. bot. mag. t. 3372. *Helianthus lævis* Linn. f. suppl. 1278. *Buphthalmum helianthoides* Linn. hort. ups. 264. L'Hér. stirp. 93. t. 45. *Silphium solidaginoides* Linn. sp. 1302? *Rudbeckia oppositifolia* Linn. sp. 1280. — Moris. ox. s. 6. t. 3. f. 69. (v. v. c. et s. sp?)

β. foliis lævissimis, ligulis pallidioribus. *Silphium helianthoides* Gron. virg. 134. ex Dun. l. c.

2. H. SCABRA (Dunal! mem. mus. h. n. Par. 5. p. 56. t. 4), caulibus foliisque scabris, foliis ovato-oblongis acuminatis serratis, squamis invol. ovato-acuminatis integris. 2 in Amer. ad flumen Missouri et (ex Hook. fl. bor. am. r. p. 310) ad ripas fl. Rubri (Red-River.). H. canescens Don in bot. reg. t. 592. excl. syn. (v. v. c. et s. sp.)

Sect. II. KALLIAS. — *Kallias* Cass. dict. 24. p. 326. 46. p. 406.

Corollæ omnes glabræ, ligulæ cum ovario continuæ nec articulatæ. Achænia subtruncata rugosa. Invol. 2-4-seriale imbricatum. — Herbae Australi-Americanæ. Folia serrata. — An fortè ex ligulis persistentibus *Phylactidi* affiniore?

3. H. BUPHTHALMOIDES (Dunal! mem. mus. Par. 5. p. 57), caulibus puberulis apice villosis, foliis pilosiscaulis ovato-acuminatis serratis, inv. squamis linearibus integris imbricatis. 2 in Peruvîâ. *Anthemis ovalifolia* Ort. dec. 6. p. 73. Desf.!

cat. ed. 2. p. 125. *Verbesina ovata* hort. Par. ex Poir. dict 8. p. 454. *Anthemis buphthalmoides* Jacq. h. Schœnbr. 2. p. 13. t. 151. Willd. sp. 3. p. 2185. excl. Cav. syn. *Acmella buphthalmoides* Pers. ench. 2 p. 473. *Kallias ovata* Cass. dict. 24. p. 327? ex syn. (v. c.)

4. *H. CANESCENS* (H. B. et Kunth nov. gen. am. 4. p. 212), caulibus sub foliorum jugis pedunculisque bifariam pilosis cæterum glabresculis, foliis latè ovatis infer. subcordatis acuminatis serratis subvillosis, invol. squamis ovalibus imbricatis. $\frac{1}{2}$ in Peruvia. *H. rugosa* hort. berol. *Helianthus buphthalmoides* hortol. *Helenomcium canescens* Willd. herb. ex Kunth. *Helianthus cordatus* Lamb. herb. ex Don. *Kallias ovata* Cass. dict. 24. p. 327? ex descr. *Kallias dubia* Cass. dict. 2. p. 333. Recept. intus cavum. Spec. à priori fortè non satis diversa. (v. c. et s. sp.)

Sect. III. *BALSAMORHIZA* Hook. fl. bor. am. 1. p. 310.

Achænia compresso-tetragona calva. Capitula ferè Helianthi. Radix crassa balsamifera. Folia radicalia petiolata pinnatilobata. — An genus proprium ex Hook. ? †

5. *H. ? BALSAMORHIZA* (Hook. l. c.), subpubescens, radice crassâ balsamiferâ, foliis radicalibus plurimis petiolatis pinnatisectis, segmentis subpinnatifidis incis, invol. squamis numerosis lanceolato-acuminatis imbricatis caulem paucifoliatum subæquantibus. $\frac{1}{2}$ in glareosis ripis fl. Columbia prope arcem Vancouver. Radix fusiformis succo terebinthinaceo. Folia 6-8 poll. longa.
6. *H. ? TEREBINTHACEA* (Hook. l. c.), pubescens, radice crassâ balsamiferâ, foliis radicalibus petiolatis ovato-lanceolatis sinuato-pinnatifidis crenato-serratis, invol. squamis numerosis lanceolato-acuminatis imbricatis caulem paucifoliatum subæquantibus. $\frac{1}{2}$ Hab. cum priore ad quam valdè accedit.

† *Species exclusa.*

H. PLATYGLOSSA Cass. = *Guizotia oleifera*.

CCCXIV. *GUIZOTIA* Cass. bull. philom. 1821. p. 187. dict. 59. p. 247. — Ramtilla DC. in Wight contrib. bot. ind. p. 18.

Omnia *Heliopsisidis*, sed cor. imâ basi et ad medium tubi pilis crassis subarticulatis annulatim instructæ, omnes super ovarium articulatæ. Achænium læve. Invol. biseriale. — Species *Gerontogea* annua. Genus inter *Heliopsidem* et *Tetragonothecam* medium. Priori affine achæniî disco minimo et invol. squamis liberis, sed distinctum corollis basi barbatis super ovarium omnibus articulatis; ad posterius accedens corollis basi barbatis et invol. biseriali, sed discrepans invol. squamis liberis et disco epigyno majore.

1. *G. OLEIFERA* (DC. mem. soc. hist. nat. Genev. v. 7. cum tab. 2), caule ad apicem pubescente, foliis seniamplexicanibus subcordato-aut ovato-lanceolatis remotè serratis subscabridis, invol. biserialis squamis ext. 5 latè ovatis foliaceis inter. superantibus. ① in Abyssiniâ ex Linn. f. sed fortè culta? in Indiâ orient. frequenter culta et verisim. etiâ spontanea! *Polymnia Abyssinica* Linn. f. suppl. (1781) p. 383. *Polymnia frondosa* Bruce voy. ed. 4^o. gall. v. 3. p. 575. *Verbesina sativa* Roxb. cat. calc. (1814) p. 62. Sims bot. mag. t. 1017. Ainsl. mat. med. ind. 2. p. 256. *Werinnua* Heyn. tract. on Ind. p. 49. ex Ainsl. *Parthenium luteum* Spreng. nov. prov. (1818) p. 31. *Heliopsis platyglossa* Cass. dict. 24. (1822) p. 332. *Tetragonotheca Abyssinica* Ledeb. ind. hort. dorp. suppl. 1824. p. 7. *Jagera Abyssinica* Spreng. syst. 3. (1826) p. 590. *Guizotia Abyssinica* Cass. dict. 59. (1829) p. 248. *Helianthus oleifer* Wall. cat. et herb. n. 3194. comp. 304. (1830) et *Bidens*? *Ramtilla* Wall. l. c. n. 3191. comp. 301. *Ramtilla oleifera* DC. diss. mss. in soc. Genev. 1833 et in Wight contrib. bot. p. 18. *Buphthalmum Ramtilla* Ham. herb. et *Anthemis Mysorensis* herb. mss. ex

Wall. Culta in Abyssiniâ et in Indiâ orient. præsertim in agro Mysorensi ad oleum ex seminibus elicendum in lampadibus et condimentis usitatissimum et olei sesamei succoedaneum. Vernac. in Indiâ dicitur *Ram-Till*, *Ram-Tilla*, *Kutrelloo*, *Kuts-Yelloo*, *Werinnua*, et in Abyssiniâ *Nook* ex Frac.

a. *sativa*, foliis elongato-lanceolatis grossè serratis, ramulis elongatis subcorymbosis. — Hinc pleraque synonyma supra citata pertinent. (v. v. c. et s. ex Ind.)

β. *angustior*, foliis lineari-lanceolatis subdentatis, ramulis axillaribus floriferis brevissimis. ① in Bengalo inferiore ad margines aquarum ductam. Plantæ sativæ stirps sylvestris Wall.† n. 394. (v. s. sp. in h. soc. Linn. Lond.)

CCCXV. TETRAGONOTHECA Dill. elth. 378. Linn. gen. ed. 1. n. 800. ed. 2. n. 976. L'Hér. stirp. 177. Cass. dict. 59. p. 319. — Polymnia sp. Linn. syst. veg. Willd.

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis 1-seriatis, disci hermaphroditis tubulosis 5-dentatis. Invol. duplex, ext. squamis 4 basi in cupulam tetragonam concretis latè ovatis acuminatis maximis, int. squamis 8 ovali-lanceolatis parvis achænia radii semiamplexantibus. Recept. convexo-conicum paleaceum, paleis membranaceis lanceolatis acuminatis. Cor. omnes secus tubum hispidæ, super ovar. articulatae, radii latæ multinerviæ, disci dentibus erectis. Styli disci rami hispidi exserti. Achænia omnia obovata vix subangulata crassa lævia calva, areolâ terminali latiusculâ, nectario elevato. — Herba Boreali-Americana oppositè ramosa. Caulis hirtellus teres. Folia opposita dentata parcè villosa. Pedunculi terminales 1-cephali. Capitula ampla. Flores lutei.

1. T. HELIANTHOIDES (Linn. sp. 1273). ① in Virginiâ (Linn.), Carolinâ (Frac.), locis arenosis siccis (Ell. sketch 2. p. 407) L'Hér. stirp. 177. Polymnia Tetragonotheca Linn. syst. veg. p. 658. Abbot insect. 2. p. 137. t. 69 ex Willd. sp. 3. p. 2116. Silphium Tetragonotheca Gærtn. — Dill. elth. 378 t. 283. f. 365. (v. s. ex herb. L'Hér.)

† Species exclusæ.

T. ABYSSINICA Ledeb. = Guizotia oleifera.

CCCXVI. FERDINANDA Lag. nov. gen. et sp. p. 31. Cass. dict. 20. p. 429. 46. p. 404. Less. syn. 224.

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis 1-serialibus, disci hermaphr. tubulosis 5-dentatis. Invol. hemisphæricum disco brevius, squamis 2-3-seriatis adpressis. Recept. conicum, paleis oblongis submembranaceis amplectentibus. Anth. subnigricantes cæcaudatæ. Styli rami pubero-hirtelli cono brevi superati. Achænia radii obconica angulata pappo 2-5-squamellato subcoroniformi minimo aut nullo superata, disci oblonga compressa sub-4-costata omninò calva. — Frutices Mexicani. Folia alterna ovata aut oblonga basi attenuata apice obtusa integerrima aut crenata triplinerviæ, suprâ pubescentia aut scabra, subtus velutino-argentea aut pubescentia. Corymbus ramosus polycephalus. Folia summa bracteiformia colorata. Flores lutei. — Genus ambigit inter Heliantheas et Anthemideas, prioribus stigmatè, posterioribus habitu toto affinis. — Toto cælo differt à Roseniâ quâcum Sprengel conjungere voluit.

1. F. ARGUSTA (Lag. l. c.), foliis basi abruptè crenatis suprâ pubes-

centibus subitè velutino-argenteis, floralibus integerrimis suprâ præcipuè aureo-flavicantibus; pappo radii squamellato irregulari, squamellis vix imâ basi concretis. \S in Mexico (Lag. Alam.!), in montibus circa urbem (Berl. n. 683 et 932), in prov. Leoninâ ad occid. urbis Guanaxuato (Mendez!). Ferdinanda velutina Deaf! h. par. Cass. dict. 20. p. 430. Variat foliis ovatis oblongisve integerrimis aut latè crenatis, semper versus basin abruptè angustatis. (v. v. c. et s. sp. et ex hort. Madr.)

2. *F. LUTESCENS*, foliis oblongis lanceolatisve obtusis integerrimis infernè sensim attenuatis suprâ vix pubescentibus subitè incanis, pappo radii nullo. \S in Mexici collibus aridis de Guadalupe prope urbem (Lallav. et Lex.). *Anthemis lutescens* Lallav. et Lex. nov. veg. 1. p. 30 ex Alam. in litt. Priori valdè affinis. Rami ex siceo subscandentes. Folia et flores fragrantissimum odorem Abrotani æmulum spirant (Lallav. et Lex.). (v. s.)
3. *F. ERINENS* (Lag. l. c.), foliis suprâ scabris subitè pubescentibus, petiolis foliaceo-alatis. \S in Mexico. Cat. lga. †.

CCCXVII. *CHRYSOPHANIA* Kunth in Less. syn. p. 224.

Capitulum radiatum, ligulis parvis rotundis fœmineis, cor. disci tubulosis hermaphr. basi obsolete puberulis. Invol. subuniseriale. Recept. conicum, paleis oblongo-ellipticis obtusis multinerviis. Stylus Zaluzaniæ. Achænia quadrangulata quadricostata conformia omninò calva, disco epigyno parvo, sectione transversali quadrata aut compresso-quadrangula. — Fruticulus Mexicanus dichotomus. Folia alterna elliptica, grossè et inæqualiter crenata, petiolata, secus petiolum obtusè decurrenti-acuminata, suprâ canescentia, subitè incana. Paniculæ terminales fastigiatae parùm ramosæ. Capitula lutea magnitudine *Spilanthi acmellæ*.

1. *C. FASTIGIATA* (Kunth l. c.). \S in Mexico prope Actopam. Folia 15 lin. longa, 9 lin. lata.

CCCXVIII. *ZALUZANIA* Pers. ench. 2 (1807). p. 473. Cass. dict. 56. p. 232 non Comm. nec Neck. (1). — Zaluzaniæ sp. Less. — *Anthemidia* sp. Ort.

Capitulum multiflorum heterogamum, fl. radii 5–7 ligulatis 1-seriatis ovalibus fœmineis, disci hermaphr. tubulosi 5-dentati. Invol. biseriale patulum, squamis sericeis, int. brevioribus cuneatis trilobis. Recept. conicum, paleis membranaceis amplectentibus cuneatis apice trilobis squamas int. referentibus. Cor. disci hispidæ, tubo faucis æquali basi achæniū obtegente. Styli rami puberuli cono brevi terminati. Achænia calva, radii subtrigona hispidæ, disci compressa glabra lævia, exangulata. — Herba perennis basi suffruticosa Mexicana ramosa villosa-tomentosa. Folia infima opposita, cætera alterna, petiolata pin-natipartita, lobis paucis obtusè sinuato-repandis sæpius trilobis. Capitula pedicellata corymbosa laxè subpaniculata. Flores lutei.

1. *Z. TRILORA* (Pers. ench. 2. p. 473). \mathcal{L} \S in Mexico (Ort.) circa Real del Monte (Berl. pl. exs. n. 396), in collibus de Los Remedios San-Angel (Lall. et Lex.). *Anthemis triloba* Ort. dec. 6. p. 72. *Anthemis sinuata* Lallav. et Lex. nov. veg. mex. 1. p. 30. *Acmella trilobata* Spreng. syst. 3. p. 591. (v. v. c. et s. sp.)

(1) *Zaluzanskia* Neck. = *Marsilea*. *Zaluzania* Comm. ined. = *Berterix* sect.

CCCXIX. CHILIOPHYLLUM. — *Anthemidis* sp. *Ort.* — *Hybridella* *Cass. dict.* 22. p. 86. — *Zaluzaniæ* sp. *Less.*

Capitulum multiflorum heterogamum, fl. radii 2-4-serialibus angustissimè ligulatis foemineis sæpè sterilibus, disci tubulosis hermaphr. 5-dentatis. Recept. convexum paleis acuminatis achænio longioribus onustum. Invol. biseriale, squamellis plurimis linearibus (an paleæ?) inter squamas majores et radii flores sitis. Ligulæ sæpè bifidæ irregulares. Tubuli imâ basi barbellati. Stylus ligularum profundè bifidus, ramis hirtellis; disci inclusus brevissimè bifidus, ramulis truncatis apice subbarbatis. Achænia calva compressa, glabra, incurva et cuneiformia ex *Ort.* — Herba Mexicana sparsè pilosa perennis, ramis apice nudis 1-cephalis. Folia bi- aut tripinnatipartita, lobis inæqualibus linearibus acutis. Capitula globosa lutea. — Ob folia Millefolium referentia antiquum Millefolii nomen huic plantæ adaptavi, ne nimis hybridum Cassinii nomen consecrare.

1. *C. GLOBOSUM*. 2. in Mexico. *Anthemis globosa* *Ort.* dec. 4. p. 46. *Jacq. hort. Schönbr.* 3. p. 64. t. 371. *Hybridella globosa* *Cass. dict.* 22. p. 86. *Acmella globosa* *Spreng. syst.* 3. p. 59a. (v. s.)

Div. II. RUDBECKIÆ *Less. syn.* p. 225. — *Rudbeckiæ* veræ *Cass. opusc.* 3. p. 59.

Capitula heterogama, fl. radii ligulatis neutris seu styliferis sterilibus. Achænia erostralia calva aut pappo coroniformi coronata.

CCCXX. ECHINACEA *Mœnch meth.* (1794). p. 591. *Cass. dict.* 35. p. 274. 46. p. 397 et 400. *Less. syn.* 225. — *Brauneria* *Neck. elem.* n. 301. — *Bobartia* *Petiv. mus. amm. herb.* 473 non *Linn.* — *Helichroa* *Raf. neogen.* (1825). n. 35. *herb.* p. 30 et in *Ser. bull.* 1830. p. 216. — *Rudbeckiæ* sp. *Linn.*

Capitulum multiflorum heterogamum, fl. radii neutris longè ligulatis 1-serialibus, disci hermaphr. regulariter 5-fidis, tubo subnullo, fauce nudâ, limbi dentibus erectis. Invol. 3-seriale, squamis lanceolatis ciliatis. Recept. ovatum, paleis rigidis supernè cartilagineis flores disci superantibus onustum. Stam. filamenta ex imâ corollâ orta. Styli rami appendiculis semilanceolatis superati. Achænia tetragona obpyramidata crassa pappo irregulariter lacero subcoroniformi deciduo coronata. — Herbæ Boreali-Americanae perennes. Folia radicalia petiolata, caulina alterna sessilia serrata aut integerrima. Rami supernè nudi 1-cephali. Capitula ampla, ligulis purpureis 2-3-dentatis 1-2 poll. longis, fl. disci obscurè virescentibus.

1. *E. PURPUREA* (*Mœnch l. c.*), caule lævissimo, foliis glabris, infer. ovatis secus petiolum longè decurrentibus, super. lanceolatis. 2. in montibus Americae bor. à Virginia ad Floridam. *Rudbeckia purpurea* *Linn. ap.* 1280. *Curt. bot. mag.* t. 2. *Schkuhr handb.* t. 259. — *Catesb. car.* 2. t. 59. *Mill. ic.* t. 224. f. 1. Ligulæ longissimæ dependentes apice acutè bidentatæ. Discus valdè convexus. (v. v. c.)
2. *E. SEROTINA*, caule hispido, foliis scabris, infer. ovatis secus petiolum vix decurrentibus, superioribus ovato-lanceolatis. 2. in planitiibus Louisianae inf. *Rudb. purpurea f. serotina* *Nutt. gen. am.* 2. p. 178. *Rudbeckia serotina* *Sweet brit. fl. gard.* t. 4. *Lodd. bot. cab.* t. 1539. *Rudb. hispida* *Hoffmann verz.* 2. 1826. p. 201. *Rudb. speciosa* *Link. h. ber. enum.* 2. p. 353 non *Schrad.* *Rhizoma* horizontale. Ligulæ 2-3-dentatæ, patulae nec dependentes, breviores quàm prioris. (v. v. c.)
3. *E. ANGUSTIFOLIA*, caule hispido, foliis omnibus lineari-lanceolatis hispidis in-

tegerimis, radicalibus subpetiolatis, caulinis sessilibus. \mathcal{L} in Mexicanâ provinciâ Texas prope Austin legit cl. Berlandier (pl. exs. n. 1692), fl. maio. Discus vix elevatus. Ligulæ breviores quàm priorum, patulæ. (v. s. sp.)

4. *E. HETEROPHYLLA* (D. Don in Sweet brit. fl. gard. ser. 2. t. 32), caule hispido, foliis radicalibus panduræformibus secus petiolum decurrentibus, caulinis lanceolatis subserratis. \mathcal{L} in Mexico. *Coreopsis heterophylla* Cav. ic. 3. p. 34. t. 268. *Simsia*? *heterophylla* Pers. ench. 2. p. 478. *Ximenesia Cavanillesii* Spreng. syst. 3. p. 605. Radix crassa. Ligulæ 15-20 purpureo-lilacinæ pollicares et ultra. Paleæ membranaceæ longitudine florum disci. Achænia 4-gona corollâ laciniatâ superata.

N. B. Species octo admisit cl. Rafinesque in herb. p. 30. sed characteribus brevissimis distinctas; conferant idèò autoptæ an *Helichroæ* Linnæana, elatior, amoena, uniflora, angustifolia, furcata et crocea sint species aut varietates.

- CCCXXI. *RUDBECKIA* Cass. dict. 46. p. 398. Less. syn. 226. Gærtn. fr. 2. p. 435. f. 172. f. 1. — *Rudbeckiæ* sp. Linn. — *Centrocarcha* Don in Sweet brit. fl. gard. ser. 2. n. 87. — *Obeliscotheca* Vaill. act. Acad. Par. 1720. — *Helio-phthallum* Raf. fl. lud. 72?

Capitulum multiflorum heterogamum, fl. radii neutris ligulatis, disci hermaphr. 5-dentatis. Invol. biseriale, squamis foliaceis patentibus. Recept. conicum, paleis acutis planis seu navicularibus onustum. Cor. disci tubo brevi, radii absque ullo genitalium vestigio. Styli rami cono brevissimo superati. Achænium tetragonum obpyramidatum, areolâ laterali, disco epigyno magno, nunc subcalvum, nunc pappo brevissimo inæqualiter dentato coronatum. — Herbæ Boreali-Americanae sæpius perennes hispido-scabræ ramosæ. Folia alterna, alia variè pinnatifidata, alia dentata aut integra. Rami nudi elongati 1-cephali. Ligulæ luteæ. Discus fusco-violaceus.

1. *R. LACINIATA* (Linn. sp. 1279), caule glabro, foliis infer. pinnatisectis, segmentis ovalibus acutis trilobis aut grossè incisis, summis ovatis indivisis, pappo inæqualiter dentato. \mathcal{L} in Amer. bor. à Canadâ ad Virginiam, ad fossas et margines paludum (Pursh), in Noveboraco (Torr.!). Schrad. neu. journ. 2. p. 61. — *Moris. oxon.* 3. s. 6. t. 6. f. 53. *Corn. canad.* 179. ic. Ligulæ patulæ sesquipollicem longæ flavæ. Discus fuscus. (v. v. c. et s. sp.)
2. *R. DIGITATA* (Mill. dict. n. 6 ex Schrad. neu. journ. 2. p. 61 non Willd.), caule lævi, foliis inferioribus pinnatisectis, segmentis pinnatifidis, superioribus simpliciter pinnatisectis, summis trifidis, pappo irregulariter dentato brevissimo. \mathcal{L} in montanis Virginie et Carolinæ. *R. laciniata*, \mathcal{L} . angustifolia. Pers. ench. 2. p. 476. — *Moris. ox.* 3. s. 6. t. 6. f. 54. — *R. lævis Hoffmans. verz.* 3. 1826. p. 58 est hujus fortè var. glabra hortensis. (v. v. c.)
3. *R. TRILOBA* (Linn. sp. 1280), caule hispidulo, foliis ciliatis pilosiusculis, infer. tripartitis, super. indivisis lato-lanceolatis hinc indè serratis, invol. squamis linearibus inæqualibus radio dimidiò brevioribus, paleis recept. cuspidatis, pappo subnullo. \mathcal{L} ? (1)? in montanis Virginie et Carolinæ (Pursh). Ker bot. reg. t. 525. *R. triloba* et *aristata* Pursh fl. bor. am. 2. p. 574 et 575. Nutt. gen. 2. p. 178. *Centrocarcha aristata* Don in Sweet br. fl. gard. ser. 2. n. 87? — Pluk. alm. t. 22. f. 2. Ligulæ luteæ 9 lin. longæ. Discus fusco-purpureus. (v. v. c.)
4. *R. TOMENTOSA* (Ell. sketch 2. p. 453), pube brevi subtomentosa, caule ramoso, ramis erectis virgatis, foliis lanceolatis acutis inciso-dentatis integrigve scabris, inferioribus trisectis, invol. squamis lineari-lanceolatis deflexis radio multò brevioribus, paleis rec. truncatis, pappo obsoleto subdentato. \mathcal{L} in Georgiæ distr. occid. *R. subtomentosa* Pursh fl. bor. am. 2. p. 575? *R. triloba* var. *tomentosa* Michx. fl. 2. p. 144? *Centrocarcha triloba* Don in Sweet l. c.? Ligulæ circ. 8-floræ. Discus fuscus.

5. *R. VULSIDA* (Aiton kew. ed. 1. v. 3. p. 251), caule hispido ramoso polycephalo, foliis oblongo-lanceolatis hispidis, infer. denticulatis, super. integerrimis, basi angustatis suboordatis apice acuminatis, invol. squamis ovali-oblongis radium sequantibus, paleis recept. lanceolatis, pappo vix ullo. $\frac{1}{2}$ in pratis et sylvis montanis à Pensylvaniâ ad Carolinam. Sims bot. mag. t. 1996. *R. chrysomela* Michx. fl. bor. am. 2. p. 143. *R. aspera* Desf. h. par. ed. 3. p. 183. *R. fulgida* et *aspera* Pers. ench. 2. p. 477. *R. fulgida* et *chrysomela* Spreng. syst. 3. p. 612. Ligulæ circ. 12. intense luteæ, 3-dentatæ. Discus atro-purpureus. (v. v. c.)
6. *R. HIRTA* (Linn. sp. 1280), caule hirsutissimo parçè ramoso oligocephalo, foliis latè ovatis aut obovatis tripli-aut quintuplinerviis serratis hirtis, invol. squamis foliaceis hirsutis radium subæquantibus, paleis recept. lanceolatis, pappo vix nullo. $\frac{1}{2}$ Desf. (1) Pursh in montanis à Virginiâ ad Floridam. Sweet brit. fl. gard. 1. t. 82. — DiH. elth. t. 218. f. 285. Pluk. alm. t. 242. f. 2. Mill. ic. t. 224. f. 1. Ligulæ flavæ subtus pubescentes. Discus atro-purpureus. (v. v. c. et s. sp.)
7. *R. GRANDIFLORA* (Gmel. hort. Carls. ex Steud.), caule ramoso angulato hispido, foliis petiolatis acutis basi attenuatis reticulato-venosis, infer. ovatis 7-nerviis crenato-dentatis, super. lanceolatis scaberrimis obsolete crenatis, invol. squamis radio quintuplò brevioribus, ligulis dorso præsertim ad basin hispidis, recept. paleis pungentibus, pappo brevissimo subintegro. $\frac{1}{2}$ in Americâ boreali. *Centrocarcha grandiflora* Don in Sweet brit. fl. gard. ser. 2. v. 1. t. 87. Rudb. nudicaulis Nutt. mas. Ligulæ flavæ 2 poll. longæ dependentes. Discus convexus atro-purpureus (v. v. in h. Gen.)
8. *R. SPECIOSA* (Wender. in florâ 1829. 1. p. 30. Schrad. ! in litt. 1832), caule hispidulo, foliis puberulis, radicalibus longè petiolatis ovatis acuminatis 5-nerviis, caulinis sessilibus lineari-lanceolatis grossè inciso-dentatis, summis integris, invol. squamis lineari-lanceolatis ciliatis radio dimidiò brevioribus, paleis lanceolatis, pappo vix ullo. — patr. ign. Folia radicalia ferè Plantaginis majoris, caulina 5 poll. longa, 6 lin. lata. Ligulæ circiter 24 luteæ 8-10 lin. longæ apice tridentatæ. Discus atro-purpureus. (v. s. et v. c.)
9. *R. MOLLIS* (Ell. sketch 2. p. 453), caule hispido-villoso ramoso, foliis sessilibus semiamplexicaulis ovali-sublanceolatis dentatis et sup. integris mollissimè tomentosis, invol. patulo villosissimo ligulis dimidiò breviori, paleis recept. lanceolatis, achæniis vix 4-dentatis. $\frac{1}{2}$ in Georgiæ distr. occidentibus. Tota subcanescens. Folia caul. 15 lin. longa, 9 lin. lata. Ligulæ 12-20 basi intense, apice pallidiùs flavæ, 6-7 lin. longæ. Discus atro-purpureus. (v. s. c. ex hort. Cels. in h. L'Hér.)
10. *R. RADULA* (Pursh fl. bor. am. 2. p. 456), caule infernè hispido supernè glabro nudiusculo, pedunculis longissimis 1-cephalis, foliis ovatis attenuatis tuberculatis hispidis, invol. squamis imbricatis ovatis acuminatis ciliatis. — in Georgiâ.
11. *R. DISCOLOR* (Pursh fl. bor. amer. 2. p. 574), caule folisque strigis rigidis basi tuberculatis hispido-scabris, foliis sessilibus spathulato-lanceolatis subintegerrimis, invol. squamis ovatis obtusiusculis radio paulò brevioribus, recept. paleis oblongis acuminatis apice fimbriatis, pappo vix ullo. — in Floridâ (Barr. ex Pursh), Georgiæ distr. occident. (Ell. sketch 2. p. 455), et in territ. Arkansano (ex sem. à cl. de Pourtales lectis). Ligulæ supernè intense luteæ, subtus pubescentes pallidæ. Discus valdè convexus atro-purpureus. Nostra meliùs convenit cum descr. Elliottii quàm Purshii. (v. s. comm. à cl. de Charpentier.)
12. *R. LÆVIGATA* (Pursh fl. bor. am. 2. p. 574), glaberrima, caule lævigato paniculato, foliis ovato-lanceolatis utrinquè acuminatis integerrimis aut subdentatis lævigatis triplinerviis, invol. squamis lanceolatis longitudine radii. $\frac{1}{2}$ in Georgiæ pinetis. Ell. sketch 2. p. 454. Ligulæ brevès pallidæ.
13. *R. GLABRA*, glaberrima, foliis caulinis distantibus sessilibus elongato-linearibus integerrimis utrinquè vix attenuatis apice callosulis, invol. squamis lanceolatis radio dimidiò brevioribus, paleis recept. lanceolatis, pappo 4-dentato. —

- in Georgiâ circa Savannah? An fortè var. *R. lævigatæ*? (Caules ramive elongati monocephali. Folia 5 polli. longa, 6 lin. lata, nervulis anastomosantibus notata. Ligulæ 12-15 lin. longæ flavæ apice acutè dentatæ. Discus fusco-purpureus. (v. s. comm. à cl. Duby.)
14. *R. SPATHULATA* (Michx. fl. bor. am. 2. p. 144), gracilis pubescens, caulibus 1-cephalis, foliis amplexicaulibus ovato-spathulatis lævibus integerrimis, involucri patulo imbricato (?), ligulis 3-dentatis. — in montanis Carolinæ (Michx) et Floridæ (Bartr.). Nutt. gen. am. 2. p. 178. De involucri imbricato silet Nuttall.
15. *R. GRACILIS* (Nutt. gen. am. 2. p. 178 non Banks), tenuis pubescens 1-3-cephala, pedunculis longissimis, foliis spathulato-ovatis remotè denticulatis, involucri patulo. — in pinetis à Virginiâ ad Floridam. Discus conicus fuscus.
- † *Species dubiæ, extra Americæ borealis confæderationem nascentes, fortè omnes à genere excludendæ.*
16. *R. ? NATIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 244), caule simplici sca-piformi piloso 1-cephalo, foliis radicalibus hastato-lyratis crenato-serratis adpresso-hispidulis, invol. squamis biserialibus lanceolatis acuminatis, paleis recept. lanceolatis scariosis, pappo paleis subulatis pluribus minutissimis constante. ? in Mexico juxta Santa-Rosa de la Sierra. An *Chrysostematis* sp.? an genus proprium?
17. *R. ? LEUCANTHA* (Lag. nov. gen. et sp. p. 3a. n. 399), foliis deltoideo-ovatis acutis serratis triplinerviis. ? in Mexico. Cæter. ign.
18. *R. ? NUDICAULIS* (Pers. euch. 2. p. 477. n. 11), caule subscapiformi hirsuto, foliis oppositis sublanceolatis obsolete dentatis glabris, pedunculis longissimis strictis nudis. — ad Monte-Video ex herb. Juss.
19. *R. ? DENSIFOLIA* (Smith in Rees cycl. 30. n. 10), caule scabro, ramis 1-cephalis, foliis oppositis petiolatis elliptico-lanceolatis, invol. squamis adpressis radio dimidiò brevioribus. — ad rupes circa Monte-Video legit Commerson. Ligulæ numerosæ latæ flavæ. Discus fusco-purpureus. Invol. squamæ scabræ fimbriatæ.
20. *R. ? DECUMBENS* (Smith in Rees cycl. 30. n. 11), caule decumbente repente, foliis oppositis petiolatis lineari-spathulatis ferè lævibus, pedunculis erectis solitariis angulatis longissimis gracilibus leviter pilosis, invol. squamis ovatis acutis adpressis radio dimidiò brevioribus. — ad rupes circa Monte-Video legit Commerson. Genus ex auct. ipso valdè dubium.
21. *R. ? BELLIDIOIDES* (Smith in Rees cycl. 30. n. 12), caule decumbente, foliis oppositis petiolatis ovatis scabris, pedunculis erectis solitariis longissimis apice inflatis et velutinis, invol. squamis ovatis adpressis radio dimidiò brevioribus. — ad Monte-Video legit Commerson.

†† *Species exclusæ.*

- R. ALATA* Jacq. = *Helenium quadridentatum*.
R. AMPLEXICAULIS Vahl. = *Dracopis amplexicaulis*.
R. AMPLEXIFOLIA Jacq. = *Dracopis amplexicaulis*.
R. ANGUSTIFOLIA Linn. = *Helianthus angustifolius*.
R. CINEREA Hort. = *Obeliscaria pinnata*.
R. CITRIODORA Hort. = *Obeliscaria pinnata*.
R. COLUMNARIS Sims. = *Obeliscaria columnaris*.
R. COLUMNIFERA Fras. = *Obeliscaria columnaris*.
R. DIGITATA Willd. = *Obeliscaria pinnata*.
R. NIAPIDA Hoffmann. = *Echinacea serotina*.
R. ODORATA Hortnl. = *Obeliscaria pinnata*.

- R. *OPPOSITIFOLIA* Linn. = *Heliopsis lævis*.
 R. *PAUCIFLORA* ex Steud. = *Zinnia pauciflora*.
 R. *PERFOLIATA* Cav. = *Dracopis amplexicaulis*.
 R. *PINNATA* Vent. = *Obeliscaria pinnata*.
 R. *PURPUREA* Linn. = *Echinacea purpurea*.
 R. *PURPUREA* β Nutt. = *Echinacea serotina*.
 R. *SEROTINA* Sweet. = *Echinacea serotina*.
 R. *TAGETES* James. = *Obeliscaria Tagetes*.

CCCXXII. *DRACOPIS* Cass. dict. 35. p. 273 et 46. p. 400. Less. syn. 226 (excl. syn. Raf. et Neck.?). — *Rudbeckia* sp. auct.

Capitulum multiflorum heterogamum, fl. radii neutris ligulatis, disci hermaphr. 5-dentatis. Invol. biseriale, squamis ext. 6-8 linearibus patulis, int. parvis adpressis. Recept. conicum, paleis squamas inv. interiores referentibus. Cor. disci subcylindraceæ, dentibus reflexis. Anth. inclusæ. Styli rami appendiculo lineari terminati. Achænium teres basi attenuatum, areolâ laterali, apice primâ fronte calvum, reverâ pappo coroniformi brevissimo integerrimo coronatum. — Herba annua Bor.-Americana glabra ramosa. Caulis ramique striato-sulcati. Folia caul. alterna cordato-amplexicaulia lævia integra acuta. Capitula terminalia solitaria, ligulis flavis, disco fusco.

1. *D. AMPLEXICAULIS* (Cass. l. c.). ① in Louisiana infer. ad ripas fl. Mississippi (Pursh) et in Mexico (Cav.). *Rudbeckia amplexicaulis* Vahl. act. soc. nat. scrut. Hafn. 2. p. 29. t. 4. Schkuhr handb. 3. t. 259. Rudb. *amplexifolia* Jacq. ic. rar. 3. t. 592. Rudb. *perfoliata* Cav. ic. 3. p. 257. t. 252. (v. v. c.)

CCCXXIII. *OBELISCARIA* Cass. dict. 46. p. 401. — *Lepachys* et *Ratibida* Raf. *journ. phys.* 1829. p. 100. — *Lepachys* Less. syn. 225. — *Rudbeckia* sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis 1-serialibus neutris, disci hermaphr. tubulosus 5-dentatis. Invol. 1-aut 2-seriale, squamis ext. paucis linearibus patulis, inter. parvis obtusis à paleis fortè non distinctis. Recept. elongatum spiciforme, paleis complicatis apice hirsutis achænia disci semiamplectentibus. Styli rami breves in appendicem semilanceolatam parçè barbellatam producti. Achænia radii trigona hispida abortiva, disci bilateraliter compressa ovalia calva anticè margine subalata aut apice unidentata. — Herba Bor.-Americana erecta speciosa glabriusculæ. Caulis striato-sulcatus. Folia alterna aut irregulariter subverticillata, pinnatisecta, petiolo striato, lobis 8-9 nunc indivisis, nunc pinnatipartitis. Rami apice longè aphylli 1-cephali. Ligulæ flavæ aut crocæ. Discus cylindricus, floribus parvis fuscis innumeris.

Sect. I. *LEPACHYS* Raf. l. c.

Invol. biseriale, serie int. brevissimâ. Ligulæ angustæ. Achænia anticè subalata apice calva.

1. *O. PINNATA* (Cass. dict. 46. p. 402), tenuissimè pubescens, foliis pinnatisectis, segmentis oblongo-lanceolatis utrinquè acuminatè parçè serratis, terminali majore trinervi, superioribus indivisis. 2. in partibus occidentalibus Carolinae et Georgiae (Pursh), in regione Illinoensi. *Rudbeckia pinnata* Vent. l. h. Cels. n. et t. 71. Michx. fl. bor. am. 2. p. 144. Smith exot. bot. 1. t. 38. Schrad. neu. *journ.* 2. p. 61. Sims bot. mag. t. 2310. Rudb. *digitata* Willd. sp. 3. p. 2247 excl. syn. *Lepachys pinnatifida* Raf. et Less. ll. cc. *Rudbeckia odorata* et R.

cinerea hortul. Flores subodorati. Recept. dissectum odorem aromaticum spirat. Ligulæ flavæ pollicem et ultrâ longæ dependentes. Achænia anticè alata. (v. v. c.)

Sect. II. RATIBIDA Raf. l. c.

Invol. uniseriale. Ligulæ elongatæ angustæ. Achænia anticè subalata apice calva.

2. *O. COLUMNARIS*, puberulo-hirtella, foliis pinnatisectis, segmentis lineari-lanceolatis acutis integerrimis, terminali cæteris subæquali, infimis gradatim subminoribus. \mathcal{L} in campis Louisianæ superioris (Pursh), ad flum. Missouri (Sims), Saskatchewan et Red-River (Hook.). *Rudbeckia columnaris* Pursh fl. bor. am. 2. p. 575. Sims bot. mag. t. 1601. Rndb. columnifera Fras. cat. 1813 ex Pursh. Ratibida sulcata Raf. l. c. Ligulæ elongatæ flavæ dependentes apice bidentatæ, interdum ex Nutt. croceo-rubræ ut in Tagete patulâ. (v. v. c.)

Sect. III. MONODONTA.

Invol. biseriale, serie int. brevissimâ. Ligulæ obovatæ. Achænia exalata ad marginem int. apice unidentata.

3. *O. PULCHERRIMA*, glabriuscula, foliis pinnatisectis, segmentis pinnatipartitis, lobis linearibus distantibus integerrimis. \mathcal{L} ? in Mexici provinc. Texas ad San-Fernando de Bejar, et in sinu Spiritus-Sancti ad lacum Sancti-Nicolai legit cl. Berlandier (pl. exs. n. 1545 et 1960). Ligulæ fusco-croceæ (ferè Tagetis seu Calliopsidis flor. æmulæ) margine et apice luteæ, patulæ demum deflexæ, 8-9 lin. longæ. (v. s. sp.)

† *Non satis nota.*

4. *O. TAGETIS*, hirsuta ramosissima, foliis radicalibus subintegris spatulatis (?) linearibus seu pinnatifidis, caulinis interruptè pinnatifidis lobis sæpius linearibus, pedunculis brevibus oligocephalis, ligulis recurvis. — in Amer. bor. ad montes Scopulosos (*Rocky-Mountain*). *Rudbeckia Tagetes* James exp. Rocky-Mount. 2. p. 353. Rndb. hirsuta Isis. 1824. p. 236. Herba pedalis multicaulis habitu *Marutæ cotulæ*. Ligulæ rubro-fuscæ. Discus fuscus minus elongatus quàm in affini *O. columnari* dicitur.

CCCXXIV. ANDRIEUXIA.

Capitulum multiflorum heterogamum, fl. radii circ. 20 uniseriatis ligulatis fœmineis sterilibus, disci hermaphr. tubulosis 5-dentatis. Invol. biseriale, squamis oblongis foliaceis vix acutis apice patentibus. Recept. convexum, paleis membranaceis acutis subconvolutis et flosculos semiinvol. Ligulæ subcoriaceæ cum ovario continuæ. Cor. disci cylindricæ glabræ. Stylus radii profundè bifidus glaber, disci rami subrevoluti apice summo barbati aut cono brevissimo superati. Achænia trigona, radii abortiva pubescentia, disci angustiora glabra omnino calva. — Herba Mexicana erecta parcè ramosa. Caulis angulatus, hinc indè subserratis piloso-hirsutus, basi glaber. Folia opposita petiolata ovata acuta serrata pubescentia. Caulis apice pedunculiformis 1-cephalus subcapitulo obconicus velutinus. Flores lutei. Capitulum ferè *Rudbeckiæ*. Genus affine *Zinnia* et *Phylactidi* ex ligulis cum ovario continuis sed differt achæniis cortice tenui donatis et epapposis: *Rudbeckiæ* et *Gymnolomati* ex styli cano brevissimo, recept. convexo, etc., sed diversum ligulis distinctè styliferis persistentibus, achæniis trigonis omnino calvis. — Dicavi cl. Andrieux qui Mexicum peragravit et plantam ipsam cum pluribus aliis hinc indè citatis detexit.

1. *A. MEXICANA*. (1) ? in Mexico ad montem San-Felipe prope Oaxacam julio

floridam legit cl. Andrieux. *Gymnolomia*? Andr.! pl. exs. n. 303. Herba sesquipedalis. Ligulæ 9 lin. longæ. Antheræ subnigricantes. (v. s.)

CCCKXV. ANOMOSTEPHIUM.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis neutris, disci hermaphrod. 5-dentatis. Invol. campanulatum pluri- (aut fortè uni-) seriale, squamis ext. ovatis acutis foliaceis laxè erectis, exterioribus oblongis membranaceis obtusis prioribus angustioribus sed longitudine æqualibus. Recept. planum, paleis membranaceis apice rufis subrotundis cæterum convolutis et achænia involventibus. Ligulæ tubo gracili longo, limbo ovali. Cor. disci tubo brevi, fauce longâ amplâ. Antheræ exsertæ. Styli disci rami puberi, saltè in flor. marginalibus obtusi, in centralibus cono acutissimo superati. Achænia lineari-subtrigona gracilia, radii glabra apice breviter bicornia, disci adpressè villosa pappo irregulari coroniformi superata. — Herbæ Brasilienses aut Caribææ basi suffruticosæ erectæ. Caulis subsimplex hispidus teres. Folia opposita sessilia utrinquè sparsè et adpressè piloso-scabra. Capitula ad apices caulis et ramorum pauca subsolitaria pedicellata. Cor. luteæ antheris nigris. Genus habitu *Wedeliæ*, sed ligulæ neutræ.

Sect. I. ANOMOSTEPHIA VERA.

Involucrum pluriserialè disco brevius.

1. *A. OVATIFOLIUM*, caule confertè hirtò, foliis sessilibus latè ovatis trinerviis basi obtusis apice vix acutis, capitulis breviter pedicellatis subconfertis. 2/ in Brasiliæ prov. Minarum-Generalium prope Sabara legit cl. Vauthier (pl. exs. n. 321). Paleæ recept. et invol. squamæ post anthesin patentim stellatæ. Antheræ exsertæ. (v. s.)
2. *A. BUPHTHALMOIDES*, caule folisque pube brevi rigidâ asperimis, foliis lanceolatis basi in petiolum attenuatis acuminatis vix subserratis penninerviis, pedicellis elongatis folium superantibus. 2/ in Guadalupâ (Bertero! Krauss!), loco dicto Pointe-à-Pitre (Perrottet!), *Buphthalmum asperimum* Spreng.! neu. entdeck. 2. p. 140 falsò demùm ab ipso auct. cum *Wedeliâ calycinâ* confusum. Antheræ inclusæ. Styli rami tennes. (v. s.)

Sect. II. ANOMOSTEPHIA DUBIA.

Involucrum subuniseriale disco longius aut æquale. — An genus diversum?

3. *A.?* *OBLONGIVOLIUM*, caule sparsè hirtò, foliis oblongo-lanceolatis utrinquè acuminatis subtriplinerviis serrulatis, infimis ellipticis, pedicello terminali foliis longiore. — in Brasiliâ prov. Minarum-Generalium ad Serra do Frio legit cl. Vauthier (pl. exs. n. 317!). Herba hispida simplex. Invol. squamæ circ. 7 oblongo-lanceolatæ disco longiores foliaceæ hirtæ. Paleæ recept. membranaceæ acutæ glabræ modicè convolutæ. Ligulæ elongatæ apice dentatæ. Antheræ inclusæ fortè subeffortæ. Stylus longè exsertus, ramis crassiusculis. Achænia juniora, radii linearia pappo bidentato, disci villosa pappo subetipitato calyciformi dentes a proferente, matura ignota et ideò de genere dubius maneo. (v. s.)
4. *A. ELLIPTICUM*, caule sparsè subhirtello, foliis ellipticis et superioribus oblongis vix subserrulatis trinerviis subscabridis ciliolatis, pedicello foliis longiore, invol. squamis oblongis disco circ. æqualibus. 2/ in Brasiliæ prov. Sancti-Pauli. Valdè affinis *A. oblongifolio* sed multò minùs hirtum, folia trinervia nec triplinervia, nervis lateralibus nempe à basi nec longè supra basin ortis. (v. s. in h. *Mss. reg. Par.* à *Mss. imp. Brasiliensi* sub n. 409 miss.)

CCCXXVI. *ASPILIA* Pet.-Th. gen. mad. p. 12. n. 40. Cass. dict. 3. suppl. p. 57.

Capitulum multiflorum radiatum, ligulis 5-10 neutris 1-serialibus apice dentatis, fl. disci tubulosis 5-dentatis hermaphr. Invol. biseriale, squamis oblongis disco vix æqualibus. Recept. planiusculum paleis oblongis complicatis achænia foveantibus apice acuminatis onustum. Styli rami cono brevi superati. Achænia linearia, adpressè villosa, superata pappo coroniformi dentato-ciliato. — Herbæ Madagascarienses procumbentes. Caules graciles puberuli. Folia opposita subsessilia lanceolata integra aut subdentata, supra sparsè, subtus ad nervos setoso-scabra. Capitula è dichotomiis longè pedicellata solitaria. Flores flavi. — Car. ex A. Bojeri, nec ab A. Thouarsii mihi ignotâ.

1. A. BOJERI, invol. squamis erectis, achæniis (immaturis) linearibus. $\frac{2}{4}$ in siccis inter dumeta per omnem insulæ Madagascar prov. Emirnensem legit cl. Bojer. Ximenesia procumbens Bojer! in litt. 1835. Ligulæ 8-10. (v. s. comm. à cl. Bojer.)
2. A. THOUARSII, invol. cylindrici squamis 5 exterioribus apice reflexis, achæniis oblongis apice latioribus. — in ins. Madagascar, verisimiliter prope Foulpointe. *Aspilium* Pet.-Th. l. c. Ligulæ 5-6. An certè prioris congener?

CCCXXVII. *GYMNOPSIS*. — *Gymnolomia* H. B. et Kunth nov. gen. am. 4. p. 217 non. Ker. — *Aldama* Lallav. et Lex. nov. gen. 1. p. 14? Less. syn. 227.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis neutris, disci tubulosis hermaphrod. Invol. biseriale, squamis ext. subfoliaceis. Recept. planum convexum aut fortè conicum paleaceum. Styli disci rami appendiculati. Achænia pappo coroniformi dentato brevissimo coronata. — Herbæ aut suffrutices Americani. Folia opposita petiolata tri- aut triplinervia dentata. Capitula pedicellata. — Genus malè definitum, differt à *Wulfiâ* achæniis non verè calvis nunquàm carnosus, à *Montagnâ* paleis apice non spinosis, ligulis homogeneis nec heterochromis, stylicque ramis longè appendiculatis. Species omnes fortè non sunt strictè congeneres nec satis notæ. — Nomen Kunthii falsum, *Lallavei* dubium, non admisi ne confusio oriatur, sed quàm minimè prius mutavi. Caract. pappi in 4 Kunthianis spec. observavit Ad. Brongniart et eorum icones humanissimè communicavit.

1. G. DENTATA, caule procumbente dichotomo tereti, foliis petiolatis sublanco-latis remotè serratis utrinquè asperis, capitulis è dichotomiis ortis longè pedunculatis, invol. squamis circ. 12 biseriatis ciliatis, ligulis 8-10 linearibus, recept. conico, paleis achænia foveantibus corrugatis apice cucullatis. — in inundatis rivuli Huehneyapa ad Sanctum-Josephum del Corval Mexic. *Aldama dentata* Lallav. et Lex. nov. veg. 1. p. 14. Flores aurei. De pappo silent cl. auct.
2. G. SCHERDEANA, caule erecto hirsuto subdichotomo, foliis breviter petiolatis scabro-hirsutis triplinerviis oblongo-ovatis acuminatis remotè serrulatis, invol. squamis biseriatis adpressis ciliatis, ext. ovatis, int. obovatis, recept. convexo, paleis achænia foveantibus carinatis, achænio subtrigono-compresso. — inter segetes ad margines dumetorum prope Jalappam. *Aldama dentata* Less. in linnæ 1830. p. 154. 1834. p. 271. Flores aurei. Achænia pappo brevissimo coroniformi superata, intima sæpè abortiva.
3. G. VERBESINOIDES, caule erecto subtetragono pubescente, foliis petiolatis ovato-lanceolatis basi cuneatis apice acuminatis vix subserratis triplinerviis utrinquè puberulis subscabridis, capitulis in dichotomiis apicibusque breviter pedicellatis, invol. biseriatis squamis paucis, ext. lanceolatis, inter. ovalibus, ligulis 5-7, paleis complicatis rectis, achæniis glabris pappo coroniformi dentato

superatis. — ad Trinitatem legit cl. Lockart (sed loci plures sunt hujus nominis?). Petioli 4 lin. longi. (v. s. comm. à cl. Hooker.)

4. *G. HELIANTHOIDES*, caule teretiusculo hirtio et aspero, foliis breviter petiolatis lanceolatis aut subpanduræformibus acuminatis triplinerviis hinc inde subdentatis utrinque strigilloso-scabris, petiolis ciliatis, capitulis ternis pedicellatis terminalibus, invol. biserialis squamis lanceolatis hirtellis, ligulis 10-12, recept. planiusculo, paleis membranaceis complicatis rectis, acheniis oblongis compressis, pappo coroniformi dentato. — ad Bonariam legit cl. Bacle. Capitula lutea. (v. s.)
5. *G. WEDDELIODES*, caule erectiusculo tereti apice hispidulo, foliis subpetiolatis oblongo-lanceolatis subacuminatis basi trinerviis et obtusis serratis supra strigillosis subtus sparse puberis, capitulis pedicellatis terminalibus, invol. squamis biserialis subæqualibus oblongis foliaceis strigillosis, paleis complicatis cuspidatis rectis glabris, ligulis 8-10, recept. vix convexo, acheniis pappo coroniformi ciliato superato. — in ins. Caribæis, fortè in Tabago (h. L'Hér.). Affine *G. tenellæ* sed formâ foliorum potius oblongâ quàm ovatâ jam primâ fronte differt. (v. s.)
6. *G. TENELLA*, caule procumbente subvolubili tereti hispidulo, foliis petiolatis ovatis acutis basi subcordatis utrinque hispidulo-scabris dentatis, capitulis pedunculatis ramos terminantibus, invol. squamis circ. 20 lineari-lanceolatis acutis laxè imbricatis, ligulis circ. 10, paleis linearibus acutis subcarinatis, recept. convexo, acheniis obovatis subtetragonis puberulis, junioribus pappo coroniformi brevissimè dentato superatis $\frac{1}{2}$ $\frac{3}{4}$ in Novâ-Granatâ prope Yaguay et Contreras. Flores flavi. *Gymnolomia tenella* H. B. et Kunth nov. gen. am. 4. p. 218. t. 373.
7. *G. RADBECKIOIDES*, volubilis, ramis sulcato-angulatis verrucosis, junioribus pilosis, foliis petiolatis ovatis acuminatis basi rotundatis crenato-serratis trinerviis utrinque hispidulo-pilosis, pedunculis axillaribus et terminalibus pilosis, invol. squamis 9-10 subbiserialis lanceolato-linearibus pilosis, paleis carinatis ovato-lanceolatis, recept. convexo, ligulis circ. 7, acheniis compresso-tetragonis, junioribus adultisque pappo vix dentato brevissimo coronatis. $\frac{1}{2}$ in temperatis prope Ayavaca Peruvianorum. *Gymnolomia radbeckioides* H. B. et Kunth nov. gen. am. 4. p. 218.
8. *G. HONKENSIIS*, caule tereti piloso, foliis petiolatis ovatis subacuminatis basi rotundatis trinerviis crenato-dentatis utrinque hispidulo-pilosis, capitulis longè pedunculatis axillaribus, invol. squamis 15-20 lanceolatis laxè imbricatis hispidis, recept. convexo, ligulis 2-3-fidis, acheniis junioribus apice pappo inæquali acutè et longiusculè dentato coronatis. $\frac{1}{2}$ $\frac{3}{4}$ in calidissimis prope Honda Novo-Granatensium. *Gymnolomia Hondensis* H. B. et Kunth nov. gen. am. 4. p. 218. Cl. auct. dubitat an à *G. tenellâ* satis differt, sed, ex icone specim. authenticæ ab amic. Ad. Brongniart missâ, satis distincta pappo videtur.
9. *G. TRIPLINERVIA*, ramis teretibus glabresculis junioribus adpresso-pilosis, foliis petiolatis ovatis angustato-acuminatis crenato-dentatis basi acutis triplinerviis utrinque adpresso-hispidulis, capitulis axillaribus longè pedunculatis, invol. squamis oblongo-lanceolatis laxè imbricatis, acheniis obovatis brevibus pappo coroniformi inæqualiter acutè et longiusculè dentato coronatis. $\frac{1}{2}$? in altâ planitie Bogotensi. *Gymnolomia triplinervia* H. B. et Kunth nov. gen. am. 4. p. 219.
10. *G. ? EUXENIOIDES*, erecta setulis adpressis scabra, foliis ovato-lanceolatis basi cuneatis acuminatis serratis, capitulis 2-3 ad apices ramorum, invol. aequali polyphylo. — in Brasiliâ ad viam Felisbertiam. *Euxenia radiata* Nees et Mart. act. soc. nat. cur. 12. p. 7. Ab *Euxeniâ* differt capitulis radiatis et ipsis docent. auct. accedit ad *Gymnolomiam*.

+ *Species exclusa.*

GYMNOLOMA MACULATUM Ker. = *Wulffia maculata*.

CCCXXVIII. *WULFFIA* Nees. elem. 1. p. 35. Cass. dict. 46. p. 398 et 403. — *Chakiatella* Cass. dict. 43. p. 37 et 46. p. 402. — *Chylodia* Rich. ex Cass. Journ. phys. 1823. mai. p. 214. non Brown. — *Gymnoloma* Ker bot. reg. t. 662. Less. syn. 227 non Kunth. — *Coreopsis*, *Helianthi*, *Verbesinæ* et *Cæsaliiæ* sp. auct.

Capitulum multiflorum sæpius heterogamum, fl. radii 1-serialibus ligulatis neutris, rarius homogamum discoideum fl. tubulosis hermaphr. Invol. subhemisphæricum, squamis laxè imbricatis sublanceolatis. Recept. subconvexum, paleis lanceolatis subcarinatis planisve persistentibus. Styli rami cono superati. Achænia radii abortiva, disci crassa obtusè tetragona plus minus carnosæ obtusæ, apice umbilicata calva glabra aut apice summo velutino-villosa. — Herbæ Austr.-Americane. Caules erecti aut subsarmentosi sæpius scabro-asperi sæpè (fortè semper) maculati. Folia opposita petiolata latè ovata acuminata serrata, nervis ad hasin folii utrinquè 3-pinnatis tertio majore et indè folia pseudo-tripinnaria. Capitula pedicellata, terminalia aut axillaria, sæpè terna, post anthesin subglobosa. Flores flavi. — Genus admodum naturale!

* *Capitulis radiatis.*

1. *W. MACULATA*, caule erecto tetragono vix subscabro maculato, foliis oblongo-lanceolatis basi cuneatis apice acuminatis serratis supernè scabris subtus glabrisculis, pedunculis ternis terminalibus, ligulis circ. 8 involvno duplò longioribus. \mathcal{L} in Brasiliâ (Ker), circa Rio de Janeiro (Lund!). *Gymnoloma maculatum* Ker bot. reg. t. 662. Petioli 3-4 lin. longi. Flores aurantiaco-flavi. (v. s.)
2. *W. STENOGLOSSA*, caule erecto subsarmentoso tetragono scabrido submaculato, foliis latè ovatis acuminatis basi vix cuneatis serratis utrinquè scabris asperimis membranaceis, pedunculis ternis ad apices ramorum et ramulorum axillarium, ligulis 8-10 invol. duplò longioribus, invol. squamis ext. ovalibus adpressè hirtellis. \mathcal{L} in Cayennâ. *Verbesina membranifolia* Poir. suppl. 3. p. 18. *Helianthus sarmentosus* Rich. act. soc. h. nat. Par. p. 105. Pers. ench. 2. p. 475. n. 9. *Verbesina oppositiflora* Poir. l. dict. 8. p. 460. *Chakiatella stenoglossa* Cass. dict. 29. p. 490 et 46. p. 406. *Chylodia sarmentosa* Rich. ined. ex Cass. *Coreopsis baccata* Linn. f. suppl. 380? *Pascalia baccata* Spreng. syst. 3. p. 602? Petioli 7-8 lin. longi. Ligulæ angustæ pallidè flavæ. (v. s.)
3. *W. PLATYGLOSSA*, caule erecto striato-sulcato adpressè hirtello, foliis petiolatis latè ovali-lanceolatis serratis utrinquè scabridis, capitulis ternis terminalibus pedicellatis, pedicello medio breviorè, invol. squamis ext. ovatis foliaceis, int. paleiformibus, paleis oblongo-lanceolatis achænia amplect., ligulis paucis invol. vix longioribus. \mathcal{L} in Guianâ gallicâ legit cl. Martin. *Crodisperma aspera* Poit. in h. Gay. ex Cass. *Chakiatella platyglossa* Cass. dict. 46. p. 403. Cor. flavæ. Ligulæ luteæ plurinerviæ. Folia petiolo excepto 5 poll. longa. 3 poll. lata. Stirps à cl. Poiteau lecta et à cl. Cassinio descripta tantum à stirpe Martini differt foliis paulò angustioribus, obs. cl. Ad. Brongniart. (v. s. comm. à Mus. reg. Par.)
4. *W. HAVANENSIS*, caule hexagono erecto scabro, foliis petiolatis latè ovatis acuminatis basi obtusis subcordatis callosè et remotè dentatis utrinquè asperis, nervis 3 inferioribus approximatis, capitulis ad apices ramorum et ramulorum axillarium ternis, invol. squamis ext. lanceolatis paleisque apice subuncinatis, ligulis 9-12. \mathcal{L} in insulâ Cubâ juxta Havanam legit cl. Ant. de La Ossa. Petioli 5 lin. longi. Limb. fol. 5-6 poll. longus, $2\frac{1}{2}$ 3 poll. latus. Ligulæ pallidè flavæ. (v. s.)
5. *W. SALZMANNI*, caule tereti striato subscabro, foliis breviter petiolatis ovatis acuminatis basi obtusis serratis utrinquè scabris, nervis 3 infer. approximatis, capitulis ad apices ramorum ternis medio brevius pedicellato, invol. squamis ob-

longis obtusiusculis, paleis oblongis glabris subcoloratis, ligulis 7-10 disco paulò longioribus. $\frac{1}{2}$ in umbrosis circa Bahiam Brasiliæ legit. cl. Salzmänn qui fructum verè baccatum in vivo vidit. Caulis 4-pedalis. Petioli vix 3 lin. longi. (v. s.)

6. W. *oblongipetala*, caule hexagono erecto subpubero, foliis petiolatis oblongo-lanceolatis acuminatis serratis basi acutis suprà sparsè puberis subtùs molliter villosis, capitulis terminalibus ternis, paleis rigidis apice uncinato-recurvus, ligulis parvis paucis albidis. $\frac{1}{2}$ circa Rio-Janeiro legit fructiferam cl. Gaudichaud et floridam ni fallor vidi in h. Mus. imp. Bras. sub. n. 120. ex prov. Mato-Grosso. Petioli 3 lin. longi. Limb. fol. 2 poll. longus, 7 lin. latus. (v. s.)

** Capitulis discoideis.

7. W. *Blanchetii*, caule teretiusculo scabro, ramis subangulatis, foliis brevissimè petiolatis ovato-lanceolatis serratis utrinquè scabris basi obtusis aut subcordatis, capitulis terminalibus ternis breviter pedicellatis, invol. squamis lanceolatis subscabris, ligulis nullis. $\frac{1}{2}$ in Brasilià circa Bahiam legit cl. Blanchet. Petioli 1-2 lin. longi. Limb. fol. 2 $\frac{1}{2}$ poll. longus, pollicem latus. Cor. intensè luteæ. (v. s.)
8. W. *scandens*, caule scandente subhexagono subaspero, foliis petiolatis ovatis valdè acuminatis subserrulatis utrinquè scabris basi obtusis, capitulis ad apices ramorum pedicellatis, invol. squamis ovalibus subacutis, paleis acutis rectis, ligulis nullis. $\frac{1}{2}$ in Peruvia legit cl. Poeppig. Cæsulia scandens Poepp. ! pl. exs. n. 24. diar. 1663. Petioli 9 lin. longi. Limb. fol. 4 poll. longus, 20 lin. latus. Cor. intensè flava. Fructum non vidi. (v. s.)

CCCXXIX. MONTAGNÆA. — *Eriocoma* H. B. et Kunth nov. gen. am. (1820) 4. p. 267. t. 396. *Cass. dict.* 15. p. 193. *Less. syn.* 226. non Nutt. (1818). — *Montanoa* Lallav. et Lex. nov. veg. mex. 2. (1825) p. 11. *Less. syn.* 420. — *Eriocarpa* Cass. dict. 59. (1829) p. 236. — *Priestleya* fl. mex. ic. ined. non DC.

Capitulum multiflorum heterogamum, fl. radii 5-10 uniseriatis ligulatis neutris, disci tubulosi 5-dentati hermaphrod. Invol. biseriale, squamis ext. 5 oblongis patulis, inter. sæpius 10 paleas referentibus. Recept. convexum, paleis basi latius apice spinoso-mucronatis achænia plùs minùs involventibus. Styli disci rami comò nunc brevi nunc longiusculo superati. Achænia radii abortiva, disci subcompressa obovata cuneata calva, nectario interdum prominulo. — Suffrutices fruticesve Americani plerique Mexicani. Rami teretes apice sæpius hirsuti. Folia opposita petiolata, ovata aut cordata, serrata aut lobata, sæpius subtùs subtomentosa. Capitula corymbosa. Ligulæ albæ (aut fortè roseæ). Discus albidus. — Nomen paulò emendavi ut melius cum nomine cl. doct. Lud. Montañæ (pronunt. Montagna) nec Montano congruat. Genus affine *Sclerocarpo*.

§. 1. *Eriocarphæ*, ligulis 5, paleis hirsutissimis.

1. M. *floribunda*, caule herbaceo? oppositè ramoso villosopubescente, foliis oppositis petiolatis deltoideo-ovatis propè basin triplinerviis subserratis suprà scabrisculis subtùs hirsutopubescentibus et canescentibus, corymbis fastigiatis polycephalis capitulis longè pedicellatis, invol. squamis biserialibus oblongo-lanceolatis obtusiusculis, paleis ovatis acuminatis albo-lanatis, ligulis 5. $\frac{1}{2}$? in Mexico inter urbem et sanctuarium Guadalupense. *Eriocoma floribunda* H. B. et Kunth nov. gen. am. 4. p. 267. t. 396.
2. M. *tomentosa* (Lallav. et Lex. nov. veg. mex. 2. p. 11), caule fruticoso, ramis villosotomentosis, foliis oppositis petiolatis latè ovato-lanceolatis basi subtruncatulis apice acuminatis sinuatis incisive propè basin triplinerviis suprà subpubescentibus subtùs petiolisque villosotomentosis, corymbo composito fastigiato, invol. squamis, 5 ext. oblongis subobtusis dorso villosis disco sub-

æqualibus, inter. paleisque dorso hirsutissimis in acumen rectum spinulosum productis, ligulis 5. $\frac{3}{4}$ in Mexico (Alam.!), circa urbem Mexici ad ripas fossarum (Lallav.), ad Real del Monte (Nées), ad Tlapujahua (Keerl. in h. Mart.!), Vulgò *Cihuapastli* Mexic. dicitur. *Priestleya corymbosa* fl. mex. ic. ined. *Eriocoma fragrans* Don in Sweet br. fl. gard. ser. 2. t. 44. *Eriocoma heterophylla* Schrad. sem. h. Goett. 1833. p. 3. (v. s.)

β. *cordifolia*, foliis basi latè et profundè cordatis, petioliis hinc indè alato-appendiculatis $\frac{3}{4}$ in horto Mexicano legit cl. Berlandier (pl. exs. n. 574!), Folia supernè glabriuscula, subtùs minùs tomentosa, magn. ferè dupla. (v. s.)

γ. *ternifolia*, foliis terno-verticillatis oblongis basi obtusis apicè acuminatis, parçè dentatis. $\frac{5}{8}$ in Mexico ad Chapolopa legit cl. Berlandier (pl. exs. n. 574!). Unicum specim. habeo et floribus junioribus donatum, undè dubius hæreo an sit spec. aut var. (v. s.)

§. 2. *Acanthocarpæ*, ligulis circ. 10. paleis parçè villosis evidentiùs mucronatis demùm spinulentibus.

3. *M. SPECIOSA*, caule herbaceo striato subangulato glabro ad apices subvillosis, foliis oppositis sessilibus basi in petiolum alatum crispum subsinuatum attenuatis longè supra basin triplinerviis irregulariter pinnatilobatis (lobis acuminatis subintegris) supra villosis subtùs molliter et densè villosis-tomentosis, ramis oppositis paniculatis nudis 1-cephalis, ligulis 10 elongatis, paleis recept. convexi basi latis apice spinulenti-subulatis $\frac{2}{3}$? in Mexico ad Cuernavaca octob. flor. legit cl. Berlandier (pl. exs. n. 1057). Folia 8-9 poll. longa, 3 poll. lata. Capitula diam. sesquipollic. (v. s.)

4. *M. GRANDIFLORA*, suffruticosa oppositè ramosa, ramis apice pubescentibus, foliis in petiolum alatum dentatum attenuatis longè supra basin triplinerviis supra strigilloso-scabris subtùs puberulis, infer. irregulariter pinnatifidis, lobis utriusquè 2-3 lato-lanceolatis subobtusis, summis ovato-lanceolatis indivisis, capitulis ad apices ramorum ternis pedicellatis, invol. squamis lineari-lanceolatis reflexis, paleis parçè villosis basi latis in spinulam productis, ligulis invol. longioribus circ. 10. $\frac{5}{8}$ in Mexico. *Eriocoma grandiflora* Alam. in litt. 1831. *Priestleya longifolia* fl. mex. ic. ined.? Ligulæ pollicem longæ apice grossè 2-3 dentatæ. (v. s. comm. à cl. Alam.)

5. *M. KARVINSKII*, caule herbaceo? tereti glabriusculo, foliis oppositis petiolatis ovatis obtusè sinuatis basi 3-5-nerviis aut subtriplinerviis parçissimè pubescentibus, paniculis laxis polycephalis, invol. squamis lineari-lanceolatis acuminatis inæqualibus, ligulis 10 obovato-oblongis $\frac{2}{3}$? in Mexico legit cl. de Karvinski. A duobus prioribus differt petiolis evidentissimis omninò immarginatis, à sequentibus foliis sinuatis seu obtusè lobatis. Ligulæ 5-6 lin. longæ, 2-3 lin. latæ. (v. s.)

6. *M. FRUTESCENS* (Maire! in litt. 1833), caule suffruticoso glabro, foliis oppositis et super. alternis breviter petiolatis lanceolatis acutis basi in petiolum non productis triplinerviis subserratis supra asperis subtùs glabris, capitulis laxè corymbosis, invol. uniseriatis squamis 5 obtusis disco brevioribus, paleis recept. glabris demùm convolutis apice recurvo rigido achenium involv. et multò superantibus, ligulis 8-10 invol. longioribus. $\frac{3}{4}$ in Mexico (Alam.! Maire!), circa urbem (Berl. pl. exs. n. 908), et in valle Tolucanà (Berl. n. 1154). *Priestleya squarrosa* fl. mex. ic. ined. *Eriocoma frutescens* Alam. in litt. Ligulæ 6 lin. longæ. Paleæ per anthesin breves, dein longè accretæ et capitulum fructiferum squarrosam formantes. (v. s.)

7. *M. ARBORESCENS*, caule fruticoso oppositè ramoso, ramis pubescenti-villosis, foliis oppositis petiolatis lanceolatis basi acutis apice acuminatis subintegerrimis subtriplinerviis supra asperis subtùs puberulo-hirtellis, corymbis laxis trifidis, capitulis longè pedicellatis, invol. squamis biseriatis oblongis disco ferè æqualibus, paleis parçè villosis mucronatis demùm apice induratis inflexis, ligulis circ. 8. $\frac{3}{4}$ in Mexico (Alam.! Maire!), ad Cordilleram Guabillaqua (Berl.)

pl. exs. n. 1006). *Eriocoma arborescens* Alam. in litt. 1831. *Montanoa floribunda* Cerv. in h. acad. Monac. Ligulæ 7-8 lin. longæ, 3 lin. latæ obtusæ. (v. s.)

8. *M. OVALIFOLIA*, caule fruticoso subscandente ad apices ramorum molliter hirsuto, foliis oppositis petiolatis ovalibus ovatisve acutis propè basin triplinerviis parçè serratis suprà petiolisque sparsè villosis subtus villosis-tomentosis canescentibus, capitulis corymbosis longè pedicellatis, invol. squamis ext. oblongis hirsutis subreflexis, inter. paleisque parçè villosis in acumen spinulosum productis, ligulis circ. 10. ♂ ad Santa-Fé de Bogota. Ligulæ ex sicco fortè roseæ, 5-6 lin. longæ. (v. s. comm. ab amico. Benj. Delessert.)

CCCXXX. *SCLEROCARPUS* Jacq. act. helv. v. 9. (1786) p. 34. t. 2. f. 1. Cass. dict. 48. p. 148. Thonn. in Schum. pl. guin. 394.

Capitulum multiflorum heterogamum, fl. radii 3 neutris minimis irregulariter ligulatis ante squamas inv. interiores sitis, disci 8-12 tubulosis hermaphroditis, centralibus sæpè abortivis. Invol. duplex, squamis ext. foliaceis subpetiolatis majoribus (fortè bracteis), int. 3 minoribus distantibus. Recept. parvum convexum, paleis disci flores involventibus et iis brevioribus. Cor. disci pubescens 5-dentata persistens. Anth. vix concretæ apice in appendicem subulatam productæ. Styli rami subulati hispidi. Achænia radii abortiva tenuissima, disci obovata lævia calva intrà paleam curvato-subcomplicatam induratum intus hiantem apice in collum contractam inclusa. — Herba Africana erecta. Caulis scaber teres apice præsertim ramosus. Folia alterna petiolata ovata acuminata serrata triplinervia suprà villosa scabra subtus puberula. Capitula ad apices ramulorum solitaria subsessilia. Cor. flavæ.

1. *S. AFRICANUS* (Jacq. ic. rar. 1. t. 176), foliis latè ovatis acutis serratis. (1) in Guineâ (Thonn.), et in cultis inundatis regionis Walo ad flum. Senegal (Perrottet!); etiâ sed verisim. introductum in Indiâ orient. legit cl. Royle et ad Baranquillas Americæ austr. cl. Bertero. Hic ultimus est *Melampodium Baranquillæ* Spreng. syst. 3. p. 619! (v. s. sp. et v. c.)

† *Species valdè dubiæ.*

2. *S. ? GRACILIS* (Smith in Rees cycl. v. 31), foliis lineari-lanceolatis integris, pedunculis 4-5 ad apices caulis foliosis. — ad oram boreali-occid. Americæ septentrionalis. Paleæ fructiferæ glandulis prominentibus onustæ. Cor. ign. et plerique car. valdè incerti. †.
3. *S. ? EXIGUUS* (Smith l. c.), foliis lineari-lanceolatis integris, pedunculis nudis folio triplò longioribus. (1) cum priore. Herba 3-4 pollic. Folia ferè *Myosotidis*. Flores et fructus vix noti. †.

CCCXXXI. *ENCELIA* Adans. fam. 2. p. 128. Cass. dict. 14. p. 447. 59. p. 139. Less. syn. 227. — *Pallasia* L'Hér. diss. 1784. Ait. kew. ed. 1. v. 3. p. 253 non Linn. — *Coreopsis* id. Jacq.

Capitulum multiflorum radiatum, ligulis 1-serialibus neutris, fl. disci hermaphr. 5-dentatis. Invol. biseriatum disco æquale subimbriatum. Recept. planum (nec conicum ut vult Less.), paleis membranaceis navicularibus. Styli rami cono superati. Achænium plano-compressum margine albo-lanatum emarginatum calvum totum villosum. — Suffrutices Americani ramosi plus minus, saltem juniores, canescentes. Folia alterna petiolata ovata aut oblonga tri- seu triplinervia integra aut subrepanda. Capitula pauca pedicellata in paniculam disposita. Cor. flavæ.

1. *E. CANESCENS* (Cav. ic. 1. p. 45. t. 61), erecta, foliis latè ovatis integris obtusis molliter canescentibus, invol. squamis villosis ciliatis. $\frac{3}{4}$ in Peruvîa (Adans. Gaud.!), circa Limam (Domb.!), nec verisim. in Mexico (ut dixit Cavanilles ex hort. Madr.) Lindl. bot. reg. t. 909. Kunth nov. gen. am. 4. p. 206. *Coreopsis Limensis* Jacq. ic. rar. 3. t. 594. *Pallasia halimifolia* L'Hér. herb. et mss. Willd. sp. 3. p. 2260. *E. alternifolia* et fortè *Eucalia canescens* Raensch. nom. 251? (v. v. c. et s. sp.)
2. *E. PARVIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 206), procumbens, ramulis sericeo-pubescentibus, foliis rhombo-ovatis acutis basi-cuneatis canescenti-pubescentibus. $\frac{3}{4}$ in Peruvîa ad littora Oceani pacifici. *Pallasia procumbens* Spreng. syst. 3. p. 610. Frutex humilis ramosissimus proximus *E. canescenti*.
3. *E. HALIMIFOLIA* (Cav. ic. 3. p. 6. t. 210), erecta, foliis ovatis subrepandis dentatisve glabrisculis margine valdè ciliatis. $\frac{3}{4}$ in Mexico ex Cav., sed verisim. potius in Peruvîa. Cass. dict. 14. p. 448. *Pallasia grandiflora* Willd. sp. 3. p. 2261. (v. v. c. in h. Par.)
4. *E. OBLONGIFOLIA*, erecta, foliis oblongis utriusque subattenuatis integerrimis canescentibus, invol. basi præsertim hirsutissimis. $\frac{3}{4}$ in Chili (Hænk.!), circa Coquimbo (Gaudichaud! Macrae.!). Ligulæ elongatæ lineares. Folia var. 1-2 pollic. longa, 4-8 lin. lata. (v. s.)

† *Species exclusæ.*

E. DENTATA Poir. = *Verbesina dentata*?

E. MEXICANA Mart. = *Simsia auriculata*.

CCCXXXII. PHILOGLOSSÆ.

Capitulum multiflorum heterogamum, fl. radii ligulatis innumeris 1-2- $\frac{1}{2}$ -seriatis neutris, disci tubulosi hermaphroditis ad medium 5-fidis. Invol. 4-5-seriale, squamis ext. inter se alternis biseriatis lanceolatis foliaceis ciliatis, inter. 3-ser. membranaceis albidis glabris lanceolatis. Recept. planum, paleis lineari-subulatis scabridis caducissimis. Ligulæ basi hirtæ angustissimæ. Cor. disci tubo brevi, lobis 5 acuminatis profundis. Anth. nigricantes. Stylus apice summo subbifidus longè hispidus. Achænia obovata lævia calva, areolâ terminali minimâ. — Herba *Limensis* villosa-hispida. Caulis adscendens. Folia caulina opposita, ramorum flor. quaterno-verticillata, ovalia aut obovata basi cuneata sessilia tenuissimè triplinervia integra aut subdentata. Pedicelli filiformes hispidi axillares terminalesque 1-cephali, anthesi ineunte folio breviores, demùm longiores. Flores lutei ferè calendulacei.

1. *P. PERUVIANA*. (1)? in Peruvîa circa Limam legit cl. Dombey et recentius cl. Abadia. (v. s.)

CCCXXXIII. CHRYSOSTEMMA Less. syn. 227. — *Coreopsis* sp. Linn. —
Anacis sp. Schrank.

Capitulum multiflorum heterogamum, fl. radii neutris 1-serialibus ligulatis, disci hermaphr. tubulosi 5-dentatis. Invol. biseriale, squamis foliaceis basi coadunatis, ext. paucis parvis linearibus patentibus, inter. erectis ovali-oblongis margine subscariosis. Recept. planum, paleis linearibus angustissimis ferè filiformibus. Styli rami cono superati pubesque descendente instructi. Achænium plano-obcompressum obovato-ellipticum angulis alatum glaberrimum apice emarginatum, pappo coroniformi lacero coronatum, disco epigyno minuto, alis albidis. — Herba Boŕeali-Americana nitida erecta. Folia opposita petiolata, imæ

subpedato-pinnatifida; super. trisecta, segmentis lanceolatis integerrimis. Pedunculi subgemini 1-cephali. Capitula radio luteo, disco brunneo.

1. *C. TAPPEZIS* (Less. l. c.). \mathcal{U} in Americæ bor. montibus altis à Virginiâ ad Carolinam (Parsh), et in Georgiâ occid. (Ell.). *Coreopsis tripteris* Linn. sp. 1282. Michx. fl. bor. am. 2. p. 138. Ell. sketch 2. p. 442. *Anacia tripteris* Schrank. in akad. Münch math. et nat. 5. p. 7. — Moris. oxon. 3. s. 6. t. 3. f. 44. (v. v. c.)

CCCXXXIV. *CALLIOPSIS* Reichenb. mag. n. 70. Cass. dict. 59. p. 321. Less. syn. 228. — *Diplosastera Tausch hort. Can. fasc. 1.* — *Calliopsisid* sp. Spreng. *Coreopsisid* sp. Nutt.

Capitulum multiflorum heterogamum, fl. radii neutris 1-serialibus ligulatis grossè 5-5-dentatis, fl. disci hermaphr. tubulosis 5-dentatis. Invol. biseriale, squamis ext. brevibus squarrosis, int. majoribus erectis coadunatis. Recept. planum paleis linearibus deciduis floribus brevioribus onustum, Stylus disci ramis truncatis apiceque solo penicillatis. Achænia obcompressa calva incurva truncata, disco epigyno minuto. — Herbæ Boreali-Americanæ elegantes glabræ. Folia opposita uni- aut bipinnatisecta, lobis integris. Capitula ramos terminantia, radio luteo basi fusco-purpureo maculato, disco luteo aut subpurpureo. — Genus vix à *Coreopsis* differt.

1. *C. TINCTORIA*, glabra, foliis radicalibus pinnatipartitis sen bipinnatipartitis lobis integerrimis, superioribus tripartito-multifidis lobis linearibus, invol. ext. brevissimo acuto, ligulis trifidis, invol. int. duplè longioribus, achæniis oblongis exalatis utrinquè minutè tuberculatis. ① in territorio Arkansano ad ripas flum. Rubri in pratis inundatis detexit cl. Nuttall. *Coreopsis tinctoria* Nutt. journ. acad. Philad. 1821 oct. p. 114. Radius schrif. ges. Leips. 1822. p. 159. t. 3. Bart. fl. am. bor. 2. t. 45. ex bot. mag. t. 2312. bot. reg. t. 846. Link et Otto abb. t. 33. Sweet fl. gard. t. 72. *Diplosastera tinctoria* Tausch hort. canal. fasc. 1. icon et descr. *Calliopsis bicolor* Reichenb. mag. t. 70. Ligulæ basi fuscæ caterum auræ. Herba elegans in hortis ubiquè culta et jam hinc inde in Europâ seminibus hortensibus sata. (v. v. c.)
2. *C. CARDAMINEFOLIA*, foliis infer. uni- aut bipinnatisectis lobis obovatis oblongis sæpè trifidis, super. bipinnatisectis lobis lineari-spathulatis, invol. ext. brevissimo obtusiusculo, ligulis trifidis invol. interiore duplè longioribus, achæniis alatis utrinquè lævibus aut minutissimè tuberculatis. ① in Mexici prov. Texas inter Bejar et flum. Trinitatis, ad Matamoros et ad lacum Sancti-Nicolai in sinu Sancti-Spiritus legit cl. Berlandier (pl. exs. n. 1668, 1956, 1478, et 2312). Ligulæ basi fuscæ apice auræ ut in *C. tinctoriâ* ad quam accedit, sed achæniis diversa. (v. s. sp.)
3. *C. ATKINSONIANA* (Hook. fl. bor. am. 1. p. 311), foliis radicalibus bipinnatipartitis lobis integerrimis, caulinis pinnatisectis lobis linearibus subspathulatis, invol. ext. squamis lineari-oblongis margine submembranaceis, ligulis invol. int. triplè longioribus obtusè 3 dentatis, achæniis alâ brevi marginatis utrinquè lævissimis. \mathcal{U} ad ripas fluminis Colombiæ Amer. bor. et insul. Mewries s. Menzies, legit cl. Douglas. *Coreopsis Atkinsoniana* Dougl. in Lindl. bot. reg. t. 1376. Similis *C. tinctoriæ* sed elatior, perennis, et achæniis distinctissima. (v. v. c.)

CCCXXXV. *PERAMIBUS* Raf. ann. nat. 1 (1820), p. 14. Cass. dict. 38. p. 416. — *Coreopsisid* sp. Nutt. — *Calliopsisid* sp. Spreng.

Omnia *Coreopsisid* sed achænia triangularia lævia nuda. — Genus valdè incertum, soli Rafinesquio notum et botanicis Americanis commendatum. †.

1. *P. HIRTUS* (Raf. l. c.), hirtus, caule angulato flexuoso ferè dichotomo, foliis semiamplexicaulibus lanceolatis acuminatis medio serratis, inferioribus ovato-lanceolatis, floribus ferè corymbosis subsessilibus, invol. squamis lineari-lanceolatis acutis, ligulis ellipticis. ② in collibus ad flum. Liking in Kentucky. *Coreopsis hirta* Raf. ex ipso. †.
2. *P. SCABRA* (Raf. l. c.), ramis remotis elongatis subulatis folisque imis oppositis, summis alternis, foliis integris (lanceolatis?) scaberrimis, achæniis oblongis angulatis. ② in Louisianâ. *Coreopsis scabra* Raf. fl. lud. 72. Herba bipedalis. An eadem ac sequens ex Raf.? †.
3. *P. ACUTUS* (Raf. l. c.), foliis ovato-lanceolatis acutis denticulatis subhirtis, capitulis corymboso-paniculatis. ② in Georgiâ. *Coreopsis acuta* Pursh fl. bor. am. 2. p. 569. Cæt. ign. †.

Div. III. *COREOPSIDÆ* Less. syn. p. 228. — *Coreopsis* et *Helianthæ* prototypæ Cass. opusc. 3. p. 58 et 59.

Capitula heterogama, fl. radii neutris. Achænia erostris pappo nunc aristis 2-4 tantum, nunc aristis et squamellis constante.

CCCXXXVI. AGARISTA.

Capitulum multiflorum radiatum, ligulis 1-seriatis neutris obovato-truncatis multinerviis, fl. disci tubulosi 5-dentati hermaphrod. Invol. campanulatum duplex, squamis acutis, ext. 4 latè ovatis basi subconcretis, inter. 8 ovali-oblongis duplò ferè longioribus submembranaceis. Recept. paleæ cum fructu deciduæ, ad latus hujus ext. applicatæ, lineari-lanceolatæ subhyalinæ. Styli rami cono superati. Achænia obcompressa, radii ovalia inania glabra calva, disci oblonga sub paleâ glabra, margine utroque et facie interiore villosissima. Pappus bisquamellatus, squamellis elongatis membranaceis corollâ parum brevioribus ex angulis achæniis ortis. — Herba Californica glaberrima annua erecta *Calliopsidis* seu *Coreopsidis* facie. Caulis teres ramosus. Folia alterna multifida rachi lobisque linearibus elongatis. Rami apice aphylli 1-cephali. Capitula erecta aurea. Genus differt ab *Heliantheis* ferè omnibus pappo ut in *Helenieis* squamellato nec aristato, sed tamen affinitate naturali ad hunc locum revocatum. — *Agarista* mythologicè erat *nympha* venustissima.

1. *A. CALLIOPSIDEA*. ① in Californiâ legit cl. Douglas. Ligulæ circ. 8 basi cuneatæ, 8 lin. longæ, 4 lin. latæ, lineis 15-20 striatulæ. (v. s. comm. ab hon. soc. hort. Lond.)

CCCXXXVII. *COREOPSIS* Linn. gen. n. 981. (excl. spec.) Less. syn. 228. — *Coreopsis* et *Coreopsoides* Mærch meth. p. 593 et 594. — *Coreopsis* et *Leachea* Cass. dict. 10. p. 419. 25. p. 388 et 59. p. 320 et 321. — *Coreopsis* et *Anacis* sp. Schrank. — *Coreopsis* et *Chrysometea* Tausch. — *Coreopsis* et *Acispermum* Neck. elem. n. 63 et 64.

Capitulum multiflorum heterogamum, fl. radii circ. 8 neutris ligulatis, disci hermaphr. 5-dentati. Invol. duplex utrumque circ. 8-phyl-lum imâ basi subcoadunatum, squamis ext. foliaceis angustioribus subpatulis, int. latioribus submembranaceis. Recept. planum paleaceum, paleis persistentibus. Styli rami apice hispidi cono superati. Achænia obcompressa-plana, rarò incurva, margine plus minus alata, apice nunc bidentata, nunc biaristata, aristis lævibus aut sursum ciliato-serratis (nec retrorsum pilosis). — Herba Americanæ rarò suffrutescentes ple-

rumquæ glabræ. Folia opposita, rarius alterna. Pedunculi 1-cephali solitarii aut corymbosi. Capituli discus luteus, radius concolor raro albus.

* *Foliis alternis.*

1. *C. ANGUSTIFOLIA* (Ait. kew. 3. p. 253), glabra, foliis alternis lineari-lanceolatis integerrimis lævibus, ligulis oblongis trifidis, laciniâ mediâ majore. 2/ in Carolinâ et Floridâ. Ait. (v. s. ?) †.
2. *C. GLADIATA* (Walt. car. p. 215), caule glabro supernè dichotomo, foliis alternis in petiolum semiamplexicaulem attenuatis angusto- aut ovali-lanceolatis puberulis glabrisve integerrimis, invol. squamis exter. lineari-lanceolatis inter. brevioribus, ligulis 3-dentatis, achæniis obovatis alatis, alis serrulatis, pappo bisetoso. 2/ in pinetis humidis et sphagnosis umbrosis à Novâ-Cæsariâ ad Carolinam (Pursh). *C. gladiata* Nutt. gen. am. 2. p. 180. Ell. sketch 2. p. 444. *C. dichotoma* Michx. fl. bor. am. 2. p. 137. Variat foliis angustioribus pilosis et latioribus glabris. (v. s. sp.)
3. *C. ASPERA* (Pursh fl. bor. am. 2. p. 570), foliis lanceolato-linearibus asperis, superioribus alternis, inferioribus oppositis, caule monocephalo. — in Maryland. Cæ. ign.

** *Foliis oppositis indivisis.*

4. *C. LINEARIFOLIA*, caule suffruticoso adscendente glabro simplici basi folioso in pedunculum nudum 1-cephalum longè producto, foliis oppositis linearibus integerrimis basi parcè ciliatis, invol. ext. squamis linearibus glabris, int. lanceolatis hirsutis, ligulis ellipticis multiradiatis nervulis alternis brevioribus, styli ramis cono acuto superatis. 2/ 3 in Mexico ad Tlapujahua legit cl. Keerl ex h. Mart! Species distinctissima, vix pedalis. Ligulæ intensè flavæ nigro striatæ. Discus fuscus. Achænia ex statu immaturo omninò calva ut in Calliopside sed styli rami cono ut in Coreopside superati. (v. s. in h. Mart.)
5. *C. VENUSTA* (H. B. et Kunth nov. gen. am. 4. p. 229), fruticosa glabra, foliis oppositis linearibus acutiusculis integerrimis uninerviis subcoriaceis, capitulis terminalibus solitariis longè pedunculatis, invol. squamis exter. oblongis obtusis patentibus, ligulis oblongis subtridentatis, achæniis (junioribus) margine albo-lanatis biaristatis, aristis adpresso-pilosis. 3/ prope Loxam et in scopulis Parami de Saragura. Ligulæ aurantiacæ, discus flavus.
6. *C. LANCEOLATA* (Linn. sp. 1283), caule tereti striato, foliis oppositis cuneato- seu oblongo-lanceolatis integerrimis basi ciliatis, infer. petiolatis, pedunculis longissimis simplicibus nudis, invol. squamis exter. lanceolatis interiores subæquantibus, ligulis apice 5-dentatis, achæniis orbiculatis alatis scabridis biaristulatis. 2/ in Carolinâ (Linn.) et Virginîâ (Pursh), ad lacum Superiorem et Huronensem (Hook.). *Leachea lanceolata* et verisim. *L. crassifolia* Cass. dict. 25. p. 388. *Chrysomelea lanceolata* Tausch hort. Canal. f. 1.
 - a. *succisæfolia*, caule erectiusculo ramoso foliisque glabris, ligulis amplissimis. — Hæc varietas hortensis videtur typus speciei et generis. Dill. elth. 55. t. 48. f. 56. Linn. hort. cliff. 420. Ligulæ amplæ 10-12 lin. longæ. (v. s. c.)
 - b. *glabella* (Michx. fl. bor. am. 2. p. 137), caule abbreviato glabro, foliis margine ciliatis utrinquè glabriusculis angustioribus, ligulis mediocribus. 2/ in humidis Carolinæ. Foliorum limbus circ. pollicem longus, petiolo ferè æqualis. Ligulæ 6-7 lin. longæ. (v. s. sp.)
 - γ. *villosa* (Michx. l. c.), caule abbreviato hirsutissimo, foliis utrinquè canescenti-villosis, pedunculis glabris. 2/ in pinetis et locis siccioribus ex Ell. *Coreopsis crassifolia* Ait. kew. 3. p. 253. Ell. sketch 2. p. 434. (v. s. comm. à cl. B. Greene.)
7. *C. INTEGRIFOLIA* (Poir. suppl. 2 (1811), p. 352), caule glabro tereti subsimplici, foliis oppositis ovatis integris utrinquè acutis glabriusculis, inf. petiolatis petiolis imâ basi connatis, summis sessilibus, capitulis paucis trichotomocorymbosis, invol. ext. brevi subpatente, ligulis trifidis, achæniis breviter

biaristatis. — in Carolinā legit cl. Bosc. Coreopsis bidenz Bosc! herb. non Walt. Capitula lutea. Habitus Bidentis sed achæniis aristæ non retrorsum scabræ. Ovaria juniora tantum vidi. (v. s.)

8. *C. OXMLEX* (Ell. sketch 2. p. 435), glabra, caule angulato, foliis oppositis sessilibus lato-lanceolatis utrinque acutis integerrimis, pedunculis axillaribus terminalibusque subdichotomè corymbosis, invol. ext. brevi, ligulis integris, achæniis cuneatis marginatis leviter dentatis. — in Carolinā ad junctionem flum. Broad et Saluda. Affinis dicitur *C. latifoliæ*.

9. *C. LATIFOLIA* (Michx. fl. bor. am. 2. p. 137), foliis oppositis ovatis acuminatis breviter petiolatis dentatis, dentibus mucronatis, ligulis integris, achæniis cuneatis-oblongis apteris apice nudis. $\frac{1}{2}$ in montibus altis Carolinæ. Habitus Silphii sed capitula minora. — An Silphium glaucum Weinm. hort. dorp. 143? sed à Coreopside et à Silphio alienum esset si recept. verè nudum ut vult Weinmann.

10. *C. ANGUSTA* (Pursh fl. bor. am. 2. p. 567), glabra, foliis oppositis petiolatis lanceolato-ovatis sensim attenuatis argutè serratis, pedunculis axillaribus terminalibusque dichotomè corymbosis. $\frac{1}{2}$ in Carolinā ex Pursh, sed ab Elliott. non reperta. An ex Nutt. var. *C. latifoliæ*? Flores aurei. *C. serrulata* Poir. suppl. 2. p. 352 huc à Spreng. admissa, videtur potius *Bidens grandiflora*.

*** *Foliis oppositis partitis sectivæ.*

11. *C. MUTICA*, suffruticulosa glabra, foliis oppositis serratis, aliis ovato-lanceolatis acuminatis serratis, aliis trisectis, segmento medio folia indivisa referente, lateralibus minoribus, capitulis ad apices ramorum pedicellatis ternis, invol. squamis ext. oblongis, ligulis multistriatis, achæniis elliptico-oblongis submarginatis apice vix ac ne vix obtusè bidentatis. $\frac{5}{8}$ in Mexico ad Tlapujhua legit F. W. Keerl ex h. Mart. Flores lutei. Paleæ membranaceæ striatæ. (v. s. in h. Mart.)

β . *subvillosa*, ramulis foliis subtus involucrique squamis exterioribus villosis. — in Mexico legit cl. de Karvinski. (v. s. in h. Acad. reg. Monac.)

12. *C. AURICULATA* (Linn. sp. 1281), caule erecto ramoso breviter puberulo, foliis oppositis aliis indivisis aliis trisectis, lobis lanceolatis acutis integerrimis utrinque brevissimè puberulis, terminali triplò quadruplòve majore, petiolo ciliato, capitulis subsolitariis pedicellatis, invol. squamis ext. lineari-oblongis subacutis glabris demum reflexis, ligulis 4-5-dentatis, achæniis suborbiculatis alatis brevissimè biaristatis. $\frac{1}{2}$ in Amer. bor. montibus Virginie et Carolinæ. Schkuhr handb. 3. t. 260 opt. *C. auriculata* var. α . Ell. sketch 2. p. 436. *Lea- chea trifoliata* Cass. dict. 25. p. 389. *Coreopsoides lanceolata* Mönch meth. 594. *Anacis auriculata* Schrank akad. Münch. nat. 5. p. 7. *Chysomelea auriculata* Tausch h. Can. f. 1. — Pluk. alm. t. 83. f. 5. (v. v. c.)

β . *glabrata*, caule glabro, foliis glabriusculis, petiolis vix basi ciliatis. $\frac{1}{2}$ cum var. α in hortis sæpius culta, fortè var. ϵ est culturā aut loco orta. Utraque var. variat achæniis plùs minùs ut in *Calliopside incurvis*, intus extusve muricato-tuberculatis aut lævibus, aristis minimis mox deciduis aut fortè nullis. (v. v. c.)

13. *C. DIVERSIFOLIA* (DC. non Jacq.), caule cæspitoso brevissimo pilis retrorsis patulive hispido, foliis radicalibus oppositivè glabriusculis, aliis ovatis indivisis, aliis trisectis lobo medio ovato lateralibus multò minoribus, petiolis hispidulis limbo multò longioribus, pedunculis longissimis glabris, invol. squamis ext. lanceolatis, ligulis 4-5-dentatis, achæniis orbiculatis alatis, alà subdentatà. (Ell.) $\frac{1}{2}$ in Carolinā mediā (Whitl. in Ell.), et boreali (B. Greene!). *Coreopsis auriculata* β diversifolia Ell. sketch 2. p. 437. — Pluk. alm. t. 242. f. 4. Habitu et car. indicatis satis à *C. auriculatà* differe videtur. In specim. meo ut in icone Pluken. folia radicalia longè petiolata indivisa, caulina trisecta, sed Elliott dicit folia inf. trisecta, sup. indivisa. *C. diversifolia* Hook. bot. mag. t. 3474 in agro Texano à cl. Drummond lecta videtur eadem. (v. s. comm. à cl. Greene.)

14. *C. SENIFOLIA* (Michx. fl. bor. am. 2. p. 138), caule erecto trichotomo, foliis

oppositis sessilibus trisectis, segmentis ovato-lanceolatis integerrimis pube brevi confertâ utrinque velutinis membranaceis, capitulis subcorymbosis, invol. squamis ext. lineari-oblongis obtusis, ligulis oblongis integris, achæniis cuneato-oblongis angustè sctis glabris integerrimis, aristis minimis deciduis. \mathcal{L} in collibus Virginie (Pursh), Carolinæ et Georgiæ (Lyon.), Kentucky (Raf.!), in Carolinæ (Fras.!) sabulosis (Michx.), et pinetis (Ell.). Pursh fl. bor. am. 508. Ell. sketch 2. p. 438. *Coreopsis major* Walt. car. 214. *C. stellata* Banks herb. ex. Pursh. (v. s.)

β. rigida (Nutt. gen. am. 2. p. 180), glabriuscula rigidula, foliis infer. indivisis, super. tripartitis aut bipartitis imò subdivisis, segmentis angustè lanceolatis suprâ scabridis. — in Georgiâ (Nutt.), in Carol. merid. ex specim. à cl. Fraser et Mitchell missis. An fortè species? (v. s. sp.)

15. *C. DELPHINIIFOLIA* (Lam. dict. 2. p. 108), glabra, foliis oppositis sessilibus à basi trisectis, segmentis indivisis seu pinnatipartitis, lobis lineari-lanceolatis integerrimis, capitulis pedicellatis, invol. squamis exter. lineari-oblongis obtusis, ligulis integris acutis, disco discolorè, achæniis obovatis subulatis brevissimè bidentatis. \mathcal{L} in montibus altis et locis siccis Amer. bor. in Virginia (Pursh), Carolinâ (Ell.), et Georgiâ (h. Duby!). *C. verticillata* var. Linn. sp. 1281 (excl. syn. Pluk.). *C. verticillata* Willd. sp. 3. p. 2251, bot. mag. t. 156. Schkuhr handb. t. 260 b. f. sin. *Ceratoceph. Delphiniifolia* Vaill. act. ac. Par. 1720. Ehret. pict. t. 9. f. 1. Folia non sunt verticillata sed opposita. Variat caule foliisque glaberrimis aut minutè subpuberulis. Discus fuscus, ligulæ flavæ. (v. v. c. et s. sp.)

16. *C. TENUIFOLIA* (Ehr. beitr. 7. p. 168. Willd. sp. 3. p. 2252), glabra, foliis oppositis sessilibus trisectis, segmentis multipartitis, lobis linearibus integerrimis, capitulis pedicellatis, invol. squamis lineari-oblongis obtusis, ligulis subintegris acutis, disco concolore, achæniis obovato-cuneatis subulatis minutè bidentatis. \mathcal{L} in montibus et excelsis siccis Virginie (Pursh) et Carolinæ (Bosc!). Schkuhr handb. 3. t. 260 b fig. dextr. *C. verticillata* Linn. sp. 1281 (excl. syn. Vaill.). *C. verticillata* Lam. dict. 2. p. 108. *C. verticillata* β tenuifolia Michx. fl. bor. am. 2. p. 139. — Pluk. mant. t. 344. p. 4. *C. delphiniifolia* affinis, sed lobi foliis angustiores semper glabri, discus luteus, etc. (v. v. c. et s. sp.)

17. *C. GRANDIFLORA* (Hogg. mss. ex Sweet brit. fl. gard. 2. t. 175), caule erecto ramoso sulcato glabro, foliis oppositis sessilibus glabris basi ciliatis, infer. biter-natisectis, sup. trisectis tripartitisve, lobis linearibus elongatis integerrimis suprâ canaliculatis, pedunculis elongatis 1-cephalis, invol. squamis ext. lanceolatis acuminatis interiores æquantibus, ligulis acutè 5-dentatis, achæniis elliptici vix alati aristis subpaleaceis cristatis. \mathcal{L} in Americâ bor. sine loc. propr. design. à Noveboraco missa. Capitulum poll. 2 ferè diam. disco radioque luteis concoloribus. (v. s. in h. Barclay ubi etiâ Sweet suam vidit, à Filio coll.)

18. *C. MITIS* (Michx. fl. bor. am. 2. p. 140), glaberrima, caule obtusè tetragono brachiato, foliis oppositis bipinnatipartitis, lobis linearibus serrulatis, capitulis paniculatis, invol. ext. squamis linearibus serrulatis, ligulis obovatis apteris obscurè dentatis, achæniis oblongis serrulatis biaristatis. (2) ex Pursh. In demissis humidis Carolinæ (Michx.) et in agro Missourienti (Lew. ex Pursh). *C. coronata* Walt. car. 2. p. 15 ex Ell. sketch 2. p. 441.

19. *C. TRICHOSPERMA* (Michx. fl. bor. am. 2. p. 139), glabra, caule obtusè tetragono, foliis oppositis, lobis lanceolato-linearibus subdentatis, superioribus in limbum lineari-lanceolatum elongatum dentatum confluentibus, capitulis subcorymbosis, invol. squamis ext. oblongo-linearibus obtusis subciliatis, ligulis integris obtusis, achæniis planis puberulis, aristis cor. disci longitudinem subæquantibus. \mathcal{L} Ell. (2) Pursh. In humidis Carolinæ super. (Michx.). Novæ-Cæsareæ (Greenel), Bostoniæ (Big.), Noveboraci et Novæ-Cæsareæ (Torr.!). Habitus omninò Bidentis sed aristæ non retrorsum scabræ. (v. s.)

20. *C. AUREA* (Ait. kew. ed. 1. v. 3. p. 252), caule glabro striato, foliis oppositis tripartitis trisectisve pubescenti-pilosis, lobis linearibus grossè serratis aliis in-

- divisis aut terminali tripartito, invol. squamis ext. linearibus patentibus, ligulis amplis obovatis integris, achæniis cuneatis glabrisculis apice bicornibus. ② in Amer. bor. fortè circa Noveboracum. Lindl. bot. reg. 1228. *C. trichosperma* β anrea Nutt. gen. am. 2. p. 180. Capitula ampla aurea.
21. *C. ARISTATA* (Michx. fl. bor. am. 2. p. 140), pubescens, foliis oppositis quinato-pinnatis-sectis, lobis serratis, ligulis integris lato-ovalibus, achæniis cuneato-obovatis biaristatis, aristis longissimis divaricatis. \mathcal{L} in Carolinâ infer. (Pursh) et agro Illinoensi (Michx.). *C. aristata* Willd. sp. 3. p. 2253. Capitula ampla. Ligulæ latissimæ.
22. *C. PUBESCENS* (Ell. sketch 2. p. 441), pubescens, ferè subtomentosa, foliis quinato-pinnatis-sectis, lobis lanceolatis obtusis integerrimis, lateralibus parvulis, invol. ext. squamis ovato-lanceolatis glabris, ligulis undulatis? achæniis orbiculatis angustè alatis emarginatis, pappo diphylo brevi subulato pubescente. \mathcal{L} in Georgiâ occidentali.
23. *C. MULTIFIDA*, herbacea glabra erecta, foliis oppositis petiolatis circumscris, suborbiculatis pinnatipartitis, lobis pinnatifidis, lobulis lanceolatis brevibus hinc indè incisis, capitulis pedunculatis solitariis paucisve corymbosis, invol. squamis lanceolatis subæqualibus, ligulis ellipticis integris, achæniis (immaturis) linearibus apice attenuatis subciliatis in aristas 2 rigidas glaberrimas abeuntibus. — in Peruvîâ? legit cl. Pavon ex h. Thibaud. Hinc refero specimina duo: unum spontaneum ætate provecinus ligulis nullis verisim. deciduis, alt. hortense junius ligulis albidis cæterum simillimum. (v. s.)
- β . *mutica*, caule magis angulato, foliorum lobulis paulò acutioribus, achæniis exaristatis. — in Peruvîâ undè semina ad hort. Par. olim misit cl. Dombey. Var. α simillima præter notas indicatas. Ligulæ ex sicco vid. roseæ. (v. s. ex h. L'hér.)
24. *C. ANTHEMOIDES*, caule herbaceo demisso ramoso radicante subpuberulo, foliis oppositis glabris multifidis, lobis linearibus acutis, pedunculis terminalibus 1-cephalis erectis, invol. squamis ext. linearibus interiores subæquantibus, ligulis ellipticis integris, achæniis (immaturis) linearibus glabris brevissimè bidentatis, \mathcal{L} ? in Mexico ad Guachilapam in Cordillera legit cl. Berlandier (pl. exa. n. 1030), fl. octob. Herba demissa habitu ferè *Anthemidis* arvensis. Flores lutei. Ligulæ 3-4 lin. longæ. (v. s.)
25. *C. NEEI*, fruticulosa? ramis teretibus glabris trichotomis, foliis oppositis basi hirsuto-ciliatis lineari-filiformibus acutis, aliis indivisis, aliis tripartitis, capitulis pedunculatis solitariis, invol. extus hirsuto, squamis subæqualibus, ligulis elliptico-oblongis integris, achæniis glaberrimis brevissimè bidentatis, \mathcal{L} ? in Amer. calid. legit cl. Née, sed locus propr. ign. Fl. flavi. Ligulæ 6-7 lin. longæ. Achænia matura ign. (v. s. ex h. Thibaud. specim. manc.)
26. *CAPILLARÆA* (H. B. et Kunth nov. gen. am. 4. p. 230), fruticulosa glabra, ramis tetragonis, foliis oppositis petiolatis tripartitis, lobis filiformibus integerrimis, capitulis solitariis pedicellatis, invol. squamis exter. paulò brevioribus lanceolato-linearibus ciliatis, ligulis lanceolato-oblongis subtridentatis, achæniis margine lanato-pilosis biaristatis, aristis corollas disci æquantibus. \mathcal{L} in Andibus Peruviz. Flores disci flavi. Ligulæ aurantiacæ.
27. *C. TRIFIDA* (Lam. ill. t. 704. f. 2. Poir. suppl. 2. p. 353), glabra, caule subflexuoso, foliis trifidis subulatis in axillâ fasciculatis, pedunculis terminalibus 1-cephalis, ligulis angustis obtusè et grossè 3-dentatis. — patr. et cæt. ign. Folia ex icone opposita, ex descr. alterna. †
- † *Species non satis notæ.*
28. *C. PALMATA* (Nutt. gen. am. 2. p. 180), caule simplici compresso sub-1-cephalo, foliis oppositis sessilibus subcoriaceis palmato-3-lobis margine scabris, lobis lineari-oblongis obtusis, invol. utroque 8-partito, achæniis oblongo-ellipticis nudis. — in plantis terris. Michiganensis, Illinoensis et in Louisiana infer. *Calliopsis palmata* Spreng. syst. 3. p. 611. Flores lutei. An *Peta-mihi* sp. ex Raf.

29. *C. NUDATA* (Nutt. gen. am. 2. p. 180), caule humili simplicifolice apice dichotomo, foliis alternis glabris subulato-linearibus remotissimis, super. minimis, radiis obsoletis 3-dentatis, achæniis nudis. — in Florida occid. circa urbem Sanctæ Mariæ. *Calliopsis nudata* Spreng. syst. 3. p. 611. Ligulæ rubræ. Invol. ext. minutum. An *Peramibi* sp. ex Raf.? †.
30. *C. ROSEA* (Nutt. gen. am. 2. p. 179), parva glaberrima, caule sæpius simplicifolice apice dichotomo oligocephalo, foliis oppositis linearibus integris, capitulis longè pedunculatis terminalibus, ligulis inæqualiter 3-dentatis, achæniis integerrimis nudis. ½ in paludosis à Novâ-Cæsareâ ad Georgiam. *Calliopsis rosea* Spreng. syst. 3. p. 611. Ligulæ rosæ. An *Peramibi* sp. ex Raf.? †.
31. *C. PAUCIFLORA* (hort. Hamb. ex linnæâ 1835. litt. p. 76), caule herbaceo simplici, foliis oppositis sessilibus tripartitis glabris, lobis lanceolatis obtusiusculis margine scabris, medio longiore, capitulis terminalibus subsolitariis. ½ in Americâ boreali.
32. *C. BITERNATA* (Lour. fl. coch. 622), caule herbaceo erecto tetragono, foliis biternatis ovato-lanceolatis serratis, paniculâ diffusâ, ligulis 6 neutris, pappo bicorni ramoso, recept. nudo. — prope Cantonem Sinarum. *Actinea biternata* Spreng. syst. 3. p. 474. Caulis integrè luteus dicitur. Genus omninò ambiguum ex recept. nudo dicto et fortè ad *Actineam* rectè relatum. †.

†† *Species exclusæ.*

- C. ACUTA* Pursh = *Peramibus acutus*.
C. ALATA Cav. = *Actinomeris tetraptera*.
C. ALATA Pursh = *Actinomeris alata*.
C. ALBA Linn. = *Bidens alba*.
C. ALTERNIFOLIA Linn. = *Actinomeris alternifolia*.
C. AMPLEXICAULIS Cav. = *Simsia amplexicaulis*.
C. ANGUSTIFOLIA Pav. = *Bidens procera*.
C. ARTEMISIFOLIA Jacq. = *Cosmos sulphureus*.
C. ATKINSONIANA Dougl. = *Calliopsis Atkinsoniana*.
C. BACCATA Linn. = *Wulffia stenoglossa*.
C. BIDENS Walt. = *Bidens chrysanthemoides*.
C. CHRYSANTHA Spreng. = *Bidens Coreopsidis*.
C. CORONATA Linn. = *Bidens leucantha*.
C. CORONATA Spreng. = *Bidens oxyodonta*.
C. DIVERSIFOLIA Jacq. = *Bidens grandiflora*.
C. FERULIFOLIA Jacq. = *Bidens Coreopsidis*.
C. FLAMMULA Banks. = *Bidens chrysanthemoides*.
C. FORTIDA Cav. = *Simsia sicifolia*.
C. FOENICULACEA fl. mex. = *Bidens foeniculifolia*.
C. FORMOSA Bosc. = *Cosmos bipinnatus*.
C. FRUTICOSA Vest. = *Bidens fruticosa*.
C. FRUTICOSA Forst. = *Campylothea micrantha* β.
C. GEORGINA NUDA Cass. = *Dahlia variabilis*.
C. GEORGINA PRUINOSA Cass. = *Dahlia coccinea*.
C. HETEROPHYLLA Cav. = *Echinacea heterophylla*.
C. HIRTA Raf. = *Peramibus hirtus*.
C. INCISA Ker. = *Bidens Coreopsidis*.
C. LEUCANTHA hort. = *Bidens leucantha*.
C. LEUCORHIZA Lour. = *Bidens leucorhiza*.
C. LIMENSIS Jacq. = *Encelia Limensis*.

- C. ODORATA Lam. = *Bidens odorata*.
 C. OVATA Cav. = *Actinomeris ovata*.
 C. PARVIFLORA Jacq. = *Cosmos parviflorus*.
 C. PERFOLIATA Bosc. = *Bidens chrysanthemoides*.
 C. PROCERA Ait. = *Actinomeris alternifolia*.
 C. PURPUREA fl. mex. = *Bidens purpurea*.
 C. REPTANS Linn. = *Bidens Coreopsisidis*.
 C. SCABRA Raf. = *Peramibus scaber*.
 C. SERRULATA Poir. = *Bidens grandiflora*.
 C. TETRAGONA Lallav. = *Bidens tetragona*.
 C. TINCTORIA Nutt. = *Calliopsis tinctoria*.
 C. TRIPTERIS Linn. = *Chrysostemma tripteris*.

CCCXXXVIII. *ACTINOMERIS* Nutt. gen. am. 2. (1818) p. 131. Less. syn. 228. — Ridan Adans. fam. 2. (1763) p. 130. — Pterophyton Cass. bull. philom. 1818. p. 76. dict. 44. (1826) p. 48. 59. p. 139. — Actimeris Raf. 1820. — Coreopsisidis sp. Linn.

Capitulum multiflorum radiatum, ligulis neutris paucis 1-serialibus elongatis, fl. disci herm. tubulosis 5-dentatis. Inv. 1-3-seriale, squamis subinæqualibus foliaceis acuminatis. Recept. convexum paleaceum, paleis achæniarum marginem amplectentibus. Styli rami appendiculati acuti. Achænia bilateraliter plano-compressa angustè alata, aristis binis triquetris sublævibus coronata. — Herbæ Boreali-Americanae scabrae biennes perennesve. Caules erecti ramosi teretes foliis longè decurrentibus alati. Folia ovata aut lanceolata serrata alterna rariùs subopposita. Capitula corymbosa. Corollæ flavæ (aut var. fortè ligulæ albæ). — Differt à Coreopside achæniis lateraliter compressis nec obcompressis, à Verbesinâ ligulis neutris.

1. *A. ALTERNIFOLIA*, caule supernè pubescente, foliis alternis lato-lanceolatis serratis scabris, paniculâ corymbosâ laxâ foliosâ, invol. patente uniseriali, recept. subglobose. ♀ in Virginiâ (Pluk. Pursh), Carolinâ (Fras.!), Georgiâ (Ell.), in territorio Arkansano è semin. à cl. Pourtales lectis. Coreopsis alternifolia Linn. sp. 1283. Jacq. hort. vind. t. 110. Ridan Adans. fam. 2. p. 130. Pterophyton alternifolium Cass. dict. 44. p. 49. Actinomeris squarrosa Nutt. gen. am. 2. p. 131. Ell. sketch 2. p. 413. Verbesina Coreopsis Michx. fl. bor. am. 2. p. 134. — Pluk. alm. t. 159. f. 3. Ligulæ 7-8 flavæ. Variat etiâ ligulis nullis ex Michx. Folia infima opposita aut subverticillata, undè verisim. Coreopsis procera Ait. kew. 3. p. 258 ab hac non differt. (v. v. c. et s. sp.)

♂? *alba* (Ell. sketch 2. p. 413), ligulis albis, foliis angustius lanceolatis magis scabris, corollis disci glaberrimis. ♀ in Carolinâ inferiore. An species propria?

2. *A. HELIANTHOIDES* (Nutt. gen. am. 2. p. 181), caule subscabrido, foliis alternis lanceolatis acutis serratis scabris subtus villis, paniculâ corymbosâ subsimplici oligocephalâ coarctatâ, invol. biseriali, recept. convexo. ♀ in Georgiâ circa Louisville (Ell. sketch 2. p. 412), in Ohio, Kentucky, Tennessee et Louisiana infer. (Nutt.). Verbesina helianthoides Michx. fl. bor. am. 2. p. 135 non Kunth. Ligulæ 10-12 aureæ elongatæ.
3. *A. TETRAPTERA*, caule subscabro, foliis oppositis et superioribus alternis ovato seu rhombeo-lanceolatis scaberrimis basi cuneatis à medio serratis, pedunculis ad capitulum usquè alatis sub corymbo, ligulis 14-15. ♀ in Mexico (Cav.). Coreopsis alata Cav. ic. 3. p. 3. t. 260 non Pursh. Helianthus tetrapterus Ort. dec. 6. p. 74. Moç. fl. mex. ic. ined. Pterophyton alatum Cass. dict. 44. p. 49. — Hern. mex. p. 189. ic. Ligulæ flavæ. (v. v. c.)

4. *A. TETRAGONA*, foliis etiã superioribus oppositis ovatis serratis basi angustatis, pedunculis alatis 1-cephalis. \mathcal{L} in Mexico circa urbem (Alam. ! Berl. ! pl. exs. n. 647), ad Temisci prœdium saccharinum ubi *Contrayerva* audit (fl. mex. mss.). *Helianthus tetragonus* Moq. fl. mex. ic. et descr. ined. Mamatzlacotl Hern. mex. p. 383. f. 2. Affinis *A. tetrapteræ* et fortè ferè varietas sed ex Hernandez et Mocino videtur diversa. Antiseptica et febrifuga ulcera putrida citò et mirificè sanat ex Moq. (v. s.)
5. *A. OVATA* (Nutt. gen. am. 2. p. 181 in adnot.), caule glabresculo, foliis alternis ovato-oblongis crenulatis, capitulis corymbosis, pedunculis apteris, invol. squamis linearibus obtusis, ligulis 8. \mathcal{L} in Mexico. *Coreopsis ovata* Cav. ic. 3. p. 41. t. 280. Ligulæ flavæ.
6. *A. PEDUNCULOSA*, foliis oppositis ovato-rhombeis dentatis triplinerviis, pedunculis terminalibus longissimis 1-cephalis omninò exalatis. \mathcal{L} in Mexico. Folia utrinquè scabro-aspera, longè secus caulem decurrentia, apice ferè obtusa. Pedunculi 7-8 poll. longi. (v. s. comm. à cl. Alaman.)
7. *A. ALATA* (Nutt. gen. am. 2. p. 181), foliis oppositis indivisis ovatis acuminatis serratis petiolatis triplinerviis. (2) in Virginiâ (ex Bartr. in h. Banks). *Coreopsis alata* Pursh fl. bor. am. 2. p. 567 non Cav.
8. *A. PAUCIFLORA* (Nutt. in Sillim. journ. 5. (1822) p. 301), foliis oppositis oblongis serrulatis, pedunculis nudis oligocephalis. — in Floridâ orientali. Caulis videtur simplex. Invol. simplex squamis lanceolatis. Achænia breviter bicornia. Flor. ign.

CCCCXXIX. *ARMANIA* Berter. herb. — *Hopkirkia* sp. Spreng.

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris 1-serialibus, disci hermaphroditis cor. tubo brevi, fauce latiore cylindricâ, limbo 5-dentato. Invol. 3-seriale, squamis imbricatis adpressis. Recept. planum paleaceum. Styli fl. hermaphrod. rami breves truncati apice barbati exappendiculati. Achænium obcompressum ellipticum cartilagineum adpressè pubescens, setis 2 rigidis anticè (nec retrorsum scabris) coronatum. — Frutex Austr.-Americanus, ramis teretibus foliisque adpressè villosis. Folia petiolata ovata acuminata subcrenata penninervia. Capitula pauca (2 in specim.) pedicellata terminalia. Flores flavi. — Genus à *Salmeâ* (quæ *Hopkirkia* Spr.) diversissimum capitulis radiatis, stigmate truncato apice barbato, recept. plano etc.; à *Bidente* differt involucrio, stigmate, formâ achæniis et habitu toto.

1. *A. FRUTICULOSA* (Bertr. ! herb.). \mathfrak{f} ad Sanctam-Martham Amer. merid. olim legit amic. Bertero. *Hopkirkia fruticulosa* Spreng. syst. 3. p. 444. (v. s.)

CCCCXL. *OYEDÆA*.

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris 1-seriatis, disci tubulosi 5-dent. hermaphr. Invol. squamæ 3-seriales subæquales foliaceæ. Recept. planum, paleis complicatis mucronatis acuminatis coriaceo-subscariosis. Cor. disci tubus gracilis, faux cylindrica, dentes 5 erecti extus puberuli. Anth. nigricantes inclusæ. Styli rami graciles appendice hispidâ superati. Achænia radii abortiva linearia pappo 2-3-aristato aristellisque minimis superata; disci compressa alata subcuneata, apice hinc sæpè gibba, pappo vario, aristis nempe 2 caducis elongatis et pluribus aliis brevibus intermediis constante. — Frutices Austr.-Americani Verbesinæ facie. Rami teretes adpressè villosi-subhirti. Folia opposita in petiolum attenuata ovata acuminata dentata supra sparsè villosa subtus villosa-subcanescentia aut tomentosa, inter. grosse triplinervia, summa subpenninervia. Capitula pedunculata co-

rymbosa. Flores lutei. — Genus à Verbesinâ ligulis neutris, ab Heliantho et Viguiérâ achæniis alatis diversum, ab Actinomeride squamellis inter duas aristas pappi intermediis. Dicavi cl. Hispano navigatori Alphonso Oyeda, qui, primus post Colombium, Americam adiit et imò in nave suâ vehebat Americum Vespuccium nimio honore postea decoratum cum Oyeda ferè oblitus. (Conf. bibl. voy. 6. p. 212.)

1. O. VERBESINOIDES, foliis subtus villosis, invol. squamis lineari-lanceolatis acuminatis pubescentibus. $\frac{3}{4}$ circa Caracas legit cl. Vargas. Folia inf. pedem longa, 7-8 poll. lata. Ligulæ pollicem longæ, apice 2-3-dentatæ. Achænia glabra sparsè maculata. (v. s.)
2. O. BUPHTHALMOIDES, foliis subtus secus nervos villosos strigosis inter nervos tomentosis, invol. squamis ovatis mucronatis, ext. dense villosis, inter. submembranaceis longè ciliatis. $\frac{3}{4}$ in Peruvia legit cl. Poeppig. Bupthalmum trifolium Poepp. pl. exs. n. 22 et 1624. Caules ramique dense hirsuti. Folia opposita, nec in meo specim. terna, 9 lin. longa. Achænia radii 3- disci 2-aristata, aristellis minimis paucis hinc inde interjectis. (v. s. comm. à cl. Poepp.)

CCCXLI. SIMSIA Pers. ench. 2 (1807). p. 478. Cass. dict. 5^e p. 136 non Brown. (1810). — Ximenesiæ sp. Kunth Less. — Simsia et Ximenesiæ sp. Loud. h. brit.

Capitulum multiflorum heterogamum, fl. radii neutris ligulatis 1-serialibus paucis (5-12), disci hermaphr. tubulosis 5-dentatis. Invol. squamæ 2-3-seriales erectæ lineares subæquales. Recept. planum, paleis membranaceis complicatis. Styli rami appendiculati. Achænia plano-compressa glabriuscula exalata apice biaristata. — Herba Mexicanæ, Viguière facie, pleræque hispida. Folia inf. opposita, sup. alternâ trinervia petiolata aut sessilia, sæpè basi auriculato-amplexicaulia. Capitula pedicellata laxè corymbosa. Flores omnes lutei. — Differt à Viguiérâ squamellis inter aristas nullis, ab Heliantho invol. non imbricato, à Ximenesiâ ligulis neutris et achæniis exalatis.

1. S. LAGASCEFORMIS, caule parçè subhispido, foliis inf. oppositis, sup. alternis petiolo exaurito longè pappo donatis ovatis acuminatis serratis utrinquè mollior subvillosis, corymbo laxo subpaniculato, invol. squamis lanceolatis, ext. paucis dorso hispidis dimidiò brevioribus, inter. glabris subcoloratis disco æqualibus, ligulis circ. 5 breviter bidentatis, achæniis glabris. — in Mexico. Species ab omnibus aliis faciliè differt invol. serie externâ dimidiò internâ brevior. (v. s. comm. à cl. Mairet.)
2. S. FICIFOLIA (Pers. ench. 2. p. 478), ramis hispidis et pube glandulosâ pulverulento-velutinis, foliis utrinquè puberulis petiolatis exauriculatis cordatis profundè trilobis dentatis, infer. oppositis, sup. alternis, capitulis paucis subpaniculatis, invol. squamis ext. hispidis, int. margine membranaceis, ligulis 8 vix denticulatis, achæniis glabris apice vix ciliatis. (1) in Mexico. Coreopsis foetida Cav. ic. 1. p. 55. t. 77. Ximenesia foetida Spreng. syst. 3. p. 606. Herba graveolens. Folia exactè Fici aut Mori. (v. s.)
3. S. HETEROPHYLLA (DC. non Cass.), ramulis hispidis, foliis inf. oppositis, sup. alternis, omnibus ovali-oblongis membranaceis utrinquè pilosiusculis denticulatis, aliis petiolatis petiolo nudo exauriculato, aliis sessilibus obtusè auriculatis, nunc variè trilobis nunc indivisis, invol. hispidi squamis lineari-subulatis, ligulis 7-8 tridentatis, achæniis glabris margine ciliatis. (2) in Mexico prope Zimapequaro (H. et B.) inter Tampico et Real del Monte (Berl.) pl. exs. n. 442) Ximenesia heterophylla H. B. et Kunth nov. gen. am. 4. p. 227. t. 380. Simsia Kunthiana Cass. dict. 5^e p. 137, sed Simsia heterophylla Pers. è genere excludenda et ad Echinaceam referenda. (v. s.)
4. S. AURICULATA, ramis hispidis, foliis utrinquè villosis latè ovatis irregulariter

trilobatis acuminatis dentatis basi breviter cuneatis, petiolo hispido immarginato basi latè auriculato, corymbo composito laxo polycephalo, invol. squamis lineari-lanceolatis acuminatis, ext. hispidis, int. margine membranaceis, ligulis 6-8 minutè 2-3-dentatis, achæniis glabriusculis. — in Mexico circa urbem legit cl. Berlandier (pl. exs. n. 850 et 927). Encelia Mexicana et Ximenesia hirta Mart. f. herb. A. S. heterophylla differt foliis multò latoribus, petiolisque basi amplè auriculatis. Variat foliis oppositis alternis, limbo basi truncato aut ovato. (v. s.)

5. *S. HISPIDA* (Cass. dict. 59. p. 137), caule apice hirtò-hispidulo, foliis alternis sessilibus ovato-oblongis obsolete serratis suprà piloso-subtùs sericeo-hispidis, invol. squamis hispidis lanceolatis, ligulis circ. 12, achæniis.... ♀ in montosis Mexici prope S. Rosa de la Sierra et Valladolid. Ximenesia hispida H. B. et Kunth nov. gen. am. 4. p. 227. *S. heterophylla* similis dicitur.
6. *S. CORDATA* (Cass. dict. 59. p. 137), ramis pilosis, foliis inf. oppositis, sup. alternis deltoideo-ovatis acuminatis cordatis basi cuneatis grossè crenatis utrinquè adpressè hispidulis, petiolis basi amplè dilatatis, invol. squamis lanceolato-linearibus piloso-hispidis, ligulis 6-8 emarginato-bidentatis, achæniis sericeo-pubescentibus. ♀ in Mexico prope Capula et Valladolid de Mechoacan. Ximenesia cordata H. B. et Kunth nov. gen. am. 4. p. 228. Ximenesiæ encelioidi proxima dicitur! An fortè idèò Ximenesiæ spec.?, sed ligulæ dicuntur neutræ et achæni alà marginalis ex descr. videtur nulla.
7. *S. DOMBEYANA*, caule tereti sparse hispido et inter setas minutè puberulo-glanduloso, foliis superioribus alternis petiolatis latè ovatis irregulariter repando-dentatis hinc indè sublobatis acutis utrinquè setis hispidulis et puberulo-glandulosis, capitulis paucis breviter pedicellatis, invol. squamis lineari-lanceolatis acuminatis disco longioribus, ligulis paucis minimis, achæniis obcordatis biaristatis margine ciliolatis. — in Amer. austr. verisim. in Peruvia legit Dombey. Petioli valdè hispidi, setis ut in totà plantà longis patulis mollioribus. (v. s. comm. à Mus. reg. Par.)
8. *S. AMPLEXICAULIS* (Pers. ench. 2. p. 478), canescens, foliis alternis subpalmato-trilobis, petiolo basi auriculato-semiamplexicauli. — in Mexico. Coreopsis amplexicaulis Cav. cors. descr. p. 226. Lag. et Rod. in annal. scienc. nat. 1802. v. 5. p. 75. Ait. kew. ed. 2. p. 134. Folia sæpius sub-5-loba, infima aspera obliqua indivisa. Flores crocei. (Pers.)

† *Species exclusa.*

S. HETEROPHYLLA Pers. = *Echinacea heterophylla*.

CCCXLII. *VIGUIERA* H. B. et Kunth nov. gen. am. 4. p. 224. — *Viguiera* Less. syn. 229.

Capitulum multiflorum heterogamum, fl. radiis neutris ligulatis, fl. disci hermaphr. 5-dentatis. Invol. semi-globosum, squamis 1-2-seriatis subæqualibus apice in appendicem foliaceam productis. Recept. paleis persistentibus achænia amplectentibus obsitum. Styli rami exserti subulati hispidi. Achænia compressa obovato-cuneata pubescentia squamellis 4 ovatis obtusis subdenticulatis aristisque 2 ex angulis ortis coronata. — Herbæ Americanæ ramosæ. Caules glabri, rarius villosi. Folia opposita aut alternà petiolata triplinervia ovato-lanceolata supernè scabra. Capitula pedicellata laxè paniculato-corymbosa. Cor. luteæ. — Genus *Heliantho* et *Leighiæ* affine, sed caractere Kunthiano parum dilatato, apprimè naturale.

§. 1. *Receptaculo acutè aut obtusè conico.*

1. *V. HELIANTHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 226. t. 379), caule glabro, foliis alternis ovatis acuminatis integerrimis trinerviis suprà scabris subtùs adpressè pilosis, petiolis villosò-ciliatis, invol. squamis uni-2-seriatis

linearibus hispidulis apice dilatato-subspathulatis, recept. longo conico cavo. ① in insulâ Cubâ propè Havanam frequens.

2. *V. SAGRÆANA*, caule glabro, foliis oppositis et alternis ovato-lanceolatis subintegerrimis triplinerviis suprà subscabrisculis subtùs petiolisque parcè pubescentibus, invol. squamis biserialibus linearibus in appendicem foliaceam acutam productis, recept. conico demùm cavo. ①? in insulâ Cubâ juxtâ Havanam legit cl. Ramon de La Sagra. Valdè affinis *V. heliantheidi*, sed folia frequentius opposita subintegra, petioli non villosi-ciliati, invol. diversa, etc. (v. s. comm. cum seq. à cl. de La Sagra).
3. *V. MICROCLINE*, caule glabro, foliis oppositis ovato-lanceolatis integerrimis trinerviis suprà pube minutissimâ subscabris subtùs petiolisque villosi-canescenscentibus, invol. squamis biserialibus in appendicem foliaceam acutam apice productis, recept. conico pleno. ①? in insulâ Cubâ prope Havanam legit cl. Ramon de La Sagra. Habitus *V. helianthoidis*, sed juxta hujus descr. et iconem facillè diversa. Ligulæ 5-7. Achænia adpressè villosa. (v. s.)
4. *V. CANESCENS*, caule villosi, foliis alternis ovalibus acuminatis in petiolum longè attenuatis integerrimis suprà pilosulo-scabris subtùs petiolisque villosi-canescenscentibus, invol. squamis biserialis dorso canescenscentibus in appendicem linearem acutiusculam productis, recept. obtusè conico. ① in Mexico prov. Leoninâ ad occid. urbis Guanaxuato legit cl. Mendez. Habitus ferè *Ximenesiæ*. Cor. croceo-flavæ. (v. s.)
5. *V. TRIQUETRA*, caule glabro, foliis serratis suprà scabris subtùs pubescentibus, infer. oppositis, summis subalternis, petiolis glabrisculis, invol. squamis linearibus acutis biserialis, recept. obtusè conico demùm cavo. ①? in valle Mexicanâ prope S. Angelo legit cl. Alaman et etiam Berlandier (pl. exs. n. 827 et 937). Panícula trichotoma, pedunculis angulatis subtriquetris. *Helianthus triquet* Ort. dec. p. 76. (v. s.)
6. *V. DENTATA* (Spreng. syst. 3. p. 615), caule glabro, foliis serratis oppositis, petiolis subciliatis, paniculâ oligocephalâ, invol. squamis 2-seriatis acutis, recept. convexo. — in Mexico. *Helianthus dentatus* Cav. ic. 3. p. 10. t. 220. Folia basi latiora apice magis acuminata quàm priorum. Pedicelli graciles vix angulati. (v. s.)

§. 2. Receptaculo planiusculo.

7. *V. PROSTRATA*, caule diffuso debili pubescente, foliis oppositis breviter petiolatis ellipticis utrinquè acutis apice subserratis triplinerviis utrinquè pubescenti-scabridis, ramis elongatis floridis apice nudiusculis 1-cephalis, invol. squamis lineari-oblongis biserialis canescenti-villosis, paleis indivisis, achæniis glabris angulatis. 2/ in Amer. bor. ex Willd., sed ab hodiernis in confed. Anglo-Americ. non reperta! fortè Mexicana? *Helianthus prostratus* Willd. sp. 3. p. 2242 non Poir. Pappi aristæ tenuissimæ, squamellæ minimæ. Caulis ex Link non verè prostratus. (v. s.)
8. *V. SESSILIFOLIA*, caule suffruticoso adscendente laxè ramoso sparsè hispido demùm glabrato, foliis inf. oppositis, rameis alternis, sessilibus ovatis acutis subintegerrimis utrinquè villosis, paniculâ laxâ oligocephalâ subnudâ, invol. squamis 2-3-seriatis oblongo-linearibus villosi-hispidis, ligulis 5-6, paleis lanceolatis acuminatis, achænio adpressè sericeo. 3 in Mexico circa Villalpando legit cl. Mendez et ad Tlapujahua cl. F. W. Keerl (h. Mart.). Paleæ recept. glabræ, apice virescentes, cor. disci æquales. Pappus squamellis 2 aristatis, 4 brevibus, omnibus ciliato-pubescentibus. Rec. planiusculum. Ligulæ pallidè flavæ nervis nigricantibus lineatæ. (v. s.)
9. *V. BREVIPES*, caule herbaceo puberulo laxè ramoso, foliis breviter petiolatis ovato-lanceolatis acuminatis subintegerrimis triplinerviis utrinquè puberis-scabris, petiolis villosi-ciliatis, paniculâ laxâ, invol. squamis 2-3-seriatis lanceolatis acutis dorso subtrinerviis, paleis lanceolatis acutis membranaceis, achæniis adpressè subpubescentibus. — in Mexico circa Bejar legit cl. Berlandier (pl. exs. n. 1905). Habitus omninò *V. helianthoidis*. (v. s.)

10. *V. GRAMMATOGLOSSA*, ramis vix subhirtellis, foliis oppositis petiolatis ovato-lanceolatis acuminatis vix subdentatis utrinque petioloque longè et adpressè villosis, pedicellis ad apices ramorum 1-2 nudis 1-cephalis, invol. squamis oblongo-linearibus acutis, ligulis latis circ. 15-nerviis, achænio villoso 1-2-aristato, aristis squamellis plurimis subvillosis. — in Mexico ad Tlapachua legit cl. F. W. Koeberl ex h. Mart. Affinis *V. brevipedis*, sed folia achæniæque multò magis villosa; *V. canescenti*, sed recept. parè convexum; à duabus et à plerisque differt ligulis latis multistriatis. (v. s. in h. Mart.)
11. *V. LAXA*, caule suffruticuloso glabriusculo, foliis alternis longè petiolatis latè ovatis basi cuneatis apice acuminatis remotè serratis triplinerviis suprà scabridis subtus pubescentibus, paniculà laxà, pedicellis pubescentibus elongatis, invol. squamis biserialibus linearibus acutis, ext. apice recurvis, ligulis 5 oblongis, recept. paleis integris acutis. ♂? in Mexico inter San-Fernando et Santander legit cl. Berlandier (pl. exs. n. 2258!). Valdè accedit ad *Helianthum quinqueradiatum* sed (si Cavan. descr. exacta) satis differt. Petioli inf. 2 et ultra pollices longi ciliati. Recept. vix subconvexum. (v. s.)
12. *V. OPPOSITIFOLIA*, caule ascendente lævi, foliis alternis longè petiolatis elliptico-ovatis basi acutis apice acuminatis subdentatis margine scabris suprà sparse puberis subtus jun. pubescenti-villosis, pedunculis folio oppositis 1-cephalis, invol. biseriali, squamis ext. patulis foliaceis linearibus disco longioribus ligulas ferè æquantibus, ligulis 5-6, recept. plano, paleis lanceolatis acutis, achænio adpressè pubescente. — in Mexico. Ligulæ ex sicco aurantiacæ. Pappus 6-squamellatus, squam. 2 aristæformibus, 4 intermediis obtusis. (v. s.)
13. *V. P. FOLIACEA* (Spreng. syst. 3. p. 616), simpliciuscula hirta, foliis oppositis petiolatis acutis basi attenuatis integerrimis, pedunculis solitariis, invol. foliaceo radium longè superante. — in Brasiliâ. Cæter. iga. An potius *Leighia* spec. ? †.

CCCXLIII. *LEIGHIA* Cass. dict. 25. p. 435. — *Helianthi* sect. Less. — *Helianthi* sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris 1-serialibus. Invol. imbricatum, squamis oblongis in appendicem foliaceam patulo-squarrosam productis. Recept. convexum, paleis lanceolatis achænia amplectentibus. Cor. et genit. *Helianthi*. Achænia compressa hispidula. Pappus squamellis 2 aristæformibus ex angulis achæniis ortis et 4 intermediis ovalibus brevibus denticulatis constans. — *Herbæ* aut suffrutices Americanæ. Folia alterna aut opposita. Capitula *Helianthi* sed minora. Genus ex pappo affine *Viguiæræ* et *Harpalii*; à priore invol. squarroso, à posteriore squamis appendiculatis diversum. Fortè *Viguiera*, *Leighia* et *Harpalium* in unicum genus congreganda, sed certè inter se potius quàm *Heliantho* affinia.

* *Foliis plerisque alternis.*

1. *L. LAXA*, caule suffruticuloso ascendente basi glabro apice sparse setoso. foliis alternis oblongo-lanceolatis acutis basi attenuatis hinc inde subdentatis penninerviis suprà sparse strigillosis subtus secus nervos hispidulis, pedicellis terminalibus alaribusque folio triplò longioribus puberulis 1-cephalis, invol. squamis 2-3-seriatis ovalibus acutis disco longioribus ciliatis dorso puberis, paleis membranaceis acuminatis, achæniis villosis biaristatis et plerisque squamellatis. 2L ♂ in Brasiliâ prov. Rio-Grande. (v. s. in h. Mus. reg. Par. ex. h. Mus. imp. Bras. sub n. 880.)
2. *L. ANCHUSÆFOLIA*, caule herbaceo tereti pilis sparsis scabro apice paniculato. foliis alternis sessilibus elongato-lanceolatis acutis triplinerviis integerrimis utrinque et margine strigilloso-hirtellis, capitulis ramos pedunculiformes terminalibus subsolitariis, invol. squamis lineari-lanceolatis acuminatis subpatulis scabris, ligulis 10-12, paleis mucronulatis, achæniis adpressè subvillosis

2L in Brasiliæ prov. Rio-Grande. Herba et pubes ferè *Anchusæ angustifoliæ*. Folia 2½ poll. longa, 4-5 lin. lata. Ligulæ flavæ 9-10 lin. longæ obtusæ. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 875.)

3. *L. NONREPOLIA*, caule suffruticoso diffuso, ramis teretibus præsertim ad apices hispidis, foliis alternis sessilibus oblongo-lanceolatis acutis serratis penninerviis utrinquè sparsè piloso-hispidis, pedicellis terminalibus et alaribus folio triplò longioribus 1-cephalis, invol. squamis 2-3-serialibus oblongis acutis foliaceis laxiusculis ciliatis et dorso puberis, ligulis circ. 12-14 invol. duplò longioribus, paleis membranaceis acutis, pappi aristis 2-3, squamellis acutis subirregularibus. 5 in Brasiliæ prov. Rio-Grande. Folia alia grossè, alia minutè serrata, 18-20 lin. longa, 5-6 lata. Flores flavi. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 877.)
4. *L. MICROPHYLLA* (Cass. dict. 25. p. 436), fruticosa ramosissima, ramis jun. cano-tomentosis, adultis glabris, foliis alternis adproximatis subsessilibus oblongis obtusis margine revolutis integerrimis subtripplinerviis coriaceo-crasiusculis, suprà strigoso-scabris, subtùs incano-tomentosis, capitulis pedicellatis solitariis cernuis, invol. squamis lineari-oblongis laxè imbricatis, ligulis circ. 16, paleis lanceolatis membranaceis, achæniis adpressè pilosis. 5 in subfrigidis Peruvix juxta Caxamarca. *Helianthus microphyllus* H. B. et Kunth nov. gen. am. 4. p. 220. t. 375. Pappus ex descr. biaristatus, ex icon. biaristatus et pluripaleaceus.
5. *L. DISSITIFOLIA*, caule erecto tereti paniculato ramoso glabro vix ad apices puberulo, foliis alternis oppositisque remotis sessilibus ovali-lanceolatis subintegerrimis triplinerviis, suprà sparsè, subtùs nervos minutè scabris, nervia basi margine contiguis, pedicellis terminalibus paucis, invol. squamis lanceolatis pubescenti-canescensibus 3-seriatis, ligulis 7-9 invol. duplò longioribus, paleis membranaceis subobtusis, pappo 2-aristato et 4-squamellato. 2L ? in Brasiliæ prov. Rio-Grande. Capitulum flavum, ligulis 5-6 lin. longis. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. n. 881.)
6. *L. LOMATONEURA*, caule erecto tereti adpressè et minutè puberulo paniculato-ramoso, foliis alternis, ramis hinc indè oppositis subsessilibus lineari-oblongis vix acutis integerrimis utrinquè adpressè puberulis et scabridis triplinerviis, nervis lateralibus margine subcontiguis, pedicellis paucis 1-cephalis subterminalibus, invol. squamis 3-serialibus foliaceis pubescentibus patulis lanceolato-linearibus, paleis vix acutis, achæniis minutissimè pubescentibus compressis, pappo aristis 1-2 parvulis et squamellis linearibus acutis plurimis constante. 2L 5 in Brasiliæ prov. Rio-Grande. Habitus ferè *L. linearis*. Flores non vidi. Pappus ferè ad *Helenias* accedit. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 878.)
7. *L. IMMARGINATA*, caule herbaceo erecto tereti adpressè et minutè puberulo, foliis alternis (rarissimè in ramulis oppositis) sessilibus elongato-lanceolato-linearibus integerrimis 1-nerviis utrinquè minutè et asperè strigillosis demùm albo-punctatis, capitulis terminalibus pedicellatis, invol. squamis lanceolato-linearibus pubescenti-canescensibus..... 2L in Brasiliæ prov. Rio-Grande. Similima *L. lomatoneuræ*, sed nervi marginales non sunt perspicui. Folia 3-poll. longa, 3 lin. lata. Fruct. non vidi. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1052.)
8. *L. LINEARIS*, caule erecto ramoso scabrido, foliis sessilibus alternis rariùs oppositis linearibus margine revolutis integerrimis uninerviis suprà et subtùs in nervis hispidò-scabris, invol. squamis oblongis apice foliaceo-appendiculatis squarroso-recurvis scabris multiserialibus, lignis circ. 20 invol. paulò longioribus. 2L 5 in Mexico juxta Hapam (Cav.), ad Guanajuato (H. et B. Alam!), Queretaro (Née!), Tlapujhua (Keer!), S. Angelo (Berl.) pl. exs. n. 924). *Helianthus linearis* Cav. ic. 3. p. 9. t. 218. Ker bot. reg. t. 523. H. squarrosus H. B. et Kunth nov. gen. am. 4. p. 222. t. 377. *Leighia elegans* Cass. dict. 25. p. 435. Achæn. hispidulum. Pappus persistens 6-8-paleolatus, 2 paleis elong. aristatis, 3-6 brevibus membranaceis denticulatis. (v. v. c. et c. sp.)

** *Folii oppositis.*

9. *L. ANGUSTIFOLIA*, caule suffruticuloso ramosissimo tereti basi glabro, ramis puberulo-hirtellis, foliis oppositis sessilibus linearibus acutis integerrimis utrinque piloso-hirtis, pedicellis terminalibus 1-cephalis, invol. squamis 2-3-seriatis, ext. foliaceis dorso hirtellis disco longioribus, intim. membranaceis, paleis apice calloso-mucronatis, ligulis invol. duplō longioribus, achæniis villosis 2-aristatis et-4 squamellatis. \mathcal{U} \mathcal{Z} in Brasiliæ prov. Rio-Grande (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 884.)

β . *breviaristata*, foliis squamisque inv. exterioribus magis acuminatis paulo rigidius strigoso-hirtis, paleis in acumen subrotundum productis, achæniis villosis, pappo breviter biaristato et minutissimè subsquamellato. \mathcal{U} \mathcal{Z} in prov. Rio-Grande. An species propria? (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. n. 1037.)

10. *L. ARILLATA*, caule basi suffruticuloso adscendenti-erecto ramosissimo puberulo scabrido tereti, foliis oppositis sessilibus lineari-lanceolatis hinc inde grossè dentatis aut minutè serratis penninerviis, suprā sparśe, subtūs secus nervos pubero-hirtellis, pedunculis terminalibus 1-cephalis, invol. squamis 3-seriatis, ext. lanceolatis dorso hirtellis, mediis latoribus submembranaceis, achæniis villosis radii abortivis 3-aristatis, disci obovato-oblongis 2-aristatis et pauci-squamellatis basi utrinque membranā adpressā subrotundā ex areoli ortā quasi arillatis! \mathcal{U} \mathcal{Z} in Brasiliæ prov. Rio-Grande. Species *L. calendulacæ* valde affinis. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. n. 883.)

11. *L. CALENDULACEA*, caule suffruticuloso? gracili adscendente subsimplici puberulo, foliis oppositis sessilibus lineari-lanceolatis acutis utrinque villosis subscabridis integerrimis, pedunculis terminalibus solitariis 1-cephalis, invol. squamis oblongo-lanceolatis dorso puberis apice squarrosis ciliatis, ligulis 9-10, paleis lanceolatis, achæniis sericeo-villosis. \mathcal{Z} \mathcal{U} in Brasiliæ littoralis parte dictā Banda orientale legit cl. Bacle. Cor. ex sicco videntur aurantiacæ. Pappi aristæ breves. (v. s.)

12. *L. BUPHTHALMIFLORA*, caulibus suffruticulosis adscendentibus setis crassis inæqualibus præsertim ad apices hirtis, foliis oppositis sessilibus elongato-lanceolatis acuminatis serratis triplinerviis suprā sparśe strigillosis secus ramulos hirtellis, ramis apice pedunculiformibus nudis 1-cephalis, invol. squamis 3-serialibus, ext. lanceolatis scabris laxis disco longioribus inæqualibus foliaceis, paleis membranaceis acuminatis, achæniis villosis biaristatis et pluri-squamellatis. \mathcal{U} \mathcal{Z} in Brasiliæ prov. Rio-Grande. Capit. magnitudine colore et habitu simile cap. *Bupthalmi salicifolii* (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 879.)

13. *L. URTICIFORMIS*, caule suffruticuloso tereti hispidissimo apice scabro, foliis oppositis sessilibus lanceolatis subserratis basi obtusis trinerviis apice acuminatis suprā scabridis subtūs villosis, capitulis terminalibus solitariis, invol. squamis appendice patulā squarrosis, exter. acuminatis, inter. obtusis, achæniis biaristatis, squamellis intermediis minimis acutis. \mathcal{Z} in Mexico circa Villalpando propè urbem Guanajuato legit cl. Mendez. Capit. flava, diam. sesquipollic. Folia 2 poll. longa, 7 lin. lata. (v. s.)

14. *L. SPÆROCEPHALA*, caule herbaceo oppositè ramoso striato pubescenti-villoso, foliis oppositis petiolatis ovato-lanceolatis acuminatis triplinerviis subdentatis, suprā scabris, subtūs secus nervos villosis, cæterum pubescentibus, capitulis ternato-corymbosis, invol. squamis villosis squarrosis acutis, paleis glabris complicatis mucrone rigido recurvo, achæniis villosis ciliatis. \mathcal{U} ? in Mexico legit cl. Alaman. Pedicelli striati. Capitula diam. pollic. ferè globosa. Cor. disci ex invol. reflexo quasi exserta. (v. s.)

† *Species subdubiæ.*

15. *L.? LEPTOCEPHALA*, fruticosa, ramis petiolisque pube brevi confertā velutinis, foliis oppositis petiolatis ovatis acuminatis basi breviter cuneatis crenatis

- utrinque sparæ et minutè puberulis, paniculâ parvâ corymbosâ, invol. oblongi squamis lineari-lanceolatis apice patenti-squarrosis parçè hirtis, lignis 5-6 apice dentatis, squamellis setiformibus plurimis inter aristas setis achæniûm coronantibus. ♀ in Mexico? Habui ex herb. Thib. sub nomine evid. falso Coreopsidis foetidæ. Habitus à cæteris Leighiis differt, sed ex caract. ad nullum aptius genus adjungere quivi. (v. s.)
16. L. ? *ECLIPSEFOLIA*, caule herbaceo erecto glabro, ramulis erectis parvis versus apices adpressè puberulis, foliis oppositis sessilibus elongato-lanceolatis linearibusve adpressè et minutissimè puberulis, aliis integerrimis, aliis utrinque unidentatis et triplinerviis, capitulis terminalibus solitariis pedicellatis, invol. squamis lineari-lanceolatis apice squarrosis foliaceis, achæniis villosis nunc breviter 1-aristatis nunc exaristatis et squamellis coroniformibus constante. (1) ? in Brasiliæ provinciâ Rio-Grande. Folia ferè *Ecliptæ* erectæ, 4 poll. longa, 4 lin. lata. Flores flavi. Pappus ex ovario judicatus à genere subdiversus. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 88a.)
17. L. ? *OVATIFOLIA*, ramis teretibus glabrisculis, foliis suboppositis alternisve sessilibus distantibus ovatis acutiusculis integerrimis suprâ sparè subatrigillois subtus velutino-hirsutis, pedicellis terminalibus et axillaribus apice puberulis, invol. squamis pluriarerialibus lanceolatis acutis ciliatis adpressiusculis, ligulis circ. 10 ellipticis invol. paulò longioribus, achæniis radii glabris abortivis, disci villosis 1-2-aristatis et squamellis plurimas parvas gerentibus. — in Brasiliâ prov. Sancti-Pauli. Ex invol. coriaceo adpresso paulò magis ad *Harpalium* fortè accedit. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 410.)
18. L. ? *SPECIOSA*, caule erecto tereti, foliis petiolatis cordatis indivisis trilobisque crenatis, pedunculo 1-cephalo supernè incrassato, invol. squamis apice foliosis acutis squarrosis, paleis acuminatissimis cuspidatis longitudine florum disci, cor. disci basi inflatis ad faucem cylindricis, achæniis obtusè 4-gonis biaristatis et brevissimè 4-squamellatis. 2/ in Mexico ad Jorullo. *Helianthus speciosus* Hook. bot. mag. t. 3295. Herba habita Tithoniæ, caractere Leighiæ. Ligulæ circ. 12-13 intensè crocæ.

† *Species exclusa.*

L. *BICOLOR* Cass. = *Helianthus angustifolius*.

CCCXLIV. *HARPALIUM* Cass. dict. 20. p. 299. 25. p. 437. — *Helianthi* sect. Less. — *Helianthi* sp. *anc.*

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris 1-serialibus. Invol. squamæ imbricatæ ovatæ inappendiculatæ obtusæ. Recept. convexum, paleis oblongis achæniâ amplectentibus. Cor. et genit. *Helianthi*. Achæniûm compressum parçè villosum. Pappus disci squamellis 2 aristæformibus ex angulis achæniî ortis et 4 brevibus obtusis intermediis. — *Herbæ Americanae* alterni- aut oppositifoliæ, habitu *Viguieræ* et *Helianthi*. Genus à priore invol. pluriseriali imbricato inappendiculato, à posteriore pappi squamellis interinediis diversum.

SECT. I. EUHARPALIUM.

Ligularum pappus nullus.

1. H. *ARGINUM* (Cass. dict. 20. p. 300), caule erecto rigidè piloso, foliis infer. oppositis triplinerviis basi attenuatis, super. remotis subsessilibus lanceolatis vix dentatis rigidis pube brevi confertâ utrinque cinereis, ramis apice nudis 1-cephalis, invol. hemisphærici squamis ovalibus obtusis imbricatis, rec. convexo, paleis obtusis, achæniis hispidis. 2/ in Americâ boreali. *Helianthus rigidus* Desf. cat. h. Par. ed. 3. p. 184. Folia subpallida seu cinerascientia. Capit. æquale cap. *Helianthi* multiflori. Cor. disci fusca. Styli ligulæque lutei. Pappi squamellæ 2 longæ aristæformes, 4 breves oblongæ. (v. v. in hort. Girod. Genev.)
2. H. *TRUXILLENSE* (Cass. dict. 25. p. 438), caule herbaceo erecto pilosiusculo, foliis

alternis sessilibus lanceolato-oblongis acutis basi rotundatis integerrimis triplinerviis rigidis scaberrimis, capitulis solitariis pedicellatis, invol. squamis adpressis lanceolatis extus hispidis, ligulis 12-15, paleis lanceolatis scariosis, achænia glabro biaristato squamellis pluribus interjectis, ½ in maritimis Peruvia juxta Truxillo. *Helianthus Truxillensis* H. B. et Kunth nov. gen. am. 4. p. 223.

3. *H. AUAUUM* (Cass. dict. 25. p. 438), caule herbaceo, ramis hispido-villosis, foliis alternis? petiolatis, oblongis utrinque acutis subintegerrimis subtripplinerviis rigidis supra pilosis subtus incanis et villosulo-lanatis, capitulis solitariis longe pedunculatis, invol. squamis lanceolato-oblongis acutis adpressis hispido-pilosis, ligulis 16-20, paleis linearicarinatis obtusis, achæniis adpresso-pubescentibus biaristatis et pluriusquamellatis, ½ in apricis prope Alansi Quitensium. *Ligulæ aurantiacæ. Helianthus aureus* H. B. et Kunth nov. gen. am. 4. p. 224.

Sect. II. HARPALIZIA.

Achænia disci sericeo-villosissima, pappo aristis 2 et squamellis latis ovatis fimbriatis aristas ferè æquantibus constante; radii abortiva gracilia glaberrima pappo amplo coronæformi dentato superata.

4. *H. ? SERICEUM*, frutescens, ramis jun. sericeo-canescens, foliis alternis petiolatis oblongo-lanceolatis basi acutis apice acuminatis integerrimis subtripplinerviis adpressis sericeo-pubescentibus, pedicellis solitariis folio æqualibus 1-cephalis, invol. squamis 3-serialibus adpressis, ext. ovatis obtusis, int. obovato-mucronatis, omnibus dorso canescenti-villosis, ligulis 9-10 multistriatis. ½ in Americâ calid. legit. cl. Née, sed locus propr. ign. Habitus ferè *Helianthi argentei*, sed achænia diversissima. Car. fract. H. aurei, sed fruticosum. (v. s.)

CCCXLV. *TITHONIA* Desf. in Juss. gen. p. 189. et in ann. Mus. h. n. Par. 1. p. 46. t. 4. Cass. dict. 35 (1825), p. 277. 54. p. 454 et 59, p. 147. B. in mag. Gart. Weim. 1828. p. 156. Less. syn. 229. non Ræusch. (1)

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris, disci hermaphr. 5-fidis. Invol. 2-3-seriale, squamis ovatis basi intus striatis subcallosis apice foliaceo appendiculatis obtusis. Recept. convexum, paleis lanceolatis membranaceis achænia plùs minùs amplectentibus. Cor. disci tubo brevissimo, fauce inflatâ multinerviâ. Styli rami subulati hispidi exserti revoluti. Achænia radii obcompressa seu triquetra, pappo brevissimo parco, disci subtetragona compressa lævia, pappo 1-seriali squamellis pluribus denticulatis, et in omnibus aut saltè in fl. centralibus setis 1-2 ex angulis fructus majoribus ortis coronata. — Herbæ annuæ ex Americâ æquin. Folia alterna serrata triplinervia. Rami apice pedunculiformes 1-cephali subnudi. Cor. crocæ seu flavæ. — Genus *Heliantho* propinquum.

* *Pedunculis apice obconicis incrassatis.*

1. *T. TAGETIFLORA* (Desf. l. ann. Mus. 1. p. 46. t. 4. cat. h. Par. ed. 3. p. 404), foliis plerisque trilobis, pedunculis apice incrassatis, invol. biseriali breviter velutino erecto, ligulis ovali-oblongis. (1) ad Vera-Cruz Mexic. legit olim. cl. Thierry de Menonville. Specimen habeo à cl. Bertero in Sancto-Domingo lectum, sed fortè cultum. Lam. ill. t. 708. mal. Poir. ill. 3. p. 284 et 519. mal. Ic. fl. Mex. ined. Cass. dict. 54. p. 456. Don in bot. reg. t. 591? Ligulæ crocæ-aurantiæ. Achænia pleraque squamellis tantum et centralia tantum setis insuper coronata. Folia triloba ferè *Fici* folia referunt. (v. v. c.)
2. *T. TURBIFORMIS* (Cass. dict. 35. p. 277 et 54. p. 454), foliis omnibus indivisis, pedunculis apice incrassatis, invol. hirsuto subpatente, ligulis oblongis. (1) in Mexico (Ort.), ad Tlapujahu (Keerl in h. Mart.). *Helianthus tubæformis*

(1) Genus sine descr. videtur *Rudbeckiæ* sp.

Ort. dec. p. 101. Jacq. h. Schoenbr. 3. p. 65. t. 375. Lindl. bot. mag. t. 1519. Tith. helianthoides B. in mag. gart. Weim. 1828. p. 156. ex bull. Fer. sc. nat. 16. p. 432. Ligulæ flavæ. Achænia pleræque biaristata. (v. v. c.)

** *Pedunculis cylindricis.*

3. *T. PACHYCEPHALA*, caule juniore hinc indè hirsutissimo demùm scabro, foliis ovato-lanceolatis serratis utrinquè scabris trinerviis reticulatis, pedunculis teretibus, invol. squamis plariserialibus, exter. ciliatis squarrosis, achæniis exterioribus uni-, interioribus biaristatis. $\frac{1}{2}$? in Mexici prov. Leoninà ad occid. urbis Guanaxuato legit cl. Mendez. Capitulum amplum crassum. Recept. hemisphæricum. Cor. omnes croceo-flavæ. Pili caulis è basi tuberculosâ fasciculati decidui. (v. s. sp.)
4. *T. EXCELSA*, caule scabro et hinc indè bifariam villosa, foliis ellipticis acutis apice serratis triplinerviis utrinquè scabris, pedunculis teretibus, invol. squamis plariserialibus ciliatis adpressis, achæniis plerisque biaristatis. $\frac{1}{2}$ in Mexico (Cav.), ad S. Angelum (Berl. pl. exs. n. 940). Helianthus giganteus Cav. ic. 3. p. 10. t. 119 non Lin. nec Lour. Helianthus excelsus Willd. sp. 3. p. 2243. Helianthus trinervis hort. Madr. et Monsp. 1813. Herba orgyalis. Capitula flava. (v. v. c. et sp.)

CCCXLVI. *HELIANTHUS* Linn. gen. n. 979. Cass. dict. 20. p. 351. 59. p. 140. — Chrys. Ren. 84. t. 83. — Corona-Solis Tourn. inst. p. 489. t. 279. — Vossian *Adans. fam.* 2. p. 130. — Helianthi sect. 1^a Less. — Helianthi sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris 1-ser., disci tubulosis hermaphroditis. Invol. irregulariter imbricatum, squamis ext. foliaceis acutis appendiculatis inappressis, intimis minoribus paleiformibus. Recept. planum aut convexum, paleis semiamplexentibus oblongis acutis. Cor. disci tubus brevis, faux amplior cylindracea, dentes 5. Styli rami appendiculati. Achænia lateraliter compressa aut subtetragona, parcè villosa aut glabra. Pappus squamellis 2 ex angulis achæniis ortis aristæformibus constans, et rarissimè 2 alias conformes ex angulis lateralibus ortas exhibens. — Herba (rariùs suffrutices) Boreali-Americanae pleræque scabræ aut villosæ. Folia opposita, super. interdum alterna, integra aut dentata, sæpiùs triplinervia. Capitula ampla solitaria rarò corymbosa. Cor. disci luteæ aut fuscae, radii luteæ aut subaurantiacæ.

* *Annui.*

1. *H. ANNuus* (Linn. sp. 1276), radice fibrosâ, caule subamplici erecto, foliis alternis petiolatis cordatis seu latè ovatis trinerviis grossè serratis, capitula maximè cernuis, invol. squamis latè ovatis abruptè acuminatis ciliatis, paleis subindivisis, achæniis disci biaristatis. ① in Peru (Lob. Dod. Cam.), Mexico? (Linn.), nunc in hortis totius ferè orbis frequens et hinc indè per campos seminibus vulgatus. Mill. illustr. ic. capituli. *H. platycephalus* Cass. dict. 20. p. 352. — Lobel. ic. t. 592. f. 2. Pappi paleolæ duæ subconcaevæ lanceolatæ denticulatæ, Herba 5-6-pedalis. (v. v. c.)

ℓ. *Indicus*, involucri squamis saltem exterioribus in folia subpetiolata expansis. — *H. indicus* Linn. mant. 117. Moustrum potiùs quàm varietas. (v. s.)

γ. *pumilus*, pumilus multò hirsutior. — *H. pumilus* Pers. ench. 2. p. 475. (v. v. c.)

2. *H. MACROCARPUS*, caule erecto vix ramoso strigoso, foliis alternis petiolatis ovatis acuminatis dentatis subscabris grossè triplinerviis, summis juxta capitulum subconferis etiàm petiolatis, invol. squamis scabris ciliatis acuminatissimis, ligulis basi pubentibus oblongo-lanceolatis, achæniis corollas disci per anthesin duplò superantibus. ① patr. ign. Vagat in hortis sub nom. *hybrida* Hel.

- longicarpi. Ligulæ citrino-flavæ, apice acutè dentatæ, 2 $\frac{1}{2}$ poll. longæ, 8-9 lin. latæ. Cor. disci basi villosæ, flavæ. Pappi et recept. paleæ pallidæ. (v. v. c.)
3. *H. OVATUS* (Lehm. cat. h. Hamb. 1828. p. 16 et in Linneæ 1830. p. 376. pl. nov. n. 7), caule erecto subsimplici atrigoso, foliis longè petiolatis ovatis acuminatis remotè subdenticulatis triplinerviis strigoso-asperis, inf. oppositis, super. alternis, pedunculis terminalibus incrassatis, invol. squamis oblongo-lanceolatis setoso-ciliatis, ligulis numerosis integris. (1) in Mexicanis terris. Vagat in hortis sub nomine evidentissimè falso *Hel. petiolaris*. Capitula 3-5 poll. diam. Corollæ disci basi aurantiacæ, lobis intus subfuscis, tubo ad partem dilatatam annulo piloso cincto. (v. v. c.)
4. *H. LENTICULARIS* (Dougl. in bot. reg. t. 1265), caule erecto hispido, foliis alternis petiolatis ovatis attenuatis apice acuminatis triplinerviis serratis, capitulis amplis subcernuis, pedunculo tereti, invol. squamis expansis ovalibus acuminatis dorso scabris, paleis acutè trifidis, achæniis biaristatis. (X) in Amer. bor. temperatis ad oram occident. et reg. interioribus legit Douglas, in regione Arkansanæ ex sem. Pourtal. Hook. fl. bor. am. 1. p. 313. Valde accedit ad *H. annuum*, sed capitula minora. Corollæ disci ad apicem purpureo-fuscæ, basi villosæ. Pappi et receptaculi paleæ apice fusco-rubentes. Achænia obovata ferè per anthesin longitudine corollæ. Ligulæ intensè flavæ 2 poll. longæ, ferè pollicem latæ. (v. v. c.)
5. *H. PATENS* (Lehm. ind. sem. h. Hamb. 1821. p. 8. ind. schol. 1828. p. 19), caule erecto asperissimo patentim ramoso, foliis alternis longè petiolatis ovato-subcordatis serratis undulatis triplinerviis scabris subitis canescentibus, pedunculis longissimis sub capitulo incrassatis, invol. squamis linearis-lanceolatis, infer. reflexis, achæniis pilosis. (1) in Americâ calid. sed loc. propr. ign. Discus fusco-purpureus. Ligulæ saturatè flavæ.
6. *H. PETIOLARIS* (Nutt. in Journ. ac. nat. sc. Philad. 1821. oct. p. 115), caule erecto ramoso-subpubescente foliis alternis longissimè petiolatis ovatis acuminatis trinerviis integrisculis scabris, invol. squamis lanceolatis acuminatis dorso breviter scabris, ligulis circ. 20 integris. (1) in arenosis ad ripas fl. Arkansas Amer. bor. detexit cl. Nuttall. Sweet! brit. fl. gard. ser. 2. v. 1. t. 75. Desf. cat. h. Par. ed. 3. p. 403. Capitula longè pedunculata. Ligulæ flavæ, magn. circ. *H. multiflori*. Discus fuscus. Achænia villosa. Pappus biaristatus. (v. v. c. et s. sp.)
- ** *Perennes, disco atro-purpureo. — Discomela Raf. neogen. 1825. p. 3.*
7. *H. ATRO-RUBENS* (Linn. sp. 1279), caule erecto pilis longis sparsis hispido, foliis oppositis oblongo-spathulatis seu ovatis crenatis triplinerviis scabris, invol. squamis ovatis acutis vix discum æquantibus glabrisculis, ligulis circ. 16, achæniis jun. villosis biaristatis. (1) in Amer. bor. à Pensylvaniâ ad Carolinam in glareosis et siccis necnon ad ripas fl. Saskatchewan (Hook.). Ker bot. reg. t. 508. *H. Missuriensis* Spr. pug. 21. n. 44. *H. diffusus* Sims bot. mag. t. 2020. — Dill. elth. 111. t. 94. f. 110. Discus atro-rubens. Capitula longè pedicellata solitaria ex icon. recentiorum, aut brevius pediculata ternato-corymbosa ex ic. Dill. Recept. dicitur convexum. Eadem fortè species ex Desf. ac *H. suus rigidus*, sed pappus ex Ell. biaristatus nec squamellatus.
8. *H. LATIFLORUS* (Pers. ench. 2. p. 476. n. 26), caule erecto aspero, foliis inf. oppositis, sup. alternis ovali-lanceolatis basi cuneato-attenuatis apice longè acuminatis serratis utrinquè asperis, pedunculis elongatis 1-cephalis nudis, invol. squamis lanceolatis acuminatis ciliatis disco ferè longioribus, ligulis 15-16, achæniis glabris biaristatis. (1) in Americâ bor. (hortul.), in Georgia occid. ? (Ell.). *H. atro-rubens* Lam. dict. 3. p. 86 non L. *H. sparsifolius* Ell. sketch 2. p. 415. Discus subatro-rubens. Paleæ integræ apice callosæ fusco-virentes. (v. v. c.)
9. *H. ORCYALIS*, caule elato lævi, foliis alternis sessilibus linearibus planis subdenticulatis vix subscabridis 1-nerviis, capitalis 5-7-corymbosis longè pedunculatis, invol. squamis linearibus acuminatis ciliolatis, recept. paleis lineari-

subcuneatis integris apice subciliatis, achænio glabro 2-3-4-aristato. \mathcal{H} (Linn.) in Amer. bor., Virginiâ (Linn.), agro Arkansasano (Pourtales?). *Helianthus angustifolius* Linn. sp. 1279 non Michx. *Coleopsis* Mill. ic. t. 224. f. 2. Sæpè cum *H. angustifolius* Michx. confusum et stricto jure angustifolium dicendum, sed nomen speciei Michx. servavi nè augeatur confusio. Herba 6-10-pedalis. Discus atro-purpureus. (v. v. c. in h. Genev. ex sem. Pourtalesianis.)

10. *H. ANGUSTIFOLIUS* (Michx.! fl. bor. am. 2. p. 141), caule erecto parè villosus, foliis sessilibus linearibus margine revolutis subsabridis 1-nerviis, inf. oppositis summis alternis, capitulis paucis longè pedicellatis, invol. squamis lanceolatis disco æqualibus, recept. paleis cuneatis apice serratis trifidis, achænio glabro biaristato. \mathcal{H} in Amer. bor. à Virginiâ ad Floridam (Michx.!), Carolinæ. humidis pinetis (Ell. Bosc!), Novâ-Cæsareâ (Torr.! Green.!). Ell. sketch 2. p. 415. Sims bot. mag. t. 2051. *Rudbeckia angustifolia* Linn. sp. 1281. excl. syn. Mill. *Leighia bicolor* Cass. dict. 25. p. 436. Discus atro-rufus. Rec. subconvexum. Ligulæ 12-15. (v. s. sp.)

11. *H. CILIARIS*, caulibus pluribus stricatis glabris, foliis oppositis lineari-lanceolatis basi attenuatis apice acuminatis undulatis integerrimis glaucis 1-nerviis utrinquè glabris margine præsertim ad basin rigidè ciliatis, capitulis solitariis terminalibus, invol. squamis imbricatis adpressis latè ovatis subacutis ciliatis, ligulis circ. 8-10, achæniis glabris biaristatis. \mathcal{H} in Mexico prope Reynosa de Tamaulipas apr. flor. legit cl. Berlandier. Rhizoma lignescens. Caules pedales. Ligulæ 3 lin. longæ, flavæ, striis circ. 12 fuscis notatæ. Discus subfuscus involucri longior. (v. s.)

*** *Perennes, disco luteo, foliis omnibus oppositis.*

12. *H. MOLLIS* (Lam.! dict. 3. (1789) p. 81 non Willd.), caule erecto piloso, foliis oppositis sessilibus semiamplexicaulibus cordato-ovatis acuminatis adpressè serratis suprâ pubero-asperis subtùs molliter villosis, invol. squamis lanceolato-linearibus hirsuto-canescens, recept. demùm convexo, paleis acuminatis integris, ligulis circ. 20, achæniis glabris biaristatis. \mathcal{H} in Americæ bor. pratis humidis, in Alabamâ, Georgiâ (Ell.). Buchoz ic. ex hort. Paris. *H. pubescens* Willd. sp. 3. p. 2240. Ker bot. reg. t. 524 (excl. syn. Vahl. ex descr. et specim. h. Hafn.). Tausch in flor. 1828. p. 503. *H. canescens* Michx. fl. bor. am. 2. p. 140. (v. v. c.)

13. *H. DORONICOIDES* (Lam. dict. 2. (1789) p. 84), caule erecto basi lævi cæterùm pubescenti-scabrido, foliis sessilibus oppositis et sup. alternis ovato-lanceolatis acutis adpressè serratis suprâ subsabridis subtùs pubescentibus, invol. squamis lineari-lanceolatis puberulis margine longè ciliatis, recept. paleis trifidis lobo medio longiore acuminato, ligulis circ. 12, achæniis glabris biaristatis. \mathcal{H} in Amer. bor. siccis subfertilibus à Pensylvaniâ ad Carolinam. *H. pubescens* Vahl symb. 2. (1791) p. 92 (ex spec. hort. Hafn. et conf. descr.). Hook. bot. mag. t. 2778 non Willd. *H. mollis* Willd. sp. 3. p. 2240 (non Lam. et excl. syn. Michx.). Ell. sketch 2. p. 418. *H. Hookeri* Don in Lond. h. brit. p. 358. *H. pilosus* Tausch in florâ 1828. p. 502? (v. v. c.)

14. *H. MACROPHYLLUS* (Willd. hort. berol. t. 70), caule erecto piloso-scabro, foliis oppositis breviter petiolatis ovato-aut oblongo-lanceolatis serratis utrinquè scabris triplinerviis, petiolo ciliato, capitulis paucis subcorymbosis, invol. squamis lineari-lanceolatis basi longè ciliatis et dorso glabriusculis apice scabridis patulo-squarrosis, paleis 3-dentatis, achæniis glabris biaristatis. \mathcal{H} in Amer. bor. An *H. strumosus* Linn. sp. 1277. Ell. sketch 2. p. 420? — Boeck. sic. t. 27. f. 4. pessim. Zan. hist. t. 56. med. *H. serotinus* Tausch in flor. 1828. p. 504? (v. s.)

15. *H. DIVARICATUS* (Linn. sp. 1279 excl. syn. Mor.), caule erecto glabro lævi, foliis oppositis subsessilibus oblongis aut ovali-lanceolatis acuminatis serratis suprâ scabris subtùs pubescentibus imâ basi ciliatis, corymbò trichotomo, invol. squamis lanceolatis brevissimè ciliatis patulo-squarrosis, paleis acutè 3-dentatis, achæniis glabris biaristatis. \mathcal{H} in Amer. bor. à Canadâ (Hook.) et

- Noveboracœ ad Carolinam. Lam. dict. 3. p. 86. Tausch in florâ 1828. p. 504. *H. hispidulus* Ell. sketch 2. p. 419? *H. leiostemon* Raf. in litt. — *Moris. ox. s. 6. t. 3. f. 62* (nec t. 7. f. 66 quâ citatione falsâ decepti plures auct. Amer. sub hoc nom. alteram spec. foliis longè petiolatis habuerunt; tales sunt Elliott, Beck, etc.). *Radix repens.* (v. v. c. et a. sp.)
16. *H. DECAPETALUS* (Linn. sp. 1277), caule erecto lævi apice scabro, foliis oppositis ovatis petiolatis acuminatis grossè serratis triplinerviis utrinquè scabris, invol. squamis lineari-lanceolatis squarrosis hispidulis ciliatis, ligulis 10, rec. paleis 3-dentatis. 2/ in Canadâ (Kalm ex L.), in siccis lapideosis sylvaticis à Canadâ ad Virginiam (Pursh). Affinis *H. multifloro*, sed caulis glaber, folia angustiora, ligulæ pauciores angustiores breviores. Rob. ic. t. 235 ex Pursh. (v. v. c.)
17. *H. BUDDLEIÆFORMIS*, caule suffruticuloso tereti pubescenti-scabrido, foliis oppositis breviter petiolatis oblongo-lanceolatis subserratis trinerviis acutis suprâ scaberrimis subtùs secus nervos hirtis, capitulis pedicellatis corymbosis plerisque ternis, invol. squamis appendiculatis, appendice patulo-reflexâ squarrosâ, ext. acuminatis, cæteris obtusissimis, achæniis biaristatis. 3/ in Mexico circa Villalpando legit cl. Mendez et Alaman. *Verbesina mollis*? Alam. in litt. Accedit ad *Leighiam* urticiformem. (v. s.)
18. *H. TRUNCATUS* (Schwein. in Ell. sketch 2. p. 416), caule gracili glabro, foliis oppositis breviter sessilibus ovatis supernè attenuatis serratis pilosis scabris, capitulis paucis, invol. squamis lanceolatis, paleis indivisis apice fimbriatis. 2/ in Carolinâ boreali (Schw.) et Georgiâ occid. (Ell.)
19. *H. LONGIFOLIUS* (Pursh fl. bor. am. 2. p. 571), glaberrimus, caule paniculato, ramis summitate oligocephalis, foliis oppositis subsessilibus lanceolatis triplinerviis integerrimis, infer. serratis, invol. squamis acutis, ext. linearibus divaricatis, ligulis circ. 10, paleis lanceolatis 3-dentatis. 2/ in humidis fertilibus Georgiæ occid. Ell. sketch 2. p. 417. Habitus *Coreopaidum aquaticarum*. Pappus subulatus caducus.
20. *H. TENUIFOLIUS* (Ell. sketch 2. p. 420), caule lævigato, foliis oppositis longè petiolatis ovato-lanceolatis acuminatis grossè serratis utrinquè scabrinusculis membranaceis, capitulis parvis paucis, invol. squamis lanceolatis ciliatis adpressis, ligulis circ. 10?, paleis 3-dentatis fimbriatis. 2/ in Georgiæ district. occidentilibus. Affinis *H. spathulato*. Pappus subulatus pubescens.
21. *H. SPATHULATUS* (Ell. sketch 2. p. 421), caule supergè scabro, foliis oppositis in petiolum attenuatis spathulato-ovatis paulò acuminatis serratis suprâ scabridis subtùs pubescentibus, ramulis oppositis, invol. squamis lanceolatis hispidis apice subulatis, ligulis 10-12?, paleis acuminatis integris. 2/ in Georgiâ occidentali. Pappus subulatus pubescens. Ligulæ pollicem longæ.
22. *H. TRICUSPIS* (Ell. sketch 2. p. 422), scaberrimus, caule oppositè ramoso, foliis oppositis oblongis ovato-lanceolatis subintegris margine subrevolutis, capitulis paucis, invol. squamis lato-subulatis ciliatis, ligulis 14-16, paleis profundè tricuspidatis. 2/ in Georgiâ occidentali. Pappus subulatus lacernus.
23. *H. DIVERSIFOLIUS* (Ell. sketch 2. p. 423), caule scabro oppositè ramoso, foliis oppositis breviter petiolatis suprâ scaberrimis subtùs pubescentibus, infer. ovato-lanceolatis acuminatis, superioribus cordato-ovatis mucronatis, invol. squamis lanceolatis ciliatis, ligulis 10-12, paleis subtridentatis apice villosis. 2/ in Georgiâ occidentali. Affinis *H. tricuspidi*. Pappus subulatus pubescens achænio longior.
24. *H. SCABERRIMUS* (Ell. sketch 2. p. 423), caule scabro parcè ramoso, foliis oppositis basi attenuatis lanceolatis utrinquè scaberrimis subintegerrimis, invol. squamis ovatis adpressis, ligulis 16-20, paleis integris dorso ciliatis. 2/ in Georgiâ occid. Pappus subulatus pubescens.
25. *H. PAUCIFLORUS* (Nutt. gen. am. 2. p. 177), caule nudo trichotomo oligocephalo, foliis oppositis lineari-lanceolatis acuminatis serratis ferè lævibus, invol. acutè imbricatis ovatis. 2/ in Louisianâ inferiore. Folia interdum ternato-verticillata, subtùs pallidiora.

**** *Perennes, disco luteo, foliis superioribus alternis.*

26. *H. PROCUMBENS* (Pers. ench. 2. p. 475), caule diffuso-procumbente vix apice pubescente, foliis alternis subsessilibus ovalibus utrinque obtusis basi triplinerviis apice subserratis, capitulis solitariis pedicellatis, inv. squamis biseriatis pubescentibus subciliatis lineari-oblongis, paleis indivisis, achæniis tenuissimè 1-2-aristatis. \mathcal{U} in Peruvia ex hort. Par. *H. prostratus* Desf. cat. ed. 2 nec ed. 3. Ligulæ 6 lin. longæ. — *Sanvitalia helianthoides* Willd. sp. 3. p. 2190 huc à Persoonio citata videtur diversa caule erecto, capitulis duplò majoribus, etc. *H. procumbens* Raf. fl. Lud. p. 70, omninò diversus à genere removendus sed vix notus. (v. v. c.)
27. *H. MICRANTHUS* (Spreng. syst. 3. p. 618), herbaceus, ramis retrorsum scabro-hirtellis, foliis inf. oppositis sessilibus lineari-oblongis obtusis subintegerrimis hispidò-scaberrimis rigidis, capitulis solitariis longissimè pedicellatis, invol. squamis 10 lanceolatis adpressis hispidis, paleis acutis, achæniis glabris biaristatis. \mathcal{U} prope Santa-Rosa Mexicanorum. *H. parviflorus* H. B. et Kunth nov. gen. am. 4. p. 222. t. 378 non Bernh.
28. *H. AMPLEXICAULIS*, caule herbaceo ramoso subscabrido, foliis semiamplexicaulibus dentatis, inf. oppositis basi obtusè auritis longè supra basin triplinerviis supra aurículas angustatis dein ovato-lanceolatis ferè panduræformibus, mediis oppositis indivisis, summis alternis, paniculâ laxâ, pedicellis birtellis, invol. squamis 2-3-seriatis lineari-lanceolatis acutis basi hirtò-ciliatis adpressis, recept. paleis acuminatis integris, achæniis aristatis. — in Mexici prov. Leoninâ ad occid. urbis Guanaxuato legit cl. Mendez. *Habitus* Vignieræ sed recept. vix convexum et pappus purè biaristatus nec squamellas ullas exhibens. (v. s.)
29. *H. TRACHELIOLIVUS* (Willd. sp. 3. p. 2241), caule scabro supernè ramoso, foliis ovato-lanceolatis acuminatis serratis triplinerviis utrinquè scaberrimis, inf. oppositis, sup. alternis, paniculâ laxâ, invol. squamis lineari-lanceolatis ciliatis, ligulis 10-12, paleis apice 3-dentatis villosis. \mathcal{U} in Amer. bor. sylvia et palud. exsiccatis in Novâ-Cæsarâ (Torr.!), à Canadâ ad Carolinam (Pursh), in montibus (Ell.). Hook. fl. bor. am. 1. p. 312. *H. gigas* Michx. fl. bor. am. 2. p. 141. (v. s. et v. c.)
30. *H. CORNIFOLIUS* (H. B. et Kunth nov. gen. am. 4. p. 223), herbaceus, ramis angulatis hispidis, foliis oppositis alternisque petiolatis oblongo-ellipticis acuminatis basi rotundatis remotè denticulatis triplinerviis rigidis suprâ asperissimis subtus hispidis, invol. squarrosis hispidis, paleis acuminato-pungentibus, achæniis adpressè pubescentibus biaristatis. \mathcal{U} in asperis Mexici prope Santa-Rosa de la Sierra.
31. *H. TOMENTOSUS* (Michx. fl. 2. p. 141? Ell. sketch 2. p. 424), caule aspero parcè ramoso, foliis ovato-lanceolatis supernè attenuatis acutis serrulatis suprâ scabris subtus tomentosis plerumquè alternis, capitulis grossè pedunculatis, invol. squamis foliaceis squarrosis lanceolatis ciliatis, ligulis 12-14, paleis trifidis. \mathcal{U} in Georgiâ occid. et inter flum. Chatahouchia et Alabama (Ell.), in pratis Illinoensium? (Michx.)
32. *H. CALIFORNICUS*, caule elato lævi apice laxè paniculato, foliis inf...., sup. alternis remotis elongato-lanceolatis integerrimis basi attenuatis subciliatis apice acuminatis utrinquè scabris triplinerviis, pedunculis scabris, invol. squamis lineari-sublanceolatis disco paulò longioribus squarroso-patulis puberulo-subhirtellis, achæniis glabris biaristatis. \mathcal{U} ? in Californiâ legit cl. Douglas. Affinis *H. giganteo* sed caulis lævis, folia angustiora integerrima, invol. squamæ vix ciliatæ, etc. (v. s. comm. ab hon. Soc. hort. Lond.)
33. *H. GIGANTUS* (Linn. sp. 1278 non Cav.), caule elato scabro, foliis sparsis lanceolatis serratis obsoletè triplinerviis scaberrimis basi attenuatis et ciliatis, corymbo stricto, invol. squamis lanceolatis linearibus disco longioribus squarroso-patulis basi villosò-ciliatis apice scabris, achæniis glabris, squamellis 2 acuminatis subfimbriatis. \mathcal{U} in Amer. bor. à Canadâ ad Carolinam. Mill. dict. n. 5. Tausch in florâ 1828. p. 501. *H. virgatus* Lam. dict. 3. p. 85. *H. al-*

- simus Jacq. h. vind. 2. t. 18a non Linn. H. giganteus et altissimus Willd. sp. 3. p. 2242. — Moris. ox. 3. s. 6. t. 7. f. 66 et 67. Herba elata in hortis frequens variat caule plus minus scabro, purpurascente aut viridi; foliis angustioribus latoribus plus minus serratis; corymbo oligo- aut polycephalo; invol. squamis in appendicem foliaceam sæpe longissimè productis, semper disco longioribus; ligulis 12-15, etc. Inter has variationes limites non satis certas video ut varietates sint tutò distinguendæ. (v. v. c. et s. sp.)
34. H. ALTISSIMUS (Linn. sp. 1278. excl. syn. Mor.), caule erecto glabriusculo, foliis alternis oblongo-lanceolatis in petiolum ciliatum attenuatis acuminatis obtusè serratis triplinerviis superne scabris. invol. squamis lanceolatis disco æqualibus dorso glabris margine longè ciliatis, achæniis glabris, squamellis 2 acuminatis fimbriatis. \mathcal{L} in Amer. bor. sed locus propr. incertus ob confusions hujus sp. cum H. giganteo. H. altissimus Tausch in florâ 1828. p. 502. H. giganteus Hill. veg. syst. 2. t. 38. f. 6^p Ligulæ 10-12. Paleæ apicè villosociliatæ acutæ integræ. Pedicelli adpressè pilosi scabriusculi. Discus luteus. (v. v. c.)
- β . foliis exsertè et argutè serratis \mathcal{L} . (v. v. c.)
35. H. MULTIFLORUS (Linn. sp. 1277), rhizomate tereti inflexo, caule erecto ramoso scabrido, foliis petiolatis dentatis triplinerviis scabris, inf. cordatis, super. ovatis, invol. squamis ext. lineari-lanceolatis ciliatis, inter. lanceolatis, ligulis oblongis numerosis. \mathcal{L} in montanis siccis sylvaticis Amer. bor. à Canadâ (Hook.), et à Pensylvaniâ ad Carolinam (Pursh). Curt. bot. mag. t. 227. Hel. polycephalus Cass. dict. 20. p. 352. — Pluk. phyt. t. 159. f. 2. Tab. ic. t. 764. Invol. patentia imbricata nec squarrosa. Capitula erecta 2 poll. diam. Folia alterna, interdum variantia opposita, imò ternato-verticillata. (v. v. c.)
36. H. TUBEROSUS (Linn. sp. 1277), radice repente tubercula oblonga gerente, caule erecto ramoso scabro, foliis alternis petiolatis triplinerviis scabris serratis, inf. cordato-ovatis, super. ovato-acuminatis, petiolis basi ciliatis, invol. squamis lineari-lanceolatis ciliatis. \mathcal{L} in Brasiliâ (C. Bauh.), nunc in Europâ ad tuberes edules cultus sub nom. *Topinambour*. Jacq. h. vind. t. 161. Cass. dict. 20. p. 353. Gis. ic. f. 1. t. 19. Schkhr. handb. 3. t. 258. — Col. ecephr. 2. p. 11. t. 13. Pappus 1-4-aristatus aristis ciliatis. Ligulæ rudimenta organ. foemineorum retinent. Variat tuberibus subrotundis et oblongis subdulcibus ex Tausch in fl. 1828. p. 506. (v. v. c.)
- β . *oblongifolius*, foliis oblongo- seu ovali-lanceolatis. — Specim. sub nom. H. tuberosi ex horto Calcuttensi leger. cl. Heyne et Wallich, cat. n. 303. Sat similis H. decapetalo sed folia alterna. Radices ign. (v. s. comm. ab hon. Cur. merc. ind. or. Lond.)
37. H. HOOKERIANUS, caule simplici striato hispido, foliis alternis oblongo-lanceolatis obtusis longissimè in petiolum attenuatis, pedunculo elongato terminali 1-cephalo, invol. squamis patentibus lato-lanceolatis ciliatis, pappo aristis solitariis binisve inæqualibus validis et margine lacero constante. — in humidis plantationibus Columbiæ ad Oceanum pacificum, in vallibus montium scopulorum et in Californiâ ad San-Francisco. Hel. longifolius Hook. fl. bor. am. 1. p. 313 non Pursh. An fortè Leighiæ seu Harpalii spec.?
- ***** *Fruticosi*.
38. H. QUINQUERADIATUS (Cav. ic. 3. p. 38. t. 272), caule fruticoso, juniore subtomentoso, foliis alternis petiolatis ovato-lanceolatis basi cuneatis serrato-crenulatis utrinquè subtomentosis, corymbo subconferto, invol. squamis biserialis acuminatis, ext. apice reflexis, ligulis 5 ovatis, paleis rec. tridentatis obtusis. \mathfrak{J} in Mexico. An hujus generis?
39. H. ARGENTEUS (H. B. et Kunth nov. gen. am. 4. p. 221. t. 376), fruticosus ramosissimus, ramulis foliisque subutis sericeo-lanatis, foliis alternis breviter petiolatis lanceolatis integerrimis trinerviis suprâ sericeo-pilosis, capitulis pedicellatis solitariis erectis, invol. squamis oblongis acutis subsquarrosis, ligulis circ. 15, paleis oblongo-lanceolatis membranaceis, achæniis glabris.

♂ in Andibus Assuayensium prov. Quitensis. Achænia dicuntur biaristata. An rectè? Habitus Leighiæ.

40. *H. VERBESINOIDES* (H. B. et Kunth nov. gen. am. 4. p. 221), fruticosus ramosissimus, ramulis lanatis, foliis alternis petiolatis ovato-lanceolatis acuminatis integerrimis trinerviis suprâ adpresso-pilosis subtus incanis et sericeo-tomentosis, capitulis subcorymbosis, invol. squamis oblongis obtusis ext. subaquarosis, ligulis circ. 10, paleis linearibus obtusis, achænio glabro. ♂ in temperatis Peruviz juxta Guacabamba. An Leighiæ sp.?

† *Species non satis notæ.*

41. *H. ? ARISTATUS* (Ell. sketch 2. p. 428), caule erecto scabro, foliis infer. oppositis arcuè sessilibus ovali-lanceolatis acutis dentatis scabris subtus pilosis, capitulis paucis corymbosis, invol. squamis ovato-lanceolatis pubescentibus, paleis ovato-oblongis, aristis 2 persistentibus. ♀ in siccis sterilibus Georgiæ occid. Car. Actinomeridis ex Ell. sed folia non decurrentia.
42. *H. ? COCHINCHINENSIS* (Pers. ench. 2. p. 476. n. 21), caule recto scabro parè ramoso, foliis alternis lanceolatis integerrimis enerviis scabris, capitulis solitariis terminalibus, invol. squamis oblongis imbricatis patenti-squarrosis, ligulis multis longissimis, pappo biaristato. — in Cochinchinâ et Chinâ cultus. *H. giganteus* Lour. fl. coch. ed. w. a. p. 623 non Linn. nec Cav.
43. *H. FRONDOSUS* (Linn. amœn. 4. p. 290), caule solitario infernè glabro, foliis ovatis serratis, pedunculis scabris, invol. squamis squarrosis undulatis frondosis ciliatis hispidis. ♀ in Canadâ. Species dubia, hodiernis ignota et fortè varietas. *H. decapetali* invol. in folium mutato. Confer Hooker fl. bor. am. 1. p. 312.
44. *H. HIRSATUS* (Raf. ann. nat. 1820. p. 14), hirsutus, caule simplici, foliis remotis breviter petiolatis lanceolatis acutissimis serratis subtus pallidis subtrinerviis, capitulis terminalibus solitariis subsessilibus, invol. squamis adpressis ovatis acutis ciliatis striatis. ♀ in collibus Kentuckensibus.
45. *H. INCANUS* (Pers. ench. 2. p. 475), incano-tomentosus, foliis ovatis integerrimis trinerviis, invol. villosis. — in Peruviz. *H. canescens* Juss. herb. Affinis dicitur *H. molli*. Cæt. ign.
46. *H. PARVIFLORUS* (Bernh. ex Spreng. syst. 3. p. 617 non Kunth), caule herbaceo ramoso glabro, foliis oppositis alternisque petiolatis ovato-oblongis acuminatis subserratis triplinerviis scabriusculis, invol. squamis linearibus laxissimis subreflexis integerrimis. — in Amer. bor. Cæt. mihi ign. Bernh. descr. in libris mihi obvis non reperire quivi.
47. *H. TRILOBATUS* (Link enum. h. ber. 2. p. 352), caule herbaceo hirsuto supernè ramosissimo, foliis longè petiolatis trilobatis scaberrimis, lobis serratis acutis, invol. squamis pilosis, exter. acutis reflexis, achænio breviter biaristato. ♀ in Mexico. Capitulum in genere minimum. An certè generis?

†† *Species exclusæ.*

- H. AUREUS* H. B. K. = *Harpalium aureum*.
H. BUPHTHALMOIDES Hort. = *Heliopsis canescens*.
H. CORDATUS Lamb. = *Heliopsis canescens*.
H. CORYMBOSUS Poepp. = *Flourensia thurifera*.
H. DENTATUS Cav. = *Viguiera dentata*.
H. EXCELSUS Willd. = *Tithonia excelsa*.
H. GIGANTEUS Cav. = *Tithonia excelsa*.
H. GLUTINOSUS Hook. et Arn. = *Flourensia thurifera*.
H. LÆVIS Linn. f. = *Heliopsis lævis*.
H. LINEARIS Cav. = *Leighia linearis*.
H. MICROPHYLLUS H. B. K. = *Leighia microphylla*.

- H. OLIVERA Wall. = *Guizotia oleifera*.
 H. PROSTRATUS Willd. = *Viguiera prostrata*.
 H. RIGIDUS Desf. = *Harpalium rigidum*.
 H. RIVULARIS Poepp. = *Leptocarpus rivularis*.
 H. SARMENTOSUS Rich. = *Wulfia membranifolia*.
 H. SPECIOSUS Hook. = *Leighia speciosa*.
 H. AQUAROSUS H. B. K. = *Leighia linearis*.
 H. TETRAPTERUS Ort. = *Actinomeris tetraptera*.
 H. THURIFERA Mol. = *Flourensia thurifera*.
 H. TRINARVIS Hort. = *Tithonia excelsa*.
 H. TRIQUETER Ort. = *Viguiera triquetra*.
 H. TRUXILLENSIS H. B. K. = *Harpalium Truxillense*.
 H. TUBÆFORMIS Ort. = *Tithonia tubæformis*.

CCCXLVII. FLOURENSIA.

Capitulum multiflorum homogamum aut heterogamum, fl. radii ligulatis sterilibus neutris seu organa fœminea rudima. gerentibus, fl. disci hermaphr. 5-dentatis. Invol. 2-3-seriale, squamis oblongis subæqualibus. Recept. subconvexum, paleis complicatis deciduis flore brevioribus. Cor. disci faux vix tubo latior. Styli rami exerti elongati ferè linguiformes obtusi extus parcè hispidi. Achænia compressa basi attenuata villosissima 1-4-aristata, aristis membranaceis lineari-subulatis ciliato-fimbriatis, 1-2 longioribus, 2 lateralibus minoribus sæpè nullis. — Suffrutices aut frutices Americani glaberrimi sæpius glutinosi. Folia alterna sessilia oblongo-lanceolata coriacea sæpius penninervia reticulata. Pedunculi axillares terminales subcorymbosi. Corollæ omnes luteæ. — Genus spectabile dicatum clar. et amicissimo Flourens, physiologo acutissimo, Acad. scient. Paris. secretario perpetuo.

* *Capitulis radiatis*, Chilenses.

1. F. THURIFERA, glaberrima, foliis breviter petiolatis oblongo-lanceolatis repando-serratis penninerviis, capitulis radiatis, invol. squamis oblongo-linearibus. 5 in Chili circa Valparaíso (Gay! Poepp.! pl. exs. n. 235) et Coquimbo (Gaudich.) in fruticetis sterilibus calidisque collinis (Bert.!). *Helianthus thurifer* Mol. chil. ed. gall. p. 336 ex Gay, Bertero et Poepp. et ex usu popul. Hel. glutinosus Hook. et Arn. pl. Beech. 1. p. 32. *Diomedea thurifera* Bert.! herb. Colla mem. acad. Turin. 38. p. 35. n. 76. t. 31. Vulgò Chilensibus *Maravilla* seu *Maravilla del Campo*. Resinam thuris succedaneam fundit. (v. s.)

β. *angustifolia*, foliis lineari-oblongis integerrimis aut vix subserrulatis, achæniis disci exaristatis aut 1-2-aristatis. — in Chili legerant cl. Hæmke, Dombey et Née. An fortè species distincta? (v. s.)

2. F. CORYMBOSA, glabrescens, foliis breviter petiolatis ovali seu oblongo-lanceolatis integerrimis triplinerviis minutissime puberulis, capitulis radiatis corymbosis, invol. squamis lanceolatis. 4 5 in Amer. australi legit cl. Poeppig. *Helianthus corymbosus* Poepp.! pl. exs. n. 791. Caulis striatus. Capitula longè pedicellata. Invol. squamæ ext. 2-3-seriales rigidæ adpressæ lanceolatæ, intimæ breviores. Recept. convexus. Paleæ subcutæ. Achænia villosissima longè biaristata, squamellis 1-2 hinc inde intermediis. (v. s. comm. à cl. Poeppig.)

** *Capitulis discoideis*, Mexicanæ.

3. F. LAURIFOLIA, glaberrima, foliis petiolatis elliptico-lanceolatis integerrimis acuminatis, capitulis discoideis erectis. 5 in Mexico inter Vittoria et Tula legit

- cl. Berlandier (pl. exs. n. 22051). Arbor 5-10-pedalis mense nov. florens. Folia juniora et involucria glutinosa. Invol. squamæ adpressæ subobtusæ disco breviores. Achænia villosissima, compressa aut subtriquetra, aristis 1-2 superata. (v. s. sp.)
4. F. CERNUA, caule ramisque subpuberis incanis, foliis subsessilibus ellipticis utrinque acutis integerrimis suprà glabris subtus pubescentibus, capitulis discoideis cernuis. 5 in Mexici provinciâ Novâ-Leoninâ ad Monterey legit cl. Berlandier (pl. exs. n. 1401). Frutex ramosissimus cinérascenti-viridis. Folia 7-9 lin. longa, 3-5 lata. Capitula ad apices ramorum subracemosa aggregata. Pedicelli foliosi apice recurvi. Achænia villosissima biaristatâ. (v. a.)

DIV. IV. BIDENTIDÆ Less. syn. 229. — Coreopsidearum gen. Cass. opusc. 3. p. 58.

Capitula heterogama fl. radii neutris, rarissimè homogama discoidea. Achænia apice rostrata, rariùs calva, sæpiùs producta in aristas 1-4 ferè semper retrorsum aculeatas.

CCCXLVIII. CAMPYLOTHECA Cass. dict. 51 (1827). p. 475. Less. in linnæâ 1831. p. 508. syn. p. 230. — Bidentis sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis multinerviis neutris, disci hermaphr. tubulosis 5-dentatis, tubo brevi, fauce longissimâ 10-nerviâ. Recept. planum paleis inter flores instructum. Invol. pauciseriale, foliolis oblongo-ellipticis demum reflexis. Achænia conformia elongata subcontorta aut curva subtetragono-obcompressa angustissimè alata, alis apice in mucrones minimos sæpè deciduos aut subnullos abeuntibus. — Frutices (aut fortè herbæ) ex insulis Polyneziæ orti, erecti trichotomi. Folia opposita ternatim biternatimve secta, segmentis ovato-lanceolatis dentatis aut integris. Capitula lutea pedicellata.

1. C. MICRANTHA (Cass. l. c.), fruticosa glabra, foliis ternatim pinnatimve sectis, segmento impari sæpè trilobo, fructibus incurvis apice biaristulatis. 5 in ins. O-Wahu (Cham.), et insulis Sandwicensibus (Gaudich.). Bidens micrantha Gaudich. bot. voy. Freyc. p. 464. t. 85. Camp. australis Less. in linnæâ 1831. p. 509. Capitula 2 lin. longa. Cor. lineam longæ.

Æ? australis, Coreopsis fruticosa Forst. prod. 91. Bidens australis Spreng. syst. 3. p. 458. An stirps australis eadem ac Sandwicensis? Bidens pinnatifida Forst. in h. Mus. Par.? Specim. valdè imperfectum. Differt toto cœlo à pl. Gaudichaudi. Caulis fruticulosus gracilis binec trichotomus. Folia superiora pinnatilobata, lobis linearibus integerrimis. Capitula videntur minora sed fl. fruct. et invol. destituta. Ex Forst. prod. crescit inter tropicos. (v. s. in h. Mus. Par.)

2. C. GRANDIFLORA, herbacea? puberulo-hirtella, foliis ternatim sectis, segmentis lanceolatis dentato-serratis, fructibus subspiraliter tortilibus nudis aut junioribus vix bisetosis. — in ins. O-Wyhee Sandwicensium ad montem Kaah legit cl. Macrae. Capitula 7 lin. longa. Cor. 4-5 lin. longæ. (v. s. comm. ab hon. soc. hortic. Londin.)

CCCXLIX. BIDENS Linn. gen. n. 932. Gartn. fr. 2. p. 412. t. 167. Less. syn. 230. — Bidens et Kerneria Mærch meth. 569 et 595. Cass. dict. 24. p. 397 et 402. 51. p. 473 et 59. p. 321, 328 et 329. — Pluridens et Edwarsia Neck. elem. n. 156 et 157. — Ceratocephalus Vaill. Rich. non Mærch.

Capitulum multiflorum, nunc homogamum discoideum, nunc in iisdem aut diversis speciebus sæpiùs radiatum, fl. radii ligulatis neutris. Invol. squamæ biserialis, seriebus similibus dissimilibusve. Recept.

planiusculum paleaceum, Styli rami cono brevi superati. Achænium plius minus obcompressum aculeatum rostro ab achænio vix diverso superatum, desinens in aristas 2-5 rigidas retrorsum piloso-scabras. — Herbarum sæpius annuæ, pleræque Americanæ, paucæ Gerontogæ. Caules tetragoni teretesve sæpius oppositè ramosi. Folia opposita, summa interdum alterna, indivisa, partita aut secta, lobis incisiss serratisve. Capitula sæpius flava, rarius radio albo, in unicâ specie disco radioque purpureis.

Sect. I. PLATYCARPÆ.

Achænia ovalia seu obovato-cuneiformia obcompressa - plana setulis omnibus aut superioribus retrorsis ad margines ciliata, imò basi vix in areolam dilatata. Capitula discoidea lutea aut radiata, ligulis flavis rarius albis. Folia sæpius indivisa rarissimè in segmenta divisa.

1. *B. TRIPARTITA* (Linn. sp. 1165), glabra, foliis tripartitis, lobis lanceolatis dentatis, capitulis discoideis basi bracteatis, bracteis basi ciliatis capitulum superantibus, invol. squamis marg. vix membranaceis, achæniis sæpius biaristatis. ① in Europæ fossis et inundatis, etiâ ad Caucasum (Stev.!), et in Sibiria et Dahuriâ (Turcz.!). Huds. fl. angl. 2. p. 355. opt. Sow. engl. bot. t. 1113. *B. frondosa* α Lam. dict. 1. p. 413. ill. t. 668. f. 4. *B. radiata* Thuil. ! fl. Par. ed. 2 p. 422. quæ discoidea in ipsissimo auct. specim. — Blakw. t. 519. *Achænia*, ut in *B. cernua*, obovato-cuneata obcompressa erecta. (v. v.)
 - ℓ. *pinnatifida* (Turcz. ! in litt.), foliis tripartitis, lobis angustis pinnatifidis. — in Dahuriâ ad flum. Argun. (v. s. sp.)
 - γ. *hybrida*, foliis sub-5-partitis, lobis longo-linearibus. — circa Parisios. *Bidens hybrida* Thuil. ! fl. Par. ed. 2. p. 422.
 - δ. *tenuis*, capitulis paucifloris, foliis indivisis aut vix basi grossius tridentatis. ① cum var α. — *Verbesina minima* Dill. ex Ray syn. angl. 188. t. 7. f. 2. *B. minima* Linn. sp. 1165. ex syn. Ray? *B. cernua tenuis* Turcz. ! in litt. Similis var. γ *B. cernua* mihi ignotæ, sed achænia bi- nec 4-aristata. Ab Hudsonio tantum ritè distincta. (v. v. et s.)
2. *B. COMNATA* (Muhlenb. in Willd. sp. 3. p. 1718), caule glabro ramoso, foliis glabris, infimis ternatisectis lobis lateralibus in petiolum decurrentibus, super. indivisis oblongo-lanceolatis, utrinquè attenuatis serratis, capitulis discoideis breviter pedicellatis, invol. squamis foliaceis disco longioribus, achænio ad margines retrorsum ciliato apice 2-3-aristato. ① in sylvis et agris Amer. bor. frequens à Canadâ ad Carolinam. Ell. sketch 2. p. 430. Affinis *B. frondosa* sed foliis et achænio faciliè differt. Varietatem foliis ternato-verticillatis notat cl. Hooker fl. bor. am. 1. p. 314. (v. s. comm. à cl. Janin.)
3. *B. FRONDOSA* (Linn. sp. 1166), caule lævi sulcato, foliis glabris, infer. pinnatisectis hijugis cum impari, mediis trisectis, lobis omnium lanceolatis serratis, capitulis pedicellatis discoideis erectis, invol. ext. squamis foliaceis basi ciliatis disco longioribus, achæniis biaristatis marginibus subciliatis, pilis paucis supernè retrorsis infernè erectis. ① in humidis Americæ borealis in Canadâ (Hook.), et à Pensylvaniâ ad Carolinam. Ell. sketch 2. p. 431. Willd. sp. 3. p. 1718. — Moris. ox. 16. t. 5. f. 21. mal. Invol. nunc paulò nunc multò disco longius. *Achænia subrugosa*. (v. s. sp. et v. c.)
4. *B. CERNUA* (Willd. sp. 3. p. 1716), glabra, foliis indivisis longè lanceolatis dentatis, capitulis cernuis basi bracteatis, bracteis subintegris capitulo longioribus, invol. squamis margine membranaceis coloratis, achæniis 4-aristatis margine retrorsum ciliatis. ① in Europæ et nunc in Amer. borealis fossis inundatis et aquaticis frequens. Sm. fl. brit. 858. DC. fl. fr. n. 3288.
 - α. *discoidea*, capitulis discoideis cernuis caule ramoso. *B. cernua* Linn. sp. 1165. Engl. bot. t. 1114. (v. v.)
 - ℓ. *radiata*, capitulis radiatis. *Coreopsis Bidens* Linn. sp. 1281. — *B. cer-*

nna Schkuhr handb. t. 234. Barr. le. t. 1209. — Interdum in eadem speciminibus discoidea et radiata. Fl. dan. t. 841. (v. v.)

7. *minima*, capitulis discoideis erectis caule subsimplici. Bid. minima Linn. sp. 1165? excl. Ray. syn. — Fl. dan. t. 312. †.

5. *B. HELIANTHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 230), erecta ramosa glabra, caule teretiusculo, foliis sessilibus lanceolatis serratis, pedunculis terminalibus et axillaribus 1-cephalis, capitulis radiatis, invol. squamis ovalibus obtusis, exter. foliaceis subpatentibus, int. subcoloratis, achæniis obcompressis oblongo-cuneatis margine retrorsum ciliato-serratis apice biaristatis. ♀ in paludosis et fossis Mexici ad montem Chapultepec (H. et B.), ad Yotla (Andr. pl. exs. n. 312), in montibus circa urbem (Berl. pl. exs. n. 605), et in Amer. austr. circa urbem Angustura (Bert.), et circa Buenos-Ayres (Bacle!). *Kernæria helianthoides* Cass. dict. 24. p. 399. *Coreopsis serrata* ic. fl. mex. ined. Flores lutei. Ab omnibus differt achæni marginibus retrorsum serrato-scabris. (v. s.)

6. *B. CHRYSANTHEMOIDES* (Michx. fl. bor. am. 2. p. 136. non Pursh), glabra erecta, foliis oblongis utrinque attenuatis dentato-serratis basi connatis, capitulis cernuis radiatis, invol. squamis ext. oblongis rigidulè ciliatis patulis, inter. ovalibus margine membranaceis, ligulis ellipticis invol. duplò longioribus, achæniis retrorsum scabris biaristatis. ① in oryzetis et sphagnosis Carolinæ (Ell. sketch 2. p. 430), et ad oram occid. Amer. bor. (Hook. fl. bor. am. 1. p. 314). *Coreopsis Bidens* Walt. car. 215 C. perfoliata Bosc! herb. Affinis B. cernue et B. quadriaristatæ, sed achænia bi- nec 4-aristata, B. helianthoidi sed invol. squamæ ext. rigidè ciliatæ. (v. s. ex h. Bosc et Fraser.)

7. *B. QUADRIARISTATA*, glabra, foliis oblongis utrinque attenuatis dentatis basi connatis capitulis erectis, invol. ext. longo patente squamis foliaceis linguiformibus basi rigidè ciliatis, ligulis invol. interiore quadruplò longioribus ovalibus, achæniis 4-aristatis. ① in Amer. bor. humidis, circa Bostoniam. *B. chrysanthemoides* Pursh fl. bor. am. 566. Rig. fl. bost. ed 2. p. 294. non Michx. Capitula amplissima aurea. Ligulæ 12-15 lin. longæ, 6-8 latæ. Paleæ rec. apice coloratæ. (v. s.)

8. *B. BECKII* (Torr. in Spreng. nep. entd. 2. p. 135), foliis infer. capillaceo-multifidis, super. lanceolatis connatis acutè serratis, pedunculis terminalibus 1-cephalis, ligulis invol. longioribus. ♀? in Amer. bor. stagnosis aquis ad Schenectady Noveboracensium (Torr.), in Noveboraco occid. (Beck), et ad flum. St.-Laurent prope Mont-Real. Gold. pl. canad. in Edimb. phil. joura. 1822. apr. Beck bot. p. 207. Folia inf. omninò Ranunculi aquatilis sed opposita, sup. sæpè variè incisa. Ligulæ oblongæ, 6 lin. longæ, 1 ½ lin. latæ. (v. s. sine fr.)

9. *B. BULLATA* (Linn. sp. 1167), caule hirtio, foliis aliis indivisis latè ovatis grossè dentatis scabris, aliis tripartitis lobis lateralibus ovali-lanceolatis parvis, terminali folia indivisa referente, capitulis terminalibus discoideis, invol. ext. squamis ovatis ciliatis disco paulò longioribus, achæniis basi sursum, apice retrorsum ciliatis, biaristatis. ① in Italiâ ad Patavum, Ticinum, etc., ex Americâ dicitur orta, tamen in floribus Americæ nullibi indicatur. Willd. sp. 3. p. 1723. Ard. specim. 2. p. 37. f. 18. optim. Cor. luteæ. Folia interdum omnia indivisa, crassa, et quasi bullata. (v. v. c.)

10. *B. MODIFLORA* (Linn. sp. 1165), caule hirtio dichotomo, foliis ovatis oblongis utrinque 1-2-dentatis, capitulis ad alas sepè apices ramorum inter foliorum par ultimum sessilibus solitariis discoideis, invol. cylindrico, achæniis aristatis 4 (2 longiores et 2 breviores) gerentibus. ① in Benghalo Willd. sp. 3. p. 1716. — Dill. elth. 5a. t. 44. f. 52.

11. *B. REPENS* (Don prod. fl. nep. 180?), glabra, radice repente, foliis indivisis tri-antè rarius pinnatipartitis, lobis oblongo-lanceolatis dentibus acutiusculis, terminali longiore, capitulis discoideis erectis sparsè bracteatis, invol. squamis acutis inter se æqualibus, achæniis apice biaristatis. ♀ in Nepaliâ B.

tripartita Roxb. h. calc. p. 60. Wall. cat. et herb. n. 297. A. excl. litt. seq. Valde accedit ad B. tripartitam, sed radice repente perenni, foliorumque lobis angustioribus differt. (v. s. comm. ab ampl. aulâ merc. angl. Ind. or.)

Sect. II. PSILOCARPÆA.

Achænia lineari-tetragona, glabra aut margine sursum ciliata, basi vix in areolam subdilata. Capitula raro discoidea flava, sæpius radiata, ligulis flavis seu in paucis albis.

* *Foliis indivisis integris dentatisve sæpius lanceolatis.*

12. B. FRUTICOSA, fruticosa glabra, ramis angulatis, foliis oppositis oblongis lanceolatis acutè dentatis in petiolum attenuatis, pedunculis axill. et termin., capitulis radiatis, ligulis 12 ovato-oblongis tridentatis, achæniis subtetragonis 4-arithatis. 5 patr. ign. Coreopsis fruticosa Vest in florâ 1820. p. 7. non Forst., sed auctor ipse dixit aristas retrò aculeatas undè ad Bidentem pertinet. Ligulæ aurantiacæ.
13. B. DECOLORATA (H. B. et Kunth nov. gen. am. 4. p. 233), caule erecto ramoso glabro tetragono, foliis petiolatis lanceolatis serratis suprâ pilosiusculis, capitulis subcorymbosis longè pedunculatis radiatis, radio heterochromo, pedunculis pubescentibus, invol. squamis linearibus ciliatis, exter. patulis, achæniis biaristatis. 2/ in Mexico inter Capula et Valladolid de Mechoacan. Ligulæ albæ obovato-ellipticæ multinerviæ 5 lin. longæ.
14. B. TETRAGONA, glabra, caule tetragono, foliis oppositis lanceolatis, infer. serratis, super. integris, breviter petiolatis subconnatis, pedunculis longissimis solitariis, capitulis radiatis, invol. squamis exter. linearibus ciliatis, inter. oblongis subcoloratis, ligulis 8 ovatis crenatis bidentatis, achæniis oblongis biaristatis. — in pratis humidis et fossis circa urbem Mexici. Coreopsis tetragona Lallav. et Lex. nov. veg. 1. p. 31. Cl. auct. ipsi dicunt aristas setis introflexis instructas et idè species Bidentis. Ligulæ flavæ.
15. B. TRIPLINERVIA (H. B. et Kunth nov. gen. am. 4. p. 231), caule erecto sulcato ramisque pubescenti-hirtis, foliis petiolatis oblongo-lanceolatis grossè serratis triplinerviis utrinquè glabris, capitulis longè pedunculatis solitariis radiatis, invol. squamis lanceolatis subhispidulis, achæniis. . . . glabris biaristatis. 2/ in Mexico circa urbem et prope St.-Augustin de Las Cuevas (H. et B.). Ligulæ flavæ ellipticæ multinerviæ subsemipollicares.
16. B. HIRTELLA (H. B. et Kunth nov. gen. am. 4. p. 232), tota pubescenti-hirta, caule erecto sulcato subsimplici, foliis petiolatis lanceolatis serratis infernè subincisis, capitulis longè pedunculatis radiatis, invol. squamis lanceolatis adpressis, achæniis compressis biaristatis. 2/ in Mexico? cum B. triplinerviâ? cui valdè affinis: differt foliis rigidioribus non triplinerviis et notis indic.
17. B. PROCUMBENS (H. B. et Kunth nov. gen. 4. p. 232), caule procumbente simplici r-cephalo bifariam piloso, foliis petiolatis lanceolatis serratis triplinerviis hispidulis, pedunculo glabro elongato, capitulo radiato, invol. . . . , achænio biaristato. 2/ prope Xalapam Mexicanorum?
18. B. ARGUTA (H. B. et Kunth nov. gen. am. 4. p. 231), erecta glabra, caule tetragono oppositè ramoso, foliis lanceolatis petiolatis argutè serrato-dentatis, capitulis terminalibus subcorymbosis longè pedunculatis radiatis, invol. subæqualis squamis lanceolatis, ext. ciliatis patulis, achæniis scabris linearibus obcompressis biaristatis. 2/ in Mexico juxta Valladolid (H. et B.), ad Toluca et in Cordillera Guichilana (Berl. pl. exs. n. 1091 et 1001), ad Tlapujahu (Keerl ex h. Mart.). Cor. radii flavæ ovals obtusissimæ multistriatæ. Ach. aristæ apice tantum glochidiatæ. (v. s. comm. à cl. Alaman et Berlandier.)
β. luxurians (Willd. enum. 847), invol. exteriore interius superante. — in Mexico (Willd.) et fortè in insulis Sandwich ex Hook. et Arn. bot. Beech. p. 86.

19. *B. LONGIFOLIA*, herbacea glabra aut subglabriuscula, caule tereti, foliis inf. oppositis, super. alternis elongato-lineari-lanceolatis utrinque acuminatis integerrimis aut vix lobulo incisis, pedunculis longis 1-cephalis subcorymbosis, capitulis radiatis, invol. squamis ext. linearibus subpatulis scabro-ciliatis, achæniis lineari-subtrigonis glabris biaristatis. — in valle Mexicanâ legit cl. Berlandier (pl. exs. n. 717) et etiam à cl. Alaman recepi. Flores flavi. Folia distantia 3 poll. longa, 3 lin. lata. (v. s.)

** *Foliis aliis indivisis ovatis lanceolatisve, aliis basi incisis aut trisectis, segmentis serratis.*

20. *B. HETEROPHYLLA* (Ort. dec. p. 99. t. 12), caule simplici glabro vix apice pubescente, foliis glabris, inf. pinnatipartitis, mediis tripartitis, summis indivisis, lobis lineari-lanceolatis serratis, terminali longissimo, capitulis subcorymbosis erectis radiatis, invol. squamis subæqualibus ciliatis aliis linearibus aliis lanceolatis, achæniis bi- (rarius 3- ex Ort.) aristatis. ♀ in Mexico. Petioli vix subconnati basi subciliati. Capitula flava. Ligulæ oblongo-cuneatæ obtusæ disco duplò longiores. (v. v. c.)

21. *B. SERRATA* (Pavon in h. L'hér.), caule tetragono ramossissimo villosio, foliis petiolatis lanceolatis utrinque acutis subpuberulis serratis aliis indivisis aliis basi incisis imò subtrilobis, capitulis pedunculatis subsolitariis radiatis, invol. 10-phylo basi imprimis piloso, squamis adpressis ext. lanceolatis int. linearibus, achænio lineari-teretiunculo bicorni. — in Peruvîâ. Coreopsis L'hér.! uss. Capit. floridum 18-20 lin. diam. Ligularum achænium calvum (L'hér.! mss.). Folia cum petiolo 2 poll. longa, 6 lin. lata. (v. s. sine fl. in h. L'hér. et flor. descr. ex L'hér. mss.)

22. *B. DICHOTOMA* (Desf. cat. h. par. 1804. p. 104), caule tetragono glabro trichotomè ramoso, foliis petiolatis glabris, aliis indivisis ovatis acuminatis serratis, aliis basi 1- seu 2-lobis accessoriis terminali minoribus acutis et bi-trisectis, capitulis subcorymbosis pedicellatis discoideis, invol. squamis inter se et disco æqualibus, ext. patulis puberis, int. glabris erectis margine scariosis, achæniis lineari-tetragonis glabris 3-4-aristatis. ♀ in Brasiliâ ex specim. Commers.! Lund! Pers. ench. 2. p. 393. Poir. suppl. 1. p. 630. An *B. reflexa* Link enim. h. berol. 2. p. 306 ex Mexico orta? Habitus ferè *B. rubifoliæ*, sed discoidea. Pili pauci in ramulis. (v. v. c. et s. sp.)

23. *B. STRIATA* (Sweet brit. fl. gard. t. 237), caule erecto subtetragono striato puberulo oppositè ramoso, foliis glabris petiolatis trisectis, segmentis ovato-lanceolatis acuminatis serratis, medio paulò majore et petiolulato, capitulis corymbosis pedicellatis radiatis, invol. squamis lanceolatis puberis exter. subreflexis, achæniis lineari-subtetragonis margine scabris biaristatis. (1) ? in Mexico. Hook. bot. mag. t. 3155. Ligulæ amplæ albæ multiradiatæ.

*** *Foliis 3-5-sectis, segmentis ovato-lanceolatis serratis non verò parvis.*

24. *B. DONDÆIFOLIA* (Less. in linnæâ 1830. p. 155), glabra, ramis subtetragonis, foliis trisectis, segmentis sessilibus obovatis acutis obtusive apice serratis basi cuneatis, summis indivisis, capitulis radiatis geminis longissimè pedunculatis, invol. squamis exter. squarrosis, achæniis biaristatis. — prope Vera-Cruz.

25. *B. PILOSA* (Linn. sp. 1166), caule erecto subtetragono glabriusculo, foliis infer. impari-pinnatisectis, super. trisectis, omnium lobis glabriusculis ovatis acutis serratis vix decurrentibus, petiolis basi ciliatis, capitulis pedicellatis subcorymbosis discoideis rariùs radiatis, invol. squamis inter se et disco subæqualibus, achæniis lineari-angulatis glabriusculis 2-3-aristatis, exter. brevioribus aristis divergentibus, centralibus longioribus aristis rectis. (1) in agris et cultis Americæ bor. à Pensylvaniâ ad Carolinam (Pursh), unde fortè migravit in Teneriffam (Bach! Brouss.), ins. Mauritiî (Bout.), Novam-Zelandiam (Forst. Lesson). — Dill. elth. 1. t. 43, sed caulis nimis hispidus. *Ceratocephalus pilosus* Rich. cat. h. med. p. 91. *Kerneria dubia* Cass. dict. 24. p. 393. *Kerneria tetragona* Mœnch meth. 595. (v. v. c. et s. sp.)

26. *B. LEUCANTHA* (Willd. sp. 3. p. 1282), caule glabro subtetragono erecto, foliis glabrisculis, infer. impari-pinnatisectis 1-3-jugis, sup. unijugis, lobis omnium ovatis acutis serratis basi subæqualibus, capitulis subcorymbosis pedicellatis radiatis, radio heterochromo, invol. squamis lanceolatis inter se et disco subæqualibus, acheniis linearisubtetragonis, aliis glabris, aliis puberulis, 2-4-angulatis. (1) in Amer. merid. (Linn.), in Chili et insulis Caribæis (Bert.!), ad Bahiam in ruderatis (Salzm.!), in Mexico circa Jalapam (Schied.). Ex Amer. migravit ad Maderam (Lowe!), insulas Mauritanas ubi dicitur *Herbe à Sornet* (Bory!) et ad Cap. Bonæ-Spei (Boiss. herb.). Ex hac regione habui specim. (Eckl. n. 872. Drege! n. 5084) huic speciei simillimum sed aut discoideum aut ligulis casu quodam orbatum. Coreopsis leucantha Linn. sp. 1282a. amæn. ac. 4. p. 291. *B. leucantha* H. B. et Kunth fortè diversa? *B. chinensis* Willd. sp. 3. p. 1719 huc forsàn revocanda? — Icon Plum. ed. Burm. t. 53. f. 2 huc certè pertinet et ideò Coreopsis coronata Linn. sp. 1281 eadem, sed species obscura remanet et *C. coronata* Spreng. Steud. et hortul. omninò diversa. (v. v. c. et s. sp.)
27. *B. SUNDAICA* (Blum.! bijdr. p. 913), glabruscula ant subpubera, foliis inf. et super. ternatisectis, mediis subpinnatisectis, segmentis ovatis acuminatis serratis, lateralibus basi obliquè attenuatis, terminali ovato-lanceolato reliquis longiore, capitulis longè pedunculatis laxè corymbosis subradiatis, invol. biserialis squamis margine membranaceis, acheniis glabris apice triaristatis. — in graminosis humidis Javæ ad Buitenzorg (Bl.!), Zeylonæ (Lesch.!) et fortè Indiæ contin. *B. tripartita* Wall. cat. et herb. n. 3187. *A. Javanica* sub nom. *Harruja* fortè non immeritò cum *B. chinensi* confusa et hujus forsàn varietas? Achenia 4-5 lin. longa. (v. s.)
28. *B. WALLICHI*, glabruscula ant subpubera, foliis infer. pinnati-, super. ternatisectis, segmentis ovatis acuminatis inciso-serratis dentatisve, lateralibus basi obliquè attenuatis, capitulis longè pedunculatis laxè corymbosis subradiatis, invol. biserialis squamis subæqualibus, inter. ovali-oblongis margine membranaceis, acheniis linearibus angulatis striatis glabris aristas 3 apice gerentibus. — in locis incultis prope Bataviam (Blume) et in Indiâ contin. ad montes Dendigal (Wight!), Prome, Tavoy, Segain, etc. (Wall.!), eandem, ni fallor in Cafrorum regione legit cl. Drege, n. 5083? *B. chinensis* Wall. cat. n. 3189. comp. 299. DC. in Wight contr. p. 19 sed species homonyma Wall. sp. 3. p. 1719 ex Chinâ et Japoniâ orta videtur diversa. — *Agrimonia* Molucca Rumph. amb. 6. t. 15. f. 2 sed ligulæ dicuntur flavæ. Ligulæ in pl. spont. paucæ albæ neutræ. Achenia 5-6 lin. longa. (v. s. comm. ab hon. aulâ merc. angl. Ind. or.)
29. *B. PEDUNCULARIS* (Gaudich. voy. (1826) p. 464), glabra, foliis ternatim pinnatimve sectis, segmentis ovatis acuminatis argutè serratis, paniculâ trichotomâ, involucrio exter. interiori æquante, acheniis aristas 3 apice tantum hamosas gerentibus, radio brevi. — in insulâ Rawak (Gaudich.!), in O-Wahu archip. Sandwicensium (h. Acad. imp. Petrop.!). *B. Sandwicensis* Less. in linnæa 1831. p. 508. Simillima dicitur *B. leucanthæ*, sed radius luteus, achenia 3 lin. tantum longa, aristæ brevissimæ achenio nempe sextuplò breviores interdum nullæ. (v. s.)
30. *B. RUBIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 237. t. 381), glabra, caule tetragono, foliis trisectis, super. indivisis, segmentis serratis amprâ nitidis, lateralibus ovatis acutis sessilibus, medio maximo ovato-oblongo apice angustato-acuminato petiolato, capitulis terminalibus corymbosis radiatis, invol. squamis ext. linearibus patentibus, int. ovato-oblongis, acheniis linearisubtetragonis ad angulos hispidulis biaristatis. 2/ 5 in provinciâ Quitensi? *Kerneria rubifolia* Cass. dict. 24. p. 399. Ligulæ flavæ. Achenia 5 lin. longa.
31. *B. TERETICAULIS*, caule subsicante tereti glabro, ramulis vix striatis, foliis petiolatis trisectis, segmentis ovato-lanceolatis acuminatis serratis sparsè subpilosis, medio petiolulato, capitulis pedicellatis subpaniculatis radiatis, invol. squamis subæqualibus acutis, ext. reflexo-squarrosis, acheniis apice ciliatis

biaristatis. — in Mexico prope Tantoyuca legit Berlandier (pl. exs. n. 2150 et 2148); eadem ni fallor adest ex Brasiliæ prov. Sancti-Pauli in herb. Mus. Par. (h. imp. Bras. n. 412). B. squarrosa Less. in Linnæa 1830. p. 156 (è sylvis Jalapensis orta) excl. utroq. Kunthii syn. Differt à B. rubifolia, caule tereti nec tetragono, à B. squarrosâ foliis glabrisculis, ab utraq. capitulo minore, foliis etiâ superis trisectis et patriâ. (v. s.)

32. B. COREOPSISIS, glabra, caule tereti scandente, foliis oppositis petiolatis pinnatim trisectis, segmentis ovato-lanceolatis grossè dentatis acuminatis, capitulis subcorymbosis breviter pedicellatis radiatis, invol. squamis ext. linearibus patulis, ligulis oblongis integris, achæniis biaristatis. ½ in Jamaicâ (Bert.) inter sepes et dumeta (Sloan.). Coreopsis chrysantha Spreng.† syst. 3. p. 614 non Linn.—Sloan. hist. 1. p. 261. t. 154. f. 2 et 3. Caulis certè scandens aut jam dixit Sloane, fortè non repens. Flores lutei. Achænia immatura vidi. Aristæ jam in flore evidenter retrorsam aculeato-pilosæ. (v. s. comm. à cl. Bert.)

Æ? reptans, glabra, foliis aliis pinnatim sectis, aliis indivisis, ligulis apice denticulatis. ½ in Jamaicâ. Coreopsis reptans Linn. amæn. ac. 5. p. 407 (excl. syn. Brownei ex Smith et Sloanei ex Ker.). Smith spicil. 2. n. et t. 22.—An hæc aut var. æ, in Barbados à cl. Maycock lecta?

γ? incisa, fruticosa, petiolis foliisque pilosiusculis, foliis pinnatim sectis, aliis uni-, aliis bijugis cum impari, ligulis integerrimis. ½ in Jamaicâ. Coreopsis incisa Ker bot. reg. t. 7. An species propria?

33. B. SQUARROSA (H. B. et Kunth nov. gen. am. 4. p. 238), erecta, ramis teretibus glabris, foliis trisectis, super. indivisis, segmentis ovato-oblongis acuminatis serratis suprâ pilosiusculis subtus pubescenti-hirtis, later. sessilibus, medio longè petiolato majore, capitulis racemoso-corymbosis radiatis, invol. squamis ext. squarroso-reflexis linearibus, int. lanceolatis, achæniis lineari-subtetragonis ciliatis biaristatis. ½ inter. Caracas et montem Buenavista. Achænia 4 lin. longa.

34. B. FLORIBUNDA (H. B. et Kunth nov. gen. am. 4. p. 238), glabra erecta, ramis tetragonis, foliis infer. tripartitis?..., sup. indivisis ovatis acuminatis basi angustato-cuneatis serratis, corymbis terminalibus paniculatis foliatis, capit., invol. squamis ext. spatulato-linearibus patulo-reflexis, int. lanceolatis, achæniis lineari-tetragonis margine piloso-hispidis biaristatis. ½ prope Quito ad radices montis Javirac. Affinis B. squarrosæ dicitur.

35. B. ANDICOLA (H. B. et Kunth nov. gen. am. 4. p. 237), caule erecto quadrangulo glabrisculo, foliis trisectis pubescenti-hispidis, segmentis oblongis acutis serratis basi cuneatis, lateralibus sessilibus, medio majore petiolato, capitulis pedunculatis radiatis, invol. squamis lanceolatis, exter. hispidulis, achæniis lineari-subtetragonis scabrisculis biaristatis. ① in declivitate montis Chimborazo. Ligulæ albæ (Kunth), flavæ (Willd.). Achænia 3 lin. longa.

36. B. SAMBUCIFOLIA (Cav. lc. 3. p. 15. t. 219), glabra, foliis decursivè pinnatipartitis, lobis oblongo-lanceolatis serratis, ultimo paulò majore, capitulis radiatis, invol. squamis æqualibus disco brevioribus, ligulis disco duplò longioribus, achæniis 4-ariatis. ½ in Peruvâ (Cav. L'her. mss.) et fortè in Mexico? Cor. coccinæ. Ligulæ interdum bilabiatae. Species lobis fol. secus petiolum decurrentibus faciliè distincta. (v. v. c.)

37. B. HISPIDA (H. B. et Kunth nov. gen. am. 4. p. 237), caule erecto tetragono sulcato piloso-hispido, foliis infer. impari-pinnatisectis, super. trisectis, lobis ovatis acuminatis crenato-serratis secus petiolum plerisque decurrentibus in petiolo nervisque subtus hirsutis, capitulis pedicellatis subcorymbosis discoideis, invol. squamis inter se et disco subæqualibus, achæniis lineari-tetragonis biaristatis glabris. ① in siccis prov. Venezuelæ (H. et B.) et olim in hortis culta sub nom. B. decussatæ Pav. ex Peruvâ. Icon Dill. elth. 1. t. 43 hæc fortè pertinet excl. fig. 6-10. (v. s. cum descr. in h. L'her. nunc meo.)

38. B. CALIFORNICA, caule tetragono oppositè ramoso basi glabro apice scabrisculo, foliis parçè pilosiusculis impari-pinnatisectis infer. bi-super. 1-jugis.

segmentis ovatis acuminatis serratis, capitulis pedicellatis subpaniculatis discoideis aut ligulas paucas parvas gerentibus, invol. squamis lanceolatis subæqualibus, ext. patulis, omnibus demum reflexis, achæniis lineari-tetragonis apice subscabris 3-4-aristatis. (I) in Californiâ legit cl. Douglas. Achænia 4 lin. longa. Caulis circ. pedalis à basi ramosus. Capitula parva. (v. s. comm. ab hon. soc. hort. Londin.)

39. *B. EXARISTATA*, caule erecto tetragono glabriusculo, ramis oppositis tetragonis, foliis glabriusculis pinnatisectis, segmentis oppositis alternisve ovalibus aut ovali-lanceolatis dentato-serratis, terminali elongato, petiolis ad basin segmentisque subciliatis, capitulis pedicellatis laxè corymbosis radiatis, invol. squamis æqualibus acutis glabris, achæniis glabris lineari-subtetragonis apice attenuatis truncatis exaristatis. — in Mexico inter Vittoriam et Tulam legit cl. Berlandier (pl. exs. n. 2220). Herba circ. bipedalis ferè omninò glabra trichotoma laxè corymbosa. Ligulæ pluristriatæ pallidæ flavæ. Achænia longitudine in iisdem capitulis varia, et etiam per anthesin calva. (v. s.)
40. *B. SUBALTERNA*, glabra erecta, caule striato, foliis inf..., sup. alternis petiolatis pinnatisectis, segmentis sessilibus circ. bijugis cum impari, subalternis ovali-lanceolatis basi cuneatis apice acuminatis serratis, infer. sæpè tripartitis, capitulis pedicellatis subcorymbosis radiatis, invol. squamis ext. brevioribus linearibus patulis pedicellisque puberis, achæniis lineari-tetragonis glabris 3-4-aristatis. (I) in locis cultis circa Bahiam legit cl. Salzmann. Ligulæ flavæ. Achænia 4 lin. longa abortiva et interdum subhirta. (v. s. comm. à cl. Salzmann.)
41. *B. QUADRANGULARIS*, caule erecto tetragono glabriusculo oppositè ramoso, foliis petiolatis infer. pinnatisectis, segmentis 3 ovato-lanceolatis serratis petiolulatis medio longius stipitato, super. pinnatipartitis lobis bijugis cum impari oblongis acutis sessilibus subserratis, capitulis pedunculatis subcorymbosis radiatis, invol. squamis ext. linearibus ciliatis subpatulis, int. lanceolatis, achæniis lineari-tetragonis glabris 4-aristatis. (I) ? circa Rio-de-Janeiro legit cl. Ponson. Ligulæ paucæ ex sicco videntur albæ. Achænia 4 lin. longa, exteriora hinc inde 2-3-aristata. (v. s. comm. à cl. inv.)
42. *B. BRACHYCARPA*, glabra, caule erecto tetragono trichotomo, foliis pinnatisectis, segmentis 3-4-jugis ovalibus utrinquè acuminatis dentato-serratis impari paulò majore, infer. basi inciso-partitis, capitulis radiatis pedunculatis laxè corymboso-paniculatis, invol. squamis subæqualibus lanceolatis margine scariosis, achæniis linearibus scabridis 2-aristatis invol. paulò longioribus. — in Mexico ad Tampico de Tamaulipas legit cl. Berlandier (pl. exs. n. 5 et 113). Ligulæ paucæ in specim. meo disco paulò longiores, flavæ? Achænia 3 lin. longa. (v. s.)
43. *B. CARACASANA*, glabra erecta ramosissima, caule sulcato-angulato, foliis petiolatis pinnato-trisectis, segmentis ovali-lanceolatis basi cuneato-attenuatis apice acuminatis serratis, impari majore, capitulis longè pedunculatis laxè paniculatis radiatis, invol. squamis int. lanceolatis margine subscariosis, ext. linearibus subreflexis, achæniis lineari-tetragonis setulis rigidis valdè echinatis apice biaristatis. (I) circa Caracas legit cl. Vargas. Ligulæ obovatæ, obtusæ in sicco pallidæ, disco elevato paulò longiores, patentes. Achænia 5-6 lin. longa. (v. s. comm. à cl. Vargas.)
44. *B. OXYODONTA*, glabra erecta, caule tetragono subramoso, foliis pinnatisectis, segmentis petiolatis lanceolatis utrinquè acuminatis argutè serratis 2-3-jugis, inferioribus sæpè trisectis, terminali indiviso aut tripartito, capitulis pedicellatis in paniculam dispositis radiatis, invol. squamis lanceolatis disco subæqualibus, ext. subpatulis, achæniis lineari-tetragonis apice subscabridis biaristatis. — in Sancto-Domingo legit Bertero. Coreopsis coronata Spreng. ! syst. 3. p. 614 sed ab icone Plum. à Linnæo citatâ certè diversa et Bidenti nec Coreopsidi adgreganda. Ligulæ pallidæ in sicco. (v. s.)
45. *B. ANTHRISCOIDES*, tota pubescenti-hirta, caule suberecto tetragono, foliis petiolatis trisectis, inferiorum segmentis ovatis acutis subæqualibus medio vix

dissimili, superiorum segmentis lateralibus ovatis medio duplò longiore petiolato lanceolato utrinque attenuato, omnibus dentatis, ramis floridis nudis elongatis 1-3-cephalis discoideis, invol. squamis disco et inter se subæqualibus dorso pubescenti-hirtellis, achæniis lineari-tetragonis apice scabris biaristatis. ① in Mexici Cordillera de Guachilagua legit cl. Berlandier (pl. exs. n. 1010 et 1152) octob. flor. Capitula parva, in sicco flavo-subfusca. Radix obliqua seu caulis stricto sensu adscendens sed ferè erecta. Achæn. 6 lin. longa. (v. s.)

β. *angustiloba*, caule foliisque glabriusculis, foliorum lobis elongato-lanceolatis in apice caulis ferè linearibus. ① in Mexico circa urbem legit etiam Berlandier (pl. exs. n. 875). (v. s.)

46. *B. PORTORICENSIS* (Spreng. l. in litt. ad Balb.), caule glabro erecto tetragono, foliis glabris petiolatis, aliis biternato-sectis, aliis impari-pinnatisectis, omnium segmentis ovatis acutis basi cuneatis dentato-serratis, capitulis pedicellatis discoideis, invol. squamis disco brevioribus acutis inter se æqualibus, achæniis lineari-tetragonis glabris 4-aristatis. ① in Porto-Ricco legit cl. Bertero. Valdè affinis *B. bipinnatæ*, sed satis distincta notis indicatis videtur. (v. s. comm. ab amic. Bertero.)

47. *B. DAUCIFOLIA*, caule suberecto tetragono sulcato pubescenti-hirtello, foliis villosa-hirsutis pinnatipartitis, lobis circ. 3-jugis cum impari lineari-lanceolatis inciso-serratis acutis, pedunculis paucis 1-cephalis, capitulis radiatis, invol. squamis disco brevioribus inter se æqualibus ciliatis, achæniis lineari-tetragonis striatis scabriusculis biaristatis. — in Mexico legit cl. Berlandier fortè in valle Toluacanà (commixt. sub n. 1138). Ligulæ disco longiores ovals obtusissimæ multistriatæ flavæ 3-4 lin. longæ. (v. s.)

48. *B. RIPARIA* (H. B. et Kunth nov. gen. am. 4. p. 236), erecta, caule ramisque glabris, foliis duplicato-trisectis suprâ pilosiusculis margine ciliato-scabris, lobis ovatis inciso-serratis, sup. trifidis, lobulis terminalibus angustato-acuminatis, capitulis pedunculatis radiatis, invol. hispidi squamis exter. linearibus duplò longioribus, achæniis lineari-tetragonis 5-aristatis, aliis glabris, aliis brevioribus scabris. ① in calidissimis ad ripam flum. Magdalene prope Badillas.

49. *B. ABADIAE*, caule tetragono sulcato hispido erecto ramoso, foliis utrinque villosa-hirsutis bipinnatisectis segmentis 2-3-jugis ovatis grossè inciso-dentatis cum terminali trilobo basi cuneato, dentibus obtusis mucronatis, capitulis pedunculatis laxè et rigide paniculatis radiatis, invol. squamis glabris lanceolatis subæqualibus disco brevioribus, achæniis lineari-subtetragonis vix involucri longioribus apice subciliatis biaristatis. — in Peruvia circa Limam legit cl. Abadia. Ligulæ ex sicco albæ 5-striatæ. Caulis apice subglabratus. (v. s. comm. à cl. Abadia.)

*** *Foliis variè sectis, segmentis angustis integris linearibus aut multifidis, lobis linearibus subintegris,*

50. *B. ANGUSTISSIMA* (H. B. et Kunth nov. gen. et. sp. 4. p. 233), erecta glabra, ramis tetragonis, foliis tripartitis, lobis angustissimè linearibus acuto-mucronatis integerrimis subcarinosis, intermedio longiore, capitulis terminalibus longè pedunculatis radiatis, invol. squamis exter. patentibus brevioribus, int. oblongis extus hispidulis, achæniis compressis biaristatis. ② in frigidis Mexici prope Los Joares et Santa-Rosa de la Sierra (H. et B.). Cor. flavæ. Achænia matura ign.

51. *B. CRITHMIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 234), caulibus diffusis procumbentibus hirtis pubescentibus, foliis pinnatum quinque-sectis glabris, lobis linearibus integerrimis, capitulis longè pedunculatis solitariis radiatis, invol. squamis exter. linearibus obtusis, inter. lanceolatis, achæniis 4-gonis? biaristatis? ②? prope Quilo. Ligulæ flavæ.

52. *B. GLABERRIMA*, herbacea erecta ex omni parte glabra, ramis (caulibusve) teretibus, foliis oppositis petiolatis pinnato-multifidis internodio multò longioribus, lobis oblongo-linearibus crassiusculis, capitulis solitariis pedicellatis terminalibus radiatis, invol. squamis ext. linearibus obtusis subpatulis, int.

oblongis, ligulis invol. triplò longioribus multinerviis, acheniis glabris biaristatis. — in Amer. calidiore legit cl. Née ex h. Thibaud, sed locus propr. ign. Ligulæ flavæ ellipticæ obtusæ, 6-7 lin. longæ, nervulis 15-16 nigrellis striatæ. (v. s.)

53. *B. HUMILIS* (H. B. et Kunth nov. gen. am. 4. p. 234), diffusa subprocumbens glabra, foliis irregulariter bipinnato-quinque-partitis, lobis lineari-lanceolatis, petiolis ciliatis, capitulis longè pedunculatis solitariis radiatis, invol. squamis ext. linearibus obtusis, int. lanceolatis, acheniis lineari-quadrangularibus scabriusculis biaristatis. $\frac{1}{2}$ in radicibus montis Cotopaxi prope Malalo Quitensium. Ligulæ flavæ oblongo-ellipticæ multinerviæ 3 lin. longæ. Achenia 2 $\frac{1}{2}$ lin. longa.
54. *B. SCANDICINA* (H. B. et Kunth nov. gen. am. 4. p. 235), caule erecto angulato glabro, ramis pilosis, foliis pinnato-quinque-partitis subtus hispidis, lobis incisis, inf. tripartitis, petiolis ciliatis, pedunculis pilosis, capitulis radiatis, invol. squamis ext. linearibus, int. lanceolatis, acheniis linearibus compresso-quadrangularibus scabriusculis 2-3-aristatis. (1) prope Llaetacunga et Hambato Quitensium. Ligulæ flavæ ex Kunth, albæ ex Bonpl. Achenia 5 lin. longa.
55. *B. ALAUSKENSIS* (H. B. et Kunth nov. gen. am. 4. p. 235), caule erecto glabro, ramis tetragonis, foliis pinnato-quinque-partitis pilosiusculis, lobis oblongis inciso-serratis, petiolis ciliatis, capitulis longè pedunculatis radiatis, invol. squamis ext. linearibus, int. lanceolatis, acheniis linearibus compresso-quadrangularibus scabriusculis biaristatis. $\frac{1}{2}$ prope Alausi Quitensium (H. et B.), et in summo Serro-Colorado Mexici (Schied.). *B. diversifolia* Willd. herb. Ligulæ flavæ. Achenia 4 lin. longa. Tota plus minus hirsuta ex Less. in Linnaea, p. 155.
56. *B. DELPHINIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 234), erecta glaberrima, caule sulcato supernè ramoso, foliis irregulariter bipinnatisectis, lobis linearibus, capitulis longissimè pedunculatis. . . , invol. . . , acheniis linearibus tetragonis scabriusculis triaristatis (1) ? ad urbem Quito? Ligulæ flavæ?
57. *B. PARVIFLORA* (Willd. enum. 840), glabra, foliis multipartitis, lobis infimorum oblongo-lanceolatis incisis, summorum linearibus, capitulis discoidis longè pedunculatis laxè corymboso-paniculatis, invol. biseriali, interiore demum exterius longè superante, acheniis sparse pubescentibus longissimas aristas 2 rectas apice gerentibus. (1) in Sibiria ad lacum Baikal (Fisch.), in ripibus ad Selengam (Turcz.!), in China (Staunt.!). *B. macrosperma* Fisch. ! hort. Gorenk. *B. multifida* Desf. ! cat. h. Par. ed. 3. p. 186. *B. pauciflora* Poir. suppl. 1. p. 630. *B. Messerschmidii* Turcz. ! in litt. *B. Chinensis* herb. acad. Petrop. sed vix Willd. Flosc. 3-6 quadridentati. Achenia inæqualia centralia majora, 8-9 lin. (aristis excl.) longa. (v. v. c. et s. sp.)
58. *B. DECOMPOSITA* (Wall. ! cat. et herb. n. 298), caule glabro tetragono ramoso, foliis bipinnatipartitis, lacinii lanceolatis basi cuneatis mucronatis utrinquè scabridis, capitulis longè pedunculatis laxè subpaniculatis discoidis, invol. biseriali demum reflexo, acheniis elongatis biaristatis, aliis brevibus strigosocaberrimis, aliis basi glabris apice subtrigosis in eodem capitulo. — in India orient. ad Monghir prov. Bahar et in Nepalia. *Coreopsis corymbifolia* Hamilt. ! ex Wall. (v. s. comm. ab hon. aulâ merc. angl. Ind. or.)
59. *B. GRANDIFLORA* (Balb. cat. h. taur. 1812. p. 19), caule suberecto tereti glabro, foliis glabris impari-pinnatisectis, segmentis 3-5-jugis, aliis oblongis serratis aut subincisis, aliis in lobulos lineares pinnatipartitis, capitulis pedicellatis radiatis, invol. squamis ciliatis disco subbrevioribus, ext. patentibus, acheniis lineari-subtetragonis glabris 2-4-aristatis. (1) in Mexico circa urbem (Berl. ! exs. n. 1168), DC. cat. h. monsp. p. 83. Horn. h. hafn. 2. p. 788. *Bidenis odorata* Dum. Coara. bot. cult. 4. p. 241 non Cav. *Coreopsis diversifolia* Jacq. eclog. 1. p. 80. t. 54. opt. *Bidenis diversifolia* hortul. B. serratulata Desf. ! cat. h. par. 1829. p. 186. *Kerneria serrulata* Cass. dict. 51. p. 474. *Cosmea lutea* Sims bot. mag. t. 1689. — Ab hac ex descr. videtur diversa *Coreopsis ser-*

- rulata Poir. suppl. 3. p. 352 etiam à cl. Desf. admissa. Caulis sæpè basi radicans. Ligulæ 9-10 lin. longæ, 3-3½ latæ, 9-10 nervulis striatæ. Variat invol. glabro et extus plus minus hirsuto. (v. v. c. et s. sp.)
60. *B. ODORATA* (Cav. ic. x. p. 9. t. 13), glabra erecta, caule tetragono ramosissimo, foliis bipinnatisectis lobis cuneato-trifidis, capitulis pedunculatis solitariis radiatis, invol. squamis lanceolatis apice ciliatis, exter. subreflexis, achæniis lineari-subtetragonis ad angulos scabris biaristatis, lateralibus brevioribus. ① in Mexico (Cav.) fortè etiam in insulis Societatis ex Hook. et Arn. bot. Beech. p. 66? *Coreopsis odorata* Lam. ill. t. 704. f. 1. Accedit ad *B. grandifloram*, sed radius albus, flores odorati, achænia scabra, et fol. lobi cuneati.
61. *B. FERULÆFOLIA*, glabra, caule herbaceo teretiusculo, foliis oppositis bipinnatisectis, lobis linearibus vix sublancoelatis integerrimis costam segmentorum latitudine non superantibus, capitulis pedicellatis laxè corymbosis radiatis, invol. squamis linearibus demùm patulis, int. ovali-lanceolatis margine membranaceis, ligulis obovatis obtusis multinerviis invol. quadruplò longioribus, achæniis linearibus glabrisculis biaristatis. ② in Mexico (Willd.), *Coreopsis ferulæfolia* Jacq. h. Schænb. 3. p. 65. t. 375. bot. mag. t. 2059. Quid sit var. odoratissima à Persoonio citata ignoro. *Kerneria ferulæfolia* Cass. dict. 5r. p. 43. Aristæ certè retrorsum scabræ in ipso specim. à Jacquinio filio comm. et apprimè iconi conveniente. An ideò aliam plantam sub oculis habuit cl. Lessing cum inter *Coreopsides* veras admisit? (v. s. et v. c.)
62. *B. PROCERA* (Don in bot. reg. t. 684), glabra striata ramosa, foliis decursivè 2-3-pinnatisectis, lobis linearibus acutis canaliculatis integris flaccidis, capitulis subcorymbosis pedicellatis amplè radiatis, invol. squamis ext. linearibus obtusis pubescentibus, achæniis glabris compresso-cuneiformibus biaristatis; ② in Mexico. *Coreopsis angustifolia* Pavon in h. Lamb. ex Don. Ligulæ flavæ 10 lin. longæ. Ex habitu et lobis foliorum integris ad Cosmum pertinere videtur, sed achænium (junius equidè scriptum) potiùs *Bidentis*.
63. *B. FœNICULIFOLIA*, caule erecto acutè 4-6-angulato glabrescens, ramis acutè tetragonis subpuberulis, foliis oppositis bi- aut tripinnatisectis glabrisculis, lobis linearibus acutis integerrimis aut hinc indè lobulatis, capitulis sæpiùs ternis pedicellatis subcorymbosis radiatis, invol. squamis linearibus ciliatis, inter. ovato-lanceolatis margine scariosis longit. æquantibus, ligulis obovatis obtusis multinerviis inv. triplò longioribus, achæniis lineari-tetragonis glabris vix apice scabridis biaristatis. — in Mexico ad occid. urbis Guanaxuato in prov. Leoninà leger. et mecum comm. cl. Mendez et Alaman. *Coreopsis fœniculacea* ic. fl. mex. ined. Achænia 3-3½ lin. longa. Flores intensè lutei. (v. s.)
64. *B. CYNAPIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 235), glabra, caule erecto subtetragono, ramis floridis divaricatis, foliis irregulariter tripinnatisectis, lobis oblongis basi cuneatis, terminali longè acuminato, capitulis longè pedunculatis breviter radiatis, invol. squamis linearibus ciliatis disco subæqualibus, exter. reflexis, achæniis lineari-tetragonis 4-aristatis, aliis elongatis glabris, aliis brevioribus pilosulis. ① in Insulâ Cubâ prope Guanaxacoa (H. et B.), circa Havanam (La Sagra!). Ligulæ parvæ angustæ, in sicco pallidæ. (v. s.)
65. *B. BIPINNATA* (Linn. sp. 1166), glabra, caule erecto obtusè tetragono, foliis bipinnatisectis, lobis lanceolatis, capitulis pedicellatis radiatis, ligulis paucis (2-4) vix disco longioribus, invol. squamis subinæqualibus disci circ. longitudine, ext. reflexis, achæniis lineari-tetragonis glabris 3-4-aristatis. ① in sicca agris Amer. bor. frequens; nunc in Senegaliâ (Pers.!), et in agrum Tyrolense quasi spontè introducta (Koch!). Ell. sketch 2. p. 432. — Herm. parad. t. 123. Moris. oxon. s. 6. t. 7. f. 23. Ligulæ flavæ, primâ fronte vix perspicuæ. (v. s. sp. et v. c.)
66. *B. CHILENSIS*, glabra, caule erecto tetragono sulcato vix ramoso, foliis subciliatis petiolatis biternatisectis seu pinnatisectis, segmentis bijugis cum impari tripartitis, lobis obovatis basi cuneatis acutis grossè serratis dentibus mucron-

tis, capitulis paucis longè pedicellatis radiatis, invol. basi barbati squamis ext. linearibus, inter. lanceolatis, ligulis invol. duplò longioribus, achæniis lineari-tetragonis apice vix scabridis biaristatis. (1) in Chili circa Quillota in pascuis apricis collium (Bert.), ad Valparaiso (Gaud.), et circa Limam, si Bidens Feuill. obs. 2. t. 33 huc ritè relata. B. bipinnata Hook. et Arn. bot. Beech. 1. p. 30. B. Valparadisiaca Colla. mem. ac Tur. 38. p. 12. t. 24. In nostris specim. ligulæ flavæ, sed ex Bert. variant albæ et crocæ. An satis a B. bipinnatâ differt? (v. s.)

β. *apiifolia*, caule glabro subsimplici striato, foliis trisectis, segmentis latè ovatis obtusis dentato-serratis suprâ sparse et adpressè pilosis capitulis pedicellatis solitariis radiatis, invol. basi pilosi squamis linearibus disco brevioribus achæniis biaristatis. (1) in Chili legit cl. Poeppig. Bidens leucantha Poepp. ! pl. exs. n. 207. Herba 6-8 pollic. Ligulæ flavæ, fortè var. albæ ex nomine Poepp., disco ferè duplò longiores. (v. s. comm. à cl. Poeppig.)

67. B. CAUCALIDÆA, caule adscendente glabriusculo subtetragono, foliis subpubescentibus petiolatis pinnatisectis, segmentis uni-aut bijugis cum impari pinnatipartitis, lobulis oblongis acutiusculis, pedunculis pubescenti-hirtellis subcorymbosis, capitulis radiatis, invol. squamis disco brevioribus inter se æqualibus erectis glabris, achæniis lineari-tetragonis scabriusculis biaristatis. (1) in Mexici valle Toluccanâ legit cl. Berlandier (pl. exs. n. 1138). Ligulæ flavæ pluristriatæ disco paulò longiores ovales 3-4 lin. longæ. Aristæ fruct. nunc breves tennes nunc longiores rigidæ. Proximè habitu accedit ad Cosmeam chrysanthemoidem Kunth. t. 482 sed poly- nec 1-cephala, pedunculi villosi nec 1-cephali, pedunculi villosi nec glabri, ligulæ minores flavæ nec violaceæ, achæmium scabrum nec glabrum. (v. s.)

Sect. III. DISCOPODA.

Achænia lineari-subtetragona, utrinquè attenuata, infernè in areolam orbiculatam obliquam dilatata. Aristæ patulæ. Capitula radiata, ligulis et radio purpureis. — An Cosmi affinium?

68. B. PURPUREA, caule adscendente erecto basi subtetragono hispido, ramis oppositis elongatis subaphyllis glabriusculis teretibus, foliis sparse hirsutis hispidis pinnatisectis, segmentis 3-4-jugis, lateralibus ovato-lanceolatis hinc indè incis, terminali basi cuneato trilobo inciso, omnibus ciliatis callosomucronatis, capitulis longè pedicellatis subcorymbosis radiatis, involucri squamis glabris subacutis squarrosis, achæniis lineari-subtetragonis lævibus biaristatis. — in Mexici Cordillerâ Gnachilaquensi legit Berlandier (pl. exs. n. 1007). Cor. disci et radii atro-purpureæ. Ligulæ 7 lin. longæ, 2 ½ lin. latæ. Fructus juniores aristis erectis. Coreopsis purpurea ic. fl. mex. ined. sed aristæ 3. (v. s.)

β. *glabriuscula*, caule foliisque glabriusculis, lobis foliorum longioribus sublanceolatis. — cum var. α legit Berlandier (pl. exs. n. 1021). Ligulæ in specim. magis provento desiderantur. Achænia matura lineari-subtetragona utrinquè subattenuata glaberrima; aristæ rigidæ retrorsum scabræ ad peripheriam capituli deflexæ. (v. s.)

+ Species non satis notæ.

69. B. GRAVEOLENS (Mart. in Isis 1824. p. 590, et in linnæa litt. 1830. p. 58), caule...., foliis oblongo-lanceolatis crenato-serratis basi cuneatis integerrimis reticulato-venosis, capitulis longè pedunculatis subpaniculatis. — in Brasiliâ.
70. B. MEGAPOTAMICA (Spreng. syst. 3. p. 454), foliis omnibus bipinnatifidis lineari-filiformibus glabris, capitulis subgeminis pedunculatis erectis discoideis, invol. colorato disco æquante. — ad Rio-Grande Brasiliæ. Achænia indeser.
71. B. PANICULATA (Hook. et Arn. bot. Beech. p. 66), suffruticosa, foliis longè

- petiolatis lato-lauceolatis acuminatis glabris serratis acuminis integerrimo, paniculâ patente terminali, capitulis parvis parçè radiatis, involucri brevi, achæniis dentibusque 2 brevibus hispidis. ζ in insulâ Otahiti. — Mixtum cum succo nucis Cocœ et ebullitione coctum est catharticum, *motu dictum*.
72. B. TENELLA (Linn. amæn. ac. afr. 47), foliis linearibus, pedunculis capillaribus, invol. sub-4-phyllis, achæniis 5 erectis. $\textcircled{1}$ ad Cap. Bonæ spei. \dagger .
73. B.? CHRYSANTHA, foliis ternatisectis segmentis ovato-oblongis serratis, capitulis longè pedunculatis subcorymbosis, radio concolore. — in insulis Caribæis. Coreopsis chrysantha Linn. sp. 1282. — Plum. ed Burm. t. 53. f. 1. Affinis videtur B. scandenti.
74. B.? LEUCORHIZA, caule herbaceo erecto tetragono, foliis quinis lanceolatis serrato-incisis, pedunculis oligocephalis, ligulis 6 ovatis integerrimis achæniis 3-cornutis. — prope Cantonem Sinarum. Coreopsis leucorhiza Lour. fl. coch. 622. Radix perpendicularis crassa alba. Capitula integrè crocea. Ligulæ neutræ. Aristæ setis inversis ramosæ, undè ad Bidentem potius pertinere videtur.
75. B. ALBA, caule debili flexuoso, foliis oppositis petiolatis trisectis rariùs 5-sectis, segmentis obovatis serratis, capitulis pedicellatis radiatis, ligulis 8 obovatis, achæniis linearibus biaristatis. — in insulâ Sanctæ-Crucis. Coreopsis alba Linn. sp. 1282. — Herm. parad. 124. t. 124. Pluk. alm. t. 160. f. 3. Ligulæ albæ. Aristæ more Bidentium ex descr. Herm. scabræ. Affinis videtur B. Portoricensis et fortè hujus varietas radiatæ. — Syn. Brownei à Linnæo citatum videtur alienum ex recept. nudo dicto.
76. B.? VERTICILLATA (Linn. sp. 1167), caulibus procumbentibus subsimplicibus, foliis inf. alternis, longè sup. oppositis breviter petiolatis lanceolato-ovatis subintegris suprà glabris subtùs tomentoso-albidis, capitulis in axillis sup. binis discoideis, achænio tenui setis 2 patulis coronato. — in Verâ-Cruce. Bidens Linn. hort. Cliff. 399. n. 4.

†† Species exclusæ.

- B. ACNELLA Lam. = *Spilanthes acmella*.
 B. ACNELLOIDES Big. = *Spilanthes oleraceus*.
 B. ANGUSTIFOLIA Lam. = *Spilanthes cereus*.
 B. APIIFOLIA Linn. = *Chrysanthellum procumbens*.
 B. ARBOREA Roxb. = *Petrobium arboreum*.
 B. ATRIPLICIFOLIA Linn. = *Isocarpha atriplicifolia*.
 B. AUSTRALIS Spreng. = *Campylothea micrantha* β .
 B. BARDANA Wall. = *Chrysanthellum Indicum*.
 B. BERBERIANA Spreng. = *Cosmos caudatus*.
 B. CROCATI Cav. = *Verbesina crocata*.
 B. FERVIDA Lam. = *Spilanthes oleracea*.
 B. FRUTESCENS Mill. = *Salmea scandens*.
 B. FUSCA Lam. = *Spilanthes oleracea fusca*.
 B. HIRSUTA Sw. = *Salmea hirsuta*.
 B. INSIPIDA Steud. = *Spilanthes insipida*.
 B. MICRANTHA Gaudich. = *Campylothea micrantha*.
 B. NIVEA Linn. = *Melanthera hastata*.
 B. NIVEA Sw. = *Melanthera deltoidea*.
 B. OCTYMFOLIA Lam. = *Spilanthes alba*.
 B. PINNATIFIDA Wall. = *Glossogyne pinnatifida*.
 B. PINNATIFIDA Forst. = *Campylothea micrantha* β .
 B. RAMTILLA Wall. = *Guizotia oleifera*.

B. SCANDENS Br. = *Salmea scandens*.

B. SCANDENS Sw. = *Salmea eupatoria*.

B. TENUIFOLIA Labill. = *Glossogyne tenuifolia*.

CCCL. COSMOS Cav. ic. 1. (1791). p. 9. t. 14 et 79. Cass. dict. 11. p. 4. — *Cosmea Willd. sp. 3.* (1803). p. 2250. *Hoffmans. verz.* 1824. p. 206. *Less. syn.* 230. — *Cosmos Pers. ench. 2. n.* 1917. — *Georgia Spreng.*

Capitulum multiflorum radiatum, ligulis neutris. Invol. duplex, utrumque squamis 8-10 basi plus minus concretis apice acuminatis. Recept. planum paleaceum, paleis membranaceis in filum elongatum productis. Styli rami apice incrassati hispidi in conum subulatum producti. Antheræ apice appendice scariosa cordatâ superati. Achænium tetragonum exalatum rostratum et interdum stipitatum, aristis 2-4 retrorsum piloso-scabris deciduis coronatum. — Herbæ Americanæ annuæ glabræ aut vix pilosulæ elatæ ramosæ. Folia bipinnatisecta, lobis linearibus lanceolatis sæpius margine integerrimis. Capitula ad apices ramorum longè nudorum solitaria. Discus intensè luteus. Radius versicolor.

1. C. BIPINNATUS (Cav. ic. 1. p. 9. t. 14), glaberrimus, foliis bipinnatisectis, lobis linearibus remotis integerrimis, invol. ext. squamis ovato-lanceolatis acuminatis, achæniis glabris breviter rostratis triaristatis. (1) in Mexico (Cav.), Georgia bipinnata Spreng. syst. 3. p. 611. *Coreopsis formosa Bonato in diss. Pisaur. p. 21. t. 2.* Ligulæ 7-9, amplæ, 7-8 lin. longæ, purpureæ rosæ seu albidæ nunquàm flavæ.

β. *exaristatus*, aristis ad apicem achæniis nullis. (1) in Mexico (Alam.) ad Pezcuaro (H. et B.), Oaxaca (Andr. ! pl. exs. n. 307). S. Angel. (Berl. pl. exs. n. 938). Hæc var. sola in hortis mihi notis nunc colitur et sola provenit in herb. Mexic. (v. s. sp. et v. c.)

2. C. PARVIFLORUS (H. B. et Kunth nov. gen. am. 4. p. 241), glaberrimus, foliis bipinnatipartitis, lobis capillaceo-linearibus integerrimis, invol. squamis ext. 8 lanceolatis acutis brevioribus, achæniis glabris ad angulos scabridis breviter rostratis 3-aristatis. (1) in Mexico (Jacq.) inter Puerto-de-Varientos et S.-Rosa de la Sierra (H. et B.). *Coreopsis parviflora Willd. sp. 3. p. 2251.* Ligulæ rosæ aut albæ, 3-dentatæ, 3 lin. longæ. (v. s. et v. c.)

3. C. TENELLUS (H. B. et Kunth nov. gen. am. 4. p. 240), foliis bipinnatipartitis glabris, lobis lanceolatis acutis, inferioribus basi subserratis, terminalibus majoribus, invol. squamis ext. lineari-spathulatis brevioribus, achæniis subscabris.... aristis.... (1) in frigidis Mexici inter urbem et Huehuetoca. Ligulæ rosæ?

4. C. CAUDATUS (H. B. et Kunth nov. gen. am. 4. p. 240), caule glabro, foliis bipinnatipartitis, lobis lanceolatis cuspidatis margine scabris, petiolis ciliatis, invol. squamis ext. lineari-lanceolatis acutis ciliatis patulis vix basi concretis interiores ferè æquantibus, achæniis glabrisculis apice scabris longissimè rostratis, aristis 2 patulis. 2/ in insulâ Cubâ juxta Havanam (H. et B.), Porto-Rico (Bert. ! Wydl. !), Sancti-Thomasii agris saccharo cultis (Less. in linnæâ 1821. p. 727), Martinicâ (herb. Duby !), etiâ in Indiâ or. (ex h. Wight) et insulâ Mauritii (Bout. !), sed introduct. *Bidens Berteriana Spr. ! syst. 3. p. 454.* DC. in Wight contrib. p. 19. Ligulæ rosæ apice trifidæ. Achænia ferè pollicem longa. Flores disci 12 ex Kunth, circ. 20 in sp. meis. (v. s. sp.)

5. C. SULFUREUS (Cav. ic. 1. p. 56. t. 79), caule piloso, foliis bipinnatipartitis, lobis lanceolatis mucronatis margine subscabridis, petiolis ciliatis, invol. squamis ext. linearibus adpressis quàm interiores adhuc brevioribus et multo angustioribus, achæniis glabrisculis apice scabris longè rostratis, aristis 2 patulis caducissimis. 2/ in Mexico (Cav.) circa Oaxacam (Andr. ! pl. ex. n. 308) ad Willalipando (Mendez !), *Cosmea sulfurea Willd. sp. 3. p. 2250.* *Coreopsis artemisiæfolia Jacq. ic. 3. t. 595.* Ligulæ sulfureæ. Cor. disci suberosæ, lobis supernè puberis. Achænium 9-10 lin. longum. (v. s. sp.)

6. *C. CHRYSANTHEMOIDES* (H. B. et Kunth nov. gen. am. 4. p. 239. t. 382), glaber, foliis pinnato-quinque-partitis, lobis obovato-cuneatis bi- aut trilobulatis, petiolis ciliatis, invol. squamis exter. lanceolato-linearibus paulò brevioribus, achæniis glabris biaristatis. \mathcal{N} in Mexico. *Cosmea chrysanthemifolia* Spreng. syst. 3. p. 615. Ligulæ violaceæ 8 lin. longæ. Achænium, ex ovario, videtur compressum subcuneatum, potiùs Bidentis quàm Cosmi.
7. *C. PILOSUS* (H. B. et Kunth nov. gen. am. 4. p. 241), caule foliisque hispidopilosis pinnatipartitis, lobis bijugis oblongo-ovatis serratis, terminali paulò majore, invol. squamis exter. linearibus obtusis brevioribus, achæniis glabris biaristatis. \mathcal{N} in aridis Mexici prope Santa-Rosa de la Sierra. Ligulæ rosæ ellipticæ 3-dentatæ 5-6 lin. longæ. Achænium junius obcompressum ut in Bidente.
8. *C. CRITHMIFOLIUS* (H. B. et Kunth nov. gen. am. 4. p. 242), glaber, foliis pinnatipartitis, lobis angustè linearibus integerrimis rigidis, infer. rariùs bifidis, invol. squamis exter. lanceolatis acutis brevioribus, achæniis... (I) ? prope Valladolid Mexicanorum (H. et B.). Ligulæ rosæ? — Hic fortè pertinet n. 1067 Berl.! pl. exs. circa Toluccam lectus. Flores in specim. desiderantur. Achænia tetragona rostrata apice subscabra 6 lin. longa, aristis 2-3-erectis superata. (v. s.)
9. *C. SCABIOSOIDES* (H. B. et Kunth nov. gen. am. 4. p. 242), caule pubescente, foliis pinnatipartitis subtus hispidulis, lobis bijugis lanceolato-oblongis acutis basi cuneatis apice suberratis, terminali majore, invol. squamis exter. lanceolatis acutis brevioribus, achæniis glabris 3-aristatis. (I) ? in Mexici montanis Mechoacanensibus prope Pezcuaro (H. et B.) et ad Tlapujahua (Keerl. in h. Mart.). *Cosmea scabiosæ* Spreng. syst. 3. p. 615. Ligulæ violaceo-purpureæ latæ obtusæ semipollicares. (v. s. in h. Mart.)

CCCLI. ADENOLEPIS Less. in *linnæa* 6. 1831. p. 510. t. 6. f. 6-10. syn. 230.

Capitulum pluriflorum heterogamum, fl. radii circ. 4 ligulatis neutris ligulâ subellipticâ emarginatâ, disci tubulosis hermaphr. 5-dentatis, limbo basi inflato dentibus revolutis. Recept. planum bracteolatum. Invol. cylindraceum disco æquale, squamis biserialibus linearibus, inter. foliaceis glandulâ magnâ terminatis. Anth. exsertæ ecaudatæ. Achænia conformia subtriangulato-obcompressa exalata, breviter rostrata seu apice attenuata omnino calva, in disco sæpè linearia sterilia. — Herba annua Sandwicensis gracilis parva apice puberula. Folia opposita petiolata ternatisecta aut superiora sessilia triloba. Capitula parva lutea terminalia solitaria.

1. *A. PULCHRELLA* (Less. ll. cc.). (I) in O-Wahu Sandwicensium detexit cl. de Chamisso.

Div. v. VERBESINÆ Less. syn. 230. — Helianthæ Spilanthæ et Verbesinæ pleræque *Cass. opusc.* 3. p. 58.

Capitula heterogama, fl. radii ligulatis fœmineis, rariùs homogama discoidea. Achænia cortice tenui donata, fere semper compressa aut obcompressa, rariùs 3-sulcata seu trigona, nunc calva, nunc sæpiùs in aristas rigidas tot quot achæniis anguli producta, rarissimè aut pappo vaginæformi aut aetis fragilibus rigidis superata,

CCCLII. LASIANTHÆA. — Lasianthus Zucc. in *h. Acad. Monac. non DC. nec Jack.*

Capitulum multiflorum radiatum, fl. radii 1-serialibus ligulatis fœmineis, disci longè tubulosis hermaphr., lobis 5 supernè pube crassâ hirsutissimis. Invol. campanulatum, squamis pluriserialibus imbricatis

ovatis obtusis, exter. ciliatis. Recept. planum, paleis membranaceis complicatis. Styli fl. herm. ramis apice summo barbatis exappendiculatis. Antheræ appendice lineari-lanceolatâ superatæ. Achænia glabra, radii triquetra trialata 3-aristata, disci compressa bialata et triaristata, aristis radii et disci basi alato-dilatatis. — Frutex Mexicanus di- seu trichotomus. Rami teretes pubero-asperi. Folia opposita breviter petiolata ovata sublanceolata, vix subdentata, suprâ sparsè strigilloso-pubera, subtus subvelutino-villosa reticulato-nervosa. Capitula circ. 5 ad apices ramorum breviter pedicellata umbellata crassa. Flores lutei. — Nomen paululum mutavi ne cum Lasiantho (sect. Gordoniæ) confundatur.

1. *L. HELIANTHOIDES* (Zucc. l. c.). 5 in temperationibus intra Saltepec et Cuito Mexicanorum legit septembr. flor. cl. de Karvinski. (v. s. in h. Acad. reg. Monac.)

CCCLIII. *PERYMENIUM* Schrad. ind. sem. hort. Gætt. 1830. et in litt. descr. et ic. — *Schistocarpa* Less. in linnæâ 1831. p. 387. syn. 387.

Capitulum multiflorum radiatum, ligulis 1-serialibus foemineis, fl. disci hermaphr. 5-dentatis. Invol. ovati aut campanulati squamæ imbricatæ ovatæ obtusæ aut subacutæ. Recept. planiusculum, paleis membranaceis complicatis. Stylus disci ramis apice breviter appendiculatis seu cono terminatis. Achænia subconformia subcompressa aut marginalia obcompressa mutica aut apice submarginata exaristata. Pappus setis rigidulis fragilibus 8-10 uniserialibus inæqualibus constans. — Suffrutices frutesce Mexicani setulis adpressis scabridi. Folia opposita plûs minûs petiolata seu triplinervia ovata aut oblonga dentata. Capitula pedicellata solitaria aut corymbosa. Flores croceo-flavi. — Genus *Verbesinæ* et *Lipochætæ* affine.

1. *P. DISCOLOR* (Schrad. l. hort. Gætt. ind. sem. 1830 et in litt. ic. et descr.), fruticosum, ramis cinereo-puberis, foliis oppositis subsessilibus oblongis vix acutis triplinerviis integerrimis margine subrevolutis suprâ et subtus in nervis adpressè villosis subtus inter nervos canescenti-velutinis, corymbis compositis trichotomis polycephalis, invol. ovatis squamis ovalibus obtusis ciliatis, ligulis ovalibus disco paulò longioribus. 5 in Mexico (Schrad.), in ditione Oaxacanâ legit cl. Andrieux (pl. exs. n. 305!). *Schistocarpa bicolor* Less. in linnæâ 1831. p. 409. Folia ferè Rosmarini. Flores croceo-flavi. Ligulæ sæpè appendice lineari auctæ subbilabiata. (v. s.)
2. *P. VERBESINOIDES*, fruticulosum, ramis striatis scabridis, foliis oppositis petiolatis ovatis basi vix acutis apice acuminatis serratis triplinerviis suprâ scabris subtus molliter pubescenti-velutinis, capitulis 7-9 pedicellatis subcorymbosis, invol. ovatis squamis ovatis acuminatis dorso villosis, ligulis 9-10 oblongo-linearibus. 5 in Mexico. Valdè simile *Verbesinæ* serratæ sed pappo et involuero certè distinctum (v. s. ex hort. Mexic. comm. à cl. Berlandier sub n. 565.)
3. *P. BERLANDIERII*, fruticulosum, ramis striatis vix subasperis, foliis oppositis brevè petiolatis ovato-lanceolatis triplinerviis remotè et parvè serratis suprâ sparsè subtusque in nervis minutissimè strigillosis, capitulis 7-9 apice in axillis supremis corymbosis breviter pedicellatis, invol. ovatis squamis ovatis subobtusis extus tenuè pubescentibus, ligulis 8-10 oblongo-linearibus. 5 in Mexici montibus Guchilapensibus legit cl. Berlandier (pl. exs. n. 1040). *Medinam* inter *P. verbesinoides* et *P. Cervantesii*, sed ab utroque distinctum. (v. s.)
4. *P. MENDEZII*, fruticosum, ramis teretibus, junioribus pedicellisque adpressè pubescentibus, foliis oppositis brevè petiolatis ovato-lanceolatis triplinerviis subintegerrimis utrinquè puberulo-scabris, capitulis pedicellatis ad apices ramorum subumbellatis, invol. squamis ovatis obtusis infer. extus pubescen-

- tibus, ligulis 3-4 oblongis. ζ in Mexico circa Villalpandæ legit cl. Mendez. Medium inter P. Berlandierii et Cervantesii. Pedicelli vix semipollicem longi. Styli rami breves cono crasso hirsutissimo terminati. Achænia matura regulariter punctis rufis maculata. Pappus 8-10-setus 1-2 subaristæformibus. (v. s.)
5. P. CERVANTESII, suffruticosum, ramis adpressè piloso-scabris, foliis oppositis ovatis acutis basi obtusiusculis brevissimè petiolatis hinc indè serratis triplinerviis suprâ piloso-scabris subtus pubescenti-hirtellis, capitulis 3-5 pedicellatis corymbosis, invol. campanulati squamis ovatis paucis dorso substrigillosis obtusis, intimis membranaceis oblongis, ligulis 5-10 ellipticis. \mathcal{U} in Mexico (Alam.), circa urbem in montanis (Berl. pl. exs. n. 634). Verbesina Mexicana Cerv. ined. ex Alam. Variat pedicellis 3 ferè bipollicaribus, aut 5 dimidiò et ultrâ brevioribus. (v. s.)
6. P. BUPHTHALMOIDES, suffruticulosum, ramis adpressè puberulis, foliis oppositis subsessilibus ellipticis-lanceolatis acuminatis basi 3-5-nerviis grossè serratis suprâ sparsè subtus præcipuè in nervis puberis, pedunculis terminalibus elongatis subsolitariis, invol. campanulati squamis oblongo-lanceolatis subacutis dorso adpressè puberis, ligulis 7-10 oblongis. ζ in Mexico legit cl. Alaman. Habitu refert Verbesinam buphthalmoidem Link (abb. t. 49), sed facillè differt. Pappus brevior quàm priorum. (v. s.)
7. P. BARCLAYANUM, suffruticosum subscondens, ramis adpressè villosis, foliis petiolatis ovatis acutis subdentatis trinerviis suprâ sparsè subtus secus nervos strigilloso-pilosis, pedicellis solitariis elongatis 1-cephalis, invol. campanulati squamis ext. latè ovatis obtusis strigillosis, ligulis 5-7 oblongis. ζ in Mexico ex hort. Barclay, ubi pro genere novo à cl. Don descriptum anno 1830 habebatur. Ligulæ croceo-flavæ 6-8 lin. longæ. Pappus 12-13-setus longior quàm priorum. (v. s. à Filio coll.)
8. P. GYMNOLOMOIDES, fruticosum scandens? ramis..... foliis....., invol. campanulati squamis ext. obtusissimis extus hirsutis, int. elliptico-obovatis, ligulis 5 disco paulò longioribus, tubo hispidiusculo, ovariis subalatis ciliatis. ζ in dumetis Misanthæ Mexicanorum. Lipotriche gymnomoloides Less. in linnæa 1831. p. 408. Simillimum dicitur Gymnomolæ microcephalæ.

CCCLIV. PSATHUROCHÆTA.

Capitulum multiflorum radiatum, ligulis 1-serialibus fœmineis, fl. disci hermaphrod. tubulosis 5-dentatis. Invol. 1-2-seriale, squamis lanceolatis disco brevioribus. Recept. convexum, paleis lanceolato-linearibus acuminatis. Cor. lobi puberuli. Stylus disci ramis cono elongato acuminato hispido terminatis. Achænia acuta tetragona subcompressa brevia, setis disci plurimis 8-10 fragilibus inæqualibus deceduis coronata, setis radii paucioribus brevioribus. — Herba Australi-Africana, Verbesinæ facie. Caulis teretiusculus sex-sulcatus, oppositè ramosus et ideò trichotomus, pube brevi scabrigo-canescens. Folia opposita petiolata utrinquè scabra trinervia sæpiùs triloba, lobo medio longiore et magis acuminato. Pedunculi ad apices ramorum alterni subterni 1-cephali nudi. Cor. intensè flavæ. — Genus affine Perymenio et Lipochætæ; à priore involucre 1-2-seriali, achæniis tetragonis, styli ramis longius acuminatis; à posteriore angulis fructûs in aristas non productis; ab utroque recept. convexo diversum. Nomen ex $\psi\alpha\theta\upsilon\rho\varsigma$ fragilis, et $\chi\alpha\iota\tau\alpha$ seta.

1. P. DREGEI. \mathcal{U} ? in Africâ australi ad Omsamcoubo, ferè ad maris altitudinem, detexit cl. Drege. An varietates, an fortè species duæ distinguendæ? nempe:

a. latifolia, foliis submembranaceis basi latioribus trilobis, lobis dentatis

lateralibus sæpe sinuato-bilobis, nervulis subtus amplè areolatis vix prominulis. (v. s.)

β. *reticulata*, foliis rigidioribus, lobis serratis, lateralibus brevibus subdivisis, nervulis subtus crebrè reticulatis elevatis. (v. s.)

CCCLV. LIPOCHÆTA. — *Lipotriche* Less. syn. 231. et in *linnaea* 1831. p. 310 non Brown. — *Zexmenia* Lallav. et Lex. nov. veg. 1. p. 13? — *Verbesinæ* sp. Gaud.

Capitulum multiflorum radiatum, fl. radii 1-serialibus fœmineis, disci hermaphr. 5-dentatis. Invol. ovati aut campanulati squamæ 2-5-seriales ovales adpressæ. Recept. planiusculum, paleis membranaceis complicatis. Stylus disci ramis apice appendiculatis. Achænia radii triquetra vix subalata, angulis sæpius ternis in aristam persistentem productis, dentibus setiformibus paucis interjectis; disci compressi 1-2-aristata ad latus interius subalata. — Suffrutices aut herbæ. Folia opposita breviter petiolata aut sessilia ovato-lanceolata subserrata triplinervia. Capitula pedicellata solitaria aut corymbosa. Flores lutei. — Habitus Verbesinarum oppositifoliarum.

* *Species Americanae.*

1. *L. MONOCEPHALA*, caule fruticoso tereti glabro, foliis petiolatis ovato-lanceolatis serratis triplinerviis utrinquè scabris, petiolis intus villosissimis, pedunculis ex axillis ramorum solitariis 1-cephalis, invol. squamis biserialis paucis ovalibus submembranaceis, ligulis 4-5, achæniis cuneatis alatis, pappo inæqualiter pluriaristato. ♂ patr. ign. olim in horto Paris. culta sub nom. Verbesinæ fruticosa et inde fortè ex ins. Caribæis orta. Flores lutei. (v. s.)

2. *L. STRIGOSA*, suffruticosa, ramis teretibus glabriusculis, foliis petiolatis ovalibus acuminatissimis serrulatis suprâ rigide et sparsè strigosis subtus adpressè pubescenti-villosis, pedicellis terminalibus 1-cephalis, invol. squamis lanceolatis biserialis, ext. foliaceis strigosis, achæniis elongatis angulatis 2-3-aristatis et paleolas totidem inter aristas gerentibus. ♂ in Mexici ditione Oaxacani australi circa Tehuantepec legit cl. Andrieux (pl. exs. n. 3131). Fol. pagina super. pili basi tuberculosi. Flores non vidi et ex fructu tantum huc refero. (v. s.)

3. *L. UMBELLATA*, fruticulosa, ramis adpressè scabridis, foliis brevissimè petiolatis ovato-lanceolatis utrinquè subacutis minutè serrulatis suprâ brevè et sparsè strigillosis subtus secus ramulos hirtellis, nervis utrinquè 3 prope basin medii ortis, petiolis subciliatis, capitulis plurimis umbellatis, pedicellis capitulo triplò longioribus, invol. ovatis squamis ext. obtusissimis, int. acutis disco dimidiò ferè brevioribus. ♂ in Mexico prope Cuernavaca legit cl. Berlandier (pl. exs. n. 1065), et ad Acapulco (h. Hænk!). Achænia radii 3-gona inæqualiter 3-aristata, disci compressa longè biaristata. (v. s.)

β. *conferta*, capitulis numerosioribus confertis, pedicellis capitulo brevioribus, invol. disco paulò breviorè. ♂ ad Cuernavaca Mexicanorum legit cl. Berlandier pl. exs. n. 1053! (v. s.)

4. *L. FASCICULATA*, caule fruticoso glabro, ramulis vix apice puberulis, foliis petiolatis oblongo-lanceolatis utrinquè acuminatis subserratis suprâ asperis subtus puberulis, pedicellis axillaribus terminalibusque, infer. solitariis, super. fasciculatis capitulo discoideo sextuplò longioribus, invol. squamis ovalis acuminatis, ext. pubescentibus, int. longioribus glabris. ♂ in Mexico legit cl. Berlandier (pl. exs. n. 2134!). Frutex orgyalis. Pedicelli inter axillam et caulem sæpius 2-3-seriatim dispositi! Ligulæ nullæ. Achænia exter. triquetra, 3 aristata, inter. compressa biaristata, squamellis intermediis omnium minimis. Species subaffinis *L. umbellatæ*. (v. s.)

** *Species Sandwichenses.*

5. *L. LOBATA*, fruticulosa, ramis pubero-scabris, foliis breviter petiolatis ovato-lanceolatis grossè dentatis sæpè tri- aut rariùs 5-lobatis 3-5-plinerviis suprà aspero-scabris subtùs secus nervos strigillos inter eos adpressè puberis, capitulis 3-5 pedicellatis ad apices ramorum subumbellatis, invol. ext. squamis latè ovatis, ligulis 5-10 disco duplò longioribus. ζ in insul. Sandwich (Gaud. Hook.), nempè Woahoo sen O-Waha (Macr. Cham.!). *Verbesina lobata* Gaud. voy. Freyc. bot. p. 464. Hook. et Arn. bot. Beech. 87. *Lipotriche australis* Less. in linnæâ 183. p. 310 (v. s. coll. à cl. Macrae comm. ab hon. hortie. soc. Lond.)
6. *L. HASTULATA* (Hook. et Arn. in Beech. bot. 87. sub *Verbesinâ*), suffruticosa, foliis oppositis brevè petiolatis ovato-lanceolatis 3-nerviis versus basin utrinquè sursum lobulatis grossè serratis suprà scabris subtùs hispidis. ζ in insulis Sandwich. Affinis *L. connatæ* dicitur, sed ex phrasi vix à *L. lobatâ* differe videtur. \dagger .
7. *L. CONNATA* (Gaudich. in bot. Freyc. voy. 464. sub *Verbesinâ*), suffruticosa, foliis sessilibus connatis rhombeo-ovatis argutè et grossè duplicato-serratis suprà scabris subtùs densè hispidis. ζ in insulis Sandwich. Cæt. ign. (v. s. in h. Mus. Par. sed non car. carp. vidi.)
8. *L. LAVARUM* (Gaudich. in bot. Freyc. voy. 464. sub *Verbesinâ*), fruticulosa, foliis lanceolatis argutè serratis utrinquè strigoso-sericeis argenteis. ζ in montanis insularum Sandwich, ubi dicitur *Kokolao* (v. s. in h. Mus. Par. sed car. non vidi.)
9. *L. SUCCULENTA* (Hook. et Arn. in Beech. bot. 87. sub *Verbesinâ*), herbacea glabra nitida succulenta, foliis oblongo-ovatis apice obtusis mucronulatis basi in petiolum attenuatis suprà medium crenulato-serratis, pedunculis solitariis, invol. squamis orbiculatis. — in maritimis Sandwichianis inter rupes volcanicas ins. Onesheow ubi dicitur *Nehe* et *Nenehe*. *Achænia compressa* seu trigona, angulo uno acutissimo interruptè subalato, aristis 2-3 brevibus coronata.

† *Species vix nota.*

10. *L. ? SERRATA*, caule sarmentoso 12-pedali et ultrà, ramis oppositis teretibus pubescentibus, foliis oppositis sublanceolatis petiolatis serratis asperis, junioribus subtùs pubescentibus, inflorescentiâ terminali, squamis inv. 10-12 biserialibus ovatis apice revolutis, paleis apice dilatatis. — in San-José del Corral Mexici, ad vias. *Zexmenia serrata* Lallav. et obs. nov. veg. 1. p. 13.

CCCLVI. DIPLOTHRIX.

Capitulum multiflorum radiatum, ligulis circ. 5 ovatis fœmineis cum ovario continuis persistentibus, fl. disci tubulosis 5-dent. hermaphr. Invol. ovatum imbricatum, squamis ovatis. Recept. paleis complicatis flores involv. onustum. Styli rami cono superati. *Achænia* radii obcompressa (seu triquetra) calva seu denticulis minimis superata, disci compressa, exter. subcalva, centralia in aristas 2 producta. — Suffrutices Mexicani ramosissimi. Folia opposita conferta acerosa acuta integerrima vix subpuberula 1-nervia, marginibus subcallosis. Capitula ad apices ramorum solitaria breviter pedicellata. Ligulæ ex sicco albidæ. — Nomen (olim ab Oppiano Pino consecratum) alludit et ad habitum foliorum ad Coniferas accedentem et ad aristas binas florum centralium.

1. *D. ACEROSA*, pedicellis involuero et folio vix longioribus, achæniis radii obcompressis. ζ in Mexici ditione Sancti-Ludovici de Potosi decemb. flor. legit cl. Berlandier (pl. exs. n. 1343!). Folia tenuia 3 lin. longa. Capitula vix 3 lin. (v. s.)

2. *D. JUNIPERIFOLIA*, pedicellis capitulo et folio longioribus, achæniis radii triquetris. ꝯ cum priore legit cl. Berlandier (pl. exs. n. 1359!), sed ab eâ differt, foliis 6 lin. longis, ad margines callosio-nervosis, pedicello 6-12 lin. longo, capitulo duplò majore. In meo specim. nimis ætate provecto caract. florum non ritè vidi. (v. s.)

CCCLVII. *SELLOA* H. B. et Kunth nov. gen. am. (1820) 4. p. 265. t. 395. Cass. dict. 55. p. 273. Less. syn. 231 non Spreng. — *Fœea* Spreng. syst. 3. (1826) p. 581 non Bory (1824.)

Capitulum multiflorum radiatum, ligulis 1-serialibus 10-15 fœmineis et filam. antheris destituta gerentibus, fl. disci hermaphr. tubulosi 5-dentatis. Invol. 2-3-seriale, squamis circ. 10-15, ext. ovato-ellipticis, int. oblongis acutis. Recept. convexum, paleis lineari-setiformibus. Achænia cuneata pentagona lævia coronata setulis 3 (5?) hispidulis caducis tubo corollæ adpressis. — Herbæ Mexicanæ subscapiformes, caule nempè basi folioso apice longè denudato 1- rariùs 3-cephalo. Folia opposita petiolata ovato-elliptica subintegerrima triplinervia glabra. Capitulum ferè Senecionis elegantis. Ligulæ albæ extùs purpurascens 9-nerviæ apice trilobæ.

1. *S. PLANTAGINEA* (H. B. et Kunth nov. gen. 4. p. 265. t. 395), caule apice nudo, foliis ovali-oblongis subdentatis triplinerviis. ꝯ in montibus Mexicanis inter Agnasarco et Ario. Fœea plantaginea Spreng. syst. 3. p. 581.
2. ? *S. LINEARIS*, caule etiàm apice foliato, foliis spathulato-linearibus enerviis integerrimis. ꝯ in Monte-Video. Fœea linearis Spreng. syst. 3. p. 581. Species ex Less. syn. p. 231 omninò obscura. †.

† Species exclusæ.

- S. CAPENSIS* Spreng. = *Denekia Capensis*.
S. GLUTINOSA Spreng. = *Gymnosperma glutinosum*.
S. NUDATA Nutt. = *Gymnosperma nudatum*.

CCCLVIII. *VERBESINA* Less. syn. 231. — *Verbesinæ* et *Siegesbeckiæ* sp. Linn. — *Verbesina* et *Platypteris* Kunth. — *Verbesina*, *Platypteris* et *Hamulina* Cass. — *Verbesina* et *Locheria* Neck.

Capitulum multiflorum, sæpiùs radiatum ligulis fœmineis, rariùs homogamum discoideum. Invol. bi- aut pluriseriale, squamis subæqualibus aut imbricatis. Recept. planum aut subconvexum, paleis plùs minùs complicatis et flores amplectentibus. Stylus disci ramis apice appendiculatis. Achænia erostria plano-compressa ad angulos alata et in aristas binas rigidas producta. — Herbæ aut suffrutices. Species genuinæ omnes Americanæ. Folia opposita aut alterna, petiolata sessilia aut decurrentia dentata aut pinnatilobata. Capitula pedicellata solitaria aut corymbosa. Flores nunc omnes lutei, rariùs omnes albi, interdum in disco flavi in radio albi. — Genus idèò multiforme sed ex caract. ægrè in plurima divellendum.

Sect. I. *VERBESINARIA*. — *Verbesina* Cass. dict. 59. p. 142.

Capitula radiata, ligulis uniseriatis. Achæniarum aristæ rectæ sæpissimè binæ æquales, unâ interdum brevior aut deficiente.

§. 1. *Aptera, nempe foliis non decurrentibus et ideo caule non alato.** *Oppositifolia.*

1. *V. SERRATA* (Cav. ic. 3. p. 7. t. 214), caule herbaceo jun. tomentoso demum subsabro, foliis oppositis breviter petiolatis ovato-lanceolatis grossè serratis suprà scabris subtus tomentosis, petiolis villosis subconnatis, capitulis compositè corymbosis, ligulis paucis aut (ex Kunth) nullis. \mathcal{L} in Mexico (Cav.), in temperatis inter Temascatio et Guanaxuato (H. et B.), in insul. Cùbà (h. Lem.). Cass. dict. 59. p. 142. Folia summa subalterna interdum integerrima. Invol. squamæ biseriales oblongæ foliaceæ disco breviores. Ligulæ breves sæpè diffformes. Achænium maturum glabrum nigricans alà albà cinctum. Variat foliis super. alternis et capitulis discoideis. (v. v. c. et s. sp.)
2. *V. CANTHOPIFOLIA* (Willd. sp. 3. p. 2225), caule erecto, foliis oppositis brevè petiolatis ovatis obtusè serratis quintuplinerviis utrinquè pilis brevibus obsitis subtus scabris, pedunculis elongatis racemoso-paniculatis. — in Mexico prope Acapulco. Cat. ign. †.
3. *V. ATRIFOLIOLIA* (Juss. et Desf. hort. Par. ! Poir. ! dict. 8. p. 460), caule fruticoso tereti apice villosa-velutino, foliis oppositis petiolatis latè ovatis triplinerviis acuminatis dentatis indivisis aut trilobatis suprà scabridis subtus petiolisque molliter villosa-velutinis, capitulis subternis pedicellatis, invol. squamis lanceolatis acutis disco brevioribus, ligulis circ. 12 apice dentatis. \mathcal{L} patr. ign. Spreng. 3. p. 577. L'Hér. ! descr. ined. Colla ! hort. rip. p. 144. t. 31. Galinsoega discolor Spreng. ! nov. prov. 19. Clerodendron phlomoides hort. ital. ! Ligulæ fortè steriles nunc stylo donatæ nunc destitutæ, albæ. Discus flavus. Achænia ex statu immatura subvillosa biaristata? Hinc fortè species duæ confusæ, sed certè ambæ à *V. Phaetusa* diversissimæ. (v. s.)
4. *V. RUPERTHALMOIDES* (Link et Otto abb. p. 105. t. 49), caule decumbente suffrutescente supernè pubescente, foliis suboppositis lanceolatis acutis serrulatis in petiolum attenuatis pubescentibus, pedunculis elongatis solitariis 1-cephalis, invol. 2-3-serialis squamis lanceolatis subæqualibus, ligulis 10-12 tridentatis, achæniis pubescentibus exalatis biaristatis. \mathcal{L} \mathcal{L} in Monte-Video. Cor. flavæ.
5. *V. AUREA*, fruticosa, ramis teretibus adpressè canescenti-villosis, foliis oppositis sessilibus ovato-lanceolatis basi obtusis apice subacuminatis serratis utrinquè pubero-scabris, capitulis ad axillas super. solitariis et terminalibus, ideo ultimis ternis, invol. squamis ext. ovalibus obtusis dorso pubero-scabris, ligulis 7-9 involucri duplè longioribus, achæniis radii obcompressis biaristatis, disci triquetris inæqualiter aristatis. \mathcal{L} in Mexico. *Wedelia aurea* Hook. bot. mag. t. 3384. Flor. radii fœminei sed fortè tamen steriles. (v. s. c. in hort. Barcl.)

** *Alternifolia.*

6. *V. HELIANTHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 204 non Michx.), caule herbaceo sulcato hirta-piloso, foliis alternis subsessilibus lanceolato-oblongis acuminatis remotè denticulatis basi cuneato-angustatis suprà scabris subtus pubescenti-pilosis, corymbis ramosis, invol. semiglobosi squamis imbricatis oblongis obtusis subscariosis, ligulis circ. 15, paleis oblongo-lanceolatis, achæniis glabris angustissimè alatis 2-3-aristatis. \mathcal{L} in regno Quitensi? Cor. flavæ? *Verbesina Humboldtii* Spreng. syst. 3. p. 577.
7. *V. SORDESCENS*, rhizomate lignoso, caulibus suffruticosis teretibus pubero-scabris apice subhirsutis, foliis alternis sessilibus oblongo-lanceolatis basi obtusis apice acutis serratis utrinquè puberulis, capitulis ramulos terminantibus in corymbum dispositis, invol. squamis oblongis obtusis disco brevioribus, paleis vix acutis, achæniis glabris angustissimè alatis aut subexalatis longè biaristatis. \mathcal{L} \mathcal{L} in Brasiliæ prov. Rio-Grande verisim. in campis. Caulis circ. pedalis obscurè viridis. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bca. vol. n. 876.)

8. *V. ACUMINATA*, caule subherbaceo glabro striato, foliis alternis subpetiolatis oblongo-ellipticis utrinquè longè acuminatis integerrimis suprâ glabris subtus pallidioribus subpuberulis, corymbo composito ramosissimo polycephalo, capitulis oblongis pedicellatis, invol. squamis paucis obtusis, ligulis 1-2, achæniis puberis angustissimè alatis. \mathcal{L} \mathfrak{z} circa Caracas legit cl. Vargas. Folia 6 poll. longa, 15 lin. lata, in sicco marginibus subrevoluta. Ligulæ albæ aut pallidæ. (v. s.)
9. *V. ELEGANS* (H. B. et Kunth nov. gen. am. 4. p. 204), caule herbaceo ramoso hirtio-villoso, foliis alternis subsessilibus oblongis acutis argutè denticulatis hirtio-pilosis corymbis subsimplicibus, invol. squamis paleisque lanceolatis acutis, ligulis circ. 15, achæniis apice ciliato-hispidis. \mathcal{L} ? in regno Quitensi? Color flor. dubius.
10. *V. DENTATA* (H. B. et Kunth nov. gen. am. 4. p. 205), caule herbaceo ramoso pubescenti-canefescente, foliis alternis subsessilibus approximatis oblongis grossè dentatis basi cuneatis subcoriaceis suprâ hirtio-pilosis subtus cano-lanatis, corymbis subsimplicibus, invol. semiglobosi imbricati squamis ext. obtusis, inter. acutis, ligulis circ. 20, paleis acutis, achæniis glabris alatis ciliolatisque subbiaristatis, aristâ alterâ corollam æquante, alterâ brevissimâ. \mathcal{L} in calidis regni Quitensis. Cor. flavæ. Pallasia dentata H. et Bonpl. pl. equin. 2. p. 10. t. III.
11. *V. SALICIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 205), caule herbaceo glabro, ramis virgatis striatis apice pedunculisque pubescentibus, foliis alternis subsessilibus lineari-sublanceolatis utrinquè angustatis, corymbo ramoso ramis foliaceis superato, pedicello centrali brevi crasso, invol. squamis lineari-oblongis, ligulis 7-8, paleis vix apice subuncinatis. \mathcal{L} in Mexico (Mairet), circa Kalapam (H. B.). Valdè accedit ad *V. virgatam*. Flores pallidius flavi. Folia 3 poll. longa, 4-5 lin. lata. (v. s.)
12. *V. PERSICIFOLIA*, suffruticosa glabra, foliis elongato-lineari-lanceolatis distanter callosa-serratis utrinquè acuminatis breviter petiolatis penninerviis, corymbo terminali ramoso polycephalo, invol. squamis lanceolatis, ligulis paucis? parvis? recept. paleis lineari-cuneatis, achænio glabro latè alato obcordato ad latus interius 1-aristato, ad exterius mutico. \mathfrak{z} \mathcal{L} in Mexico inter Santander et Victoria legit cl. Berlandier (pl. exs. n. 2209). Capitula matura ferè globosa. Affinis *V. virgatæ*, sed folia nec auriculata nec decurrentia. Folia 5 poll. longa, 10-18 lin. lata. (v. s.)
13. *V. TOMENTOSA*, suffruticosa, ramis teretibus alternis apice cano-tomentosis, foliis alternis petiolatis ovato-lanceolatis sublobatis inæqualiter dentato-serratis suprâ araneoso-pubescentibus subtus petiolisque cano-tomentosis, corymbo composito confertè polycephalo ramoso, invol. squamis lineari-oblongis, ligulis..., achæniis glabris latè alatis emarginatis biaristatis. \mathfrak{z} in Amer. austr.? legit cl. Née, sed locus propr. ign. Cor. disci cylindricæ basi tomentosæ. (v. s. fructifer.)
14. *V. OAXACANA*, caule fruticoso apice summo pedicellis subpubescentibus, foliis alternis in petiolum attenuatis ellipticis utrinquè acuminatis serrulatis penninerviis suprâ glabris subtus molliter cano-sabtomentosis, corymbo subsimplici 5-8-cephalo, capitulis subsphæricis, invol. squamis biseriatis apice foliaceis subobtusis, ligulis plurimis brevibus obtusis, paleis submembranaceis acutiusculis, achæniis glabris. \mathfrak{z} in Mexici ditione Oaxacanâ prope Tehuantepec angust. flor. legit cl. Andrieux (pl. exs. n. 301) et fortè ad Acapulco ex h. Hænke! Flores lutei. (v. s.)
15. *V. SUBCORDATA*, fruticosa, ramis pube brevi confertâ velutinis, foliis alternis sessilibus oblongo-lanceolatis utrinquè attenuatis basi obtusè subcordatis grossè dentatis subrepandis suprâ scabris subtus velutinis, pedicellis velutino-tomentosis 1-cephalis subcorymbosis, invol. squarrosis, ligulis 10-12, achæniis pubescentibus angustatis vix marginatis biaristatis. \mathfrak{z} circa Buenos-Ayres legit cl. Bacle. Flores flavi. (v. s. comm. à cl. Iuv.)

§. 2. *Pterocaulæ*, *nempè foliis in alas foliaceas secus caulem decurrentibus.*

* *Foliis pinnatilobatis alternis.*

16. *V. GIGANTEA* (Jacq. ic. rar. 1. t. 175. coll. 1. p. 53), caule fruticoso simpliciter glabro tereti, foliis alternis infer. decurrentibus, sup. sessilibus auritis amplexicaulibus basi cuneatis suprâ pinnatipartitis lobis utrinquè 3-4 lanceolatis subintegerrimis suprâ pubescentibus subtus molliter villosos-canescents, corymbo composito, invol. squamis lanceolatis, ligulis paucis aut nullis, achænio glabro. § in insulis Caribæis (Jacq.), fruticetis montosis Jamaicæ (Swartz fl. Ind. occ. 3. p. 1368). Capitula ovato-oblonga circ. 12-flora. Cor. albæ. (v. s.)
17. *V. PINNATA* (Clark. ex herb. et mss. L'Hér.), caule suffruticoso tereti molliter villosos alato, foliis alternis confertis breviter decurrentibus pinnatipartitis basi cuneatis, junioribus subvillosis, adultis utrinquè pubescentibus, lobis utrinquè 4-5 lanceolatis serratis. § ? patr. ign. ex L'Hér. mss. Corymbus compositus villosus confertus. Flores albi, ligulis 2-3 sæpè nullis. Achænia jun. subvillosa. Invol. subimbricatum. Confusa cum *V. gigantea*, sed differt foliis etiâ summis evidenter decurrentibus multò minùs villosis, caule villosos, etc. (v. s. sine flor. ex h. L'Hér.)
18. *V. TURBACENSIS* (H. B. et Kunth nov. gen. am. 4. p. 203), caule suffruticoso ramosissimo apice cano-tomentoso, foliis (superioribus) alternis decurrentibus sinuato-lacinatis basi valdè angustatis suprâ piloso-hirtis subtus tenuiter canescenti-tomentosis, corymbis ramosissimis, invol. campanulatis squamis spatulato-lanceolatis, ligulis 6, achæniis glabris. § ad Turbaco Novo-Granaten-sium. Herba procera affinis *V. giganteæ*. Cor. flavæ.
19. *V. SINUATA* (Ell. sketch 2. p. 411), caule herbaceo striato pubescente infernè alato, foliis alternis, sup. sessilibus inf. decurrentibus, sinuatis pinnatifidisve basi attenuatis apice acutis suprâ scabris subtus pubescentibus, corymbo composito, invol. squamis subimbricatis, ligulis 3-5, achæniis subulatis. ¶ in Carolinâ merid. (Nutt.), in arenosis insularum maritimarum Edding et Hilton-Head (Ell.). *V. laciniosa* Nutt. gen. am. 2. p. 170 excl. syn. Lam. Flores albi. — Fortè ex Ell. hinc referenda *Siegesbeckia laciniosa* Poir. dict. 7. p. 158 à cl. Bosc in Carolinâ lecta?
20. *V. DIVERSIFOLIA*, caule herbaceo glabriusculo, foliis alternis decurrentibus ovatis acuminatis basi longè cuneatis suprâ sparse puberulis subtus molliter pubescenti-velutinis, aliis grossè hinc indè incisis, aliis trilobis lobis lanceolatis, summis sessilibus indivisis, corymbo composito polycephalo, invol. oblongi squamis lineari-lanceolatis, ligulis paucis, achæniis glabris. ¶ circa Bahiam Brasiliæ leg. cl. Blanchet et Salzmann. Affinis *V. pinnatifidæ* sed folia alterna, *V. giganteæ* sed suprema etiâ decurrentia. Flores albi. Caulis 5-6-pedalis. Var. habeo foliis indivisis. (v. s.)

** *Foliis pinnatilobatis oppositis.*

21. *V. PINNATIFIDA* (Cav. ic. 1. p. 67. t. 100 non Swartz), caule herbaceo tomentoso alis 4 aucto, foliis oppositis longè decurrentibus pinnatifidis basi longè cuneatis utrinquè præsertim in nervis puberulo-hirtis, lobis utrinquè 2-4 lanceolatis hinc indè subincisis, paniculâ oppositè ramosâ apice corymbosâ, invol. squamis lineari-lanceolatis acutis, ligulis circ. 12 oblongis, achæniis glabris alatis. ¶ in Mexico (Cav.). Flores pallidè flavi. Invol. nigricantia. Achænia radii interdum 3-alata ex Cav. (v. s.)

ℓ. *undulata*, alis caulinis valdè undulato-crispatis, caule minùs tomentoso basi suffruticoso, foliorum lobis regulariter dentatis. — Jacq. h. Schœnbr. 3. t. 305.

*** *Foliis indivisis oppositis.*

22. *V. PHARTUSA* (Cass. dict. 51. p. 476 et 59. p. 149), caule herbaceo alis 4 foliaceis integerrimis aucto, foliis oppositis decurrentibus ovato et fere rhombeo-lan-

ceolatis triplinerviis glabrinusculis serratis, paniculâ trichotomâ polycephalâ apice corymbosâ, invol. squamis paucis obtusis, ligulis 1-5 homochromis, achæniis villosis nunc exaristatis nunc 1-2-aristatis pilis nonnullis intermediis. \mathcal{L} in Amer. bor. sed locus propr. ign. Frequenter in hortis bot. Eur. culta ubi abortu subdioica evadit. Phaetusa Americana Gärtn. fr. 2. p. 465. t. 169. Lem. syn. 216 excl. syn. Poir. Siegesbeckia occidentalis Linn. spec. 1269. Phaetusa borealis Spreng. syst. 3. p. 59r. — Pluk. mant. t. 342 fig. penult. Certè specificè differt à V. Siegesbeckiâ Michx. Foliis grossè triplinerviis et à Verbesinæ caract. paulò recedit ob achænia inter aristas seriem pilorum brevium gerentia, sed non admittere queo sententiam cl. Lessingii, qui Phaetusæ genus in tribu Verbesinæ omninò alienâ collocavit. (v. s. c.)

23. V. SIEGESBECKIA (Michx. fl. bor. am. 2. p. 134), caule herbaceo glabro alis 4 foliaceis integerrimis aucto, foliis oppositis decurrentibus ovali-lanceolatis utrinquè acuminatis penninerviis glabrinusculis grossè serratis, paniculâ trichotomâ polycephalâ apice corymbosâ, invol. squamis paucis obtusis, ligulis 1-5 homochromis, achæniis exalatis subpuberis, aristis rigidis, pilis nullis intermediis. \mathcal{L} in umbrosis Amer. bor. sylvis à Virginiâ ad Carolinam (Pursh, Bosc!) et in Georgiâ (h. Duby!). Willd. sp. 3. p. 2224. Verbesina occidentalis Walt. fl. car. p. 213. Siegesbeckia Gron. virg. 128 sed Siegesbeckia occidentalis Linnæi qui expressis verbis dixit folia triplinervia ad V. Phaetusam potius referenda. (v. s. sp. nec nunquàm in hort. Eur. cult. vidi.)

**** Foliis indivisis alternis.

24. V. MICROPTERA, caule suffruticoso apice pubescenti-velutino, foliis alternis decurrentibus ovatis basi longè petioliformibus cuneato-angustatis obtusis obtusè sinuato-dentatis suprâ scabridis subtus vix pubescentibus, ramis flor. subnudis, capitulis 3-10 paniculatis, invol. squamis linearibus, ligulis 6-7, achæniis cuneato-linearibus vix alatis. \mathcal{L} \mathfrak{z} in Mexico inter Laredo et Bejar febr. florentem legit cl. Berlandier (pl. exs. n. 1442). Flores ex sicco vid. carnei. Capitula subrotunda. Ligulæ obovatæ disco paulò longiores. (v. s.)
25. V. VIRGATA (Cav. ic. 3. p. 38. t. 273), caule suffruticoso glabro, ramulis pedicellisque pubescentibus, foliis alternis lineari-lanceolatis utrinquè attenuatis serratis brevissimè petiolulatis, aliis basi nudis, aliis auriculâ subrotundâ utrinquè acutis, aliis breviter decurrentibus, corymbo terminali, pedicellis centralibus brevissimis, invol. squamis lanceolatis acutis, ligulis 10-12, paleis lanceolatis, jun. apice rigidè uncinatis. \mathfrak{z} in Mexico undè habui lectam à cl. Née, Alaman, Mairet. Flores lutei. Achænia pubescenti-hispida margine alata ciliolata biaristata. (v. s.)

\mathcal{L} ? conyzoides, foliis suboppositis basi auriculatis. V. conyzoides fl. mex. icon. ined. (v. ic.)

26. V. POLYCEPHALA, caule herbaceo? brevissimè pubescenti-velutino, foliis alternis decurrentibus ellipticis acuminatis basi longè attenuatis integerrimis suprâ sparsè scabridis subtus molliter pubescenti-velutinis subincanis, corymbo composito polycephalo, invol. oblongi squamis lineari-lanceolatis puberis, ligulis 3-4, achæniis striatis hispidulis latè alatis. \mathcal{L} ? \mathfrak{z} ? in Mexico prope Bejar angusto flor. legit cl. Berlandier (pl. exs. n. 1813). Caulis 4-6-pedalis. Flores albi. Pedunculi albido-tomentosi. (v. s.)
27. V. PTEROCAULA (fl. mex. icon. ined.), caule herbaceo glabrinusculo, foliis alternis longè decurrentibus lineari-lanceolatis utrinquè acuminatis acutis serratis glabris, corymbo 10-12-cephalo, invol. squamis linearibus obtusis, ligulis 10-12, achæniis angustè alatis et ciliatis. \mathcal{L} ? in Mexico. Cor. luteæ. Habitus V. virgatæ sed folia omnia decurrentia. (v. ic.)
28. V. VIRGINICA (Linn. sp. 1270), caule herbaceo pubescente infernè subalato apice aptero, foliis alternis penninerviis lanceolatis seu obovato-lanceolatis subserratis glabrinusculis, infimis decurrentibus, corymbo composito, invol. squamis subobtusis pubescentibus ligulis 3-4 heterochromis, achæniis puberis angustè alatis. \mathcal{L} in sylvis umbrosis à Pensylvaniâ ad Carolinam (Pursh, Bosc!),

in Kentucky orient. (Raf.!), Georgia (Ell. sketch 2. p. 410). *Anomantia* Raf. ! in litt. 1831. *V. paniculata* Poir. ! dict. 8. p. 456. Ligulæ albæ ovales. Achæniis aristæ 2 pilis nullis intermediis. (v. s. sp.)

29. *V. mollis* (H. B. et Kunth nov. gen. am. 4. p. 203), caule herbaceo simpliciter dense hirtio, foliis alternis decurrentibus lanceolatis obtusè serratis suprâ densè piloso-hirtis subtus albo-lanatis, capitulis ternis subeorymbosis, invol. semiglobosi squamis ligulisque lineari-lanceolatis, achænio glabro margine ciliato, aristis valdè inæqualibus. ① in Mexico inter Guanaxuato et Villalpando. Flores flavi. *V. alata* ex radice annuâ et aristis inæqualibus affinis.

Sect. II. HAMULIUM. — *Hamulium* Cass. dict. 20. p. 260.

Capitulum radiatum, ligulis numerosis subbiseriatis. Achæniis aristæ 2 inæquales, minore rectâ majore apice uncinatâ. — Herba annua, foliis decurrentibus.

30. *V. ALATA* (Linn. sp. 1270), caule herbaceo glabriusculo alato, foliis alternis longè decurrentibus oblongis ovatisve obtusè undulatis et sinuato-dentatis glabriusculis, capitulis subglobosis solitariis terminalibus, invol. squamis oblongis, achæniis hispiduli alati ciliolati aristis inæqualibus, majore apice uncinatâ. ② in Curaçò (Volk. nor. t. 106), Surinamo (Comm. h. amst. 1. t. 3), Jamaica (Sloan. Br.), in Mexico (fl. mex. ined. ic.), ins. Cubâ (H. B.). Swartz obs. p. 313. Sims bot. mag. t. 1716. *Hamulium alatum* Cass. dict. 20. p. 261. — Magn. hort. t. 40. Cor. crocæ, ligulæ numerosæ subbiseriales breves. (v. v. c. et s. sp.)

Sect. III. PLATYPTERIS. — *Platypteris* H. B. et Kunth nov. gen. am. 4. p. 200. Cass. dict. 50. p. 259.

Capitula discoidea, ligulis planè nullis.

§. 1. *Involucro imbricato, foliis decurrentibus oppositis, caule herbaceo.*

31. *V. CROCATÀ* (Less. syn. 232), herbacea, ramis tetrapteris apice hispido-scabris, foliis oppositis decurrentibus irregulariter pinnatilobatis, lobis ovalibus eroso-dentatis, terminali subdeltoideo, capitulis pedunculatis solitariis, invol. squamis imbricatis paleisque lineari-subulatis, ligulis nullis, achæniis glabris latè alatis biaristatis. ② in Mexico propè Tusco (H. et B.) et alibi. *Bidens crocata* Cav. ic. 1. p. 66. t. 99. *Spilanthes crocata* Sims bot. mag. t. 1627. *Platypteris crocata* H. B. et Kunth nov. gen. am. 4. p. 201. Cass. dict. 50. p. 259. Capitula globosa crocea. (v. s.)

§. 2. *Invol. circ. bi- seu triseriali non sensim imbricato, foliis alternis non decurrentibus, caule fruticoso.*

32. *V. AURICULATA*, caule suffruticoso tereti brevissimè et confertè velutino, foliis alternis obtusè auriculatis sessilibus ellipticis acuminatis basi longè attenuatis penninerviis suprâ scabris subtus molliter et sordidè tomentosis subdentatis, capitulis plurimis breviter pedicellatis ad apices ramorum racemoso-corymbosis, invol. squamis ovali-oblongis subacutis 2-3-serialibus, rec. paleis elongatis rigidis apice subulatis, ligulis nullis. ⑤ in Mexici ditione Oaxacanâ ad Tehuantepec august. flor. legit cl. Andrieux (pl. exs. n. 302). Folia 7 poll. longa, 2 poll. lata. (v. s. florib. non satis evol.)
33. *V. ARBOREA* (H. B. et Kunth nov. gen. am. 4. p. 202), arborea, ramis teretibus tenuiter fusciscenti-villosis, foliis alternis petiolatis oblongis acuminatis integerrimis suprâ hirtis subtus lanato-tomentosis et fusciscenti-incanis, corymbis ramosissimis foliatis, invol. imbricati campanulati squamis obtusis oblongo-linearibus, ligulis nullis, floeculis 10-12, paleis obtusis. ⑤ in frigidis montosis agri Quitensis præes. ad montem Pichincha.

† *Verbesinæ non satis notæ.** *Americanæ.*

34. V. *ÆSTUANS* (Lam. dict. 3. p. 262), caule simplici, foliis oppositis petiolatis subspathulatis acutis penninerviis dentatis subtus tomentosis, capitulis pedunculatis, invol. ovato adpressè imbricato, ligulis linearibus numerosissimis. — in Amer. calid. Plum. ed. Burm. t. 41. f. 2. Asteris sp. Tourn. Helenii sp. Vaill. Inula æstuanus Linn. sp. 1236. Propius ad Verbesinam quàm ad alia genera citata accedere videtur, sed car. flor. et fruct. vix noti. †.
35. V. *FOLIACEA* (Spreng. syst. 3. p. 578), herbacea, foliis oppositis petiolatis ovatis crenatis pubescentibus, pedunculis adgregatis axillaribus abbreviatis hirsutis, invol. foliaceo. — in Guadalupà (Bert.), in Brasilià (Sell.). An plantæ duæ diversæ? An generis?
36. V. *ILICIFOLIA* (Poir. dict. 8. p. 460), caule...., ramis teretibus glabris, foliis oppositis petiolatis coriaceis elliptico-lanceolatis utrinquè acutis grossè incisodentatis triplinerviis suprâ glabris subtus petiolisque sparsè et minutè puberulis, capitulis pedicellatis subsolitariis, invol. squamis lanceolatis.... — in Sancto-Domingo. V. fruticosa Linn. sp. 1271? non Willd. V. frutescens Plum. ed. Burm. t. 52? Cor. luteæ. Cæt. car. ign. An Flourensia species? (v. s. folium.)
37. V. *MEGAPOTAMICA* (Spreng. syst. 3. p. 578), caule herbaceo simplici oligocephalo, foliis oppositis breviter petiolatis lanceolatis integerrimis substrigillosis, pedunculis axillaribus elongatis, radio angusto. — in Brasiliæ provinciâ Rio-Grande.
38. V. *MONTVIDENSIS* (Spreng. syst. 3. p. 578), caule herbaceo simplici 1-cephalo hispido, foliis oppositis subfasciculatis lanceolatis integerrimis hispidissimis, capitulo terminali solitario. — in Monte-Video. Affinis ex phrasi vid. V. buphthalmoides sed descr. incompl.
39. V. *TRIDENTATA* (Spreng. syst. 3. p. 577), fruticosa, foliis oppositis cuneatis in petiolum decurrentibus 3-dentatis subserratis scabris triplinerviis, pedunculis 1-cephalis, invol. simpliusculo. ♀ in Amer. merid. (Willd.), in Martinicâ (Spr.). V. fruticosa Willd. sp. 3. p. 577 excl. syn. et Spreng. †.

** *Gerontogea.*

40. V. *CHINENSIS* (Linn. sp. 1270), subtomentosa fruticosa, caule tereti, foliis alternis petiolatis ovato-lanceolatis serratis, capitulis terminalibus solitariis pedunculatis, invol. hemisphærico, achænio 4-gono setis 4 lævibus coronato. ♀ in Chinâ. Fl. lutei.
41. V. *SPICATA* (Lour. fl. coch. 2. p. 620), herbacea ramosa, foliis alternis longè petiolatis serratis subtus tomentosis, inf. palmato-incisis, sup. lanceolatis, capitulis spiciformibus congestis terminalibus, invol. squamis subulatis biserialibus, ligulis 5 unilateralibus invol. æqualibus, achænio calvo. — Culta in hortis Cochinchinæ et Chinæ. Cum acetariis mensis apponitur. Flores albiissimi radiati. A Verbesinâ videtur omninò aliena! sed locum aptiorem ignoro.

†† *Species exclusæ.*

- V. *ACMELLA* Linn. = *Spilanthes Acmella*.
 V. *ALBA* Linn. = *Eclipta erecta*.
 V. *ASTEROIDES* Linn. = *Amellus Lychnitis*.
 V. *BIFLORA* Linn. = *Wollastonia biflora*.
 V. *BIFLORA* Blume. = *Wollastonia scabriuscula*.
 V. *BIFLORA* var. Wall. = *Wollastonia urticæfolia* β.
 V. *BOSWALLIA* Linn. f. = *Glossocardia Boswallæ*.
 V. *CALENDULACEA* Linn. = *Wedelia calendulacea*.

- V. CANESCENS Gaudich. = *Wollastonia canescens*.
 V. CONNATA Gaudich. = *Lipochæta connata*.
 V. CORCOPSIS Michx. = *Actinomeris alternifolia*.
 V. DICHOTOMA Murr. = *Blainvillea rhomboidea*.
 V. DICHOTOMA Wall. = *Blainvillea latifolia*.
 V. HASTULATA Hook. = *Lipochæta hastulata*.
 V. HELIANTHOIDES Michx. = *Actinomeris helianthoides*.
 V. LANCEOLATA Poir. = *Blainvillea rhomboidea* β .
 V. LAVARUM Gaudich. = *Lipochæta lavarum*.
 V. LAVENIA Linn. = *Adenostemma Lavenia*.
 V. LOBATA Gaudich. = *Lipochæta lobata*.
 V. MEMBRANIFOLIA Poir. = *Wulffia membranifolia*.
 V. MOLUCCANA Blume. = *Wollastonia Moluccana*.
 V. MONTANA Blume. = *Wollastonia montana*.
 V. MUTICA Linn. = *Chrysanthellum procumbens*.
 V. MUTICA Sw. = *Chrysanthellum Swartzii*.
 V. NODIFLORA Linn. = *Synedrella nodiflora*.
 V. OPPOSITIFLORA Poir. = *Wulffia membranifolia*.
 V. OVATA hort. Par. = *Heliopsis bupthalmoides*.
 V. PROSTRATA Linn. = *Eclipta prostrata*.
 V. PSEUDO-ACMELLA Linn. = *Spilanthes Pseudo-Acmella*.
 V. FUSILLA Poir. = *Eclipta pusilla*.
 V. REPANDA Pers. = *Lepidophorum repandum*.
 V. SATIVA Roxb. = *Guizotia oleifera*.
 V. SCABERRIMA Wall. = *Wedelia urticæfolia* γ .
 V. SUCCULENTA Hook. = *Lipochæta succulenta*.
 V. STRIGULOSA Gaudich. = *Wollastonia aristata*.
 V. URTICÆFOLIA Blume. = *Wedelia urticæfolia* γ .

CCCLIX. DITHRICHUM *Cass. bull. philom.* 1817. febr. dict. 13. p. 272. —
Spilanthes sp. *Less.*

Capitulum pluriflorum homogamum, fl. hermaphr. tubulosis, cor. extus pilosis. Invol. floribus longius cylindraceum biseriale, squamis ext. brevissimis laxis, int. longis erectis coriaceis acuminatis. Recept. planum, paleis in appendicem subulatam productis. Achænia compressa obovata glabra. Pappus biaristatus, aristis ex angulis ortis crassis subbarbellulatis. — Herba elata. Caulis teres pubescens. Folia alterna sessilia subdecurrentia oblongo-lanceolata irregulariter sinuato-lobata scabra ferè pedalia. Capitula paniculato-corymbosa. Flores flavi.

1. D. MACROPHYLLUM (Cass. ll. cc.). — Patr. ign. verisim. hab. in Americâ; confusa enim latebat in herb. Juss. sub nomine *Conyze lobatæ*.

CCCLX. MICRACTIS.

Capitulum multiflorum, fl. radii minutissimè ligulatis fœmineis, disci tubulosis 4-dentatis hermaphroditis, centralibus fortè sterilibus. Invol. 1-2-seriale, squamis oblongis subconcavis dorso subpuberulis. Recept. paleæ etiâ complicato-concavæ sed glabræ, ut squamæ inv. achænia semi-involventes. Stylus radii bifidus, ligulâ integrâ concavâ

paulo longior, disci inclusus. Anth. 4 ecaudatæ. Achænia tetragonocompressa ferè obovata glabra omninò calva. — Herba Madagascariensis annua erecta minutissimè puberula. Folia opposita oblonga acuta vix subcrenata scabra, super. integra. Capitula ad apices ramorum terna congesta breviter stipitata. Achænia nigra inter paleas persistentes seu tardè deciduas semioccultata. Flores minutissimi flavi. — Ambigit inter capitula monoica (ex Bojer) et heterogama, sed potiùs ad hæc meo sensu referenda. An potiùs ad Millerieas seu Silphieas rejicienda?

1. M. BOJERI. ① in ins. Madagascar legit cl. Bojer. Habitus plebeius sed structura singularis præsertim ob ligularum minutiam, et florum disci stylum exiguum fortè sæpiùs abortivum. (v. s. comm. à cl. Bojer.)

CCCLXI. SPILANTHES Jacq. Amer. 212. Schreb. gen. 1266. — *Spilanthos* Linn. mant. 475. Gærtn. fr. 2. t. 167. Less. syn. 232. — *Pyrethrum* Medic. act. pal. 3. p. 237 non Willd. — *Ceruchis* Gærtn. mss. ex Schreb. — *Spilanthos* et *Acmella* Rich. in Pers. ench. 2. p. 293 et 472. Cass. dict. 59. p. 137 et 138. 50. p. 257 et 258. p. 24. 328. — *Spilanthus* et *Athronia* Neck. elem. 1. p. 87 et 32.

Capitulum multiflorum, nunc heterogamum fl. radii ligulatis fœmineis sæpiùs parvis, nunc homogamum fl. omnibus hermaphrod. tubulosis 4-5-dentatis. Invol. biseriale adpressum disco brevius, squamis ext. subfoliaceis, inter. submembranaceis complicatis. Styli fl. herm. rami apice truncati et penicillati. Anth. nigricantes. Achænia disci compressa erostria sæpiùs ad latera ciliata seu exaristulata, radii (ubi adest) triquetra seu obcompressa. — Herbæ sæpiùs annuæ interdum perennes aut suffruticulosæ dichotomæ aut vagè ramosæ, Geronto- et sæpiùs Neogææ. Folia opposita subintegra. Pedunculi terminales aut è dichotomiis orti erecti 1-cephali. Capitula ovata aut conica. Cor. flavæ rariùs albæ, ligulis disco concoloribus, rariùs discoloribus. — Herbæ pleræque gustu fervidæ et salivariæ.

Sect. 1. Acmella. — *Acmella* Rich. in Pers. ench. 2. p. 472. Cass. dict. 24. p. 531. — *Athronia* Neck. elem. 1. p. 32.

Capitula radiata, ligulis basi hispidis. Achænia radii sæpiùs trigona triaristata seu rariùs calva, aut angulo interiore nudo biaristata, aut angulo inter. obliterato ferè obcompressa.

1. S. ARNICOIDES, caule repente adscendente hirtello parcè ramoso, foliis ad basin subconfertis oppositis in petiolum longè attenuato-cuneatis ellipticis ovato-ve sublanceolatis subdentatis glabriusculis, pedunculo longissimo 1-cephalo erecto sparsè puberulo, capitulo subgloboso, invol. squamis circ. 12 ovali-oblongis obtusis, ligulis 12-15 invol. triplò longioribus, achæniis radii abortivis, disci ciliatis biaristatis. ♀ in Brasiliæ prov. Rio-Grande. Cor. intense luteæ ferè crocææ, in specimine majori pallidiùs flavæ, in altero (verisim. conspecifico) capitulum majus et magis depressum quàm in cæteris sp. (v. s. in h. Mus. reg. Par. à Mus imp. Bras. sub n. 1023 miss.)
2. S. DORONICOIDES, caule basi subrepente adscendente basi glabro inter folia confertè hispido, foliis lanceolatis obtusis basi longè cuneato-attenuatis subpetiolatis integerrimis hinc indè ad nervum et margines subsetosis, pedunculo elongato rigido 1-cephalo subangulato adpressè puberulo, capitulo ovato obtuso, invol. squamis ovali-oblongis obtusis, ligulis triplò longioribus obtusè 3-dentatis, achæniis subciliatis, radii 3-, disci 2-angulatis aristatisque. ♀ in Brasiliæ prov. Rio-Grande. Affinis S. arnicoidi. Cor. flavæ. Herba florens semi-

- pedalis, pedunc. 4-poll. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1038.)
3. *S. LONGIFOLIA*, caule erecto simplici hispido ad medium foliato supernè in pedunculum longissimum 1-cephalum sub capitulo hirsutum producto, foliis oblongo-lanceolatis basi attenuatis integerrimis medio subtriplicinerviis, infer. brevioribus glaberrimis, super. longioribus hispidis, invol. squamis linearibus circ. 15-16, ligulis invol. duplò longioribus.... 2½ in Brasiliæ prov. Rio-Grande. In specim. malè servato car. fruct. non vidi, sed ex analogiâ cum *Sp. arnicoide* de genere non dubito. Folia 3 poll. longa, 5 lin. lata. Ligulæ crocææ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1029 miss.)
4. *S. BLEPHARICARPA*, caule basi repente adscendente mox erecto infra foliorum juga hirsuto cæterum glabriusculo, foliis lineari-lanceolatis obtusis basi longè cuneato-attenuatis integerrimis glabris, pedicellis elongatis striatis 1-cephalis, capitulis ovatis obtusissimis, invol. squamis ovalibus substriatis, ligulis invol. duplò longioribus, acheniis radii 3- disci 2-angulatis aristatisque. angulis densè et longè ciliatis. 2½ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1030!), et fortè eandem in prov. Sancti-Pauli legit cl. Lund. Herba pedalis ad *S. macropodam* accedens. Flores flavi. (v. s.)
5. *S. MACROPODA*, caule herbaceo stricto subsimplici glabriusculo, foliis oppositis petiolatis oblongo-lanceolatis subacutis basi longè attenuatis integerrimis glabris margine scabris, pedicellis longissimis paucis 1-cephalis apice subpuberis, capitulis ovato-globosis, invol. squamis ovatis circ. 10 dorso trinerviis, ligulis circ. 10 elongatis 3-dentatis, acheniis glabris radii 3- disci 2-aristatis. — in Brasiliæ prov. Rio-Grande. Caulis circ. pedalis. Pedicelli 10-12 poll. longi. Cor. flavæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1049 miss.)
6. *S. LEPTOPHYLLA*, glaberrima, caule basi repente mox adscendente basi folioso-parcè ramoso, foliis linearibus integerrimis rigidulis, pedunculis nudis 1-cephalis caule longioribus, invol. squamis 10-12 oblongis, capitulis ovato-globosis, ligulis invol. triplò ferè longioribus, acheniis.... 2½ in Brasiliæ prov. Rio-Grande inter muscos. Car. fruct. in specim. carie subdestructo non vidi. Affinitas summa cum *S. macropoda*. Flores flavi. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1026.)
7. *S. STOLONIFERA*, caule à basi ramoso stolonifero, ramis floridis adscendentibus hinc indè hispidulis, foliis lanceolatis basi attenuatis subobtusis integerrimis triplinerviis glabris basi vix subsetoso-ciliatis, pedicellis terminalibus elongatis apice adpressè pubescentibus, capitulis globosis, invol. squamis elliptico-oblongis subacutiusculis, ligulis invol. duplò triplòve longioribus 3-dentatis, acheniis ciliatis aristulatis. 2½ in Brasiliæ prov. Rio-Grande. Habitus ferè *Hieracii Auriculæ*. Flores flavi. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1077.)
8. *S. SPHÆROCEPHALA*, glabra debilis, caule tereti gracili repente, foliis oppositis in petiolum attenuatis oblongis basi trinerviis vix subserratis, capitulis ramos apice nudos terminantibus globosis aut vix ovatis, invol. squamis lanceolatis, ligulis 8-10 invol. duplò longioribus apice dentatis, acheniis ciliatis, radii triquetris 3-aristatis, disci compressis biaristatis. — in Brasiliæ prov. Rio-Grande. Flores flavi. Caules 8-12 poll. longi. Foliorum juga remota. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1075 miss.)
9. *S. FIMBRIATA* (H. B. et Kunth nov. gen. am. 4. p. 208), caulibus erectis vel procumbentibus subsimplicibus glabris, foliis oppositis petiolatis ovatis acutis basi subrotundatis dentatis glabris, pedicellis terminalibus demùm alaribus elongatis, capitulis breviter radiatis, invol. squamis circ. 7 oblongis adpresso-pilosis, acheniis radii triangularibus, disci oblongis, omnibus ciliatis 2-3-? aristulatis. ① in Andibus Novo-Granatensium prope Ybague et Passo del Machin. *Acmella fimbriata* Cass. l. c. Flores lutei.
10. *S. CILIATA* (H. B. et Kunth nov. gen. am. 4. p. 208), caule erecto? tetragono glabro, ramis junioribus hispidulis, foliis oppositis petiolatis ovatis ac-

minatis basi paulò angustatis grossè dentatis glabris petiolisque ciliatis, pedicellis terminalibus demum alaribus folio longioribus, capitulis breviter radiatis, invol. squamis 6-7 ovato-oblongis acutis glabris, achæniis lævibus? ciliatis hiaristulatis. ① in Amer. merid. prope Chipò et Santa-Fé de Bogota. *Acmella ciliata* Cass. l. c. Flores flavi. Affin. *Sp. Mutisii* et *fimbriatæ* dicitur.

11. *S. MUTISII* (H. B. et Kunth nov. gen. am. 4. p. 209), caulibus erectis aut procumbentibus subdichotomis glabriusculis, foliis oppositis petiolatis ovatis acutis basi subrotundatis grossè dentatis glabris, pedicellis subhispidulis terminalibus folio longioribus, capitulis subrotundo-ovatis radiatis, invol. squamis circ. 10 ovato-oblongis pubescentibus, ligulis 10 invol. longioribus acutè 3-dentatis, achæniis exaristatis, radii subverrucosis ciliolatis, disci lævibus glabris. ①? prope Santa-Fé de Bogota. *Athemis Americana* Mutis in Linn. f. suppl. 378. Anth. occidentalis Willd. spec. 3. p. 2183. Anth. oppositifolia Lam. dict. 1. p. 576. *Acmella occidentalis* Pers. ench. 2. p. 473. *Heliopsis*? *dubia* Dunal mem. Mus. Par. 5 p. 58? *Acmella Mutisii* Cass. dict. 24. p. 33 r. Ligulæ tubo piloso majores quàm in cæt. spec., corollæque disci flavae. (v. s.)
12. *S. GRANDIS*, caule erecto ramoso hispidulo, foliis oppositis ellipticis in petiolum cuneato-attenuatis apice obtusis triplinerviis subintegerrimis glabris, pedunculis folio triplò longioribus 1-cephalis striatis rigidulis, capitulis subglobosis, invol. squamis ellipticis obtusis, ligulis invol. duplò triplòve longioribus apice dentatis, achæniis subciliatis aristatis. — in Brasiliæ prov. Rio-Grande. Herba sesquipedalis. Folia 3 poll. longa, 8-10 lin. lata. Flores flavi, ligulis ex sicco subtus striato-purpurascentibus. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub. n. 1005.)
13. *S. BECCABUNGA*, caule basi repente glabro dein adscendenti - erecto apice subhispido, foliis oppositis petiolatis latè ovatis subacutis repandulo-subdentatis triplinerviis glabriusculis, pedicellis terminalibus glabriusculis folio multò longioribus, capitulis rotundo-ovatis radiatis, invol. squamis circ. 12 ovali-oblongis, ligulis circ. 10 invol. longioribus sub-3-dentatis, achæniis glabris exaristatis. 2/ in Mexico in valle Mexici et Toluçæ legit cl. Berlandier (pl. exs. n. 720 et 1160!). Flores latei. Capitula ferè magnit. *S. oleracei*. Fol. sesquipoll. longa, pollicem lata, petiolo 3-4 lin. (v. s.)
14. *S. PÖPPIGII*, caule erecto subtetragono glabriusculo, foliis oppositis petiolatis ovatis subacutis dentatis trinerviis suprà glabris subtus secus nervos sparse strigilloso-pilosis, petiolis ciliatis, pedicellis terminalibus demum alaribus petioli longitudine, capitulis ovatis radiatis, invol. squamis ext. circ. 8 ovalibus ciliatis, ligulis parvis, achæniis submarginatis ciliatis exaristatis aut aristulam minimam gerentibus. — in Peruvia legit cl. Pöppig. (pl. exs. n. 32. diar. n. 1205!). Cor. disci 4-dentatæ. Ligulæ flavae. (v. s.)
15. *S. LUNDII*, caule è nodis radicante demisso, ramis erectis subpuberulis, foliis oppositis petiolatis ovato-lanceolatis dentatis glabriusculis, petiolis parè ciliatis, pedicellis terminalibus folio longioribus, capitulis ovato-conicis radiatis, invol. squamis ext. 7-9 ovato-lanceolatis eciliatis, ligulis totidè invol. paulò longioribus, achæniis ciliatis tenuè aristulatis. 2/ ad rivulos circa Rio-Janeiro legit cl. Lund. Ligulæ et cor. disci flavae. (v. s.)
16. *S. SUBHIRSUTA*, caulibus repentibus adscendentibus vix ramosis petiolis pedicellisque hispidò-subhirsutis, foliis oppositis petiolatis ovatis aut summis ovato-lanceolatis subacutis acutè serratis sparse hispidulis, pedicellis terminalibus folio longioribus, capitulis subrotundo-ovatis parè radiatis, invol. squamis circ. 10 lanceolatis, ligulis obovatis 3-dentatis invol. paulò longioribus, achæniis glabris aristatis. 2/ in Mexico circa Tampico de Tamaulipas legit cl. Berlandier (pl. exs. n. 149!). Ligulæ corollæque disci flavae. (v. s.)
17. *S. CÆSPITOSA*, caule à basi ramoso subradicante procumbente cæspitoso glabro, foliis oppositis confertis petiolatis lanceolatis basi acutis apice acuminate vix subdentatis glabris, pedicellis terminalibus folio multò et caule ferè longioribus, capitulis subrotundis ovatis parè et minutè ligulatis, invol. squa-

mis 8-10 lanceolatis, achæniis ciliatis exaristatis. \mathcal{U} in Brasiliæ prov. Minarum-Generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 3081). Affinis S. Salzmanni, sed habitus diversissimus et achænia aut ad latus superius aut utrinquè ciliata. Basis caulis ferè suffrutescit. (v. s.)

18. S. KARVINSKIANA, caule imà basi suffruticuloso ramoso erecto diffuso glabro sub nodis hirsutulo, foliis oblongo-lanceolatis sessilibus acutis subintegerrimis glabriusculis setis longis hinc indè ciliatis, pedicellis axillaribus 1-cephalis folii longitudine et ex dichotomiis folio longioribus, invol. squamis oblongo-linearibus extus hirtellis, ligulis parvis oblongis indivisis, achæniis glabris calvis angulatis. \mathcal{U} inter plantas Mexicanas Karvinskii sine loci indicatione hanc video. Ligulæ parvæ flavæ subtus virescentes. Valdè affinis S. Mariannæ. (v. s. in h. acad. reg. Monac.)

19. S. MARIANNÆ, glabra herbacea erecta debilis, foliis oppositis in petiolum attenuatis ovali-lanceolatis utrinquè acuminatis triplinerviis ferè quintuplinerviis subtus glabris suprà et margine junioribus subpuberis, pedicellis axillaribus 1-cephalis folio longioribus, invol. squamis oblongo-linearibus hirtellis, ligulis minimis linearibus indivisis bipartitisve, achænio glabro calvo angulato. $\textcircled{1}$ in Brasiliæ prov. Minarum-Generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 322). Capitula parva. Flores flavi. Achænia nigra. (v. s.)

20. S. CAULIRHIZA, caule prostrato sub petiolis radicante, foliis breviter petiolatis ovali-rhombeis subcrenatis ramisque glabriusculis, pedunculis è dichotomiis aut apice ramorum ortis scabridis folio triplò duplòve longioribus, capitulis subglobois, ligulis obovatis 3-dentatis involucri longioribus, achæniis glabris calvis. — in Sennaar Africæ borealis. *Acmella caulirhiza* Caill. et Delil. pl. afr. p. 45. t. 64. f. 7.

β . *Madagascariensis*, glabriuscula foliis ovatis paulò longius petiolatis, capitulo paulò majore, recept. breviori. — in insulâ Madagascar legit cl. Bojer. Fortè species propria? (v. s. comm. à cl. Bojer.)

21. S. ACMELLA (Linn. syst. veg. 610), caule basi subradicante erecto aut adscendente, foliis petiolatis ovato-lanceolatis subintegerrimis dentatisve glabriusculis, pedunculis folio triplò longioribus, capitulis ovatis, ligulis 5-6 minimis, achæniis ciliatis 1-2-aristatis. $\textcircled{1}$ in Indiâ orient. *Verbesina Acmella* Linn. mant. 475. *Acmella* Linnæi Cass. dict. 24. p. 330. et fortè *Acm. brachyglossa* v. 50. p. 258 ex ipso in Desf. cat. ed. 3. p. 186. Omnino refert S. pseudo-acmellam et fortè hujus mera varietas. Synonyma omnia confusa videntur. Icones omnes ad alteram citatæ ob ligularum minutiam etiâ hic citari possent. (v. s.)

22. S. REPENS (Michx. fl. bor. am. 2 p. 131), caule repente ramoso sparse subhirtello, foliis oblongo-lanceolatis brevissimè petiolatis acutis subintegerrimis, pedunculis terminalibus demùm alaribus folio longioribus, capitulis ovatis radiatis, invol. squamis circ. 12 lanceolatis, ligulis totidè sub-3-dentatis invol. duplò longioribus, achæniis scabridis exaristatis ad angulos eciliatis. \mathcal{U} in Carolinæ humidis. *Anthemis repens* Walt. car. 211. Pursh fl. 2. p. 562. *Acmella repens* Pers. ench. 2. p. 473. Ell. sketch 2. p. 406. Discus radiusque flavus. Folia pollicem longa, 2-3 lin. lata. (v. v. c.)

23. S. SALZMANNI, caule imà basi breviter adscendente radicante cæterùm erecto glabriusculo, foliis oppositis petiolatis ovatis aut ovato-lanceolatis subdentatis, pedicellis terminalibus demùm alaribus folio longioribus, capitulis ovato-conicis parcè radiatis, invol. squamis 8-10 lanceolatis, ligulis invol. æquantibus, achæniis glaberrimis eciliatis exaristatis. $\textcircled{1}$ in paludosis circa Bahiam Brasiliæ legit cl. Salzmann. Habitu accedit à S. uliginosam, sed major, cor. omnes flavæ et præ. achænia diversa. (v. s.)

24. S. AFRICANA, caule adscendente hinc indè setulis aspero, foliis oppositis petiolatis ovato-lanceolatis acutè et inæqualiter dentatis margine rigidè et brevè ciliatis, pedicellis folio duplò longioribus, capitulis ovato-conicis, ligulis involucri paulò longioribus, invol. squamis oblongis vix acutis, achæniis glabris

eciliatis exaristatis. — in Africâ austr.-orient. ad Zw. Omtata et Omsamwubo legit cl. Drege n. 5086! Valdè affinis *S. Salzmanni*, sed ciliis foliorum præcipuè diversa. Folia petiolo incluso var. 1 $\frac{1}{2}$ -2 $\frac{1}{2}$ poll. longa. (v. s.)

25. *S. ULIGINOSA* (Swartz prod. 110. flor. Ind. occ. 1291), caule erectiusculo dichotomo glabro aut apice subpuberulo, foliis petiolatis ovato- aut oblongo-lanceolatis subdentatis glabrusculis, pedicellis terminalibus demùm alaribus folio longioribus, capitulis jnn. subrotundis demùm ovato-conicis, invol. squamis circ. 10-12 exter. ellipticis, ligulis paucis minimis obtusis, achæniis ciliatis breviter biaristatis. (I) in uliginosis argillosis Jamaica australis (Swartz). Porto-Ricci (Ledru!), Sanctæ-Crucis? (Vest.). *Acmella uliginosa* Cass. dict. 24. p. 531. Cor. disci 4-dentatæ flavæ, ligulæ albæ. (v. s.)

26. *S. TENELLA* (H. B. et Kunth nov. gen. am. 4. p. 211), caule adscendente dichotomo glabro, foliis oppositis petiolatis ovato-oblongis acutis basi paulò angustatis subintegerrimis glabris margine scabrusculis, pedicellis folio longioribus angulatis pilosiusculis, capitulis parvis breviter radiatis, invol. squamis circ. 5 oblongis acutis, achæniis ciliatis 2-aristulatis, radii triangulari-compressis, disci compressis. ☉ circa Caracas. *Acmella tenella* Cass. l. c. Cor. flavidæ. An var. *Sp. debilis*?

27. *S. DEBILIS* (H. B. et Kunth nov. gen. am. 4. p. 210), caule adscendente dichotomo glabro, foliis oppositis petiolatis ovatis acutis basi rotundatis subintegerrimis glabris margine scabrusculis, pedicellis terminalibus et alaribus longissimis, capitulis radiatis, invol. squamis 5-6 elliptico-oblongis ciliatis, ligulis obovatis brevissimis obtusè 3-dentatis, achæniis ciliatis biaristulatis. (I) ad ripam fluminis Orinoci Amer. merid. *Acmella debilis* Cass. l. c. Cor. flavæ. Affinis dicitur *Sp. exasperatæ*.

28. *S. INTERMEDIA*, caule procumbente glabro, foliis ovato-lanceolatis dentatis, pedunculis terminalibus. — in Americâ. *Acmella intermedia* Rich. in Pers. ench. 2. p. 472. Car. mancus omnibus speciebus radiatis ferè conveniens. †

Sect. II. SALIVARIA — *Spilanthus* Cass. dict. 50. p. 257. — *Spilanthus* Pers. ench. 2. p. 493.

Capitula discoidea. Achænia omnia conformia compressa.

29. *S. RADICANS* (Schrader! cat. h. Goett. 1831. p. 5. non Jacq.), caulibus ramosis procumbentibus basi radicanibus, ramis adscendentibus magis minùsve pilosis, foliis oppositis petiolatis ovatis basi cuneatis à medio crenato-serratis margine scabrusculis, pedicellis terminalibus folio longioribus, capitulis ovato-oblongis, invol. squamis 12-15 elliptico-oblongis subciliatis, corollis radii elongatis tubulosis. ♀ patr. incert. Flores lutei. Capitula subradiata dixit cl. Schrader et reverà fl. marginales elongatos non verò ligulatos, sed verisim. steriles in specim. video. (v. s. comm. à cl. Schrader.)

30. *S. OLERACEA* (Jacq. hort. Vind. 2. t. 135), caule ramoso diffuso, foliis oppositis petiolatis latè ovatis basi obtusis truncatis aut subcordatis repando-subdentatis, pedicellis terminalibus 1-cephalis folio longioribus, capitulis crassis ovatis discoideis, invol. squamis 15-16 ovali-oblongis, achæniis ad angulos ciliatis biaristatis muticisve. (I) in Indiâ orientali? (Willd.), sed potius in Americâ merid. (Lam. Desf. Plum. sp. 10). Vulgò *Cresson de Para* hortul. *Pyrethrum Spilanthus* Medic. act. pal. 3. phys. p. 242 et 273. *Bidens acmeloides* Berg. act. holm. 1768. *Bidens fervida* Lam. dict. 1. p. 415. Herba dilutè virens, corollis etiã in medio disci flavis. Folia, imò herba tota, gustu fervida et intensè salivaria! (v. v. c.)

♀? *fusca*, corollis in medio disci fuscis, foliis obscure viridibus. (I) patr. ign. fortè *Brasilis* ex nom. sp. *Brasiliensis* ab hortul. olim admissio. *Spilanthus fusca* Hort. Par.! *Bidens fusca* Lam. dict. 1. p. 416. An suad. cl. Lamarck et Desf. species distincta? Constans enim à dimidio sæculo in hortis culta servatur, sed caract. excepto colore nulli. (v. v. c.)

31. *S. ALBA* (Willd. sp. 3. p. 1714), caule ramoso adscendente, foliis oppositis et alternis petiolatis ovatis subrepandis, pedicellis terminalibus folio longioribus, capitulis ovatis discoideis, invol. squamis 15-20 oblongo-linearibus, achæniis ad latera ciliatis biaristatis. ① in Peruvîa legit cl. Dombey! *Spilanthus albus* L'Hér. l. stirp. p. 7. t. 4. *Spil. Salivaria* Murr. comm. Gœtt. 1783. p. 3. t. 1. *Bidens ocyimifolia* Lam. dict. 1. p. 416. ill. t. 668. f. 3. Corollæ albæ. Sapor et virtus *Sp. oleraceæ*. Variat in iisdem aut diversis individuis foliis longiusculis aut brevissimè petiolatis. (v. s.)
32. *S. LEUCANTHA* (H. B. et Kunth nov. gen. am. 4. p. 210. t. 370), caule volubili? sexangulato glabro, foliis oppositis petiolatis ovato-oblongis acuminatis basi rotundatis dentatis rariùs integerrimis quintuplinerviis supra margineque scabriusculis, pedicellis terminalibus aut è dichotomiis ortis folio longioribus, capitulis ovatis discoideis, invol. squamis circ. 15 ovatis-lanceolatis, achæniis glaberrimis eciliatis exaristatis. ① in collibus siccis prope Alausi Quitensium. Cor. albidæ 5-dentatæ.
33. *S. URENS* (Jacq. Amer. 212. t. 126. f. 1), caule prostrato aut adscendente basi radicante, foliis oppositis lanceolatis integerrimis glabris basi attenuatis, pedunculis terminalibus folio longioribus glabriusculis, capitulis ovato-globosis discoideis, invol. squamis ovatis sublanceolatis ciliatis, achæniis ciliatis biaristatis. ② in arenosis maritimis Carthagenæ (Jacq.), insulis Caribæis (Plum.), Sancto-Domingo (Bert. 1), in Peruvîa prope Truxillo (H. B. et Kunth nov. gen. am. 4. p. 211). *Bidens angustifolia* Lam. dict. 1. p. 416. et ill. t. 668. f. 2. (var. angustifolia). *Sp. repens* Spr. 1 in h. Balb. *Spil. urens* et verisim. *S. Portoricensis* Spreng. syst. 3. p. 444. *Cotula Spilanthus* Linn. mant. 1. p. 116. *Spilanthus urens* et *Cotula Spilanthus* Willd. Cor. albidæ. Anthere nigricantes. (v. s.)
ℓ. hispidula, foliis utrinquè pedunculisque hirtellis. — circa Limam legit cl. D'Urville. (v. s.)
34. *S. MAURITIANA*, caule repente adscendente, foliis petiolatis ovali-lanceolatis subdentatis utrinquè ramisque subpuberulis, pedunculis folio duplò longioribus, capitulis ovato-globosis discoideis, achæniis glabris calvis. — in insulâ Mauritii (Belang. 1), in graminosis montium excelsorum (Bout. 1), speciatim ad montem Pollicis (Bojer 1). Folii petiolus 3 lin., limbus 6 lin. long., 3 lin. latus. Capitulum parvum. *Acmella Mauritanica* Rich. in Pers. ench. 2. p. 472. (v. s.)
35. *S. PSEUDO-ACMELLA* (Linn. syst. veg. 610), caule erecto subpubero, foliis petiolatis ovalibus lanceolatisve dentato-serratis asperiusculis acuminatis pedunculatis folio triplò longioribus, capitulis ovatis discoideis, achæniis lævibus ciliatis biaristatis. ① in Indiâ orient. frequens. *Verbesina pseudo-Acmella* Linn. sp. ed. 2. p. 1270. mant. 475. *Pyrethrum Acmella* Medic. act. pal. phys. 3. p. 243. t. 19. *Sp. Acmella* Blum. 1 bijdr. 912. *Wight herb.* 2. n. 30. — *Sebastes*. 1. p. 19. t. 10 et 11. *Rumph. amb.* 6. t. 65. *Pluk. alm.* t. 159. f. 4. (v. s. sp. et v. c.)
36. *S. CALVA* (DC. in Wight contrib. p. 19), caule adscendente basi repente apice hirsutulo, foliis petiolatis ovatis obtusiusculis serratis crenatisve glabriusculis basi ciliatis, pedunculis folio triplò longioribus, capitulis ovatis discoideis, achæniis glabris calvis. — in Indiæ orientalis agris Mysorensi et Travacorensi legerunt cl. Heyne et Wight (fl. ind. 2. n. 29!). *Sp. Acmella* Wall. 1 cat. et herb. n. 295. F. H. An à *S. pseudo-Acmella* satis differt? (v. s.)
37. *S. PANICULATA* (Wall. 1 cat. et herb. n. 296), caule erecto ramosissimo glabriusculo, foliis breviter petiolatis ovatis acutis dentatis, pedunculis innumeris folio triplò longioribus, capitulis ovatis discoideis, achæniis ciliatis biaristatis ad utrumque latus setoso-scabris. — in Indiâ orient. ad Martaban legit cl. Wallich. (v. s. comm. à cl. Cur. merc. angl. Ind. or.)
38. *S. RUGOSA* (Blum. 1 in litt. 1831), caule adscendente subpubero, foliis breviter petiolatis lanceolatis parèc et grossè dentatis rugosis supra sparsè setoso-scabris subtus secus nervos subscabris, pedunculis folio quadruplò longioribus, capitulis discoideis ovatis, achæniis glabris calvis. — in Javæ orientalis prov. Bon-

juerangie legit cl. Blume et necnun comm. Hirsutie et foliis rigidioribus obtusioribus grossius serratis à *S. pseudo-Acmellâ* differt. (v. s.)

39. *S. LIXIOCARPA*, caulibus diffuso-procumbentibus, foliis oppositis sessilibus obovato-oblongis integris glabris tenuissimè triplinerviis, pedicellis terminalibus solitariis vix folio longioribus, capitulis globosis subdepressis, invol. squamis ovatis obtusis, achæniis glabris ciliatis inæqualiter biaristatis. ? in Peruvia (Dombey!), ad sinum Chorillo (Macrae!). Recept. paleæ obtusæ. (v. s. comm. à Mus. reg. Par. et Soc. hort. Lond.)

40. *S. EXASPERATA* (Jacq. ic. rar. 3. t. 584), caule dichotomo adscendente basi radicante exasperato glabriusculo, foliis oppositis petiolatis ovatis repandis, pedicellis terminalibus folio longioribus, capitulis ovatis discoideis, invol. squamis ext. 5 lanceolatis puberis, inter. obovatis subconvolutis apice truncatis denticulatis, achæniis ciliatis biaristatis. ① in Venezuelâ. Spil. radicans Jacq. coll. 3. p. 229 non Schrad. Cor. albæ sæpius 4-fidæ. Achæniis superficie utraque ex icone glabra.

β. *Cayennensis*, petiolis sparsè pilosiusculis, foliis ovato-lanceolatis acuminateis, achæniis utrinquè sparsè puberulis. ① in Cayennâ legit cl. Patris. Pars inf. plantæ mihi ignota. (v. s.)

41. *S. ? INSIPIDA* (Jacq. amer. 216. t. 126. f. 2. folium), caule frutescente procumbente ramoso, foliis oppositis obovatis subdentatis sessilibus glaberrimis. 5 in Havanæ rupestribus maritimis. Pedunculi et capitula omninò referant *S. exasperatam* ex Jacq. Cæt. ign. †.

42. *S. BRASILIENSIS* (Spreng. syst. 3. p. 444), caule erecto, foliis ovatis acutis basi attenuatis mucronato-dentatis glabris, pedunculis solitariis, capitulis discoideis. — in Brasiliâ. Car. mancus ferè omnibus speciebus discoideis conveniens. †.

† *Species dubia.*

43. *S. NITIDA* (Lallav. et Lex. nov. veg. mex. 1. p. 28), caule volubili suffruticuloso, foliis oppositis brevè petiolatis ovato-lanceolatis subintegerrimis glabris, corymbis terminalibus axillaribusve, invol. imbricati calyculati squamis ovatis ciliatis, capitulis discoideis, achæniis ciliatis et biaristatis. 5 in Mexico circa S. José del Corral ad vias in montosis. In specim. siccis folia juniora olivaceo et cuprino nitore splendent. — An *Salmeæ* potius species?

†† *Species exclusæ.*

S. ARBOREA Forst. = *Petrobium arboreum*.

S. CROCATÀ Sims. = *Verbesina crocata*.

S. PSEUDO-GUMMIFERA Forst. mss. = *Petrobium arboreum*.

S. TETRÀNDRA Roxb. = *Petrobium arboreum*.

S. TINCTORIA Lour. = *Adenostemma tinctorium*.

CCCLXII. DUNANTIA.

Capitulum multiflorum homogamum, fl. omnibus hermaphr. tubulosis 5-dentatis. Invol. squamæ 1-seriales adpressæ lanceolatæ aut oblongæ. Recept. elongatum. Paleæ exteriores squamis invol. simillimæ, interiores sensim tenuiores submembranaceæ. Styli rami exserti in appendicem filiformem hispidam longè producti. Achænia tetragona gracilia erostria glabra calva. — Herbæ Mexicanæ. Caulis erectus teres pubescens. Rami sæpius oppositi subdivaricati apice nudi. Folia opposita brevissimè petiolata lanceolata basi acuta triplinervia pubescentia integerrima. Capitula ovata ad apices ramorum solitaria bina aut terna, lateralia ubi adsunt subsessilia. Cor. ex sicco albidæ. — Genus affiné *Spilanthi* sed distinctissimum, involucri stigmatibus et achæniis

structurâ, præ cæteris analogiam inter squamas invol. et paleas recept. summam exhibet. — Genus dixi in honorem amic. Phil. Dunant itineris Berlandierani fautoris, qui herbarium ditissimum sollicitè curat et botanicis humaniter communicat !

1. *D. ACHYRANTHES*, caule pube basi confertâ velutinâ, foliis acuminatis, invol. squamis et recept. paleis acuminatis indurato-subspinescentibus. \mathcal{L} ? in Mexico ad Tantoyuca legit cl. Berlandier (pl. exs. n. 2155!) et in Panamaide cl. Hænke! Caulis 1-2-pedalis, habitu ferè *Achyranthis porrigentis*. (v. s.)
2. *D. MICROCEPHALA*, caule vix pubescente, foliis oblongis subacutis, invol. squamis et recept. paleis obtusis. \mathcal{L} in Mexico legit cl. Hænke. Folia pollicem longa, 3 lin. lata. Capitula ovata. (v. s. in h. Hænke à cl. de Sternberg comm.)

CCCLXIII. *XIMENESIA* Cav. ic. 2. p. 60. t. 178. — *Ximenesiæ* sp. Less. syn. 232. — *Ximenesiæ* sp. Kunth.

Capitulum multiflorum heterogamum, fl. radii fœmineis ligulatis 1-serialibus, disci hermaphr. tubulosis 5-dentatis. Invol. circ. biseriale disco ferè longius, squamis patulis angustis acutis foliaceis subæqualibus. Recept. plano-subconicum, paleis amplexantibus membranaceis lanceolatis. Cor. omnium tubus hispidus. Styli disci ramis apice appendiculatis. Achænia radii plano-compressa alata apice profundè emarginata subpilosa biaristata, aristis cum lateribus alarum subconcretis. — Herbæ annuæ pleræque Mexicanæ plûs minûs canescentes. Folia opposita aut alterna in petiolum sæpiûs subalatum et basi auriculatum attenuata, cordata ovata aut oblonga, dentata. Capitula laxè et irregulariter corymbosa. Corollæ luteæ. — Genus affinè *Verbesinæ*, semper enim vidi stylos in ligulis et embryonem in achæniis radii, sed ignoro an radius sit fertilis?

1. *X. ENCELIOIDES* (Cav. ic. 2. p. 60. t. 178), achæniis disci villosiusculis undiquè alâ cinctis, apice emarginatis, radii apteris rugosis, ligulis profundè tridentatis. ① *Pallasia serratifolia* Smith in Rees cycl. v. 26. n. 3. Hûc quatuor stirpes fortè in posterum distinguendas adhûc servo, nempe :
 - a. *hortensis*, foliis infer. cordatis, super. latè ovatis, alâ achæniî pellucidâ, lobis acutis. — Frequens in hortis botanicis sed nunquàm specimen certè spontaneum vidi. (v. v. c.)
 - β. *pachyptera*, foliis infer. cordatis, super. latè ovatis, alâ achæniî crassiusculâ opacâ, lobis acutiusculis. — in Mexico legit cl. Berlandier (pl. exs. n. 2068 et 2286!). (v. s. sp.)
 - γ. *oblongifolia*, foliis infer. ovatis, super. oblongis, alâ achæniî subopacâ parè ciliatâ, lobis subobtusis. — in Senegaliâ ad ripas flum. regionis Walo legit Perrotet, in insulis Mauritanis Sieber (fl. maur. 2. n. 118!) et Belanger, et in hort. Calcuttensi Wallich. (v. s.)
 - δ. ? *cana*, foliis omnibus ovali-oblongis valdè cano-villosis, achæniî villosissimi alis opacis obtusis. — in Mexico circa Laredo legit cl. Berlandier (pl. exs. n. 2068 et 2074). Facile ut species propria admittenda videtur. (v. s.)
2. *X. MICROPTERA*, achæniis disci glabriusculis apice summo tantùm parè alatis et brevissimè aristatis. — circa Buenos-Ayres legit cl. Bacle. Folia latè ovata subtus pubescentia suprâ scabriuscula. Petioli alii nudi, alii ad basin brevè et acutè auriculati. Ligulæ minutè tridentatæ. Achænium sæpè hinc tantùm ad apicem subalatum. (v. s.)

† *Species exclusæ.*

X. CAVANILLESII Spr. = *Echinacea heterophylla*.

X. CORDATA H. B. K. = *Simsia cordata*.

X. FORTIDA Spr. = *Simsia ficifolia*.

X. *HETEROPHYLLA* H. B. K. = *Simsia heterophylla*.

X. *HIRTA* Mart. = *Simsia auriculata*.

X. *HISPIDA* H. B. K. = *Simsia hispida*.

CCCLXIV. *SANVITALIA* Gault. in *Lam. journ. h. nat.* 2. (1792) p. 176. t. 33. *illustr.* (1797) t. 686. *Cav. ic.* 4. (1797) t. 351. *Cass. dict.* 47. p. 294. *Less. gn.* 232. — *Lorentea* *Art. dec.* 4. (1797) p. 42. t. 5 non *Less.*

Capitulum multiflorum radiatum, ligulis fœmineis persistentibus 1-serialibus, fl. disci hermaphr. tubulosis 5-dentatis. Invol. 2-3-seriale. squamis adpressis subimbricatis, inter. paulò longioribus disco æqualibus. Recept. conicum convexumve, paleis oblongis semiamplectentibus. Cor. disci super ovarium articulata pilifera. Styli rami cono superati. Achænia radii grossa triquetra lævia aristis 3 crassis lævibus conicis divergentibus coronata, disci compressa biformia, exter. muricata fere calva, interiora alata ciliata aristellis 2 minimis superata. — Herba Americanæ trichotomæ annuæ sæpius villosæ. Folia opposita ovata integra triplinervia in petiolum ciliatum attenuata. Capitula solitaria terminalia inter ultima folia bracteantia sessilia. Ligulæ luteæ. Discus atro-purpureus, convexusque ut in *Rudbeckia*.

1. *S. PROCUMBENS* (Lam. *journ.* 2. p. 176. t. 33. *ill.* t. 686), caule procumbente aut diffuso, foliis ovatis, ligulis aristas achæniis superantibus, achæniis disci exterioribus alâ continuâ ciliatâ cinctis et aristellis binis superatis. ① in Mexico (Cav. *Ort. Née*), circa urbem (Berl. pl. exs. h. 872). *Sanvitalia villosa* Cav. *ic.* 4. t. 351. *Lorentea atro-purpurea* *Art. dec.* 4. p. 42. t. 5. — *Pluk. phyt.* t. 342. *fig.* antepenult. (v. v. c. et s. sp.)

fl. *oblongifolia*, foliis oblongis, petiolis vix subciliatis, capitulis dimidiò minoribus, ligulis 6-7. — in Mexico ad Chapultepec legit Berlandier (pl. exs. n. 372). An species propria? sed fructus non satis vidi. (v. s.)

2. *S. ACINIFOLIA*, caule procumbente aut diffuso, foliis oblongis, ligulis aristas achæniis superantibus, achæniis disci exter. muricatis, inter. lævibus alâ dentatâ et ciliatâ cinctis vix aristellatis. ① in Mexico legit cl. Alaman. *Habitus* prioris. Ligulæ magis ovales. (v. s.)

3. *S. OCYMOIDES*, caule diffuso erectiusculo, foliis ovatis, aristis radii ligulas superantibus, achæniis disci compresso-tetragonis, exter. muricatis exalatis brevissimè biaristatis, inter. lævioribus subalatis paulò longius aristatis. ① in Mexici prov. Texas legit cl. Berlandier (pl. exs. n. 21021). Herba et folia fere *Ocymi*. Petioli ciliati. Ligulæ minimæ. Achænia longiora quàm priorum. (v. s.)

4. *S. TRAGIÆFOLIA*, caule procumbente aut diffuso, foliis ellipticis, aristis radii ligulas superantibus, achæniis disci fere omnibus compresso-tetragonis calvis exalatis aut vix subalatis. ① in Mexico ad San-Fernando de las Pressas legit cl. Berlandier (pl. exs. n. 22331) et in ditione Oaxacanâ Andrieux (pl. exs. n. 3001). Folia supernè magis scabra quàm priorum.

5. *S. HELIANTHOIDES* (Rich. in Willd. sp. 3. p. 2190), caule erecto, foliis ovato-oblongis apice serratis. — in Peru. Spreng. *syst.* 3. p. 580 excl. *syn.* Capitula magnitudine *Helianthi* gigantei, pedunculata. An certè congener? *Cst.* ign. †

CCCLXV. ANAITIS.

Capitulum multiflorum radiatum, ligulis fœmineis multistriatis subpersistentibus, fl. radii herm. tubulosis.... Invol. campanulatum 2-5-seriale, squamis imbricatis obtusis. Recept. convexum, paleis deciduis complicatis calloso-acuminatis. Styli radii brevissimè exserti, disci.... Achænia glabra minutissimè et parçè muriculata, radii trigona rostrata calva, disci plano-compressa subalata emarginata aut bidentata. —

Suffruticulus Acapulcensis ramis teretibus divaricatis, junioribus pubescentibus. Folia opposita oblonga basi attenuata integerrima pubescentia. Rami apice longè aphylli 1-cephali. Flores radii lutei, disci subfusi. — *Anatidis* deæ pudicitæ nomen selegi ob stylos florum fœmineorum vix exsertos.

1. *A. ACAPULCENSIS*. $\frac{5}{8}$ in Mexico occidentali circa Acapulco legit cl. Hænke. Folia 5-6 lin. longa, 1-1 $\frac{1}{2}$ lata. Pedunculi circ. 5 pollic. (v. s. in h. Hænke à cl. de Sternberg comm.)

CCCLXVI. OLIGOGYNE.

Capitulum pluriflorum heterogamum, fl. radii 1-5 fœmineis ligulatis, disci 6-8 hermaphr. tubulosis. Invol. 1-seriale erectum, squamis 5 ovali-oblongis foliaceis. Recept. planum, paleis lineari-lanceolatis planis. Styli fl. herm. rami exserti hispidi. Achænia obcompressa seu teretiuscula submuriculata glabra aristis 2-4 conicis rigidis superata. — *Herbæ Americanæ* habitu *Sanvitaliæ*, diffusæ, puberulæ. Folia opposita petiolata ovata serrata subtriplinervia. Pedicelli axillares 1-cephali. Capitula parva. Flores flavi. — Genus affine *Sanvitaliæ*.

1. *O. BAHIENSIS*, pedicellis capitulo ferè brevioribus, achæniis teretiusculis. — circa Bahiam Brasiliæ legit cl. Blanchet (pl. exs. n. 1706 l). Caules debiles pedales glabrinusculi. Folia sparse puberula 2 poll. longa, pollicem lata, petiolo 8-10 lin. (v. s.)
2. *O. MEXAPOTAMICA*, pedicellis capitulo brevioribus, achæniis obcompressis. — in Brasiliæ provinciâ Rio-Grande (h. Mus. imp. Bras. n. 874). Valdè affinis *O. Bahiensi*. Caules basi subrepentes adscendentes ad apicem adpressè pubescentes. Involucrum adpressè villosum. Ligulæ 3-4 pallidæ. (v. s. in h. Mus. reg. Par.)
3. *O. TAMPICANA*, pedicellis capitulo multò longioribus, achæniis exter. obcompressis. — circa Tampico de Tamaulipas Mexicanorum legit cl. Berlandier (pl. exs. n. 611). Folia 9-12 lin. longa, 6-9 lata, petiolo 4 lin. longo. Caules 6-9 poll. longi, ramosissimi, adpressè pubescentes. (v. s.)

CCCLXVII. *SYNEDRELLA* Gært. fr. 2. p. 456. t. 171. Rich. in Pers. ench. 2. n. 1901. Cass. dict. 51. p. 470. Less. syn. 233. — *Ucaou* sp. Adams. — *Verbesinæ* sp. Linn.

Capitulum pluriflorum heterogamum, fl. radii circ. 8 subbiseriatis ligulatis fœmineis, fl. disci circ. 8 hermaphr. tubulosis. Invol. oblongum oligophyllum erectum. Recept. parvum planum paleaceum. Ligulæ disco circ. æquales. Achænia radii obcompressa ovalia biaristata aut mutica margine alata, alis laciniatis, disci linearia subtriquetra integra 2-3-aristata. — *Herba Americana* dichotoma. Caulis teres glabriusculus. Folia opposita petiolata ovalia utrinquè attenuata integra parcè villosa. Capitula in axillis ramorum aut foliorum superiorum sessilia subglomerata. Flores lutei.

1. *S. NODIFLORA* (Gært. l. c.). (†) in littore Mexicano prope Campèche (H. et B.), ins. Cubâ (La Sagr.), Jamaicâ (Sl.), Guianâ (Perr.) et verisim. introducta in Senegaliâ (Perr.). *Verbesina nodiflora* Linn. auct. 4. p. 290. Willd. sp. 3. p. 2226. — Dill. elth. 53. 1. 45. f. 53. Sloan. hist. 1. t. 154. (v. s.)

CCCLXVIII. *CALYPTOCARPUS* Less. syn. 221 et in *linnaea* 1834. p. 269.

Capitulum multiflorum heterogamum, fl. radii ligulatis 1-seriatis fœmineis, disci hermaphrod. tubulosis 5-dentatis. Invol. 5-phyllum.

Recept. paleaceum. Achænia conformia erostris plano-obcompressa biartistata interruptè et angustè alata. — Herba Mexicana gracilis dichotoma annua procumbens. Folia decussata petiolata scabra dentata. Capitula solitaria lutea brevissimè radiata pedunculata terminalia demùm alaria. — Genus proximum Heterospermo, Synedrellæ, etc., sed achæniarum cortice crasso donatum.

1. *C. VIALIS* (Less. ll. cc.). ① ad vias prope Jalapam Mexic. Caulis basi radicans. Folia ovata rhombea albo-hirsuta scabra. Invol. squamæ acuminatæ æquales. Paleæ scariosæ floribus breviores.

CCCLXIX. ELECTRA DC. non Panz (1).

Capitulum pluriflorum heterogamum, fl. radii circ. 5 ligulatis fœmineis, disci circ. 8-10 tubulosis hermaphr. 5-dentatis. Invol. duplex, squamis exter. 5 linearibus distantibus, 5 inter. ovali-oblongis submembranaceis basi inter se et ferè cum ext. concretis. Recept. angustum planum, paleis oblongo-linearibus coloratis persistentibus flores ferè superantibus onustum. Cor. disci glabræ abruptè in faucem dilatata. Styli rami cono hispido terminati, fl. radii glabri. Achænia obcompressa plana elliptico-oblonga utrinquè subangustata apice obtusè truncata glaberrimà calva. — Fruticulus Mexicanus glaber erectus. Rami juniores tetragoni, adulti teretes. Folia opposita in petiolum brevem angustata oblongo-lanceolata acuminata grossè serrata penninervia. Capitula pedicellata sæpius terna corymbosa ovali-oblonga. Cor. et recept. paleæ luteæ, undè nomen ad electrum seu succinum alludens. Electra uxor Atlantis alludit etiam ad plantam transatlanticam.

1. *E. MEXICANA*. 5 in Mexico circa Villalpando ultra Guanajuato leger. et mecum comm. cl. Mendéz et Alaman. Etiam habui sine loci design. ex itin. Nezi. (v. s.)

CCCLXX. CHRYSANTHELLUM Rich. in Pers. ench. (1810) 2. p. 471. Cass. dict. 9. p. 150. Less. syn. 234. — Chrysanthellina Cass. dict. 25. (1822) p. 390. — Sebastiania Bertol. opusc. 1822. p. 37 non Spreng. — Collæa Spreng. syst. 3. (1826) p. 622 non DC. 1825. — Anthemidis Bidentis et Verbesinæ sp. Linn.

Capitulum pluriflorum radiatum, ligulis fœmineis 1-serialibus linearibus bidentatis, disci hermaphr. 5-dentatis. Invol. biseriale, ext. squamis 5 paulò brevioribus erectis, int. squamis 10 subcoloratis. Recept. planum, paleis linearibus. Styli fl. herm. ramis in appendicem productis. Achænia calva, exter. oblonga truncata emarginata aut truncata alà minimà crassà cartilagineà utrinquè aucta. — Herbæ annuæ glabræ parvulæ, ramulis subnudis apice 1-cephalis. Folia inter se diversa, radicalia petiolata ovalia basi cuneata, summa oblongo-linearia apice tridentata. Capitula parva. Flores lutei. — Genus ex habitu naturale, ex caract. faciliè dividendum, sed tunc genera tot quot sunt species admittenda, meo sensu invitâ naturâ.

1. *C. PROCUMBENS* (Rich. l. c.), achæniis omnibus glabris lævibus, exter. tertiusculis collo brevissimo superatis, inter. obcompressis truncatis, ① in pascuis subhumidis insularum Caribæarum nempe in Jamaicâ? (Sloan?), Santo-Domingo (Bert.!). Anthemis Americana Linn. sp. ed. 1. p. 895. Bidens apiifolia Linn. avcen. 5. p. 403. Bupthalmum Linn. hort. cliff. 414. Verbesina mutica Linn. sp. ed. 2. p. 1273. Lam. ill. t. 686. f. 2 bona. Chrysanthellina

(1) Electra Panzer = Schismus ex Steud.

- fasciculata et fortè *C. gracilis* Cass. dict. 25. p. 391? *Sebastiania heterophylla* Bertol. opusc. scient. 1822. p. 37. Collæa procumbens Spreng. syst. 3. p. 622. — Sloan. jam. 1. t. 155. f. 1? Plum. ed. Burm. t. 86. f. 2. Variat caulibus erectis procumbentibusve et formâ foliorum. (v. s. comm. à cl. Bertero.)
2. *C. SWARTZII* (Cass. dict. 25. p. 392. sub *Chrysanthellinâ*), achæniis omnibus obcompressis, exter. glabris, inter. ad angulos ciliatis. ① in insulis Caribæis nempe in pascuis subhumidis Jamaicæ (Sw.) et ins. Cubæ circa Havanam (R. de La Sagra!). Verbesina mutica Swartz obs. 314. t. 8. f. 1. Herba variat etiam procumbens diffusa et erectiuscula et foliis polymorphis. (v. s.)
3. *C. SENEGALENSE*, achæniis omnibus obcompressis glabris, exter. tuberculis immarginatis, inter. lævibus alâ marginatis. ① in demissis humidis Senegambiæ circa Casam Senegalem dictam, april. fl. legit cl. Perrottet. *Bidens marginata* Perrottet in litt. 1833. Folia hinc inde opposita multipartita. (v. s. sine flor.)
4. *C. INDICUM*, achæniis omnibus obcompressis apice brevissimè emarginatis, margine callosis glabris, hinc lævibus planis, illinc convexis striatis. ① in Indiâ orient. ad Gojpur et Sukanaghur. *Bidens?* *Bardana* Wall. cat. et herb. n. 323. comp. n. 401. *Achillea?* *Bardana* Hamilt. herb. ex Wall. Ligulæ paucæ parvæ. (v. s. comm. ab hon. Cur. Ind. or. et à cl. Royle.)

CCCLXXI. NEURACTIS Cass. dict. 34. p. 496. Less. syn. 232.

Capitulum pluriflorum heterogamum, fl. radii 3 ligulatis fœmineis, disci hermaphr. tubulosis 5-dentatis. Invol. subbiseriali disco multò brevius, squamis oblongis, exter. brevioribus. Recept. planum medio nudum, paleis ad margines oblongo-linearibus planis. Ligulæ amplæ ultrâ medium trifidæ. Styli fl. herm. rami exserti revoluti in appendicem gracilem longissimam producti. Achænia obcompressa? oblonga glabra apice breviter in collum producta. Pappus nullus aut margo incompletus dentatus. — Herba Javanica glabra diffusa ramosa multifloris. Folia alterna petiolata bipinnatilobata, lobis oblongis integris mucronatis. Capitula pedicellata solitaria. Cor. flavidæ. Ligulæ nervis fuscis lineatæ. Genus *Chrysanthello* valdè (fortè nimis) affîne, sed mihi ignotum.

1. *N. LESCHENAUULTII* (Cass. l. c.). ① in Javâ.

CCCLXXII. GLOSSOCARDIA Cass. dict. 19. p. 62. Less. syn. 233. — Verbesinæ sp. Linn. f. — Zinnia sp. Retz.

Capitulum pluriflorum, fl. radii 1-2 (imò interdum nullis) ligulatis fœmineis, disci circ. 5 hermaphr. tubulosis 4-dentatis. Invol. oblongum, squamis paucis subimbricatis, externis brevissimis. Recept. paleis linearibus deciduis membranaceis inter flores onustum angustissimum. Stylus fl. herm. ex Less. apice appendiculatus. Achænum disci plano-obcompressum in angulis 4 villosissimum, subbialatum, aristas binis rigidis lævibus ex angulis lateralibus ortis coronatum. — Herba Indica annua multicaulis diffusa. Folia alterna basi linearia. Capitula pedunculis nudis brevibus insidentia solitaria flava.

1. *G. BOSWALLEA* (DC. in Wight contrib. p. 19). ① in Indiâ orient. (Kœn. ex Linn. f.), in dumetis aridis circa Madras (Wight!), in Carnatic (Belang.). *Verbesina Boswallæ* Linn. suppl. (1781) p. 379. *Verbesina Boswallia* Willd. sp. 3. p. 2223. Linn. herb. et Roxb. fl. ind. 3. p. 443 ex Wight! *Glossocardia linearifolia* Cass. dict. 19. p. 62. *Pectis meifolia* Wall. cat. et herb. n. 308. Folia non linearia sed lineariloba! Herba esculenta odore et sapore *Fœniculi* (Kœn. in Linn. f.). (v. s. comm. ab ampl. Aulâ merc. angl. Ind. or.)

CCCLXXIII. *HETEROSPERMUM* Willd. sp. 3. (1803) p. 2129. Cass. dict. 21. p. 128. H. B. et Kunth nov. gen. am. 4. p. 245. Less. syn. 233. — *Heterosperma* Cav. ic. 3. (1794) p. 34. t. 267.

Capitulum pluriflorum heterogamum, fl. radii 1-3 ligulatis foemineis, disci 5-15 tubulosis hermaphr. Invol. duplex utrumque oligophyllum, squamis erectis oblongis, exter. ciliatis. Recept. angustum, paleis oblongo-linearibus planis obtusis. Styli rami apice appendiculati. Achænia obcompressa glabra alis incrassatis cincta difformia, radii ovalia erostria calva, disci lineari-oblonga rostrata biaristata, aristis retrorsum aculeolatis. — Herbae Americanæ annuæ oppositè ramosæ. Caules glabri aut bifariam pilosi. Folia opposita dentata lobata aut pinnatipartita glabra. Pedunculi ex dichotomiis aut apicibus ramorum graciles 1-cephali. Cor. flavæ.

1. *H. PINNATUM* (Willd. sp. 3. p. 2129), caule glabro, foliis pinnatipartitis, lobis linearibus acutis integris aut apice dentatis. ① in Mexico (Cav.) circa urbem et in valle Toluacanâ (Berl. pl. exs. n. 576 et 1081!), in prov. Leoninâ ad occid. urbis Guanajuato (Mend.!). *Heterosperma pinnata* Cav. ic. 3. t. 267. (v. s. sp. et v. c.)
2. *H. DIVERSIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 226. t. 384), caule bifariam piloso, foliis aliis obovato-cuneatis indivisis apice crenato-serratis, aliis trifidis. ① in aridis prope urbem Quito et in convalli Guallagambensî. (v. s.)
3. *H. MARITIMUM* (H. B. et Kunth nov. gen. am. 4. p. 245. t. 383), caule bifariam piloso, foliis obovato-cuneatis indivisis apice crenato-serratis. ① in maritimis Peruvîæ prope Truxillo. *Heterosperma ovatifolia* Cav. dem. bot. p. 204? *H. ovatum* Willd. l. c.?

CCCLXXIV. *GLOSSOGYNE* Cass. dict. 51. p. 475 non Less. *Bidentis* sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis, disci hermaphrod. tubulosis 5-dentatis. Recept. planum demum convexum paleolatum. Invol. 2-seriale breve adpressum. Stylus fl. herm. brevis in stigmata 2 hirta longissima productus. Achænia linearia angulata aristas 2 retrorsum setosas gerentia. — Herbae glabræ erectæ, nunc basi suffruticosæ dichotomæ, nunc frutescentes pedunculos plurimos nudos simplices gerentes. Folia alterna ad collum conferta brevia pinnatipartita, lobis linearibus acutis integerrimis, summa pauca parva linearia indivisa. Capitula erecta ebracteata lutea. — Cassinius *Bidentis*, Lessingius *Salmæ* affinem judicant.

1. *G. PINNATIFIDA* (DC. in Wight contrib. p. 19), caulibus dichotomis, achæniis aristis valdè divaricatis. ½ in Indiâ orient. ad Gornepore, Ramispur, etc. legerunt cl. Heyne et Wight. *Bidens pinnatifida* Wall. cat. et herb. n. 300. *Coreopsis*? *Bacana* Heyne! herb. *Zinnia Bidens* Retz obs. 5. p. 28 ex Roxb. fl. ind. 3. p. 435 teste Wight. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et à cl. Wight.)
2. *G. TENUIFOLIA* (Cass. et Less. ll. cc.), caulibus dichotomis, achæniis aristis erectis. ½ in Novâ-Caledoniâ (Labill.!), Novâ-Hollandiâ ad Port-Jackson et insulâ Guam Marianarum (Gaudich.), *Bidens tenuifolia* Labill.! sert. caled. p. 44. t. 45. (v. s. comm. à cl. Labill.)
3. *G. PEDUNCULOSA*, caulibus fruticosis basi simplicibus apice pedunculos plurimos longissimos nudos monocephalos gerentibus, achæniis aristis erectis. ½ in Novâ-Hollandiâ parte tropicâ in collibus rupestribus ad Caput Cleveland legit cl. Cunningham. Caulis crassitie pennæ anserinæ. Folia elongata linearia pinnatipartita, lobis paucis alternis integerrimis. Pedunculi pedales erecti.

Capitulum fructigerum et cor. disci omnino generis sed ligulas in specim. deflorato non vidi. (v. s. comm. à cl. Cunningham.)

N. B. Coreopsis Tannensis Forst. ex Spreng. syst. 614 ad hoc genus pertinet, ex patria ad secundam speciem, ex testimonio Sprengelii (in litt. ad Wall.) ad primam referenda. Glossogyne Chinensis Less. syn. 212 mihi omnino ignota. *Bidens Chinensis* mihi et ut antumo Botanicis cognita, est certe *Bidens* species!

CCCLXXV. DELUCIA.

Capitulum multiflorum radiatum, fl. radii 1-seriatis fœmineis ligulatis, fl. disci hermaphr. tubulosis 5-dentatis. Invol. duplex subæquale, squamis lineari-lanceolatis erectis, exter. foliaceis, inter. coloratis. Recept. planum paleaceum, paleis linearibus. Anth. nigricantes. Styli fl. herm. rami cono superati. Achænia disci et radii obcompressa linearia aristis 3 retrorsum hispidis coronata. — Herba Mexicana subsimplex vix pedalis glabra. Folia opposita petiolata trisecta, segmentis ovatis utrinque acutis grosse dentato-serratis. Capitula solitaria longè pedicellata lutea. — Genus affiné *Bidenti* sed ligulæ fœmineæ fertiles, Narvalinæ sed ligulæ plures, aristæ achæni 3 persistentes et achænia linearia immarginata. Genus montanum dixi in honorem cl. geologi et physici Th. Andr. Deluc, qui laboribus circa barometri et thermometri usum geographiæ botanicæ progressus egregiè paravit.

1. *D. ostruthioides*. ? in Mexici montanis circa urbem legerunt cl. Berlandier (pl. exs. n. 920), Alaman, Mairet, etc. Folia ferè omnino *Imperatoriae Ostruthii* sed opposita. Capitulum è longinquo ferè *Trollii Asiatici*. (v. s.)

CCCLXXVI. *NARVALINA* Cass. dict. 38. p. 17. 59. p. 320. Less. syn. 234. — *Needhamia* Cass. dict. 34. p. 335 non Scop. nec Brown.

Capitulum pluriflorum radiatum, ligulis solitariis (aut 2-3) fœmineis, fl. disci circ. 8 hermaphr. 5-dentatis. Invol. biseriale, exter. breve squamis circ. 3 foliaceis acutis, inter. longius disco brevius squamis circ. 5 oblongis erectis apice coloratis. Recept. planum, paleis squamas inter. referentibus. Ovaria obcompressa ovalia amplè latè marginata ciliata. Pappus caducus biaristatus, aristis crassis fragilibus coloratis triquetro-subulatis secus angulos spinellas strigosas gerentibus. — Frutex Domingensis glaberrimus. Folia opposita petiolulata obovata serrata coriacea. Capitula corymbosa. Flores lutei.

1. *N. Domingensis* (Cass. dict. 38. p. 17). ♂ in Sancto-Domingo. *Needhamia Domingensis* Cass. dict. 34. p. 335. — Ramum huic speciei pertinentem in horto Paris. lectum recepi à cl. Adolph. Brongniart qui cum specimine Cassinii comparavit. Differt à descr. Cass. tantum ligulis 2-3 nec solitariis (v. s.)

CCCLXXVII. *THELESPERMA* Less. in *linnæa* 1831. p. 511. syn. p. 234.

Capitulum multiflorum homogamum. Invol. biseriale utrumque circ. 8-phyllosum, exter. breve squamis lanceolatis vix inter se concretis, inter. campanulatum squamis longè concretis striatis. Recept. planum, paleis oblongis medio nervigeris marginibus latè albo-membranaceis. Styli rami apice appendiculati. Achænia obcompressa-teretiuscula erostrata. Pappus biaristatus, aristis oppositis retrorsum fimbriatis trigono-conduplicatis subulatis persistentibus. — Herba Brasiliensis glabra perennis. Caulis teres oppositè et ideò trichotomè ramosus, rigidus. Folia

opposita biternatim seu-bipinnatim secta, lobis linearibus elongatis acutis integerrimis. Capitula ramos apice nudos terminantia. Cor. obscurè luteæ ex sicco videntur. — Ex invol. gamophyllo subglanduloso et stigmatum structurâ ad Tagetines accedit, sed pappus Helianthearum.

1. *T. SCABIOSOIDES* (Less. l. II. cc.). \mathcal{L} in Brasiliâ (Sellow.!), in provinciâ Rio-Grande (h. imp. Bras. n. 910). Herba bipedalis habitu ferè Scabiosæ columbariæ Gramuntiae sed glabra et tenuius secta. (v. s. comm. à Mus. reg. Berol.)

CCCLXXVIII. ISOSTIGMA Less. in *linnæa* 1831. p. 513. syn. 235.

Capitulum multiflorum radiatum, ligulis fœmineis oblongis 1-serialibus, fl. disci hermaphr. tubulosis 5-dentatis. Invol. campanulatum 2-3-seriale, squamis liberis, exter. brevioribus angustioribus. Recept. planum paleaceum. Styli rami in fl. \mathcal{Q} et etiam \mathcal{Q} in appendicem lineari-filiformem longissimam et hispidam producti. Achænium lineare striatum plano-obcompressum subalatum rostro achænio simili terminatum, alis in mucrones aristasve 2 divergentes læves productis. — Herbæ Brasilienses perennes glaberrimæ. Rhizoma lignosum subhorizontale. Caules erecti nudi 1-cephali simplicissimi. Folia ad collum orta alterna, nunc indivisa, sæpius biternatim seu-bipinnatim secta, lobulis integerrimis. Capitula magna speciosa, in sicco fusca, cocta aquam colore luteo tingentia. — Ligulæ in herbar. sæpè nullæ, verisim. deciduæ.

1. *I. SIMPLICIFOLIUM* (Less. in *linnæa* 1831. p. 513), caule apice esquamato tereti, foliis integris linearibus basi angustatis apice subdenticulatis rariùs trifidis, achæniis aristis latitudinem ovarii æquantibus. \mathcal{L} in Brasiliâ. (Less.)
2. *I. PEUCEDANIFOLIUM* (Less. l. c. p. 514), caule tereti esquamato, foliis multipli-pinnatisectis, segmentis trigonis suprâ canaliculatis, achæniis aristis divergentibus latitudinem ovarii non æquantibus, interdum subnullis. \mathcal{L} in Brasiliâ (Sellow!), in prov. Rio-Grande (h. Mus. imp. Bras. n. 1060!), *Tragoceras peucedanifolium* Spreng. syst. 3. p. 576. Caulis 2-3-pedalis. Capitula 15-16 lin. diam. (v. s. comm. à Mus. reg. Berol.)
3. *I. SPECIOSUM* (Less. l. c. p. 515), caule apice incrassato squamigero, foliis ternatim seu pinnatim sectis bisectisve, segmentis subtrigonis crassiusculis, achæniis aristis vix dimidiam ovarii latitudinem æquantibus. \mathcal{L} in Brasiliâ cum præcedente. (Less.), in prov. Rio-Grande ex h. Mus. imp. Bras. n. 1059. (v. s. in h. Mus. reg. Par.)
4. *I. CRITHMIFOLIUM* (Less. l. c. p. 515), caule tereti esquamato, foliis ternatim biternatimve sectis, segmentis planis striatis, achæniis aristis latitudinem ovarii superantibus. \mathcal{L} in Brasiliâ cum præcedentibus (Less.), in prov. Rio-Grande ex h. Mus. imp. Bras. n. 1058! (v. s. in h. Mus. reg. Par.)

Subtrib. III. *FLAVERIÆ* Less. syn. 233. — Gen. Helianthearum Millenarum Cass.

Capitula nunc uni- vel pauciflora, densè aggregata, alia heterogama fl. uno fœmineo et reliquis hermaphroditis, altera intermixta homogama, nunc multiflora fl. fœmineis pluriserialibus reliquis hermaphroditis. Achænia calva exalata teretiuscula basi longè attenuata. Styli fl. hermaphr. rami sæpius exappendiculati. Recept. in capitulis paucifloris epaleaceum, in multifloris paleaceum. — Herbæ glabriusculæ Austr.-Americanae. Folia opposita tri- seu multiplinervia. Flores flavi.

CCCLXXIX. FLAVERIA Juss. gen. (1789), p. 186. Cass. dict. 17. p. 127. Less. syn. 233. — Vermifuga Ruiz et Pav. syst. p. 210. prod. fl. per. t. 24. — *Milleria* sp. Cav.

Capitula pauciflora glomerato-fasciculata, nunc omnia homogama, nunc heterogama, fl. unico ligulato foemineo, cæteris hermaphrod. 5-dentatis. Invol. 3-4-phyllum connivens oblongum, squamis longitudine æqualibus, unâ exter. majore subconcavâ. Recept. parvum epaleaceum. Styli fl. herm. rami apice exappendiculati. Achænia oblonga calva striata glabra. — Herbæ annuæ Americanæ glabræ aut vix secus ramos puberulæ. Folia opposita sessilia interdum basi attenuata dentata aut integra sæpius trinervia. Capitula secus ramulos unilateralia plûs minus conferta et ideò cymas glomerulosve referentia. Cor. luteæ.

1. F. CONTRAYERBA (Pers. ench. 2. p. 489), erecta, foliis lanceolatis serratis trinerviis basi attenuatis, capitulis confertissimis, omnibus flosculosis aut uniligulatis, ligulâ discum non superante. ① in Amer. austr. regionibus occidentalis nempe in Chili (Fenill.), Peruvîâ (Ruiz et Pav. Domb. Gaudich. etc.), nec non in Mexico inter Fernando et Santander (Berl. pl. exs. n. 2259), Tampico et Real del Monte (Berl. n. 38), et ad occid. urbis Guanaxuato (Mend.), demum in insulâ Cubâ (Rio de La Sagra). Sims bot. mag. t. 2400. *Milleria Contrayerba* Cav. ic. 1. p. 2. t. 4. M. contrahierba Lam. dict. 4. p. 183. illustr. t. 710 f. 4. Vermifuga corymbosa R. et Pav. syst. 210. *Flaveria capitata* Juss. ex Smith in Rees cycl. v. 14. n. 1. *Ethulia Bidentis* Linn. mant. p. 110 ex Smith. — Fenill. obs. 3. t. 14. Herba alexipharmaca et colore luteo pannos inficiens, vulgò *Contrayerba* et *Matagusano* dicta. (v. v. c. et s. sp.)
2. F. ANGUSTIFOLIA (Pers. ench. 2. p. 489), erecta, foliis lineari-lanceolatis sessilibus trinerviis, capitulis corymbosis uniligulatis, ligulâ discum superante. ① in Mexico (Cav. Berl. pl. exs. n. 640). *Milleria angustifolia* Cav. ic. 3. p. 12. t. 223. Variat foliis apice serrulatis et integerrimis, ligulis majoribus et dimidiò minoribus, corymbis fastigiatis terminalibus et etiam ad apices lateraliû ramorum. An fortè duæ species hic confusæ? Flav. integrifolia et Fl. radicans icon. fl. Mex. ined. An icon Cav. 1. t. 4 huc etiam, saltèm quoad florum analyses, pertinet? Sæpè vidi in hortis sub nom. Fl. *Contrayerba* cultam. (v. v. c. et s. sp.)
3. F. LINEARIS (Lag. nov. gen. (1815), p. 33. n. 407), procumbens, foliis sessilibus connato-vaginantibus linearibus integerrimis carnosissimis, capitulis confertè corymbosis uniligulatis. ① in insulâ Cubâ maritimis prope Havanam. Fl. maritima H. B. et Kunth nov. gen. am. 4 (1820), p. 285.
4. F. BONARIENSIS, erecta, foliis sessilibus subconnatis linearibus acuminatis membranaceis tenuissimè trinerviis, capitulis cymoso-corymbosis. ① circa Bonariam legit cl. Bacle. Flores ign. (v. s.)

† *Species exclusæ.*

F. PERUVIANA Juss. = *Piqueria artemisioides*.

F. REPANDA Lag. = *Broteroa trinervata*.

CCCLXXX. BROTEROA. — *Brotera* Spreng. in *Schrad. journ.* 1800. p. 2. p. 186. t. 5. Pers. ench. n. 1959. Cass. dict. 34. p. 304. 59. p. 236 non alior. (1). — *Nauemburgia* Willd. sp. 3 (1803), p. 2393. pessimè! *Hoffmans. verz.* 1824. p. 237. Less. syn. 232 non Mærch (1802). — *Flaveria* sp. Spreng. (1826). Lag.

Capitula in glomerulum ovatum densè spicato-conferta sessilia inter se varia, alia biflora heterogama seu homogama, alia 1-flora foemi-

(1) *Brotera* Cav. (1799). = *Pentapteris* sp. — *Brotera* Willd. (1803). = *Cardopatum*. — *Brotera* Spreng. (1802). = *Hyptis* sp.

nea aut hermaphrodita. Invol. squamæ nunc solitariæ concavæ amplæ, nunc 2-3, alteræ amplæ concavæ, alteræ parvæ. Recept. punctiforme nudum. Cor. tubulosæ extus pilis articulatis confertis hispidæ 5-fidæ, fœmineæ filiformes subligulatæ? Styli fl. herm. ramis apice exappendiculatis. Achænia obcompressa obovato-oblonga glabra calva, fl. herin. minora quàm fl. fœm. ex Cass. — Herba Austr.-Americana glabra, ramis paucis divaricatis. Folia opposita sessilia ovali-lanceolata acuminata trinervia integerrima serrata aut subrepanda. Glomeruli ad apices ramorum aut axillas super. pauci subumbellati. Cor. flavæ. — Nomen paulò mutavi ut melius cum nomine ill. viri congruat et à diversis homonymis distinguatur.

1. *B. TRINERVATA* (Pers. ench. 2. p. 498). ①? 2l? in America austr. circa Caracas legit cl. Vargas, in Mexico ubi dicitur *Azumiatas* (Lag.). Brotera Contrayerva Spreng. in Schrad. journ. 1800. p. 2. p. 186. t. 5. excl. syn. Cav. et Feuill. Brotera Sprengelii Cass. dict. 34. p. 304. Naumburgia trinervata Willd. sp. 3. p. 2393. Flaveria repanda Lag. nov. gen. p. 33 ex Spreng. syst. 3. p. 500. neu. entd. 3. p. 42. Herba basi indurata ferè lignosa et verisim. perennis, etiam ex hortis Europæis annua dicatur. Variat foliis terno-verticillatis. (v. s. sp. comm. à cl. Vargas.)

CCCLXXXI. CLAIRVILLEA.

Capitula fasciculato-capitata pluriflora heterogama, fl. radii. ligulatis circ. 5 uniserialibus fœmineis, fl. disci 4-5 tubulosis hermaphr. 5-dentatis. Invol. ovatum, squamis ovatis arcte imbricatis. Recept. angustum paleis oblongo-linearibus in margine onustum, centro epaleaceum. Styli fl. herm. rami apice appendiculati hispiduli. Achænia ovata glabra calva. — Frutex Austr.-Americanus. Rami teretes, juniores densè albo-tomentosi, adulti glabri. Folia opposita connato-vaginantia ovato-lanceolata acuminata serraturis distantibus callosospinellosis serrata, suprâ 5-nervia, juniora araneosa, adulta rugosa glabriuscula, subtus densè albo-tomentosa. Capitula subsessilia in glomerulum ovatum terminalem densè aggregata, centralia præciora. — Genus Brotera affine sed ligulis circ. 5, styli ramis appendiculatis et disposit. capitulorum faciliè diversum. Dixi in memoriam cl. de Clairville Galli botanici et entomologici de historiâ naturali Helveticâ benè meriti.

1. *C. QUINQUENERVIA*. ♂ in Amer. australi legit ill. Dombey, sed locus in herb. non design. verisim. in Peruvîâ (v. s. comm. à Mus. reg. Par.)

CCCLXXXII. ENHYDRA. — Enydra Lour. fl. coch. (1790), ed. Willd. 2. p. 624. — Meyera Schreb. gen. (1791), n. 1318. Less. syn. 236. — Sobrya R. et Pav. fl. per. prod. (1794) p. 109. t. 24. — Enydra, Meyera et Sobrya Pers. ench. n. 1964, 1896 et 1966. — Hingcha Roxb. (1814) sine descr. — Tetraotis Reinw. (1824), in Blum. bijdr. 892. — Phyllimena Blum. in litt. — Meyeræ sp. Brown. — Enydræ sp. Cass.

Capitulum multiflorum heterogamum, fl. omnibus paleâ complicatâ involutis, ext. pluriserialibus fœmineis, int. hermaphr. sterilibus. Recept. parvum planiusculum. Paleæ quasi bivalves, subcoriaceæ arcte conniventes apice hirtellæ. Invol. 4-phyllum, squamis foliaceis, 2 oppositis majoribus. Cor. tubo radii filiformi, disci subconico, limbo radii subligulato 3-4-fido, disci infundibuliformi 5-fido. Stigm. 2 exserta revoluta versùs apicem hispidula. Achæmium obcompressum calvum erostre paleâ omninò involutum. — Herbæ Asiaticæ et Ameri-

canæ fortè etiam Australasicæ, aquaticæ, radicantes, oppositifoliæ, *Cryphiospermum* referentes. Folia oblonga interdum biauriculata. Capitula ad axillas superiores solitaria. Flores albi (an in omnibus?)

* Asiaticæ, foliis brevissimè petiolulatis.

1. *E. FLUCTUANS* (Lour. fl. coch. 2. p. 625), tota glabra, foliis subhastatis serratis brevissimè petiolatis, capitalis pedunculatis. — in aquosis paludibus Cochinchinæ. *Meyera fluctuans* Spreng. syst. 3. p. 602. excl. syn. R. et Pav. (v. s. ?)
2. *E. LONGIFOLIA*, caule glabro apice summo pubernulo, foliis basi subhastatis vix petiolatis distanter et minutè serrulatis glabris margine scabris, capitulis sessilibus. $\frac{1}{2}$ in paludosis ins. Javæ prope Bataviam. *Tetraotis paludosa* Reinw. in Blum. ! bijdr. 892. *Phyllimena* Blum. ! in litt. Folia lineari-lanceolata, 2 poll. longa, 3 lin. lata. (v. s. in h. Blum.)
3. *E. PALUDOSA*, caule hispido, foliis basi subhastatis vix petiolatis grossè serratis asperis, capitulis sessilibus. $\frac{1}{2}$ in paludosis ins. Javæ prope Bataviam. *Tetraotis paludosa* Reinw. in Blum. ! bijdr. 892 ! *Phyllimena paludosa* Blum. ! in litt. Capitula interdum terminalia. Folia prioris sed pilis scabrida et grossius serrata. (v. s. in h. Blum.)
4. *E. HELONCHA*, caule basi glabro, apice hirtello, foliis basi subhastatis vix petiolatis serratis glabris, capitulis sessilibus. $\frac{1}{2}$ in aquosis India or. in prov. Silhet (Wall.), ad Dumdumma et Golpara (Ham.). *Meyera Heloncha* Wall. ! cat. n. 3195. comp. 305. *Hingcha repens* Roxb. cat. calc. 62. Duabus prioribus valdè affinis. Vernac. dicitur *Heluncha* et *Helumochika*. (v. s.)

** Americanæ, foliis sessilibus.

5. *E. SOBRYEA*, glabra, foliis sessilibus basi sagittatis lanceolatis serratis, capitulis. . . $\frac{1}{2}$ in Peruviz stagnosis. *Sobryea sessilifolia* fl. per. syst. 1. p. 197. *Sobrya sessilifolia* Pers. ench. 2. p. 473.
6. *E. OBLONGA*, erecta glabra, foliis sessilibus oblongis basi rotundatis obscure dentatis, capitulis breviter pedunculatis. $\frac{1}{2}$ in Peruviz stagnosis. *Sobryea oblonga* fl. per. syst. 1. p. 197. *Sobrya oblonga* Pers. l. c. (v. s. comm. ex herb. Dombey à Mus. reg. Par.)
7. *E. SESSILIS*, caule erecto glabro sub foliorum jugis pubernulo, foliis sessilibus oblongis basi rotundatis apice subacutis parcè serratis glabris, capitulis sessilibus. $\frac{1}{2}$ in aquosis Jamaicæ australis (Sw.), Sancti-Domingi (Bert.!). *Eclipta sessilis* Swartz prod. 114. *Meyera sessilis* Swartz fl. Ind. occ. 1364. Spreng. syst. 3. p. 602. Variat foliis subhirsutis ex Sw. (v. s.)
8. *E. MARITIMA*, glabra, foliis oblongis acutis basi rotundatis obsolete serratis, subcarnosis, capitulis sessilibus. $\frac{1}{2}$ in littore Oceani pacifici prope El Callao de Lima (H. et B.). *Meyera maritima* H. B. et Kunth nov. gen. am. 4. p. 269. Willd. herb. ex Less.
9. *E. SUBCUNEATA*, demissa glabra, foliis sessilibus oblongis basi cuneato-atenuatis apice vix acutis obtusè et parcè dentatis, capitulis ad axillas solitariis sessilibus. $\frac{1}{2}$ in fossis circa Rio-Janeiro speciatim ad viam Brocco leger. cl. Lund et Gandichaud. A quatuor prioribus facilè differt foliis basi nec sagittatis nec auriculatis nec obtusis. An fortè *E. maritima* Less. in linnzæ 1831. p. 159 à cl. Chamisso ad fretum Sanctæ-Catharinæ lecta, huc ex loco natali et foliis oblongo-obovatis referenda? (v. s.)

† *Species exclusæ.*

ENYDRA CÆSULIOIDES Cass. = *Cryphiospermum repens*.

MEYERA GUINEENSIS Spr. = *Cryphiospermum repens*.

MEYERA CAPITATA Spreng. quæ *Tilesia capitata* Mey. esseq. 251. omninò aliena.

TETRAGONOTHECA L'Hér. etiam à Spreng. admissa est omninò diversa.

Subtrib. IV. *TAGETINÆ*. — Tagetineæ *Cass. dict.* 20. p. 367. 59. p. 61. *ann. sc. nat.* 1829. p. 12. *opusc.* 3. p. 54. — Tagetineæ, Pectideæ et pars Chrysocomearum *Less.*

Capitula multiflora, nunc heterogama fl. radii fœmineis ferè semper ligulatis, nunc homogama discoidea. Invol. nunc uniseriale squamis plùs minùs concretis, nunc pluriseriale paleis exterioribus bractæformibus liberis interioribus concretis. Recept. planum epaleaceum. Achænium erostre striatum elongatum basi attenuatum. Pappus aristis squamellis setisve constans. — *Herbæ* interdum suffruticosæ sæpiùs annuæ Americanæ glandulas pellucidas grossas in contextu foliorum et involucrorum gerentes et ideò sæpè odoriferæ. Folia opposita aut alterna. — Pectideæ à cl. Cassini hùc fortè non immeritò adnexæ, ad Vernoniaceas à cl. Lessing rejectæ, nexum inter duas tribus indicant.

Div. I. *TAGETÆ*. — Tagetineæ, Dyssodieæ et Archetypæ *Cass. dict.* 59. p. 61. — Tagetineæ et pars Senecionearum *Less. syn.* p. 236 et 388.

Capitula sæpiùs heterogama radiata. Invol. squamæ saltèm interiores inter se concretæ. Cor. disci regulariter 5-dentatæ. Styli rami cono superati. Pappus inæqualis 1-2-serialis aristis squamellisve constans.

CCCLXXXIII. *ADENOPHYLLUM Pers. ench.* 2. (1807) n. 1870. — Willdenow *Cav. ic.* 1. p. 61. t. 89 non *Thunb.* — *Schlechtendalia Willd. sp.* 3 (1803) p. 2125 non *Less.* 1830. nec *Spreng.* 1829 (1). — *Dyssodæ sp. Lag.* — *Tagetis sp. Less.*

Capitulum radiatum. Invol. squamæ lineares concretæ apice productæ in aristam subulatam basi 1-glandulosam, extùs cinctum bracteis sub-1-serialibus aristato-subulatis basi uniglandulosis. Recept. alveolatum seu fimbrilliferum. Achænia elongata. Pappus duplex, ext. paleæ breves truncatæ, inter. elongatæ apice acutè subtrifidæ. — *Herbæ Mexicanæ Tagetis facie.* Folia opposita et superiora alterna pinnatisecta, segmentis ovatis aut ferè linearibus aristato-dentatis, rachi inter segmenta aristas hinc indè gerente. Pedunculi 1-cephali sub capitulo incrassati. Cor. crocæ seu coccineæ, in disco dilutiores.

1. *A. COCCINEUM* (Pers. l. c.), foliorum segmentis aliis ovatis dentatis, aliis intermediis capillaceo-subulatis. — in Mexico juxta Doscaminos oppidum legit *Née* (*Lag.*). Willdenow glandulosa *Cav. ic.* 1. p. 61. t. 89. *Lam. ill.* t. 685. *ic. Cav. Schlechtendalia glandulosa Willd. sp.* 3. p. 2125. *Tagetes punctata fl. Mexic. ic. ined.* *Bœbera Cavanillesii Spreng. syst.* 3. p. 544 ex hæc et *Dysodia porophylla* mixtis facta videtur. *Dyssoda Cavanillesii Lag. elench. h. Madr. p.* 29. n. 371. Variat statu abnormali nempe floribus disci 6-8-fidia et stylo trifido, sed potius monstrum quàm varietas et perperam *Cavanillesius* hanc in caract. gener. recepit. Confer *Lag.* et *Cass. ll. cc.*

2. *A. CAPILLACEUM*, foliorum lobis omnibus capillaceo subulatis, aut in lobulos subulatos ad basin usque fissis. — in Mexico. *Schlechtendalia capillacea fl. Mex. ic. ined.* An prioris var.? (v. *ic. pict.*)

(1) Stricto jure nomen Willdenovianum servandum esset, sed Personianum admisi, ne augeatur synonymia et ne botanici peritissimi nomen generi subdubio sit consecratum.

CCCLXXXIV. *LEBETINA* Cass. dict. 25 (1822), p. 394. 59. p. 68. — *Bœbera* Less. syn. 237 non Willd. — *Dyssodia* Willd. enum. 900 non Cav.

Capitulum radiatum. Invol. cylindraceo-campanulatum 1-seriale, squamis circ. 20 internè concretis apice variè liberis, bracteis 1-2-seriatis pinnatipartitis lobis apice subulato-piliformibus cinctum. Recept. convexum fimbriiferum. Achænium cylindraceum minutissimè pubescens. Pappus duplex, ext. paleis 10 oblongis integris, int. 10 setis basi cuneato-paleatis apice ramoso-pinnatis filiformibus. — Herba Mexicana glabra, Tagetis seu *Dyssodiæ* facie, erecta ramosa. Folia alterna pinnatipartita, lobis serratis, serraturis in pilum longum desinentibus. Rami apice subnudi, foliis supremis paucis in bracteas desinentibus, 1-cephali. Cor. flavo-aurantiacæ.

1. *L. CANCELLATA* (Cass. dict. 25. p. 394). ① in Mexico ad occid. urbis Guanaxuato legit cl. Mendez, circa Tamanlipas cl. Berlandier (pl. exs. n. 2140!), et ad Tlapujahua F. W. Keerl (h. Mart.!). *Dyssodia* Porophyllum Willd. enum. p. 900 non Cav. *Bœbera* Porophyllum Less. syn. 237. excl. syn. Habitus *Dyssodiæ* porophyllæ, sed ligulæ multò majores invol. superantes, recept. magis convexum longius fimbriiferum et pappus duplex! (v. s.)

CCCLXXXV. *DYSODIA* DC. non Lour. (1). — *Dyssodia* Cav. anal. scienc. nat. 6. (1802) p. 334. Cass. dict. 25. p. 396. 59. p. 61. Less. syn. 388. — *Bœbera* Willd. sp. 3. (1803) p. 2125. Cass. dict. 5. suppl. p. 2. non Less. — *Dyssodiæ* sp. Lag.

Capitula radiata aut rarissimè discoidea. Invol. proprium 1-seriale, squamis plùs minùs inter se concretis, rariùs nudum, sæpè extùs bracteis 1-2-seriatis calyculatum. Recept. planum brevissimè hirsuto-fimbriiferum aut subalveolatum. Styli rami cono pubero superati. Achænium elongatum tetragono-compressum. Pappus 1-serialis, paleis imà basi indivisis et in pilos setaceos scabros aut barbellato-palmatim fissis, et ideò quasi polyadelphus dicendus. — Herbae Mexicanæ Tagetis facie. Folia opposita aut rariùs alterna, indivisa aut sæpiùs pinnatipartita, serraturis acutis aut in pilos abeuntibus.

Sect. 1. EUDYSODIA.

Involucrum bracteatum, bracteis pinnatipartitis, lobis in pilos abeuntibus. Recept. densè hirsuto-fimbriiferum. — Folia alterna pinnatisecta.

1. *D. POROPHYLLA*, erecta glabra, foliis alternis pinnatipartitis, lobis extrorsum latoribus dentatis, dentibus pilo longo apiculatis, invol. bracteis pinnatipartitis apice setigeris obvallato, ligulis invol. vix longioribus aut nullis. ① in Mexico. *Dyssodia* porophylla Lag. elench. h. madr. 29. *Dyssodia* glandulosa Less. syn. 388. excl. syn. *Bœbera* Porophyllum Kunth nov. gen. am. 4. p. 198. (excl. Willd. syn.) Spreng. neu. entd. 3. p. 44. Habitus *Lebetinæ* sed pappus omninò diversus et ligulæ multò minores oblongæ aut nullæ.

a. radiata, ligulis oblongis involucrum paulò superantibus. Dyss. porophylla Cav. anal. 6. p. 334. *Bœbera* alternifolia fl. mex. ic. t. 636. (v. v. c.)

β. discoidea, ligulis nullis. *Pteronia* Porophyllum Cav. ic. 3. p. 13. t. 225. Willd. sp. 3. p. 1782. (v. s. sp. in pl. exs. Berl. n. 957.)

(1) *DYSODIA* Lour. = *Serissa*.

Sect. II. BOEBERA. — *Boebers Willd. Pers.*

Invol. bracteatum, bracteis oblongis linearibusve indivisis. Recept. brevissime fimbriatiferum aut subnudum. — Folia opposita pinnatisecta.

2. *D. CHRYSANTHEMOIDES* (Lag. elench. h. madr. p. 29. n. 364), adscendens aut diffusa subpuberula, foliis oppositis subpinnatisectis, lobis linearibus apice dentatis, capitulis solitariis, pedunculis foliosis, invol. 8-phyllo squamis basi concretis constante bracteis circ. 8 linearibus basi ciliatis cincto, ligulis pauci involucri paulò superantibus. (Y) in Americà frequens ad ripas flum. Missouri et Mississipi (Pursh), in Mexico (Alam.! Mair.!), ad Santa-Fe Bogotensium ubi ex odore dicitur *Ruda* (Lag.). *Tagetes papposa* Vent. l. h. cels. t. 36. Michx. fl. 2. p. 132. *Tagetes pumila* hortul. *Dysodia glandulosa* Cav. dem. bot. 202. *Boebers chrysanthemoides* Willd. sp. 3. 2125. *Boebers glandulosa* Pers. ench. 2. p. 459. Variat caule adscendente aut diffuso, glabriusculo aut puberulo, foliis variè incisis angustissimis aut sublatisioribus, bracteis plàs minus ciliatis. An fortè species plures hic confusæ? in omnibus video recept. nudum aut vix alveolatum, achænia compressa tetragona glabriuscula pappo longiora. (v. v. c. et s. sp.)
3. *D. FASTIGIATA*, erecta puberula, foliis oppositis pinnatipartitis, lobis linearibus apice serratis, capitulis corymbosis, pedunculis 1-cephalis subaphyllis, invol. 8-phyllo bracteis 6-8 indivisis cincto, ligulis circ. 8 invol. paulò longioribus. (Y) in Mexici montanis (Alam.!) prope Rio Pazcuaro et lacum Cuiseo (H. et B.). *Dysodia tagetiflora* Lag. elench. h. madr. p. 29? *Boebers tagetiflora* Spreng. l. c.? Valdè accedit ad *D. chrysanthemoides*. Fæniculum redolet ex Kunth. (v. s. comm. à cl. Alaman et Berl.)
4. *D. PUBESCENS* (Lag. elench. h. madr. p. 29. n. 365), caule pubescente, foliis pinnatisectis, lobis lineari-cuneatis apice acutè 3-5-inciso-serratis cæterum integerrimis, pedunculis 1-cephalis foliosis, invol. squamis 8 basi concretis bracteis 6-8 indivisis ciliolatis cincto. — in Mexico (Lag. Hænke!), in Leon prope Guanaxuato (Mend.!). *Aster pinnatus* Cav. ic. 3. p. 6. t. 212. *Boebers pubescens* Spreng. syst. 3. p. 544. Variat caule patulo et erecto. Species *D. chrysanthemoides* proxima, sed lobis latioribus subcuneatis faciliè distincta. (v. s.)
5. *D. INCAUA*, caule erecto hirsuto, foliis pinnatisectis infernè subhirsutis, lobis linearibus acutis canaliculatis, aliis integerrimis, aliis subcuneatis trifidis, terminali à cæteris non distincto, pedunculis 1-cephalis. — in Mexico (Lag. Alam.!), circa urbem (Berl.!) pl. exs. n. 631. *Boebers incaua* Lindl. bot. rep. t. 1602. opt. *Dysodia subintegerrima* Lag. elench. p. 29. n. 366. *Bœb. subintegerrima* Spreng. l. c. Nomen Lag. vetustius sed decipiens et subdubium. Pappus fusco-nigricans (v. s.)
6. *D. APPENDICULATA* (Lag. elench. h. madr. p. 29. n. 368), glabra, foliis oppositis pinnatisectis, lobis 5-7 ovato-sublanceolatis inæqualiter argutè serratis glandulâ oblongâ apice appendiculatis, invol. bracteato. (Y) in Mexico prope Acapulco legit. cl. Née et ad occid. urbis Guanaxuato cl. Mendez. Pappus *D. chrysanthemoides*. Invol. *D. porophyllæ* si cilia exterioris exceperis. (v. s. comm. à cl. Mendez et Alaman.)

Sect. III. BOEBEROIDES.

Invol. bracteatum, bracteis ovatis acutis subseratis. — Folia opposita indivisa.

7. *D. GRANDIFLORA*, erecta glabra, foliis oppositis ovali-lanceolatis argutè serratis acuminatis, capitulis ad apices ramorum solitariis, involuacro bracteis ovatis subdentatis acuminatis cincto, ligulis invol. multò longioribus, achænio pubescenti-villoso. — in Mexico. *Boebers grandiflora* ic. fl. mex. ned. t. 635. Paulò refert *Zinniam coccineam*. (v. s. in herb. Hænke. à cl. de Sternberg comm.)

Sect. IV. GYMNOLENA.

Invol. ebracteatum omnino nudum. Recept. subalveolatum. — Folia opposita indivisa.

8. *D. serratifolia*, erecta glaberrima, foliis oppositis indivisis oblongo-lanceolatis utrinque attenuatis argutè serratis minutè punctatis, capitulis longè pedicellatis corymbosis, invol. nudo, ligulis oblongis tubulosisque involucri ovali-oblongo longioribus, achænio adpressè villosa. — in Mexici prov. Oaxaca ad Atitla et locos vicinos. Habitus ferè Tagetis lucidæ, sed folia ciliata et pappus diversissimus. Recept. vix subalveolatum. (v. s. comm. à cl. Alamani.)

Sect. ? v. ACIPHYLLÆ.

Invol. nudum inter folia suprema acerosa subbracteiformia sessile. Recept. minutè alveolatum. — Folia opposita integerrima. — Habitu à genere valdè recedit sed cum caract. generico quadrat.

9. *D. acerosa*, suffruticosa ramosissima erecta aut subdescendens glabra, foliis oppositis distinctis subulatis acutis, capitulis inter folia suprema subbracteantia sessilibus solitariis, involucri disco breviorè obtusè 8-dentato, ligulis 5-6 oblongis, achænio glabro. § in Mexici prov. Sancti-Ludovici de Potosi decemb. fl. legit cl. Berlandier (pl. exs. n. 13401). Fruticulus vix semipedalis. Folia 3-4 lin. longa erecta rigidula integerrima. Invol. oblongo-cylindricum subcoctatum. Pappus sordidè subgriseus. (v. s.)

† *Species exclusa.*

BORBERA POROPHYLLUM Less. = *Lebetina cancellata*.

CCCLXXXVI. *CLOMENOCOMA* Cass. dict. 9. p. 416. v. 59. p. 61 et 66. — *Bartolina* Adans. fam. 2. p. 24 (ex car. et syn. Houst.) non Broom.

Capit. multiflorum radiatum, ligulis 1-serialibus fœmineis, fl. disci tubulosis 5-dent. hermaphr. Invol. imbricatum, squamis linearibus acutis dorso versùs apicem glandulam grossam gerentibus. Recept. tenuiter fimbriatiferum. Achæmium elongatum multistriatum glabriusculum. Pappus squamellis 10 uniseriatis basi polyadelphis supernè lineari-setiformibus. — Herbæ Austr.-Americane Tagetis facie. Folia opposita pinnatisecta. Capitula terminalia pedunculata solitaria. Flores aurantiaci. — Genus recognoscendum. Differt à *Dysodiæ* involucri imbricato nec uniseriali.

1. *C. aurantia* (Cass. dict. 9. p. 416. 59. p. 66), foliis ad basin petioli lobulos setaceos utrinque gerentibus, segmentis bijugis cum impari ovalibus acutis serratis. — in Verà-Cruce (Houst.). *Aster aurantius* Linn. sp. 1232. *Inula aurantia* Spreng. syst. 3. p. 524. — *Aster*, etc. Houst. reliq. t. 18. Caulis dicitur basi radicans.

† *Species dubia.*

2. *C. ? pinnata*, foliorum segmentis multi- (5-7) jugis, omnibus æqualibus linearibus basi integris apice latioribus inciso-dentatis. — in Novâ-Hispaniâ, prope Hapam. *Aster pinnatus* Cav. ic. 3. p. 6. t. 212. Habitus prioris sed pappus videtur setis paucis simplicibus constans. Fortè genericè separanda, sed *Clomenocoma* certè multò magis affinis quàm *Asteri*. †

CCCLXXXVII. *HYMENATHERUM* Cass. bull. philom. 1817 jan. et 1818 decemb. dict. 22. p. 313. Less. syn. 236.

Capitulum multiflorum radiatum, ligulis 10 fœmineis ovalibus suprâ velutinis. Invol. turbinatum campanulatum disco brevius squamis

10-14 uniseriatis concretis obtusis grossè glandulosis constans. Recept. planum omnino nudum. Cor. subinæqualiter 5-dentatæ. Styli rami elongati. Achænia gracilia striata. Pappus paleis 10 rarius 5 subuniseriatis infernè indivisis membranaceis supernè trifidis, lobis nunc omnibus rigidulis aristatis scabris, nunc lateralibus membranaceis medio aristato, nunc alternè trifidis aristatis, alternè oblongis membranaceis. — Herba annuæ, omnes fortè Chilenses; diffusæ. Caules angulati. Folia saltè inferiora opposita. Capitula ramos terminantia solitaria verisim. lutea.

§. 1. *Pappo 1-seriali, foliis pinnatisectis.*

1. *H. P. Næxi*, foliis infer. oppositis, super. alternis pinnatipartitis, lobis filiformibus integerrimis, pappi paleis 10 basi scariosis trifidis, lobis lateralibus membranaceis, medio elongato aristæformi scabro. ① in Chili? legit cl. Née. Herba vix 5-6-poll. apice ramosa, ramis foliosis 1-cephalia. Invol. campanulatum cinctatum 12-14-dentatum. Flosculi 70 quorum 10 ligulati. An potius *Dysodia* species? (v. s.)
2. *H. tenuifolium*, caule herbaceo duriusculo erecto aut adscendente ramoso, foliis pinnatipartitis subpuberulis, lobis subulatis acutis integerrimis, pedunculis elongatis 1-cephalis subnudis, pappi 1-serialis squamellis 10, omnibus basi submembranaceis apice trifidis, lobo medio setiformi scabro, lateralibus brevibus submembranaceis. ① in Mexico circa Bejar legit cl. Berlandier (pl. exs. n. 2063!). Etià habeo sine loci indic. ex itin. Næxi. *H. tenuifolium* Cass. dict. 22. p. 313 dictum cum dubio è Chili ortum, huc fortè pertinet. (v. s.)
3. *H. BERLANDIERI*, caule suffruticuloso erecto ramosissimo, foliis confertis puberulis pinnatipartitis, lobis subulatis acutis integerrimis, pedunculis elongatis 1-cephalis squamulas paucas gerentibus, pappo 1-seriali, squamellis alternè 5 ovali-oblongis obtusis membranaceis, 5 duplè longioribus in aristas scabram productis basi margine scariosis. ② in Mexico inter Santander et Vittoria novemb. fl. legit cl. Berlandier (pl. exs. n. 2253!). Simillimum *H. tenuilobo*, sed pappo, facillè distinctum. (v. s.)
4. *H. PENTACTETUM*, caule suffruticuloso erecto ramoso, foliis pinnatipartitis puberulis demum glabratiss, lobis subulatis acutis, infer. brevioribus approximatis, pedunculis elongatis 1-cephalis subnudis, pappi squamellis 5 basi submembranaceis subtrifidis, lobis lateralibus brevibus aut nullis, medio setiformi elongato scabro. ③ in Mexico circa Monterey legit cl. Berlandier jan. flor. (pl. exs. n. 1382!). Facillè cum *H. Berlandieri* et *tenuilobo* primâ fronte confundendum, sed pappo 5- nec 10-seto distinctum. (v. s.)

§. 2. *Pappo biseriali, foliis indivisis.*

5. *H. P. KÖNTZII* (Less. syn. 236), foliis in vaginam connatis linearibus integerrimis pilosiâculis. ① in Chili. Herba gracilis uliginosa paucè-poll. earis simplex aut parùm ramosa. Folia decussata, 6-9 lin. longa, vix semilineam lata. Invol. 6-8-dentata, dentibus ciliatis subæqualibus. Confer cum *Coilopodio* Chilensi.

COCLXXXVIII. TAGETES *Tourn. inst. 488. t. 278. Linn. gen. n. 964. Carn. fr. 2. p. 437. t. 172. Less. syn. 236.* — *Tagetes*, *Diglossus* et *Eulcidea* Cass. dict. 59. p. 62 et 69-70.

Capitulum radiatum aut rarissimè subdiscoideum. Invol. squamæ 1-seriales concretæ in cupulam campanulatam cylindricamve apice dentatam. Recept. subalveolatum. Achænia elongata basi attenuata, compresso-tetragona. Semen in achænio pedicellatum. Pappus simplex, paleis inæqualibus, aliis obtusis subconcretis, aliis aristatis elongatis liberis. — Herba annuæ Americanæ sæpius foetidæ rarò suaveolentes.

glandulis in foliorum laminis subrotundis pellucidis et in invol. linearibus plūs minus obsitæ. Folia opposita aut alterna, indivisa serrata aut pinnatisecta, segmentis serratis integerrimisve. Capitula solitaria aut fasciculata plūs minus pedicellata. Cor. crocæ seu flavæ. Ligulæ sæpè (præsertim in capitulis plenis) bilabiatae.

§. 1. Pectidiformes, *foliis indivisis oppositis serratis, involucri cylindrico.*

1. T. LUCIDA (Cav. ic. 3. p. 33. t. 264), caule erecto parè ramoso, foliis lanceolatis argutè serratis basi aristato-serratis, invol. cylindraceo, ligulis 2, pappo 5-mero, paleis 3 obtusis, setis 2 scabris cæteris superantibus. (I) in Mexico (Cav.), ad montes circa urbem (Berl. pl. exs. n. 22, 709 et 2132), ad Tlapujahua (Keerl. in h. Mart.), ad Tamaulipas (Berl. n. 2132). Vild. sp. 3. p. 2126. Variat foliis inf. obtusis, super. acutis, angustioribus aut latioribus. Nunquam folia vidi connata ut vult Cavan. (v. v. c. et s. sp.)
2. T. FLORIDA (Sweet brit. fl. gard. ser. 2. t. 35), caule erecto ramoso, foliis lanceolatis argutè serratis basi aristato-serratis, ligulis sæpius 3, achænio paleis membranaceis 2-3 æqualibus aristisque nullis coronato. (I) in Mexico (Sw.). Similis T. lucidæ sed ligulæ majores et aristæ pappi nullæ. (v. s. ex hort. Barcl. in quo etiã Sweet descr.)

§. 2. Macrocephalæ, *foliis pinnatisectis, capitulis multi- (30 et ultra) floris, ligulis subrotundis amplis, invol. ovatis campanulatisve.*

3. T. ELLEPTICA (Smith in Rees cycl. v. 35. n. 7), erecta ramosa, foliis pinnatisectis, segmentis 3-4-jugis cum impari ellipticis subintegerrimis mucronatis. infer. minoribus, summis gradatim majoribus, corymbo subpaniculato, capitulis pedicellatis bracteolis subulatis 2-2 stipatis, invol. lævibus obovato-oblongis, ligulis amplis. — in Peruvia (Domb. l). Tagetes n. 27. Poepp. pl. exs. Peruv. Folia Rotam pinnatam subreferunt. Glandulæ foliorum rotundæ, involucri lineares. Spec. distinctissima. (v. s. ex coll. Domb. et Poepp.)
4. T. ZYPAQUIRENSIS (Humb. et Bonpl. pl. eq. 2. p. 18. t. 73), caule erecto paniculato, ramis oppositis subfastigiatis, foliis oppositis pinnatisectis, segmentis 4-6-jugis cum impari oppositis oblongis obtusis serratis subtus pubescentibus, rachi puberulâ hinc inde setigerâ, capitulis longè pedunculatis solitariis subcorymbosis, involucri oblongo-campanulato, ligulis oblongis 10-11 elongatis. (I) in Novâ-Granatâ propè urbem Zypaquira. H. B. et Kunth nov. gen. am. 4. p. 195. Ligulæ flavæ. Pappi squamæ circ. 5 lineari-subulatæ subinæquales.
5. T. PATULA (Linn. sp. 1249), caule erecto, ramis patulis, foliis pinnatisectis, segmentis lineari-lanceolatis serratis, serraturis præsertim superiorum aristatis, pedunculis 1-cephalis elongatis subcylindricis, involucri lævibus. (I) in Mexico et nunc in hortis totius fere orbis culta. Eandem aut valde affinem circa Quito legit. cl. Humb. et Bonpl. Sims bot. mag. t. 150. — Dill. elth. 273. t. 279. f. 361. Dod. pempt. 255. Hern. mex. p. 156. f. 1 et forte 2 et 3? Knorr del. 1. t. S. 8. Cor. aureo-fusæ sen fulvæ. Capitula duplò minora et longius pedunculata quàm T. erectæ, eodem modo variantia. (v. v. c.)
6. T. ERRECTA (Linn. sp. 1249), caule ramisque erectis, foliis pinnatisectis, segmentis lanceolatis serrulatis, serraturis præsertim superiorum aristatis, pedunculis 1-cephalis apice ventricoso-incrassatis; invol. subangulatis. (I) in Mexico sponte et nunc in hortis orbis fere totius culta. Lam. ill. t. 684. Cempoaxochilt Hern. mex. 154. T. major Gärtn. fr. 2. p. 434. t. 172. — Knorr del. 1. t. S. 6. Cor. unicolores citrino-flavæ. Capit. duplò majus quàm T. patulæ, brevius pedunculatum. Variat capitulis proliferis, corollis exter. tantum ligulato-subbilabiatis, aut omnibus ligulato-subbilabiatis, etc. Odor herbæ foetidus. (v. v. c. et s. sp. ex Berl. pl. exs. n. 737.)
7. T. CORYMBOSA (Sweet brit. fl. gard. ser. 1. t. 151), caule ramisque erectis sub-

cato-angulatis, foliis oppositis alternisque pinnatisectis, segmentis 6-8-jugis cum impari lineari-lanceolatis acuminatis serratis punctatis, fol. sup. serratis aristatis, rachi inter segmenta hinc inde setigerà, pedunculis 1-cephalis subnudis corymbosis, invol. oblongo-subturbato 5-angulato sub-50-floro, ligulis 20 amplis subrotundis. (1) in Mexico (Sweet), ad Tlapujahua (Keerl! in h. Mart.). Tag. tenuifolia H. B. et Kunth nov. gen. 4. p. 196 non Cav. Ligulæ supernè fusco-croceæ. Achænia glabra nigra. Pappi palææ subconcretæ obtusæ, sæpius libera longior acuminata. In hortis occurrit sæpè sub nom. falso T. Bonariensis Pers. (v. s. c.)

8. T. PEDUNCULARIS (Lag. et Rod. in annal. scienc. nat. 1802. v. 5. p. 74. Cav. cors. p. 201), caule erecto subdiffuso, ramis folisque inf. oppositis, sup. alternis, foliis pinnatisectis 5-6-jugis, segmentis lineari-oblongis grossè et acutè serratis, serraturis hinc inde setigeris, ramis longè denudatis 1-cephalis, invol. oblongo 5-dentato circ. 30-floro, ligulis 5 amplis subrotundis. (1) in Mexico circa urbem leger. cl. Alaman, Mairet et Berl. ! pl. exs. n. 1183. T. elongata Willd. sp. 3. p. 127? Ligulæ flavæ nec fusco maculatæ. Pappi palææ membranacæ obtusæ concretæ, 1 duplò longior libera lineari-subulata. (v. s. sp.)

9. T. PAUCILOBA, glabra erecta, caule basi suffruticoso, foliis oppositis 1-2-jugis, rachi lobisque linearibus acutis integris grossè glandulosis, capitulis nudis solitariis terminalibus pedicellatis, invol. oblongo 5-dentato, ligulis circ. 5 subrotundis. 3 inter Chilenses herb. Hænkeani servatur. Folia 1 $\frac{1}{2}$ -2 poll. longa, lobis 6-8 lin. longis, lineam latis, terminali pollicem longo. Invol. 7 lin. longum, 3 lin. latum. (v. s. in h. Hænke.)

10. T. TENUIFOLIA (Cav. ic. 2. p. 54. t. 169), erecta vix ramosa, foliis oppositis alternisque pinnatisectis, segmentis 6-8-jugis cum impari linearibus serratis, fol. summorum serraturis aristatis, pedunculis 1-cephalis subnudis paucis subcorymbosis, invol. obovato levi disco breviorè, ligulis amplis subrotundis. (1) in Peru, ubi vulgò *Chinchimali* dicitur. T. tenuifolia Pers. ench. n. 8 (excl. syn. Ort.). Corollæ flavæ concolores immaculatæ. Achænia glabra nigra. Pappus palæis 5, quarum 4 concretæ obtusæ et quinta sublibera longior acuminata. (v. v. c. in h. Madr. et Par.)

§. 3. Leptocephalæ, foliis pinnatisectis, rariùs bipinnatisectis, capitulis paucis- (5-20) floris, invol. cylindraceis, ligulis parvis aut nullis.

* Radiatæ.

11. T. GRAVIOLENS (L'Hér. ! herb. et mss.), caule erecto subtetragono oppositè ramoso, foliis oppositis pinnatisectis, segmentis 10-11-jugis cum impari elongato-lanceolatis serratis, super. decurrentibus subconfluentibus, infimis parvis in lobos aristatos incisus ad fol. conjunctionem confertis, paniculis laxis foliaceis, capitulis pedicellatis, invol. subtetragono 4-5-dentato, ligulis 4-5 patentibus invol. paulò longioribus. (1) ? in Peruvia olim legit Pavon. Herba 3-orgyalis graveolens. Folia ferè Ebuli, segmenta 2 poll. longa, 4 lin. lata. Pappi palææ inæquales, una subulata longissima. Species ex foliis distinctissima. Flores in herb. desiderantur, sed in mss. L'Hér. descr. (v. s. sine fl.)

12. T. GLANDULIFERA (Schränk pl. rar. h. Mon. 2. t. 54), caule erecto, ramis erectis brevibus, foliis alternis pinnatisectis, segmentis lineari-sublanceolatis utrinquè acuminatis serratis 6-8-jugis cum impari, super. subdecurrentibus, inf. cujusque folii parvis pinnatifidis lobis aristatis, capitulis corymboso-fasciculatis ad apicem ramorum plerisque subsessilibus circ. 7-floris, invol. cylindrico, ligulis 3 parvis. (1) in ruderatis et fossis ad San-Jago (Bert.!) et Valparaiso Chilensium (Gaud.!), Bonariâ (Dill.), Brasiliæ merid. Banda orient. (Bacle!) et prov. Rio-Grande (h. Mus. imp. Bras. n. 987). T. minuto flore Ventill. obs. 3. p. 64. Dill. elth. 2. p. 374. t. 280. f. 362. T. minuta Linn. sp. 1250. T. Bonariensis, Pers. 2. p. 459. n. 5. T. glandulosa Link enum. 2 p. 339. Colla mem. ac. Tur. 33. p. 165. app. 3. p. 53. T. montana hort.

Lugd. 1821. Ligulæ nunc albicantes ut vult Dill. sed pallidæ flavæ. Caulis nunc semipedalis et nomini Linnæano sat conveniens, nunc sæpius 1-2-pedalis. Odor gravis. Segmenta foliorum sesquipollicem ferè longa, 2 lin. lata. Glandulæ in foliis sparsæ grossæ. Ligulæ nunc integræ subrotundæ, nunc apice 2-3-dentatæ. Achænium subpubescens. Pappi palææ membranacæ breves obtusæ, una aristata longa corollâ tamen brevior. (v. s. sp. et v. c.)

13. *T. DAUCOIDES* (Schrad. ! ex sem. hort. Gœtt. 1833. p. 5. et in Linnæâ 1835: litt. p. 71), erecta alternè ramosa, foliis alternis pinnatisectis 10-11-jugis elongato-linearilanceolatis grossè et argutè inciso-serratis, lacinulis hinc indè serratis, segm. inf. parvis in dentes longè setigeros fissis, pedunculis axill. terminalibusque sæpius trifidis, pedicellis involuero oblongo 5-dentato brevioribus, flosculis circ. 15-18, ligulis 5 parvis subrotundis. ① patr. ign. Affinis *T. foetidissima* et glandulifera. Folia parcè glandulosa 3-4 poll. longa; segm. 15 lin. longa, 2-2 $\frac{1}{2}$ lin. lata, rarissimè pinnati lobata. Pappus ligul. palæis 2 elongatis et cæt. obtusis, disci palæâ 1-aristatâ, cæteris obtusis. (v. v. c.)
14. *T. FOETIDISSIMA* (hortul.), caule erecto paniculato-ramoso, foliis alternis pinnatisectis, segmentis oblongo-lanceolatis serratis 10-14-jugis, summis decurrenti-subconfluentibus, infer. parvis in lobos aristatos incisis, capitulis pedicellatis solitariis subcorymbosis circ. 12-floris, invol. ovato cylindraceo, ligulis 5 parvis oblongis. ① in Mexico (Alam.!), ad Guchilacæ Cordilleram (Berl. ! pl. exs. n. 1043). Affinis *T. glandulifera* et cum eâ sæpè sub nom. *T. glandulosa* confusa. Herba 1-3-pedalis foetidissima. Segm. fol. 6 lin. longa, 2 ferè lata. Pappi palææ 3 membranacæ breves, 2 aristatæ corollam ferè superantes. (v. s. sp. et v. c.)
15. *T. TERNIFLORA* (H. B. et Kunth nov. gen. am. 4. p. 196), caule erecto ramoso sulcato apice corymboso polycephalo, ramis patulis oppositis, foliis oppositis pinnatisectis, segmentis subsexjugis cum impari oppositis lanceolatis argutè serratis, capitulis subternis breviter pedicellatis, invol. clavato tubuloso 9-10-floro 5-dentato. ① prope Quito ad radices montis Pichinchæ. Pappi palææ 3 minimæ membranacæ, 2 linearilanceolatis. Ligulæ indescer.
16. *T. MULTIFLORA* (H. B. et Kunth nov. gen. am. 4. p. 197), caule erecto ramoso tereti apice corymboso polycephalo, foliis alternis pinnatisectis, sequentibus subquadrijugis subalternis lanceolatis serratis, capitulis fasciculatis longè pedunculatis, invol. tubuloso-subventricosio 14-15-floro, ligulis 4-5 subrotundis. ① prope Quito in radicibus montis Pichinchæ. Pappi duplicis palææ 5 exteriores lineares subæquales, 5 interiores minimæ obtusæ.
17. *T. MULTISETA*, caule erecto alternè ramoso, foliis alternis bipinnatisectis, lobis linearilanceolatis ferè omnibus in setam elongatam productis, pedunculis solitariis longis nudis 1-cephalis, invol. oblongo 5-dentato, flosculis 15-20, ligulis 5 parvis retusis obcordatisve. ① in Mexico (Alam. ! Hænke!), ad Cærnavacam (Berl. ! pl. exs. n. 969). An *T. subulata* Lallav. et Lex. nov. veg. mex. 1. p. 31 sed descr. non suff. Herba 6-12 poll. Folia hinc glandulas grossas gerentia, inf. setis destituta. Pedunc. 2-4 pollic. longi. Pappi palææ ferè omnes concretæ, 2 longiores subliberæ. (v. s.)
18. *T. GRACILIS*, erecta apice tantum parcè ramosa, foliis oppositis pinnatisectis, segmentis oblongo-linearibus grossè pauci serratis subalternis, inf. parvis in dentes setigeros incisis, pedunculis trifidis corymbosis, pedicellis involuero cylindrico lævi longioribus, flosculis 10-12, ligulis paucis vix invol. longioribus. ① in Peruvîâ? Radix tenera. Caulis pedalis teres. Folia distantia 1-1 $\frac{1}{2}$ poll. longa, segmentis glandulosis 4-5-serratis 4 lin. longis, lineam sesquilineamve latis. Pappi palææ 3 breves membranacæ, 2 aristatæ elongatæ. Affinis vid. *T. ternifloræ* et multifloræ. (v. s. comm. à Mus. reg. Par. ex coll. Donbeyi sine loci propr. design.)
19. *T. OLIGOCEPHALA*, caule erecto simplici aut infernè oppositè, supernè alternè, ramoso, foliis oppositis, summis alternis, pinnatisectis 5-7-jugis cum impari, sequentibus linearilanceolatis acuminatis serratis, infer. foliis et omnibus ferè ad apices caulis serrato-longè-aristatis, pedunculis paucis elongatis 1-cephalis,

- invol. oblongo circ. 20-floro, ligulis nullis aut fortè hinc inde minimis. (1) in Mexici Cordilleris Guachilapensibus legit cl. Berlandier (pl. exs. n. 1033 et 1197) Pappi paleæ 3 membranaceæ et 2 elongatæ aristatæ. Folia vix pollicem longa, segm. 4-6 lin. longis, 1-1 $\frac{1}{2}$ latis. (v. s.)
20. *T. FUSILLA* (H. B. et Kunth nov. gen. am. 4. p. 194), procumbens ramosa, foliis oppositis alternisve bipinnatisectis, lobis linearibus integerrimis, pedunculis solitariis 1-cephalis, invol. tubuloso-clavato, flosc. 12-13, ligulis 1-3 parvis. (1) prope Chillo Quitensium ubi dicitur *Anicillo*, herba enim anisum redolet. Ligulæ albidae. Pappi paleæ 3 obtusæ membranaceæ breves, 3 aristæformes alternantes. (v. s. sine flor.?)
21. *T. ANGUSTIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 194), caulibus diffusis subdichotomis, foliis alternis pinnatisectis, segmentis linearibus integerrimis, terminali subdentato, capitulis terminalibus solitariis longè pedunculatis circ. 15-floris, invol. clavato-tubuloso. (1) in frigidis montium Mexicanorum prope Toluca. Pappi paleæ 2 longiores lanceolatæ, 3^a truncata. Ligulæ indescrite, an nullæ?
22. *T. CARACASANA* (Humb. in Willd. sp. 3. p. 2128), caule erecto sulcato apice corymboso, foliis pinnatisectis, segmentis lanceolatis apice serratis, pedunculis elongatis erectis 1-cephalis alternis, invol. cylindræcis. (1) ad Caracas. Cæ. ign. †.
23. *T. DIANTHIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 197), caule erecto ramoso striato-sulcato apice corymboso polycephalo, foliis alternis impari-pinnatisectis, segmentis trijugis oppositis lanceolatis subintegerrimis, pedunculis 1-cephalis, invol. ventricosotubuloso 7-9-dentato, flosc. circ. 20, ligulis 7 lanceolato-oblongis. (1) in altis Peruviae inter flum. Catamayo et villam Lucarque. Ligulæ flavæ. Pappi squamæ 3 exter. subulatæ, 3 inter. alternantes minimæ.
24. *T. CORONOPIFOLIA* (Willd. enum. suppl. p. 60), caule ramisque erectis teretibus, foliis oppositis alternisve pinnatisectis, segmentis 6-8-jugis cum impari linearibus integerrimis rariùs dentatis, pedunculis solitariis 1-cephalis nudis, invol. oblongo-cylindræco, ligulis brevissimis subrotundis. (1) in Mexico. Jacq. eclog. 1. p. 118. t. 80. T. clandestina Lag. elench. h. Madr. p. 28. n. 362? Less. in Linnæa 1830. p. 160. Enalceida pilifera Cass. dict. 14. p. 443. Affinis *T. multifida*, sed capitula solitaria nec sensim corymbosa. Capit. 5-6 lin. longa, 1 $\frac{1}{2}$ lata. Pappi paleæ membranaceæ apice dentatæ concretæ sæpiùs inæquales, sed nullæ aristatæ. Achænia ad angulos puberula.
25. *T. FORNICULACRA* (Poepp. pl. exs. peruv. n. 26), erecta stricta parcè ramosa, foliis oppositis à basi pinnatisectis, segmentis 6-8-jugis cum impari linearibus integerrimis, pedunculis solitariis 1-cephalis nudis, invol. cylindrico, ligulis subsolitariis parvis. (1) in Peruvia legit cl. Poeppig. Herba ferè sesquipedalis glabra omninò media inter *T. micrantham* et *T. multifidam*. Diglossus variabilis Cass. dict. 13. p. 241 ad hanc (fortè cum aliâ specie mixtam?) pertinere videtur. (v. s.)
26. *T. MIGRANTHA* (Cav. ic. 4. p. 31. t. 352), erecta aut diffusa oppositè ramosa, foliis oppositis pinnatisectis, segmentis 1-2-jugis cum impari linearibus subulatis integerrimis, pedunculis solitariis 1-cephalis apice nudis, involucro cylindrico, ligulis 1-3 parvis obtusissimis. (1) in Mexico circa urbem (Berl. pl. exs. n. 472) et ad Queretaro (Née). Herba semipedalis tenuis glaberrima. Cor. flavæ parvæ. Pappus paleis 2-3 membranaceis truncatis inæqualibus et aristis 2-3 scabris rigidulis longis alternantibus constans. (v. s.)
- * * *Flosculosæ.*
27. *T. MULTIFIDA*, erecta ramosa, foliis alternis uni- aut bipinnatisectis, lobis linearibus acutis in setas non productis, capitulis corymbosis breviter pedicellatis, invol. oblongo subtrubinato lævi, flosculis 20-25 omnibus tubulosis. (1) in Mexico ad Lerma (Berl. pl. exs. n. 1145). Variat segmentis 4-6-jugis integerrimis dentatis aut pinnatilobatis. Pappi paleæ 3 aut 4 membranaceæ truncatæ dentatæ, 2 seu 1 liberæ lineares elongatæ non verè aristatæ. An *T. fil-*

folia Lag. elench. 25, sed ligulæ dicuntur minutæ lunulatæ dùm in meâ nullæ? (v. s.)

28. *T. FLOSCULOSA* (Spreng. syst. 3. p. 571), canle...., foliis pinnatifidis filiformibus elongatis integerrimis summis indivisis, pedunculis terminalibus solitariis, invol. turbinato, radio obsoleto. — in Monte-Videq. Cæt. ign. †.

† *Species non satis notæ.*

29. *T. LUNULATA* (Ort. dec. p. 44), canle erecto subangulato, ramis patulis, inf. oppositis, sup. subcorymbosis, foliis pinnatisectis, segmentis alternis oppositisque lineari-lanceolatis obtusè serratis, intimis minoribus eciliatis, pedunculis longissimis 1-cephalis, ligulis 5 obcordatis. (I) in Mexico. Ligulæ maculâ sanguineâ lunatâ maculatæ. Pappi paleæ a lineari-subulatæ, reliquæ brevès truncatæ. Numerus florum, eorum proportio, et forma invol. ignorantur.

30. *T. SUBVILLOSA* (Lag. elench. h. Madr. p. 29. n. 363), canle...., foliis pinnatisectis villosulis, segmentis lanceolatis serratis, capitulis laxè corymbosis. (I) in Mexico. Cæt. ign. †.

31. *T. VERTICILLATA* (Lag. et Rod. in annal. scienc. nat. 1802. v. 5. p. 73), canle erecto ramosissimo, foliis ramisque verticillatis ternis, capitulis ebrymosis, ligulis brevissimis. — ad Camanam.

Quid *T. ozolis* Spreng. à Steud. citata?

†† *Species exclusæ.*

T. FRUILLÆ Bert. = *Lasthenia Chilensis*.

T. PAPPOSA Vent. = *Dysodia chrysanthemifolia*.

T. PUMILA hortul. = *Dysodia chrysanthemifolia*.

T. PUNCTATA fl. mex. = *Adenophyllum coccineum*.

CCCLXXXIX. *THYMOPHYLLA* Lag. nov. gen. in elench. h. Madr. p. 25.

Capitulum multiflorum homogamum discoideum. Invol. campanulati apice dentati squaræ concretæ. Recept. nudum. Pappus paleis 5 truncatis brevibus 1-serialibus. — Suffrutex Mexicanus, ramis oppositis. Folia opposita conferta subsetacea parva (vix sesquilineam longa) tomentosa. Pedunculi terminales 1-cephali 1-2-phylli villosuli. Invol. extus pulverulento-tomentosum achæniis æquale. Cor. ex sicco atropurpureæ. — Genus mihi ignotum ad Tagetineas à cl. Cassini et Lessing relatum, sed tomentum fol. obstat et de glandulis silet cl. Lagasca qui solus plantam vidit.

1. *T. SETIFOLIA* (Lag. l. c. n. 325). — in Mexico.

Div. II. *POROPHYLLÆÆ*. — Tagetinearum et Pectidearum pars. Cass. opusc. 3. p. 55.

Capitulum semper homogamum discoideum. Invol. squaræ liberæ aut vix basi concretæ. Cor. disci inæqualiter fissæ subbilabiata. Styli rami in appendicem subulatam producti. Pappus setosus pluriserialis.

CCCXC. *POROPHYLLUM* Vaill. act. acad. Par. 1719. p. 407. Linn. hort. cliff. 494. Adams. fam. 2. p. 122. Cass. dict. 43. p. 56. — Kleinia Jacq. amer. 215. Willd. spec. 3. p. 1738. Less. syn. 196 non Juss. nec Haw. (1). — *Cacalia* sp. Linn.

Capitulum multiflorum homogamum. Recept. nudum punctatum. Invol. 1-seriale, squamis 5- (in spuris 9-12) oblongo-linearibus, ju-

(1) Ex juribus collectis prioritatis, facilitatis, etymologiæ, nomen *Vaillantii* hic

nioribus basi subconcretis. Cor. tubulosæ regulariter 5-dentatæ aut dente unico profundius diviso sublabiatæ. Styli rami exserti, basi intus series stigmaticas 2 gerentibus, apice undiquè (ferè ut in Vernoniaceis) hispidi acuti. Achænia elongata angulata sæpissimè scabra, apice plùs minùs in rostrum attenuata. Pappus pluriserialis pilosus scaber. — Herbæ aut suffrutices. Species genuinæ omnes glabræ Americanæ. Folia alterna aut rarius opposita, sæpiùs glauca, ovata aut oblonga, integra aut dentata, sæpiùs glandulis pellucidis amplis oblongis notata. Pedicelli elongati 1-cephali. Invol. sæpè cæruleo-purpureove colorata et glaucescentia. Cor. sordidè purpureæ. — Affinitas subdubia; an Senecioneis, an Vernoniaceis propinqua?

Sect. 1. EUPOROPHYLLUM.

Invol. squamæ 5. — Folia sæpè poris glandulosis notata. — Species Americanæ.

§. 1. *Capitulis cylindraceis elongatis, invol. squamis mucrone calloso apiculatis, cor. regulariter 5-dentatis, achæniis evidentius rostrato.*

1. *P. MACROCEPHALUM*, herbaceum erectum, foliis longè petiolatis ellipticis obtusis crenatis ad sinus pellucido-glandulosis, limbis petiolo brevioribus, pedunculis inflato-turbinatis, invol. squamis mucrone calloso apiculatis. (1) in Mexico circa Villalpando legit cl. Méndez. *Tsinoma* Hern. mex. 434. f. 3. *Kleinia glandulosa* flor. mex. icon. ined. nimis fortè *P. elliptico* affine, differt petiolis longioribus, foliis parvis 6 lin. longis, 4 lin. latis, pedunculis crassioribus, invol. pollicem longis distinctis glandulosis. Vulgò in Mexico dicitur *Tzinoma* ex Hern., *Popoloquolite* ex Moç. (v. s.)
2. *P. ELLIPTICUM* (Cass. dict. 43. p. 46), herbaceum erectum, foliis petiolatis ellipticis oblongisve obtusis mucronatisve crenatis ad sinus aut in disco pellucido-glandulosis, invol. squamis mucrone calloso acuminatis. (1) in Peruvia (Cav.), ad Turbaco Novo-Granatensium (H. et B.), in Brasiliâ (Cham.!), ad Bahiam (Salzm.), in Santo-Domingo (Menz. in h. Lindl.), Martinicâ (Sw.), in aridis et argillosis. *Cacalia Porophyllum* Linn. sp. 1169. (excl. syn.) Cav.! ic. 3. p. 11. t. 222. *Kleinia Porophyllum* Willd. sp. 3. p. 1738. — Pluk. alm. a. 161. f. 1. mal. Variat achæniis in diversis aut iisdem individuis hirtellis, puberis aut glabris! (v. v. et s.)
- f. intermedium*, foliis ellipticis oblongisve crenatis opacè glandulosis aut subglandulosis. (1) in Brasiliâ (Gaudich.), ad Bahiam (Blanch.), in Serrâ dos Orgaos prov. Rio-de-Janeiro (Vauth. ! pl. exs. n. 305). Valdè accedit ad *P. ruderale*, sed huc potiùs referendum videtur. Variat ut var. *a* achæniis hirtis et lævibus. An species propria? (v. s.)
3. *P. RUDERALE* (Cass. dict. 43. p. 56), herbaceum erectum, foliis petiolatis elliptico- aut lanceolato-oblongis acuminatis subintegerrimis impunctatis. (1) in ruderatis et cultis insularum Caribæarum, Jamaicæ (Sw.), Sancti-Domingi (Jacq.), Martinicæ (Jacq.), Guadalupæ (Perr.), Porto-Ricci (Wydl.!), Cubæ (La Sagr.!) *Cacalia ruderalis* Sw. ! prod. 110. *Kleinia ruderalis* Jacq. am. 215. t. 127. (v. s.)
4. *P. VIRIDIFLORUM*, fruticosum erectum, foliis breviter petiolatis obovatis subundulato-repandis aut summis integerrimis, ad margines glandulas pellucidas paucas gerentibus. 3 locis temperatis Mexici juxta Walladolid (H. B. K.) et Cuernavaca (Berl. ! pl. exs. n. 956). *Kleinia viridiflora* H. B. et Kunth nov.

admissi, nam Kleinie nomen generi in quo *Cacalia* *Kleinia* Linn. militabat, evidenter præcæntibus Haworthio et ipso Linnæo servandum erat.

KLEINIA Juss. = *Jaumea*.

gen. am. 4. p. 157. Frutex orgyalis affinis P. elliptico. Cor. virescentes. Invol. squamæ obtusiusculæ glandulis biseriatis notatæ. (v. s.)

5. P. NUMMULARIUM, herbaceum erectum, ramis foliisque oppositis, foliis petiolatis elliptico-orbiculatis obtusissimis subcrenatis ad sinus opacè glandulosi, invol. squamis mucrone calloso apiculatis dorso glandulosi. — in Mexici umbrosis ad Tehuantepec et loc. vicin. Herba fetida angusto florens (Alam.). Rami primâ fronte videntur folia pinnata! Invol. 4-5 lin. longa. Limbus folii 8 lin. longus, 6 lin. latus. Petioli 4-lin. (v. s. comm. à cl. Alaman.)

β. foliis obovatis. — (v. s. in h. Hænke.)

6. P. JORULLENSE (Cass. dict. 43. p. 57), fruticosum ramosissimum, foliis longè petiolatis oblongis obtusis subrepandis eglandulosi, invol. squamis lanceolato-linearibus acutis. ♀ in Mexici montis ignivomi Jorullo declivitate orientali. Kleinia Jorullensis H. B. et Kunth nov. gen. am. 4. p. 156. t. 356.

7. P. LANCEOLATUM, glabrum strictum, caule imâ basi suffruticuloso, foliis alternis lanceolatis basi attenuatis subacutis subingerrimis eglandulosi, ramis erectis apice nudis 1-cephalis, achænio scabro. ♀ in Brasiliæ prov. Rio-Grande. Affine P. prenanthoidi, sed folia alterna, pedunculi strictè erecti, achænia scabra. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 936. miss.)

8. P. PRENANTHOIDES, glabrum erectum, caule virgato basi suffruticuloso, foliis oblongo-lanceolatis utriusquè attenuatis subintegerrimis impunctatis, infer. oppositis, super. alternis, paniculâ laxâ, capitulis nutantibus oblongis. ♀ in prov. Brasiliæ Sancti-Pauli. Habitus ferè Prenanthis. Achænia scabriuscula. An capitula exsiccatione aut naturâ propriâ nutant? (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 402. miss.)

9. P. HÆNKI, fruticosum erectum, ramis foliisque oppositis, foliis lineari-lanceolatis utriusquè attenuatis eglandulosi, capitulis breviter pedicellatis, achæniis scabris. ♀ in Cordilleris Chilensibus ex herb. Hænk. Affine P. lineari, sed folia latiora omnia opposita, pedunculi breviores, capitula paulò minora. An P. oppositifolium Spreng., sed patria diversa et folia non brevia dicenda cum 15 lin. longa, 2-3-lata? (v. s. in Hænk. à cl. Sternb. miss.)

10. P. LINEARE, fruticosum erectum glabrum, foliis alternis oppositisque linearibus integerrimis acutis eglandulosi, invol. oblongi squamis apice breviter mucronatis, achæniis scabris. ♀ in Brasiliæ prov. Rio-Grande. Folia sesquipollic. longa, lineam lata. Pedunculi post anthesin incrassati striati subtrabinati. Capit. 6 lin. longum. (v. s. in h. Mus. Par. à Mus. imp. Bras. sub n. 933 et fortè 934.)

11. P. LINIFOLIUM, suffruticosum glabrum multicaule subcæspitosum, foliis alternis linearibus integerrimis subacutis eglandulosi, invol. ovali. ♀ in Brasiliâ (Ard.), ad Monte-Video (Comm.), in prov. Rio-Grande (h. Mus. Bras.), fortè in Peruvia et Mexico (h. Hænk.). Cacalia linifolia Linn. mant. 109. Kleinia suffruticosa Willd. sp. 3. p. 1739 non Lodd. Cacalia Linaria var. β pumila Poir. suppl. 2. p. 6. — Lam. ill. t. 673. f. 4. Valdè accedit ad P. Linariam, sed multò minus et magis cæspitosum. Folia 4-6 lin. longa. Achænia glabriuscula. (v. s. in h. Mus. reg. Par. ex Mus. imp. Bras. sub n. 924, 935 et fortè 940 miss.)

§. 2. Capitulis ovali-oblongis brevioribus, invol. squamis obtusis margine scariosis, corollis lobo uno profundiore sublabiatis (an semper?), achænio apice non sensim attenuato. — Hunteria fl. mex. icon. non Roxb.

12. P. LINARIA, suffruticosum glabrum erectum, ramis virgatis, foliis angustè linearibus acutis integerrimis glandularum pellucidarum serie duplici notatis, invol. squamis obtusis coloratis. ♀ in Mexico (Cav.) Cacalia Linaria Cav. ic. 3. p. 29. t. 257 non Poir.? Hunteria columbina icon fl. mex. Invol. ovali-

oblongum 5 lin. longum. 4 lin. latum, colore fusco cærulescente substatibili decorum. Achænium suberostre. (v. s. ex hort. Mex. contr. à cl. Berl. pl. exs. n. 881.)

13. *P. TAGETOIDES*, herbaceum? erectum, foliis breviter petiolatis lanceolato-linearibus, acutis subintegerrimis margine pellucido-punctatis subtus glaucescentibus, invol. squamis ovato-oblongis obtusis externè pruinoso-glaucescens. (1) in hortis Mexicanis leger. cl. Humb. et Bonpl. *Kleinia tagetoides* H. B. et Kunth. nov. gen. 4. p. 158. Valdè affine dicitur *P. linifolio*.
14. *P. DECUMBENS*, suffruticosum multicaule decumbens subcæspitosum, foliis linearibus-oblongis basi attenuatis obtusiusculis integerrimis glaucissimis apice glanduloso-callosis cæterum eglandulosis, invol. squamis vix acutis. 2/ 5 in Mexico (Lodd.). *Kleinia suffruticosa* Lodd. bot. cab. t. 1561 non Willd. *Kleinia linearis* hort. Par. 1817! Capitula ferè *P. Linariæ*, folia *P. colorati* sed obtusiora, caules *P. linifolii*. (v. v. c. et s.)
15. *P. COLORATUM*, herbaceum erectum, ramis virgatis, foliis linearibus-lanceolatis subintegerrimis margine hinc indè pellucido-punctatis, invol. squamis obtusiusculis coloratis. (1) in Mexico circa Guanajuato et Temascalco (H. et B.), ad Tlapujahu (Keerl). *Kleinia colorata* H. B. et Kunth nov. gen. am. 4. p. 158. Mart. amœn. monac. p. 23. t. 15. *Hunteria fuscata* icon. flor. mex. Invol. fusco-sen purpureo-cærulescens. Folia pollicem longa, sequilineam lat.
16. *P. OBTUSIFOLIUM*, suffruticosum multicaule, caulibus erectis striatis, foliis oblongis obtusis basi attenuatis subintegerrimis glandulas pellucidas paucas ad margines gerentibus, nervo medio albo prominulo, invol. squamis obtusis coloratis. 5 in Mexico propè Villalpando legit cl. Mendez. Stirps circ. pedalis. Folia sup. ferè linearia subacuta. Invol. ovalia pulchrè cærneo-purpurascens. (v. s.)

Sect. II. CUSIMBUA.

Invol. squamæ 9-12. — Folia poris glandulosis destituta (saltèm in descr. omissa). — Species Indicæ. — An genus proprium?

17. *P. CUSIMBUA*, basi fruticosum erectum, caulibus sulcatis apice paniculatis. foliis semiamplexicaulibus oblongo-lanceolatis longè acuminatis argutè serratis basi integerrimis, pedicellis squamulosis capituli longitudine, invol. cylindrici squamis plurimis (ultra 9) linearibus. 5 in Nepaliâ ubi Nawaria dicitur *Cusimba-Swa*. *Cacalia Cusimbua* Don prod. fl. nep. 179. *Kleinia Cusimbua* Less. in Linnæâ 1831. p. 133. Folia ex Don subtus pubescentia, ex Less. pallidiora. Ovaria glaberrima.
18. *P. CACALIOIDES*, herbaceum erectum, foliis petiolulatis carnosulis oblongo-ovatis acuminatis retrorsum mucronulato-denticulatis, capitulis longè pedunculatis racemoso-subcorymbosis, invol. 9-phylo cylindraceo, squamis linearibus. — in insulâ Luzoniâ. *Kleinia cacalioides* Less. in Linnæâ 1831. p. 132. Cor. apice purpureæ? limbo à tubo non distincto. Stylus basi bulbosus. Affine dicitur *P. hieracioidi*.
19. *P. HIERACIOIDES*, herbaceum erectum simplex, foliis petiolatis obovatis dentatis, capitulis subcorymbosis breviter pedunculatis, invol. squamis 12 lanceolatis. (1) in Indiâ orientali. *Cacalia hieracioides* Willd. sp. 3. p. 1731? *Kleinia hieracioides* Less. in Linn. 1831. p. 133. Stirps Lessingiana videtur diversa à Willdenowianâ. Folia ex Willd. obtusa, ex Less. longè acuminata. †.

Sect. III. KUGAIA.

Invol. basi calyculatum. — Folia pinnatifida. — Species Japonica. — An genus proprium?

20. *P. JAPONICUM*, herbaceum, caule debili carnosio villosio, foliis petiolatis incisopinnatifidis glabris, lobis oblongis acutis dentatis, capitulis paniculatis, pedicellis subsquamosis, involuero calyculato. 2/ in Japoniâ insulis circa Nagasaki ubi

vulgò *Kwugai Sai* dicitur. Senecio Japonicus Thunb. fl. jap 315. Willd. sp. 3. p. 1978. *Kleinia Japonica* Less. in linnæa 1831. p. 134. syn. 133.

† *Species vix notæ et verisimiliter à genere excludendæ.*

21. P. ? *OPPOSITIFOLIUM* (Spreng. syst. 3. p. 438, sub *Kleinia*), suffruticosum, caule ramisque filiformibus, foliis oppositis sessilibus linearibus brevibus involucrique pubescenti-pulverulentis, capitulis solitariis pedicellatis. 3 ad Buenos-Ayres. *Cacalia oppositifolia* Poir. l suppl. 2. p. 7. (v. s. sine fl.)

22. P. ? *SERLOI* (Spreng. l. c. sub *Kleinia*), fruticosum simplicissimum, foliis decussato-oppositis sessilibus lanceolatis obtusiusculis crenato-serratis eglandulosis hirsutis, pedunculis racemosis terminalibus. 5 ad Rio-Grande Brasiliæ. Cæst. ign. †.

23. P. ? *FILIFOLIUM*, (Spreng. l. c. sub *Kleinia*), herbaceum ramosum, foliis fasciculatis lineari-filiformibus glabris, pedunculis 1-cephalis terminalibus. — in Monte Video. Cæst. ign. An eadem ac *P. linifolium* ? †.

24. P. *OBSCURUM* (Spreng. l. c. sub *Kleinia*), herbaceum elatum supernè divinum glaucum, foliis alternis remotis linearibus glabris, pedunculis axillaribus terminalibusque 1-cephalis. — ad Rio-Grande Brasiliæ. Cæst. ign. †.

Subtrib. V. *HELENIEÆ* Cass. dict. sc. nat. 1829. p. 13. opusc. 3. p. 36. Less. syn. 237

Capitula sæpius heterogama, fl. radii ligulatis (rarò tubulosis ampliatis) femineis neutrisve 1-serialibus, reliquis hermaphr., rariùs homogama, nunquam dioica. Antheræ sæpè nigricantes, basi subproductæ nec caudatæ, articulo antherifero tereti. Styli rami nunc apice truncati, nunc cono imò appendice supra pilos collectores primarios gerente aucti. Achænium erostre. Pappus multipaleaceus, squamellis planis distinctis 1-serialibus scariosis obtusis seu acuminatis, rarissimè aut abortu nullis, aut in vaginulam concretis. Invol. squamæ plurimæ liberæ. — Folia sæpius alterna interdum opposita. Species pleræque Americanæ.

Div. I. *GAILLARDIÆ*. — *Galardiæ* Nutt. gen. am. 2. p. 177. — *Galardiæ* et *Rosilleæ* Less. syn. 237 et 245.

Pappi paleæ membranacæ sæpiissimè integræ. Recept. epaleaceum nudum aut alveolatum fimbriliferumve. — Species notæ omnes Americanæ.

Subdiv. I. *EUGAILLARDIÆ*.

Ligulæ neutre.

CCCXCI. *GAILLARDIA* Foug. mem. acad. sc. Par. 1786. p. 1 et 6 (sub *Gail-lardâ*). Cass. dict. 18. p. 17. 55. p. 264. — *Galardia* Lam. dict. 2. p. 285. illustr. t. 708. Less. syn. 237. — *Calonnea Buchoz* icon. 1786 t. 126. — *Virgilia L'Hér. diss. cum ic. non Lam.* — *Galordia Rœusch. nom.* 251.

Capitulum multiflorum radiatum, ligulis neutris multinerviis glandulosus apice palmato-trifidis, fl. disci hermaphroditis, tubo brevi, limbo pilis articulatis hispido 5-fido. Recept. fimbriliferum convexum. Invol. squamæ 2-3-seriales, appendice foliaceâ acuminatâ disco longiore. Styli rami appendice longâ subulatâ hispidâ terminati. Achænia oblonga villosa. Pappi paleæ uninerviæ in aristam longè acuminatæ. — Herbhæ Boreali-Americanæ erectæ subpilosæ. Folia alterna, super. integra, sessilia aut semiamplexicaulia. Rami longè denudati 1-cephali. Capituli discus fuscus seu flavus, ligulæ flavæ seu basi crocæ.

§. 1. Annuæ, invol. squamis interioribus abruptè in appendicem lineari-subulatam angustatis.

1. *G. DRUMMONDII*, annua subpubera, foliis angustè lanceolatis indivisis aut incisis, invol. squamis basi brevissimâ glabrâ appendice subulatâ scabro-ciliatis, corollis hirsutis pappo longioribus, pappi paleis basi latè dilatatis, ligulis basi cuneatis involucri paulò longioribus. (1) in campis Mexicanæ prov. Texas ad San-Fernando de Bexar (Berl. pl. exs. n. 2027 et 2067), et ad Rio-Brazos (Drumm.). *Gaillardia bicolor* var. β Drummondii Hook. bot. mag. t. 2368. Ligulæ 5 lin. longæ, purpureæ, vix apice flavæ. Recept. fimbriellæ distantes rigidæ. Folia alia indivisa, alia pinnatilobata, parcè pubera. (v. s. sp.).
2. *G. PULCHRELLA* (Fong. l. c. Cass. dict. 18. p. 17), annua subhirsuta, invol. squamis basi glabriusculis appendice abruptè angustatâ rigidè ciliatâ, corollis disci parcè villosis pappi longitudine, pappi paleis basi angustis rigidulis. ligulis basi valdè angustatis maculatis. (1) in Louisiana olim legit com. d'Es-sales et in hortos Gallicos introduxit anno 1786 sed anno (ni fallor) 1791 semina omnia ad maturitatem non pervenerunt et species in hortis amissa fuit. *Gaillardia bicolor* Lam. dict. 2. p. 585. ill. t. 708. *Calonnea pulcherrima* Bouchoz icon. t. 126. *Virgilia helioides* L'Hér. diss. cum. icon. 2. Garten mag. 1810 p. 42. t. 6. *G. amara* Raf. fl. lud. p. 69? *G. alternifolia* Ræusch. l. c. Ligulæ 8-9 lin. longæ, circ. 3 lin. latæ. (v. s. ex h. L'Hér.)

§. 2. Perennes, invol. squamis lineari-lanceolatis appendice vix à basi distinctâ.

3. *G. LANCEOLATA* (Michx. fl. bor. am. 2. p. 142), perennis, invol. squamis extus villosis-subhirsutis lineari-lanceolatis, corollis disci hirsutis pappi circ. longitudine, pappi paleis basi subdilatatis, ligulis obovato - cuneatis basi maculatis. \mathcal{L} à Carolinâ ad Floridam in aridis (Michx.), in pinetis siccis Georgiæ mediæ (Ell.), ad fl. Saskatchewan et in montibus scopulosis (Hook. fl. bor. am. t. p. 315). *Gaillardia rustica* Cass. dict. 18. p. 20. *Gaillardia bicolor* Ell. sketch 2. p. 449. Sims bot. mag. t. 1602. (excl. syn. ferè omn.) Ligulæ 7 lin. longæ, 5 lin. latæ, basi crocæ, cæterum flavæ. Cor. disci crocæ. (v. v. c.)
4. *G. ARISTATA* (Pursh fl. bor. am. 2. p. 573), perennis, invol. basi hirsutissimis squamis lineari-lanceolatis, cor. disci hirsutissimis pappo longioribus, pappi paleis basi latè dilatato-membranaceis, ligulis basi longè cuneatis unicoloribus. \mathcal{L} in collibus siccis ad Rocky-Mountains (Pursh), in collibus herbosis ad fl. Missouri (Nutt.). Lindl. bot. reg. t. 1186. Hook. bot. mag. t. 2940. fl. bor. t. p. 315. *G. bicolor* var. β . Nutt. gen. am. p. 175. Magis quàm priores hirsuta. Ligulæ 12-14 lin. longæ, 5 lin. latæ, flavo-subaurantiæ non maculatæ. Cor. disci fusco-purpureæ, lobis elongatis. Achenium rufo-villosissimum. Pappus albus, hanc et 2 antecedentes sub nom. *G. bicoloris* confudit Spreng. syst. 3. p. 618. excl. syn. Walt. (v. v. c.)

+ *Species exclusæ.*

G. ACAULIS Pursh = *Actinella acaulis*.

G. FIMBRIATA Michx. = *Leptopoda puberula*.

CCCXCII. BALDUINA Nutt. gen. am. 2. p. 175. Ell. sketch 2. p. 447. Less. syn. 238.

Capitulum multiflorum radiatum, ligulis neutris apice tridentatis aut trifidis, fl. disci hermaphrod., cor. tubulosis 4-5-dentatis extus glanduliferis basi corneis. Invol. imbricatum campanulatum. Recept. convexum alveolato-fimbriiferum. Anthæræ basi breviter bisetosæ! Styli rami inclusi vix appendiculati. Achænia in foveolis immersa se-

ricea obconica. Pappi paleæ membranaceæ breves. Herbæ Boreali-Americanae perennes glabriusculæ. Caules teretes simplices aut ramosi, ramis apice nudis 1-cephalis. Folia alterna sessilia integra. Cor. flavæ. — Affinitas ex antheris bisetosis dubia!

Sect. I. BALDUINA *Ell. in obs. 2. p. 447.*

Pappus achænium æquans biserialis paleaceus, paleis inter. acutis, exter. sub-fimbriatis. Invol. plariserialæ.

1. B. UNIFLORA (Nutt. et Ell. ! ll. cc.), caule simplici 1-cephalo pubescente, foliis angustè obovatis. \mathcal{U} in humidis et paludum marginibus Carolinæ et Georgiæ. (v. s. comm. à cl. Elliott.)

Sect. II. ACTINOSPERMUM *Ell. sketch p. 448. in observ.*

Pappus brevis 12-14-paleaceus, paleis in cupulam patentem dispositis obtusis. Invol. biserialæ.

2. B. MULTIFLORA (Nutt. et Ell. ll. cc.), caule ramoso glaberrimo polycephalo, foliis linearibus ferè setaceis. \mathcal{U} in collibus arenosis secus Altamaha ad acidulam Barrington.

CCCXIII. LEPTOPODA *Nutt. gen. am. 2. p. 174. Ell. sketch 2. p. 445. Cass. dict. 26. p. 79. 55. p. 264 et 272. Less. syn. 238.*

Capitulum multiflorum; fl. radii circ. 20 ligulatis trifidis neutris, fl. disci hermaphr. tubulosi 3-5-dentatis. Invol. biserialæ, squamis exter. lineari-subulatis, int. sublanceolatis. Recept. convexum nudum. Styli rami breves exappendiculati. Achænia subcylindracea. Pappus paleis membranaceis circ. 8 brevibus fimbriatis in aristam non productis. — Herbæ Boreali-Americanae erectæ simplices 1-cephalæ. Folia alterna lineari-lanceolata subdentata. Capitula terminalia solitaria. Flores flavi. — Habitus Helenii sed ligulæ neutræ.

1. L. PUBERULA (Macbr. in Ell. sketch 2. p. 445), caule viscido-pubescente fistuloso stricto, foliis semiamplexicaulibus glabris punctatis, caulinis incisedentatis, achæniis villosis subturbinatis. \mathcal{U} in paludosis apricis Carolinæ et Floridæ. *Galaridia fimbriata* Michx. fl. bor. am. 2. p. 142. *Helonium vernale* Walt. fl. car. p. 210.
2. L. HELENIUM (Nutt. gen. am. 2. p. 174), caule glaberrimo solido, foliis decurrentibus glabris denticulatis, achæniis glabris cylindricis. \mathcal{U} in humidis pinetis Carolinæ et Georgiæ. *Lept. helenioides* Cass. dict. 26. p. 80. *Lept. decurrens* Macbr. in Ell. sketch 2. p. 446.

Subdiv. 2. EUHELENIÆ.

Ligulæ fœminæ seu nullæ.

CCCXIV. GUTIERREZIA *Lag. elench. h. madr. p. 30.*

Capitulum pluri- (8) florum radiatum, ligulis 3 fœmineis. Invol. imbricatum, squamis apice reflexis. Recept. favosum dentato-paleaceum. Pappus polyphyllus paleaceus. — Planta Mexicana suffruticosa glabra resinosa. Folia sparsa sessilia linearia acuta integerrima. Corymbus terminalis oligocephalus. Cæt. ign. — Genus mihi ignot. et dubium. A Galinsogâ, quâcum junxit Sprengel, differt recept. favoso nec ut videtur ex descr. Lag. verè paleaceo, invol. imbricato nec 1-seriali, etc. †.

1. *G. LINEARIFOLIA* (Lag. l. c.). ♂ ♀ in Mexico. *Galinsoga linearifolia* Spreng. syst. 3. p. 579. Folia pollicem et amplius longa, vix lineam lata, sup. breviora.

CCCXCV. *ACHYROPAPPUS* H. B. et Kunth nov. gen. am. 4. p. 257. t. 390 nec Bieb. — *Achyropappi* sect. Less. syn. 239. — *Chamæstephanum* Willd. in Berl. mag. 1807. p. 140. ex Kunth.

Capitulum multiflorum radiatum, fl. radii 3-5 fœmineis. Invol. squamæ quinque 1-seriales ovales obtusissimæ. Recept. parvum nudum alveolatum. Cor. disci 5-fidæ vix pappo longiores. Styli rami breviter appendiculati. Achænia lineari-cuneata 3-4-gona. Pappus paleis 6 membranaceis obovato-muticis. — Herbæ Mexicanæ annuæ vix subpuberulæ erectæ. Folia opposita et super. alterna, multipartita, lobipinnatipartitis oblongis seu linearibus obtusis. Capitula longè pedicellata laxissimè corymbosa. Cor. flavæ (nec radio albo ut sphalmate legitur in opere Kunth). Genus vix à *Schkuhria*, rectè monente cl. Nees del. sem. Bonn. 1831, satis diversum.

1. *A. ANTHEMOIDES* (H. B. et Kunth 4. p. 259. t. 390), caule foliisque pubescentibus, fol. laciniis lineari-sublancoatis, ligulis 5. ① in Mexici agris frigidis circa Toluccam ad pagum Islahuaca. (v. s. in h. acad. reg. Monac.)
2. *A. SCHKUHRIOIDES* (Link et Otto abb. neu. gew. gart. Berl. n. et t. 30), caule foliisque subscabris, fol. laciniis linearibus, ligulis 1-3. ① in Mexico (Link). ad occid. urbis Guanaxuato (Mendez!). *Schkuhria senecioides* Nees del. ex. h. Bonn. 1831. An à priore satis differt? (v. v. c. et s. sp.)

† Species exclusa.

A. PEDATUS Less. = *Florestina pedata*.

CCCXCVI. *SCHKUHRIA* Roth cat. bot. 1. (1797), p. 116. Cass. dict. 48. p. 87. 55. p. 263. Less. syn. 239. non Mœnch. — *Tetracarpum* Mœnch suppl. (1802), p. 241. — *Pectidis* sp. Lam. Ort. — *Mieria* Lallav. et Lex. nov. reg. mex. 2. p. 9. nec Link.

Capitulum pauciflorum, ligulis 1-2 fœmineis, fl. disci 5-8 tubulosi 4-5-dentatis brevibus. Invol. squamæ circ. 5 uniseriales obovatæ apice submembranaceo coloratæ. Recept. parvum nudum. Styli rami breviter appendiculati. Achænia tetragona basi attenuata. Pappi paleæ 8, 4 ex angulis achæniæ ortis acuminato-aristatis, 4 alternis nunc priores referentibus nunc iis brevioribus et obtusioribus margine subfimbriatis. — Herbæ Americanæ annuæ ramosissimæ glabræ. Folia alterna tri- seu pinnatisecta, segmentis linearibus tenuissimis integerrimis. Capitula ovata solitaria longè pedicellata. Cor. lutæ.

1. *S. OCTOARISTATA*, pappi paleis 8 æqualibus acuminato-aristatis corollas disci superantibus, ligulâ involucri longiore. ① in Peruvîâ legit cl. Dombey. Sequenti simillima et cum eâ in herb. interdum confusa, sed pappo (an satis?) diversa. (v. s.)
2. *S. ABROTANOIDES* (Roth cat. 1. p. 116), pappi paleis 4 acuminato-aristatis, 4 alternis brevioribus obtusatis omnibus corollis disci brevioribus, ligulâ involucri longiore. ① In Peruvîâ (Dombey!), ad Huanacum urbem (Ort.) nec in Mexico. — *Schkuhr* handb. 3. t. 250. b. *Pectis* pinnata Lam. Journ. h. nat. 2. p. 150. t. 31. *Pectis* multifida Ort. dec. p. 45. *Tetracarpum pinnatum* Mœnch l. c. In Peruvîâ dicitur *Asha-Pichana* id est scopus amaræ (Ort.). Caulis in specim. Dombeyano basi induratus suffrutescens. (v. v. c. et s. sp.)
3. *S. VIRGATA*, pappi paleis 4 acuminato-aristatis, 4 alternis brevissimis obtus-

satis, ligulâ involuero brevior, interdum nullâ. ① in Mexico circa urbem (Berl. pl. exs. n. 708), ad colles Taenbaya et aliis locis Mexico vicinis (Lallav.), ad occid. urbis Guanaxuato (Mend.). *Mieria virgata* Lallav. et Lex. nov. veg. mex. 2. p. 9. Invol. basi squamulis accessoriis duabus instructum, nec ut in aliis nudum. (v. s. sp.)

† *Species exclusa.*

S. *SENECIONOIDES* Nees = *Achyropappus* Schkumrioides.

CCCXCVII. FLORESTINA Cass. bull. philom. 1815. p. 175. *journal. phys.* 1816. p. 144. 1817. p. 11. *dict.* 17. p. 155 et 55. p. 266. *atl.* 1816. f. 3. t. 8. *opusc. phyt.* 1. p. 197. t. 8. — *Lepidopappus fl. mex. ic. ined.* — *Stevia* sp. Cav. — *Agerati* sp. Ort. — *Hymenopappi* sp. Lag. — *Achyropappi* sect. Less.

Capitulum homogamum discoideum. Invol. floribus brevius, squamis 8 uniserialibus oblongis obtusis. Recept. parvum planum subalveolatum. Cor. tubo brevi, lobis 5 inæqualibus papilloso-hirtellis. Styli rami in appendicem subulatam hirtam producti. Achænia subtetragona oblonga hispidula. Pappus paleis 8-12, uniseriatis scariosis suborbiculatis nervo saltē basi opaco. — Herbæ Mexicanæ, *Stevia* facie, pube adpressâ brevissimâ subcanescentes, affines *Hymenopappo*. Folia alterna pedatipartita seu indivisa. Capitula laxè corymbosa seu subpaniculata. Cor. albidæ aut purpurascens. — Si ut sectio habenda, potius meo sensu *Hymenopappo* quàm *Achyropappo* adjungenda.

1. F. PEDATA (Cass. ll. cc.), foliis petiolatis pedatipartitis, lobis 5 oblongis integris, summis indivisis. ① in Mexico (Cav.), circa urbem in montanis (Berl. pl. exs. n. 707 et 743). *Stevia pedata* Cav. ic. 4. p. 33. *Ageratum pedatum* Ort. dec. p. 38. *Hymenopappus pedatus* Lag. nov. gen. p. 28. n. 356. *Achyropappus pedatus* Less. syn. 239. Invol. sæpè purpurascens. (v. s. sp. et v. c.)
2. F. TRIPTERIS, foliis petiolatis trisectis, segmentis ovatis aut ovali-oblongis aut in apice caulis linearibus, omnibus integerrimis. ① in Mexico ad Laredo (Berl. pl. exs. n. 2077). *Lepidopappus tripteris* fl. mex. ic. ined. Pappi paleæ acutæ, margine hyalinæ, nervo medio crassiusculo. (v. s. sp.)
3. F. CALLOSA, foliis linearibus integerrimis, ramis subdichotomis glandulosomuricatis. ① ? in territorio Arkansano Amer. bor. *Stevia callosa* Nutt. ! *journal. ac. Philad.* 1821. p. 121. et in linnæa litt. 1829. p. 40. Annua à Nuttallio dicitur, sed in specim. ab eo misso caulis suffruticulosus videtur. Pappi paleæ membranaceæ obtusæ nervo non perspicuo. Cor. purpurascens. Styli rami longè appendiculati, appendice hispida stylum *Vernoniæ* ferè simulant. (v. s. comm. à cl. Nutt.)

CCCXCVIII. ACTINOLEPIS.

Capitulum pluriflorum, fl. radii 3-5 ligulatis foemineis, disci tubulatis 5-dentatis bisexualibus, stylo abortivo masculis? Invol. ovato-oblongum bracteis foliaceis paucis cinctum, squamis paucis obtusis adpressis dorso molliter tomentosis. Recept. angustum nudum. Ligulæ latæ breves 2-3-dentatæ, stylo bifido exserto. Tubuli tubo tereti, fauce dilatata, antheris albis, stylo simplici apice capitellato aut subnullo? Achænia oblonga subangulata, radii pubentia pappo squamellis circ. 5 scariosis acuminatis constante superata, disci glabra calva. — Herba annua tenella multicaulis Californica. Caules tenues apice subaraneosi. Folia opposita parva latè obovata, basi cuneata et integra, apice obtusissima 5-dentata. Capitula in axillis supremis approximatis solitaria

subcongesta parva. Flores flavi. — An *Helenia* ex pappo radii squamellato? An *Parthenia* si discus verè masculus?

1. *A. MULTICAULIS* ① in Californiâ legit cl. Douglas. Radix exilis simplex. Caulis circ. 2-pollic. Folia 2-3 lin. longa et lata, superiora araneosa. Capitula cano-tomentosa. *Achænia* nigra. (v. s. comm. ab hon. soc. hortie. Londin.)

CCCXCIX. BAHIA. — Bahia et *Eriophyllum* Lag. nov. gen. in *elench. h. madr.* 1815. p. 28. et 30. Less. syn. p. 238. — *Trichophyllum* Nutt. gen. am. (1818) 2. p. 166. nec Ren. — *Phialis* Spreng. gen. 2 (1831), n. 631. — *Helenii* sp. Spreng. syst.

Capitulum multiflorum radiatum, ligulis fœmineis 5-11, fl. disci hermaphr. 5-dentatis extus glanduloso-subsetosis. Invol. ovati seu campanulati squamæ adpressæ 1-2-seriales. Recept. epaleaceum nudum aut parcè alveolato-fimbrilliferum. Styli disci rami obtusi exappendiculati. *Achænia* turbinato- aut lineari-tetragona, ferè semper glabra. Pappi paleæ 4-8 ovales aut oblongæ obtusiusculæ membranacæ. — Herbæ perennes aut suffrutices Americani. Folia opposita aut alterna, integra aut partita, suprâ glabra aut araneosa, subtus tomentosa. Pedunculi nunc elongati 1-cephali, nunc apice corymbosi. Flores lutei, saltèm in genuinis, purpurei in specie dubiâ. — Genus fortè dividendum, sed car. generum nunc propositorum non sufficiunt.

§. 1. *Foliis alternis indivisis.*

1. *B. ANGUSTIFOLIA* (Less. in linnæâ 6. 1831. p. 519 sub *Trichophyllo*), erecta, ramis hirsutis, foliis alternis subsessilibus lanceolato-linearibus subserratis scabris, pedunculis elongatis glabris. 2? in Monte-Video. Ageratum angustifolium Spreng. syst. 3. p. 446. Capitula radiata lutea. An sequentium, ipso monente Less., omninò congener?
2. *B. STÆCHADIFOLIA*, suffrutescens?, foliis alternis linearibus obtusis margine subrevolutis suprâ araneosis glabrativè subtus cano-tomentosis, capitulis corymbosis, invol. squamis et pappi paleis 8-10. 2? in Mexico (Lag.), prope Real del Monte (Hænk.). *Eriophyllum* stæchadifolium Lag. *elench. h. Madr.* p. 28. n. 361. *Helenium* stæchadifolium Spreng. syst. 3. p. 574. Folia ex Lag. interdum opposita, in specim. obvio alterna, sed partes inferiores desid. *Achænia* elongatum subvillosum. Pappi paleæ oblongæ vix obtusæ. (v. s. in h. Hænk. à cl. de Sternb. miss.)
- β. Californica*, capitulis paucioribus, invol. pedicellisque cano-tomentosis. — in Novâ-Californiâ legit Douglas. An fortè distinguenda? (v. s. comm. ab hon. soc. hort. Lond.)
3. *B. INTEGRIFOLIA* (Hook. fl. bor. am. 1. p. 316. sub *Trichophyllo*), foliis inferioribus obovato-spathulatis, superioribus lineari-spathulatis integris integerrimisque vel rarissimè apice subtrifidis utrinquè albo-lanatis. — in Amer. bor. occid. cum *B. lanatâ*.

§. 2. *Foliis oppositis trilobis tripartitisve.*

4. *B. SINUATA* (Less. in linnæâ 1831. p. 160), fruticosa, foliis oppositis triplinerviis sæpius trilobis basi truncatis suprâ pubescentibus subtus incano-tomentosis. 2? in Mexici rupibus prope Hacienda de La Laguna.
5. *B. OPPOSITIFOLIA* (Nutt. gen. am. 2. p. 167 sub *Trichophyllo*), decumbens ramosissima pubescens, foliis oppositis palmato-trifidis, lobis ligulatis simplicibus vel divaricato-subdivisis, pedunculis filiformibus plurimum dichotomis vix folio longioribus. 2? in collibus sterilibus Amer. bor. circa Fort-Mandan. *Helenium* oppositifolium Spreng. syst. 3. p. 573. Caulis sulcatus. *Herba* amara (nec aromatica), 6-12-pollic. Ligulæ brevissimæ.

6. *B. ARTEMISIOIDES* (Lag. l. c.), fruticosa, foliis oppositis tripartito-multifidis pubescentibus, lobulis subobtusis. ♂ in Chili fruticetis rupestribus maritimis (Bert.!), circa Conceptionem (Lag.), Valparaiso (Bert. Gay! Macr. Gaud.!). Vulgò *Manzanilla-Simarona* dicitur. *Achyropappus maritimus* Poepp. pl. exs. n. 238. Ligulæ albæ elliptico-oblongæ discum longè excedentes. (v. s.)

§. 3. *Foliis alternis pinnatilobatis.*

7. *B. ARTEMISIÆFOLIA* (Less. in Linnæa 1829. p. 160. 1831. p. 258), suffruticosa, foliis alternis pinnatipartitis supra subglabris subtus cano-tomentosis, lobis 2-3-jugis integris, capitulis corymbosis, invol. ovati glabriusculi squamis oblongis. ♂ in Californiâ septentr. (Ménz.), ad Sanctum-Franciscum (Chamisso!). Ligulæ ovales parvæ. Invol. cylindraceum 1-2-seriale, squamis subæqualibus oblongis subconvolutis 2 lin. longis. Recept. planum alveolato-fimbriatiferum. Pappi paleæ lineares. (v. s.)

♂. *Douglasii*, pedicellis involucrisque magis tomentosis, capitulis duplò ferè majoribus, ligulis etiã majoribus. ♂ in Californiâ legit Douglas. (v. s. comm. ab hon. Soc. hort. Lond.)

8. *B. CONFERTIFLORA*, caule herbaceo erecto araneoso aut densè brevissimèque tomentoso, foliis alternis pinnatipartitis, lobis linearibus integris subtus sub-tomentosis, capitulis confertè corymbosis, invol. ovati squamis obovatis retusis arachnoideo-tomentosis. ♀ in Novâ-Californiâ legit cl. Douglas. Folia ferè *B. lanatæ*, sed angustiora. Capitula confertè corymbosa nec solitaria, duplò minor, magis ovata. Ligulæ paucæ obovatæ. (v. s. comm. ab hon. Soc. hort. Londin.)

9. *B. TENUIFOLIA*, herbacea erecta tota pube adpressâ aut subaraneosâ canescens, foliis alternis apice pinnatipartitis margine revolutis, rachi lineari integrâ, lobis 1-3-jugis linearibus dentatis integrisve, corymbo laxo, invol. tomentoso, ligulis obovatis inv. duplò longioribus. ♀ in Californiâ legit cl. Douglas. Habitus lanugine exceptâ *Senecionis artemisiæfoliæ*. Capitula parva aurea. Pappi squamellæ oblongæ vix acutæ. (v. s. comm. ab hon. Soc. hort. Lond.)

10. *B. LANATA* (Nutt. gen. am. 2. p. 168 sub *Trichophyllo*), decumbens seu erecta parçè ramosa, caule tereti foliisque subtus lanuginoso-tomentosis, foliis alternis pinnatifidis, summis linearibus integris, pedunculis elongatis 1-cephalis tomentosis. ♀ ad fontes flum. Columbia (Pursh! Dougl.) et Kosokosky (Hook.). *Actinella lanata* Pursh fl. bor. am. 2. p. 560. *Eriophyllum cæspitosum* Dougl. in Lindl. bot. reg. t. 1167. *Helenium lanatum* Spreng. syst. 3. p. 574. *Phialis* Spreng. gen. 2. p. 634 excl. car. Involucrum enim non verè gamophyllum sed squamis liberis 1-seriatis. Herba pedalis *Hymenopappum artemisiæfolium* aut *Bahiam artemisiæfoliam* referens. Ligulæ magnæ. (v. s. comm. ab hon. Soc. hort. Lond.)

11. *B. LEUCOPHYLLA*, tota cano-lanuginosa, caule herbaceo erecto, foliis oppositis et alternis obtusè sinuato- aut pinnatilobatis, lobis foliisque summis dentatis, pedunculis elongatis 1-cephalis, capitulis multi-(25-30) floris, achæniis glaberrimis tetragonis, pappi paleis octo, 4 lanceolatis acutis, 4 minimis alternantibus. ① ♂ ad Nootka et Mullgrave Amer. occid.-bor. leg. cl. Née! et Hænke! Ex specim. malè servato priori videtur affine. Herba 8-10-pollicaris *Seneciones* incanos referens, tota, etiã in invol., lanugine densâ vestita. (v. s.)

12. *B. ACHILLÆOIDES*, basi fruticosa ramosa, ramis virgatis teretibus tomentosis apice nudis parçè ramosis, foliis alternis basi cuneato-attenuatis apice pinnatipartitis, lobis 2-4-jugis approximatis dentatis supra araneosis subtus tomentosis, capitulis ramulos terminantibus solitariis, invol. subgloboso lanato. ♂ in Californiâ legit cl. Douglas. Folia 7-9 lin. longa apice cristato-partita, lobulis apice callosis. Ligulæ oblongæ flavæ. Pappus squamellatus parvulus. (v. s. comm. ab hon. Soc. hort. Lond.)

13. *B. TROLLIIFOLIA*, tota subincana, caule herbaceo, foliis alternis, infer. tripartitis multifidis dentatis, super. indivisis linearibus, pedunculis elongatis

1-cephalis, invol. 9-11-phyll. pappi paleis 4 lanceolatis acutis canaliculatis. 2½ in Mexico legit cl. Née. *Eriophyllum trolliifolium* Lag. *elench. h. Madr.* p. 28. n. 360. *Helenium trolliifolium* Spreng. *syst. 3. p. 573. Cor. 9-11 purpureæ* (Lag.). An verè congener?

CD. HYMENOPAPPUS *L'Hér. diss. cum icon. sine anno. Cass. dict. 55. p. 266 et 270. Less. syn. 238. — Rothia Lam. journ. h. nat. 1792. v. 1. p. 16. t. 1 non Schreb. — Hymenopappi sp. Juss.*

Capitulum homogamum. Recept. parvum nudum foveolatum. Invol. squamæ 2-3-seriales elliptico-oblongæ obtusissimæ, interiores interdum majores petaloideæ. Cor. tubo dilatato extus glanduloso-setigero, limbo campanulato 5-fido. Anth. exsertæ. Styli lobi apice breviter appendiculati. Achænia turbinata striata basi attenuata subpedicellata, matura subtetragona. Pappus paleis circ. 12-16 membranaceis brevibus elliptico-oblongis obtusis 1-serialibus. — Herbæ Carolinianæ perennes. Caules sulcato-angulati ramosi infernè glabri, supernè tomentosi aut velutini. Folia alterna pinnatipartita, lobis sæpè incisis aut multifidis. Capitula corymbosa pedicellata. Corollæ et inv. squamæ majores albidæ.

1. *H. tenuifolius* (Pursh fl. bor. am. suppl. p. 742), foliis glabriusculis subbipinnatipartitis, lobis linearibus integris, corymbo composito subconferato, invol. squamis 5-6, interioribus vix flosculis superantibus, achæniis villosissimis. ② (Nutt.) in Louisiana superiore (Pursh), ad Red-River (Nutt. ! herb.) et ad ripas fl. Missouri in collibus glareosis (Nutt. gen. am. 2. p. 139). A sequente certè videtur distinctus. (v. s.)
2. *H. scabiosæus* (L'Hér. ! l. c.), foliis subtus caudicanti-lanuginosis pinnatipartitis, lobis lineari-oblongis vix subdentatis, corymbo simpliciusculo laxo, invol. squamis interioribus 10 majoribus membranaceo-petaloideis flosculos superantibus, achæniis parçè villosis. 2½ (Ell.) in Carolinâ (Michx. ! Fras. !) et Georgiâ (Ell. sketch 2. p. 313). Michx. ! fl. bor. am. 2. p. 104. Pursh fl. 2. p. 519 et 742. *Rothia Carolinensis* Lam. l. c. et ill. t. 667. *Hym. laxiflorus* L'Hér. ! mss. (v. s.)
3. *H. artemisiæfolius*, foliis subtus cano-tomentosis, infer. in petiolum longè attenuatis oblongis sinuato-dentatis, cæteris sessilibus pinnatipartitis, lobis 3-4-jugis integris acutis cum terminali majore, corymbo composito laxo, invol. squamis ovalibus disco vix longioribus, achæniis villosissimis. ② in Mexicanâ prov. Texas legit cl. Berlandier (pl. exs. n. 153a). Bracteæ secus pedicellos obovatæ membranaceæ obtusæ. Cor. albidæ. Folia *Artemisiæ campestris* aut *Bahizæ artemisiæfoliæ*. (v. s.)
4. *H. douglasii* (Hook. fl. bor. am. 1. p. 316), lanâ albidâ deciduâ tomentosâ, ramis elongatis subcorymbosis, foliis bipinnatifidis, lobis linearibus sinuatis, capitulis solitariis, invol. squamis sub-20 linearibus pubescentibus disco brevioribus, corollis filiformibus apice pilosis pappi paleas lineares obtusas 10-12 vix superantibus. — in siccis arenosis Columbiz à loco dicto Great-Falls ad Rocky-Mountains. *H. scabiosæus* Dougl. mss. non L'Hér. (v. s. comm. sine nom. ab hon. Soc. hort. Lond.)
5. *H. filifolius* (Hook. fl. bor. am. 1. p. 317), lanâ albâ deciduâ tomentosâ, ramis paniculatis sparsis, foliis bipinnatifidis, lobis filiformibus elongatis integerrimis, capitulis solitariis, invol. squamis sub-12-ellipticis pubescentibus disco subbrevioribus, corollæ tubo supernè campanulato, pappo brevissimo. — in aridis flum. Columbiz ad Wella-Wellah et ad ripas flum. Spokan et Flat-Head Amer. bor.-occid. *H. tenuifolius* Dougl. mss. non Pursh.

† *Species exclusæ.*H. ANTHEMOIDES Juss. = *Hymenoxys anthemoides*.H. GLAUCUS Spreng. = *Cephalophora glauca*.H. PEDATUS Lag. = *Florestina pedata*.

CDI. CHÆNACTIS.

Capitulum multiflorum homogamum, fl. radii amplis basi tubulosis sursum subbilabiatis aut subpalmatis, disci tubulosis regulariter et obtusè 5-dentatis. Invol. campanulatum circ. 20-phyllum, squamis erectis 1-2-seriatis linearibus acutiusculis. Recept. epaleaceum alveolatum. Cor. tubo glabro, lobis hirsutulis. Styli disci rami hispidi cono obtuso brevi superati. Achænia lineari-tetragona basi attenuata subpedicellata adpressè pubescenti-villosa. Pappus paleis 5-6 membranaceis, radii brevibus obtusis, disci longioribus subacutis corollæ tubi dimidiam partem æquantibus. — Herbæ Californicæ erectæ habitu *Hymenopappum tenuifolium* referentes. Folia alterna pinnatipartita lobis linearibus integris. Rami apice nudi monocephali. Flores lutei. — Genus inter *Hymenopappum* et *Bahiam medium*; à priore fl. radii majoribus hiantibus, pappo squamellis 5-6 nec 15-16; à posteriore fl. radii non ligulatis nec fœmineis; ab utroque, pappo radii et disci diverso.

1. C. GLABRIUSCULA, caule ramoso, foliis involucriisque glabriusculis seu hinc inde subaraneosis, fl. radii palmatifidis expansis disco evidenter longioribus. 2/ 3 in Novâ-Californiâ legit cl. Douglas. Herba pedalis, junior araneoso-villosa, adulta glabra. Foliolorum lobi utrinquè 5-6 obtusiusculi. Folia summa linearia integerrima. Rami ad apicem per intervallum 2 1/2 poll. aphylli. (v. s. comm. ab hon. Soc. horticult. Lond.)

2. C. LANOSA, caulibus è collo plurimis foliis involucriisque cano-lanatis, floribus radii tubuloso-infundibuliformibus vix disco majoribus. (1) in Novâ-Californiâ etiam legit cl. Douglas. Radix exilis simplex. Folia rarissimè glabrata cæterum cano-lanata, summa linearia parva. Ramorum pars nuda 6 pollic. circ. longa. (v. s. comm. cum prior.)

CDII. POLYPTERIS Nutt. gen. 2. p. 139? Ell. sketch 2. p. 313? non Less.

Capitulum homogamum. Invol. squamæ biseriales foliaceæ ovali-lanceolatæ breviter acuminatæ, jun. erectæ demum reflexæ. Recept. hemisphæricum omninò nudum. Cor. tubo brevi, fauce amplâ inflatâ, lobis 5 subulatis hirtellis. Styli rami in appendicem linearem hirtam longè producti. Achænia turbinata sericeo-villosa. Pappi paleæ 9 achænio ferè longiores basi angustatæ membranacæ grossè uninerviæ, nervo in aristam longam rigidam producto. — Herba Boreali-Americana erecta ramosa pubescenti-scabra. Folia alterna integra oblongo-linearia sessilia. Capitula ramos terminantia solitaria. Cor. fuscae aut pallidè flavæ, lobis fuscis. Achæniî setæ et pappi aristæ fuscae. — Genus forsàn à *Cercostylo* non satis diversum.

1. P. INTEGRIFOLIA. 2/ ? in Carolinâ meridionali olim legit Fraser. P. integrifolia Nutt. à nostrâ differt invol. squamis oblongo-ellipticis membranaceis albidis, achænio subtetragono scabro nec villosa, pappi paleis 12-14. An verè ut vult Less. ad *Palafoxiam* referenda? P. integrifolia Ell. differt tantum achænio leviter scabro. Nostra certè ad *Palafoxiam* non referenda. Ex notulâ herb. L'Hér. ad *Hymenopappum* referebat cl. Michaux sed in Michx. fl. desideratæ. (v. s.)

† *Species exclusa.*

P. BRASILIENSIS Less. = *Cercostylos Brasiliensis*.

CDIII. ESPEJOA.

Capitulum 7-florum homogamum, fl. tubulosis 5-fidis glabris hermaphrod. Invol. 4-seriale, squamis 8 oppositis latè ovatis imbricatis apice mucronatis. Recept. alveolatum. Styli rami obtusi parcè puberuli. Anth. breves inclusæ. Achænia obcompressa-cuneata villis sericeis rufis confertissimis onusta, disco epigyno lato orbiculari. Pappus paleis circ. 16 uniserialibus membranaceis ovali-oblongis 1-nerviis mucronulatis corollæ ferè longitudinem adæquantibus, villis sericeis achæniæ basi cinctis. — Herba glabriuscula erecta dichotoma Mexicana. Caulis subangulatus versus apicem pilosiusculus. Folia opposita ovali-oblonga sessilia integerrima. Pedunculi axillares 1-cephali. Flores ex sicco flavo-subpurpurascens. — Genus Mexicanum dicavi cl. navigatori Hispano Antonio de Espejo qui anno 1533 plagas Mexicanas primus adiit.

1. *E. MEXICANA*. (1)? in Mexico ad Divum-Dionysium septemb. flor. legit cl. amic. Alaman. Stobæa glabrata var. Alam. in litt. 1831. (v. s.)

CDIV. *CERCOSTYLOS* Less. syn. 239. — *Polypteris* Less. in *linnæa* 1831. p. 518 nec Nutt.

Capitulum homogamum. Invol. campanulatum biseriale, squamis longè acuminatis. Recept. convexum epaleaceum densè paleaceo-fimbriliferum. Cor. 5-dentatæ. Styli rami appendice lineari angustissimâ. Achæmium turbinatum villosum. Pappi paleæ decem 1-nerviæ apice lacere in aristulam acuminatæ. — Herba Brasiliensis basi lignescens, foliis sessilibus vel decurrentibus integerrimis alternis pilosiusculocanescentibus, capitulis solitariis caulem ramosque elongatos et superne aphyllous superantibus. Cor. luteæ. — Unica species adhucdum descr.

1. *C. BRASILIENSIS* (Less. syn. 239). ♂ in Brasiliâ. *Polypteris Brasiliensis* Less. in linn. 1831. p. 518. Capitul. magn. Gaillardie lanceolatæ. Folia oblongo-obovata punctata, 1½-2 poll. lata, vix semilineam lata.

CDV. HOPKIRKIA. DC. non Spreng (1).

Capitulum 3-florum homogamum. Invol. oblongum 4-5-phyllum, squamis obovato-oblongis subscariosis, bracteolis minimis 2-3 basi stipatum. Recept. nudum angustum. Cor. tubulosæ pappo breviores inæqualiter 4-5-dentatæ. Styli rami breves apice capitellati. Achænia tetragona basi attenuata substipitata, angulis densè villosociliatis. Pappi squamellæ 8 scariosæ medio 1-nerviæ apice subtrifidæ nervo in aristulam productæ. — Herba Mexicana ramosa adscendens glabra. Folia alterna pinnatipartita, lobis 1-2-jugis cum impari angustè linearibus elongatis integerrimis. Capitula solitaria pedicellata terminalia parva. Flores lutei. — Cum *Hopkirkia* Spreng. sit omninè delenda, dicavi hoc novum genus cl. Thom. Hopkirk auctori Floræ anomoiæ, nè honore debito fraudetur.

1. *H. ANTHEMOIDEA*. — in Mexici agro Regiomontano legit cl. Henke. (v. s. in h. Henk. à cl. de Sternberg comm.)

(1) *Hopkirkia* Spreng. (nov. prov. 1818) est *Salmea* DC. (cat. h. monsp. 1813).

CDVI. HYMENOXYS *Cass. dict. 55. p. 278.*

Capitulum multiflorum discoideum, homogamum seu ligulis 8-10 uniserialibus fœmineis, fl. disci hermaphr. tubulosis 5-dentatis. Invol. biseriale, squamis rigidulis adpressis, interioribus longioribus. Recept. conicum epaleaceum alveolatum, glandulis parvis subonustum. Stylus disci ramis apice truncatis barbellatis. Achænia conformia turbinata villis sericeis erectis villosissima. Pappus paleaceus, paleis 5-8 inæqualibus membranaceis ovali-lanceolatis aristato-acuminatis erectis. — Herbæ Americane erectæ ramosæ glabriusculæ, tritæ odorem Anthemidearum spirans. Folia alterna aut in eodem caule rarius opposita, in lobos lineares multipartita et Schkuhriæ aut Achyropappi æmula. Rami apice nudi pedunculiformes. Cor. luteæ.

Sect. I. EUHYMENOXYS. — *Hymenoxys Cass. l. c.*

Capitula discoidea homogama.

1. *H. ANTHEMOIDES* (Cass. dict. 55. p. 278), invol. squamis obovatis subspathulatis. — in Bonariâ (Comm.). *Hymenopappus anthemoides* Juss. ann. mus. Par. 2. p. 426. Incautæ, ex accuratâ Cass. descr., cl. Jussieu pappum duplicem et Persoon folia decurrentia dixerunt. Capitula egregiè nota, sed herbæ deser. non sufficiens. — *Cephalophora anthemoides* Less. syn. 240 et in linnæâ 1831. p. 518. videtur ex capitulo radiato diversa.
2. *H. HÆNKELIANA*, invol. squamis ellipticis obtusiusculis. — in Peruvîâ legit cl. Hænke. An ab *H. anthemoides* mihi ignotâ satis distincta? Caulis teres divaricato-ramosus. Folia pinnatipartita, rachi lobisque linearibus integerrimis. Rami apice nudi et pedicelli ex axillis supremis orti 1-cephali apice striati. Capitula globosa. Recept. in centro conicum. Achænia tereti-turbinata longè sericea villosa. Pappi paleæ basi ovales in aristam cuspidatæ. Corollæ viz pappo longiores tubo brevitereti. Styli inclusi. (v. s. in h. Hænke, à cl. de Sternberg comm.)

Sect. II. OXYPAPPUS. — *Cephalophoræ* sect. 3. *Less. syn. 240.*

Capitula radiata.

3. *H. ODORATA*, invol. squamis lanceolatis acuminatis. ? in Mexico legit cl. Berlandier, sed sched. deperditâ, locus proprius ign. *Cephalophora anthemoides* Less. in Linn. 1831. p. 518. syn. 240 excl. syn. Juss. Folia multò tenuius pinnatipartita quàm prioris. Pedunculi foliis duplò triplòve longiores. Ligulæ 5-6 flavæ apice grossè tridentatæ. Pappi squamellæ lanceolatæ acuminatæ. Capitulum tritum valdè odorum. (v. s.)
4. *H. CHRYSANTHEMOIDES*, caule herbaceo? ramoso polycephalo glabro apice villosopubescente, foliis petiolatis subpubescentibus subbipinnatifidiatis multipartitis, lobis linearibus, ramis apice nudis 1-cephalis, invol. squamis biserialibus, pappi paleis 5 acuminato-subulatis, ? propè Moran Mexicanorum (H. B.) et in diluione Chalco (Andr. ! pl. exs. n. 316). Actinea chrysanthemoides H. B. et Kunth nov. gen. am. 4. p. 298. t. 411. Capitula flava magnitudine Leucanthemi. Achænia juniora lineari-cuneata sericeo-hirsuta. Ligulæ 12-15 apice 3-lobæ (v. s.)

CDVII. CEPHALOPHORA *Cav. ic. 6. p. 79. t. 599.* — *Grammia Hook. exot. fl. t. 189.* — *Cephalophoræ* sect. 1a. et 2a. *Less. syn. 240.*

Capitulum sphaericum homogamum aut radiatum, ligulis fœmineis tridentatis. Invol. squamæ biseriales reflexæ lineari-lanceolatæ flosculis æquales aut iis breviores. Recept. nudum foveolare. Cor. extus hispidula, tubo brevissimo, fauce obovoideâ tubulosâ, limbo 5-dentato

connivente subcalloso. Anth. inclusæ. Stylus filiformis. Styli rami exserti patulo-revoluti, apice hispiduli exappendiculati. Ovarium turbinato-4-gonum pilis squamosis adpressis tectum. Pappi paleæ 5-8 membranaceæ margine erosæ apice apiculatæ. — Herbaræ aut suffrutices Chilenses erecti glauci, caule striato ramoso brevissimè puberulo, foliis alternis oblongo-linearibus acutis. Capitula ramos nudiusculos terminantia. Cor. flavæ, juniores apice fuscæ. Involucri squamæ cum flosc. externis interdum mixtæ.

Sect. I. EUCEPHALOPHORA.

Capit. homogamum.

1. *C. GLAUCA* (Cav. l. c.), caule suffruticoso ramoso, foliis oblongo-lanceolatis, infer. inciso-serratis, summis integerrimis. 3 et ex Bert. variat ② in Chili propè Cucha-Cucha (Née!), in pascuis et fruticetis collium propè Valparaíso (Bert.!). *Ethulia repanda* Née! herb. *Hymenopappus glaucus* Spreng. syst. 3. p. 449. *Santolina tinctoria* Mol. chil. ed. gall. p. 336. *Santolinoides* vulgò *Poquill.* Feuill. obs. 3. t. 45. f. 2. Pedunculi teretes in omnibus specim. sylvestr. et etiã in Neëanis cultis, nec ut in ic. Cav. incrassati. Foliis exsiccatis ad colorem luteum utuntur Chilenses. (v. v. et s. sp.)
2. *C. PLANTAGINEA*, caule herbaceo sparsè ramoso, foliis radicalibus in petiolum attenuatis oblongo-lanceolatis basi cuneatis apice obtusis repando-dentatis, caulinis distantibus linearibus acutis integerrimis. ② in Chili? Radix subsimplex elongata. Rami circ. 3 valdè elongati. Pappus rufo-fuscus. (v. s. domin. à Mus. reg. Par. ex coll. Dombey sine loci design.)
3. *C. AROMATICA* (Schrad. ind. sem. h. Goett. 1830. Less. syn. 240), caule herbaceo ramosissimo, foliis lineari-lanceolatis subundulatis semiamplexicaulibus, infer. pinnatifidis, sup. vix subdentatis. ① in Chili (Hook.) in pascuis et fruticetis collium propè Valparaíso (Bert.!) ubi (ex odore aromatico ad Chamomillam accedente) dicitur *Manzanilla-del-Campo*. *Græmia aromatica* Hook. exot. bot. 2. t. 189. opt. (non Poepp. quæ *Euxenia Mitiqui* var. 2). *Cephaloph. tenera* Cass. opusc. 3. p. 93. Capitula dimidio quàm *C. glauca* minora cæterum similia. (v. s. sp. et v. c.)

Sect. II. *ACTINELLA* Less. syn. 240. — *Actinella Pers.* 2. p. 469. n. 1888. — *Actinea Juss. ann. mus. Par.* 2. p. 426. — *Actinea* et *Dugaldea Cass. dict.* 55. p. 270. — *Actinæ* sp. *Spreng.*

Capitula radiata, ligulis fœmineis apice 3-dentatis.

§. I. *Invol.* 1-3-seriali adpresso disco breviori aut subæquali. *Achæniis turbinatis.* — *Folia non decurrentia.*

* *Caule fruticoso folioso.*

4. *C. HETEROPHYLLA* (Less. syn. 240), caule fruticoso ramoso rufescente, foliis sessilibus lineari-lanceolatis, infer. grossè serrato-incisus, super. integerrimis, ramis apice nudis 1-cephalis, invol. squamis 1-2-serialis lanceolatis disco brevioribus, pappi paleis 8 oblongis aristatis. 3 in Bonariâ. *Actinea heterophylla* Juss. ann. Mus. 2. p. 425. t. 61. f. 1. *Actinella heterophylla Pers. ench.* 2. p. 469. *Actinea alternifolia Spreng. syst.* 3. p. 574. *Achænia oblonga villosa*, pilis biapiculatis. Ligulæ (albæ?) apice trilobæ Cass. dict. 55. p. 270.
5. *C. RADIATA* (Less. in linnæâ 1831. p. 516), caule basi fruticuloso ramosissimo glabro, foliis sessilibus linearibus integerrimis punctatis, ramis apice aphyllis 1-cephalis, invol. biserialis squamis ext. ovato-linearibus acuminateis, inter. oblongis vix mucronatis, pappi paleis oblongo-ellipticis aristatis. 2. 3 in campis ad Rio-Negro et Rio-Grande Brasiliæ. Folia pollicem longa,

lineam lata. Capit. magnitudine Anthemidis arvensis. Recept. hemisphæricum. Cor. disci limbo apice purpureo, ligulæ pallidè luteæ: Achænia turbinata fusco-villosa. Anth. obsol. caudatæ. (v. s. in h. Mus. reg. Par. ex. h. Mus. imp. Bras. sub n. 1013. 1042. 1051 et 1053.)

** *Caule herbaceo aphylo scaposo, foliis radicalibus.*

6. *C. ACAULIS*, scapo nudo elongato 1-cephalo, foliis radicalibus lineari-spathulatis integris punctatis villosis, invol. squamis circ. biseriatis oblongo-lanceolatis, pappi paleis 5-6 apice aristatis. \mathcal{L} in collibus Missouriensibus ad Fort-Mandan Amer. bor. Galaria acaulis Pursh. fl. bor. am. 2. p. 743. Actinella acaulis Nutt. gen. am. 2. p. 173. Capitula magnit. Chamomillæ, latè flava. Ligulæ 8-10 apice 3-dentatæ.

7. *C. SCAPOSA*, villosa, caule aphylo stricto simplici 1-cephalo, foliis radicalibus lineari-lanceolatis basi attenuatis, aliis integerrimis, aliis pinnatifidis lobis paucis acutis, invol. ext. squamis disco brevioribus obtusis adpressis. \mathcal{L} in Mexicanæ prov. Texas district. orientalibus legit cl. Berlandier (pl. exs. n. 1867 et 1882). Scapus 8-12 poll. longus. Folia 2-3 poll. longa, 2-3 lin. lata, acuta. Ligulæ 4-nerviæ 3-dentatæ. Pappi paleæ uninerviæ, margine scariosæ, ovals, abruptè aristatæ, cor. disci paulò breviores. (v. s.)

§. 2. *Invol. biseriale, squamis exter. linearibus reflexis. Achænia tetragona. — Folia ut in Helenio decurrentia:*

8. *C. DECURRENS* (Less. in linnæâ 1831. p. 517), caule herbaceo dichotomo pubero, foliis decurrentibus oblongis subrepandis puberis, infer. obtusis, super. acuminatis, ramis apice aphyllis 1-cephalis, invol. squamis linearibus reflexis, pappi paleis 6 ellipticis pinnatifidis aristatis. \mathcal{L} in Californiâ. An ex Less. Helenium pubescens Ait. kew. 3. p. 227? Ligulæ disci diametro breviores. Achænia tetragona. Anthæræ obsolete caudatæ.

† *Species exclusæ.*

C. ANTHEMOIDES Less. = *Hymenoxis odorata*.

ACTINELLA COMMUTATA Spreng. = *Calea pinnatifida*.

A. LANATA Pursh. = *Bahia lanata*.

CDVIII. *JAUMEA* Pers. ench. 2. p. 397. n. 1816. — *Kleinia* Juss. ann. Mus. 2. p. 424. t. 61. f. 1. non Linn. nec Willd.

Capitulum multiflorum discoideum homogamum. Invol. latum patens imbricatum, squamis subrotundis triplici ordine dispositis. Recept. nudum. Anth. ecaudatæ appendice ovato-lanceolatâ terminatæ. Styli rami vix ac ne vix subpapilloso cono brevi terminati. Achænia teretiuscula costis minimis longitudinalibus notata. Pappi squamellæ 8-10 breves inæquales denticulatæ acutæ. — Suffrutex Bonariensis glaber. Folia opposita connata linearia carnosâ integra. Capitula terminalia solitaria nutantia. Car. ex Juss. descr. et ic. nec non ex notis et iconè ineditis à cl. Adr. de Jussieu et Decaisne juxta specim. ipsum Commers. paratis et mecum benevolenter communicatis.

1. *J. LINEARIFOLIA* (Pers. l. c.). \mathcal{L} in Bonariâ ad ostium flum. Plata. — Flores ex sched. Commers. vix comparebant lutei.

CDIX. BURRIELIA.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœminicis obovatis, disci plurimis tubulosis 5-fidis hermaphroditis aut interdum abortu sterilibus. Involucri patenti-campanulati squamæ ovales acumi-

natæ disco paulò longiores uni- aut biseriales subalternæ æquales. Recept. epaleaceum. Cor. disci lobis extus tenuè barbellatis. Styli disci ramis cono brevi terminatis. Achænia gracilia lineari - subtetragona, radii ferè obcompressa pappo longè bi- seu triaristato, disci pappo 3-4-phylo, paleis lanceolato-aristatis corollam longitudine subæquantibus. — Herbæ Californicæ annuæ graciles parcè puberulæ aut glabræ subsimplices erectæ. Folia opposita linearia longa integerrima. Capitula ramos caulesve apice nudos terminantia. Flores aurei — Dixi in honorem cl. Joh. Marc. Burriel qui anno 1758 iter Venegasii in Californiam edidit et auxit.

1. B. GRACILIS, ligulis et invol. lobis 10-12. ① in Novâ-Californiâ legit cl. Douglas. Herba 5-6 pollic. Folia 2 poll. longa, lineam lata. Capitulum fere Coreopsisidis Bidentisive. (v. s. comm. cum seqq. ab hon. Soc. hort. Lond.)
2. B. TENERRIMA, ligulis et invol. lobis circ. 5. ① in Novâ-Californiâ legit cl. Douglas. Herba 3-4-pollic. Magis adhuc glabra et tenera quàm prior. Folia 6 lin. longa, vix $\frac{1}{2}$ lin. lata. Achænia disci 3-aristata. (v. s.)
3. B. MICROGLOSSA, ligulis et invol. squamis circ. 3. ① in Novâ-Californiâ legit cl. Douglas. Herba habitu B. tenerrimæ sed villosiuscula et primâ fronte discoidea. Folia $\frac{1}{2}$ -1 lin. lata. Ligulæ 2-3 brevissimæ integræ stylo suo superatæ. Achænia omnia 2-3-aristata. (v. s.)

CDX. LASTHENIA Cass. opusc. 3. p. 88 (1834). DC. in Lindl. bot. reg. t. 1780. — Rancagua Poepp. et Endlich. nov. gen. ac spec. pl. amer. x. p. 15. t. 24 a 25 (1835).

Capitulum multiflorum subhomogamum, fl. radii nempè fœmineis paucis parvis tubulosis obliquè truncatis (ex Cass.). Recept. conicum epaleaceum papillosum. Invol. squamæ uniseriales concretæ in cupulam 5-8-15-dentatam, dentibus acutis ciliatis. Corollæ breves, fauce inflatâ, limbo 5-dentato. Anth. inclusæ. Styli in radio exserti, in disco subinclusi, rami breves extus arcuati cono brevi superati. Achænia compressa adpressè pubescentia, areolâ basilari corneâ cavâ. Pappus paleis 5-10 apice dentato-laceris acutis corollam subæquantibus aut nullis. — Herbæ Occid.-Americana, glabræ aut subpilosiusculæ, erectæ, ramosæ, in uliginosis nascentes, Enhydræ ferè habitu. Folia opposita linearia subintegerrima ad apicem limbi et lobulorum callosobtusata. Rami nudi apice 1-cephali sub capitulo in conum inversum cavum dilatati. — Genus affine Gamolepidi ex involuero gamophyllo, Coinogyni ex foliis oppositis, ab utroque distinctum capitulo subhomogamo et pappo paleaceo; à Tagetineis differre videtur glandulis nullis.

§. 1. Pappo paleaceo.

1. L. GLABERRIMA, tota glaberrima, involuero 15-dentato, pappo 5-paleaceo. ① in uliginosis Californiæ legit cl. Douglas. Lasthenia Californica (DC. in Lindl. bot. reg. n. 1780 excl. car.) Cum error quidam per litteras in textu Lindleyano irrepserit et nunc altera species in Californiâ detecta fuerit, nomen deleo nè confusio persistat. (v. s. comm. ab hon. Soc. hortie. Londin.)
2. L. OBTUSIFOLIA (Cass. opusc. 3. p. 89), glaberrima, involuero 5-8-dentato, pappo 10-paleaceo. ① in Chili ad fossarum margines locis udis propè Rancagua legit cl. Bertero. Tagetes Feuillei Bert. ! pl. exs. n. 680. Colla mem. ac Turin 38. p. 34. n. 15. t. 30. sed à Tagete Chilensi Feuill. omnino differe videtur. Rancagua Feuillei Poepp. et Endlich. nov. gen. ac sp. amer. 1. p. 15. t. 24. Folia 2 poll. longa, integerrima vix obtusa (undè nomen spec. Cassinianum minùs aptum), summa 2 lin. longa dente grosso utrinquè mediâ

aucta. — Herba procumbens. Invol. 5-dentatum. Pappi paleæ lineares, corollâ breviores, apice lacero bi- vel trifida. Poepp. et Endlich. l. c. (v. s.)

β. *Bridgesii*, erecta, pappi corollam æquantis paleis ovato-lanceolatis duplicato-serratis. Rancagua Bridgesii Poepp. et Endlich. l. c. t. 25. An species distincta?

§. 2. *Pappo nullo.*

3. L. GLABRATA (Lindl. bot. reg. t. 1780), ramulis pedicellisque tenuissimè pubescentibus, invol. 15-dentato, pappo nullo, achæniis apice mucronulatis. ① in Californiâ legit cl. Douglas. Similis L. glaberrimæ sed non omnino glabra, capitula paulò majora, invol. basi vix attenuatâ et potiùs campanulatâ. (v. s. comm. ab hon. soc. hort. Lond.)

CDXI. PICRADENIA Hook. fl. bor. am. 1. p. 317. t. 108.

Capitulum multiflorum radiatum, ligulis 8-10 fœmineis obovato-oblongis grossè tridentatis, fl. disci hermaphr. 5-dentatis extûs glandulosis. Invol. subhemisphærici squamæ imbricatæ, externæ basi coadunatæ. Recept. conicum nudum. Styli disci rami inclusi breviusculi apice incrassati dorso hirsuti. Achænia oblonga villosa. Pappi paleæ 6-7 erectæ ovatæ acuminatissimæ nudæ membranacæ. — Herba Boreali-Americana. Radix crassa. Folia alterna lineari-filiformia rigida pinnatifida glanduloso- et pellucido-punctata. Capitula terminalia flava. Glandulæ cor. disci amarissimæ undè nomen. — Cl. auctor ad Helenieas suum refert genus et hîc ex ejus sententiâ admisi sed ex stigmate ab ipso Hookero delineato potiùs ad Astereas Achyrideas pertinere videtur.

1. P. RICHARDSONI (Hook. l. c. p. 317. t. 108). 2/ in Amer. bor. circa Carlton-House ad Saskatchewan.

CDXII. HECUBÆA.

Capitulum multiflorum, fl. radii ligulatis fœmineis 3-5-fidis, disci planiusculi fl. hermaphroditis 5-fidis, lobis brevibus fusco-hirsutis. Invol. biseriale, squamis exterioribus foliaceis patulis radiûm æquantibus, int. brevioribus linearibus discum vix paulò superantibus. Recept. nudum planiusculum. Styli rami apice truncati subcapitati hispiduli. Achænia angulata glabra glandulosa calva. — Herba Mexicana glabra pedalis simplex, caule tereti ad medium folioso apice nudo pedunculiformi 1-cephalo, pedunculo sub capitulo obconico fusco-subvillosa. Folia alterna decurrentia oblongo-lanceolata acuminata integerrima. Capitula majora et planiora quàm Helenii, ligulis flavis, disco fusco-purpurascente. — Genus Helenio valdè affine Hecubæam, ex Hecubâ Heleni matre, ob analogiam summam dixi.

1. H. SCORSONERÆFOLIA. 2/ in Mexico legit cl. Alaman qui cum plur. aliis communicavit et genus novum in schedulâ indicavit. (v. s.)

CDXIII. HELENIUM Linn. gen. n. 961. Lam. ill. t. 688. — Helenia Lin. hort. cliff. 418. Manch meth. 589. Gærtn. fr. 2. p. 438. t. 169. — Helenium et Tetrodus Cass. Less.

Capitulum multiflorum radiatum, fl. radii fœmineis 1-seriatis ligulatis aut rarò tubulosis subirregulariter 3-5-fidis, fl. disci hermaphrod. tubulosis brevissimè 4-5-dentatis. Recept. convexum globosum aut oblongum, nudum. Invol. squamæ biseriales, exter. plurimæ foliaceæ elongatæ sublineares reflexæ aut patentissimæ, inter. subpauciores.

breviores acuminatæ recept. paleas simulantes, sed extra flores sitæ. Styli rami obtusi exappendiculati. Achænia turbinato-obovata multifariâ villosa. Pappi paleæ 5-6 membranaceæ plus minus apiculatæ. — Herbæ Boreali-Americanae glabræ aut pubescentes. Folia alterna decurrentia subtus punctata integra dentata aut subpinnatifida. Capitula ad apices caulis et ramorum solitaria. Cor. luteæ, extus puberulæ.

Sect. I. HELENIA Linn. *Gærtn. Mænnch II. cc.* — *Helenium Cass. dict. 55. p. 262. Less. syn. 240.*

Cor. disci 5-dentatæ. Pappi paleæ apice aristatæ aut distinctè apiculatæ. Recept. convexum vel globosum.

1. H. AUTUMNALIS (Linn. spec. 1120), glabrum, caule erecto ramoso polycephalo, foliis lanceolatis serratis longè decurrentibus, ligulis planis 3-5-fidis, pappi paleis ciliato-subfimbriatis in aristam desinentibus cor. disci triplò brevioribus. $\frac{1}{2}$ in humidis secus rivulos frequens à Canadâ ad Carolinam. Schkuhr handb. 3. t. 250. Lam. ill. t. 688. Hook. bot. mag. t. 2994. *Helenia decurrens Mœnnch meth. 589. Helenia autumnalis Gærtn.* — *Corn. canad. p. 63. ic. (v. c. et s. sp.)*

β . *pubescens* (Pers. ench. 2. p. 475), totum brevissimè pubescens, caule erecto ramoso polycephalo, foliis lanceolatis grossè serratis decurrentibus, ligulis planis 3-fidis, pappi paleis basi ciliatis in aristam desinentibus cor. disci triplò brevioribus. $\frac{1}{2}$ in Amer. bor. H. pubescens Ait. kew. ed. 1. v. 3. p. 287? An, suadente cl. Persoon, mera var. hortensis H. autumnalis? An spec. propria? (v. s. c.)

2. H. MEXICANUM (H. B. et Kunth nov. gen. am. 4. p. 299?), glabriusculum, caule erecto ramoso polycephalo, foliis lineari-lanceolatis elongatis integerrimis decurrentibus, ligulis planis trifidis subtus pubescentibus, pappi paleis scariosis basi ovatis apiculatis cor. disci quintuplò brevioribus. $\frac{1}{2}$ in Mexici ditione Oaxaca legit cl. Andrieux (pl. exs. n. 3171). Achænia nervoso-angulata, secus angulos pilis albis adpressis vestita. Ligulæ disco breviores. (v. s.)

3. H. TUBULIFLORUM, minutè et sparè puberulum, caule erecto apice ramoso, ramis subnudis apice 1-cephalis, foliis membranaceis elliptico-lanceolatis subdentatis, ligulis tubulosis inæqualiter 5-fidis, pappi squamis lineari-lanceolatis acuminatis scariosis basi ciliatis cor. disci dimidiò brevioribus. — Ex Amer. bor. occid. semina in hortos Anglicos introduxit cl. Douglas. (v. s. in hort. Soc. hortic. lect. 1830.)

4. H. CANALICULATUM (Lam. journ. h. nat. 2. p. 213. t. 35), glabriusculum, caule erecto simplici 1-cephalo, foliis lanceolatis dentatis decurrentibus, ligulis concavo-canaliculatis apice tridentatis, pappi paleis acuminatis cor. disci triplò brevioribus. — in Amer. bor.?

Sect. II. TETRODUS. — *Tetrodus Cass. dict. 55. p. 262. Less. syn. 240.*

Cor. disci 4-dentatæ. Pappi paleæ breves. Recept. ovatum aut oblongum.

5. H. QUADRIDENTATUM (Labill. act. soc. h. n. Par. 1. p. 22. t. 4), glabriusculum, caule erecto apice ramoso polycephalo, foliis decurrentibus oblongo-linearibus, infer. pinnatifidis, super. integerrimis, ligulis glabris, invol. brevissimo, achæniis squamellis vix apiculatis. (†) in Amer. bor. nempe in Carolinâ (Fras.), Louisianâ (Labill.), Mexico (Berl. ! pl. exs. 436). Lam. ill. t. 688. f. 2. Don in Lindl. bot. reg. t. 598. *Rudbeckia alata Jacq. ic. rar. 3. t. 593. coll. 5. p. 154. Hel. Milleri Schult. ex rec. Salz. Mesodetra alata Raf. fl. ind. suppl. p. 141. (v. v. c. et s. sp.)*

6. H. VARIUM (Schrad. ! hort. Gœtt. sem. 1831. et in litt. 1832), glabrum, caule erecto subsimplici sublaæ ramoso, foliis decurrentibus subtrinerviis, caulinis

- integerrimis, infer. . . , invol. squamis radio brevioribus, ligulis trifidis sub-
tus vix subpuberulis, pappi squamellis aliis obtusis aliis brevissimè apiculatis.
① in Mexico (Alam.), eùam in horto Mexicano (Berlandier! pl. exs. n. 436),
et Gœttingensi (Schrad.), cultum. Folia 2-2 $\frac{1}{2}$ poll. longa, basi 4-8 lin. lata.
Capitula rotunda 5-6 lin. diam. (v. s.)
7. *H. PUBERULUM*, tenuissimè puberulum, caule erecto ramoso polycephalo,
foliis decurrentibus oblongo-lanceolatis integerrimis, infimis incisis, ligulis
trifidis pubero-velutinis, invol. brevissimo, pappi squamellis acuminatis. ① ?
in Californiâ legit cl. Douglas. Affine *H. quadridentato*, sed majus et ex omni
parte puberulum. Capitulum globosum 6 lin. diam. Ligulæ parvæ. Pedicelli
elongati striati. (v. s. comm. ab hon. Soc. hort. Londin.)
8. *H. HETEROPHYLLUM*, glabrum, caule erecto apice corymboso-ramoso polyce-
phalo, foliis decurrentibus, inf. obtusè pinnatifidis, mediis sinuatis serratisve,
summis integerrimis acutis, invol. squamis apice subulatis radio ferè longiori-
bus, ligulis planis 3-fidis, pappi paleis ovatis breviter aristatis obtusiusculis
cor. disci triplò duplòve brevioribus. ① in Mexico prov. Tamaulipas circa
Tampico legit cl. Berlandier (pl. exs. n. 107, 190 et 2113). Herba pedalis. Ca-
pitulum ovato-globosum diam. 3-4 lin. (v. s.)
9. *H. ELEGANS*, glabrum, caule erecto stricto apice corymboso-ramoso oligoce-
phalo, foliis decurrentibus lanceolato-linearibus integerrimis, utrinquè glandu-
loso-punctatis, infer. . . , invol. squamis apice subulatis radio subbrevio-
ribus, ligulis basi villosis apice obtusè 3-4-dentatis, pappi squamellis brevissimis
obtusis. ① ? in Mexico circa Bejar legit cl. Berlandier (pl. exs. n. 1651!) Li-
gulæ ex sicco basi purpureo-croceæ, cæterum luteæ. (v. s.)
10. *H. MICROCEPHALUM*, glabrum, caule erecto ramosissimo polycephalo, foliis
decurrentibus lanceolatis integerrimis subpunctatis, infer. . . , invol. brevis-
simo, ligulis tridentatis glabris, pappo brevissimo obtuso. — in Mexico inter
Lando et Bejar legit cl. Berlandier (pl. exs. n. 1429!). Ligulæ flavæ. Capitulum
subrotundum 3 lin. diam. (v. s.)

† *Species non satis notæ.*

11. *H. LONGIFOLIUM* (Smith. in Rees cycl. n. 2), foliis lineari-lanceolatis inte-
gris lævissimis, pedunculis undis elongatis. — in Amer. boreali. *H. autumnale*
Mill. dict. ed. 8. ex Sm. †.
12. *H. PUMILUM* (Willd. enum. h. Ber. suppl. p. 60), foliis oblongis subinteger-
rimis, invol. squamis lanceolatis patentibus. 2℥ patr. et cæt. car. ign. in h.
Willd. non reperitur (Link): †.
13. *H. QUADRIPARTITUM* (Link enum. h. Ber. 2. p. 338), foliis lanceolatis de-
currentibus, cor. radii quadripartitis. ② patr. et cæt. car. ign. An var. *T. qua-*
dridentati seu *Hel. autumnalis* ligulas sæpè quadrifidas gerentia? †.

†† *Species exclusæ.*

- H. LANATUM* Spreng. = *Bahia lanata*.
H. OPPOSITIFOLIUM Spreng. = *Bahia oppositifolia*.
H. PUBESCENS Ait. = *Actinella decurrens*?
H. STORCHADIFOLIUM Spreng. = *Bahia stœchadifolia*.
H. TROLLIIFOLIUM Spreng. = *Bahia trolliifolia*.

CDXIV. AMBLYOLEPIS.

Capitulum multiflorum radiatum, fl. radii fœmineis 1-serialibus
ligulatis apice 3-fidis, fl. disci hermaphrod. tubo brevi, fauce amplâ,
lobis 5 lanceolatis callosio-apiculatis nervo marginatis. Invol. biseriale,
squamis ser. ext. circ. 6-7 ovatis acutis foliaceis adpressis disco æqua-
libus, ser. int. obovato-rotundis obtusissimis hyalinis enerviis. Recept.

planiusculum alveolatum. Antheræ longè et acutè appendiculatæ. Styli rami exappendiculati. Achænium turbinatum adpressè villosissimum. Pappi paleæ 5 uniseriales obtusissimæ enerviæ corollæ tubo æquales. — Herba Mexicana annua erecta simplex, hinc indè secus caulem margines foliorum et squamarum invol. exteriorum pilis longis ciliatopilosa. Folia alterna semiamplexicaulia nec decurrentia ovali-lanceolata subintegerrima. Capitula terminalia solitaria flava, ferè Pyrethri Myconis magnit. et facie. — Nomen (ex ἀμβλυσ obtusa et λεπίς squama) alludit ad squamas invol. exteriores et ad paleas pappi obtusas.

1. A. SETIGERA. ① in Mexicanâ prov. Texas inter. Bejar et Austin legit cl. Berlandier (pl. exs. n. 16131). Herba pedalis paucifolia. Folia inf. obtusa, sup. acuminata. (v. s.)

CDXV. ARGYROXIPHIMUM.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci hermaphrod. tubulosis 3-dent. Recept. nudum planum. Invol. campanulatum, squamis 2-3-serialis lanceolato-linearibus subæqualibus disco paulò brevioribus. Ligulæ obovato-cuneatæ apice latè incisæ. Styli stigm. 2 divergenti-revoluta subfiliformia, in disco apice pilosiuscula. Antheræ ecaudatæ apice appendiculatæ. Achænium elongatum glabrum ad angulos 2 vix subciliatum compressum, latere uno alterove nervo seu angulo subxserto instructum (in specim. mihi obv. cassum). Pappus persistens 1-serialis paleaceus radii auricularis ad latus exterius situs acuminatus integer, disci paleis 2-5 irregularibus serratis subfoliaceis rigidis. — Herba Sandwicensis crassicaulis Tussilagines, Cinerarias seu Ligularias Europæas texturâ referens. Folia alia è basi subrigida elongata subensiformia integerrima villis sericeis adpressis copiosissimis argentea, alia basi teneriora virescentia pube glandulosâ instructa. Involucra etiâ hirta et fol. supremis texturâ similia. Pedunculi ferè ex axillis foliorum orti solitarii glanduloso-hirti subaphylli 1-cephali. Capitula subglobosa nutantia.

1. A. SANDWICENSIS. 2 in insulis Sandwicensibus ad Owhyhee floridam mense junio legit cl. Macrae. Herba mirabilis et distinctissima circ. bipedalis. Folia argyræa pedem longa, 3-4 lin. lata. Capitula ferè diam. pollic., ex siccò fortè purpurascencia. (v. s. in herb. cl. et am. Lindley.)

CDXVI. ROSILLA Less. syn. 245.

Capitulum radiatum, ligulis fœmineis. Invol. cylindracei squamæ scariosæ, exter. sublineares. Recept. epaleaceum. Styli rami appendiculati. Achænium obcompressum basin versus angustatum. Pappus paleis plumoso-fimbriatis. — Herba annua Mexicana, floribus luteis.

1. R. LUTEA (Less. l. c.) ① in Mexico. Cæt. ign.

Div. II. GALINSOGÆ. — Galinsogæ et Caleinæ Cass. dict. 55. p. 264 et 265. Less. syn. 241 et 245.

Pappus 1-serialis, paleis integris aut plumosis constans, rarissimè nullus. Recept. paleaceum. Invol. squamæ planiusculæ, achænia radii non involventes.

Subdiv. 1. EUGALINSOGÆ.

Ligulæ fœmineæ sen nullæ. Pappi squamellæ planiusculæ nec æstivato-contortæ.

CDXVII. LEMMATIUM. — *Caleacte* Less. in *linnæa* 1830. p. 158. syn. 248. non Brown.

Capitulum 6-7 florum heterogamum, fl. radii circ. 3 ligulatis foemineis, disci etiā 3-4 tubulosi hermaphrod. 5-fidis, lobis longiusculis acutis glabris. Invol. cylindricum, squamis imbricatis oblongis, exter. subfoliaceis, inter. membranaceis. Recept. angustum, paleis convolutis lineari-oblongis flores involventibus et eos æquantibus. Styli rami exappendiculati. Achænia trigono-prismatica ad angulos villosa. Pappus membranaceus vaginans apice dentatus, squamellis concretis verisim. constans, cor. faucem adæquans. — Frutex Brasiliensis. Caulis densè et breviter hirsutus teres seu obtusè angulatus. Folia opposita sessilia subrotunda basi cordata crenato-dentata coriacea, suprā puberula, subtus tri- aut quintuplinervia reticulata villosa-scabra. Capitula subsessilia in corymbum amplum valdè compositum fastigiatum digesta, bracteis paucis foliaceis instructa. Flores lutei. — Genus habitu et stigmatibus *Lasiantheæ* affine sed achæniis et corollæ lobis diversum, et potiùs *Calydermo* et *Caleæ* affine, ab omnibus *Helenieis* pappi squamellis in vaginam concretis distinctum, sed tamen meo sensu iis potiùs quàm *Chrysantheméis* adjungendum. Nomen ex λεμματιον vaginula.

1. L. ROTUNDFOLIUM. $\frac{5}{2}$ in Brasiliæ prov. Minarum-Generalium legit cl. Vauthier (pl. exs. n. 284). *Caleacte rotundifolia* Less. ll. cc. Folia 2 $\frac{1}{2}$ poll. diam. (v. s.)

CDXVIII. CALYDERMOS Lag. nov. gen. p. 24. — *Calydermos* et *Calebrachys* Cass. dict. 55. p. 265 et 277. Less. syn. p. 225 et 241. — *Caleæ* sp. Kunth.

Capitulum multiflorum homogamum. Invol. oblongum imbricatum, squamis subscariosis obtusissimis. Recept. angustum conicum, paleis linearibus acutis. Cor. tubulosæ 5-fidæ. Styli rami cono terminati. Achænium 4-5-angulatum pilosum. Pappi paleæ 1-seriales lineares obtusæ 1-nerviæ aut nullæ. — Herba Mexicanæ erectæ scabræ aut pilosæ parvæ ramosæ. Folia opposita sessilia ovata aut elliptico-acuminata grossè serrata trinervia. Rami longè aphylli, apice breviter trifidi, capitula fasciculato-corymbosa gerentes. Invol. sublutescentia. Flores lutei.

Sect. I. EUCALYDERMOS. — *Calydermos* Lag. Cass. et Less. ll. cc.

Achænia calva.

1. C. SCABER (Lag. nov. gen. et sp. n. 323), caule hispidulo, foliis ovato-lanceolatis utrinquè scaberrimis subtus præcipuè villosis, achæniis calvis. $\frac{2}{2}$ in Mexici montanis circa Guanajuato, Acapulco et Real del Monte (Lag.), Villalpando (Ménd.), et urbem Mexici (Berl.) pl. exs. n. 646). Herba pedalis et sesquipedalis. An fortè *Caleæ* peduncularis var. epapposa H. B. et Kunth 4. p. 296. t. 408. f. 5 huc referenda? (v. s.)
2. C. LONGILOBIUS (Lag. nov. gen. et sp. n. 324), caule subpiloso, foliis oblongo-lanceolatis lævisculis. $\frac{2}{2}$ in Mexico. Variat foliis oppositis et ternato-verticillatis.

Sect. II. CALEBRACHYS. — *Calebrachys* Cass. Less. ll. cc.

Achænia pappo 1-seriali paleis 10-12 oblongo-linearibus obtusis coronato.

3. C. PEDUNCULARIS, caule hirtio, foliis inf. brevissimè petiolatis, sup. sessilibus ovato aut oblongo-lanceolatis utrinquè pubescenti-hirtis, achæniis paleis 8-10

coronato, stylo basi bulboso. 4? ①? in Mexici monte ignivomo Jorullo. *Calea peduncularis* H. B. et Kunth nov. gen. am. 4. p. 295. t. 408 (excl. fig. 5). Less. in linnæa 1830. p. 157 et 158. *Calebrachys peduncularis* Cass. et Less. Herba pappo excepto primæ speciei simillima et forsân à nonnullis cum eâ confusa. (v. s. ex h. Née.)

4. *C. SALMERFOLIUS*, caule suffruticoso, ramis teretibus glabriusculis, foliis brevissimè petiolatis ovatis acuminatis trinerviis acutè pauciserratis glabris suprà lævisculis subtùs crebrè glandulosis, capitulis pedicellatis corymbosis ovatis 8-10-floris, invol. squamis obtusis, exter. brevissimis adpressis. 3 in Mexico (Hænk.! herb.), ad Tamaulipas (Berl.! pl. exs. n. 2135). Caulis ex sicco fortè subscandens. (v. s.)
5. *C. rugosus*, ramis glabriusculis vel ad apicem densè hirtellis, foliis subsessilibus ovatis acutis serratis reticulato-rugosis vix suprà subpuberulis subtùs hirtello-velutinis parcissimè glandulosis, capitulis pedicellatis corymbosis ovato-oblongis 4-5-floris, invol. squamis subacutis, exter. brevibus. 3 in Mexico (Hænk.!), ad Cuernavaca (Berlandier pl. exs. n. 1061!). Frutex 6-8-pedalis, octob. flor. Variat foliis acuminatis et vix acutis. (v. s.)

† *Species exclusa.*

C. JAMAICENSIS Spreng. = *Eupatorium Dalea*.

C. REPENS Spreng. = *Oiospermum involucreatum*.

CDXIX. MEYERIA DC. non Schreb.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis. Invol. campanulatum seu ovatum, squamis pluriserialibus adpressè imbricatis ovatis. Recept. planiusculum, paleis apice sæpè tridentatis complicatis flosculos amplectentibus. Styli disci ramis cono terminatis. Achænia prismatica villosa. Pappus squamellis plurimis 1-serialibus oblongis ellipticisve obtusis membranaceis constans. — Frutices suffruticesve Brasilienses. Folia opposita. Capitula solitaria aut rariùs corymbosa. Flores lutei. — Cum Meyera Schreberi ex jure prioritatis in Enhydram ceciderit, nolui sine honore debito gentem naturæ curiosorum plurimorum et inclytorum feracem relinquere, et hoc novum et pulchrum genus collectivè dicavi: 1° Joh. Meyer cui genus Schreberianum ex Boehmer (lex. bot. p. 134) dictum erat. 2° Frid. Alb. Ant. Meyer entomologo celebri et corticis Angusturæ descr. 3° Georg. Frid. Guil. Meyer lichenographo et floræ Essequiboensis auctori, 4° Car. Ant. Meyer Caucasicæ floræ et famil. Cyperacearum scrutatori, 5° Ern. Meyer (cui jam Ernestia dicata) floræ Labradoricæ et familie Juncorum illustratori.

Sect. I. HOLOPHYLLÆ.

Capitula terminalia solitaria. Involucrum campanulatum. Folia integerrima.

1. *M. MYRTIFOLIA*, glaberrima, foliis brevissimè petiolatis ovato-lanceolatis trinerviis basi obtusis apice acutis utrinquè lævibus, pedicellis folio longioribus, invol. squamis ovatis, exter. foliaceis brevibus. 3 in Brasiliæ prov. Minas-Generaliam prope Tejuco legit cl. Vauthier (pl. exs. n. 370!). Pedicelli sub capitulo minutissimè puberuli. Folia magnit. Myrti communis majoris. Achænia radii 3-gona, disci 4-gona, glabra, angulis radii lateralibus, disci interioribus ciliatis. (v. s.)
2. *M. PARVIFOLIA*, glaberrima, foliis sessilibus oblongo-lanceolatis acutis basi obtusis suprà lævibus subtùs reticulato-venosis, ramis ad apicem ferè foliosis, invol. squamis ovatis, exter. foliaceis interiores subæquantibus. 3 in Brasiliæ

prov. Sancti-Pauli. Folia 5-6 lin. longa, 2-3 lin. lata. Ramuli floridi tenuissimè puberuli. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 406.)

3. *M. LONGIFOLIA*, glaberrima, foliis sessilibus lanceolatis subobtusis utrinquè lævibus uninerviis erectis, ramis ad apicem usquè foliosis, invol. squamis inter se longitudine æqualibus, exter. lanceolatis foliaceis, inter. elliptico-oblongis. § in Brasiliæ prov. Sancti-Pauli. Rami (in sicco!) glaucescentes. Folia sesquipollicem longa, 4 lin. lata. (v. s. in h. Mus. reg. Par. ex h. imp. Bras. sub n. 407.)

Sect. II. GLUPHIPHYLLÆA.

Capitula in corymbum oligocephalum conferta. Involacrum ovatum laxum. Folia grossè et obtusè crenata.

4. *M. HISPIDA*, caule ramosissimo hispidissimo, foliis sessilibus latè ovatis subrotundis grossè crenatis utrinquè sparsè puberis, ramis ad apicem usquè foliosis, capitulis subcorymbosis brevissimè pedicellatis, invol. squamis exter. foliaceis subrotundis interiores adæquantibus. § in Bras. prov. Sancti-Pauli campis editis legit cl. Lund. Species rara novemb. florens. Folia diam. pollicari crenis latibus obtusis utrinquè 5-7 notata. (v. s. comm. à cl. Lund et ex h. imp. Bras. sub n. 405 ?)

CDXX. CALLILEPIS.

Capitulum multiflorum heterogamum, fl. radii plurimis 1-seriatis ligulatis fœmineis, disci tubulosis 5-dentatis hermaphr. Invol. campanulatum, squamis pluriserialibus lanceolatis subæqualibus. Recept. planum, paleis coriaceo-scariosis complicatis acuminatis flores includentibus. Styli rami cono brevi terminati. Achænia glabra, radii trigono-compressa disci plano-compressa. Pappus squamellis 3 scariosis acuminatis inæqualibus persistentibus, adjectis in disco hinc indè 1-2 minimis. — Suffrutices in Africâ austr.-orient. indigeni erecti pedales. Caules à basi plurimi aut solitarii. Rami apice 1-cephali. Folia inf. opposita, super. alterna, lanceolata acuminata integerrima aut subserrata. Capit. diametro ferè pollicari. Ligulæ 20-30 (ex sicco albæ?), planæ.

1. *C. GLABRA*, glaberrima, caule ramosissimo, foliis plerisque oppositis, summis tenuiter serrulatis. § in Africâ austr.-orient. ad Omsanculo legit cl. Drege! n. 5115. Rami tennes striati. Folia pollicem longa modicè acuminata. (v. s.)
2. *C. LAUREOLA*, ramis molliter hispidulis, foliis glabris plerisque alternis integerrimis, super. valdè acuminatis. § cum priore legit cl. Drege! n. 5115. An hujus varietas? sed præter characteres indicatos differt caule simplici aut parcè ramoso, invol. squamis exter. lineari-subulatis longioribus. Folia 20-24 lin. longa, 6 lin. lata, erecta, approximata. (v. s.)
3. *C. HISPIDA*, caulibus à collo plurimis simplicibus molliter villosis-hispidis, foliis margine subpilosis integerrimis plerisque oppositis, super. alternis remotis acuminatissimis angustissimis. § in Africâ austr.-orient. ad Zw. Omsanculo et Omtendo legit cl. Drege! n. 5114. Folia sesquipollicem longa, 5 lin. lata. (v. s.)

CDXXI. *CALEA* Brown trans. lin. soc. 13. p. 109. Less. syn. 241. — *Calea* sp. Linn. Swartz. Kunth. — *Moçinna* Lag. nov. gen. p. 31 non Spreng. nec Ort.

Capitulum multiflorum nunc homogamum nunc radiatum, ligulis fœmineis. Invol. ovati aut rariùs campanulati squamæ imbricatæ sæpiùs obtusæ siccæ. Recept. plùs minùs conicum, paleaceum. Styli 4.

hermaphr. rami exappendiculati. Achænium teres compressum angulatumve. Pappus multipaleaceus, paleis 5-20 sæpius circ. 10 linearilanceolatis scariosis acuminatis uninerviis subæqualibus. — Frutices (rarius herbæ) ex Americâ æquinoctiali indigeni. Folia opposita, rarisimè terna, sæpius 3-nervia, scabra integra aut dentata. Capitula terminalia aut axillaria. Flores lutei.

Sect. I. AMPHICALÆA.

Capitula discoidea 4-flora. Receptaculum epaleaceum. — An genus proprium? sed habitus valdè ad *C. cymosam* accedit. Fortè flores omnino marginales et ideo paleæ ut flores centrales ipsi deficientes?

1. *C. GENTIANOIDES*, erecta herbacea glaberrima, caulibus teretibus striatis simplicibus, foliis elliptico-oblongis sessilibus basi attenuatis apice obtusissimis integerrimis prope basin triplinerviis, pedunculis oppositè corymboso-paniculatis capitula 3 pedicellata discoidea 4-flora gerentibus, invol. squamis oblongis adpressis, recept. nudo! pappi paleis angustis acuminatis. $\frac{1}{2}$ in Brasiliæ prov. Sancti-Pauli. Herba gentianoidea circ. pedalis. Inv. squamæ ext. parvæ, int. submembranaceæ multò majores. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 406.)

Sect. II. DISCOCALEA. — *Calea Brown l. c. Cass. dict. 55. p. 265.*

Capitulum homogamum discoideum multiflorum.

2. *C. CORNIFOLIA* (Swartz fl. ind. occ. 3. p. 1326), fruticosa, ramis oppositis divergentibus teretibus glabris, foliis petiolatis cordatis acuminatis serratis trinerviis scabris, corymbis terminalibus et axill. 3-8-cephalis, capitulis ovatis discoideis 16-18-floris, invol. squamis membranaceis obtusis, bracteolis lanceolatis acutis, achæniis compressis, pappi paleis circ. 20 striato-penninerviis lanceolatis cor. disci æqualibus. $\frac{1}{2}$ in frutetis montium Jamaicæ australioris. Flores luteo-purpurascetes.
3. *C. JAMAICENSIS* (Linn. sp. 1179), fruticosa, ramis teretibus laxis pubescentibus, foliis petiolatis ovatis acuminatis subserratis trinerviis supra scabris subtus hirtis, capitulis pedicellatis ad apices ramulorum ternis ovatis discoideis, invol. squamis ovatis extus hirtis, paleis lanceolatis, pappi paleis serratis. $\frac{1}{2}$ in Jamaicâ. Swartz fl. ind. occid. 1327. obs. 302. *Santolina Jamaicensis* Linn. amœn. 5. p. 404. — Sloan. jam. 1. p. 257. t. 151. f. 3.
4. *C. BERTERIANA*, suffruticosa? ramis teretibus hirtellis demùm glabrescentibus, foliis petiolatis ovatis acutis grossè dentatis subtrinerviis supra sparsè puberis subtus densè villosis-hirtis, capitulis pedicellatis ad apices ramorum umbellatis discoideis ovatis, invol. squamis ext. acutis subfoliaceis, int. obtusis submembranaceis, achænio puberulo squamellis circ. 20 linearibus acuminatis, receptaculo planiusculo. $\frac{1}{2}$ in Amer. austr. ad Sanctam-Martham legit Bertero. *Marschallia australis* Spreng. in h. Balb. 1822 sed in ejus oper. desid. Valdè affinis *C. prunifoliæ*. (v. s.)
5. *C. ZACATECHICHI* (Schlecht. in linnæâ 1834. p. 589), caule fruticoso tereti subtrichotomo, foliis petiolatis ovatis acutis subtriplinerviis grossè crenatis supra petiolisque hispidulis, ramulisque pubescentibus, capitul. 12-floris in cymis irregularibus dispositis, invol. cylindracei squamis concavis obtusis margine scarioso-undulatis, achænio turbinato, pappo brevi. $\frac{1}{2}$ in collibus apricis Mexici prope Hacienda de La Laguna et Jalapam. Vulg. Mexic. dicitur *Zacatechichi*.
6. *C. PRUNIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 294. t. 406), fruticosa, ramis junioribus subangulatis hirtis-scabris, foliis petiolatis ovatis remotè crenato-serratis utrinquè scabris, capitulis longè pedicellatis ad apices ramorum umbellatis ovatis discoideis, invol. squamis ext. acutis, interioribus bracteolisque obtusis, achænio compresso-tetragono hispido, pappi paleis linearibus

acuminatis achænio paulò longioribus. $\frac{3}{4}$ in ripâ fluminis Magdalenæ prope urbem Honda. Fl. centrales masculi aut steriles. Recept. oblongum.

7. *C. TERNIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 294), fruticosa, ramis subteretibus hirtis-pubescentibus, foliis terno-verticillatis petiolatis ovatis acutis basi angustatis serratis suprâ scabris subtus pubescentibus, capitulis sessilibus trichotomis, capitulis discoideis circ. 7-floris, invol. campanulati squamis 10 ovatis, extimis minutis hirtellis, bracteolis oblongis, achænio hispidulo, pappi paleis lanceolatis margine fimbriatis. $\frac{3}{4}$ in ripâ flum. Magdalenæ prope Honda.
8. *C. SOLIDAGINEA* (H. B. et Kunth nov. gen. am. 4. p. 295. t. 407), suffruticosa, ramulis pubescentibus demum glabris, foliis petiolatis ovato-oblongis acuminatis basi angustatis obsolete serratis suprâ scabris subtus pubescentibus, capitulis breviter pedicellatis ovatis discoideis ad apices ramorum et ramulorum axillarum umbellatis, invol. squamis ovatis subacutis, bracteolis ovato-oblongis acutis, achænio compresso-tetragono apice subhirsuto, pappi paleis linearisubulatis corollas disci æquantibus. $\frac{3}{4}$ $\frac{1}{2}$ in declivitate montis Silla de Caracas.
9. *C. PEDUNCULOSA*, caule basi suffruticoso demisso glabro, ramis dense hirtis basi foliosis apice in pedicellum elongatum 1-cephalum puberulum productis, foliis sessilibus latè ovatis acutis basi obtusis grosse serratis utrinquè hirtellis et impresso-punctatis, invol. campanulati squamis obtusissimis, exter. foliaceis dorso subpuberulis, inter. membranaceis lineato-striatis, achænio angulato villosulo, pappi squamellis linearis-lanceolatis fimbriatis acuminatis achænio longioribus. $\frac{1}{2}$ $\frac{3}{4}$ in Brasiliæ prov. Rio-Grande. Habitus *C. oligocephala*. Recept. paleæ paucæ lineares complicatæ acuminatæ. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1050 miss.)
10. *C. SESSILIFLORA* (Less. in linnæâ 1830. p. 158), fruticosa..., foliis basi obtusissimis et subcordatis, capitulis discoideis cylindraceis circ. 10-floris..., invol..., bracteolis muticis. $\frac{3}{4}$ in Mexico. *Chrysosphaerium gnaphalioides* Willd. herb. Cæt. ign. \dagger .
11. *C. SERRATA* (Less. in linnæâ 1830. p. 158), caule herbaceo..., foliis subtus hirsuto-pubescentibus, capitulis discoideis plurimifloris..., bracteolis muticis. $\frac{1}{2}$ in Brasiliâ. Cæt. ign. \dagger .

Sect. II. CALEACTE. — *Caleacte Brown l. c. Cass. dict. 55. p. 265 non Less.*

Capitulum radiatum. Fl. radii 5-10. — Invol. squamæ siccæ nec exteriores foliaceæ bracteæformes.

12. *C. AXILLARIS*, ramulis teretibus petiolisque subhirsutis, foliis petiolatis ovato-lanceolatis basi breviter attenuatis apice longè acuminatis grosse serratis suprâ vix scabris subtus villosulo-hirsutis, capitulis pedicellatis radiatis ex axillis apicibusque plurimis ortis subumbellatis, invol. squamis siccis, exter. brevioribus acutis, inter. obtusis, bracteolis scariosis oblongis aristatis disco brevioribus, achænio subangulato apice hispidulo, pappi paleis numerosis linearibus acuminatis cor. disci paulò brevioribus. $\frac{3}{4}$? in Amer. calid. (Née!), in Mexico (Hænke). Spec. *C. solidagineæ* affinis. *Mocinna serrata* Lag. nov. gen. p. 31. n. 384 quæ *Galinsogea serrata* Spreng. syst. 3. p. 579, à Næo in Mexico circa Salmanticam urbem lecta, videtur nostra. (v. s.)
13. *C. VERBENÆFOLIA*, ramis teretibus pubescenti-hirtis, foliis petiolatis ovatis acutis serratis, suprâ sparse puberulis, subtus trinerviis reticulatis sparse et præcipuè in nervis puberulo-scabris, capitulis breviter pedicellatis ad apices ramulorum dispositis in thyrsum terminalem digestis ovatis 5-radiatis, invol. squamis siccis obtusis, exter. subacutis, bracteolis linearibus obtusis, pappi paleis 3-4 deciduis, achænio lineari glabro circ. æqualibus. $\frac{3}{4}$? in Amer. calid. legit cl. Née, sed locus propr. in herb. Thibaud desideratur. (v. s.)
14. *C. BRACHIATA*, fruticosa, ramis brachiatis, foliis ovatis obtusè mucronatis, capitulis fasciculatis plerisque terminalibus. $\frac{3}{4}$ in Panamaide. *Mocinna bra-*

- ehiata Lag. nov. gen. p. 31. n. 385. *Galinsogea brachiata* Spreng. syst. 3. p. 579. Cat. ign. †.
15. *C. CACOSMIODES* (Less. in linnæa 1830. p. 157), ramis...., foliis petiolatis ovato-lanceolatis argutè serratis in petiolo non productis subglabris supra lucidis, capitulis radiatis cylindraceis circ. 12-floris, bracteolis latis apice laniatis acuminatis aristatis. — in collibus apricis prope Jalapam.
16. *C. URTICÆFOLIA*, caule tereti piloso, foliis breviter petiolatis ovato-lanceolatis acutis crenatis scabris, racemis brevissimis lateralibus, capitulis radiatis. — ad Vera-Cruz. *Caleacte urticæfolia* Brown in trans. lin. soc. 13. p. 109. *Solidago urticæfolia* Mill. dict. n. 30.
17. *C. NITIDA* (Less. in linnæa 1830. p. 158), ramis florigeris usque ad apicem foliatis, foliis...., capitulis radiatis...., bracteolis oblongo-subellipticis apice integris obtusis flores disci æquantibus squamas invol. intimis superantibus. 3 in Brasiliâ.
18. *C. PINNATIFIDA* (Less. in linnæa 1830. p. 158), glabra, caule suffruticoso scandente, ramis sexangulatis, foliis breviter petiolatis trinerviis ovatis acuminatis grossè inciso-crenatis supra scabriusculis, capitulis pedicellatis paucis terminalibus plurifloris radiatis, invol. squamis obtusissimis. 3 in Brasiliâ (Sellow!), in prov. Rio-Grande (h. imp. Bras. n. 8641). *Caleacte pinnatifida* Brown trans. lin. soc. 13. p. 109. *Mocinna Brasiliensis* Spreng. nen. endt. 1. p. 139. *Actinea commutata* Spreng. syst. 3. p. 574. *Ligulæ* 5. *Achænia scabra*. Pappi paleæ elongatæ acuminatæ. Recept. paleæ muticæ. (v. s.)
19. *C. GLABRA*, tota glabra, ramis scandentibus? hexagonis, foliis petiolatis ovato-lanceolatis basi obtusis apice acuminatis integerrimis, capitulis pedicellatis radiatis, aliis axillaribus solitariis, aliis terminalibus 4 umbellatis, invol. campanulatis squamis siccis obtusis, bracteolis membranaceis lineari-oblongis apice eroso-dentatis, achænio apice glanduloso-scabrido, pappi paleis linearibus acuminatis achænio paulò longioribus. 3 in Brasiliâ ad fretum S. Cathariæ legit cl. d'Urville. *Ligulæ* in specim. meo paucæ, aliæ deciduæ? Pappus albus corollis disci brevior. (v. s. commun. à cl. d'Urville.)
20. *C. ELEGANS*, ramis teretibus glabris, foliis subsessilibus ovato-lanceolatis basi obtusis subcordatis apice longè acuminatis 3-5-nerviis reticulatis subserrulatis glabris secus nervos villosis, capitulis ad apices ramorum pedicellatis ternis 10-radiatis, invol. squamis obtusissimis ovatis glabris, bracteolis membranaceis subacutis floribus brevioribus, achænio glabro angulato, pappi paleis angustissimis caducis achænio paulò brevioribus. 5? in Mexico legit et necum comm. cl. Alaman. (v. s.)
21. *C. CYMOSA* (Less. in linnæa 1830. p. 158), caule herbaceo erecto simplici sparse hispido seu hirsuto basi folioso longè supernè aphylo, foliis inf. oppositis ovalibus basi 5-nerviis, sup. sæpè terno-verticillatis lanceolatis basi cuneatis 3-nerviis apice acuminatis, omnibus dentatis scabridis ad margines et subtus secus nervos ciliato-hispidis, capitulis pedicellatis in corymbum umbelliformem dispositis, invol. squamis acuminatis, recept. paleis muticis, pappi paleis achænio longioribus integris acuminatis. 2? in Brasiliâ (Sellow!), campis editis prov. Sancti-Pauli ad Mugi das Cruzes rara (Lund!), in prov. Rio-Grande (h. imp. Bras. n. 1056). (v. s. commun. à Mus. reg. Berol.)
22. *C. CUNEIFOLIA*, caule simplici minutissimè puberulo-velutino apice aphylo, foliis obovato-cuneatis basi integris 3-5-nerviis apice grossè dentatis subobtusis utrinquè minutissimè puberulis, capitulo solitario terminali, invol. squamis omnibus glabris obtusis multinerviis, pappi squamellis lineari-lanceolatis margine ciliato-fimbriatis achænio duplò longioribus. — in Brasiliâ prov. Sancti-Pauli campis editis legit cl. Lund flor. novemb. Valdè affinis *C. uniflora*, sed notis indic. sat differt. Achænium pubescens. Pappus rufescens corollà longior. (v. s. commun. à cl. Lund.)
23. *C. UNIFLORA* (Less. in linnæa 1830. p. 159), caule simplici basi suffruticulosulo glabriusculo striato folioso apice nudo scabro 1-cephalo, foliis elliptico-

lanceolatis sessilibus grossè et pauciserratis basi trinerviis utrinquè subscabridis, capitulo radiato, invol. squamis exter. bracteisve 2 dorso subscabris foliaceis, mediis ovali-oblongis obtusis multistriatis glabris, recept. paleis acuminatis flores disci superantibus, pappi paleis achæni longit. æqualibus margine eroso-ciliatis. ♂ ♀ in Brasiliâ (Sellow!), in prov. Rio-Grande (h. imp. Bras. n. 1040!). (v. s. comm. à Mus. reg. Berol.)

24. *C. MULTIPLINEARIA* (Less. in linnæa 1830. p. 159), caule simplici superne aphylo, capitulo solitario radiato, invol. squamis exter. acuminatis. ♀ in Brasiliâ. Cæt. ign.

Sect. III. *LEONTOPHTHALMUM* Less. syn. 241. — *Leontophthalmum* Willd. mag. nat. Berol. 1807. p. 140. *H. B. et Kunth nov. gen. am.* 4. p. 296. *Cass. dict.* 25. p. 471. 55. p. 265.

Capitula radiata, ligulis 15-25. Invol. basi bracteis foliaceis puberis cinctum.

25. *C. OLIGOCEPHALA*, caule suffruticuloso pluricipiti, ramis presertim ad basin hirsutis basi foliosis cæt. nudiusculis, foliis ellipticis basi cuneatis sessilibus grossè crenatis utrinquè sparsè hispidis penninerviis, capitulis 1-3 longè pedicellatis radiatis, ligulis circ. 15, invol. squamis ext. seu bracteis subrotundis hirsutis, int. longioribus lævibus obtusis, recept. paleis achænio trigono brevioribus subintegerrimis. ♀ ♂ in Brasiliæ prov. Minarum-Generalium ad Serra do Frio legit cl. Vauthier (pl. exs. n. 318!). Habitus ferè *C. unifloræ* sed fortè ex achænio trigono à genere removenda. (v. s.)

26. *C. LEONTOPHTHALMUM* (Less. syn. 241), fruticosa, ramis teretiusculis villosis, foliis oppositis petiolatis ovato-ellipticis crenato-serratis suprâ hirtocanescentibus subtus canescenti-villosis, capitulo solitario pedicellato subterminali bracteis 4 foliaceis patulis cincto radiato, ligulis circ. 25, invol. squamis obtusissimis glabris, bracteolis lanceolatis apice acutè dentatis, pappi paleis circ. 20 linearibus subulatis cor. paulò brevioribus, achænio glabro. ♂ in temperatis Peruviae. *Leontophthalmum Peruvianum* H. B. et Kunth nov. gen. am. 4. p. 296. t. 409. Capitulum $1\frac{1}{2}$ -2 poll. diam. Buphthalmi facie.

27. *C. CORIACEA*, fruticosa, ramis teretibus, junioribus breviter puberulis, foliis brevissimè petiolatis ovatis seu ovalibus basi obtusis apice subacutis grossè dentato-serratis penninerviis reticulatis coriaceis utrinquè glabris subtus punctatis, corymbo oligocephalo, invol. campanulatis squamis ovatis siccis obtusis, recept. semigloboso, achæniis subtetragonis secus angulos pilosiusculis, pappi paleis acuminatis linearibus achænio ferè longioribus. ♂ in Peruvia legit cl. Hænke ex h. acad. reg. Monac. Flores non vidi. Species tamen distinctissima ad hanc sect. ex habitu relata. (v. s. in h. suprâ cit.)

+ *Species vix nota.*

28. *C.?* *CORDATA* (Lour. fl. coch. 2. p. 595), suffruticosa, foliis alternis cordatis subcrenatis subtus tomentosiss, capitulis pedicellatis ternis quaternisve discoideis. ♂ in Cochinchinâ. Ex foliis alternis, floribus purpureis, et pappo piloso evidenter à genere removenda, sed vix nota! †.

++ *Species exclusæ.*

- C. ACULEATA* Labill. = *Cassinia aculeata*.
C. AMELLUS Linn. — *Salmea scandens*.
C. ASPERA Jacq. = *Melanthera deltoidea*.
C. LEPTOPHYLLA Forst. = *Cassinia leptophylla*.
C. LOBATA Gertl. = *Neurolæna lobata*.
C. OPPOSITIFOLIA Linn. = *Isocarpa oppositifolia*.
C. PEDUNCULARIS H. B. K. = *Calydermos peduncularis*.

C. PINIFOLIA Forst. = *Oreothamnus pinifolius*.

C. SCOPARIA Linn. = *Baccharis scoparia*.

C. SPECTABILIS Labill. = *Apalochlamys spectabilis*.

CDXXII. ALLOCARPUS *H. B. et Kunth nov. gen. am. 4. p. 291. t. 405. Cas. dict. 55. p. 265. Less. syn. 242. — Alloispermum Willd. mag. naturf. Berl. 1807. p. 139. ex Cass.*

Capitulum multiflorum radiatum, fl. radii circ. 5 ligulatis foemineis, disci hermaphr. tubulosis 5-dentatis. Invol. hemisphærici squamæ laxè imbricatæ, circ. 10 scarioso-membranaceæ. Recept. subplanum paleis lanceolatis persistentibus scariosis onustum. Styl. fl. hermaphrod. rami exappendiculati. Achænia radii cuneata compressa calva, disci cuneato-cylindracea pappo 1-seriali multipaleaceo coronata. Paleæ pappi acuminatæ, ex icon. pectinatæ ex descr. integræ. — Herbae Americanæ ramosæ. Folia opposita breviter petiolata, tripli- seu 5-7-nervia, ovalia utrinquè acuminata integra villosa. Capitula pedicellata flava aut fortè albida.

1. A. CARACASANUS (*H. B. et Kunth l. c. t. 405*), caule erecto hirsuto, ramis apice corymbiferis, foliis ovali-lanceolatis acuminatis hinc indè dentatis triplinerviis villosis. 2½? in temperatis propè urbem Caracas. *Alloispermum divaricatum* Willd. l. c. *Galinsogea allocarpa* Spreng. syst. 3. p. 579. Pappi paleæ margine fimbriatæ.
2. A. INTEGRIFOLIUS, caule herbaeo tereti glabro virgato, foliis breviter petiolatis ovali-lanceolatis acuminatissimis integerrimis 5-7-plinerviis utrinquè strigillis brevibus sparsis scabridis, corymbo composito polycephalo. — inter plantas Mexicanas à cl. Karvinski lectas video in h. acad. reg. Monac. ad locum non indicatur. Herba fortè subscandens? Folia 5 pollic. longa, pollicem lata, membranacea subnitida. Capitula fortè albida, dimidiò minorà quàm prioris, circ. 15-flora. Pappi paleæ vix margine scabræ. (v. s. in h. cit.)
3. A. SARAZIOIDES (*Less. syn. 242 et in linnæa 1834. p. 590*), herbacea procumbens monocephala, foliis petiolatis ovatis acutis crenatis triplinerviis hirsutis. bracteolis scariosis acuminatissimis bidentatis. — in Mexico prope San-Miguel del Soldado et La Joya.

CDXXIII. VARGASIA.

Capitulum heterogamum, fl. radii ligulatis paucis 1-serialibus inter invol. squamas et rec. paleas ext. et majores occultatis, disci tubulosi 5-dentatis. Recept. convexum paleaceum, paleis membranaceis angustissimis, ext. majoribus. Invol. hemisphæricum, squamis subbiserialibus ovalibus mucronulatis. Ligulæ breves obovatæ apice tridentatæ. Stylus brevis inclusus, ramis cono brevi terminatis? Achænia radii obcompressa inter squamam int. et paleam ext. inclusa, disci subtri-gona, omnia basi attenuata subpuberula. Pappi paleæ variæ, radii setiformes parvæ, disci 5 ovatæ erectæ serrato-ciliatæ acutæ corollâ triplò longiores. — Herba Austr.-Americana æquinoctialis annua erecta gracilis, parcè dichotomè ramosa, sparsè hispida. Folia opposita brevissimè petiolata ovata acuminata dentato-serrata. Pedunculi è dichotomis orti, 1-cephali ebracteati. Capitula subglobosa, invol. glabro. Flores pallidè purpurascentes, à *Galinsogâ* differt ferè ut *Allocarpus* à *Caleâ*. — Genus affine *Galinsogæ* dixi in honorem cl. doct. Vargas Caracasani rerum naturalium sagacis observatoris.

1. V. CARACASANA. (1) ad Caracas legit cl. Vargas. (v. s. comm. à cl. inventore.)

CDXXIV. GALINSOGA R. et Pav. *fl. per. prod.* (1794) p. 110. t. 24. *Cass. dict.* 18. p. 96. — Galinsogæ Zucc. in *florâ* 1821. p. 612. — Wiborgia Roth *cat.* 2. (1800) p. 112. *Less. syn.* 245 non Spreng. Thunb. Mænnch (1). — Galinsogæ spec. Cav. Willd.

Capitulum pluriflorum radiatum, ligulis 4-5 fœmineis suborbiculatis, fl. disci hermaphr. tubulosis 5-dent. Invol. 1-seriale, squamis 5 ovatis margine hyalinis. Recept. conicum paleis lineari-lanceolatis acutis integris trifidisve onustum. Styli rami acuti vix appendiculati. Achænium angulatum adpressè hispidulum. Pappus 1-serialis, paleis oblongis brevibus plumoso-fimbriatis. — Herbæ Americanæ annuæ trichotomoramosissimæ. Folia opposita ovata triplinervia serrata. Pedunculi axillares et terminales 1-3-cephali. Capitula parva, disco luteo, radio albo.

1. G. PARVIFLORA (Cav. *ic.* 3. p. 41. t. 281), glabriuscula, receptaculo conico, pappi paleis 8-16. ① in Peruvîæ uliginosis et ruderatis (R. et Pav. Cav.), in Chili ad Coquimbo (Hook. et Arn.), in Novâ-Granatâ circa Ybague (H. B.), Mexico (Keerl!), nunc circa Erlang, etc., seminibus ex hort. bot. egressis quasi spontanea (Zucc.). Gal. quinqueradiata R. et Pav. *syst.* 198. Wiborgia Acnella Roth *cat.* 2. p. 112. — Feuill. *per.* 2. p. 744. t. 32. Vulgò dicitur Peruvianis Paica-Jullo (Feuill.) seu Pacoyuyu-Fino (R. et Pav.). Vulneraria et antiscorbutica habetur. (v. v. c.)

2. ligulis 3-4. — Galinsoga quadriradiata R. et Pav. *syst.* 198. — in Peruvîâ ad Limam et Chancay. Vulgò Pacoyuyu-Cimarron ex R. et P. Vidi in hortis specim. 4-ligulata certè à var. æ non diversa. (v. v. c.)

3. hispida, caule foliisque sparsè hispidis, ligulis 5. ① in Mexico circa urbem (Berl. pl. exs. n. 615), in Chilensibus montibus (h. Hanke!). Caulis crassior. Folia majora. Pedunculi abbreviati. (v. s.)

2. G. ? ORLONGIFOLIA (Hook. *misc. bot.* 2. p. 226 sub Wiborgia), tota pilis adpressis albis aspera, foliis oblongo-lanceolatis triplinerviis grossè serratis basi attenuatis subsessilibus, invol. polyphylo, rec. paleis linearibus hispidis, pappo piloso brevissimo. — ad Lurin prope Limam legit cl. Cruikshanks. Pedicelli binociales. Ligulæ bidentatæ. Videtur præsertim ex pappo è genere exclud. et novum verisim. constituendam.

+ Species vix notæ.

3. G. ? ANGUSTIFOLIA (Spreng. *neu. entd.* 2. p. 138. *syst.* 3. p. 580), caule basi suffruticoso hirsuto, ramis glabriusculis angulatis, foliis oppositis lineari-lanceolatis dentatis margine revolutis, infer. hirsutis, super. glabris integris, pedicellis terminalibus 1-cephalis, invol. squamis subrotundis coloratis, ligulis crenatis, pappo squamellato. 3 in Brasiliâ. Flores lutei.

4. G. ? MEGAPOTAMICA (Spreng. *syst.* 3. p. 580), herbacea glabra, foliis linearibus tripartitis integerrimis, ramis ternis 1-floris rigidis. — in Brasiliâ ad Rio-Grande. Cæt. ign.

5. G. ? UNIFLORA (Spreng. *syst.* 3. p. 580), caule herbaceo simplicissimo 1-cephalo supernè hirsutulo, foliis sparsis linearibus mucronatis margine scabris. — ad Monte-Video. Cæt. ign.

6. G. ? RESINOSA (Hook. et Arn. *bot. Beech.* 1. p. 32), caule fruticoso angulato ramoso, ramulis foliisque angustissimè linearibus resinoso-glandulosis, capitulis paniculato-corymbosis, invol. imbricato inæquali viscoso, recept. nudo. 5 in Chili ad Coquimbo et Valparaiso. Species ex recept. nudo potius Bahiæ quàm Galinsogæ referenda videtur. Valdè dubia nec satis descr. †.

(1) Wiborgia Thunb. seu Viborgia Spreng. et DC. est genus in prod. jam inter Leguminosas admissum (2. p. 135). Wiborgia Mænnch (1794) est sect. Cytisi mihi Tubocytisus. Wiborquia R. (1797) est Varennea DC. *prod.* 2. p. 522.

2. *B. SCABER* (Hook. l. c.), foliis angusto-linearibus scabris, involucria subbescentibus, glandulis pedicellatis nullis, acheniis omnibus villosis, paleis pappi breviusculis superne præcipue fimbriato-ciliatis. ① hab. cum. priore.

CDXXIX. *MARSHALLIA* Schreb. gen. (1791), n. 1762. Cass. dict. 55. p. 261. Less. syn. 241. — *Persoonia* Michx. fl. bor. am. 2. p. 104 non Sm. — *Phytomopsis* Juss. ined. Michx. l. c. icon. — *Trattenickia* Pers. ench. n. 1825. n. Willd. — *Athanasia* Walt. fl. car. 201. non Linn.

Capitulum homogamum. Invol. 1-2-serialis squamæ lineari-lanceolatæ disco æquales. Recept. convexum paleis linearibus acutis onustum. Cor. tubulosæ 5-dentatæ extus hirsutæ. Styli rami exappendiculati. Achenium turbinato-subquingulum pubescens seu villosum. Pappi paleæ 5 enerviæ ellipticæ acuminatæ integræ. — *Herbæ* Boreali-Americanae perennes. Folia alterna integra. Capitula terminalia solitaria purpurascentia non malè *Scabiosarum* capit. referentia. — An satis ab hoc genere differt *Therolepta* Raf. nov. gen. 1825. p. 3?

1. *M. CÆSPITOSA* (Nutt. ! in litt. 1825), glabra cæspitosa, caule simplici aphylo 1-cephalo, foliis elongato- et lato-linearibus subobtusis integerrimis, invol. squamis oblongo-linearibus obtusis, paleis recept. linearibus. ② in Amer. bor. ad Red-River legit cl. auctor. Habitus ferè *Armeriæ*. Caulis 8-10-pollic. ter striatus. Cor. ex sicco albæ, fortè subroseæ. Achenium villosum. (v. s.)
 2. *M. LANCEOLATA* (Pursh fl. bor. Am. 2. p. 519), caule simplici inferne foliis superne nudato, foliis glabris, radicalibus petiolatis obovatis triplinerviis, caulinis sessilibus oblongo-lanceolatis, invol. squamis ovali-oblongis obtusis, recept. paleis lineari-spathulatis. ② in Carolinae montanis. Ell. sketch 2. p. 315. *Persoonia* lanceolata Michx. fl. bor. am. 2. p. 105. *Trattenickia* lanceolata Pers. ench. 2. p. 403. *Athanasia* obovatæ Walt. car. p. 201. (v. s.)
 3. *M. LATIFOLIA* (Pursh l. c.), caule simplici, foliis glabris lanceolato-ovalibus acuminatis trinerviis sessilibus, infimis vaginantibus, invol. squamis acutis, recept. paleis angusto-linearibus. ② in Carolinae montanis. *Persoonia* latifolia Michx. fl. bor. amer. 2. p. 105. t. 43. *Athanasia* trinervia Walt. car. 201. *Trattenickia* latifolia Pers. l. c. *Marshallia* Schreberi Tratt. arch. gew. 1. p. 104. n. 123. cum icon. Michx. mut.
 4. *M. ANGUSTIFOLIA* (Pursh l. c. p. 520), caule ramoso vix ad apices pubescente, foliis infer. angusto-lanceolatis, super. linearibus obscure 3-nerviis glabris, invol. squamis acuminatis rigidulis, recept. paleis linearibus. ② in Tennessee (Michx.), in pinetis humidis Carolinae et Georgiae (Ell. Fras.!). *Persoonia* angustifolia Michx. l. c. *Trattenickia* angustifolia Pers. ench. 2. p. 403. *Athanasia* graminifolia Walt. car. p. 200. (v. s.)
- ℓ. *cyananthera* (Ell. sketch 2. p. 317), caule simplici apice magis pubescente, foliis evidentiis 3-nerviis, invol. pubescente. ② in Georgia prope St.-Mary. (v. s.?)

† *Species exclusæ.*

M. ALIENA Spreng. = *Heterothalamus brunoides*.

M. AUSTRALIS Spreng. = *Calea Berteriana*.

M. BULBOSA Raf. fl. lud. 61 et ideò Spreng. syst. = *Liatridia* spec. ?

CDXXX. *DUBAUTIA* Gaudich. in voy. Freyc. bot. 468. t. 84. Hook. et Arn. bot. Beech. 88. Less. syn. p. 247.

Capitulum multi- (8-10) florum homogamum. Recept. Invol. campanulatum, squamis circ. 8 liberis 1-serialibus sibi mutuo incumben- tibus. Cor. tubulosæ regulares 5-dentatæ. Antheræ ecaudatæ breviter appendiculatæ. Filam. apice vix incrassata. Stigmata dilatata acuta

ciliata. *Achænium hirsutum oblongum seu obpyramidatum*, areolâ terminali. Pappus 1-serialis paleaceus, paleis angustis ciliato-plumosis. — Suffrutices Sandwichenses, ramis strigilloso-pilosis teretibus apice foliosis, infernè cicatricato-annulatis. Folia opposita sessilia rigida subparallelè venosa dentata. Capitula in racemos paniculatos adgregata. — Genus affine *Tridaci* ex Less. et Hook. aut *Railliardæ*.

1. *D. PLANTAGINEA* (Gandich. bot. Freyc. voy. 468. t. 84), foliis amplexicaulis lineari-lanceolatis acuminatis crenatis glabris, paniculâ foliatâ, capitulis fasciculatis. ♂ in insul. Sandwich. montanis ad 550-600 hexap. altit., in ins. O-Wahu ex Less. in linnæâ 1831. p. 164. Facies *Echli fruticosi*! (v. s. comm. à cl. de Chamisso.)
2. *D. LAXA* (Hook. et Arn. bot. Beech. 87.) foliis oblongo-lanceolatis basi attenuatis argutè serratis subtus strigoso-hispidis, paniculâ corymbosâ laxâ nudiusculâ. ♂ in insulis Sandwich.

Subdiv. 2. SPHENOGYNEÆ.

Ligulæ neutræ. Pappi palææ obovatæ obtusissimæ more corollarum contortarum convolutæ.

CDXXXI. SPHENOGYNE *R. Brown h. Kew. ed. 2. v. 5. p. 142. Cass. dict. 50. p. 204. Less. syn. 242. — Oligæria Cass. dict. 2. p. 75. 29. p. 187. — Thelythamnus Spreng. f. suppl. 25. — Spermothylla Neck. elem. 1. p. 2. excl. sp. — Arctotidis sp. Linn. Thunb. — Ursiniæ sp. Lam. ill. t. 716. f. 1-2. Spreng. syst. 3. p. 409.*

Capitulum multiflorum radiatum, fl. radii ligulatis 1-serialibus neutris, disci tubulosis 5-dentatis hermaphroditis. Recept. paleaceum, paleis scariosis flores amplectentibus sæpius apicè truncatis. Invol. campanulatum multiseriale, squamis imbricatis, inter. majoribus apice amplè scariosis. Stylus in ramos apice truncatos ferè cuneatos desinens. Anthæræ appendice terminali subcordiformi superatæ. *Achænium* pilis mollibus è basi ortis cinctum, cylindraceum vix basi angustatum. Pappus 1-serialis paleaceus, laminis obovatis cuneatisve obtusis ante anthesin contortarum more spiraliter dispositis, junioribus pellucidis, adultis lacteis opacis. — Herbæ aut suffrutices Capenses *Anthemidis* facie. Folia alterna sæpius pinnatiloba, rariùs trifida, imò integerrima. Rami apice pedunculiformes nudi 1-cephali. Cor. flavæ aut apice fuscæ, ligulis interdum subtus rubris.

Sect. 1. THELYTHAMNOS.

Involucri squamæ exteriores absque margine scarioso manifesto.

§. 1. *Species annuæ.* — *Anthemoides* Less. syn. 242.

1. *S. ANTHEMOIDES* (*R. Brown h. Kew. ed. 2. v. 5. p. 142*), herbacea glabra, foliis irregulariter bipinnatipartitis laciniis lineari-filiformibus, pedicellis elongatis nudis, recept. paleis conformibus truncatis, invol. squamis externis acutiusculis, ligulis discoloribus, paleis truncatis. (†) ad Cap. Bonæ-Spei. Less. syn. 242. *Arctotis anthemoides* Linn. sp. 1307? Berg. cap. 324. Sims bot. mag. t. 544 bona. *Arctotis pinnata* et *A. cernua* Thunb. fl. cap. 711 et 712. *Arctotis paradoxa* Ait. h. Kew. ed. 1. v. 3. p. 373. *Ursinia anthemoides* Poir. dict. 8. p. 257. (v. s. c.)
2. *S. VERSICOLOR*, herbacea multicaulis diffusa glabrinscula, junior subpubescens, foliis bipinnatipartitis lobis linearibus, omnibus integris linearibus vix acutis, infer. brevioribus, pedicellis elongatis nudis, invol. squamis ext. ovatis albis

villosis, ligulis paulò magis pallidis, pappo dimidiò breviorè. \mathcal{L} \mathfrak{z} ad Lederbergen alt. 3-400 ped. legit cl. Drege pl. exs. n. 6008! An fortè specis distinguenda? (v. s.)

β . *Foliis glabriusculis, lobis brevibus dentatis, dentibus apice setigeris.*

17. *S. DENTATA* (R. Brown h. Kew. ed. 2. v. 5. p. 143), fruticosa glabra, foliis pinnatipartitis, lobis brevibus integris aut trifidis, dentibus setigeris, pedunculis valdè elongatis, invol. squamis exter. ovatis acutis, radii ligulis discoloribus. \mathfrak{z} ad Cap. Bonæ-Spei circa urbem. Burch! cat. Afr. austr. n. 438. Drege! n. 158 Eckl. n. 1920. 460. 858. *Arctotis dentata* Linn. sp. 1307. N. L. Burm. prod. 44. non Thunb. nec Willd. *Arctotis piligera* Berg. cap. 325. *Arctotis pectinata* Thunb. fl. cap. 710. *Ursinia dentata* et pilifera Poir. dict. 8. p. 256. Spreng. syst. 1. c. — Burm. Afr. t. 64. Pluk. alm. t. 276. f. 2. et fortè 382. f. 5. (v. s.)

β . *marginata*, invol. exter. nigro marginatis. \mathfrak{z} ad Cap. Bonæ-Spei Burch! cat. Afr. austr. n. 6698. Drege! pl. exs. Eckl. 1257? 1541. 1602. (v. s.)

18. *S. SETIGERA*, suffruticosa, caule inter folia puberulo, foliis pinnatisectis glabris, segmentis utrinquè 4-5-apice 2-4-fidis, dentibus longè setigeris, pedunculo elongato folia sparsa linearia integra apice setigera gerente, invol. squamis exter. lanceolatis acutis nigro-marginatis, ligulis discoloribus invol. fere triplò longioribus, recept. paleis truncatis. \mathfrak{z} in Africâ austr. ad Cap. Agullas legit cl. Drege pl. exs. n. 6009! Valdè accedit ad *S. dentatam*, sed capitulum duplò majus et setæ fol. frequentiores et longiores. (v. s.)

γ . *Foliis glabris seu glabriusculis, lobis apice non setigeris.*

19. *S. NUDICAULIS* (Less. 1. c.), glaberrima, basi suffruticosa, foliis subradicalibus pinnatisectis, lobis 6-7-jugis, inferioribus brevibus indivisis, superioribus longioribus rarissimè partitis, lobis lobulisque linearibus vix acutis, scapo foliis multò longiore nudo stricto, invol. squamis extimis ovatis acutis, ligulis concoloribus invol. duplò longioribus. \mathcal{L} ad Cap. Bonæ-Spei (Burch. l. n. 632), ad montem Tabularem alt. 4-500 ped. (Drege! n. 6007. Eckl. n. 1918. 659. 1706). *Arctotis nudicaulis* Thunb. fl. cap. 711. Variat parte caulis fruticosâ nunc demissâ aut rhizoidè adscendente, nunc erectâ et verè caulescente. (v. s.)

20. *S. SCAPIFORMIS*, glaberrima, rhizomate suffruticoso, foliis subradicalibus pinnatipartitis, lobis 10-12-jugis, inf. parvis indivisis acuminatis, sup. longioribus in lobos lineares 2-3 partitis, pedicello longissimo stricto nudo 1-cephalo, invol. squamis exter. ovato-lanceolatis acutis, ligulis concoloribus invol. circ. longitudine. \mathcal{L} \mathfrak{z} ad Cap. Bonæ-Spei loco dicto Langekloof. alt. 2-300 ped. legit cl. Drege pl. exs. n. 6004 et ad Tulbagh cl. Ecklon n. 1362. et ad Worcester n. 1446! An mera *S. nudicaulis* var.? Differt præter notas indic. capitulo dimidiò minore. Stirps Eckl. n. 1743, 1047, 213 ad Worcester lecta utrasque ferè conjungit. (v. s.)

21. *S. CHAMOMILLÆFOLIA*, glaberrima, caule imò basi suffruticuloso radicante subadscendente aut erecto, foliis pinnatipartitis, lobis linearibus acutis 8-9-jugis, lob. inf. indivisis, sup. 2-3-lobatis longioribus, pedicellis valdè elongatis nudis 1-cephalis, invol. squamis ext. minimis acutis, mediis ovali-oblongis obtusis, summis scarioso-appendiculatis, ligulis discoloribus, paleis truncatis subdenticulatis. \mathcal{L} \mathfrak{z} in Cap. Bonæ-Spei ad Draakensteensberg alt. 3-400 ped. legit cl. Drege pl. exs. n. 6006! ad Worcester (Ecklon! n. 913). Accedit ad *S. scapiformem*. (v. s.)

22. *S. ANETHIFOLIA* (Less. syn. 243), fruticosa erecta glabra, foliis subcoriaceis punctatis pinnatisectis, lobis lineari-filiformibus acutis, inf. integris, super. apice 2-3-fidis, pedicellis elongatis 1-cephalis, invol. squamis ext. ovatis acutis, ligulis discoloribus invol. paulò longioribus. \mathfrak{z} ad Cap. Bonæ-Spei (Less.). Lobi fol. utrinquè 7-8. Ligulæ 3 lin. longæ. An certè nostra eadem ac Lessingiana ex descriptionis brevitate dubia? (v. s. comm. à cl. Boissier ex coll. Verreaux.)

23. *S. ANETHOIDES*, fruticosa erecta glabra aut subpuberula, foliis subcoriaceis pinnatisectis, lobis lineari-filiformibus acutis subpunctulatis, infer. integris, super. apice 2-3-fidis, pedicellis elongatis 1-cephalis, invol. squamis exter. lanceolatis acutis, ligulis concoloribus invol. duplò longioribus, paleis truncatis. ζ ad Cap. Bonæ-Spei prope Breederivier et Draakensteensberg legit cl. Drege alt. 1-200 ped. pl. exs. n. 6015, 6019, 6020! et ad Worcester cl. Ecklon n. 1058 592. 212. 1860! Proximè accedit ad *S. anethifoliam*. (v. s.)
- β . *brachyglotta*, ligulis vix invol. longioribus. ζ ad Piquetberg alt. 2-300 ped. legit cl. Drege pl. exs. n. 6012! (v. s.)
- γ . *ramosissima*, caule laxè ramosissimo, foliis brevioribus brevius lobatis. ζ ad Giftberg alt. 2-300 ped. legit cl. Drege pl. exs. n. 6011! (v. s.)
24. *S. GRACILIS*, fruticplosa erecta virgata glabriuscula, foliis pinnatipartitis brevibus, lobis inf. indivisis, sup. subternis 3-dentatis vix 3-lobatis, omnibus lineari-filiformibus mucronulatis brevibus, ramis apice longe denudatis, invol. squamis exter. paucis minimis acutis, cæteris obtusis, ligulis concoloribus vix invol. longioribus, paleis truncatis. ζ in Africâ australi ad Zeederbergen alt. 4-500 ped. legit cl. Drege pl. exs. n. 6010! Affinis *S. anthemoidi*, sed capit. dimidiò minorâ et ligulâ minimâ. Villi baseos fructûs longissimi. (v. s.)
25. *S. CHAMÆMELOIDES*, suffruticosa glabra, foliis pinnatisectis, lacinii linearibus acutis, superiorum integris, inferiorum sæpè trifidis, invol. squamis exter. triangularibus mucronulatis aristatis, radii ligulis discoloribus. ζ ad sinum Algoa Africæ australis (Forbes!). Rami floriferi parcè foliosi 1-2 poll. longi. Invol. squamæ plurimæ scarioso-membranacæ. Ligulæ 5-7-nerviæ. (v. s.)
26. *S.?* *RIGIDULA*, suffruticosa erecta aut adscendens glabra, foliis pinnatipartitis subapinescentibus, inf. brevibus integris, sup. longioribus trilobatis, pedicellis elongatis 1-cephalis, invol. squamis ext. ovali-oblongis margine scariosis subobtusis, ligulis discoloribus, paleis obtusis. ζ in Africâ austr. ad Piquetberg alt. 2-300 ped. legit cl. Drege pl. exs. n. 6013! Aff. *S. chamæmelifoliæ* et cum eâ fortè ad *Urnium* repellenda? Fruct. ign. Pedicelli 3-6 poll. longi. Folia 10-12 lin. longâ. (v. s.)
27. *S. BREVIFOLIA*, fruticosa erecta ramosissima puberula aut glabriuscula, foliis rigidulis glaucescentibus pinnatipartitis paucijugis, lobis mucrone subuncinato-acutis, infer. 2-3-jugis minimis interdum nullis, super. 3-5 approximatis longioribus, pedicellis elongatis folia linearia integerrima erecta basi gerentibus, invol. squamis ext. ovali-oblongis apice subscariosis obtusis rufò marginatis, ligulis discoloribus, paleis obtusis. ζ in Africâ Capensis distr. Uitenhage et Caledon legit cl. Ecklon n. 275. 258. 963. 1431 et 1257. Fruct. ign. Folia 5 lin. longâ. (v. s.)
28. *S. PALEACEA* (Less. syn. 244), fruticosa glabriuscula, foliis pinnatisectis, segmentis margine revolutis, invol. squamis ext. numerosis angustè linearibus acuminatis conduplicatis subpatulis, radii ligulis discoloribus. ζ ad Cap. Bonæ-Spei. Arctotis paleacea Thunb. fl. cap. 712.
29. *S. GRANDIFLORA*, fruticosa glabra, foliis punctatis pinnatisectis, lacinii linearibus acutis, inf. brevioribus, super. longioribus fastigiatis, pedunculo foliis duplò longiore, invol. squamis linearibus vix acutis, radii ligulis concoloribus squamis inv. inter. duplò longioribus. ζ ad Cap. Bonæ-Spei in Hottentots-Holland (ex h. Deless.) ad Draakensteensberg alt. 3-400 ped. legit cl. Drege! n. 6018. Eckl.! n. 257. Valdè affinis *S. punctatæ* et forsân cum eâ à plurimis conjuncta, sed ligulâ 8-9 lin. longæ, 2 lin. forè latæ, 5-7-nerviæ, evidentiùs quàm prioris glandulosæ. (v. s.)
30. *S. DISCOLOR* (Less. syn. 243), fruticosa tomentosa (Th.), foliis coriaceis pinnatisectis segmentis integerrimis, invol. squamis extimis minutis ovatis acutis, radii ligulis discoloribus. ζ ad Cap. Bonæ-Spei. Arctotis dentata Thunb. fl. cap. 712. excl. syn. ex Less. Capitula solitaria (Less.). Fol. segmenta subulata brevina. (Thunb.)
31. *S. CRITHMIFOLIA* (R. Brown h. Kew. ed. 2. v. 5. p. 142), fruticosa glabra,

foliis pinuatisectis, lobis lineari-filiformibus acutis, pedicellis elongatis folia linearia indivisa hinc inde gerentibus, invol. squamis extimis linearibus acuminatis, mediis lanceolatis, intimis in appendicem scariosam amplam ovalem productis, ligulis concoloribus, recept. paleis truncatis. ζ ad Cap. Bonæ-Spei (Burm. Th.), in monte Tabulari (Dreg. l. n. 76). *Arctotis crithmoides* Berg. cap. 326 non Jacq. *Arctotis paleacea* Linn. amœn. 6. p. 107 ex ic. cit. — Burm. afr. t. 65. f. 1. Foliorum lobi intervallo longiores, 3 ultimi approximati. Folia super. apice trifida, summa indivisa. Achænium albidum striato-sulcatum. Pappi laminæ obovatæ lacteæ. (v. s.)

β . *trifurcata*, foliis multò brevioribus, 3 summis approximatis. ζ *Arctotis trifurcata* Burm. l. prod. fl. cap. p. 28. (v. s.)

32. S. PUNCTATA (Less. l. c.), fruticosa glabra, foliis punctatis pinnatisectis, lacinis linearibus acutis, inf. brevioribus, summis longioribus fastigiatis, pedunculo folio triplò quadruplòve longiore, invol. squamis extimis acutis, radii ligulis concoloribus involucri squamis inter. paulò longioribus. ζ ad Cap. Bonæ-Spei. *Arctotis punctata* Thunb. fl. cap. 711. (v. s.)

33. S. PAUCILOBA, fruticosa ramosa erecta glabra, foliis coriaceo-carnosulis pinnatilobatis, lobis sæpiùs 5 obtusiusculis integris, terminalibus 3 approximatis, summis linearibus indivisis, pedicellis modicè elongatis ramisque junioribus subaraneosis, invol. squamis ext. oblongis obtusiusculis nigro-marginatis, ligulis concoloribus invol. paulò longioribus, paleis truncatis. ζ in Cap. Bonæ-Spei ad Zwellendam am Berge alt. 4-500 ped. legit cl. Drege pl. exs. n. 1758. (v. s.)

34. S. SUBFLOSCULOSA (Erm. Mey. l. in Drege herb. n. 1757), fruticulosa glabra, foliis crassiusculis punctatis, infer. pinnatipartitis bijugis cum impari, mediis apice trifidis, summis linearibus integerrimis, ramis apice nudis pedicelliformibus 1-cephalis, invol. squamis exter. linearibus acuminatis, inter. ad apicem parçè scariosis oblongis, ligulis invol. ferè brevioribus, recept. paleis apice obtusis subdentatis. ζ ad Cap. Bonæ-Spei ad Zeederbergen alt. 3-400 ped. legit cl. Drege! Achænia teretia striata albida imà basi pilifera. Pappi laminæ albæ obovatæ subenerviæ. (v. s.)

35. S. PINNATA (Less. syn. 243. E. Mey. l. in Drege pl. exs. n. 1728), fruticosa glabra, foliis sessilibus subamplexicaulibus pinnatipartitis, lobis lineari-filiformibus mucronatis, infimis in parte amplectente sitis ferè stipulæformibus, ramis floridis folia parva indivisa gerentibus ramoso-paniculatis, capitulis ovatis, invol. squamis ext. oblongis vix acutis, ligulis parvis discoloribus linearibus, recept. paleis obtusis. ζ ad Cap. Bonæ-Spei (Thunb. Burch. l. n. 7597 et 8626), locis dictis Draakensteensberg et Giftberg (Dreg. l.), in distr. Caledon Eckl. l. n. 731. 193 et 339. *Relhania pinnata* et *Arctotis pinnata* (ex parte) Thunb. fl. cap. 641 et 711. *Ursinia filiformis* Griesel. in linnæa 1830. p. 414. *Thelythamos filiformis* Spreng. f. suppl. 25. (v. s.)

* * *Foliis tri- aut quinquelobatis.*

36. S. QUINQUEPARTITA, suffruticosa glabra ramosa, ramis usque ad apicem confertè foliosis, foliis ferè à basi 5-partitis, lobis lineari-filiformibus integerrimis acutis, parte indivisâ brevi latiusculâ transversè demùm sectâ et infernè persistente, capitulis breviter pedicellatis solitariis, invol. squamis extimis lineari-lanceolatis, ligulis discoloribus. ζ ad Cap. Bonæ-Spei in Stellenbosch leg. cl. Ecklon et Zeyher. Ligulæ in iisdem capitulis, aliæ neutra, aliæ foemineæ fertiles. Disci flor. sæpiùs abortivi. (v. s. comm. à cl. Ecklon.)

37. S. TRIPARTITA, fruticulosa glabra, ramis ad apicem usquè confertè foliosis, foliis ferè à basi tripartitis, lobis lineari-filiformibus acutis integerrimis, parte indivisâ latiusculâ demùm transversè sectâ et infernè persistente, capitulis breviter pedicellatis, invol. squamis extimis lanceolatis, ligulis concoloribus. ζ ad Cap. Bonæ-Spei (Lamb. l. herb.) in distr. Zwellendam (Eckl. l. et Zeyh.). loco dicto Hottentots-Holland (Dreg. l.). Folia summa subpubentia. Ligulæ in

specim. simillimis nunc steriles, nunc plures fertiles. Disci flor. sæpè steriles. (v. s.)

38. *S. HETERODONTA*, suffruticosa glabra, foliis trifidis, nunc lobis 2 lateralibus lineari-filiformibus medio dentiformi brevissimo, nunc in parte superà 2 lateralibus dentiformibus dente medio elongato, nunc rarius subæqualibus, capitulo longè pedunculato, invol. squamis extimis acuminatis adpressis, radii ligulis discoloribus. ♂ ad Cap. Bonæ-Spei in coloniæ distr. orient. legit cl. Burchell (cat. geogr. n. 6266), in Uitenhage (Eckl. ! n. 1198. 105). (v. s.)

39. *S. TRILOBA*, fruticulosa, foliis glabriusculis punctulatis lineari-cuneatis apice breviter et obtusè trifidis subtus bisulcis, capitulis supra folia supremata pedicello foliis duplò longiore donatis, invol. squamis extimis lanceolato-linearibus acuminatis, ligulis concoloribus, recept. paleis obtusis apice dentatis cor. tubi duplò ferè brevioribus. ♂ ad Cap. Bonæ-Spei in distr. Zwelendama legit cl. Burchell pl. exs. n. 7362 et cl. Ecklon n. 593. Diff. à *S. trifida* (juxta Less. descr.), radio concolore nec discolore, à descr. Thunb. foliis nec indivisis nec 5-fidis. (v. s.)

40. *S. TRIFIDA* (Less. syn. 244), caule fruticuloso, foliis apice trilobis, invol. squamis extimis linearibus acuminatis paucis planis adpressis, radio discolore. ♂ ad Cap. Bonæ-Spei. *Arctotis trifida* Thunb. fl. cap. 711. Cl. Less. specim. Thunb. descripsit. Caulis ex Thunb. pedalis foliorum casu tuberculatus. Folia 2-3-5-fida tenuissimè pubescentia subtus bisulca unguicularia. Pedicelli capit. vix pollicares.

*** *Foliis indivisis integerrimisve.*

41. *S. LEPTOGLOSSA*, suffruticosa tenuis erecta ramosa, ramis subdichotomis glabriusculis, foliis linearibus acutis punctato-glandulosis glabris integerrimis, capitulis terminalibus subsessilibus, invol. squamis exterioribus oblongo-linearibus, ligulis angustissimis subintegris parcè glandulosis. ♂ ad Cap. Bonæ-Spei in distr. Zwelendama legit cl. Burchell et mecum sub n. 6897 cat. geogr. communic. Habitus à genere alienus. Folia 5 lin. longa, $\frac{1}{2}$ lin. lata. Capitula 5-6 lin. diam. (v. s.)

42. *S. TENUIFOLIA*, fruticosa subcæspitosa glabra, foliis lineari-filiformibus integerrimis, pedunculis folio duplò longioribus, invol. squamis extimis oblongis acutis apice subscariosis, radii ligulis concoloribus. ♂ ad Cap. Bonæ-Spei (Forb. ! Eckl. ! n. 1338). *Arctotis tenuifolia* Linn. mant. 288. Linn. f. suppl. 385. Burm. ! herb. Ursinia tenuifolia Poir. dict. 8. p. 259. Spreng. syst. 3. p. 409. Involucro ad *S. nudicaulem* accedit, ferè ab omnibus foliis integerrimis distinctissima. (v. s.)

43. *S. CILIARIS*, suffruticulosa glabra, foliis ad basin confertis lato-linearibus basi distanter et rigidè ciliatis, cæterum integerrimis aut hinc indè serrato-ciliatis, ramis erectis scapiformibus aphyllis aut parcè foliolatis 1-cephalis, invol. squamis exter. oblongo-linearibus fusco-marginulatis, radio amplo discolore, recept. paleis apice denticulatis. ♂ in Africà austr. extrà-trop. ad caput Aghillas ferè ad maris alt. legit cl. Drege pl. exs. n. 3689! Caulis basi erectus aut adscendens, circ. pedalis, in parte aphyllà junior subaraneosus. Fol. 2-2 $\frac{1}{2}$ poll. longa. (v. s.)

Sect. II. XEROLEPIS Less. syn. 244.

Involucri squamæ, etiam extimæ, margine latè scariosæ. — Herbæ perennes aut frutices.

44. *S. FILIPES* (E. Mey. ! in Drege pl. exs. n. 1748), herbacea ramosissima glabra, foliis sessilibus oblongo-lingulatis acutè et profundè inciso-serratis punctulatis, pedicellis axillaribus terminalibusque gracilibus elongatis, invol. squamis ext. lanceolatis acutis margine angustissimè subscariosis, ligulis concoloribus, recept. paleis suberosis, pappi laminis lacteis obovatis. (1) ? ad Cap. Bonæ-Spei in Draakensteensberg alt. 3-400 ped. legit cl. Drege. Folia ferè Sph. serratæ sed invol. diversum. Capitula triplò ferè minora. (v. s.)

45. *S. serrata*, suffruticosa puberula, foliis semiamplexicaulis oblongis lanceolatis grosse serratis, pedunculis pluribus 1-cephalis subcorymbosis folio triplo longioribus, radii ligulis concoloribus. \S ad Cap. Bonæ-Spei Burch.! n. 5665. Eckl. n. 1351. *Arctotis serrata* Thunb. fl. cap. 705. Linn. f. suppl. 385. *Ursinia serrata* Poir. et Spreng. ll. cc. *Sphen. Sonneratii* Cass. dict. 50. p. 205. (v. s.)
46. *S. incisa*, suffruticosa puberula, foliis elongato-oblongis apice subdilatis inciso-serratis, serraturis acuminatis, inferioribus parvis distantibus, super. approximatis longioribus, ramis in pedunculos 1-cephalos in parte nudâ folio quadruplo longioribus, radii ligulis concoloribus. \S ad Cap. Bonæ-Spei in distr. Zwellendam legit cl. Burchell et mecum sub n. 7008 cat. geogr. comm. Folia sesquipoll. longa, basi 2-3 apice 4-5 lin. lata. (v. s.)
47. *S. pilifera* (Ker bot. reg. t. 604), fruticosa ramoso-diffusa, foliis pinnatisectis subcarunculatis patentibus et brevibus hispidulis, lobis linearibus setulâ terminatis, pedicellis elongatis subhispidulis, invol. squamis inf. margine scarioso angusto cinctis obtusis, ligulis discoloribus, paleis truncatis, pappi lamina obovatis achænio paulo brevioribus. \S ad Cap. Bonæ-Spei. Less. syn. 244. *Arctotis pilifera* Thunb. fl. cap. 712 non Berg.
- β . *crithmoides*, foliis glabrescentibus, lobis brevioribus. \S *Arctotis crithmoides* Jacq. h. Schœnbr. 2. t. 155 non Berg. *Ursinia crithmoides* Poir. dict. 8. p. 258.
- γ . *subcanescens*, ramulis foliisque novellis pube adpressâ plus minus canescentibus, foliorum lobis abbreviatis. \S ad Zwellendam legit cl. Drege, pl. exs. n. 6367. (v. s.)
48. *S. scariosa* (R. Brown et Less. ll. cc.), fruticosa glabra, foliis pinnatisectis, segmentis lineari-filiformibus integerrimis mucronulatis, pedunculis folio duplo triplo longioribus, invol. squamis extimis omnino scariosis subrotundis, radii ligulis subscloribus, rec. paleis truncatis, achæniis molliter et parce villosis. \S ad Cap. Bonæ-Spei. *Arctotis scariosa* Willd. sp. 3. p. 2360 excl. Thunb. syn. *Ursinia scariosa* Poir. dict. 8. p. 257. — Burch.! cat. afr. austr. n. 5451. Sæpe in hortis sub nom. *Arctotidis* paleaceæ vagat. (v. s. sp. et c.)
- β . *brevipes*, capitulis supra ultima folia brevissimè pedicellatis. \S ad Cap. Bonæ-Spei Drege! pl. exs. n. 6016. (v. s.)
49. *S. odorata* (R. Brown l. c.), fruticosa glabra, foliis planis apice inciso-pinnatifidis, pappo obsoleto, radii ligulis discoloribus. \S ad Cap. Bonæ-Spei. *Anthemis odorata* Ait. kew. 3. p. 238. Willd. sp. 3. p. 2186. *Achænia glabra* ex Br.

CDXXXII. *URSINIA* Gærtn. fruct. 2. p. 462. t. 174. Less. syn. 244. — *Ursinie* sp. Poir. Spreng. — *Arctotidis* sp. auct.

Omnia *Sphenogynes*, sed 1°. achæmium obovatum turbinatum basi distinctè attenuatum glaberrimum nervulis 5 minimis subcostatum. 2°. Pappus duplex, ext. lamina 5 amplius obovatis lacteis spiralerè æstivato-contortis ut in *Sphenogyne*, int. filis 5 concoloribus angustissimis imâ basi subconcretis. — Herbæ aut frutices Capenses. Folia uni- aut bipinnatisecta. Invol. squamæ pleræque apice in appendicem scariosam productæ.

§. 1. *Species perennes aut frutescentes.*

1. *U. MONTANA*, fruticosa adscendens humilis parcè ramosa, caule ramisve hinc inde subpuberis, foliis confertis pinnatifidis crassiusculis glabris puberis, lobis inf. minimis acutissimis, sup. linearibus mucronulatis, pedicellis nudis longitudine canlis, invol. ovatis squamis ext. ovato-lanceolatis obtusis margine nigrello et scarioso albo cinctis, ligulis parvis discoloribus, paleis truncatis. \S in Cap. Bonæ-Spei ad Willbergen alt. 8-900 ped. legit cl. Drege, pl. exs. n. 6027. Fruct. matur. non vidi. (v. s.)

2. *U. APICULATA*, fruticosa cæspitosa demissa glabra, foliis pinnatipartitis, lobis omnibus apiculo albo mucronatis seu albo-sphacelatis, sup. linearibus indivisis longis trilobisve, pedicellis caule ferè longioribus, invol. squamis ext. ovatis nigro marginatis, ligulis discoloribus, paleis obtusis subdentatis. $\frac{1}{2}$ $\frac{1}{2}$ in Cap. Bonæ-Spei ad Sneeuwbergen alt. 5-600 ped. legit cl. Drege, pl. exs. n. 2132. Habitus *U. montanæ*, sed ab hac et ab omnibus facilè differt fol. lobis apice mucrone albiissimo terminatis. (v. s.)
3. *U. TENUILOBA*, glaberrima, caule suffruticuloso subdemisso brevi parçè ramoso, foliis pinnatipartitis, lobis linearibus aristato-acuminatis, infer. brevioribus integris, super. 2-3-lobis, pedicellis caule longioribus, paleis apice obtusè dentatis, pappo int. laminis exterioribus paulò brevior. $\frac{1}{2}$ $\frac{1}{2}$ in Cap. Bonæ-Spei ad Knt-Riviersberg alt. 5-600 ped. legit cl. Drege, pl. exs. n. 3698 et in distr. Uitenhage prope Zwartkop-Rivier cl. Eckl. n. 107. *Ursinia fœniculacea* Eckl. l. c. non Spreng. Folia ferè *Santolinæ* viridis. Habitus *Achillæ* moschatae. Pappus ext. amplus. (v. s.)
4. *U. PARADOXA* (Gærtn. et Less. ll. cc.), suffruticulosa adscendens glabriuscula, foliis pinnatisectis, lobis inf. brevibus integris, sup. longioribus 2-3-fidis, lobulis oblongo-linearibus apice piliferis, ramis apice nudis pedicelliformibus, invol. squamis omnibus margine scariosis obtusis, ligulis discoloribus, pappo int. exteriore brevior. $\frac{1}{2}$ $\frac{1}{2}$ ad Cap. Bonæ-Spei legit cl. Drege, pl. exs. n. 531. *Arctotis paradoxa* Linn. amœn. 4. p. 330 (excl. syn.). Thunb. fl. cap. 712. *Sphenogyne anthemoides* E. Mey. in Drege coll. non Brown. Habitus omnino *Sphen. anthemoidis*, sed differt caract. gen. (achæn. glabro et pappo duplice) nec non fol. apice piliferis, et invol. squamis omnibus margine scarioso albo cinctis. Capitulum non malè refert *Anacyclum Pyrethrum* seu icon. bot. mag. t. 544. (v. s.)
5. *U. ALBICAULIS*, suffruticulosa erecta, ramis sublanatis, foliis pinnatipartitis glabriusculis, lobis linearibus mucronatis integerrimis aut apice 2-3-fidis, ramis apice nudis pedicelliformibus glabriusculis, invol. squamis ext. in appendicem scariosam lanceolato-acuminatam productis, pappo inter. externum longitudine subæquante. — ad Cap. Bonæ-Spei ad Zwellendam t'Kouze et Gariep, alt. 2-300 ped. legit cl. Drege. Pedicelli circ. 2-3 pollic. longi. Ligulæ intense flavæ subdiscolores 9-10 lin. longæ. Pappus lacteus, exterior amplus. (v. s.)
6. *U. PEDUNCULOSA*, subherbacea, ramis laxè sublanatis, foliis glabris bipinnatisectis, lobis linearibus mucronatis, ramis longè denudatis glabris, invol. squamis ext. in appendicem scariosam lanceolato-acuminatam productis. — ad Cap. Bonæ-Spei loco dicto Klein-Namaqualand alt. 2-300 ped. legit cl. Drege, pl. exs. n. 2807. An ab *U. albicauli* certè diversa? Fruct. matur. ign. tamen in specim. quodam (Eckl. n. 322) huc ex invol. et foliis relato vidi achænia turbinata lævia alba pappo (verisimiliter deciduo) omnino destituta. (v. s.)
7. *U. RADICANS*, caule subherbaceo decumbente basi repente glabriusculo, foliis pinnatipartitis, lobis inf. simplicibus, super. 2-3-lobatis acutè subdenticulatis, omnibus mucrone apiculatis, pedicellis modicè elongatis 1-cephalis, invol. squamis ext. lanceolatis appendice brevi scariosâ subapiculatis, ligulis discoloribus, paleis obtusis apice subdentatis, pappo int. exteriore paulò brevior. $\frac{1}{2}$ ad Cap. Bonæ-Spei in Algoa-bay. verisim. in arenosis maritimis legit cl. Drege, pl. exs. n. 2131. Habitus *Pyrethri* maritimi. Accedit ad *U. speciosam*, sed habitu decumbente et repente facilè diversa. (v. s.)
8. *U. FILICAULIS*, herbacea glabra, caule filiformi prostrato hinc indè radicante, foliis pinnatipartitis, lobis acutis rachique linearibus, infer. indivisis, super. trifidis, pedicellis adscendentibus 1-cephalis, invol. campanulati, squamis ext. in appendicem scariosam acuminatam productis, ligulis discoloribus, paleis obtusis. $\frac{1}{2}$ ad Cap. Bonæ-Spei loco dicto Kaapsche-Vlakte legit cl. Drege, pl. exs. n. 6021 in distr. Caledon? (Eckl. n. 454?). Car. gener. non vidi sed huc refero ex involucrio, achænio juniore, et analogiam maximam cum *U. radicante*. Caules circ. 2 ped. longi graciles. Folia remota. Ligul. crocæ, subtus subrubræ, 7-8 lin. longæ. (v. s.)

§. 2. *Species annuæ.*

9. *U. SPECIOSA*, herbacea erecta ramosa glabra, foliis pinnatipartitis, lobis inf. brevibus integris, sup. longioribus incisus pinnatifidiæ, lobulis omnibus seta apiculatis, pedicellis nudis elongatis, invol. squamis ext. obtusiusculis scarioso-marginatis, ligulis concoloribus, paleis truncatis, pappo intern. laminis externis paulò brevior. ①? in Cap. Bonæ-Spei loc. dict. Zwellendam t'Kouze et Gariep, alt. 2-300 ped. legit cl. Drege. Herba et infl. ferè *Chrysanthemi* carinati.

α. *croceo-flava*, ligulis croceo-flavis. Drege! pl. exs. n. 2745. (v. s.)

β. *albida*, ligulis albidis. Drege! pl. exs. n. 6368. (v. s.)

10. *U. CAKILEFOLIA*, herbacea erecta ramosissima glabra, foliis crassiusculis pinnatipartitis, lobis linearibus acutis, inf. integris, sup. partitis, pedicellis elongatis nudis, invol. squamis ext. in appendicem subscariosam acuminatam productis, ligulis concoloribus, paleis ext. 3-dentatis, pappo inter. externi longitudinem æquante. ① in Cap. Bonæ-Spei ad Giftberg, alt. 2-300 ped. legit cl. Drege, pl. exs. n. 6023! Capitula parva. Flores flavi. (v. s.)

β. *crocea*, ligulis croceis paulò majoribus, invol. squamis appendicibus subnigricantibus, paleis maturis rufidulis. ① ad Cap. Bonæ-Spei legit cl. Drege, pl. exs. n. 6025! (v. s.)

11. *U. INDECORA*, herbacea ramosissima glabrescens, foliis crassiusculis pinnatipartitis, lobis linearibus, inf. integris, sup. tri- aut pluripartitis, pedicellis nudis 1-cephalis elongatis, invol. squamis ext. ovatis subacutis, ligulis concoloribus invol. paulò longioribus, paleis truncatis, pappo int. externum longitudine æquante. ① ad Cap. Bonæ-Spei in Carro, alt. 3-400 ped. legit cl. Drege, pl. exs. n. 772! Habitus *Senecionis*. Affinis *U. cakilefoliæ*. Ligulæ 3-4 lin. longæ. (v. s.)

12. *U. NANA*, herbacea glabra pluricaulis subdiffusa, foliis crassiusculis pinnatipartitis, lobis linearibus obtusiusculis, aliis integerrimis, aliis 2-3-lobatis, ramis apice pedicelliformibus 1-cephalis, invol. squamis ext. nigro marginatis vix subacutis int. apice albo-scariosis rotundatis, ligulis concoloribus invol. paulò longioribus, pappi int. setis laminis ext. brevioribus. ① in Cap. Bonæ-Spei ad Silver-Fontain, alt. 2-300 ped. legit. cl. Drege, pl. exs. n. 2782! Habitus ferè *Sphenogynes* pusillæ, sed magis diffusa et ligulæ minores. *Achaënia* albida glabra basi attenuata nervulis 5 costulata. An huc aut ad sequentem referenda *Urs. annua* Less. in *linnæa* 1831. p. 165, nomine tantum nota? (v. s.)

13. *U. PYGMÆA*, herbacea glabra pluricaulis suberecta, radice exili, foliis pinnatipartitis, lobis paucis lineari-filiformibus, capitulis pedicellatis solitariis, invol. squamis ext. acuminatis. ① in Africæ Capensis distr. Namaqualand legit cl. Ecklon, n. 27. Specimen mancum tantum vidi sed ex fructu ad *Ursiniam* referendum; ex invol. squamis acuminatis ab *U. nanâ* distinguendum. Herba pollicaris. (v. s.)

† *Species exclusæ.*

U. ABROTANIFOLIA Spreng. = *Sphenogyne abrotanifolia*.

U. ANTHEMOIDES Poir. = *Sphenogyne anthemoides*.

U. CRITHMOIDES Poir. = *Sphenogyne pilifera* β.

U. DENTATA Poir. = *Sphenogyne dentata*.

U. FILIFORMIS Griess. = *Sphenogyne pinnata*.

U. FOMICULACEA Poir. = *Sphenogyne fomiculacea*.

U. PILIFERA Poir. = *Sphenogyne dentata*.

U. SCARIOSEA Poir. = *Sphenogyne scariosa*.

U. SERRATA Poir. = *Sphenogyne serrata*.

U. TENUIFOLIA Poir. = *Sphenogyne tenuifolia*.

DIV. III. MADIÆ (1).

Receptaculum margine aut ferè totum paleaceum. Invol. squamæ 1-2-seriales convolutæ aut complicatæ achænia radii semper calva includentes. Achænia disci pappo paleaceo donata aut calva, fertilia aut sterilia. — Folia alterna. Flores flavi.

CDXXXIII. MADIA *Mol. chil. ed. germ.* 113. *ed. gall.* 336. *Cav. ic.* 3. p. 50. — *Madia et Biotia Cass. dict.* 34. p. 308. — *Madia* sp. *Don, Less.*

Capitulum multiflorum heterogamum, fl. radii 10-12 uniserialibus fœmineis ligulatis aut monstrosè irregulariter et amplè tubulosis, fl. disci hermaphr. tubulosis regulariter 5-dentatis. Invol. subglobosum 1-seriale, squamis (tot quot ligulæ) dorso carinatis complicatis achænia radii involventibus. Recept. planum paleas inter radium et discum 1-2-seriales gerens, in centro omnino nudum. Ligulæ basi glabræ invol. vix longiores. Styli fl. disci rami infernè ad latus internum series duas stigmatico-glandulosas gerentes, apice pilis rigidis erectis onusti. Achænia compressa recta basi attenuata glabra calva utrinquè in medio nervo longitudinali aucta et ideò 4-5-gona. — Herba Chilensis annua erecta villosa et apice præsertim glanduloso-pilosa. Folia infima opposita, sup. alterna, semiamplexicaulia oblonga integra. Capitula breviter pedicellata in racemum digesta foliis parvis sæpiùs bracteata. Cor. luteæ. Semina oleosa.

1. *M. SATIVA* (Mol. et Don II. cc.). (1) in Chili spontanea et ad oleum è seminibus eliciendum culta sub nom. vernaculis *Madi* et *Melosa* ex Bertero! Eamdem fortè cultam in Californiâ legit cl. Douglas. *M. sativa*, *M. viscosa* et *M. mellosa* Willd. sp. 3. p. 1051 vix varietates ex Don. *M. viscosa* Cav. ic. 3. p. 50. t. 298. *M. mellosa* Jacq. hort. Schœnbr. 3. p. 29. t. 302. *M. sativa* DC. not. jard. gen. VII. n. 6. mem. soc. Gen. 7. p. 277. (v. v. c. et s. sp.)

* *monstrosa*, ligulis in tubulos ampliatis irregulariter mutatis. — *Biotia* Cass. l. c. (v. v. c.)

† *Species exclusa.*

M. ELEGANS Don. = *Madaria elegans*.

CDXXXIV. MADARIA DC. mem. soc. Gen. v. 7. p. 280. not. pl. rar. jard. Gen. VII. n. 6. — *Madia* sp. *Don.*

Capitulum multiflorum radiatum, ligulis 10-12 uniserialibus fœmineis, fl. disci tubulosis 5-dentatis masculis seu sterilibus. Invol. subglobosum 1-seriale torulosum, squamis complicatis achænia radii omnino involventibus. Recept. subconvexum, margine inter ligulas et discum squamas 1-2-seriales gerens, medio epaleaceum densè et breviter fimbriatifero-hirsutum. Ligulæ involucri duplò longiores et corollæ tubulosæ basi puberulo-hirtellæ. Styli rami apice hispiduli. Achænia radii compressa nervis in latere utroque aucta et ideò sub-4-5-gona, glabra, calva basi rostello incurvulo donata, disci omnino abortiva. — Herba Californicæ annuæ erectæ villosæ et sæpè glandu-

(1) Genera sex hic habitu toto consociata inter se characteribus normalibus valdè differunt, et strictissimo sensu Hartmannia et Madaroglossa fortè ad Galinsogea, Hemizonia et fortè Madaria ex disci achæniis calvis et abortivis ad Parthenias, Madia et Oxyura ex disci achæniis calvis et fertilibus ad Heliopsideas essent referenda? sed in ordine naturali, ut in civili, summum jus sæpè summa injuria.

loso-pilosæ. Folia sessilia lanceolata aut linearia integra: Capitula in corymbum laxum disposita, longius pedicellata ebracteata. Flores flavi. — Nomen ex *μάδαρος* calvus.

1. *M. ELEGANS*, caule foliisque pilis apice glandulosis et setis eglandulosis mixtis hispido. ① in Californiâ detexit cl. Douglas. *Madia elegans* Don in Lindl. bot. reg. t. 1458. Ligulæ basi purpureo-maculatæ, lineari-cuneatæ, apice acutè trifidæ. Achænia radii intrâ inv. squamas inclusa subbiseriata nigra compresso-trigona, in quoque latere uninervia, undè ferè 5-gona apice truncata omninò calva. (v. s. sp. et v. c.)
2. *M. CORYMBOSA*, caule involucriisque pilis apice glandulosis et setis eglandulosis hispido, foliis linearibus villosis subhispidis eglandulosis. ① in Amer. bor.-occid. ad Columbia-Rivier legit cl. Douglas. Variat pilis glandulosis plurimis aut paucissimis. Achænia compresso-trigona, in specim. fortè non omninò matura, rufa submaculata. (v. s.)

ℒ? *hispida*, caule foliis involucriisque pilis omnibus eglandulosis patulis hispidis. ① in Californiâ legit cl. Douglas. An fortè species propria? (v. s. comm. cum var. *α* ab hon. Soc. hort. Lond.)

CDXXXV. HEMIZONIA.

Capitulum multiflorum radiatum, ligulis 1-serialibus foemineis invol. paulò longioribus, fl. disci tubulosis 5-dentatis masculis seu sterilibus. Invol. squamæ 1-seriales concavæ et achænia radii semi-involventes. Recept. planum paleaceum aut saltè inter radium et discum paleas 1-seriales gerens et medio epaleaceum punctatum. Cor. basi glabræ. Styli rami obtusiusculi. Achænia radii obovata ferè obcompressa extus convexa intus plana glabra calva, disci abortiva. — Herbe Californicæ erectæ ramosæ basi suffruticulosæ pilosiusculæ interdum glanduliferæ. Folia alterna conferta lanceolata aut linearia, integra aut serrata. Capitula terminalia solitaria aut pauca, bracteis foliaceis plurimis cincta. Flores flavi. — Genus omninò medium inter Madariam et Hartmanniam. Nomen ex *ἡμιζωνος* (semicinctus) alludit ad achænia invol. squamis semicincta.

Sect. I. MADARIOIDES.

Recept. margine paleas 1-seriales gerens, medio nudum planum punctatum.

1. *H. CONGESTA*, caule herbaceo erecto ramoso, ramis pilis glandulosis et eglandulosis molliter hispidulis, foliis lineari-lanceolatis parè villosis-glandulosis, capitulis bracteatis congestis, achæniis obovatis apice obtusis. — in Californiâ legit cl. Douglas. Achænia nigra, potiùs obcompressa quàm lateribus compressa, matura basi producta in stipitem rostelliformem insigniter recurvum ac areolam albam gerentem. Recept. paleæ subcoriaceæ foliaceæ. (v. s. comm. ab hon. Soc. hort. Lond.)
2. *H. ANGUSTIFOLIA*, caule ramosissimo imò basi anfruticuloso, ramulis foliis involucrique minutè puberulis, foliis bracteisque angustè linearibus integerimis, capitulis bracteatis subsolitariis, achæniis obovatis apice cono brevi acuto mucronatis. 3 in Californiâ legit cl. Douglas. Recept. paleæ angustæ membranaceæ. Achænia estipitata areolâ obliquâ, supernè conulo brevi mucronata. Eandem ni fallor video in h. Hænke, sed Regiomontana dicitur? (v. s. comm. cum priorib.)

Sect. II. OLOCARPHA.

Paleæ recept. discum totum occupantes.

3. *H. LUXULÆFOLIA*, caule molliter hirsutulo, foliis inferioribus oppositis, super. alternis, elongato-linearibus plurinerviis subdenticulatis molliter villosis eglan-

dulosi, corymbo laxo, invol. squamis hispidulis, achæniis radii muticis. — in Novâ-Californiâ legit cl. Douglas. Ligulæ 15-20. Capitula parvissime bracteata. Achænia estipitata areolâ laterali. (v. s. comm. ab hon. Soc. hortic. Lond.)

4. *H. MACRADENTIA*, caule suffruticoso ramosissimo, ramulis foliisque pilosis, foliis linearilanceolatis, infer. parce serratis, cæteris integris, rameis super. bracteisque glandulâ grossâ truncatâ terminatis, invol. squamis grossè et callosè inciso-serratis, paleis apice subulatis ciliatis, achæniis obovato-turbinatis apice incurvis rostro subadscendente. ♀ in Californiâ legit cl. Douglas. Species priori primâ fronte simillima sed verè distinctissima et structurâ in ordine anomala spectabilis. Affinis Calycadeniæ sed ligulæ planæ, etc. (v. s. comm. ab hon. Soc. hortic. Lond.)

CDXXXVI. *OXYURA DC. in Lindl. nat. syst. bot. ed. 2. p. 256.*

Capitulum multiflorum radiatum, ligulis 10-12 uniserialibus fœmineis, fl. disci tubulosi 5-dentati bisexualibus, centralibus plurimis abortivis. Invol. squamæ 1-seriales basi rigide ciliatæ convolutæ et achænia radii omnino involventes, supernè in appendicem linearem foliaceam productæ. Recept. totum paleis membranaceis inter flores onustum. Ligulæ basi pubentes apice grossè dentatæ. Tubuli obcompressi ad lobulos pubentes. Anth. nigricantes. Styli disci rami in appendicem hispidam acutissimam producti. Achænia omnia obcompressa obovato-oblonga glabra calva, centralia sæpè abortiva. — Herba Californica annua erecta *Chrysanthemum* segetum habitu referens. Caulis teres purpurascens glaber. Folia alterna ciliata pinnatifida, lobis 5-8-jugis integris. Capitula ramulos apice nudiusculos hirsutos terminantia. Flores flavi. Bractææ paucæ ciliatæ. — Nomen ex οξύς, acutus, et ὤρα, cauda.

1. *O. CHRYSANTHEMOIDES*. ① in Novâ-Californiâ legit cl. Douglas. Radix simplex perpendicularis. Caulis circ. pedalis. (v. s. comm. ab hon. Soc. hortic. Lond.)

CDXXXVII. *HARTMANNIA*.

Capitulum pluri- aut multiflorum heterogamum, fl. radii uniserialis fœmineis ligulatis grossè apice dentatis, disci hermaphr. tubulosi 5-dentatis. Invol. squamis oblongo-lanceolatis acutis convolutis flores radii semiamplectentibus, bracteis nonnullis linearibus extus cinctis. Recept. paleaceum, paleis squamas referentibus sed planiusculis. Cor. disci vix apice puberulæ. Antheræ nigricantes. Stylus, disci ramis cono acuto apiculatis. Achænia radii tri- sæpè tetragona obovata subpedicellata, dorso subconvexa; apice calva, areolâ terminali obliquâ interiùs sitâ interdum papilloso-exsertâ, disci ovali-oblonga obcompressa rarò calva sæpius paleis 5 membranaceis enerviis obtusis corollâ dimidiò brevioribus coronata. — Herbæ Californicæ erectæ. Caulis pilis patulis hispidus simplex apice corymboso-ramosus. Folia alterna, infima interdum opposita, sessilia lineari-oblonga, pinnatifida aut serrata pubera. Capitula ad apices ramorum corymbosa aut fasciculata. Cor. flavæ. — Genus meo sensu ex habitu naturale, sed in sectiones tot quot species faciliè distinguendum, Helenicis ex pappo paleaceo et antheris nigris pertinet, sed transitum ad Anthemicas exhibet! — Genus distinctissimum dixi in honorem cl. peregrinatoris Jos. Adolph. Hartmann qui, anno 1739, Californiam adiit et ejus descripsit. edidit.

1. *H. FASCICULATA*, foliis inciso-dentatis imâ basi obtusè et brevè auriculatâ.

capitulis ad apices ramorum dense fasciculatis 10-11-floris, ligulis 5 obovato-cuneatis, achæniis lævibus, radii calvis, disci paleaceo-papposis. 2P in Novâ-Californiâ legit cl. Douglas. Habitus et inflorescentia ferè Lini flavi aut campanulati. Recept. paleæ inter radium et discum in verticillum ferè disposita. (v. s. comm. ab hon. Soc. hortie. Lond.)

2. *H. corymbosa*, foliis pinnatipartitis basi attenuatis petiolatisve, lobis linearibus 5-9-jugis integris, capitulis ad apices ramorum subsolitariis multifloris, invol. pedicellisque dense hispidulis, ligulis 15-20 oblongis, achæniis pedicellatis subtragonis dorso muricatis, radii calvis, disci exter. calvis, cæteris paleaceo-papposis. ① in Novâ-Californiâ legit cl. Douglas. Caules et folia pubera. Habitus Anthemidearum. Areola fructus terminalis valdè lateralis exserta! (v. s. comm. cum priore.)
3. *H. ciliata*, caule glabrinsculo, apice laxè ramoso vix corymboso, foliis pinnatipartitis, lobis integris 5-8-jugis, infer. in petiolum attenuatis, super. semiamplexicaulibus floralibusque ciliatis, capitulis multifloris, ligulis 10-12 obovatis amplis, achæniis omnibus lævibus calvis. ① in Novâ-Californiâ legit cl. Douglas. Habitus omninò prioris, sed ex achæniis omnibus calvis areolâ terminatis, et cor. tubo plano-obcompresso proximè ad tribum sequentem accedit. Capitulum ferè Chrysanthemi Myconis. (v. s. comm. cum prioribus.)

CDXXXVIII. MADAROGLOSSA.

Capitulum multiflorum radiatum, ligulis 1-seriatis foemineis, fl. disci hermaphr. tubulosis breviter 5-dentatis. Invol. squamæ uniseriales acuminatæ basi convolutæ. Recept. paleis lineari-lanceolatis submembranaceis onustum. Styli disci ramis inclusis. Achænia radii linearia basi attenuata calva glabra intra squamas inv. omninò inclusa, disci lineari-clavata pappo 8-10 aristato, aristis ad basin ciliato-plumosis apice scabris. — Herbæ Californicæ erectæ diversè piloso-scabræ verisim. annuæ aut biennes. Caulis teres. Folia alterna sessilia oblonga acuta inciso-dentata aut integra. Capitula ramulos terminantia subcorymbosa. Discus radiusque flavi. — Ex pappo ciliato accedere videtur ad Blepharipappum, sed differt radio calvo, fl. disci apice non longè barbatis, styli ramis disci non exsertis et floribus homochromis. — Nomen ex *μαδαρος* calvus et *γλωσσα* lingua.

1. *M. hieracioides*, setis patentibus sparsis rigidulis undiquè hispida, caule subsimplici, foliis oblongo-lanceolatis grossè inciso-serratis, ramulorum floralium linearibus integris, ligulis circ. 15 ovali-oblongis invol. paulò longioribus integris. — in Californiâ legit cl. Douglas. Herba pedalis habitu ferè Echii. Setæ à basi nigræ ortæ. Folia 2 poll. longa, 6-7-lin. lata. Achænia disci villosiuscula, pappi aristis fuscis basi villosis. (v. s. comm. ab hon. Soc. hortie. Lond.)
2. *M. angustifolia*, tota setis rigidis patulis confertis infernè hispida versus apicem puberulo-scabrida, caule ramoso, foliis linearibus integris, ligulis 12-13 trifidis invol. duplò ferè longioribus. ① in Californiâ legit indefessus Douglas. Herba pedalis ferè Echii facie. Radix simplex tortuosa. Folia 1-1½ pollicem longa, 1-1½ lin. lata, inferiora hispidissima. Flores pallidè flavescens. Achænia radii adpressè villosa. Pappus albissimus, setis longè ad basin ciliato-plumosis. (v. s.)
3. *M. heterotricha*, pilis sparsis brevibus remotis aliis pubiformibus aliis glandulâ nigrâ terminatis subscabrida, caule ramoso, foliis lineari-oblongis obtusis integris aut vix subserratis, ligulis apice trifidis invol. duplò ferè longioribus. — in Californiâ legit cl. Douglas. Caulis pedalis et ultrâ. Folia 2 poll. longa, lineam lata. Ligulæ 12-14 amplè multiveniæ. Achænia disci glabra. Pappus albissimus setis basi longè ciliato-plumosis. (v. s. comm. cum prioribus.)

CDXXXIX. CALYCADENIA.

Capitulum multiflorum, fl. radii 1-serialibus 3-5 fœmineis corollâ irregulari ampliata intus fissâ, disci masculis plurimis tubulosis. Invol. extus bracteolatum 1-2-seriale, squamis basi concavis et achænia radii semi-fœventibus. Recept. margine inter radium et discum paleas 1-seriales gerens, medio nudum. Achænia radii trigona subobcompressa glabra calva, disci linearia abortiva. — Herba Californicæ erectæ strictæ. Caulis teres simplex. Folia infer. opposita, super. alterna, rigida linearia margine revoluta integerrima floralia, secus ramulos florigeros conferta, apice glandulam omnino Calycii formam referentem gerentia. Capitulum nunc ad ramulorum axillarium apicem solitaria, nunc ad apicem caulis congesta. Flores lutei. — Nomen alludit ad formam glandularum omnibus speciebus solemniū.

1. *C. TRUNCATA*, caule virgato glaberrimo, foliis linearibus minutissimè subscabridis, infer. apice mucronatis, summis floralibusque apice glandulam subsessilem calyciformem gerentibus, ramulis axillaribus brevissimis 1-cephalis bracteolas plurimas apice glandulosas gerentibus, invol. squamis dorso subscabris apice truncato-retusis, pappo disci squamellis 7-8 brevissimis apice dentato-incisis. — in Californiâ legit cl. Douglas. Caulis nitidus rubens. Capitula remota. Achænia radii subrugosa. (v. s. comm. cum sequentibus ab hon. Soc. hort. Lond.)
2. *C. VILLOSA*, caule villosa strieto, foliis linearibus obtusiusculis margine revolutis basi ciliatis floralibusque ciliato-subhirsutis apice glandulam pedicellatam calyciformem gerentibus, ramulis axillaribus brevibus 1-cephalis. ① in Californiâ legit cl. Douglas. Radix exilis. Capit. specim. mei juniora non ritè describenda, sed ex habitu species evidentissimè congener. (v. s.)
3. *C. MULTIGLANDULOSA*, caule stricto sparsè hispidulo, foliis linearibus margine revolutis subacutis basi ciliato-scabris, ramulis axillaribus brevissimis monocephalis, foliis floralibus plurimis secus ramulos quasi fasciculatis rigidulè ciliatis apice et secus margines prope apicem glandulas pedicellatas plurimas gerentibus, invol. squamis acutis. — in Californiâ legit cl. Douglas. Species glandulis numerosis calyciformibus spectabilis. Pappi disci squamellæ 5 acuminatæ. (v. s.)
4. *C. CEPHALOTES*, caule stricto præsertim apice pubescente, foliis infer. oppositis, super. alternis, elongato-linearibus glabriusculis glabris basi ciliatis, capitulis ad apicem caulis adgregatis subsessilibus, foliis floralibus paucis brevibus glandulâ calyciformi superatis. ① in Californiâ legit cl. Douglas. Achænia radii trigona læviuscula, disci squamellis acuminatis coronata. (v. s.)

ADDENDA ET EMENDANDA.

- P.* 30, *lin.* 12, *PENTACANTHA*, *lege* *PENTACONTA*.
P. 30, *lin.* 19, *hexacantham*, *lege* *hecatantham*.
P. 65, *lin.* 45, 46 et 48, et *p.* 66. *lin.* 8, *Echinopodium*, *lege* *Chresta*.
P. 66, *lin.* 47, *sub* *D. reticulato*, *adde syn.*: *Rolfinkia centauroides* Zenk. *pl. ind.* 2. *p.* 14. *t.* 14.
P. 109, *lin.* 7, *t.* 3?, *lege* *t.* 5?
P. 110, *lin.* 7, *MIKANOIDES*, *lege* *MIKANOIDES*.
P. 125, *lin.* 11, *dele syn.* *Nuttallii*.
P. 255, *lin.* 27, *adde syn.*: *Galatella punctata* Lindl. *bot. reg.* *t.* 1818.
P. 275, *lin.* 25, *dele syn.* *Lessingii*.
P. 283, *lin.* 8, *adde* *DC.* non *Bess.* — *Heterochaeta* *Bess.* est *Avena tenuis*.
P. 295, *lin.* 9, *adde*: *Erigeron filiforme* Spreng. est ex *Schlect. in litt.* *Diplopappus pinifolius* Less. *ined.*
P. 298, *lin.* 14, *disci*, *lege* *radii*.
P. 313, *lin.* 38, *adde syn.*: *DC.* *pl. rar. jard. Gen. not. VII.* *p.* 1. *t.* 1.
P. 374, *lin.* 20, *LESTADIA*, *lege* *LÆSTADIA*.
P. 376, *lin.* 27, *LÆNNECIA*, *lege* *LÆNNECIA*.
P. 398, *lin.* 8 à *fine*, *Tursonia*, *lege* *Tursenia*.
P. 415, *lin.* 11, *dele syn.* *Sprengelii ex Schlect. in litt.*, *ad* *Luciliam acutifoliam referendum*.
P. 428, *lin.* 8 à *fine*, *t.* 27, *lege* *t.* 47, et *dele* *Chrysocoma punctulata* *t.* 49.
P. 449, *lin.* 41, *dele* et *Chlænobolus*.
P. 485, *lin.* 47, *adde syn.*: *Hook. bot. mag.* *t.* 3466.
P. 503, *lin.* 45, *Petrantha*, *lege* *Tetrantha*.
P. 511, *lin.* 2, *Guardiola*, *lege* *Guandiola*.
P. 521, *lin.* 35, *Poir.*, *lege* *Poiteau*.
P. 524, *lin.* 37, *ACANTHOLÆNA*, *lege* *CENTROLÆNA*.
P. 531, *lin.* 11, *κονία*, *pulveris*, *lege* *κονις*, *pulvis*.
P. 543, *lin.* *ult.*, *Jægera* *Raensch. et Gisek.* est *genus inter Scitamineas dubium*.
P. 551, *lin.* 38, *adde syn.*: *DC.* *pl. rar. jard. Gen. not. VII.* *p.* 5. *n.* 2. *t.* 2 et 3.
P. 585, *lin.* 49, *adde syn.*: *DC.* *pl. rar. jard. Gen. not. VII.* *p.* 8. *n.* 3.
P. 586, *lin.* 46, *adde syn.*: *DC.* *pl. rar. jard. Gen. not. VII.* *p.* 9. *n.* 4.
P. 606, *lin.* 6, *Georgla*, *lege* *Georgiæ spec.*

N. B. Hujus voluminis per morbum gravem auctoris impressi Omissis et Sphalmatibus indulgeat benevolus Lector! Species addendæ, ad calcem ordinis Compositarum seu voluminis sexti brevi tempore edendi, reperiendæ sunt.

INDEX NOMINUM ET SYNONYMORUM

PARTIS QUINTÆ.

Obs. Nomina admissa litteris romanis, Synonyma italica traduntur, nempe in utrâque classe:

Ordinalia majoribus,
Tribualia majusculis,
Generica vulgaribus,
Sectionalia minoribus.

<i>Acantholæna DC.</i>	460	<i>Albertinia Spreng.</i>	80
<i>Acanthosperma Arrab.</i>	3	ALBERTINIEÆ DC.	78
<i>Acanthospermum Schrank.</i>	521	<i>Alcina Cav.</i>	520
<i>Acanthoxanthium DC.</i>	523	<i>Alcina DC.</i>	<i>ibid.</i>
ACHYRIDEÆ DC.	212, 312	<i>Alcinia Steud.</i>	<i>ibid.</i>
<i>Achyrocoma Cass.</i>	15	<i>Alciope DC.</i>	209
<i>Achyropappus H. B. K.</i>	654	<i>Aldama Lallav. et Lex.</i>	561
<i>Acicarpha Juss.</i>	3	<i>Alibum Less.</i>	97
<i>Acilepis Don.</i>	15	<i>Alix Comm.</i>	319
<i>Acilepis DC.</i>	<i>ibid.</i>	<i>Allagopappus Cass.</i>	477
<i>Aciphyllæa DC.</i>	641	<i>Allagopappus DC.</i>	<i>ibid.</i>
<i>Acispermum Neck.</i>	569	<i>Allocarpus H. B. K.</i>	676
<i>Acisella Rich.</i>	620	<i>Alloispermum Willd.</i>	<i>ibid.</i>
<i>Acmella DC.</i>	<i>ibid.</i>	<i>Alomia H. B. K.</i>	105
<i>Acosta DC.</i>	90	ALOMIÆ Less.	103
<i>Actinæa Juss.</i>	662	<i>Alymnia Neck.</i>	515
<i>Actinella Pers.</i>	<i>ibid.</i>	<i>Alymnia DC.</i>	<i>ibid.</i>
<i>Actinella Less.</i>	<i>ibid.</i>	<i>Amblyolepis DC.</i>	687
<i>Actinolepis DC.</i>	655	<i>Ambrosia Linn.</i>	525
<i>Actimertis Raf.</i>	575	AMBROSIACÆ Less.	506
<i>Actinomeris Nutt.</i>	<i>ibid.</i>	AMBROSIAE Cass. Less.	498, 522
ACTINOPHYTUM Neck.	4	AMELLEÆ DC.	211, 213
<i>Actinoptera DC.</i>	543	AMELLOIDÆ Less.	213.
<i>Actinospermum Ell.</i>	653	<i>Amellus Adans.</i>	226
<i>Adenocaulon Hook.</i>	207	<i>Amellus Cass.</i>	213
<i>Adenocyclus Less.</i>	10	<i>Amellus P. Br.</i>	544
<i>Adenolepis Less.</i>	607	<i>Amellus Linn.</i>	96, 213
<i>Adenophyllum Pers.</i>	630	<i>Ampherephis H. B. K.</i>	70
<i>Adenostemma Forst.</i>	110	<i>Amphiachysis DC.</i>	313
ADENOSTYLEÆ Cass. DC.	126	<i>Amphicalea DC.</i>	672
<i>Adenostyles Cass.</i>	203	<i>Amphirhapis DC.</i>	343
<i>Agarista DC.</i>	569	<i>Amphistephium DC.</i>	273
<i>Agathæa Cass.</i>	223	<i>Anacis Schrank.</i>	567
AGERATEÆ Less.	107	<i>Anaglypha DC.</i>	311
<i>Ageratoides DC.</i>	108	<i>Anaëtis DC.</i>	628
<i>Ageratum Linn.</i>	<i>ibid.</i>	<i>Andrieuxia DC.</i>	559
<i>Ageratum Auct.</i>	107, 108	<i>Andromachia Cass.</i>	96
<i>Aglossa DC.</i>	543	<i>Andromachia H. B. K.</i>	95
<i>Agrianthus Mart.</i>	125	<i>Andryaloides DC.</i>	377
<i>Aiolothea DC.</i>	508	ANGIANTHÆÆ DC.	500
<i>Alarçonia DC.</i>	537	<i>Anhebecarpa DC.</i>	221

<i>Anisochæta</i> DC.	109	BELLIEAE DC.	212, 302
<i>Anisotrichia</i> DC.	80	<i>Bellis</i> Linn.	304
<i>Anomostephium</i> DC.	560	<i>Bellium</i> Linn.	303
<i>Anonymos</i> Walt.	128	<i>Berlandiera</i> DC.	517
ANTENNARIEAE.	500	<i>Berthelotia</i> DC.	375
<i>Anthalmum</i> Neck.	487	<i>Bidens</i> Linn.	593
ANTHEMIDEAE.	499	BIDENTIDAE Less.	498, 593
<i>Anthemiopsis</i> Boj.	547	<i>Bigelowia</i> DC.	329
<i>Anthemoides</i> Less.	681	<i>Biotia</i> DC.	264
<i>Anvillea</i> DC.	487	<i>Biotia</i> Cass.	691
<i>Apalus</i> DC.	507	<i>Blainvillea</i> Cass.	492
<i>Aphanostephus</i> DC.	310	<i>Blanchetia</i> DC.	75
<i>Aplodiscus</i> DC.	350	<i>Blepharipappus</i> Hook.	679
<i>Aplopappus</i> Cass.	345	<i>Blepharispermum</i> Wight, DC.	368
<i>Arctogeron</i> DC.	260	<i>Blepharodon</i> DC.	346
<i>Arctotis</i> Auct.	681, 688	<i>Blumea</i> DC.	432
<i>Argophyllæa</i> DC.	266	<i>Bobartia</i> Petiv.	554
<i>Argyrochæta</i> Cav.	532	<i>Boebera</i> Less.	639
<i>Argyrochæta</i> DC.	ibid.	<i>Boebera</i> Willd.	ibid.
<i>Argyroxiphium</i> DC.	668	<i>Boebera</i> DC.	640
<i>Armania</i> Bert.	576	<i>Boeberoides</i> DC.	ibid.
<i>Arrhenachne</i> Cass.	398	<i>Bojeria</i> DC.	94
ARTEMISIEAE Less.	499	BOJERIEAE DC.	ibid.
<i>Artemisioides</i> DC.	105	<i>Boltonia</i> L'Hér.	301
<i>Ascaricida</i> Cass.	15, 60	<i>Bombycillæna</i> DC.	460
<i>Ascaricida</i> DC.	60	BOOPIDEAE Cass.	1
<i>Aspilina</i> Pet.-Th.	561	<i>Boopis</i> Juss.	2
<i>Astemma</i> Less.	502	<i>Borrichia</i> Adans.	488
<i>Aster</i> Adans.	253	<i>Botryadenia</i> F. et M.	309
<i>Aster</i> Nees.	226	<i>Brachycome</i> Cass.	305
ASTEREAE.	211	<i>Brachycome</i> Gaud.	280
ASTERINEAE DC.	211, 213	<i>Brachycome</i> Cun.	305
ASTEROIDEAE Less.	8, 211	<i>Brachyglossa</i> DC.	265
<i>Asteriscus</i> Moench.	486	<i>Brachyglottis</i> Forst.	210
<i>Asteroides</i> Tourn.	483, 485, 486	<i>Brachylæna</i> Brown.	429
<i>Asteromea</i> Blum.	302	<i>Brachyris</i> Nutt.	312
<i>Asteropsis</i> Less.	260	<i>Brachyscoma</i> Cass.	305
<i>Athanasia</i> Walt.	680	<i>Brauneria</i> Neck.	554
ATHANASIEAE.	499	<i>Brickellia</i> Ell.	182
<i>Athenæa</i> Adans.	11	<i>Brotera</i> Spreng.	635
<i>Athroisma</i> DC.	368	<i>Brotera</i> DC.	ibid.
<i>Athronia</i> Neck.	620	<i>Babonium</i> DC.	463
<i>Aurelia</i> Cass.	314	<i>Bulbostylis</i> DC.	138
BACCHARIDEAE Less.	212, 368	BUPHTHALMEAE Less.	213, 483
BACCHAREAE DC.	398	<i>Buphthalmum</i> Cass.	483
<i>Baccharis</i> Linn.	ibid.	<i>Barriella</i> DC.	663
<i>Baccharoides</i> Linn.	60	<i>Bustamenta</i> bot. mex.	166-168
<i>Bahia</i> Lag.	656	<i>Bustia</i> Adans.	483
<i>Bailliera</i> Cass.	505	<i>Caecalia</i> Tourn.	203
<i>Balbisia</i> Willd.	679	<i>Caecomia</i> H. B. K.	97
BALBISIEAE DC.	500	<i>Cæsulia</i> Roxb.	482
<i>Balduina</i> Nutt.	652	CAESULIAE DC.	ibid.
<i>Balsamorhiza</i> Hook.	551	<i>Calea</i> R. Br.	671
<i>Baltimore</i> Linn.	509	<i>Calea</i> Linn.	ibid.
BALTIMOREAE Cass.	500	<i>Caleacte</i> R. Br.	673
<i>Bartolina</i> Adans.	641, 679	<i>Caleacte</i> Less.	669
<i>Bechium</i> DC.	70	<i>Caleacte</i> DC.	673
<i>Bellidiastrum</i> Michx.	226	<i>Calebrachys</i> Cass.	669, 669
BELLIDEAE DC.	212, 304	<i>Calebrachys</i> DC.	669

NOMINUM ET SYNONYMORUM.

699

<i>CALEINEÆ</i> Cass.	668	<i>Chromolæna</i> DC.	133
<i>Caliceras</i> Cav.	2	<i>Chronolæna</i> DC.	397
<i>Calimeris</i> Cass. <i>Nées.</i>	258	<i>Chronopappus</i> DC.	84
<i>Callilepis</i> DC.	671	<i>Cryptopetalum</i> Cass.	101
<i>Calliopsis</i> Reichenb.	568	<i>Cryptopetalum</i> DC.	ibid.
<i>Callistemma</i> Cass.	274	<i>Chrysactinium</i> H. B. K.	97
<i>Callistephus</i> Cass.	ibid.	<i>Chrysanthellina</i> Cass.	630
<i>Callophora</i> DC.	542	<i>Chrysanthellum</i> Rich.	630
<i>Calonnea</i> Buchoz.	651	CHRYSANTHEMEAE.	499
<i>Calotis</i> B. Br.	302	<i>Chrysis</i> Ren.	585
<i>Calycadenia</i> DC.	695	<i>Chrysocoma</i> Cass.	353
<i>Calycera</i> Rich.	2	<i>Chrysocoma</i> Linn.	351, 353
CALYCERAE R. Br.	1	CHRYSOCOMEAE DC.	212, 310
<i>Calycium</i> Ell.	316	<i>Chrysogonum</i> Linn.	510
<i>Calydermos</i> Lag.	669	<i>Chrysomelea</i> Tausch.	569
<i>Calydermos</i> Cass.	ibid.	<i>Chrysophania</i> Kunth.	553
<i>Calydermos</i> Spreng.	11	CHRYSOPSIDEAE DC.	212, 326
<i>Calypocarpus</i> Less.	629	<i>Chrysopsis</i> Nutt.	326
<i>Campuloclinium</i> DC.	136	<i>Chrysosphaerium</i> Willd.	673
<i>Campylothea</i> Cass.	593	<i>Chrysostemma</i> Less.	567
<i>Camutia</i> Bonat.	520	<i>Chthonia</i> Cass.	98
<i>Cappa</i> DC.	469	<i>Chylodia</i> Rich.	563
<i>Carelia</i> Less.	125	CICHORACEAE Vaill.	4, 9
<i>Carelia</i> Ponted.	108	<i>Clairvillea</i> DC.	636
<i>Carphephorus</i> Cass.	132	<i>Clavigera</i> DC.	127
<i>Carphostephium</i> Cass.	678	<i>Clibadium</i> Linn.	505
<i>Curtisia</i> Cass.	71	<i>Clomenocoma</i> Cass.	641
<i>Carthamus</i> Hill.	ibid.	<i>Cœlestina</i> Cass.	107
CASSINIEÆ.	500	<i>Coleosanthus</i> Cass.	136, 138
<i>Celmisia</i> DC.	209	<i>Collœa</i> Spreng.	630
<i>Celmisia</i> Cass.	ibid.	<i>Commidendrum</i> DC.	344
<i>Centauropsis</i> Bojer.	93	COMPOSITAE Vaill.	4
<i>Gentipeda</i> Less.	371	COMPOSITIFLORÆ Gært.	ibid.
<i>Centropalus</i> Cass.	15, 60	<i>Coniothele</i> DC.	531
<i>Centratherum</i> Cass.	70	<i>Conoclinium</i> DC.	135
<i>Centratherum</i> Less.	69	<i>Contarena</i> Adans.	88
<i>Centrocarpha</i> Don.	555	<i>Conyza</i> Less.	377
<i>Centrolæna</i> DC.	524	<i>Conyza</i> Ait. Lam.	69
<i>Centrospermum</i> Kunth.	521	<i>Conyza</i> Linn.	377, 398
<i>Cephalophora</i> Cav.	661	<i>Conyza</i> Wall.	77
<i>Ceratocephalus</i> Vaill.	593	CONYZEAE Less.	212, 368
<i>Ceratoclæna</i> DC.	522	COREOPSIDEAE Less.	498, 569
<i>Cercostylus</i> Less.	660	<i>Coreopsis</i> Linn.	569
<i>Ceruana</i> Forsk.	488	<i>Coreopsis</i> Auct. 562, 568, 569,	575
<i>Ceruchis</i> Gært.	620	<i>Coreopsoides</i> Mœnch.	569
<i>Chænactis</i> DC.	659	<i>Corethrogyne</i> DC.	215
<i>Chætactis</i> DC.	317	<i>Gorona-solis</i> Tourn.	585
<i>Chætopappa</i> DC.	301	<i>Corvisartia</i> Merat.	463
<i>Chætophora</i> Nutt.	ibid.	<i>Corvisartia</i> DC.	ibid.
<i>Chakiatella</i> Cass.	563	CORYMBIFERE Vaill.	4
<i>Chamæstephanum</i> Willd.	654	<i>Corymbium</i> Linn.	88
<i>Charieis</i> Cass.	300	<i>Cosmea</i> Willd.	606
<i>Chiliadenus</i> Cass.	476	<i>Cosmos</i> Cav.	ibid.
<i>Chiliadenus</i> DC.	ibid.	<i>Cosmus</i> Pers.	ibid.
<i>Chiliophyllum</i> DC.	554	COTULEAE Less.	499
<i>Chiliotrichum</i> Cass.	216	<i>Crassicola</i> DC.	373
<i>Chlænobolus</i> Cass.	449, 453	<i>Crassina</i> Scop.	554
<i>Chlænobolus</i> DC.	453	<i>Crinita</i> Mœnch.	351
<i>Chresta</i> Arrab.	85	<i>Crinitaria</i> Cass. Less.	ibid.
<i>Chromochæta</i> DC.	351	<i>Critonia</i> P. Br.	140

<i>Critonia</i> Gærtn.	126	<i>Dubaldea</i> DC.	366
<i>Crodisperma</i> Poit.	563	<i>Dunantia</i> DC.	626
<i>Cryphiospermum</i> Beauv.	497	<i>Dysodia</i> DC.	639
<i>Cryptocarpa</i> R. Br.	3	<i>Dysodium</i> Rich.	519
<i>Cusimbua</i> DC.	650	<i>Dyssodia</i> Cav.	639
<i>Cuspidella</i> DC.	370	<i>Dyssodia</i> Willd.	ibid.
<i>Cyanopis</i> Blum.	69		
<i>Cyanthillium</i> Blum.	ibid.	<i>Echinacea</i> Mœnch.	554
<i>Cyathocline</i> Cass.	373	<i>Echinolema</i> Jacq.	3
<i>Cyathophora</i> DC.	538	<i>Eclypta</i> Linn.	489
<i>Cylindrocline</i> Cass.	458	<i>ECLYPTEAE</i> Less.	213, 488
<i>Cynara</i> Thunb.	71	<i>Edwardsia</i> Neck.	593
<i>CYNAREAE</i> Less.	9	<i>Elachothamnus</i> DC.	368
<i>CYNAROCEPHALÆ</i> Vail.	4	<i>Electra</i> DC.	630
		<i>ELEPHANTOPEAE</i> Cass.	10, 85
<i>Dahlia</i> Cav.	494	<i>Elephantopus</i> Cass.	85
<i>Decachæta</i> DC.	133	<i>Elephantopus</i> Linn.	85, 87
<i>Decaneurum</i> DC.	66	<i>Elephantosis</i> Less.	87
<i>Déliia</i> Spreng.	502	<i>Eleutheranthera</i> Poit.	545
<i>Delucia</i> DC.	633	<i>Eleutherogeron</i> DC.	293
<i>Demetria</i> Lag.	314	<i>Elphegea</i> Cass.	318
<i>Denekia</i> Thunb.	462	<i>Elphegea</i> Less.	ibid.
<i>Denira</i> Adans.	529	<i>Elvira</i> Cass.	502
<i>Detridium</i> Nées.	223	<i>Enalcida</i> Cass.	542
<i>Detridium</i> DC.	ibid.	<i>Encelia</i> Adans.	566
<i>Detrioides</i> Less.	ibid.	<i>Enhydra</i> DC.	636
<i>Detris</i> Adans.	ibid.	<i>Enydra</i> Lour.	ibid.
<i>Detris</i> DC.	ibid.	<i>Enula</i> Duby.	464
<i>Dialesta</i> H. B. K.	76	<i>Epaltès</i> Cass.	461
<i>Dichrocephala</i> DC.	371	<i>Erato</i> DC.	317
<i>Diglossus</i> Cass.	640	<i>ERECTITEAE</i> DC.	500
<i>Dimorphanthès</i> Cass.	377	<i>Eremanthus</i> Less.	82
<i>Dimorphanthès</i> DC.	ibid.	<i>Eremia</i> DC.	77
<i>Diomedeæ</i> Cass.	488	<i>ERIGEREAE</i> DC.	211
<i>Diotstephus</i> Cass.	510	<i>Erigeron</i> Linn.	283
<i>Diplogon</i> Raf.	326	<i>Erigeron</i> Don.	432
<i>DIPLOPAPPEAE</i> DC.	211, 271	<i>Eriocarpa</i> Cass.	564
<i>Diplopappus</i> Cass. DC.	275	<i>ERIOCEPHALEAE</i> .	499
<i>Diplosastera</i> Tausch.	568	<i>Eriocoma</i> H. B. K.	ibid.
<i>Diplostephium</i> Cass.	272	<i>Eriophyllum</i> Lag.	656
<i>Diplostephium</i> Kunth.	273	<i>Eschembachia</i> Mœnch.	377
<i>Diplothrix</i> DC.	611	<i>Espejoa</i> DC.	660
<i>Discocalea</i> DC.	672	<i>Espeletia</i> Mutis.	516
<i>Discomela</i> Raf.	586	<i>Ethulia</i> Cass.	12
<i>Discopoda</i> DC.	604	<i>Ethulia</i> Gærtn.	461
<i>Distasis</i> DC.	279	<i>Ethulia</i> Nées.	11, 12
<i>Distephanus</i> Cass.	74	<i>ETHULIÆ</i> DC.	10
<i>Distreptus</i> Cass.	87	<i>Euageratum</i> DC.	108
<i>Districhum</i> Cass.	619	<i>EUAMELLEAE</i> DC.	211
<i>Dizonium</i> Willd.	482	<i>EUANTHEMIDEAE</i> DC.	499
<i>Dœllingeria</i> DC.	263	<i>Euplopappus</i> DC.	347
<i>Dœllingeria</i> Nées.	263, 272	<i>EUASTEREAE</i> DC.	211
<i>Donia</i> R. Br.	314	<i>Eubrachs</i> DC.	312
<i>Donioides</i> DC.	477	<i>Eucalia</i> Rœusch.	567
<i>Doria</i> Adans.	330	<i>Eucalydermos</i> DC.	669
<i>Dracopis</i> Cass.	558	<i>Eucephalophora</i> DC.	652
<i>Drimyphyllum</i> Bursh.	501	<i>Euclibadium</i> DC.	505
<i>Dubautia</i> Gaudich.	680	<i>EUCONYZEAE</i> DC.	375
<i>Duchesnia</i> Cass.	475	<i>Eudiplostephium</i> DC.	272
<i>Dugaldea</i> Cass.	662	<i>Eudysodia</i> DC.	632

NOMINUM ET SYNONYMORUM.

701

<i>Enclypta DC.</i>	490	<i>Francoeuria Cass.</i>	475
<i>Euerigeron DC.</i>	283	<i>Franseria Cav.</i>	524
<i>Euevax DC.</i>	458	<i>Fresenia DC.</i>	328
<i>EUGALINSOGEAE DC.</i>	499, 668	<i>Fulcadea Less.</i>	77
<i>Euganelia fl. mex.</i>	502	<i>Fullartonia DC.</i>	281
<i>Euharpalum DC.</i>	583		
<i>EUHELENIEAE DC.</i>	653	<i>Gaillardia Foug.</i>	651
<i>Euheliopsis DC.</i>	550	<i>GAILLARDIEAE DC.</i>	499, 651
<i>Euhymenoxys DC.</i>	661	<i>Galardia Lam.</i>	651
<i>EUINULEAE DC.</i>	463	<i>Galatea Cass.</i>	254
<i>Eujasonia DC.</i>	476	<i>Galatella Cass.</i>	ibid.
<i>Euliatris DC.</i>	128	<i>Galinsoga R. et P.</i>	677
<i>Eulychnophora DC.</i>	79	<i>Galinsogaea Zucc.</i>	ibid.
<i>Eumelampodium DC.</i>	518	<i>Galinsoga Kunth.</i>	678
<i>Eumonosia DC.</i>	77	<i>GALINSOGEAE DC.</i>	499, 668
<i>EUPATORIACEAE Less.</i>	8, 103	<i>Galordia Ræusch.</i>	651
<i>EUPATORIEAE DC.</i>	103	<i>Gamocarpha DC.</i>	2
<i>EUPATORIEAE Less.</i>	106	<i>Garuleum Cass.</i>	309
<i>Eupatorium Linn.</i>	141	<i>Geigera Less.</i>	482
<i>Eupatorium Auct.</i>	141, 187	<i>Geigeria Griess.</i>	ibid.
<i>Eupatoriophalacron Vaill.</i>	489	<i>Georgia Spreng.</i>	494
<i>EUPECTIDEAE DC.</i>	98	<i>Georgina Willd.</i>	ibid.
<i>Euphania DC.</i>	114	<i>GLOSSARIPHYTUM Neck.</i>	4
<i>Euphrosyne DC.</i>	530	<i>Glossocardia Cass.</i>	631
<i>Eupueria DC.</i>	104	<i>Glossogyne Cass.</i>	632
<i>Euporophyllum DC.</i>	648	<i>Gluphiphyllaea DC.</i>	671
<i>Eupyrocoma DC.</i>	350	<i>Glutinaria Comm.</i>	318
<i>Eurybia Cass.</i>	265	<i>Glutinaria DC.</i>	ibid.
<i>Eurybia Nées.</i>	264	<i>GNAPHALIEAE.</i>	500
<i>Eurybiopsis DC.</i>	260	<i>Gnaphalium Vaill.</i>	458
<i>EUSENECIONEAE DC.</i>	500	<i>Gnaphalodes Tourn.</i>	460
<i>Eusphæranthus DC.</i>	369	<i>Graenia Hook.</i>	661
<i>Euthamia Cass.</i>	330	<i>Grangea Adans.</i>	372
<i>Euthamia DC.</i>	341	<i>GRANGEINEAE DC.</i>	371
<i>Euthamia Nutt.</i>	ibid.	<i>Grindelia Willd.</i>	314
<i>Eutarchonanthus DC.</i>	431	<i>Guardiola Steud.</i>	511
<i>Eutragoceras DC.</i>	533	<i>Guardiola H. B.</i>	ibid.
<i>EUTUSSILAGINEAE DC.</i>	207	<i>Guizotia Cass.</i>	551
<i>EUVERNONIEAE DC.</i>	10	<i>Gundelia Tourn.</i>	88
<i>Euxanthioides DC.</i>	524	<i>Gundelsheimera Cass.</i>	ibid.
<i>Euxanthium DC.</i>	523	<i>Guttierrezia Lag.</i>	653
<i>Euxenia Cham.</i>	501	<i>Gymnactis DC.</i>	317
<i>EUXENIEAE DC.</i>	498, 500	<i>Gymnanthemum Cass.</i>	15, 67
<i>Evax Gärtn.</i>	458	<i>Gymnanthemum DC.</i>	67
		<i>Gymnarhena Desf.</i>	374
<i>Fœna Spreng.</i>	612	<i>Gymnocoronis DC.</i>	106
<i>Felicia Cass.</i>	218	<i>Gymnolæna DC.</i>	641
<i>Felicia DC.</i>	ibid.	<i>Gymnoloma Ker.</i>	563
<i>Ferdinanda Lag.</i>	552	<i>Gymnolomia H. B. K.</i>	561
<i>Filago Willd.</i>	458	<i>Gymnopsis DC.</i>	561
<i>Fimbrillaria Cass.</i>	387	<i>Gymnosperma Less.</i>	311
<i>Fimbrillaria DC.</i>	ibid.	<i>GYMNOSPERMEAE DC.</i>	212, 310
<i>Flaveria Juss.</i>	635	<i>Gymnostephium Less.</i>	300
<i>Flaveria Juss.</i>	104	<i>Gynema Raf.</i>	450
<i>FLAVERIEAE Less.</i>	498, 605	<i>Gyneheteria Willd.</i>	456
<i>FLOSCULOSÆ Tourn.</i>	4	<i>Gyneteria Spreng.</i>	ibid.
<i>Florestina Cass.</i>	655		
<i>Flourensia DC.</i>	590	<i>Hacub Vaill.</i>	88
<i>Fougéria Moench.</i>	509	<i>Hamulium Cass.</i>	ibid.
<i>Fougerouxia DC.</i>	ibid.	<i>Hamulium DC.</i>	ibid.

<i>Haplostephium Mart.</i>	78	<i>Hysterophorus Vaill.</i>	531
<i>Harpalium Cass.</i>	583		
<i>Harpalixia DC.</i>	584	<i>Ichthyothere Mart.</i>	504
<i>Hartmannia DC.</i>	693	<i>Inula Gaertn.</i>	464
<i>Haxtonia Hortul.</i>	271	<i>Inula Linn.</i>	463, 466, 467
<i>Haynea Spreng.</i>	16	INULEAE <i>Cass.</i>	213, 462
<i>Haynea Willd.</i>	14	<i>Inulopsis DC.</i>	349
<i>Hebecarpa DC.</i>	219	<i>Iphiona Cass.</i>	475
<i>Hebeclinium DC.</i>	136	<i>Isocarpha R. Br.</i>	106
<i>Hebephora DC.</i>	453	<i>Isocarphoides DC.</i>	107
<i>Hecubaea DC.</i>	665	<i>Isoneria Wight.</i>	23
<i>Hectorea DC.</i>	95	<i>Isonema Cass.</i>	66
<i>Heleastrum DC.</i>	263	<i>Isostigma Less.</i>	634
HELENIEAE <i>Cass.</i>	499, 651	<i>Isotrichia DC.</i>	82
<i>Helenia Linn.</i>	666	<i>Iva Linn.</i>	529
<i>Helenia DC.</i>	<i>ibid.</i>	IVEAE <i>Cass.</i>	498, 527
<i>Helenium Cass.</i>	<i>ibid.</i>	<i>Ixauchenus Cass.</i>	308
<i>Helenium Linn.</i>	665		
HELIANTHEAE <i>Less.</i>	498, 534	<i>Jægeria H. B. K.</i>	543
<i>Helianthus Linn.</i>	585	<i>Jasonia Cass.</i>	476
<i>Helichroa Raf.</i>	554	<i>Jaumea Pers.</i>	663
HELICHRYSÆAE.	500		
HELIOPSIDAE <i>DC.</i>	458, 534	<i>Kahiria Forsk.</i>	12
<i>Heliopsis Pers.</i>	550	<i>Kalimeris Cass.</i>	258
<i>Helioptalum Raf.</i>	555	<i>Kallias Cass.</i>	550
<i>Hemiachyris DC.</i>	313	<i>Kallias DC.</i>	<i>ibid.</i>
<i>Hemizonia DC.</i>	692	<i>Karelinia Less.</i>	375
<i>Henanthus Less.</i>	356	<i>Kaulfussia Nées.</i>	300
<i>Henotogyna DC.</i>	432	<i>Keerlia DC.</i>	309
<i>Henricia Cass.</i>	262	<i>Kerneria Mönch.</i>	598
<i>Herderia Cass.</i>	13	<i>Kleinia Jacq.</i>	647
<i>Heterochæta DC.</i>	282	<i>Kleinia Juss.</i>	663
<i>Heterocoma DC.</i>	14	<i>Kugaia DC.</i>	650
HETEROCOMEAE <i>DC.</i>	<i>ibid.</i>	<i>Kuhnia Linn.</i>	156
HETEROPAPPEAE <i>DC.</i>	212, 297	<i>Kyberia Neck.</i>	304
<i>Heteropappus Less.</i>	297		
<i>Heterosperma Cav.</i>	632	<i>Laennecia Cass.</i>	376
<i>Heterospermum Willd.</i>	<i>ibid.</i>	<i>Laestadia Kunth.</i>	374
HETEROTHALAMEAE <i>DC.</i>	216	<i>Lagasca DC.</i>	91
<i>Heterothalamus Less.</i>	<i>ibid.</i>	<i>Lagasca Cav.</i>	<i>ibid.</i>
<i>Heterotheca Cass.</i>	316	<i>Lagascea H. B. K.</i>	<i>ibid.</i>
HETEROTHECEAE <i>DC.</i>	212, 316	<i>Lagenifera Cass.</i>	307
<i>Hidalgoa Less.</i>	511	<i>Lagenophora Cass.</i>	<i>ibid.</i>
<i>Hidalgoa Lallav.</i>	521	<i>Lasianthæa DC.</i>	607
<i>Hingcha Roxb.</i>	636	<i>Lasianthus Zucc.</i>	<i>ibid.</i>
HIPPIEAE.	499	<i>Lasthenia Cass.</i>	664
<i>Hololepis DC.</i>	16	<i>Latreillea DC.</i>	504
<i>Holophyllæa DC.</i>	670	<i>Lavenia Swartz.</i>	110
<i>Homochroma DC.</i>	324	<i>Lavenia Wall.</i>	280
<i>Homogyne Cass.</i>	204	<i>Laxmannia Forst.</i>	501
<i>Horebeckia Corn.</i>	346	<i>Leachea Cass.</i>	560
<i>Hoplophyllum DC.</i>	73	<i>Lebetina Cass.</i>	639
<i>Hopkirkia DC.</i>	660	<i>Leichæmium DC.</i>	348
<i>Hopkirkia Spreng.</i>	493	<i>Leighia Cass.</i>	580
<i>Hunteria fl. mex.</i>	649	<i>Leighia Scop.</i>	12
<i>Hunteria DC.</i>	<i>ibid.</i>	<i>Lejica Hill.</i>	634
<i>Hybridella Cass.</i>	554	<i>Leiogonia DC.</i>	120
<i>Hymenatherum Cass.</i>	641	<i>Lemmatium DC.</i>	669
<i>Hymenopappus L'Hér.</i>	658	<i>Leontophthalmum Less.</i>	675
<i>Hymenoxys Cass.</i>	661	<i>Leontophthalmum Willd.</i>	522


<i>Lepachys</i> Raf.	558	<i>Meyeria</i> DC.	670
<i>Lepachys</i> DC.	558	<i>Micractis</i> DC.	619
<i>Lepidaploa</i> Cass.	15, 26	<i>Micrelum</i> Forsk.	489
<i>Lepidaploa</i> DC.	26	<i>Microglossa</i> DC.	320
<i>Lepidopappus</i> fl. mex.	655	<i>Microgyne</i> Less.	296
<i>Lepidophyllum</i> Cass.	314	<i>Micropsis</i> DC.	459
<i>Lepidopogon</i> Tausch.	458	<i>Micropus</i> Linn.	460
<i>Leptocarpha</i> DC.	495	<i>Microtrichia</i> DC.	366
<i>Leptocoma</i> Less.	280	<i>Mieria</i> Lallav.	654
<i>Leptoderis</i> DC.	373	<i>Mikania</i> Willd.	187
<i>Leptogyne</i> Ell.	450	<i>Milleria</i> Cass.	503
<i>Leptopoda</i> Nutt.	653	<i>MILLERIEE</i> DC.	498, 502
<i>Leptospermoides</i> DC.	17	<i>Minuria</i> DC.	298
<i>Leptosyne</i> DC.	531	<i>Mocinna</i> Lag.	671
<i>Leptothamnus</i> DC.	367	<i>Molina</i> R. et Pav.	398
<i>Lessingia</i> Cham.	351	<i>Molpadia</i> Cass.	485
<i>Leucopsis</i> DC.	348	<i>Monactis</i> H. B. K.	546
<i>LEYSSERIEE</i> .	500	<i>Monarrhenus</i> Cass.	457
<i>LIABEAE</i> Cass.	94	<i>Monenteles</i> Labill.	455
<i>Liabum</i> Adans.	96	<i>Monodonta</i> DC.	559
<i>Liatrix</i> Schreb.	128	<i>Monogyria</i> DC.	325
<i>Limbarda</i> Adans.	470	<i>Monosis</i> DC.	77
<i>Limbarda</i> DC.	ibid.	<i>Montagnæa</i> DC.	564
<i>Linosyris</i> DC.	351	<i>Montanoa</i> Lallav.	ibid.
<i>Lipochæta</i> DC.	610	<i>Munychia</i> Cass.	222
<i>Lipotriche</i> R. Br.	544	<i>MUTISIAEAE</i> .	9
<i>Lipotriche</i> Less.	610	<i>Myriactis</i> Less.	308
<i>Locheria</i> Neck.	612	<i>Myriadenus</i> Cass.	476
<i>Lorentea</i> Less.	101, 102	<i>Nardosmia</i> Cass.	205
<i>Lorentea</i> Lag.	98	<i>Narvalina</i> Cass.	633
<i>Lorentea</i> Ort.	628	<i>NASSAUVIACEAE</i> .	9
<i>Lychnocephalus</i> Mart.	83	<i>Nauemburgia</i> Willd.	635
<i>Lychnophora</i> Mart.	78	<i>Nauplius</i> Cass.	486
<i>Lychnophoroides</i> DC.	ibid.	<i>Needhamia</i> Cass.	633
<i>Machæranthera</i> Nées.	261	<i>Neja</i> Don.	325
<i>Madaria</i> DC.	691	<i>Neuractis</i> Cass.	631
<i>Madarioides</i> DC.	692	<i>NEUROLAENEAE</i> .	500
<i>Madaroglossa</i> DC.	694	<i>Nidorella</i> Cass.	321
<i>Madea</i> Sol.	301	<i>Niebuhria</i> Scop.	509
<i>Madia</i> Mol.	691	<i>Nocca</i> Cass.	92
<i>Madia</i> auct.	691-692	<i>Noccea</i> DC.	ibid.
<i>MADIEAE</i> DC.	499, 691	<i>Noccea</i> Jacq.	91
<i>Mahometia</i> DC.	467	<i>Nolletia</i> Cass.	366
<i>Mairia</i> DC.	217	<i>Noticastrum</i> DC.	279
<i>Mairia</i> Nées.	ibid.	<i>Nothites</i> Cass.	186
<i>Margarita</i> Gaud.	226	<i>Obeliscaria</i> Cass.	558
<i>Marshallia</i> Schreb.	680	<i>Obeliscotheca</i> Vail.	555
<i>Matamoria</i> Lallav.	87	<i>Odontocarpha</i> DC.	71
<i>Meisteria</i> Scop.	14	<i>Odontoloma</i> H. B. K.	11
<i>MELAMPODIEAE</i> Less.	498, 517	<i>Odontospermum</i> Neck.	486, 488
<i>MELAMPODINEAE</i> Less. DC.	497, 500	<i>Ogiera</i> Cass.	545
<i>Melampodium</i> Linn.	517	<i>Ogiera</i> Spreng.	501
<i>Melanthera</i> Rich.	544	<i>Oiospermum</i> Less.	11
<i>Melanthera</i> Rohr.	ibid.	<i>Olearia</i> Manch.	271
<i>Melanodendron</i> DC.	279	<i>Oligactis</i> Cass.	95
<i>Mendezia</i> DC.	532	<i>Oligactis</i> DC.	ibid.
<i>Meratia</i> Cass.	502	<i>Oligarion</i> Cass.	681
<i>Mesoretra</i> Raf.	666	<i>Oliganthes</i> Cass.	15
<i>Meyera</i> Schreb.	636		

<i>Oligogyne</i> DC.	629	<i>Pingraea</i> Cass.	398, 420
<i>Oligolepis</i> Cass.	370	<i>Pinillosia</i> Ossa.	528
<i>Olocarpha</i> DC.	692	<i>Piptocoma</i> Cass.	74
<i>Ooclinium</i> DC.	133	<i>Piqueria</i> Cav.	104
<i>Orsinia Bertol.</i>	104	<i>Piquerioides</i> DC.	105
<i>Orsina</i> Bertol.	476	<i>Pirarda</i> Adans.	12
<i>Oswalda</i> Cass.	505	<i>Pithecoseris</i> Mart.	84
<i>Oswalda</i> DC.	ibid.	<i>Platclypta</i> DC.	491
<i>Oxylobus</i> Moq.	115	<i>Platycarpæa</i> DC.	594
<i>Oxylobus</i> DC.	ibid.	<i>Platycarpha</i> Less.	71
<i>Oxypappus</i> DC.	661	<i>Platypteris</i> H. B. K.	617
<i>Oxyura</i> DC.	693	<i>Platypteris</i> DC.	ibid.
<i>Oyedæa</i> DC.	576	<i>Pleionactis</i> DC.	95
<i>Pachyderis</i> Cass.	360	<i>Pluchea</i> Cass.	449
<i>Pachyderis</i> DC.	ibid.	PLUCHEINEAE Cass.	432
<i>Pacourina</i> Aubl.	14	<i>Pluridens</i> Neck.	503
<i>Pacourinopsis</i> Cass.	ibid.	<i>Podanthus</i> Lag.	501
<i>Palafoxia</i> Lag.	124	<i>Podocoma</i> Cass.	260
<i>Paleolaria</i> Cass.	ibid.	<i>Podoneja</i> DC.	325
<i>Pallasia</i> L'Hér.	566	<i>Pollalesta</i> H. B. K.	15
<i>Pallenis</i> Cass.	487	<i>Polylepis</i> Less.	218
<i>Paquerina</i> Cass.	307	<i>Poloa</i> DC.	275
PARTHENIEAE Less.	498, 530	<i>Polyactidium</i> D. C.	281
<i>Partheniastrium</i> Niss.	531	<i>Polyactis</i> Less.	ibid.
<i>Partheniastrium</i> DC.	532	<i>Polyarrhena</i> Cass.	218
<i>Parthenichæta</i> DC.	ibid.	<i>Polycephalos</i> Forsk.	370
<i>Parthenium</i> Linn.	531	<i>Polycephalos</i> DC.	ibid.
<i>Pascalina</i> Ort.	549	<i>Polymnia</i> Linn.	514
PECTIDIEAE DC.	10, 94	<i>Polymniastrium</i> Lam.	514, 515
PECTIDIEÆ Less.	98	<i>Polypappus</i> Less.	398
<i>Pectidium</i> Less.	ibid.	<i>Polypteris</i> Nutt.	659
<i>Pectidopsis</i> DC.	ibid.	<i>Polypteris</i> Less.	660
<i>Pectis</i> Less.	ibid.	POROPHYLLEAE DC.	499, 647
<i>Pectis</i> Linn.	ibid.	<i>Porophyllum</i> Vaill.	ibid.
<i>Pectis</i> Cass.	ibid.	<i>Praxelis</i> Cass.	133
<i>Pectinellum</i> DC.	109	<i>Priestleya</i> fl. mex.	564
<i>Pegolettia</i> Cass.	481	<i>Pronacron</i> Cass.	508
<i>Pentameranthes</i> DC.	495	<i>Proteopsis</i> Mart.	16
<i>Pentanema</i> Cass.	474	<i>Proustia</i> Spreng.	30
<i>Peramibus</i> Raf.	568	<i>Psathurochæta</i> DC.	609
<i>Persoonia</i> Michx.	680	<i>Pseudo-elephantopus</i> Rohr.	87
<i>Perymenium</i> Schrad.	608	<i>Pseudevax</i> DC.	459
<i>Petrobium</i> R. Br.	501	<i>Psiadia</i> Jacq.	318
PETASITEAE DC.	204	PSIADIEAE DC.	212, 317
<i>Petasites</i> Tourn.	206	<i>Pailocarpæa</i> DC.	596
<i>Phagnalon</i> Cass.	396	<i>Pterigeron</i> DC.	203
<i>Phalacræa</i> DC.	105	<i>Pterocaulon</i> Ell.	453
<i>Phalacroloma</i> Cass.	207	<i>Pterocaulopsis</i> DC.	454
<i>Phalacromesus</i> Cass.	457	<i>Pteronia</i> Linn.	356
<i>Phania</i> DC.	114	<i>Pteronia</i> Linn. f.	73
<i>Phialis</i> Spreng.	656	<i>Pteropappus</i> Less.	217
<i>Philactis</i> Schrad.	534	<i>Pteropappus</i> DC.	ibid.
<i>Philoglossa</i> DC.	567	<i>Pterophora</i> Neck.	364
<i>Phyllinema</i> Blum.	630	<i>Pterophorus</i> Cass.	ibid.
<i>Phyllocephalum</i> Blum.	66	<i>Pterophorus</i> DC.	ibid.
<i>Phyllocephalum</i> DC.	ibid.	<i>Pterophyton</i> Cass.	575
<i>Phylloneja</i> DC.	325	<i>Ptilostephium</i> H. B. K.	678
<i>Phyteumopsis</i> Juss.	680	<i>Pycnocephalum</i> DC.	83
<i>Pioradenia</i> Hook.	685	<i>Pyrarda</i> Cass.	372
		<i>Pyrarda</i> DC.	373

<i>Pyrethrum</i> Med.	620	<i>Skirroglyphis</i> DC.	533
<i>Pyrochæta</i> DC.	349	<i>Sobreyra</i> R. et P.	636
<i>Pyrocoma</i> Hook.	350	<i>Sobrya</i> Pers.	ibid.
<i>Pulicaria</i> DC.	477	<i>Sogalgina</i> Cass.	678
<i>Pulicaria</i> Gærtn.	478	<i>Sogaligna</i> Steud.	ibid.
		<i>Solenogyne</i> Cass.	367
<i>RADIATÆ</i> Tourn.	4	<i>SOLENOGYNEÆ</i> DC.	212, 365
<i>Ramilla</i> DC.	551	<i>SOLIDAGINEÆ</i> DC.	212, 328
<i>Rancagua</i> Poepp. et Endl.	664	<i>Solidago</i> Linn.	330
<i>Ratibida</i> Raf.	558	<i>Sommea</i> Bory.	3
<i>Ratibida</i> DC.	559	<i>Sommerfeltia</i> Less.	302
<i>RELHANIEÆ</i> Less.	500	<i>Sparganophorus</i> Vaill.	11
<i>Rhanterium</i> Desf.	463	<i>Sparganophorus</i> Michx.	114
<i>Rhinactina</i> Less.	275	<i>Spermophylla</i> Neck.	681
<i>Rhynchospermum</i> Reinw.	206	<i>SPHAERANTHÆÆ</i> DC.	212, 368
<i>Ridan</i> Adans.	575	<i>Sphæranthoides</i> Cunn.	456
<i>Riencourtia</i> Cass.	503	<i>Sphæranthus</i> Linn.	369
<i>Rochonia</i> DC.	345	<i>Sphenogyne</i> R. Br.	681
<i>Rolandra</i> Rottb.	90	<i>SPHENOGYNEÆ</i> DC.	499, 681
<i>Rolfinkia</i> Zenk.	696	<i>Spilanthus</i> Jacq.	620
<i>Rosalesia</i> .	183	<i>Spilanthus</i> Cass.	624
<i>Rosilla</i> Less.	668	<i>Spilanthus</i> Linn.	620
<i>Rothia</i> Lam.	658	<i>Spilanthus</i> Pers.	624
<i>Rudbeckia</i> Cass.	555	<i>Spiracantha</i> H. B. K.	90
<i>Rudbeckia</i> Auct.	554, 558	<i>Spixia</i> Schrank.	70
<i>RUDBECKIÆÆ</i> Cass.	498, 554	<i>Spongotrichum</i> Nées.	265
<i>Rumfordia</i> DC.	549	<i>Spongotrichum</i> DC.	266
		<i>Stachyanthus</i> DC.	84
<i>Sabazia</i> Cass.	496	<i>Stammarium</i> Willd.	102
<i>Salivaria</i> DC.	624	<i>Stammarium</i> DC.	ibid.
<i>Salmea</i> DC.	493	<i>Starkia</i> Juss.	96
<i>Sanvitalia</i> Gualt.	628	<i>Starkea</i> Willd.	ibid.
<i>Sarcanthemum</i> Cass.	367	<i>Steiractis</i> DC.	345
<i>Scepinia</i> Neck.	356	<i>Stemmodontia</i> Cass.	538
<i>Scepinia</i> DC.	ibid.	<i>Stenactis</i> Less.	298
<i>Schizogyne</i> Cass.	473	<i>Stenactis</i> Cass.	281
<i>Schlechtendalia</i> Willd.	638	<i>Stephananthus</i> Lebm.	308
<i>Schistocarpa</i> Less.	608	<i>Stevia</i> Cav.	115
<i>Schkuhria</i> Roth.	654	<i>Stilpnopappus</i> Mart.	75
<i>Schkuhria</i> Mœnch.	496	<i>Stobæa</i> Spreng.	71
<i>Sclerocarpus</i> Jacq.	566	<i>Stokesia</i> L'Hér.	ibid.
<i>Sclerolepis</i> Cass.	114	<i>Strabonia</i> DC.	481
<i>Scolospermum</i> Less.	509	<i>Strigia</i> DC.	126
<i>Sebastiania</i> Bertol.	630	<i>Strobocalyx</i> Blume.	21
<i>Selloa</i> H. B. K.	612	<i>Strophopappus</i> DC.	75
<i>SEMIFFLOSCULOSÆ</i> Tourn.	4	<i>Struchium</i> P. Br.	11
<i>SENECIONEÆ</i> Less.	500	<i>Stylimnus</i> Raf.	460
<i>SENECIONIDEÆ</i> Less.	8, 497	<i>Stylimnus</i> DC.	ibid.
<i>Sergilus</i> Gærtn.	398	<i>Suprago</i> Gærtn.	128, 129
<i>Sericocarpus</i> Nées.	261	<i>Suprago</i> DC.	129
<i>SERIPHIEÆ</i> DC.	500	<i>SYNANTHERÆ</i> Rich.	4
<i>Shawia</i> Forst.	78	<i>SYNGENESIA POLYGAMIA</i> Linn.	ib.
<i>Sideranthus</i> Fras.	347	<i>Synchodendron</i> Bojer.	92
<i>Siegesbeckia</i> Linn.	495	<i>Synedrella</i> Gærtn.	629
<i>SILPHIÆÆ</i> Less.	498, 510		
<i>Silphium</i> Linn.	511	<i>TAGETEÆ</i> DC.	499, 638
<i>Sinblocline</i> DC.	297	<i>Tagetes</i> Linn.	642
<i>Simsia</i> Pers.	577	<i>TAGETINEÆ</i> Cass.	499, 638
<i>SIPHONIPHYTUM</i> Neck.	4	<i>TARCHONANTHÆÆ</i> Less.	213, 499

<i>Tarchonanthus</i> Linn.	431	<i>Urolepis</i> DC.	136
<i>Tecmarsi</i> DC.	93	<i>Ursinia</i> Gärtn.	688
<i>Telekia</i> Baumg.	485	<i>Ursinia</i> Auct.	681, 689
<i>Tetramolopium</i> Nées.	262		
<i>Tephrodes</i> DC.	24	<i>Vanillosma</i> Less.	18
<i>Tessaria</i> R. et P.	456	<i>Vargasia</i> DC.	676
<i>Tetracarpum</i> Mönch.	654	<i>Varthemia</i> DC.	473
<i>Tetragonotheca</i> Dill.	552	<i>Verbesina</i> Less.	613
<i>Tetrantha</i> Poit.	503	<i>Verbesinaria</i> DC.	ibid.
<i>Tetranthus</i> Swartz.	528	<i>VERBESINEAE</i> Less.	498, 607
<i>Tetrodus</i> Cass.	666	<i>Vermifuga</i> R. et Pav.	
<i>Tetrodus</i> DC.	ibid.	<i>Vernonia</i> Schreb.	15
<i>Tetraotis</i> Blum.	637	<i>Vernonia</i> Less.	14, 74
<i>Thelesperma</i> Less.	633	<i>VERNONIACEAE</i> Less.	8, 9
<i>Thelythamnus</i> Less.	683	<i>VERNONIEAE</i> Cass.	9
<i>Thelythamnus</i> DC.	681	<i>VERNONIEAE</i> DC.	10
<i>Therogeron</i> DC.	283	<i>Vicoa</i> Cass.	474
<i>Thespis</i> DC.	375	<i>Viguiera</i> H. B. K.	578
<i>Thouarsia</i> Vent.	318	<i>Viguiera</i> Less.	ibid.
<i>Thymophylla</i> Lag.	647	<i>Villanova</i> Ort.	531
<i>Tilesia</i> Mey.	549	<i>Virga-aurea</i> Tourn.	330
<i>Tithonia</i> Desf.	584	<i>Virgaurea</i> DC.	ibid.
<i>Tragantes</i> Wallr.	176	<i>Virgilia</i> L'Hér.	651
<i>Tragoceras</i> Less.	533	<i>Vittadinia</i> Rich.	280
<i>Tragoceros</i> H. B. K.	ibid.	<i>Viviania</i> Willd.	95
<i>Trattenickia</i> Pers.	680	<i>Vosacan</i> Adans.	585
<i>Trianthæa</i> DC.	23		
<i>Trichogonia</i> DC.	126	<i>Wahlenbergia</i> Schum.	497
<i>Trichophyllum</i> Nutt.	656	<i>Webbia</i> DC.	72
<i>Trichospermum</i> Beauv.	531	<i>Wedelia</i> Jacq.	538
<i>Trichospira</i> H. B. K.	90	<i>Wedelia</i> Auct.	545, 546
<i>Trichostemma</i> Cass.	538	<i>Wiborgia</i> Roth.	677
<i>Trichostephium</i> Cass.	540	<i>Wickstræmia</i> Spreng.	140
<i>Tridax</i> Linn.	679	<i>Wildenowia</i> Cav.	638
<i>Trigonospermum</i> Less.	508	<i>Wollastonia</i> DC.	546
<i>Trilisa</i> Cass.	128, 131	<i>Woodvillea</i> DC.	318
<i>Trilisa</i> DC.	131	<i>Wulfia</i> Neck.	563
<i>Trimeranthes</i> Cass.	496		
<i>Trimeranthes</i> DC.	ibid.	<i>Xanthioides</i> DC.	521
<i>Trimorphæa</i> Cass.	290	<i>Xanthisma</i> DC.	94
<i>Trimorphæa</i> DC.	ibid.	<i>Xanthium</i> Linn.	522
<i>Tripolium</i> Nées.	253	<i>Xanthocoma</i> H. B. K.	310
<i>Trixidium</i> DC.	606	<i>Xantholepis</i> Willd.	97
<i>Trixis</i> Swartz.	505	<i>Xenismia</i> DC.	509
<i>Tropidolepis</i> Tausch.	216	<i>Xerolepis</i> Less.	687
<i>Tubifera</i> DC.	386	<i>Xerolepis</i> DC.	ibid.
<i>Tubilium</i> Cass.	480	<i>Ximenesia</i> Cav.	627
<i>Tubilium</i> DC.	480	<i>Ximenesia</i> Less.	ibid.
<i>Turczaninovia</i> DC.	257	<i>Xerothermus</i> DC.	311
<i>Turpinia</i> Lallav.	77	<i>Zaluzania</i> Pers.	553
<i>Tursenia</i> Cass.	398	<i>Zarabellia</i> Cass.	519
<i>TUSSLAGINEAE</i> Less.	103, 204	<i>Zarabellia</i> DC.	ibid.
<i>Tussilago</i> Tourn. Cass.	208	<i>Zexmenia</i> Lallav.	610
<i>Tussilago</i> Linn.	204, 208	<i>Zeyhera</i> Spreng.	482
		<i>Zinnia</i> Linn.	534
<i>Ucacea</i> Cass.	492	<i>Zyrphelis</i> Nées.	217, 218
<i>Ucacou</i> Adans.	629	<i>Zyrphelis</i> DC.	218
<i>Unxia</i> Linn. f.	507		





Stanford University Libraries



3 6105 011 548 166

